1   Type of article: research article
2
3   Running header: A globally invasive sponge
4
5   **The marine sponge *Hymeniacidon perlevis* is a globally-distributed invasive species**
6

7   **Thomas L. Turner**
8   Ecology, Evolution, and Marine Biology Department
9   University of California, Santa Barbara
10   Santa Barbara, CA 93106
11
12   tlturner@ucsb.edu
13
ORCiD: 0000-0002-1380-1099

Abstract
In Elkhorn Slough, a tidal estuary draining into Monterey Bay, California, the intertidal is
dominated by an orange sponge known by the name *Hymeniacidon sinapium*. This same
species is common in the rocky intertidal in California, and fresh collections described
here find this species throughout mainland kelp forest in Southern California. Farther
afield, morphologically and ecologically indistinguishable sponges are among the most
conspicuous occupants of estuaries and intertidal areas in Asia, Europe, South America,
and Africa. Here I use morphological, ecological, and genetic data to show that these
incredibly abundant sponges are all members of the same globally-distributed species,
which should be known by the senior synonym *H. perlevis*. Though previous authors
have remarked upon the morphological, ecological, and/or genetic similarity of various
distant populations, the true scope of this sponge's distribution appears to be
unrecognized or unacknowledged in the literature. Limited larval dispersal, historically
documented range expansion, and low genetic variation all support a hypothesis that this
sponge has achieved its extraordinary range via human-mediated dispersal, making it the
most widely-distributed and abundant invasive sponge known to date.

32
33   **Keywords**
34   Porifera, sponges, invasive species, kelp forest, intertidal, estuaries
35

36   **Declarations**
37   Conflicts of interest/Competing interests: none to declare
38
Availability of data and material: All raw data is included as supplementary files;
georeferenced collection data is available as a supplementary .xls file; genetic data are
archived at Genbank; specimen vouchers are archived at the California Academy of
Sciences and at the Natural History Museum of Los Angeles; specimen photos will be
made available as supplementary files, are also archived by the associated museums in
GBIF, and are posted as georeferenced data on iNaturalist.org.
45
46   Code availability: n/a

Introduction

In coastal marine ecosystems, filter-feeding marine invertebrates are among the most important invasive species in terms of species diversity, biomass, and ecological impacts (Ruiz et al. 2000; Bax et al. 2003; Byrnes and Stachowicz 2009). Sponges (phylum Porifera) are a diverse group of filter-feeding organisms that are found in all marine environments. They provide unique ecosystems services (and potential disruptions) because they preferentially consume the smaller size fractions of the plankton, such as bacterial plankton (Reiswig 1971; Maldonado et al. 2012). They can also have major affects on nutrient cycling, as some sponges convert dissolved nutrients into particulate matter available to other animals (de Goeij et al. 2013).

Our understanding of invasive sponges has been limited by an incomplete taxonomy. Sponges have simpler morphologies than most animals, confounding traditional classification schemes (Morrow and Cárdenas 2015). Many species were initially described as having a wide geographic range, but in recent decades these taxa have been recognized as clades comprised of multiple species with similar morphologies (Knowlton 1993; Xavier et al. 2010). This is consistent with what is known about larval dispersal in sponges. All known sponge larvae are lecithotrophic, meaning that they have no ability to feed until they settle and develop into juveniles (Maldonado 2006). They have a short planktonic life, lasting from minutes to a few days (Maldonado 2006). Some sponges, however, do seem to have broad geographic ranges, and this is likely due to human-mediated transport. Carballo et al. (2013) list 7 species thought to have recent range expansions, including two that have moved between the Pacific and Atlantic basins (Carballo et al. 2013). Some of these species are likely to have been accidentally introduced with aquaculture (Henkel and Janussen 2011; Fuller and Hughey 2013). Trawling, hull-fouling, and other actives also likely play a role (Carballo et al. 2013).

In the current work, I describe what appears to be the most common and widely distributed invasive sponge known to date. Genetic and morphological data support a distribution that includes Europe, the Atlantic coasts of North and South America, the Pacific coast of North America, and Asia. Morphological data suggest it is also present in New Zealand, Southwest Africa, and the Pacific coast of South America, but genetic data are not yet available from these populations. In much of this range, it is among the most common sponges in multiple habitats. In Europe, this species is known as *Hymeniacidon perlevis* (Montagu 1814). The range of *H. perlevis* was already thought to be substantial: from Norway in the North to the Macronesian Islands off Africa in the South (Erpenbeck and Van Soest 2002). Within this range it is found in diverse habitats, including both the intertidal and the subtidal, and it can grow buried in sediment or on hard substrate (Erpenbeck and Van Soest 2002). It is often abundant in these habitats, and is considered to be one of the most common sponges in Europe (Erpenbeck and Van Soest 2002). It has been described by other taxonomists as also occurring in New Zealand (Bergquist 1961, 1970) and as the most abundant intertidal sponge in Western South Africa (Samaai and Gibbons 2005), but these records were rejected from the consensus view (Van Soest et al. 2020a), probably because limited dispersal ability seemed to make such a range implausible. Sponges from additional parts of the globe have been described as morphologically indistinguishable from *H. perlevis,* but in these cases taxonomists put

forth other names for these distant populations. For example, de Laubenfels described a
sponge he named *Hymeniacidon sinapium* from California in 1930 (de Laubenfels 1930,
1932). He acknowledged that "it is doubtful whether this is a new form", and went so far
as to suggest that species with the names "*sanguinea, luxurians, caruncula, heliophila,*
sinapium, and perhaps even more species" are in fact synonyms. Consistent with this
prediction, the European species *sanguinea* and *caruncula* have been synonymized with
H. perlevis (Van Soest et al. 2020a). The status of *H. luxurians* is unclear (Van Soest et al.
2020b), but the other two species, *H. sinapium* and *H. heliophila,* are still considered
valid. In the current work, I will present evidence that *H. sinapium* is a junior synonym of
H. perlevis, and that most sponges placed under the name *H. heliophila* are also *H.*
perlevis.
104
When describing *H. sinapium* in California, de Laubenfels remarked on its impressive
ecological breadth. He described it as abundant in the "surf-beaten" intertidal throughout
Southern California, but also the most abundant sponge on the oyster beds in Newport
Bay (de Laubenfels 1932). (He reported only one sample from the subtidal, but his
subtidal sampling was limited, primarily via trawling.) In contrast to this abundance in
Southern California, de Laubenfels was only able to locate a single specimen of this
species in Central California, so he described it as common only in the South. This is
notable because he was based at Hopkins Marine Station in Monterey Bay (Central
California), and this was the area that he studied most comprehensively. This makes it
unlikely that this species was present in large numbers in Central California in the 1930s.
Subsequently, however, it has become a dominant species in intertidal portions of
Elkhorn Slough, which drains into Monterey Bay (Wasson et al. 2001), and it is also
known from San Francisco Bay and Tomales Bay in Northern California (Wasson et al.
2001; Fuller and Hughey 2013). Morphological (Sim 1985) and genetic (Hoshino et al.
2008) comparisons later confirmed that a common *Hymeniacidon* in Korea, Japan, and
China were the same species as those in California, so it was proposed that *H. sinapium*
was introduced to California from Asia with oyster mariculture (Fuller and Hughey 2013).
Though this is certainly possible, the data I compile here illustrates that it may also be
non-native in Asia. This species has been said to occur in the Mexican Pacific (Hofknecht
1978) and the Galapagos Islands (Desqueyroux-Faúndez and Van Soest 1997) as well,
but genetic data are not yet available from those populations.
126
The final species to consider, *H. heliophila* (Wilson 1911), is ascribed a substantial range
in the Western Atlantic, from the Gulf of Maine to Argentina (Weigel and Erwin 2016;
Van Soest et al. 2020c). Originally described as the most abundant sponge in Beaufort
Harbor North Carolina (Wilson 1911), it is also said to be very common in the Caribbean
(Diaz et al. 1993), and was recently described as the most common sponge in the
intertidal in the Bahía San Antonio, Argentina (Gastaldi et al. 2018). In the latter case,
however, the authors opted to refer to their samples by the name *H. perlevis,* as the
Argentinian samples were indistinguishable from ones in Northern Europe in genotype,
habitat, and morphology.
136
Here, I build on these results by 1) analyzing additional samples from Southern
California, which contains the type locality for *H. sinapium,* and 2) compiling all publicly

available genetic data (from 20 publications and several unpublished datasets). When
presented together, the data provide a compelling case for a single species ranging across
both the Atlantic and Pacific basins and the Northern and Southern hemispheres. Given
the limited dispersal capabilities of the species (Xue et al. 2009), the limited genetic
variation over most of its range (see below), and the historically documented range
expansion in California, these data are most consistent with an invasive spread via human
facilitation.
146
147 **Methods**
148
Collections
Sponges were located while SCUBA diving. Effort was made to photo-document all
sponge morphotypes present at each dive site, so that data on presence vs. absence could
be compiled. It should be noted, however, that search time was higher at some sites then
others, as shown in supplementary table 1. The search times listed are the total dive time,
cumulative across all dives at a site. This only approximates search effort, as some dives
were spent mainly searching and photographing sponges, while on others considerable
time was spent collecting samples. Collections were made by hand with a small knife.
Samples were placed individually in ziplock bags while underwater, accompanied with
copious seawater. These bags were put on ice until preservation, which was generally
within 2-5 hours, but sometimes up to 12 hours. Samples were moved directly from
seawater to 95% ethanol; in most cases, the preservative was changed with fresh 95%
ethanol after 1-3 days, and sometimes changed again if it remained cloudy. Most samples
were photographed underwater with an Olympus TG5 before collection and
photographed again in the lab. These photos accompany this paper as supplementary data
and are also posted as georeferenced records on the site iNaturalist.org. Two samples
were collected during the "LA Urban Ocean Bioblitz", and are present as vouchers at the
Natural History Museum of Los Angeles. Three other samples were vouchered with the
California Academy of Sciences in San Francisco. Voucher numbers are shown in
supplementary table 1. This table lists all samples known or suspected to be
Hymeniacidon sinapium: note that the standard of evidence is variable in each case, as
indicated in the table (e.g. some were photographed but not collected, and the ID is
therefore tentative; see results for further details).
172
Spicules
Sponge spicules were examined by digesting soft tissues in bleach. A subsample of the
sponge was chosen, taking care to include both the ectosome and choanosome. This was
placed in a 1.5 ml microcentrofuge tube with household bleach for several hours, until
tissue appeared to be dissolved. With the spicules settled at the bottom of the tube, the
bleach was then pipetted off and replaced with distilled water; this was repeated several
times (I found that 2-3 times was sufficient for visualizing spicules with a light
microscope, but removing all the salt from the sample for other downstream applications
required 5 or more rinses and worked best when the final ones were done with absolute
ethanol). In some cases, samples were centrifuged at low speed to reduce settling time
between rinses, though this increased the proportion of broken spicules.
184

Spicules were imaged using a compound triocular scope and pictures were taken using a
D3500 SLR camera (Nikon) with a NDPL-1 microscope adaptor (Amscope). Pictures of
a calibration slide were used to determine the number of pixels per mm, and 20-30
spicules were then measured using ImageJ (Schneider et al. 2012). Effort was made to
select sets of spicules without respect to their size, so as to get an unbiased estimate of
size distributions. All raw data are available as supplementary table 2. Spicule photos are
available in the supplementary data that accompanies this paper, and have been linked to
georeferenced records at iNaturalist.org.
193
I also imaged the spicular architecture in cleared tissue sections. I used a razor blade to
cut perpendicular sections that were as thin as possible by hand, and removed the
ectosome by hand for a surface view. These sections, already in 95% ethanol, were
soaked in 100% ethanol for a short time and then cleared for several hours in Histoclear
(National Diagnostics). These photos are available in the supplementary data that
accompanies this paper, and have been linked to georeferenced records at iNaturalist.org.
200
Genotyping
DNA was extracted from small subsamples, taking care to minimize contamination by
the many sponge-associated organisms that are often present. Some samples were
extracted with the Qiagen Blood & Tissue kit while the others were extracted with the
Qiagen Powersoil kit. The "barcoding" region of the cox1 gene was amplified using the
Folmer primers LCO1490 (GGTCAACAAATCATAAAGAYATYGG) and HCO2198
(TAAACTTCAGGGTGACCAAARAAYCA) (Folmer et al. 1994). A single attempt was
made to amplify a longer region using the primers from Rot et al. (Rot et al. 2006):
LCO1490 and COX1-R1 (TGTTGRGGGAAAAARGTTAAATT) but without success. A
portion of the 18S locus was amplified using the primers SP18aF
(CCTGCCAGTAGTCATATGCTT) and 600R18S (CGAGCTTTTTAACTGCAA)
(Redmond et al. 2013); the C2-D2 region of 28S was amplified using primers C2
(GAAAAGAACTTTGRARAGAGAGT) and D2 (TCCGTGTTTCAAGACGGG)
(Chombard et al. 1998). All primer sequences are listed 5' to 3'. PCR was performed in a
Biorad T100 thermocycler with the following conditions: 95C for 3 min, followed by 35
cycles of 94C for 30 sec, 52C for 30 sec, 72C for 1 min, followed by 72C for 5 minutes.
The C2-D2 28S region was amplified with a 57C annealing temperature instead of 52C.
PCR was performed in 50 ul reactions using the following recipe: 24 μl nuclease-free
water, 10 μl 5x PCR buffer (Gotaq flexi, Promega), 8 ul MgCl, 1 μl 10mM dNTPs
(Promega), 2.5 μl of each primer at 10 μM, 0.75 bovine serum albumin (10 mg/ml, final
conc 0.15 mg/ml), 0.25 μl Taq (Gotaq flexi, Promega), 1 ul template. ExoSAP-IT
(Applied Biosystems) was used to clean PCRs, which were then sequenced by Functional
Biosciences using Big Dye V3.1 on ABI 3730xl instruments. All PCR products were
sequenced in both directions, and a consensus sequence was constructed using Codon
Code v.9 (CodonCode Corporation). Blastn was used to verify that the resulting traces
were of sponge origin; all sequences have been deposited in Genabank as accessions
MT007958-MT007960 (cox1), MT001298 (18S), and MT006362 (28S). See
supplementary table 3 for additional details and information.
229
Genetic analysis

I retrieved all sequences with high sequence similarity to *H. perlevis* from Genbank. I
used the NCBI taxonomy browser to compile all data from samples identified as *H.*
perlevis, H. sinapium, H. heliophila, and *H. flavia.* Together, these data are from 20
different publications and several unpublished datasets (Erpenbeck et al. 2002, 2004,
2005, 2006, 2007; Park et al. 2007; Hoshino et al. 2008; Turque et al. 2008; Erwin et al.
2011; Alex et al. 2012, 2013; Morrow et al. 2013; Redmond et al. 2013; Thacker et al.
2013; Fuller and Hughey 2013; Jun et al. 2015; Miralles et al. 2016; Weigel and Erwin
2016; Gastaldi et al. 2018; Regueiras et al. 2019). I also retrieved all samples identified as
Hymeniacidon sp. and checked these and other sequences using blastn for similarity to *H.*
perlevis/sinapium/heliophila. Only four of these (JN093018 and KU697715-KU697717),
all identified as *Hymeniacidon sp.*, were closely related to the other samples, and all
appeared to be identical to others sequences within the *H. perlevis* clade. These four
unidentified samples were not included in downstream analyses.
244
I was not able to use all sequences in every analysis because of differences in the
sequenced portion of the gene or a lack of information regarding collecting location.
Importantly, no samples were excluded simply because they showed discordant patterns
of sequence variation. Supplementary table 3 lists every Genbank accession found,
indicates which were included in each analysis, and explains the reasons why any were
excluded. Some reads were included in the phylogenetic analysis, which could account
for unequal read lengths, but not the haplotype network, which included only samples
with complete data over the entire alignment. Sequence alignments were produced in
Codon Code v.9 (CodonCode Corporation). Haplotype networks were produced using the
minimum spanning method (Bandelt et al. 1999) as implemented in Popart (Leigh and
Bryant 2015). Phylogenies were estimated with maximum likelihood using IQ-Tree
(Nguyen et al. 2015; Trifinopoulos et al. 2016). I used a GTR model of sequence
evolution and used the Ultrafast bootstrap (Hoang et al. 2018) to measure node
confidence. Phylogenies were produced from the IQ-Tree files using the Interactive Tree
of Life webserver (Letunic and Bork 2019). Figures were made ready for publication
using R (r-project.org) and/or Gimp (gimp.org).
261
262   **Results**
263
Status in Southern California
Little data has been published about the distribution of *H. sinapium* in California outside
of bays and estuaries. Past surveys have focused on intertidal habitat and/or subtidal
sampling via deep water trawl (de Laubenfels 1932; Sim and Bakus 1986; Bakus and
Green 1987; Green and Bakus 1994). Is was therefore unknown if *Hymeniacidon*
sinapium is present in kelp forest ecosystems. I searched for it using SCUBA at 47 sites
in Southern and Central California (table S1). Subtidal sites were shallow rocky reefs
except for two locations which were oil platforms. Subtidal sites include four marine
protected areas along the mainland coast and three marine protected areas within the
Channel Islands National Marine Sanctuary. Six of the sites are also field sites within the
Santa Barbara Coastal Long-Term Ecological Research Network (sbclter.msi.ucsb.edu).
Though the survey was focused on kelp forest habitats, I also checked two intertidal sites
and floating docks in two harbors, as shown in table S1.

277
The distribution of *H. sinapium* in the Channel Islands region, where sampling was most
comprehensive, is shown in figure 1. I found the sponge at 8 of 19 mainland reefs in
Southern California, including both mainland marine protected areas investigated in
Southern California (Naples and Campus Point Marine Protected Areas). In only one
location (Carpinteria Reef) did I find *H. sinapium* growing on rock; in all other locations
it was largely buried in sediment, with projections extending into the water column. In
contrast to its prevalence on the Southern California mainland, I did not find it at any
island sites. This difference seems unlikely to be due to dispersal limitation because
island and mainland sites have high connectivity (Watson et al. 2010). It is more likely
due to the ecological differences between sites: none of the island sites investigated had
areas with the fine silty sediment where the sponge was most common on the mainland.
Though silty sites at the islands may simply have been unsampled in this survey, it is
likely they are less common than on the mainland. For example, satellite data show that
particles at the islands are less prone to resuspension by wave action (Freitas et al. 2017).
An intertidal survey of island sites in the 1970s did find *H. sinapium* at both San Miguel
and Santa Rosa Islands (Bakus and Green 1987). It has also been reported from the more
Southern islands of Catalina and San Clemente, which were barely sampled by my survey
(Sim and Bakus 1986; Bakus and Green 1987). To the North, in Central California, I only
surveyed three subtidal sites. I did not find *H. sinapium* in any of these Central California
sites, nor did I find it at the few intertidal sites, floating docks, or oil rigs that were
checked.
299
Together, my recent collections and the published intertidal and bay surveys in California
produce a portrait of a species that thrives in a wide variety of conditions, from bays to
the rocky intertidal to the kelp forest (Lee et al. 2007). It seems to dominate the intertidal
in some bay habitats with a muddy substrate and high sedimentation, and seems more
common in the kelp forest where fine sediment is found. These data are completely
consistent with the published descriptions of habitat preferences for *H. perlevis* in Europe
(Erpenbeck and Van Soest 2002) and *H. heliophila* in the Western Atlantic (Weigel and
Erwin 2016).
308
Gross Morphology of kelp forest samples
All but one of the newly collected samples were found embedded in sediment with
irregular projections extending into the water column. These projections varied from
stout cone-shaped or bag-shaped oscula to long, tendril-like digitations. One sponge was
found unburied, growing on rock. It lacked projections and instead resembled
Halichondria panicea (its identity was confirmed with spicule and DNA data, presented
below). All samples had a fleshy consistency, with the rock-dwelling sponge somewhat
more firm. Color varied from yellow to yellowish-orange in the field. Field photos are
available for 8 samples in the supplementary data accompanying this paper, and are also
available at iNaturalist.org.
319
I was interested in whether these sponges could be identified in the field and therefore
monitored using roving diver surveys or photo transects. These samples were collected as
part of an ongoing project to characterize the diversity of kelp forest sponges, with over

500 samples collected to date. This is the first survey of sponges in California via
SCUBA, and the first with extensive field photos of specimens that have also been
analyzed morphologically. Though the bulk of these data will be published elsewhere,
comparisons to date indicate that *H. sinapium* is the only sponge in these habitats that
grows by extending irregularly shaped projections out of silty sediment. Though this
morphology is certainly known from other species, *H. sinapium* was the only sponge with
this morphology found within the sampling effort shown in supplementary table 1. This
indicates that this morphology, when found in the Southern California kelp forest, is
strongly suggestive of the presence of this species. The most similar species found to date
is *Polymastia pachymastia*: as the name suggests, this sponge is covered in nipple-like
projections. This sponge was also found covered in sediment, with only the projections
visible. However, these projections tend to be uniform in shape and regularly spaced in *P.
pachymastia*, which contrasts with the irregularly spaced and morphologically various
projections seen in *H. sinapium*. The projections are also nearly white in *P. pachymastia*,
while they vary from yellow to nearly orange in *H. sinapium* (field photos of both species
have been deposited at iNaturalist.org). The rock-dwelling *H. sinapium* found at
Carpenteria Reef, however, would be more challenging to identify from field photos, as it
is very similar to other Halichondridae found in the survey.
341
Spicular morphology
I characterized the spicules of 9 samples to confirm their identity and compare them to
published data. All spicules were styles: tapered to a point at one end, and rounded at the
other. Width was usually uniform over the entire length, but a small minority had faint
swelling at or near the rounded end. This was manifest as a very weak swollen head
including the end (similar to the head of a match), or more commonly as a swollen band
near the head end (like a bead on a string). Most were somewhat curved or bent. The
skeleton of one sample was investigated further using hand-cut sections cleared with
Histoclear. Spicules in perpendicular sections through the choanosome formed wavy,
meandering tracts, the largest of which were about 30 μm wide. Spicules were also found
outside the tracts pointing in all directions (referred to as a "confused" arrangement in
sponge taxonomy). Surface sections revealed that the surface layer (ectosome) of the
sponge was filled with spicules that appeared to be tangential (parallel to the sponge
surface) and also "paratangential" (at an angle to the surface of less than 90 degrees).
These spicules were in messy bundles that formed an approximate mesh on the surface of
the sponge. Table 1 shows measurements of spicules as compared to values published in
other studies of *Hymeniacidon*. Newly collected data are consistent with published data
from *H. sinapium, H. perlevis,* and *H. heliophila,* as well as *H. fernandezi* (Thiele 1905)
from Chile (for which no genetic data is yet available). The arrangement of spicules in
cleared sections is also congruent with the spicular architecture described for *H. perlevis*
and other species (Erpenbeck and Van Soest 2002). Photos of tissue sections and spicules
are available as supplementary data.
364
Genetic analysis
I sequenced the newly collected samples at the cox1 locus (3 samples), the 18S rDNA (1
sample) and the 28S rDNA (1 sample). The *Hymeniacidon sinapium* holotype was also
loaned to me by the Smithsonian Natural History Museum: despite repeated attempts, I

was unable to amplify DNA from this sample. This is not surprising, as it was collected
in 1926 and little is known about its initial preservation.
371
I mined Genbank for all DNA data available for *H. perlevis, H. sinapium,* and *H.*
heliophila (see methods). I generated sequence alignments for four loci: cox1, 18S rDNA,
28S rDNA, and a locus spanning the first intergenic transcribed spacer, the 5.8S rRNA,
and the second intergenic transcribed spacer (hereafter referred to as the ITS). No other
locus had more than 2 sequences available in Genbank from any of these taxa.
Preliminary phylogenies indicated that sequences of *Hymeniacidon flavia* were more
closely related to the clade containing my target species than anything else in Genbank.
When available, these sequences were included for comparison.
380
Figure 2 shows the haplotype networks for the three loci with the most data. A large
dataset was available for 226 base pairs at the ITS locus. A set of 271 sponges from Japan
and Korea contained little genetic variation, as previously described (Park et al. 2007;
Hoshino et al. 2008). Samples from Northern, Central, and Southern California were all
identical to the most common Asian haplotype, as were 9 of 10 samples of *H. heliophila*
from the Eastern United States. These include samples from Alabama, Florida, and,
importantly, North Carolina. This last sequence read is the only one available that is
identified as coming from this state, which contains the type location for *H. heliophila*.
As a useful comparison to the diversity in 298 samples of *perlevis/heliophila/sinapium*, a
population sample of 212 *H. flavia* are shown. These are all from Japan, yet they contain
a similar amount of diversity as the worldwide sample from the other species.
392
A large mitochondrial dataset is also available at the Folmer barcoding region of the cox1
locus (571 bp; fig. 2). A single haplotype was shared among populations from China,
Korea, Southern California, Florida, and Portugal. Samples from Argentina contained
only 1-2 differences (99.6% identity) compared to this world-wide haplotype. The only
sample that is more than 0.5% divergent from this common haplotype, out of all *H.*
perlevis/sinapium/heliophila available, is a single sequence from Rio de Janeiro, Brazil
(1.2% divergent; top of fig. 2). No morphological description of this sample is available
in the related publication (Turque et al. 2008), but it states that vouchers were deposited
in the Museu Nacional, Universidade Federal do Rio de Janeiro. This sample is discussed
further below.
403
Less data is available for the 18S and 28S rDNA loci, but the 18S locus once again
illustrates the genetic similarity of Atlantic *H. perlevis/heliophila* populations and Pacific
H. sinapium populations (fig). Over the aligned 1,642 bp, samples from China shared an
identical haplotype with samples from Argentina. A sample of *H. perlevis* from Ireland,
the type locality for that species, differs by only a single base pair. Only a single data
point has any notable divergence: a sponge identified as *H. heliophila* from the USA.
This sample is separated from all others by 12 substitutions (0.7% divergence). I created
a phylogeny including selected other Halichondridae to place this divergence in context
(figure 3). While all other sequences of *H. heliophila/sinapium/perlevis* form a clade, this
USA sample is as divergent as other distinct species. The interior nodes of this phylogeny
are not well resolved, but it is clear that this sequence is an outlier and likely from a

different species. This sample (NCI217, Smithsonian voucher #0M9G1074-H) is part of a
collection of sponges for the National Cancer Institute deposited at the Smithsonian
Museum of Natural History (Redmond et al. 2013). It was collected by the Coral Reef
Research Foundation in the Florida Keys (Key Largo, from mud substrate), and identified
by Belinda Glasby (William Moser, Smithsonian Museum of Natural History, pers.
comm.). It is discussed further below.
421
The D3-D5 region of 28S also allowed for an interesting comparison (figure 3). The only
data available at this locus is from two European samples and two from the Florida Keys,
USA. One of the Florida sequences is from the same isolate as the outlier at 18S
(NCI217), while the other sample is from the same collection (NCI083, Smithsonian
voucher #0M9G1369-A) (Thacker et al. 2013). It was collected in the Florida Keys
(Marquesas Key, sand substrate; William Moser, pers. comm.) In agreement with the 18S
data, these samples do not appear to be from the same species as the European ones.
429
Discussion
Genetic data provide strong support for the synonymy of *H. perlevis* and *H. sinapium*.
The type locality for *H. sinapium* is Newport Bay, in Southern California (de Laubenfels
1932). Previously, the only genetic data from Southern California was from a single
sample from Mission Bay, roughly 140 km to the South of the type location. To this I
have added additional data from Santa Barbara County (200 km North of the type
location). All of these samples are identical to samples of *H. perlevis*. Indeed, there is no
appreciable genetic divergence between any sample from California, Japan, Korea or
Europe. I therefore formally propose that *H. sinapium* de Laubenfels (1930) be
considered a junior synonym of *H. perlevis* Montagu (1814).
440
It is likely that *H. heliophila* is also a junior synonym of *H. perlevis,* but some ambiguity
remains. Genetic data illustrate that the majority of samples identified as *H. heliophila*
are in fact *H. perlevis*, including the only one from North Carolina, which contains the
type location. The two National Cancer Institute vouchers from Florida, however, appear
to be from a different species. One cox1 sequence from Brazil is also modestly divergent,
and could be from another species. It is possible that these samples were merely
misidentified, as no public data exist on their spicular morphology, and I have not
examined the vouchers. Another possibility is that there are two morphologically similar
Hymeniacidon within the range ascribed to *H. heliophila*, mirroring the case of *H.*
sinapium and *H. flavia* in Japan. Though further work will be required to determine if the
name *H. heliophila* is valid, is it clear that the most common sponge matching its
description is in fact *H. perlevis*, whose range therefore includes North Carolina, Florida,
Alabama and Argentina at the very least.
454
Ecological and morphological data also support these far-flung populations being within
the same species. It should be noted, however, that there is another species that is
morphologically similar yet genetically distinct. The genetic outgroup to the *H.*
perlevis/heliophila/sinapium clade is *H. flavia*, known from Japan and Korea (Park et al.
2007; Hoshino et al. 2008). This species is sympatric with *H. sinapium* in Japan, and
cannot be distinguished from it based on spicular morphology. These species can only be

identified using genetic data, color when alive, or larval morphology (Sim and Lee 2003;
Hoshino et al. 2008). This illustrates that it may be difficult to resolve the taxonomy of
Hymeniacidon without genetic data. As pointed out by Gastaldi et al. (Gastaldi et al.
2018), there are additional species with morphological descriptions matching *H. perlevis*,
but the existence of *H. flavia* illustrates that confidently determining which are synonyms
will require DNA data.
467
Data on larval biology are available for *H. perlevis*, and they support a hypothesis of
recent range expansion via human facilitation. Larvae of *H. perlevis* are non-tufted
parenchymella, and do not appear to differ in their dispersal time compared to other
studied species (Xue et al. 2009). In the lab, all larvae stopped swimming and were
exploring the benthos by 19 hours after release, and all had settled by 43 hours. In
unfavorable conditions the larvae may travel farther: under high artificial illumination
(which increased mortality), sponge larvae swam for a maximum of 24 hours and some
were still exploring the benthos when the experiment was terminated at 68 hours. These
data are consistent with the larval ecology of other sponges (Maldonado 2006). It
therefore seems unlikely that the larvae of this species are exceptional compared to other
sponges, and could be naturally responsible for its exceptional range.
479
The data do not seem sufficient to form a strong hypothesis about the native range of this
species. It seems unlikely to be California, as almost no genetic variation was found at
the ITS or cox1 loci. Moreover, the species has likely undergone range expansion in
California between the 1920s and the present. Max de Laubenfels did extensive
collecting in California in the 1920s, naming many new species. He was based at
Stanford's marine station on Monterey Bay, but was only able to locate a single sample of
H. sinapium in the area. The species is now the "dominant" intertidal organism in
Elkhorn Slough, which drains into Monterey Bay; if this was true in the 1920s it seems
unlikely to have been missed by de Laubenfels. Europe is perhaps the most likely source,
as we know it was present there in the early 1800s. The genetic diversity at cox1 in
Portugal seems notably higher than the diversity at the ITS locus in Asia, though better
support would clearly come from comparing data from the same locus.
492
Future work will be needed to understand what impact this species has on the ecosystem
services provided by the communities in which it resides. Its exuberant abundance in
many habitats seems to make impacts likely, if for no other reason than the occupation of
space (Wasson et al. 2001). It is also notable that it has successfully colonized the kelp
forests in California, which have been relatively resistant to invasion (Steneck et al.
2002). Within these kelp forests, my observations suggest that this species can be
monitored, if imperfectly, using roving diver surveys or photo transects. Though some
sponges would require follow-up confirmation in the lab, this might allow the existing
large-scale monitoring efforts in California to include this species (Claisse et al. 2018;
Miller et al. 2018).
503
The work presented here builds on the excellent previous work of many authors, and
aspects of the pattern I describe have certainly been recognized by others. Hoshino et al.
(2008) and Gastaldi et al. (2018) both remarked upon the similarity of *Hymeniacidon* in

the *H. perlevis* clade, though in both cases they referred to this clade as a species complex
that might be synonymized in the future. Others have simply started referring to their
samples by the senior synonym *H. perlevis*, even if they are within the range ascribed to
one of the other taxa (Xue et al. 2009; Gastaldi et al. 2018). I build on these earlier efforts
by adding data from Southern California and, for the first time, presenting all the genetic
data from other projects in one analysis. I recommend synonymizing *H. sinapium* and *H.*
perlevis, and recognizing that *H. perlevis* is an invasive species with a global distribution.
My hope is that recognition of the unusual distribution and abundance of this species
motivates further work into its ecology and ecological impacts.
516

Acknowledgements
I am grateful for the help and support of many people in UCSB's Marine Science Institute
and Diving & Boating Program, especially Robert Miller, Clint Nelson, Christoph Pierre,
Frankie Puetzer, Christian Orsini, H. Mark Page, and Alecia Dezzani. Steve Lonhart
(NOAA) and Shannon Myers (UCSC) were instrumental in facilitating collections in
Central California, and the Natural History Museum of Los Angeles' DISCO program
facilitated collections in Los Angeles County.
524
Funding declaration
Financial support was provided by UCSB and by the National Aeronautics and Space
Administration Biodiversity and Ecological Forecasting Program (Grant
NNX14AR62A); the Bureau of Ocean Energy Management Environmental Studies
Program (BOEM Agreement MC15AC00006); the National Oceanic and Atmospheric
Administration in support of the Santa Barbara Channel Marine Biodiversity Observation
Network; and the U.S. National Science Foundation in support of the Santa Barbara
Coastal Long Term Ecological Research program under Awards OCE-9982105, OCE-
0620276, OCE-1232779, OCE-1831937. The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.
535

References

Alex A, Silva V, Vasconcelos V, Antunes A (2013) Evidence of unique and generalist
microbes in distantly related sympatric intertidal marine sponges (Porifera:
Demospongiae). *Plos One* 8(11): e80653

Alex A, Vasconcelos V, Tamagnini P, Santos A, Antunes A (2012) Unusual Symbiotic
Cyanobacteria Association in the Genetically Diverse Intertidal Marine Sponge
Hymeniacidon perlevis (Demospongiae, Halichondrida). *Plos One* 7(12): e51834

Bakus GJ, Green KD (1987) The distribution of marine sponges collected from the 1976-
1978 Bureau of Land Management Southern California Bight program. *Bulletin of*
the Southern California Academy of Sciences 86(2): 57–88

Bandelt H, Forster P, Röhl A (1999) Median-joining networks for inferring intraspecific
phylogenies. *Molecular Biology and Evolution* 16(1): 37–48

Bax N, Williamson A, Aguero M, Gonzalez E, Geeves W (2003) Marine invasive alien

species: a threat to global biodiversity. *Marine Policy* 27(4): 313–323

Bergquist PR (1961) A collection of Porifera from Northern New Zealand, with
descriptions of seventeen new species. *Pacific Science* 15(1): 33–48

Bergquist PR (1970) The marine fauna of New Zealand: Porifera, Demospongiae. Part 2
(Axinellida and Halichondrida). *Bulletin New Zealand Department of Scientific and*
Industrial Research [Memoir New Zealand Oceanographic Institute] 51: 9–85

Byrnes J, Stachowicz JJ (2009) Short and long term consequences of increases in exotic
species richness on water filtration by marine invertebrates. *Ecology Letters* 12:
830–841

Carballo JL, Aguilar-Camacho JM, Knapp IS, Bell JJ (2013) Wide distributional range of
marine sponges along the Pacific Ocean. *Marine Biology Research* 9(8): 768–775

Chombard C, Boury-Esnault N, Tillier S (1998) Reassessment of homology of
morphological characters in tetractinellid sponges based on molecular data.
Systematic Biology 47: 351–366

Claisse JT, Blanchette CA, Dugan JE, Williams JP, Freiwald J, II DJP, Schooler NK,
Hubbard DM, Davis K, Zahn LA, Williams CM, Caselle JE (2018) Biogeographic
patterns of communities across diverse marine ecosystems in southern California.
Marine Ecology 39(S1): e12453

Desqueyroux-Faúndez R, Soest RWM Van (1997) Shallow waters demosponges of the
Galapagos Islands. *Revue Suisse de Zoologie* 104: 379–467

Diaz MC, Pomponi S, Soest R van (1993) A systematic revision of the Central West
Atlantic Halichondrida (Demospongiae, Porifera). Part III. Description of valid
species. *Sciencia Marina* 57(4): 283–306

Erpenbeck D, Breeuwer JAJ, Parra-Velandia FJ, Soest RWM van (2006) Speculation
with spiculation?—Three independent gene fragments and biochemical characters
versus morphology in demosponge higher classification. *Molecular Phylogenetics*
and Evolution 38: 293–305

Erpenbeck D, Breeuwer JAJ, Soest RWM van (2005) Implications from a 28S rRNA
gene fragment for the phylogenetic relationships of halichondrid sponges (Porifera:
Demospongiae). *Journal of Zoological Systematics and Evolutionary Research*
43(2): 93–99

Erpenbeck D, Breeuwer JAJ, Velde HC van der, Soest RWM van (2002) Unravelling
host and symbiont phylogenies of halichondrid sponges (Demospongiae, Porifera)
using a mitochondrial marker. *Marine Biology* 141: 377–386

Erpenbeck D, Duran S, Rützler K, Paul VJ, Hooper JNA, Wörheide G (2007) Towards a
DNA taxonomy of Caribbean demosponges: a gene tree reconstructed from partial
mitochondrial CO1 gene sequences supports previous rDNA phylogenies and
provides a new perspective on the systematics of Demospongiae. *Journal of the*
Marine Biological Association of the United Kingdom 87(6): 1563–1570

Erpenbeck D, McCormack GP, Breeuwer JAJ, Soest RWM van (2004) Order level
differences in the structure of partial LSU across demosponges (Porifera): new
insights into an old taxon. *Molecular Phylogenetics and Evolution* 32: 388–395

Erpenbeck D, Soest RWM Van (2002) Family Halichondridae Gray, 1867. Systema
Porifera: A Guide to the Classification of Sponges. pp 787–815

Erwin PM, Olson JB, Thacker RW (2011) Phylogenetic diversity, host-specificity and
community profiling of sponge-associated bacteria in the northern Gulf of Mexico.
Plos One 6(11): e26806

Folmer O, Black MB, Wr H, Vrijenhoek RC (1994) DNA primers for amplification of
mitochondrial Cytochrome C oxidase subunit I from diverse metazoan invertebrates.
Molecular marine biology and biotechnology 3(5): 294–299

Freitas FH, Siegel DA, Maritorena S, Fields E (2017) Satellite assessment of particulate
matter and phytoplankton variations in the Santa Barbara Channel and its
surrounding waters: Role of surface waves. *JGR Oceans* 122(1): 355–371

Fuller TL, Hughey JR (2013) Molecular investigation of the invasive sponge
Hymeniacidon sinapium (de Laubenfels, 1930) in Elkhorn Slough, California.
Aquatic Invasions 8(1): 59–66

Gastaldi M, Paula TS De, Narvarte MA, Lôbo-Hajdu G, Hajdu E (2018) Marine sponges
(Porifera) from the Bahía San Antonio (North Patagonian Gulfs, Argentina), with
additions to the phylogeography of the widely distributed Cliona aff. celata and
Hymeniacidon perlevis, and the description of two new species. *Marine Biology*
Research 14(7): 682–716

Goeij JM de, Oevelen D van, Vermeij MJA, Osinga R, Middelburg JJ, Goeij AFPM de,
Admiraal W (2013) Surviving in a marine desert: The sponge loop retains resources
within coral reefs. *Science* 342(6154): 108–110

Green KJ, Bakus GJ (1994) Taxonomic atlas of the benthic marine fauna of the Western
Santa Maria Basin and the Western Santa Barbara Channel. Volume 2: The Porifera.
82 pp

Henkel D, Janussen D (2011) Redescription and new records of Celtodoryx
ciocalyptoides (Demospongiae: Poecilosclerida) A sponge invader in the north east

Atlantic Ocean of Asian origin? *Journal of the Marine Biological Association of the*
UK 91: 347–355

Hoang DT, Chernomor O, Haeseler A von, Minh BQ, Vinh LS (2018) UFBoot2:
Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution*
35: 518–522

Hofknecht G (1978) Descriptions and Key to the Intertidal Sponges of the Puerto
Peñasco Area in the Gulf of California. *Journal of the Arizona-Nevada Academy of*
Science 13(2): 51–56

Hoshino S, Saito DS, Fujita T (2008) Contrasting genetic structure of two Pacific
Hymeniacidon species. *Hydrobiologia* 603: 313–326

Jun J, Yu JN, Choi EH (2015) Complete mitochondrial genome of Hymeniacidon
sinapium (Demospongiae, Halichondriidae). *Mitochondrial DNA* 26(2): 261–262

Knowlton N (1993) Sibling species in the sea. *Annual Review of Ecology and Systematics*
24: 189–216

Laubenfels M de (1930) The Sponges of California. *Stanford University Bulletin* 5(98):
24–29

Laubenfels M de (1932) The marine and fresh-water sponges of California. *Proceedings*
of the US National Museum 81(4): 1–140

Lee WL, Elvin DW, Reiswig HM (2007) The Sponges of California: A Guide and Key to
the Marine Sponges of California. Monterey Bay Sanctuary Foundation, 130 pp

Leigh J, Bryant D (2015) PopART: Full-feature software for haplotype network
construction. *Methods in Ecology and Evolution* 6(9): 1110–1116

Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research* 47(W1): W256–W259

Maldonado M (2006) The ecology of the sponge larva. *Canadian Journal of Zoology* 84:
175–194

Maldonado M, Ribes M, Duyl FC van (2012) Nutrient fluxes through sponges: Biology,
budgets, and ecological implications. *Advances in Marine Biology* 62: 113–182

Miller RJ, Lafferty KD, Lamy T, Kui L, Rassweiler A, Reed DC (2018) Giant kelp,
Macrocystis pyrifera, increases faunal diversity through physical engineering.
Proceedings of the Royal Society B 285: 20172571

Miralles L, Ardura A, Arias A, Borrell YJ, Clusa L, Dopico E, Rojas AH de, Lopez B,
Muñoz-Colmenero M, Roca A, Valiente AG, Zaiko A, Garcia-Vazquez E (2016)

Barcodes of marine invertebrates from north Iberian ports: Native diversity and
resistance to biological invasions. *Marine Pollution Bulletin* 112(1-2): 183–188

Montagu G (1814) An Essay on Sponges, with Descriptions of all the Species that have
been discovered on the Coast of Great Britain. *Memoirs of the Wernerian Natural*
History Society 2(1): 67–122

Morrow C, Cárdenas P (2015) Proposal for a revised classification of the Demospongiae
(Porifera). *Frontiers in Zoology* 12(7): 1–27

Morrow CC, Redmond NE, Picton BE, Thacker RW, Collins AG, Maggs CA, Sigwart JD,
Allcock AL (2013) Molecular phylogenies support homoplasy of multiple
morphological characters used in the taxonomy of Heteroscleromorpha (Porifera:
Demospongiae). *Integrative and Comparative Biology* 53(3): 428–446

Nguyen LT, Schmidt HA, Haeseler A von, Minh BQ (2015) IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum likelihood phylogenies.
Molecular Biology and Evolution 32: 268–274

Park MH, Sim CJ, Baek J, Min GS (2007) Identification of genes suitable for DNA
barcoding of morphologically indistinguishable Korean Halichondriidae sponges.
Molecules and Cells 23(2): 220–227

Redmond NE, Morrow CC, Thacker RW, Diaz MC, Boury-Esnault N, Cárdenas P, Hajdu
E, Lobo-Hajdu G, Picton BE, Pomponi SA, Kayal E, Collins AG (2013) Phylogeny
and systematics of Demospongiae in light of new small-subunit ribosomal DNA
(18S) sequences. *Integrative and Comparative Biology* 53(3): 388–415

Regueiras A, Alex A, Costa MS, Pereira S, Vasconcelos V (2019) Diversity of intertidal
marine sponges from the western coast of Portugal (North-east Atlantic). *Journal of*
the Marine Biological Association of the United Kingdom 99(6): 1253–1265

Reiswig HM (1971) Particle feeding in natural populations of three marine Demosponges.
Biological Bulletin 141(3): 568–591

Rot C, Goldfarb I, Ilan M, Huchon D (2006) Putative cross-kingdom horizontal gene
transfer in sponge (Porifera) mitochondria. *BMC Evolutionary Biology* 6: 71

Ruiz GM, Fofonoff PW, Carlton JT, Wonham MJ, Hines AH (2000) Invasion of coastal
marine communities in North America: Apparent patterns, processes, and biases.
Annual Reviews of Ecology and Systematics 31: 481–531

Samaai T, Gibbons MJ (2005) Demospongiae taxonomy and biodiversity of the Benguela
region on the west coast of South Africa. *African Natural History* 1: 1–96

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of

image analysis. *Nature methods* 9(7): 671–675

Sim CJ (1985) A systematic study on the marine sponges from the South Sea and the
Yellow Sea of Korea. *Korean Journal of Systematic Zoology* 1: 21–30

Sim CJ, Bakus GJ (1986) Marine sponges of Santa Catalina Island, California. *Allan
Hancock Foundation Occasional Papers (New Series)* 5: 1–23

Sim CJ, Lee KJ (2003) A new species of the genus hymeniacidon (Demospongiae:
Halichondrida: Halichondriidae) from Korea. *Korean Journal of Biological Sciences*
7(3): 187–189

R.W.M. Van Soest, N. Boury-Esnault, J.N.A. Hooper, K. Rützler, N.J. de Voogd, B.
Alvarez, E. Hajdu, A.B. Pisera, R. Manconi, C. Schönberg, M. Klautau, M. Kelly, J.
Vacelet, M. Dohrmann, M.-C. Díaz, P. Cárdenas, J.L. Carballo, P. Ríos, R. Downey,
C.C. Morrow World Porifera Database. Hymeniacidon Perlevis (Montagu, 1814)
(2020a). World Porifera Database.

R.W.M. Van Soest, N. Boury-Esnault, J.N.A. Hooper, K. Rützler, N.J. de Voogd, B.
Alvarez, E. Hajdu, A.B. Pisera, R. Manconi, C. Schönberg, M. Klautau, M. Kelly, J.
Vacelet, M. Dohrmann, M.-C. Díaz, P. Cárdenas, J.L. Carballo, P. Ríos, R. Downey,
C.C. Morrow World Porifera Database. Hymeniacidon Luxurians (Lieberkühn,
1859) (2020b).

R.W.M. Van Soest, N. Boury-Esnault, J.N.A. Hooper, K. Rützler, N.J. de Voogd, B.
Alvarez, E. Hajdu, A.B. Pisera, R. Manconi, C. Schönberg, M. Klautau, M. Kelly, J.
Vacelet, M. Dohrmann, M.-C. Díaz, P. Cárdenas, J.L. Carballo, P. Ríos, R. Downey,
C.C. Morrow World Porifera Database. Hymeniacidon Heliophila (Wilson, 1911)
(2020c).

Steneck RS, Graham MH, Bourque BJ, Corbett D, Erlandson JM, Estes JA, Tegner MJ
(2002) Kelp forest ecosystems: biodiversity, stability, resilience and future.
Environmental Conservation 29(4): 436–459

Thacker RW, Hill AL, Hill MS, Redmond NE, Collins AG, Morrow CC, Spicer L,
Carmack CA, Zappe ME, Pohlmann D, Hall C, Diaz MC, Bangalore P V. (2013)
Nearly complete 28S rRNA gene sequences confirm new hypotheses of sponge
evolution. *Integrative and Comparative Biology* 53(3): 373–387

Thiele J (1905) Die Kiesel- und Hornschwämme der Sammlung Plate. 407-496 pp

Trifinopoulos J, Nguyen LT, Haeseler A von, Minh BQ (2016) W-IQ-TREE: a fast
online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research*
44(W1): W232–W235

Turque AS, Cardoso AM, Silveira CB, Vieira RP, Freitas FAD, Albano RM, Gonzalez
AM, Paranhos R, Muricy G, Martins OB (2008) Bacterial communities of the
marine sponges Hymeniacidon heliophila and Polymastia janeirensis and their
environment in Rio de Janeiro. *Marine Biology* 155: 135–146

Wasson K, Zabin CJ, Bedinger L, Diaz MC, Pearse JS (2001) Biological invasions of
estuaries without international shipping: the importance of intraregional transport.
Biological Conservation 102: 143–153

Watson JR, Mitarai S, Siegel DA, Caselle JE, Dong C, McWilliams JC (2010) Realized
and potential larval connectivity in the Southern California Bight. *Marine Ecology*
Progress Series 401: 31–48

Weigel BL, Erwin PM (2016) Intraspecific variation in microbial symbiont communities
of the sun sponge, Hymeniacidon heliophila, from intertidal and subtidal habitats.
Applied and Environmental Microbiology 82: 650–658

Wilson H V. (1911) Development of sponges from dissociated tissue cells. *Bulletin of the*
Bureau of Fisheries 30: 1–30

Xavier JR, Rachello-Dolmen PG, Parra-Velandia F, Schönberg CHL, Breeuwer JAJ,
Soest RWM van (2010) Molecular evidence of cryptic speciation in the
"cosmopolitan" excavating sponge Cliona celata (Porifera, Clionaidae). *Molecular*
Phylogenetics and Evolution 56: 13–20

Xue L, Zhang X, Zhang W (2009) Larval release and settlement of the marine sponge
Hymeniacidon perlevis (Porifera, Demospongiae) under controlled laboratory
conditions. *Aquaculture* 290: 132–139

745   **Table 1. Morphological data from newly collected samples with comparison data**
746   **from the literature.**
747

Sample	Date	Collection location	N	Style length	Style width
TLT550	1/7/20	Arroyo Quemado	25	151 - 262 - 336	3 - 5 - 7
TLT79	7/1/19	Tajigus	26	121 - 223 - 322	3 - 5 - 7
TLT87	7/1/19	Tajigus	21	147 - 216 - 326	2 - 5 - 8
TLT109	7/1/19	Tajigus	32	138 - 289 - 415	4 - 6 - 8
TLT339	8/30/19	Coal Oil Point	25	160 - 283 - 376	2 - 5 - 6
TLT247	7/31/19	Carpinteria	20	129 - 223 - 319	4 - 5 - 7
TLT129	7/31/19	Carpinteria	36	116 - 246 - 334	3 - 5 - 7
TLT15955	8/22/19	Redondo Barge	25	127 - 235 - 407	3 - 5 - 7
TLT349	8/23/19	Resort Wall	23	163 - 264 - 351	2 - 5 - 12
H. sinapium[1]	-	Elkhorn Slough, CA	-	115 - 460	3 - 12
H. perlevis[2]	-	Wales	-	152 - 475	3 - 12
H. perlevis[3]	-	South Africa	-	155 - 337	7
H. perlevis[4]	-	New Zealand	-	189 - 329	2 - 10
H. heliophila[5]	-	Caribbean	-	130 - 450	3 - 10
H. fernandezi[6]	-	Chile	-	200 - 340	3 - 10

748   Date = collection date, N = number of spicules measured, spicule dimensions given in
749   format min - mean - max; when only two numbers are shown they are min - max.
750   Sources: 1: Lee et al. 2007; 2: Erpenback & Van Soest 2002; 3: Samaai & Gibbons 2005;
751   4: Bergquist 1970; 6: Thiele 1905.
752

753 **Figure 1.** Collection locations in the Southern California Channel Islands region. Sites
754 where *H. sinapium* were found (black) and not found (white) are shown. The two sites
755 away from the coastline are oil platforms. Collection sites in Central California are not
756 shown.
757

758
759
760
761

762 **Figure 2.** Minimum-spanning haplotype networks for three loci. Samples are coded by
763 collection location, regardless of whether they were identified as *H. perlevis, H. sinapium,*
764 or *H. heliophila*. Closely related *H. flavia* are shown for comparison where available; all
765 data for this species is from Japan and Korea.
766

772 **Figure 3.** Phylogenies for the 18S and 28S loci. All samples identified as *H. perlevis, H.*
773 *sinapium* and *H. heliophila* are shown in bold with localities. Selected other
774 Halichondridae are shown for comparison, with *Suberites domuncula* specified as the
775 outgroup. Genbank accession numbers are also shown. Ultrafast Bootstrap support is
776 shown for all nodes with > 50% support.
777

778
779