Exposure to Mites Sensitizes Intestinal Stem Cell Maintenance, Splenic Marginal Zone B Cell Homeostasis, And Heart Development to Notch Dosage and Cooperativity.

Francis M. Kobia[1†*], Kristina Preusse[1†], Quanhui Dai[1,2], Nicholas Weaver[3], Sarah J. Stein[4], Warren S. Pear[4], Zhenyu Yuan[5], Rhett A. Kovall[5], Yi Kuang[1], Natanel Eafergen[6], David Sprinzak[6], Brian Gebelein[1], Eric Brunskill[1], Raphael Kopan[1,‡]

[1] Division of Developmental Biology, Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA.

[2] State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, Shanghai, China

[3] Immunology Graduate Program, University of Cincinnati College of Medicine, Cincinnati, OH, USA.

[4] Department of Pathology and Laboratory Medicine, Abramson Family Cancer Research Institute, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA 19104, USA

[5] Department of Molecular Genetics, Biochemistry and Microbiology, University of Cincinnati College of Medicine, Cincinnati, OH, USA

[6] School of Neurobiology, Biochemistry, and Biophysics, The George S. Wise Faculty of Life Sciences Tel Aviv University, Israel

† The authors made equal contribution

‡ Author for correspondence: Raphael.Kopan@CCHMC.ORG

* Present Address: Mount Kenya University, Department of Research and Innovation,

P. O. Box 342-01000, Thika - Kenya.

Key words: Notch, heart, intestine, stem cells, MZB, Lymphoma, SMZL, ectoparasite

Summary

Cooperative DNA binding is a key feature of transcriptional regulation. Here we examined the role of cooperativity in Notch signaling by CRISPR-mediated engineering of mice in which neither Notch1 nor Notch2 can homo- or heterodimerize, essential for cooperative binding to sequence paired sites (SPS) located near many Notch-regulated genes. While most known Notch-dependent phenotypes were unaffected in Notch1/2 dimer-deficient mice, a subset of tissues proved highly sensitive to loss of cooperativity. These phenotypes include heart development, compromising viability in combination with low gene dose, and the gut, developing ulcerative colitis in response to 1% DSS. The most striking phenotypes – gender imbalance and splenic marginal zone B cell lymphoma – emerged in combination with dose reduction or when challenged by chronic fur mite infestation. This study highlights the role of the environment in malignancy and colitis, and is consistent with Notch-dependent anti-parasite immune responses being compromised in the dimer deficient animals.

Highlights

- Notch dimerization has an *in vivo* role in contributing to intestinal homeostasis
- Loss of cooperativity can manifest as Notch gain or loss of function phenotypes
- Mite infestation exacerbates all phenotypes, triggers MZB hyperproliferation in mutant animals
- Mite-infested mutant mice develop SMZL with age

Introduction:

The evolutionarily conserved Notch receptors and ligands influence metazoan development and adult tissue homeostasis by directly translating an *inter*-cellular interaction into *intra*-cellular transcriptional outputs that control cell fate, proliferation, differentiation, and apoptosis (Artavanis-Tsakonas et al., 1999; Bray, 2006; Kopan and Ilagan, 2009). Mammals possess four Notch receptors (N1 to N4) and five Delta/Jagged ligands; all of which are Type I transmembrane proteins. The Notch pathway stands out relative to other signaling pathways in lacking signal amplification: canonical Notch signaling is initiated when a ligand on one cell engages a Notch receptor on a neighboring cell. This interaction unfolds the receptor's juxtamembrane region enabling cleavage by the metalloprotease ADAM10. The truncated, cell membrane bound polypeptide is then cleaved by the γ-secretase complex freeing the Notch Intracellular Domain (NICD), which subsequently translocates into the nucleus (Kopan and Ilagan, 2009; Kovall et al., 2017). NICD associates with the DNA-binding protein CSL (CBF1/Suppressor of Hairless/LAG-1, also known as RBPJ in vertebrates) and recruits the coactivator Mastermind; thereby assembling a Notch Transcription Complex (NTC) that activates Notch target gene expression (Bray, 2016; Gordon et al., 2008; Kovall et al., 2017).

The Notch pathway plays complex and context-dependent roles during development and adult tissue homeostasis. Perturbations in the Notch pathway are associated with developmental syndromes (Masek and Andersson, 2017) and cancers (Aster et al., 2017; Siebel and Lendahl, 2017). For example, *Notch1* promotes T-cell development (Pui et al., 1999; Radtke et al., 2010; Radtke et al., 1999), while *Notch2* is indispensable for Marginal Zone B-cell (MZB) development (Gibb et al., 2010; Hozumi et al., 2004; Liu et al., 2015; Maillard et al., 2004; Moran et al., 2007; Saito et al., 2003; Tanigaki et al., 2002; Witt et al., 2003). Accordingly, elevated Notch1 signaling is oncogenic in T-cells driving Acute Lymphoblastic Leukemia (T-ALL) (Ellisen et al., 1991; Weng et al., 2004) while increased Notch2 signaling is associated with splenic marginal zone B-cell transformation (Arcaini et al., 2016; Piris et al., 2017; Santos et al., 2017). Inversely, where Notch signals promote differentiation, the pathway can have a tumor suppressor function with diminished Notch signaling being associated with cancer (Demehri et al., 2009; Koch and Radtke, 2007).

How can the Notch pathway control multiple, dissimilar outcomes? Because each Notch receptor is consumed as it generates a signal and cannot be reused, signal strength has emerged as a key in controlling outcome. Indeed, some Notch-dependent processes are exquisitely sensitive to dosage and manifest both haploinsufficient and triplomutant effects in tissues such as the fly wing (Bray, 2006; Guruharsha et al., 2012). In mammalian tissues, signal

strength, defined as the sum of NICD released from all ligand-bound Notch receptors on the cell surface, is a far more important determinant of Notch signaling outcome than NICD composition (i.e. N1ICD vs N2ICD) (Liu et al., 2015; Liu et al., 2013; Ong et al., 2006). Several mechanisms have been found to modulate signal strength including receptor glycosylation (Haltiwanger, 2008; Kakuda and Haltiwanger, 2017), force-generating ligand endocytosis (Gordon et al., 2015; Langridge and Struhl, 2017), the contact interface between cells (Shaya et al., 2017), and the ratio of receptor/ligand constitution within the cell (Sprinzak et al., 2010). More recently, ligand-dependent signal dynamics were demonstrated to be another key determinant of signaling outcome (Nandagopal et al., 2018).

Interestingly, NICD can assemble into dimeric, cooperative NTCs (Arnett et al., 2010) on Sequence Paired Sites (SPSs), first described in the regulatory regions of the *Drosophila* E(spl) (enhancer of split) locus (Bailey and Posakony, 1995). SPSs consist of two DNA binding sites orientated in a head to head manner (Bailey and Posakony, 1995; Nellesen et al., 1999) separated by 15-17 nucleotides (Nam et al., 2007). NICD dimerization is facilitated via a conserved interface in NICD's ANK (Ankyrin repeat) domain. In human N1ICD, this interface consists of Arg^{1985}, Lys^{1946} and Glu^{1950}. Dimerization in Notch1 is effectively abolished by mutating Arg^{1985} into Ala^{1985} ($R^{1985}A$), Lys^{1946} into Glu^{1946} ($K^{1946}E$) or Glu^{1950} into Lys^{1950} ($E^{1950}K$), and loss of dimerization results in reduced activation of dimer-dependent targets (Arnett et al., 2010; Liu et al., 2010a; Nam et al., 2007).

Given the conservation of the dimer interface in most Notch receptors (*C. elegans* being the exception), the conservation of SPSs near known Notch targets (Hass et al., 2015; Liu and Posakony, 2012, 2014), and the ability of synthetic SPSs to regulate NICD levels (Kuang et al., 2019), we hypothesized that the precise mixture of agnostic and dimer-sensitive targets in a given cell will couple with Notch signal strength and dynamics to shape the responses to Notch signals and contribute to the context-specificity of Notch-related pathologies. To test this hypothesis in a mammalian species, we evaluated mice homozygous or hemizygous for dimerization-deficient alleles of *Notch1* (*N1^{RA}*) and *Notch2* (*N2^{RA}*) and found that dimerization contributes to context-specific Notch activity. We report that in the hemizygous state, dimer deficient Notch molecules are haploinsufficient in the heart and intestine, with lethal consequences when modified by the presence of the ectoparasite, *Demodex musculi* (fur mites). Mice homozygous for a dimer deficient *N1$^{RA/RA}$* allele displayed a female-biased, mild cardiac phenotype consistent with Notch loss-of-function. Conversely, *N2$^{RA/RA}$* mice displayed a striking over proliferation of marginal zone B cells in parasite-infested mice, but not in mite-free mice. In time, the cell-type specific gain of function activity of *N2$^{RA/RA}$* produced a splenic

marginal zone lymphoma-like phenotype in mite-exposed mice. Mechanistically, these effects reflect a shift in target amplitude favoring monomer-driven targets, and reduced negative feedback.

Results

Generation of N1 and N2 dimerization deficient mice

To interrogate the physiological role of dimerization, we first quantified the effective binding cooperativity of NTC to SPS and CSL sites using purified RBPj, the N1 RAM-ANK domain, and a MAML protein in electrophoretic mobility shift assays (EMSA; see supplemental information). We compared the mobility of NTC complexes containing either wild type N1ICD or a mutant version with the substitution of a single critical Arginine residue (Arg^{1974}) located in the mouse N1 ANK domain to alanine (Figure 1A-B, Supplemental Figure S1A-D). Quantifying the intensity levels of the bands in the image and fitting the data to a binding model that takes into account cooperative binding (see supplemental methods), we measured the cooperativity contributed by NTC binding to either an SPS probe (1xSPS) or a CSL probe containing two sites (2xCSL). We find that both wild type and mutant NTC had no cooperative binding to the 2xCSL probe. When the 1st site was already bound, the wild type NTC displayed a 5-fold stronger binding to the 2nd site in the 1xSPS probe compared to the binding to the 1st site. Importantly, NTC containing N1^{RA}ICD had no cooperative binding to the 2nd site in the 1xSPS probe (Fig. 1A, Supplemental Figure S1A-D).

To ask if loss of cooperativity impacts function *in vivo*, we used CRISPR-Cas9 to introduce amino acid substitutions at Arg^{1974} and Arg^{1934} in N1 and N2, respectively. Following the generation of a double strand break in Exon 32 of N1 or N2, a short oligonucleotide harboring two linked mutations was homologously recombined. Nucleotides coding for Arg^{1974} in N1 (N1R^{1974}) and Arg^{1934} in N2 (N2R^{1934}) were changed to code for Ala, creating the N1^{R1974A} (N1^{RA}) and N2^{R1934A} (N2^{RA}) mutations, respectively (Figure 1B, E). To facilitate genotyping of N^{RA} animals silent mutations were introduced to abolish a BglII restriction site while creating an XbaI restriction site in N1^{RA} (Figure 1B), or to generate a BglII restriction site in N2^{RA} (Figure 1E). Animals that carry one (N1$^{+/RA}$ or N2$^{+/RA}$) or two (N1$^{RA/RA}$ or N2$^{RA/RA}$) mutated chromosomes were first identified by digesting PCR products of the targeted chromosomal regions in Exon 32 with XbaI and/or BglII. XbaI cuts the N1^{RA} chromosome but not the wild type PCR product; BglII cuts the *N2^{RA}* but not the N2 wild type PCR products (Fig1 D, G). The presence of each respective RA mutation in *N1^{RA}* and *N2^{RA}* founders was subsequently confirmed by direct DNA sequencing

(Fig1 C, F). Animals deficient for N1 dimerization ($N1^{RA/RA}$), N2 dimerization ($N2^{RA/RA}$) or both N1 and N2 dimerization ($N1^{RA/RA}$; $N2^{RA/RA}$) were generated by crossing founders. All were viable, fertile and without any overt phenotype at P360 in mixed or C57BL6 background (unless otherwise stated, all subsequent results were generated in the mixed background). Because Notch1 signaling plays a central role in T-cell development (Radtke et al. 1999, 2010), we expected that loss of Notch1 dimerization would negatively impact the T-cell compartment. However, an analysis of the T-cell compartments in the thymus and spleen as well as thymic T-cell sub-compartments (DN, SP and DP), revealed a normal T-cell compartment in N1$^{RA/RA}$ mice relative to wild type controls (Supplemental Figure S2). This finding and the normal lifespan of $N1^{RA/RA}$; $N2^{RA/RA}$ animals, was surprising since we anticipated many dimerization-dependent genes (e.g., *Nrarp, Hes1, Hes5* and *Myc*; (Arnett et al., 2010; Hass et al., 2015) would be negatively impacted as seen when constitutively active wild type and dimer mutant Notch proteins lacking the extracellular domain (NΔE and N^{RA}ΔE) are overexpressed in cell culture (Supplemental Figure S1E).

Homologous $N1^{RA/RA}$; $N2^{RA/RA}$ mice display barrier defects in the colon and reduced proliferation in the crypt when challenged with 1% DSS

Notch1 and Notch2 act redundantly during the development and maintenance of the gut to block Klf4-induced niche exit of intestinal stem cells (Pellegrinet et al., 2011; Riccio et al., 2008), and Math1-induced differentiation of secretory cells (Noah and Shroyer, 2013; Shroyer et al., 2007). However, no pathology was revealed in histological examination of $N1^{RA/RA}$; $N2^{RA/RA}$ intestines (see below). Several studies have shown that a chronic decrease in Notch signaling compromised the intestinal barrier and exacerbated colitis in different mouse models (Mathern et al., 2014; Obata et al., 2012; Okamoto et al., 2009). Under the assumption that a compromised barrier may increase sensitivity to dextran sulfate sodium (DSS)-induced colitis, we asked if intestinal homeostasis and intestinal barrier were robust enough in $N1^{RA/RA}$; $N2^{RA/RA}$ by exposing mice to DSS. When given 2.5% DSS in their drinking water, wild type mice developed colitis as evident by a mild weight loss whereas co-housed $N1^{RA/RA}$; $N2^{RA/RA}$ mice exhibited severe weight loss necessitating euthanizing within one week (Figure 2A). To ask if $N1^{RA/RA}$; $N2^{RA/RA}$ were predisposed to develop colitis we exposed them to 1% DSS, which does not affect the weight of control littermates even after repeated exposures (Figure 2B, C). By contrast, co-housed $N1^{RA/RA}$; $N2^{RA/RA}$ mice experience significant weight loss, some had blood in stool at end of the third and fourth DSS cycle, and one had to be removed during the first cycle of DSS following severe weight loss and blood in stool. However, all remaining mice recovered

well when DSS was removed from the drinking water, even after multiple cycles of treatment (Figure 2B). Histological analysis of the colons revealed injury in wild type mice and a complete disruption of colonic epithelium of $N1^{RA/RA}$; $N2^{RA/RA}$ mice treated with 2.5% DSS (Figure 2D, top). After four periods of 1% DSS exposure $N1^{RA/RA}$; $N2^{RA/RA}$ mice had a more severe injury than controls or 2.5% DSS exposed wild type mice (Figure 2D). Relative to baseline (Figure 2E), proliferation in wild type crypt epithelia (Krt8/18 positive) was elevated in 1% or 2.5% DSS treated controls (Figure 2F). By contrast, we noticed a surprising decrease in proliferation competence (less Ki67) (and mitosis (Phospho-H3 positive cells)) in colonic epithelium following the fourth 1% DSS treatment in crypts of $N1^{RA/RA}$; $N2^{RA/RA}$ mice (Figure 2G) relative to wild type or untreated crypts, suggesting a role for Notch cooperativity within colonic stem cells recovering from injury. Finally, RA-specific while changes in barrier markers were not detected, a significant increase in IL4, Th17A and IL1ß was observed in DSS-treated $N1^{RA/RA}$; $N2^{RA/RA}$ relative to wild type (Figure 2H) consistent with enhanced inflammatory response driving colitis in these animals. Thus, these results are consistent with $N1^{RA/RA}$; $N2^{RA/RA}$ alleles being hypomorphic loss-of-function alleles in maintaining intestinal homeostasis and immune response.

$N1^{RA}$; $N2^{RA}$ hemizygotes display impaired intestinal cell proliferation and fate allocation in the crypt and cause ventricular septum defects in the heart.

The wild type and R-A mutant monomeric NICD protein bind CSL sites similarly (Figure 1A, Supplemental Figure S1), supporting the hypothesis that the signal generated in $N1^{RA/RA}$; $N2^{RA/RA}$ mice activated sufficient non-SPS gene targets to a level needed to support Notch-dependent decisions. If the Notch RA alleles are hypomorphic, challenging R-A mutant mice by lowering protein levels might reveal additional phenotypes. To test this idea, we generated an allelic series in which one copy of $N1$ or $N2$ was deleted in the $N1^{RA/RA}$; $N2^{+/+}$ or $N1^{+/+}$; $N2^{RA/RA}$ backgrounds, respectively, as well as deleting one copy of $N1$ and one copy of $N2$ deleted in the $N1^{RA/RA}$; $N2^{RA/RA}$ background. Single N1 or N2 RA hemizygotes resembled R-A mutants; however, crossing $N1^{RA/RA}$; $N2^{RA/RA}$ with $N1^{+/-}$; $N2^{+/-}$ animals generated significantly fewer $N1^{RA/-}$; $N2^{RA/-}$ pups at birth than expected ($p<3\times10^{-9}$, Chi-squared distribution analysis, Supp Table S1).

$N1^{+/-}$; $N2^{+/-}$ double hemizygous animals have a normal lifespan and no overt phenotype under normal housing conditions, but have been reported to develop mild cardiac phenotypes (Garg et al., 2005; Nigam and Srivastava, 2009). Given the critical role the circulatory system plays during gestation, we suspected that the poor representation of $N1^{RA/-}$; $N2^{RA/-}$ pups at birth was due to defects in cardiac development. Analysis of E16.5 embryos born to mite-free dams

revealed highly penetrant and severe Ventricular Septal Defects (VSD) in the hearts of $N1^{RA/-}$; $N2^{RA/-}$ embryos (Figure 3), compromising viability. Milder VSD was also observed with lower penetrance in N1$^{RA/-}$; N2$^{+/RA}$ embryos (Figure 3). Around this time, the colony became infected with fur mites, but the infestation was not immediately detected on the sentinels. Strikingly, when we retroactively segregated the fecundity data in our colony management software, we observed a significant (p=0.02, χ^2) skewing of the gender ratio in C57BL6 $N1^{RA/RA}$ carriers, with male:female ratio nearing 2:1 in fur mite infested mice, as opposed to the normal 1:1 ratio in mite-free mice (Supp Table S2).

The longest-lived $N1^{RA/-}$; $N2^{RA/-}$ pup born to mite infested, mixed background dams was much smaller than its litter mates and was sacrificed at P30 when it became moribund. Necropsy revealed a striking and complete loss of the intestinal crypts and villi (Carulli et al., 2015; Riccio et al., 2008; van Es et al., 2005; Vooijs et al., 2007). Since complete failure to form an intestine would have not permitted survival outside the womb, we assumed this phenotype reflected loss of intestinal stem cells (Figure 4A, A') consistent with the loss of Ki67 in DSS-challenged $N1^{RA/RA}$; $N2^{RA/RA}$ mice (Figure 2G). To investigate this phenotype further, we assessed proliferation and fate allocation in the developing intestine of E18-P1 animals with various allele combinations. To examine fate allocation, we stained sections with alcian blue and analyzed the number of secretory Goblet cells, which are known to expand in Notch hypomorphic background (Noah and Shroyer, 2013; Vooijs et al., 2007). The small intestines of control or $N1^{+/-}$; $N2^{+/-}$ P0 pups were indistinguishable (Figure 4B, C). By contrast, alcian blue staining revealed a significant expansion in the goblet cell compartment at P0 in $N1^{RA/-}$; $N2^{RA/-}$ pups born to fur mites infested dams (Figure 4D). In mice free of mites, the overall morphology appeared normal with few regions displaying excess goblets. To assess proliferation, we stained adjacent intestinal sections with antibodies against Ki67 and Phospho-H3. Strikingly, whereas trans-hemizygote $N1^{+/-}$; $N2^{+/-}$ intestines resembled wild type (Figure 4E-F), the $N1^{RA/-}$; $N2^{RA/-}$ intestine from mite-infested dames contained very few Ki67 positive cells (Figure 4G-H). We also noted a reduction in the number of cells undergoing DNA replication in particular, and competence to self-renew in general, in $N1^{+/-}$; $N2^{RA/RA}$ intestines (Supplemental Figure S3C-C', G,G'). However, upon eradication of *Demodex musculi* with permethrin, all E18.5 and all but 1/8 surviving P0 $N1^{+/-}$; $N2^{+/-}$ pups had at least some regions within the small intestine in which Ki67-positive tissue could be observed (Figure 4J, the percentage of Ki67+ crypts; Supplemental Figure S3I-J'; note reduced proliferation in Ki67-positive crypts of a P1 pup post-mite eradication). Overall, these data reflect an impaired ability to maintain intestinal stem cells in $N1^{RA/-}$; $N2^{RA/-}$ mice, which was exacerbated by *Demodex musculi* infestation. Because these

phenotypes can reflect complex interaction between multiple tissues in the $N1^{RA/RA}$; $N2^{RA/RA}$ and $N1^{RA/-}$; $N2^{RA/--}$ mice, we deferred the mechanistic analysis of this gene-environment interaction in ISC maintenance to future investigation.

The N2^{RA} allele drives expansion of MZ B-cells in *Demodex musculi* infested mice.

Marginal zone B-cells (MZB) are exquisitely sensitive to *N2* dosage (Gibb et al., 2010; Hozumi et al., 2004; Liu et al., 2015; Maillard et al., 2004; Saito et al., 2003; Tanigaki et al., 2002; Witt et al., 2003). Even a mild reduction in N2ICD can generate a noticeable decline in MZB cell numbers (Liu et al., 2015). We used FACS to analyze the splenic B cell population and asked if $N2^{RA}$ displayed haploinsufficient phenotypes in the dose-sensitive MZ compartment. Intriguingly, we observed a significant increase in the MZB compartment in $N2^{RA/RA}$ spleens in a mixed-background mouse exposed to *Demodex musculi* (Figure 5A). The size of other splenic B-cell subsets, including the immediate MZB precursors (MZP), Follicular B-cells (FoB) and transitional type-2 cells (T$_2$) did not change significantly in either $N2^{RA/RA}$ (Figure 5A), $N2^{+/-}$ or $N2^{RA/-}$ (Figure 5B), inconsistent with a fate switch.

MZB cells reside in the splenic marginal zone where they surveil for blood-borne pathogens. Upon encountering antigens, they rapidly differentiate into IgM producing plasmablasts that secrete vast amounts of IgM before undergoing apoptosis (Cerutti et al., 2013). To test IgM levels *in vivo*, we performed ELISA on serum collected from unstimulated $N2^{RA/RA}$ and littermate controls and detected no significant difference in the levels of circulating IgM (Figure 5C). To test for the possibility that loss of N2 dimerization negatively impacted differentiation into IgM producing plasmablasts upon stimulation, we FACS isolated MZB and FoB cells from mite-exposed $N2^{RA/RA}$ and wild type spleens. We then stimulated these cells *in vitro* with LPS for 5 days followed by imaging to assess for proliferation, and by ELISA to quantify the IgM levels secreted into the medium. Relative to wild type MZB cells, there was no appreciable difference in the ability of N2$^{RA/RA}$ MZB cells to respond to LPS stimulation (Supplemental Figure S4A), and ELISA analysis detected no significant difference in the amount of IgM secreted by $N2^{RA/RA}$ MZB cells relative to wild type controls (Figure 5C). Finally, we tested if loss of dimerization affected the development of B1 B cells, which predominantly reside in the peritoneal cavity (Witt et al., 2003). FACS analysis revealed no significant changes in the levels of either the B1a or B1b subsets of B1 B-cells between $N2^{RA/RA}$ and wild type animals (Figure 5D). Together, these data suggest that MZB cell numbers did not increase due to a fate switch or a failure to differentiate properly, and are consistent with a proliferative phenotype triggered by the N^{RA} mutation.

In the course of these experiments, the *Demodex musculi* infestation was eradicated with permethrin. Within a few months, as we saw with other phenotypes, mite-free $N2^{RA/RA}$ animals no longer showed MZB expansion (Figure 5E, post-permethrin, PP). Since permethrin has been reported to suppress the immune system in mice (Blaylock et al., 1995; Punareewattana et al., 2001), we assumed permethrin transiently suppressed MZB proliferation. Multiple subsequent analyses over many months, however, failed to detect a difference in MZB numbers between $N2^{RA/RA}$ and control mice post eradication and permethrin treatment, suggesting that this allele was indistinguishable from wild type in untreated and pathogen free animals. Accordingly, we compared MZB numbers between $N2^{+/-}$ spleens, which have decreased MZB numbers due to haploinsufficiency, with $N2^{RA/-}$ spleens and found no further reduction in MZB numbers (Liu et al., 2015; Saito et al., 2003). These data suggest that the $N2^{RA}$ has neither a haploinsufficient character for the $N2^{RA}$ allele in the MZB linage (Figure 5B), nor a proliferative phenotype triggered cell-autonomously by the N2^{RA} protein. To ask if environmental stimuli interacted with N2^{RA} within the spleen, mice were exposed to bacterial lipopolysaccharide (LPS) or to dermatitis, produced by twice weekly exposure of the ear epidermis to dust mite extract (HDM, *Dermatophagoides farinae*) for 6 weeks (see methods). MZB cell numbers were analyzed 8 weeks after the first LPS injection or HDM extract application. LPS induced transient IgM production but did not change MZB census in the spleen at 8 weeks (Figure 5E, LPS). Importantly, HDM-induced dermatitis elevated MZB numbers only in the $N2^{RA/RA}$ mice but not in controls (Figure 5E, HDM). Collectively, these experiments establish that specific gene-environment interactions produce the MZB phenotype in $N2^{RA/RA}$ mice.

To assess the rate of splenic B-cell proliferation in mite-infested mice, we stained C57BL/6J $N2^{RA/RA}$ and WT spleen sections with antibodies against Ki67, B220 (to mark the B-Cells) and Moma-1 (to mark the marginal zone). We detected defuse B220 staining with elevated Ki67 in marginal zone cells in the $N2^{RA/RA}$ spleen (Figure 5F), as well as in germinal centers (rich in FoB-cells and strongly B220 positive, Supplemental Figure S4), not seen in WT spleen of co-housed animals (Supplemental Figure S4B-D). Caspase 3 activation was not observed in $N2^{RA/RA}$ sections (not shown). Increased marginal zone proliferation in $N2^{RA/RA}$ relative to wild type control was also detected in spleens after six weeks of HDM treatment (Figure 5Fd, d'). Combined, these data suggest that mite infestation or prolonged HDM exposure leads to enhanced MZB proliferation.

Aged *N2^RA/RA* mice have enlarged spleens but only *Demodex musculi* infested mice progress to a Splenic Marginal Zone B-cell Lymphoma (SMZL)-like state

As noted above, we observed no change in lifespan in *N2^RA/RA* or *N1^RA/RA; N2^RA/RA* animals. However, as old animals were culled, we noticed enlarged spleens in animals carrying the *N2^RA/RA* allele post mite eradication (Supplemental Figure S5A). Spleens from mice infested by *Demodex musculi* were much larger, the most severe splenomegaly exhibited by *N2^RA/RA; N1^+/RA* (Figure 6A-E), perhaps because they contained a higher contribution of the inflammation-prone FVB strain (Quigley et al., 2011; Quigley et al., 2009). Histologically, spleens from mite-infested *N2^RA/RA* carriers lost their typical architectures with *N2^RA/RA; N1^+/RA* spleens showing the greatest disruption. Splenic morphology showed an expansion of the white pulp at the expense of the red pulp. In the most severe cases, the spleens appeared to consist of white pulp only (Figure 6F-K). In addition, two *N2^RA/RA; N1^+/RA* animals had multiple enlarged abdominal lymph nodes. H/E staining revealed a striking morphological similarity to the enlarged spleens from the same animals, suggesting infiltration by a splenic population, or lymphoproliferation within the LN (Figure 6L-L"). To further characterize the nodes, we stained the enlarged spleens and lymph nodes for the B-cell markers B220 and Pax5. Both were positive for the B-cell markers (Figure 6M-N'). These features are reminiscent of human SMZL (Splenic Marginal Zone B-cell - Lymphoma), a clonal B-cell neoplasm that manifests itself in elderly patients and consists of small lymphocytes that infiltrate the lymph nodes and other organs (Arcaini et al., 2016; Piris et al., 2017; Santos et al., 2017). As would be expected, the enlarged lymph nodes and spleens were highly proliferative as indicated by Ki67 and phospho-H3 staining (Supplemental Figure S5B,D). Paradoxically, human SMZL are associated with hyperactive NOTCH2 signaling due to an N2ICD-stabilizing truncation upstream to the WSSSP sequence in the PEST domain of NOTCH2 (Kiel et al., 2012; Peveling-Oberhag et al., 2015; Rossi et al., 2012), whereas the *N2^RA* alleles are hypomorphic in expression assays and in the context of the heart, crypt, and gut barrier. To confirm that this line of CRISPR-modified mice did not inherit a truncated *Notch2* allele, we sequenced Exon 34 DNA isolated form lymph node filled with B-cells and found a perfect match to the wild type *Notch2* allele (ENSMUSG00000027878).

Loss of Notch dimerization in *Demodex musculi* infested mice did not stabilize N2ICD in MZB cells but promoted their proliferation of MZB cells via the Myb-FoxM1 axis.

Recently, we identified an unanticipated consequence of increasing the number of SPS sites in the *Drosophila* genome, namely, accelerated degradation of phospho-NICD (Kuang et al., 2019). Further, we demonstrated that this enhanced degradation affects some, but not all,

Notch dependent decisions in *Drosophila.* Bristle precursor cells requiring a pulse of Notch were refractory to the destabilizing effect, whereas wing margin cells reliant on prolonged Notch signals were sensitive to NICD (Kuang et al., 2019). To test if the similarity of our phenotype to SMZL might in part be reflective of N2ICD stabilization, we assessed N2ICD levels in nuclear extracts from wild type and $N2^{RA/RA}$ MZB cells. We did not detect a significant accumulation of nuclear N2ICD in sorted $N2^{RA/RA}$ MZBs (Supplemental Figure S6), suggesting that the gain-of-function phenotypes observed in $N2^{RA/RA}$ MZB cells is not due to enhanced N2ICD stabilization. Thus, unlike in many human SMZL patients, increased N2ICD stability is unlikely to be the cause of the SMZL phenotype in the $N2^{RA/RA}$ mice.

In $N2^{RA/RA}$ mice, Notch pathway activity in T-cells and skin is intact as it relies on *Notch1*. Assuming the MZB effect is cell autonomous we addressed the mechanism involved in enabling a haploinsufficient allele to drive MZB expansion via transcriptome and chromatin analyses. We sorted and performed RNAseq on MZB cells from spleens of three $N2^{RA/RA}$ and three wild type littermates infested with *Demodex musculi*. The data were mapped to the mouse genome (mm10), and PCA analysis indicated that one of the triplicates in each group was an outlier. To stabilize the variance across replicates a two-way normalization strategy was used (see methods). Differentially expressed transcripts were identified using EdgeR. 1207 differentially expressed transcripts (>1 LogTPM, >1 log fold changes, 0.1 FDR, <0.05 pVal) were clustered into upregulated (270) and downregulated (937) $N2^{RA/RA}$ genes (Figure 7A). Down regulated genes included the validated Notch MZB targets *Hes1*, *Hes5*, and *Dtx1* and upregulated genes included *FoxM1*, *E2F1*, and *Myb* (Figure 7B). These three genes are known to promote cell cycle, DNA replication and DNA repair in B-Cells (Figure 7C, (Lefebvre et al., 2010; Lei et al., 2005)). Indeed, all the GO term for biological process for the upregulated genes were associated with the cell cycle (Figure 7B'; Table S3), with clear enrichment for E2F1 targets (Table S3). The downregulated genes did not enrich for any significant GO terms.

Given that Myb can act as an inducer of FoxM1 and E2F1 (Lefebvre et al., 2010), we confirmed its transcript levels were elevated by RT-qPCR (Figure 7D). *Fos* and *Pax5* were also found to be increased within $N2^{RA/RA}$ MZB. *Myb* could be activated by altered accessibility or by increased burst frequency and/or size, affecting amplitude. To assess accessibility, we performed ATACseq on MZB cells isolated from wild type and $N2^{RA/RA}$ mice (4 biological replicates each). We mapped reads under the peaks and used EdgeR to identify differential accessibility across the genome between samples (see methods for detail). 89059 peaks were mapped, of which 87575 were present in both $N2^{RA/RA}$ and wild type MZB cells with all eight samples being highly related (R>92%; Spearman Correlation; Figure 7G). Hence, genome

accessibility changed minimally between these MZB cells with only 984 peaks (1.1%) enriched in $N2^{RA/RA}$ and 500 (0.56%) in wild type. We next used GREAT to assign genes to all peaks. Based on these assignments, 14698 peaks mapped near differentially expressed (DE) genes, and of these only 108 were unique to wild type and 122 were unique to $N2^{RA/RA}$. Importantly, none of the differently enriched peaks were present at the *Myb*, *FoxM1*, *Dtx1*, *Hes1* or *Hes5* loci (Supplemental Figure S7). Next, we analyzed DNAse hypersensitive peak upstream of the *Myb* locus in the ENCODE datasets, and identified an enhancer accessible in several cell types, including the kidney. The genome of a kidney-derived cell line used to overexpress constitutively active NΔE and a dimer deficient version, NΔERA (Supplemental Figure S1) was examined by SplitDamID (Hass et al., 2015). We detected strong binding to the *Myb* enhancer using Notch/RBP complementing pairs, but not by Notch dimers. This data suggested that the *Myb* enhancer responds to monomeric NTC but not to dimeric NTC. Collectively, these data suggest that the changes in MZB gene expression in the $N2^{RA/RA}$ animals are not due to dramatic changes in chromatin accessibility but are rather due to changes in gene expression caused by the loss of Notch dimer NTC complexes at some other loci, most likely the SPS-dependent negative regulators *Hes1*, *Hes5,* and *Nrarp* (Hass et al., 2015; Severson et al., 2017).

.

Discussion

The possibility that DNA binding site architecture contributes to Notch signaling outcomes has been considered in past studies (Liu and Posakony, 2012, 2014). NTCs can bind to enhancers with CSL binding sites as monomers that function independently of one another. In such a model, the probability of target activation will depend upon the relative abundance of nuclear NICD and the number of CSL sites at a given enhancer (Gomez-Lamarca et al., 2018; Kueh et al., 2016; Schroeter et al., 1998) and/or burst size (Falo-Sanjuan et al., 2019; Lee et al., 2019). SPSs are found in enhancer regions of up to 30% of mammalian Notch targets including *Nrarp/NRARP*, *Hes1/HES1*, *Hes5*, and *Myc* (Arnett et al. 2010, Cave et al. 2005, Liu et al. 2010, Ong et al. 2006, Severson et al. 2017, Yashiro-Ohtani et al. 2014). We have previously documented that ~2500 SPSs were bound by N1ICD dimers in the mouse genome, and that dimerization-deficient Notch molecules bind poorly to SPSs *in vivo*, even at high NICD concentrations (Hass et al., 2015). In the murine kidney cell line, mK4, ≃15% of Notch targets required NICD dimerization. SPSs and cooperativity also proved critical for the oncogenic activity of N1ICD in murine T-cells, where NICD dimerization at a distant enhancer was required for *Myc* activation (Liu et al., 2010b). Thus, Notch targets are either agnostic to dimerization or sensitive to its presence, leading to the hypothesis that cooperativity may modulate Notch responses at physiological NICD concentrations. Recently, evidence that SPS contributed to increased probability of activation and increased duration during *Drosophila* embryonic development were reported (Falo-Sanjuan et al., 2019).

While our study concludes that embryonic development in the mouse proceeds normally in the presence of Notch receptors lacking cooperativity, it nonetheless uncovered important contributions of cooperativity to mammalian development and homeostasis under environmental stress. Notch^{RA} phenotypes are greatly enhanced by reduced gene dosage in Notch1 (causing ventricular septal defects (Donovan et al. 2002, MacGrogan et al. 2010, Sakata et al. 2002)) or in both Notch1 and Notch2 (gut). The developmental phenotype uncovered when N1^{RA} dose is reduced (in *N1$^{RA/-}$*) resembles other Notch pathway deficiencies known to cause a wide range of heart defects including ventricular septal defects (VSD) (Donovan et al. 2002, MacGrogan et al. 2010, Sakata et al. 2002), and are milder than a weak hypomorphic Notch1/2 allele on the B6 background (Liu et al. 2015). Interestingly, dosage alone did not impact the ability of the *N2^{RA}* allele to control development and homeostasis of the dosage-sensitive marginal zone B cells in parasite-free mice, even in the hemizygote state (*N2$^{RA/-}$*).

Strikingly, loss of cooperativity-dependent contributions to homeostasis becomes acute when mutant animals are burdened with exoparasites, namely fur mites. Interestingly, when the

environment includes unmanaged exoparasite infestation, dimerization-deficient cooperativity mutant receptors can behave as hypomorphic (loss-of-function) alleles in some contexts (intestinal barrier formation, ventricular septum formation, intestinal stem cell self-renewal, female survival), remain indistinguishable from the wild type allele in many other tissues, or can drive the same disease as a gain-of-function allele (splenic marginal zone lymphoma, SMZL) in a cell type that otherwise seemed to be unaffected by the mutation. Note that in $N2^{RA/RA}$ mice, Notch pathway activity in T-cells and skin, where *Notch1* is present, are identical to the wild type. This provided the rationale for the molecular analysis – the defect is more likely to reside in the B-cell linage than elsewhere, though this will remain to be examined more fully in the future. We propose these behaviors can be explained by assuming that the balance between targets agnostic to dimerization (*Myb*) and those dependent on cooperativity on SPS sites (*Hes1*, *Hes5*, *Nrarp*) varies by cell type (Figure 8). Moreover, these differences in dependence upon SPS vs non-SPS target gene regulation between tissues can be further exposed by additional stressors such as environmental insults (i.e. exoparasites) or changes in gene dose. We have proposed before that accessible SPS can act as a "sink", holding on to NICD that otherwise will be available to regulate monomer-dependent enhancers. If true, more N2^{RA}ICD may available to regulate *Myb*. Importantly, this insight may translate to other transcription factors where variants of uncertain significance are associated with developmental syndromes or neoplastic disease. Such mutations may control molecular behaviors integrating the environment with specific cooperative interactions in the affected tissues.

The diverse cell types responding to a cutaneous parasite must reflect a systemic change driven by the parasite. Since we could replicate the impact on MZB proliferation in short exposures to dust mite extract, live parasite is not needed and the response is likely mediated by an immune cell (or cytokine) intermediate. Dermatitis is typically associated with both skin-produced cytokines (e.g., IL-33, -17C and TSLP), and a skewing of CD4+ T-cells into T Helper 2 (Th2) and their associated cytokines (e.g., IL-4,-5,-13, (Huang et al., 2003)). Similarly, fur mite infestation has been shown to increase the level of different cytokines (e.g. IL-6 and IL-12; (Johnston et al., 2009)), including Il-4 and -5 (Moats et al., 2016; Pochanke et al., 2006). While the exact axis linking anti-parasite immune responses and splenic marginal zone lymphoma, intestinal homeostasis, female/male ratios and heart development are yet to be identified, this study highlights the role of the environment in chronic disease, including ulcerative colitis and a smoldering malignancy such as SMZL. Combined with the significantly better outcomes in SMZL patients with activated *Notch2* alleles (Rossi et al., 2012), one has to wonder about the

role of the immune environment in onset and progression of human SMZL patients with *Notch2* mutations.

Acknowledgments

The authors wish to thank Dr. Matt Kofron with help with imaging, Dr. Yueh-Chiang Hu and the genome editing core for generating the NRA alleles, and members of the Kopan lab for comments and encouragement during the study. Dr. Stephan Waggoner, Kelli VanDussen and members of the DB and Notch community for valuable advice. Many thanks to Dr Sai Tammula and the veterinarians at CCHMC. Cell sorting was supported by NIH S10OD023410 and 01DK106225 to the CCHMC Research Flow Cytometry Core. BG and DS were supported at a grant from the National Science Foundation and the Binational Science Foundation (NSF/BSF - 1715822). SJS was supported by NIH T32CA009140 and the American Cancer Society PF-15-065-01-TBG. WSP was supported by NIH R01 CA215518 (WSP). RK, FMK, KP, NW, and EB were supported by MIH RO1 CA163353 (RK). RK and QD were supported by the William K. Schubert Endowed Chair at Cincinnati Children's Hospital Medical Center. QD receives a small stipend from the Chinese government.

Authors contributions

FMK generated most of the data in mite infested mice and contributed to writing of the manuscript; KP contributed most of the data in the mite-free colony and supervised QD during the DSS experiments. NW contributed the initial establishment of the MZB phenotype. RAK/ZY purified the proteins, YK, NE, DS and BG analyzed cooperativity, with YK/BG performing EMSA experiments and NE/DS calculating the cooperativity factor. SJS and WSP analyzed $N1^{RA}$ T-cells. EB performed the RNA and chromatin analyses and supervised NW. BA performed the RNAseq clustering and enrichment analysis. EB/RK conceptualized the study and analyzed the data with input from KP, FMK, and QD. RK wrote the manuscript. All authors read and edited the final version.

Figure Legends:

Figure 1. Generation of Notch dimerization deficient mice. (A) Electrophoresis mobility shift assay for purified proteins binding to CSL (magenta) or SPS (green) probes. Balls mark occupancy of one or two sites. Note that cooperative binding by wild type NTC, but not the RA mutant NTC, specifically depletes the SPS probe, but not the CSL probe (compare lanes 8 and 13). See text and Supplemental Figure S1 for detail. (B-C) CRISPR-Cas9-mediated double-strand break was used to mediate homologous recombination of a short oligonucleotide into Exon 32 substituting Arg $N1^{R1974}$ (B-B") and $N2^{R1934}$ into Ala (C-C"), generating $N1^{RA/RA}$ and $N2^{RA/RA}$ animals. To facilitate genotyping, two silent mutations (in red) were included in the oligo, abolishing the BglII restriction site while generating an XbaI site in $Notch1$ (B'). In Notch2, a silent mutation (in green) was included to create a BglII site (C'). Sequencing PCR products containing these regions confirmed the presence of the Arg to Ala substitution in founders; digestion of these PCR products with XbaI (for N1, B") and BglII (for N2, C") confirmed the presence of one (N1) or two (N2) mutant alleles.

Figure 2. Notch dimerization deficient mice are sensitized to DSS-induced colitis. (A) All $N1^{RA/RA}$; $N2^{RA/RA}$ mice exposed to 2.5% DSS treatment had to be euthanized due to severe weight loss before day seven. (B) Daily weight measurements of wild type (blue) or mutant (red) mice treated with alternate cycles of 1%DSS (gray sections) or no DSS (white sections). (C) Survival curve of 1% DSS treated mice. Three of four mutant mice had to be euthanized due to sever weight loss by the fourth cycle. (D) Hematoxylin-eosin staining of colonic tissue from DSS treated mice. Dashed black boxes are enlarged below. Yellow dashed box showed a section with crypts in an otherwise injured colon in 1% DSS treated $N1^{RA/RA}$; $N2^{RA/RA}$ mice. (E) Ki67 staining of untreated mice with the indicated genotypes, crypt regions (Ki67+) are boxed. (F) Increased proliferation competence in colon of DSS treated wild type mice (Krt8/18+, Ki67+ staining outside the box). (G) Decreased proliferation competence in colon of $N1^{RA/RA}$; $N2^{RA/RA}$ mice exposed to four rounds of 1% DSS. (H) qPCR on RNA extracted from distal colon of $N1^{RA/RA}$; $N2^{RA/RA}$ and WT mice treated for 11 days with 1% DSS; n=3. (*, $p<0.05$).

Figure 3. Notch1^{R1974A} substitution compromised ventricular septum development. (A) Normal heart with complete septum. (B) $N1^{RA/-}$; $N2^{RA/-}$ heart with a severe, highly penetrant VSD. The penetrance by age shown below the image. (C) $N1^{RA/-}$; $N2^{+/RA}$ heart with a milder, less penetrant VSD.

Figure 4. Notch dimerization deficient mice are hypomorphic for Notch activity in the gut. (A-C) Alcian blue staining of $N1^{RA/RA}$; $N2^{RA/RA}$ and $N1^{+/-}$; $N2^{+/-}$ adult intestines. (D) Deletion of one N1 and one N2 allele in the RA background ($N1^{RA/-}$; $N2^{RA/-}$) caused lethality; loss of the intestine (presumably due to stem cell exhaustion) was evident in the longest surviving pup at P30. (E-G) Alcian blue analysis of P0 intestines detected an increase in goblet cell numbers in $N1^{RA/-}$; $N2^{RA/-}$ (G) relative to wild type or $N1^{+/-}$; $N2^{+/-}$ intestines (E-F), quantified in (H). (I-J) Ki67 staining in P0 intestine from control $N1^{+/-}$; $N2^{+/-}$ (I) or $N1^{RA/-}$; $N2^{RA/-}$ (J) newborn. (K) quantification of Ki67-positive area at the indicated age. Among surviving $N1^{RA/-}$; $N2^{RA/-}$ pups during (n=2) and after (n=12) fur mite infestation.

Figure 5. Loss of Notch2 dimerization expands the splenic MZ B-cell compartment in the presence of mites. (A) FACS analysis of $N2^{RA/RA}$ fur mite infested mouse spleens revealed an expansion the MZ B-cell compartment relative to wild type (+/+). Values for MZ P, F0, and T2 splenic B-cell populations were not significantly different; n=11. (B) The MZB-cell compartment in heterozygotes. (C) LPS-induced differentiation and IgM production *in vitro* and serum levels of IgM *in vivo;* n=4. (D). Peritoneal B1a B-cell compartment in mite infested wild type and $N2^{RA/RA}$ mice. (ns=not significant, n=2). (E). The MZB-cell compartment in $N2^{RA/RA}$ mice after treatment with permethrin, an immunosuppressant used to manage fur mite infestation (PP), after LPS, or after 6 weeks of treatment with house dust mites (HDM) extract. (*, $p<0.05$; ns=not significant). In (A-E) P-values generated by two-tailed t-test. Error bars represent s.d. (F). Proliferation in the splenic marginal zone of mite infested mice: (Fa) The spleen from a mite infested control animal or (Fb) from $N2^{RA/RA}$ mice. Boxed region magnified in Fc and again, in Fc', asterisk and dot provided for orientation. Magnification noted. (Fd, d') 60x view of marginal zone from control and HDM treated $N2^{RA/RA}$ mice.

Figure 6. Aged $N2^{RA/RA}$ and $N2^{RA/RA; N1+/RA}$ mice develop severe splenomegaly and tumors reminiscent of Splenic Marginal Zone B-cell Lymphoma (SMZL). (A-E) Images of spleens from mice P600-P700 with indicated genotypes. (F-K) Severely enlarged spleens are associated with disorganized morphology and the appearance of cells with large nuclear:cytoplasmic ratio (genotypes indicated, see low magnification image in Supplemental Figure S6E). (L) large enteric lymph nodes detected in the individual shown in (J), densely populated with cells with large nuclear:cytoplasmic ratio (L'-L"). Staining the spleen (M) and lymph node (N) with Ki67 and the B-cell markers B220 and Pax5 identifies infiltrating cells as spleen-derived B cells. Magnification: 10x.

Figure 7. Mite infested *N2^{RA/RA}* activate a proliferation module in MZB. (A) RNAseq analysis identifies enrichment for genes trending down or up in *N2^{RA/RA}* MZB realative to wild type controls. (B) Down trending genes include classical dimer-dependent Notch targets in MZB, *Hes1* and *Hes5,* and *Dtx*. Up regulated genes include *FoxM1, Myb*, and *E2F1*. (B') clusters 6 and 7 are highly enriched for GO terms associated with replication. (C) A schematic of the replication module controlled by *Myb*. (D) QPCR validation for *Fos, Myb*, and *Pax5.* (F) The *Myb* locus analyzed for chromatin accessibility (Encode DHS and ATAC-seq data generated in this study), transcription (RNA), and DNA methylation by DAM methyl transferase complementation (SplitDAM). Control D/AM haves methylation patterns are compared to methylation by Notch/RBPj pairs (both dimers and monomers) and Notch/Notch pairs (only dimers). See text and (Hass et al., 2015) for additional detail. (G) Spearman correlation table measures ATACseq peaks similarity matrix between all replicates. All are >92% similar.

Figure 8. A schematic summary of the findings with a hypothetical mechanism. *Notch* integrates environmental cues (mites, cytokines) to drive proliferation in MZB (*Notch2)* and block differentiation in intestinal stem cells (ISC. *Notch*1 & 2). Activation of SPS-dependent *Hes* repressors creates a negative feedback tuning of transcription amplitude in MZB, and driving the response magnitude in ISC. In *Notch^{RA/RA}* mutants, SPS-dependent *Hes* gene expression is dampened, and increased availability of NICD^{RA} to monomer driven target due to reduce trapping on accessible SPS may also contribute. Repeated DSS treatment, or hemizygosity in *Notch^{RA/RA}* mutants, exposed insufficient blockade of pro-differentiation signals in ISC, most likely delivered via immune cells or cytokines (IL1ß, 4, or 17A) in DSS treated mice. Fur mite infestation creates an environment for runaway proliferation in MZB, and enhanced dosage effects in *Notch^{RA/RA}* mutants leading to complete loss of ISC postnatally even without DSS challenge.

References

Arcaini, L., Rossi, D., and Paulli, M. (2016). Splenic marginal zone lymphoma: from genetics to management. Blood *127*, 2072-2081.

Arnett, K.L., Hass, M., McArthur, D.G., Ilagan, M.X., Aster, J.C., Kopan, R., and Blacklow, S.C. (2010). Structural and mechanistic insights into cooperative assembly of dimeric Notch transcription complexes. Nat Struct Mol Biol *17*, 1312-1317.

Artavanis-Tsakonas, S., Rand, M.D., and Lake, R.J. (1999). Notch signaling: cell fate control and signal integration in development. Science *284*, 770-776.

Aster, J.C., Pear, W.S., and Blacklow, S.C. (2017). The Varied Roles of Notch in Cancer. Annu Rev Pathol *12*, 245-275.

Bailey, A.M., and Posakony, J.W. (1995). Suppressor of Hairless directly activates transcription of Enhancer of split complex genes in response to Notch receptor activity. Genes and Development *9*, 2609-2622.

Blaylock, B.L., Abdel-Nasser, M., McCarty, S.M., Knesel, J.A., Tolson, K.M., Ferguson, P.W., and Mehendale, H.M. (1995). Suppression of cellular immune responses in BALB/c mice following oral exposure to permethrin. Bull Environ Contam Toxicol *54*, 768-774.

Bray, S.J. (2006). Notch signalling: a simple pathway becomes complex. Nat Rev Mol Cell Biol *7*, 678-689.

Bray, S.J. (2016). Notch signalling in context. Nat Rev Mol Cell Biol *17*, 722-735.

Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. Current protocols in molecular biology / edited by Frederick M Ausubel [et al] *109*, 21 29 21-21 29 29.

Carulli, A.J., Keeley, T.M., Demitrack, E.S., Chung, J., Maillard, I., and Samuelson, L.C. (2015). Notch receptor regulation of intestinal stem cell homeostasis and crypt regeneration. Dev Biol *402*, 98-108.

Cerutti, A., Cols, M., and Puga, I. (2013). Marginal zone B cells: virtues of innate-like antibody-producing lymphocytes. Nat Rev Immunol *13*, 118-132.

Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A., Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., *et al.* (2017). An improved ATAC-seq

protocol reduces background and enables interrogation of frozen tissues. Nat Methods *14*, 959-962.

Demehri, S., Turkoz, A., and Kopan, R. (2009). Epidermal Notch1 Loss Promotes Skin Tumorigenesis by Impacting the Stromal Microenvironment. Cancer Cell *16*, 55-66.

Ellisen, L.W., Bird, J., West, D.C., Soreng, A.L., Reynolds, T.C., Smith, S.D., and Sklar, J. (1991). TAN-1, the human homolog of the Drosophila notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms. Cell *66*, 649-661.

Falo-Sanjuan, J., Lammers, N.C., Garcia, H.G., and Bray, S.J. (2019). Enhancer Priming Enables Fast and Sustained Transcriptional Responses to Notch Signaling. Developmental Cell *50*, 1-15.

Garg, V., Muth, A.N., Ransom, J.F., Schluterman, M.K., Barnes, R., King, I.N., Grossfeld, P.D., and Srivastava, D. (2005). Mutations in NOTCH1 cause aortic valve disease. Nature *21*, 180-184.

Gibb, D.R., El Shikh, M., Kang, D.J., Rowe, W.J., El Sayed, R., Cichy, J., Yagita, H., Tew, J.G., Dempsey, P.J., Crawford, H.C.*, et al.* (2010). ADAM10 is essential for Notch2-dependent marginal zone B cell development and CD23 cleavage in vivo. J Exp Med *207*, 623-635.

Gomez-Lamarca, M., Falo-Sanjuan, J., Stojnic, R., Abdul Rehman, S., Muresan, L., Jones, M.L., Pillidge, Z., Cerda-Moya, G., Yuan, Z., Baloul, S.*, et al.* (2018). Activation of the Notch Signaling Pathway In Vivo Elicits Changes in CSL Nuclear Dynamics. Dev Cell *44*, 611-623 e617.

Gordon, W.R., Arnett, K.L., and Blacklow, S.C. (2008). The molecular logic of Notch signaling--a structural and biochemical perspective. J Cell Sci *121*, 3109-3119.

Gordon, W.R., Zimmerman, B., He, L., Miles, L.J., Huang, J., Tiyanont, K., McArthur, D.G., Aster, J.C., Perrimon, N., Loparo, J.J.*, et al.* (2015). Mechanical Allostery: Evidence for a Force Requirement in the Proteolytic Activation of Notch. Dev Cell *33*, 729-736.

Guruharsha, K.G., Kankel, M.W., and Artavanis-Tsakonas, S. (2012). The Notch signalling system: recent insights into the complexity of a conserved pathway. Nat Rev Genet *13*, 654-666.

Haltiwanger, R.S. (2008). Unsweetened Notch leads to myeloproliferation. Blood *112*, 214-215.

Hass, M.R., Liow, H.H., Chen, X., Sharma, A., Inoue, Y.U., Inoue, T., Reeb, A., Martens, A., Fulbright, M., Raju, S.*, et al.* (2015). SpDamID: Marking DNA Bound by Protein Complexes Identifies Notch-Dimer Responsive Enhancers. Mol Cell *59*, 685-697.

Hozumi, K., Negishi, N., Suzuki, D., Abe, N., Sotomaru, Y., Tamaoki, N., Mailhos, C., Ish-Horowicz, D., Habu, S., and Owen, M.J. (2004). Delta-like 1 is necessary for the generation of marginal zone B cells but not T cells in vivo. Nat Immunol.

Huang, C.H., Kuo, I.C., Xu, H., Lee, Y.S., and Chua, K.Y. (2003). Mite allergen induces allergic dermatitis with concomitant neurogenic inflammation in mouse. J Invest Dermatol *121*, 289-293.

Jin, M., Choi, J.K., Choi, Y.A., Kim, Y.Y., Baek, M.C., Lee, B.H., Jang, Y.H., Lee, W.J., Lee, S.J., Kim do, W.*, et al.* (2016). 1,2,4,5-Tetramethoxybenzene Suppresses House Dust Mite-Induced Allergic Inflammation in BALB/c Mice. Int Arch Allergy Immunol *170*, 35-45.

Johnston, N.A., Trammell, R.A., Ball-Kell, S., Verhulst, S., and Toth, L.A. (2009). Assessment of immune activation in mice before and after eradication of mite infestation. J Am Assoc Lab Anim Sci *48*, 371-377.

Kakuda, S., and Haltiwanger, R.S. (2017). Deciphering the Fringe-Mediated Notch Code: Identification of Activating and Inhibiting Sites Allowing Discrimination between Ligands. Dev Cell *40*, 193-201.

Kiel, M.J., Velusamy, T., Betz, B.L., Zhao, L., Weigelin, H.G., Chiang, M.Y., Huebner-Chan, D.R., Bailey, N.G., Yang, D.T., Bhagat, G.*, et al.* (2012). Whole-genome sequencing identifies recurrent somatic NOTCH2 mutations in splenic marginal zone lymphoma. J Exp Med *209*, 1553-1565.

Koch, U., and Radtke, F. (2007). Notch and cancer: a double-edged sword. Cell Mol Life Sci *64*, 2746-2762.

Kopan, R., and Ilagan, M.X. (2009). The Canonical Notch Signaling Pathway: Unfolding the Activation Mechanism. Cell *137*, 216-233.

Kovall, R.A., Gebelein, B., Sprinzak, D., and Kopan, R. (2017). The Canonical Notch Signaling Pathway: Structural and Biochemical Insights into Shape, Sugar, and Force. Dev Cell *41*, 228-241.

Kuang, Y., Golan, O., Preusse, K., Cain, B., Salomone, J., Campbell, I., Okwubido-Williams, F.V., Hass, M.R., Eafergan, N., Moberg, K.H.*, et al.* (2019). Enhancer architecture sensitizes cell specific responses to Notch gene dose via a bind and discard mechanism. bioRxiv.

Kueh, H.Y., Yui, M.A., Ng, K.K., Pease, S.S., Zhang, J.A., Damle, S.S., Freedman, G., Siu, S., Bernstein, I.D., Elowitz, M.B.*, et al.* (2016). Asynchronous combinatorial action of four regulatory factors activates Bcl11b for T cell commitment. Nat Immunol *17*, 956-965.

Langridge, P.D., and Struhl, G. (2017). Epsin-Dependent Ligand Endocytosis Activates Notch by Force. Cell *171*, 1383-1396 e1312.

Lee, C., Shin, H., and Kimble, J. (2019). Dynamics of Notch-Dependent Transcriptional Bursting in Its Native Context. Dev Cell *50*, 426-435 e424.

Lefebvre, C., Rajbhandari, P., Alvarez, M.J., Bandaru, P., Lim, W.K., Sato, M., Wang, K., Sumazin, P., Kustagi, M., Bisikirska, B.C.*, et al.* (2010). A human B-cell interactome identifies MYB and FOXM1 as master regulators of proliferation in germinal centers. Mol Syst Biol *6*, 377.

Lei, W., Liu, F., and Ness, S.A. (2005). Positive and negative regulation of c-Myb by Cyclin D1, cyclin-dependent kinases and p27 Kip1. Blood.

Liu, F., and Posakony, J.W. (2012). Role of architecture in the function and specificity of two Notch-regulated transcriptional enhancer modules. PLoS Genet *8*, e1002796.

Liu, F., and Posakony, J.W. (2014). An enhancer composed of interlocking submodules controls transcriptional autoregulation of suppressor of hairless. Dev Cell *29*, 88-101.

Liu, H., Chi, A.W., Arnett, K.L., Chiang, M.Y., Xu, L., Shestova, O., Wang, H., Li, Y.M., Bhandoola, A., Aster, J.C.*, et al.* (2010a). Notch dimerization is required for leukemogenesis and T-cell development. Genes Dev *24*, 2395-2407.

Liu, H., Chi, A.W., Arnett, K.L., Chiang, M.Y., Xu, L., Shestova, O., Wang, H., Li, Y.M., Bhandoola, A., Aster, J.C.*, et al.* (2010b). Notch dimerization is required for leukemogenesis and T-cell development. Genes Dev *24*, 2395-2407.

Liu, Z., Brunskill, E., Varnum-Finney, B., Zhang, C., Zhang, A., Jay, Y.P., Bernstein, I., Morimoto, M., and Kopan, R. (2015). The intracellular Domains of Notch1 and 2 Are Functionally Equivalent During Development and Carcinogenesis. Development *142*, 2452-2463.

Liu, Z., Chen, S., Boyle, S., Ilagan, M.X., Zhu, Y., Zhang, A., and Kopan, R. (2013). The Extracellular Domain of Notch2 Increases its Cell surface Abundance and Ligand Responsiveness During Kidney. Dev Cell *25*, 585-598.

Maillard, I., Weng, A.P., Carpenter, A.C., Rodriguez, C.G., Sai, H., Xu, L., Allman, D., Aster, J.C., and Pear, W.S. (2004). Mastermind critically regulates Notch-mediated lymphoid cell fate decisions. Blood *104*, 1696-1702.

Masek, J., and Andersson, E.R. (2017). The developmental biology of genetic Notch disorders. Development *144*, 1743-1763.

Mathern, D.R., Laitman, L.E., Hovhannisyan, Z., Dunkin, D., Farsio, S., Malik, T.J., Roda, G., Chitre, A., Iuga, A.C., Yeretssian, G., *et al.* (2014). Mouse and human Notch-1 regulate mucosal immune responses. Mucosal Immunol *7*, 995-1005.

Moats, C.R., Baxter, V.K., Pate, N.M., and Watson, J. (2016). Ectoparasite Burden, Clinical Disease, and Immune Responses throughout Fur Mite (Myocoptes musculinus) Infestation in C57BL/6 and Rag1(-/-) Mice. Comp Med *66*, 197-207.

Moolenbeek, C., and Ruitenberg, E.J. (1981). The "Swiss roll": a simple technique for histological studies of the rodent intestine. Lab Anim *15*, 57-59.

Moran, S.T., Cariappa, A., Liu, H., Muir, B., Sgroi, D., Boboila, C., and Pillai, S. (2007). Synergism between NF-kappa B1/p50 and Notch2 during the development of marginal zone B lymphocytes. J Immunol *179*, 195-200.

Nam, Y., Sliz, P., Pear, W.S., Aster, J.C., and Blacklow, S.C. (2007). Cooperative assembly of higher-order Notch complexes functions as a switch to induce transcription. Proc Natl Acad Sci U S A *104*, 2103-2108.

Nandagopal, N., Santat, L.A., LeBon, L., Sprinzak, D., Bronner, M.E., and Elowitz, M.B. (2018). Dynamic Ligand Discrimination in the Notch Signaling Pathway. Cell *172*, 869-880 e819.

Nellesen, D.T., Lai, E.C., and Posakony, J.W. (1999). Discrete enhancer elements mediate selective responsiveness of enhancer of split complex genes to common transcriptional activators. Developmental Biology *213*, 33-53.

Nigam, V., and Srivastava, D. (2009). Notch1 represses osteogenic pathways in aortic valve cells. J Mol Cell Cardiol *47*, 828-834.

Noah, T.K., and Shroyer, N.F. (2013). Notch in the intestine: regulation of homeostasis and pathogenesis. Annu Rev Physiol *75*, 263-288.

Obata, Y., Takahashi, D., Ebisawa, M., Kakiguchi, K., Yonemura, S., Jinnohara, T., Kanaya, T., Fujimura, Y., Ohmae, M., Hase, K., *et al.* (2012). Epithelial cell-intrinsic Notch signaling plays an essential role in the maintenance of gut immune homeostasis. J Immunol *188*, 2427-2436.

Okamoto, R., Tsuchiya, K., Nemoto, Y., Akiyama, J., Nakamura, T., Kanai, T., and Watanabe, M. (2009). Requirement of Notch activation during regeneration of the intestinal epithelia. Am J Physiol Gastrointest Liver Physiol *296*, G23-35.

Ong, C., Cheng, H., Chang, L.W., Ohtsuka, T., Kageyama, R., Stormo, D.G., and Kopan, R. (2006). Target selectivity of vertebrate Notch proteins: collaboration between discrete domains and CSL binding site architecture determine activation probability. J Biol Chem *281*, 5106-5119.

Pellegrinet, L., Rodilla, V., Liu, Z., Chen, S., Koch, U., Espinosa, L., Kaestner, K.H., Kopan, R., Lewis, J., and Radtke, F. (2011). Dll1- and Dll4-mediated Notch signaling is required for homeostasis of intestinal stem cells. Gastroenterology *140*, 1230-1240.

Peveling-Oberhag, J., Wolters, F., Doring, C., Walter, D., Sellmann, L., Scholtysik, R., Lucioni, M., Schubach, M., Paulli, M., Biskup, S.*, et al.* (2015). Whole exome sequencing of microdissected splenic marginal zone lymphoma: a study to discover novel tumor-specific mutations. BMC Cancer *15*, 773.

Piris, M.A., Onaindia, A., and Mollejo, M. (2017). Splenic marginal zone lymphoma. Best Pract Res Clin Haematol *30*, 56-64.

Pochanke, V., Hatak, S., Hengartner, H., Zinkernagel, R.M., and McCoy, K.D. (2006). Induction of IgE and allergic-type responses in fur mite-infested mice. Eur J Immunol *36*, 2434-2445.

Pui, J.C., Allman, D., Xu, L., DeRocco, S., Karnell, F.G., Bakkour, S., Lee, J.Y., Kadesch, T., Hardy, R.R., Aster, J.C.*, et al.* (1999). Notch1 expression in early lymphopoiesis influences B versus T lineage determination. Immunity *11*, 299-308.

Punareewattana, K., Smith, B.J., Blaylock, B.L., Longstreth, J., Snodgrass, H.L., Gogal, R.M., Jr., Prater, R.M., and Holladay, S.D. (2001). Topical permethrin exposure inhibits antibody production and macrophage function in C57Bl/6N mice. Food Chem Toxicol *39*, 133-139.

Quigley, D.A., To, M.D., Kim, I.J., Lin, K.K., Albertson, D.G., Sjolund, J., Perez-Losada, J., and Balmain, A. (2011). Network analysis of skin tumor progression identifies a rewired genetic architecture affecting inflammation and tumor susceptibility. Genome Biol *12*, R5.

Quigley, D.A., To, M.D., Perez-Losada, J., Pelorosso, F.G., Mao, J.H., Nagase, H., Ginzinger, D.G., and Balmain, A. (2009). Genetic architecture of mouse skin inflammation and tumour susceptibility. Nature *458*, 505-508.

Radtke, F., Fasnacht, N., and Macdonald, H.R. (2010). Notch signaling in the immune system. Immunity *32*, 14-27.

Radtke, F., Wilson, A., Stark, G., Bauer, M., van Meerwijk, J., MacDonald, H.R., and Aguet, M. (1999). Deficient T cell fate specification in mice with an induced inactivation of Notch1. Immunity *10*, 547-558.

Ray, A., and Dittel, B.N. (2010). Isolation of mouse peritoneal cavity cells. Journal of visualized experiments : JoVE.

Riccio, O., van Gijn, M.E., Bezdek, A.C., Pellegrinet, L., van Es, J.H., Zimber-Strobl, U., Strobl, L.J., Honjo, T., Clevers, H., and Radtke, F. (2008). Loss of intestinal crypt progenitor cells owing

to inactivation of both Notch1 and Notch2 is accompanied by derepression of CDK inhibitors p27(Kip1) and p57(Kip2). EMBO Rep *9*, 377-383.

Rossi, D., Trifonov, V., Fangazio, M., Bruscaggin, A., Rasi, S., Spina, V., Monti, S., Vaisitti, T., Arruga, F., Fama, R.*, et al.* (2012). The coding genome of splenic marginal zone lymphoma: activation of NOTCH2 and other pathways regulating marginal zone development. J Exp Med *209*, 1537-1551.

Saito, T., Chiba, S., Ichikawa, M., Kunisato, A., Asai, T., Shimizu, K., Yamaguchi, T., Yamamoto, G., Seo, S., Kumano, K.*, et al.* (2003). Notch2 is preferentially expressed in mature B cells and indispensable for marginal zone B lineage development. Immunity *18*, 675-685.

Santos, T.S.D., Tavares, R.S., and Farias, D.L.C. (2017). Splenic marginal zone lymphoma: a literature review of diagnostic and therapeutic challenges. Rev Bras Hematol Hemoter *39*, 146-154.

Schroeter, E.H., Kisslinger, J.A., and Kopan, R. (1998). Notch-1 signalling requires ligand-induced proteolytic release of intracellular domain. Nature *393*, 382-386.

Severson, E., Arnett, K.L., Wang, H., Zang, C., Taing, L., Liu, H., Pear, W.S., Shirley Liu, X., Blacklow, S.C., and Aster, J.C. (2017). Genome-wide identification and characterization of Notch transcription complex-binding sequence-paired sites in leukemia cells. Sci Signal *10*.

Shaya, O., Binshtok, U., Hersch, M., Rivkin, D., Weinreb, S., Amir-Zilberstein, L., Khamaisi, B., Oppenheim, O., Desai, R.A., Goodyear, R.J.*, et al.* (2017). Cell-Cell Contact Area Affects Notch Signaling and Notch-Dependent Patterning. Dev Cell *40*, 505-511 e506.

Shroyer, N.F., Helmrath, M.A., Wang, V.Y., Antalffy, B., Henning, S.J., and Zoghbi, H.Y. (2007). Intestine-specific ablation of mouse atonal homolog 1 (Math1) reveals a role in cellular homeostasis. Gastroenterology *132*, 2478-2488.

Siebel, C., and Lendahl, U. (2017). Notch Signaling in Development, Tissue Homeostasis, and Disease. Physiol Rev *97*, 1235-1294.

Sprinzak, D., Lakhanpal, A., LeBon, L., Santat, L.A., Fontes, M.E., Anderson, G.A., Garcia-Ojalvo, J., and Elowitz, M.E. (2010). Cis Interactions beween Notch and Delta generate mutally exclusive signaling states. Nature *465*, 86-90.

Stemmer, M., Thumberger, T., Del Sol Keyer, M., Wittbrodt, J., and Mateo, J.L. (2015). CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target Prediction Tool. PLoS One *10*, e0124633.

Tanigaki, K., Han, H., Yamamoto, N., Tashiro, K., Ikegawa, M., Kuroda, K., Suzuki, A., Nakano, T., and Honjo, T. (2002). Notch-RBP-J signaling is involved in cell fate determination of marginal zone B cells. Nat Immunol *3*, 443-450.

van Es, J.H., van Gijn, M.E., Riccio, O., van den Born, M., Vooijs, M., Begthel, H., Cozijnsen, M., Robine, S., Winton, D.J., Radtke, F*., et al.* (2005). Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and adenomas into goblet cells. Nature *435*, 959-963.

Vooijs, M., Ong, C.T., Hadland, B., Huppert, S., Liu, Z., Korving, J., van den Born, M., Stappenbeck, T., Wu, Y., Clevers, H*., et al.* (2007). Mapping the consequence of Notch1 proteolysis in vivo with NIP-CRE. Development *134*, 535-544.

Weng, A.P., Ferrando, A.A., Lee, W., Morris, J.P., IV, Silverman, L.B., Sanchez-Irizarry, C., Blacklow, S.C., Look, A.T., and Aster, J.C. (2004). Activating Mutations of NOTCH1 in Human T Cell Acute Lymphoblastic Leukemia. Science *306*, 269-271.

Witt, C.M., Won, W.J., Hurez, V., and Klug, C.A. (2003). Notch2 haploinsufficiency results in diminished B1 B cells and a severe reduction in marginal zone B cells. J Immunol *171*, 2783-2788.

A. Cooperativity

B. Notch1 → Notch1+/RA

WT 5'-ATCCTGCTCCGGAAC**AGG**GCCAC**A**GATCTGGATGCCCGAATGCAT
Bgl II

5'-ATCCTGCTCCGGAAC**gca**GCCACtgaTCTaGATGCCCGAATGCAT
Xba I

B'.

B".

C. Notch2 → Notch2RA/RA

WT 5'-tctagATTCTGATCCGCAAC**CGT**GTAAC**C**GATCTGGATGCCAGAATG

Mut 5'-tctagATTCTGATCCGCAAC**gcg**GTAAC**a**GATCTGGATGCCAGAATG
Bgl II

C'.

C".

N1$^{+/-}$, N2$^{+/-}$	N1$^{RA/-}$, N2$^{RA/-}$	N1$^{RA/-}$, N2$^{+/RA}$

E16.5

5/7 VSD

3/5 VSD

N2+/+	N2+/RA	N2RA/RA	N1+/RA; N2RA/RA	N1RA/RA; N2RA/RA
A 1.25cm	B 1.5cm	C 1.5cm	D 2.6cm	E 1.8cm

WT	N2+/RA	N1RA/RA	N2RA/RA	N1+/RA;N2RA/RA	N1RA/RA;N2RA/R
F	G	H	I	J	K

L   L'   HE   L''

N1+/RA; N2RA/RA spleen

N1+/RA; N2RA/RA lymph node

B220 DAPI   M   M'   N   N'

PAX5 DAPI   O   O'   P   P'

Marginal Zone

Mites

N2^{RA} N2

Hes1/5

Amplitude   Amplitude

Myb, FoxM1   Myb, FoxM1

MZB   MZB

SMZL

Intestine

N1^{RA}, N2^{RA} N1, N2

Hes1/5

1%DSS

IL1ß   IL1ß
IL4   IL4
IL6   IL6
IL17A   IL17A

Klf4, Atoh1

N1$^{RA/-}$, N2$^{RA/-}$

ISC Exhaustion