**Title:** A GT-seq panel for walleye (*Sander vitreus*) provides a generalized workflow for efficient

development and implementation of amplicon panels in non-model organisms.

**Running head**: Guide to develop and implement GT-seq panels

**Authors**: Matthew L. Bootsma[1*], Kristen M. Gruenthal[2], Garrett J. McKinney[3], Levi Simmons[1],

Loren Miller[4], Greg G. Sass[5], Wesley A. Larson[6]

**Affiliations**

[1] Wisconsin Cooperative Fishery Research Unit, College of Natural Resources, University of

Wisconsin-Stevens Point, 800 Reserve St., Stevens Point, WI 54481, USA,

mbootsma@uwsp.edu, lsimm290@uwsp.edu

[2] Office of Applied Science, Wisconsin Department of Natural Resources, Wisconsin

Cooperative Fishery Research Unit, College of Natural Resources, University of Wisconsin-

Stevens Point, 800 Reserve St., Stevens Point, WI 54481, USA,

kristen.gruenthal@wisconsin.gov

[3] NRC Research Associateship Program, Northwest Fisheries Science Center, National Marine

Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd E,

Seattle, WA 98112, USA, garrett.mckinney@noaa.gov

[4] Minnesota Department of Natural Resources, University of Minnesota, 135 Skok Hall, 2003

Upper Buford Circle, St. Paul, MN 55108, USA, loren.miller@state.mn.us

[5] Escanaba Lake Research Station, Office of Applied Science, Wisconsin Department of Natural

Resources, 3110 Trout Lake Station Drive, Boulder Junction, WI 54512, USA,

gregory.sass@wisconsin.gov

[6] U.S. Geological Survey, Wisconsin Cooperative Fishery Research Unit, College of Natural

Resources, University of Wisconsin-Stevens Point, 800 Reserve St., Stevens Point, WI 54481,

USA, wes.larson@uwsp.edu

**\*Corresponding author**

**Keywords:** Amplicon sequencing, GT-seq, microhaplotype, parentage analysis, genetic stock

identification, walleye

**Abstract (250 words or less)**

   Targeted amplicon sequencing methods, such as genotyping-in-thousands by sequencing (GT-seq), facilitate rapid, accurate, and cost-effective analysis of hundreds of genetic loci in thousands of individuals, but studies describing detailed workflows of GTseq panel development are rare. Here, we develop a dual-purpose GT-seq panel for walleye (*Sander vitreus*) and discuss trade-offs associated with different development and genotyping approaches. Our GT-seq panel was developed using restriction site-associated DNA data from 954 individuals sampled from 23 populations in Minnesota and Wisconsin, USA. We then conducted simulations to test the utility of loci for parentage analysis and genetic stock identification and designed 600 primer pairs to maximize joint accuracy for these analyses. We conducted three rounds of primer optimization to remove loci that overamplified and our final panel consisted of 436 loci. Optimization focused on reducing variation in amplification rate among loci and minimizing the proportion of off-target sequence, both of which are important considerations for developing large GT-seq panels. We also explored different approaches for DNA extraction, multiplexed polymerase chain reaction (PCR) amplification, and cleanup steps during the GT-seq process and discovered the following: (1) inexpensive Chelex extractions performed well for genotyping, (2) the exonuclease I and shrimp alkaline phosphatase (ExoSAP) procedure included in some current protocols did not improve results substantially and was likely unnecessary, and (3) it was possible to PCR amplify panels separately and combine them prior to adapter ligation. Well-optimized GT-seq panels are valuable resources for conservation genetics and our findings should aid in their construction in myriad taxa.

**Introduction**

The development of genotyping-by-sequencing (GBS) methods has allowed collection of

data from thousands of markers across a genome, enabling research that was not possible using

traditional genetic approaches (Davey et al., 2011; Narum et al., 2013). For example, studies

using thousands of markers genotyped with restriction site-associated DNA (RAD) sequencing

have shown improved sensitivity for detecting inbreeding depression (Hoffman et al., 2014),

increased resolution for determining complex phylogenies (Wagner et al., 2013), and allowed

researchers to observe selection on introduced alleles (Bay et al., 2019). Many genetic analyses,

however, can be conducted efficiently with genotypes from tens to hundreds of single nucleotide

polymorphisms (SNPs) (Anderson & Garza, 2006), making more expensive approaches such as

RAD-seq unnecessary (Meek & Larson, 2019). Two such analyses that have been widely used in

conservation genetics and molecular ecology for decades, are parentage analysis and genetic

stock identification (GSI).

Parentage analysis involves assigning offspring to putative parents by comparing

genotypes at multiple loci, while GSI infers the natal origins of individuals by leveraging

baseline allele frequency estimates from populations or reporting groups. These techniques were

first conducted using allozyme markers genotyped with protein electrophoresis. Although these

analyses were groundbreaking, they often lacked statistical power except in cases of highly

diverged stocks or simple pedigrees. The adoption of highly variable microsatellite markers in

the 1990s greatly increased statistical power, allowing these two techniques to become widely

adopted (Luikart & England, 1999). Despite the advances made possible by microsatellites,

problems associated with homoplasy (Garza & Freimer, 1996), locus discovery (Navajas et al.,

1998), and reproducibility among laboratories led researchers to explore the potential of biallelic

SNPs for GSI and parentage analysis (Seeb et al., 2011).

Although SNPs are less powerful than microsatellites on a per marker basis, SNPs are

more abundant in the genome, generally have low genotyping error rates, and can be genotyped

using SNP panels capable of efficiently screening a large number of samples (Brumfield et al.,

2003; Morin et al., 2004).  Early SNP panels were constrained, however, in the availability of

molecular markers suitable for genotyping and genotyping costs associated with 5' exonuclease

chemistry (Seeb et al., 2011). These constraints were significantly lessened with the proliferation

of next-generation sequencing (NGS) technology. For example, methods such as RADseq

facilitate quick and affordable discovery of thousands of candidate loci, which can then be

selected among for specific purposes.

As SNP discovery has become less prohibitive, methods of selecting the most

informative SNPs for a given study have advanced (Storer et al., 2012). Previous research has

shown that information content will vary among SNPs depending on the context within which

they are applied and location within the genome (i.e. coding or non-coding regions). For

example, Ackerman et al. (2011) found that SNPs under diversifying selection provide increased

accuracy and precision in GSI of sockeye salmon (*Oncorhynchus nerka*) from the Copper River,

Alaska. In general, previous studies have shown that GSI accuracy is generally positively

correlated with differentiation (e.g., $F_{ST}$) and, to a lesser extent, diversity (e.g., heterozygosity)

(Ackerman et al., 2011; Bradbury et al., 2011; Storer et al., 2012). Studies of SNP selection

methods for parentage analysis, however, have found that high diversity is the most important

attribute to consider when creating a panel (Baetscher et al., 2018). More recently, analytical

techniques have shifted towards consideration of closely linked SNPs (i.e. microhaplotypes),

which effectively increases the diversity at a locus and has proven useful for parentage and GSI

tests (Baetscher et al., 2018; McKinney, Seeb, et al., 2017; Reid et al., 2019). While obtaining

microhaplotypes using previous 5' exonuclease methods would require independent assays for

each SNP at a locus and statistical phasing, NGS technology has enabled the joint genotyping of

multiple SNPs within single reads, making microhaplotype data easily obtainable through a

simple modification in analytical approach.

One recently developed GBS method that improves upon previous high-throughput

genotyping technologies, such as 5' exonuclease chemistry, is Genotyping-in-Thousands by

sequencing (GT-seq). This method enables genotyping hundreds of SNPs in thousands of

individuals on a single NGS lane through the use of highly-multiplexed polymerase chain

reaction (PCR) (Campbell et al., 2015). GT-seq does not require an allele-specific probe, can

genotype multiple SNPs within an amplicon using a single primer pair, and is substantially less

expensive than 5' exonuclease chemistry, especially in the context of genotyping thousands of

individuals.

Despite its benefits, GT-seq is not yet widely used outside of salmonids. Early

applications to non-model organisms, however, have shown great promise for this method's

versatility, including the ability to reveal dispersal and mating patterns in a complex environment

(Baetscher et al., 2019), provide insight to the ecological and evolutionary dynamics of

secondary contact (Reid et al., 2019), and understand population diversity in systems that are

heavily influenced by climate change (Pavinato et al., 2019). Pedigree analysis in wild

populations is highly dependent upon the ability to genotype large sample sizes to increase the

likelihood of detecting kin relationships, toward which GT-seq is ideally suited. Moreover, GT-

seq has proven capable of generating high-quality genotypes from low-quality DNA samples

(Natesh et al., 2019; Schmidt et al., 2019), making it a viable approach for monitoring

endangered or elusive species.

127            While GT-seq panels have been developed to maximize accuracy for GSI (McKinney et

al., 2019) or parentage (Baetscher et al., 2018) analyses, the potential for developing dual-

purpose panels is largely unexplored. Moreover, developing GT-seq panels is a relatively

involved task and, to this point, there are limited resources providing standardized workflows

and guidelines for efficient panel construction (but see Campbell et al., 2015; McKinney et al.,

2019).  At a basic level, panel construction involves SNP discovery, SNPs selection, primer

design, and panel optimization (see Baetscher et al., 2018; McKinney et al., 2019; Schmidt et al.,

2019); however, within this general framework there are many decision points in panel

development related to primer selection, multiplexing approaches, laboratory protocols, and

analysis parameters that have yet to be addressed. We used walleye (*Sander vitreus*) from

Minnesota and Wisconsin, USA, as a test case to investigate various tradeoffs associated with

GT-seq panel development and optimization and leveraged our collective experience to provide

guidelines for researchers developing GT-seq panels.

140            Walleye are an apex predator and one of the most prized sportfish throughout their native

and introduced range. Recently, many walleye populations have declined across the Midwestern

United States (Embke et al., 2019; Hansen et al., 2015; Rypel et al., 2018), prompting increases

in stocking efforts relative to already large and long-term regional stocking programs that have

existed for decades. Genetic studies have been used to guide these efforts by informing

broodstock selection and general stocking practices. Genetic variation in walleye from this

region was first characterized by Fields et al. (1997), who found geographic-based patterns of

genetic structure, but limitations related to sample size and molecular marker choice resulted in

the use of contemporary watershed boundaries as genetic management units. This research was

later expanded upon by Hammen and Sloss (2019), who attempted to further define genetic

structure in the Ceded Territory of Wisconsin, approximately the northern third of the state, and

test whether significant genetic structure existed between distinct hydrological basins within this

region. Once again, constraints associated with available molecular markers used in a system

with not only low differentiation, but also extensive stocking precluded definition of fine-scale

structure. This system provides an excellent model for applying genomic techniques to

discriminate populations and evaluate hatchery programs using parentage analysis.

Like many intricacies of genomics research, GT-seq panel development is a process that

is at once broadly generalizable to non-model organisms and highly specific to the taxa it is

applied to. While the overarching steps (Fig. 1) will remain constant, there are many decision

points within that will require informed thought and decision. Using walleye, a species with few

well-established genomic resources, as a model, we examined the methods inherent to GT-seq

panel development in a manner that identifies critical decision points in the process and

illuminates the nuances associated with them. Our overarching goal was to design a dual-purpose

GT-seq panel optimized for parentage analysis and GSI in walleye. The creation of this panel

allowed us to address the following specific objectives: (1) investigate the tradeoffs between

choosing markers for parentage analysis versus GSI, (2) explore the most efficient way to design

an optimized panel, and (3) evaluate various laboratory approaches to maximizing the efficiency

of GT-seq genotyping. We provide an in-depth discussion of our experiences designing the panel

and outline important topics that should aid researchers in designing future GT-seq panels.

**Materials and Methods**

*Sample collection*

Tissue samples were collected from adult walleye from 23 inland lakes across Wisconsin,

Minnesota, and the St. Louis River (border water) (Fig. 2a, Table 1) and stored in 95% ethanol

until DNA extraction. We obtained samples from as many major drainages as possible across the

two states, with an emphasis on the Wisconsin and Chippewa River drainages in Wisconsin,

which were difficult to differentiate using microsatellites (Hammen & Sloss, 2019); in

Minnesota, sampling focused primarily on major sources of wild broodstock for stocking

programs. Samples were collected by the Wisconsin and Minnesota Departments of Natural

Resources using fyke nets or electrofishing. Sampling took place during the spring spawning

runs of April 2015 and 2017 and fall surveys in August and September of 2015 and 2017.

Stocked individuals may be tagged, or fin clipped; we inspected all sampled individuals for tags

or fin clips to avoid as many individuals as possible that were of stocked origin as possible.

*Preparation of RAD sequencing libraries*

Genomic DNA was extracted in a 96-well format with Qiagen DNeasy Blood and Tissue

Kits. Extracted DNA was quantified using a Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen,

Waltham, MA) and normalized to 20ng/µl. DNA was then prepared for RADseq library

preparation following the BestRAD protocol (Ali et al., 2016). Briefly, DNA was digested in a 2

µl reaction with the restriction enzyme *SbfI*, and biotinylated barcode adaptors were ligated to

the 5' cut ends. DNA shearing was conducted using a 12.5 µl fragmentase reaction. Library

preparation was conducted using an NEBNext Ultra DNA Library Prep Kit for Illumina (NEB,

Ipswich, MA), with a 12-cycle PCR enrichment. RAD library quality was inspected on a 2%

agarose gel before undergoing a final AMPure XP (Beckman Coulter, Indianapolis, IN)

purification and quantification on a Qubit 2.0 Fluorometer (ThermoFisher Scientific, Waltham,

MA). Libraries were sequenced using paired-end (PE) 150 technology on a HiSeq 4000

(Illumina, San Diego, CA) at the Michigan State University Genomics Core Facility or

Novogene Corporation, Inc. (Davis, CA). Sequencing was conducted to achieve a target of over

one million retained reads per individual.

*Analysis of RAD data to discover SNPs*

Loci were identified and genotyped in STACKS v.2.2 (Rochette et al., 2019) without

using gapped alignments. Raw reads were demultiplexed and barcodes were trimmed in

*process_radtags* (parameter flags: -e *SbfI*, -c, -q, -filter_illumina, -r, --bestrad). RAD-tags were

assembled into putative RAD loci with *ustacks* using the bounded model (bound_high =  0.05, --

disable-gapped) and allowing for a maximum of three nucleotide mismatches (-M = 3) and four

stacks per locus (-max_locus_stacks = 4), as well as a minimum depth of three (-m = 3). The

calling of haplotypes from secondary reads was disabled (-H). A catalog of consensus loci was

assembled in *cstacks* using the two individuals with the highest number of retained reads from

each population, allowing a maximum of three mismatches between sample loci (n = 3, --

disable-gapped). After matching all samples against the catalog in *sstacks* (--disable-gapped),

data were oriented by locus with *tsv2bam*, and individual genotypes were called in *gstacks*, with

paired-end reads incorporated. Genotypes were exported in variant call format (vcf) using

*populations*, with loose filtering parameters (SNPs present at > 5% of individuals, minimum

minor allele frequency of > 0.005).

Comprehensive filtering of individuals and genotypes was conducted in vcftools v0.1.15

(Danecek et al., 2011) by: 1) removing individuals missing > 20% of SNP calls, 2) removing

SNPs that were missing in > 20% of individuals, and 3) removing SNPs that were not in the first

140 base pairs of the RAD-tag, effectively reducing the dataset to include SNPs detectable using

single-read (SR) 150 sequencing to simplify downstream amplicon design; to control for

genotyping error, SNPs with a minor allele count $\leq 3$ were also removed. Putative duplicated loci

were identified in HDplot (McKinney, Waples, et al., 2017) ($H > 0.5$, $-7 < D < 7$) and removed

with vcftools. Retained individuals and SNPs were used to form whitelists for input into

*populations* that output a filtered vcf of multi-SNP haplotypes, which was then filtered to remove

loci with more than 10 alleles and used in simulations for locus selection. We also estimated

single-SNP $F_{IS}$ across all populations using diveRsity v1.9.90 (Keenan et al., 2013) and excluded

any SNPs with $F_{IS}$ values $> 0.2$ or $< -0.2$ from locus selection. Additionally, loci with a SNP in

the first 10 base pairs of the RAD-tag were excluded to allow room for forward primer design.

*Analysis of population structure, locus selection, and panel assessment*

To understand population structure in our system and ensure that selected loci could

facilitate accurate parentage assignment and GSI, we evaluated patterns of genetic divergence

using pairwise $F_{ST}$ (Table S1) estimated in Arlequin v3.5.2 (Excoffier & Lischer, 2010) and

constructed a dendrogram (Fig. 2b) using Nei's distance in poppr v2.8.2 (Kamvar, Tabim, &

Grünwald, 2014). These analyses facilitated identification of population pairs that would be

challenging to discriminate and supported historical data suggesting several populations were

founded from hatchery sources located outside of their drainage basin (Escanaba Lake, Sanford

Lake, and Lake Millicent in Wisconsin); these populations were removed from simulations of

panel accuracy to ensure that selected loci would best represent the natural genetic patterns of the

region.

After initial population genetic analyses, loci were selected for primer development by

constructing several test panels from the RAD data and simulating assignment accuracy for

parentage and GSI. Previous research suggested that choosing loci with greater genetic

differentiation (e.g., $F_{ST}$) should maximize accuracy for GSI (Ackerman et al., 2011; Storer et

al., 2012), while choosing loci with higher diversity (e.g., heterozygosity and number of alleles)

maximizes accuracy for parentage (Baetscher et al., 2018). We therefore constructed the test

panels using single-SNP $F_{ST}$ estimated in diveRsity v1.9.90 (Keenan et al., 2013) as well as

expected heterozygosity at a multi-SNP haplotype ($H_{E\_mhap}$) and the number of alleles at a locus

estimated in adegenet v2.1.1 (Jombart & Ahmed, 2011). All simulations were conducted with

genotypes coded as multi-SNP haplotypes.

GSI accuracy for each panel was assessed via 100% simulations implemented in rubias

(Moran & Anderson, 2018) using the *assess_reference_loo* function (mixsize = 200, reps =

1000). Populations were aggregated into reporting units based on hydrological basins (Table 1).

Collections within a simulation were drawn from a Dirichlet distribution with all parameters

equal to 10 (i.e., each simulation's prior contained approximately equal proportions of each

population for the given reporting unit). Individuals were assigned to reporting groups if they

had a cumulative probability of > 70%. Unfortunately, limited sample sizes in some reporting

units prevented creation of separate training and holdout datasets as suggested by Anderson

(2010), thus assignment accuracies presented here may be upwardly biased and would need to be

reassessed more thoroughly for populations involved in an applied study.

Parentage simulations were run in CKMRsim (Anderson,

https://zenodo.org/record/820162), which employs a variant of the importance-sampling

algorithm of Anderson and Garza (2006) that allows for more accurate estimates of very small

false-positive rate (FPR: per-pair rate of truly unrelated individuals being inferred as related)

relative to those obtained using standard Monte Carlo methods (Baetscher et al., 2018).

Parentage analyses were conducted following the methods of Baetscher et al. (2018), whereby

log-likelihood ratios between a tested relationship and the hypothesis of no relationship are

computed from the calculated probabilities of genotype pairs for related individuals simulated

from allele frequency estimates. Distributions of simulated log-likelihood ratios are then used to

compute FPRs. Using this approach, we estimated FPRs for parent-offspring (PO), full-sibling

(FS), and half-sibling (HS) relationships at false-negative rates (FNR: per-pair rate of truly

related individuals being inferred as unrelated) ranging from 0.01 to 0.1.

Panels of 600 unique loci were iteratively selected, choosing loci based first on rank

$F_{ST}$ then rank $H_{E\_mhap}$, and their utility was tested by conducting GSI tests and parentage

simulations. We ultimately defined three panels of 600 loci that best described the tradeoffs

between markers selected based on $F_{ST}$ and heterozygosity. Loci in these panels were chosen by

selecting 1) the top 600 loci based on $F_{ST}$, 2) the top 300 loci based on $F_{ST}$ and 300 based on

$H_{E\_mhap}$, and 3) the top 600 loci based on $H_{E\_mhap}$. These panels are hereafter referred to as

$F_{ST\_600}$, Composite$_{600}$, and Diversity$_{600}$, respectively. Through further testing, we determined

that a variation of the Composite$_{600}$ panel, with 250 loci based on $H_{E\_mhap}$ and 350 loci based on

$F_{ST}$, delivered optimal performance for GSI and parentage analyses and proceeded to design

primers for the selected loci.

*Primer Design*

To design PCR primers for the selected loci, their consensus sequences were subset

from the STACKS catalog into a FASTA file for import into Geneious Prime® 2019.1.1

(https://www.geneious.com). The vcf file produced in the vcftools step containing all SNPs and

alleles within a consensus sequence was included to ensure primers were properly designed (i.e.,

should a SNP fall within a primer binding region, a degenerate nucleotide could be inserted or

the primer re-designed). Primer pairs were iteratively designed, with optimal target parameters

defined as a primer length of 20 bp, product size of 140 bp to facilitate genotyping with SR

chemistry, $T_m$ of 60° C, GC content of 50%, and no more than four of the same base repeated

consecutively (i.e., poly-X repeats). Primers identified as matching one or more off-target sites,

which could lead to amplification of multiple products, were redesigned. Given that not all 600

candidate loci initially identified were suitable candidates for primer development, we continued

to iteratively select loci and design associated primers until we reached our target of 600 loci.

Unfortunately, the loci selected for primer design were based on data containing a subset of

individuals with discordant encoded and true identities as a result of transposition of barcodes

during demultiplexing. Despite these discrepancies, the effect was likely minor as only 8% of

individuals were incorrectly assigned to reporting units prior to simulation. Simulation results

shown here were conducted using corrected data.

*GT-seq optimization*

297           GT-seq was conducted following the methods of Campbell et al. (2015), with

modification to the multiplex thermal cycling conditions (95 °C hold for 15 min; five cycles of

95 °C for 30 s, 5% ramp to 57 °C for 2 min, 72 °C 30 s; and 10 cycles of 95 °C for 30 s, 65 °C

for 30 s, and 72 °C 30 s) and post-normalization dual-sided SPRI size-selection and purification

(0.6X plus 0.4X) to further restrict the product size range (e.g., primarily  toward removal of

primer inter-hybridization). Final library quality control consisted of confirmation of

amplification and barcoding by SYBR Green-based RT-qPCR (Stratagene Mx3005P QPCR

System, Agilent, Santa Clara, CA), visualization on a 2% agarose E-Gel (Invitrogen, Carlsbad,

CA), and quantification using picogreen. Libraries were then sequenced at the University of

Wisconsin-Madison Biotechnology Center (UWBC) DNA Sequencing Facility on a MiSeq

(Illumina) using $2 \times 150$ bp flowcells.

Demultiplexed amplicon sequencing data were processed using *GTscore v1.3*

(McKinney et al., 2019). *GTscore* generates *in-silico* primer-probe sequences from a catalog of

loci generated in STACKS, that are then matched to amplicon sequences and call genotypes for

individual SNPs as well as multi-SNP haplotypes. *GTscore* also enables separation of on-target

sequence reads (i.e., reads containing both an *in-silico* primer and associated probe) from reads

produced as a result of primer cross-hybridization. Primer-probe file development was

accomplished with *sumstatsIUBconvert.pl* by obtaining the IUB code information for each SNP

from the sumstats.tsv file produced in the STACKS pipeline, converting catalog sequences

produced in the STACKS pipeline to FASTA sequences using *catalog2fasta.pl*, and merging

IUB code information with the catalog.fasta using *fasta2IUB.pl*. This primer-probe file was then

input for *AmpliconReadCounter.pl*, along with an individual's fastq file, to produce read count

summaries of primers and probes.

Overall, we conducted three rounds of panel optimization to identify and remove loci

that had disproportionately high amplification rates (i.e., "overamplifiers") and ensure that our

panel was capable of delivering a high proportion of on-target reads for each locus as well as

homogeneous amplification rates among loci. The first round of optimization used DNA from a

single walleye from Sanford Lake, WI, while the second and third rounds were conducted on

subsets of 24 individuals from each of four populations (96 individuals total) originally included

in the RADseq study: Delavan Lake, Medicine Lake, and the Wolf River in Wisconsin and the

Pine River in Minnesota. Upon completing the final optimization, the characteristics of retained

loci were compared to those of loci culled from the panel. This was done by performing a

Welch's two sample t-test ($\alpha = 0.05$) between the GC:AC ratio of primers that were retained and

those culled and between the GC:AC ratio of DNA templates retained and culled, based on the

first 140 bp of the template as this was the region in which SNPs were targeted.

GT-seq libraries from each round were collectively analyzed for PCR accuracy

and uniformity. Accuracy was measured by calculating the proportion of reads containing *in-*

*silico* primer sequences (total reads) relative to those that also contained *in-silico* probes.

Uniformity of amplification among loci was determined by calculating the proportion of total

reads that were allocated to the top 10% of loci, based on locus read counts (prop_reads_T10); if

amplification was perfectly uniform across loci, we would expect prop_reads_T10 to account for

exactly 10% of total reads. Given that amplification rates vary substantially within a panel, we

compared among locus performance by plotting the relative $\log_{10}$ abundance of total and on-

target reads at each locus in descending order, which facilitated visual identification of

overamplifiers. As among-locus amplification rates evened out after the first optimization, the

on-target proportion of reads at each locus became a factor in retaining or excluding loci during

the second optimization.

*Testing methodological modifications and performance analysis*

During panel optimization, we compared the quality of GT-seq libraries prepared

from DNA extracted with Qiagen DNeasy and a more cost-effective chelating resin-based

procedure. Performance of libraries was compared using Bonferroni corrected ($\alpha = 0.016$)

Tukey's HSD for the number of on-target reads and the proportion of total reads that were on-

target, after determining whether significant differences existed among libraries via a one-way

ANOVA ($\alpha = 0.05$). DNA was extracted from the 96 test individuals twice, first using Qiagen

DNeasy and again with a 10% Chelex 100 (200-400 mesh; Bio-Rad, Hercules, CA) solution

containing 1% each of Nonidet P-40 and Tween 20 (Millipore Sigma, St. Louis, MO).

Additionally, we aimed to further reduce the cost per sample by evaluating the need for certain

library preparation steps. Specifically, we compared results with and without the exonuclease I

and shrimp alkaline phosphatase (ExoSAP) procedure included in Campbell et al. (2015) to

remove PCR inhibitors and free nucleotides. GT-seq was therefore conducted on all individuals

in triplicate: 1) Qiagen with ExoSAP, 2) Chelex with ExoSAP, and 3) Chelex without ExoSAP,

and all tests were sequenced on the same MiSeq lane. Finally, we tested whether the number of

loci that could be genotyped simultaneously could be increased by conducting multiple PCRs.

We accomplished this by dividing our optimized primer panel into two non-overlapping primer

pools before multiplex PCR amplification. We then merged PCR products from the separate

pools prior to the barcoding PCR. The sequencing performance of this joint panel was then

compared to the single multiplex containing the full panel using a Welch's two sample t-test ($\alpha$ =

0.05).

We examined genotype concordance between RADseq and GT-seq across GT-seq

read depths using the fully optimized panel in the third round. Genotypes were called using

*PolyGen* (McKinney et al., 2018), an extension of the *GTscore* pipeline that uses the same

maximum-likelihood algorithm as STACKS v1 for diploid, bi-allelic loci. Because low read

depths can lead to high estimates of genotyping error, thereby increasing rates of allelic dropout

(Catchen et al., 2013), genotypes were only compared if they had greater than 60× coverage in

RADseq. We then modeled the relationship between GT-seq read depth and genotype

concordance using only read depths with more than 30 genotypes to ensure that estimates of

genotype concordance at a given depth had adequate sample sizes.

As a final proof of concept, we tested the optimized panel on a sample of 570 walleye

obtained from Escanaba Lake, WI, using the methods described above to estimate the variance in

read depth among loci within a pool. We retained only loci present in more than 70% of

individuals and individuals genotyped at more than 70% of loci.

**Results**

*Analysis of ascertainment dataset*

A total of 954 individuals from 23 populations were RAD sequenced, with an average of

42 individuals per population (Table 1). Sequencing yielded 1,313,358 retained reads on average

per individual (range = 8,941 - 8,176,163). Initial sequence data were used to identify 682,223

putative SNPs. After passing sequence data through quality filters, 839 individuals and 20,597

SNPs were retained (Table S2).

Population estimates of $H_O$ (0.144 - 0.179), allelic richness (1.498 - 1.674), and $F_{IS}$ (-

0.050 - 0.017) were relatively similar across locations (Table 1). Populations from Minnesota

had slightly lower diversity, which may be due to ascertainment bias as 14 of the 23 populations

were from Wisconsin. The highest genetic differentiation was observed between populations

from Minnesota and Wisconsin, with further structuring by drainage basin within each state (Fig.

2b, Table S1). Structuring was higher in Minnesota, with most populations showing a relatively

high degree of isolation (average $F_{ST}$ = 0.07, Table 2). Structure in Wisconsin was shallower

(average $F_{ST}$ = 0.03, Table 2) and only loosely correlated with drainage basins. From these

results, we constructed 13 reporting groups to facilitate GSI to identifiable genetic units (Table

1). All the reporting groups from Minnesota contained single populations, whereas in Wisconsin,

while the Rock-Fox and Wolf River groups contained single populations, the Wisconsin and

Chippewa River groups each contained five populations. Some single populations in the

Wisconsin and Chippewa Rivers were distinctly identifiable (e.g., Eau Claire River, Medicine

Lake), but we grouped these populations within their drainage basin of origin as the panel will

likely be used this way for management purposes.

*Locus selection and panel assessment*

GSI accuracy was similar among the three panels, with < 1% difference in average

accuracy between the panel with loci chosen based solely on differentiation ($F_{ST}\_600$) and the

panel based solely on diversity (Diversity_600) (Fig. 3, Table 3). Average assignment accuracy

was > 90% for nine of the 13 reporting units in all panels (Fig. 3a). The remaining four reporting

units had average assignment accuracies ranging from 78% to 86%. Three of these units (upper

Chippewa River, WI; St. Louis River, MN/WI; and Red Lake, MN) are known to have admixed

stocking histories, while the fourth, North Fork Crow River, MN, included Lake Koronis, which

had the fewest individuals retained after filtering (n = 15). Misassigned individuals from the St.

Louis River, MN, and Red Lake, MN groups primarily assigned to the Pike River, MN, an

unsurprising result given that fish from the Pike River contributed to the recovery of the

collapsed walleye fishery in Red Lake (Logsdon et al., 2016) and fish in the St. Louis River

watershed. Misassignments from the Upper Chippewa basin primarily assigned to the Upper

Wisconsin basin due to the lower differentiation described previously.

The populations with the lowest assignment accuracies were found in the Chippewa

River and Wisconsin River reporting groups (Table S3, S4, S5), particularly in northern

Wisconsin near the headwaters of the Chippewa and Wisconsin River drainages, and included

Big Arbor Vitae Lake ($F_{ST}\_600$ accuracy = 74%), Manitowish Lake ($F_{ST}\_600$ accuracy = 58%), and

Turtle Flambeau Flowage ($F_{ST}\_600$ accuracy = 63%). A large portion (> 10%) of the simulated

individuals from these populations could not be assigned to any population, providing further

support for the genetic similarity of these two reporting groups. A high proportion of individuals

from Big Arbor Vitae Lake were assigned to Manitowish Lake (12%) and vice versa, from

Manitowish Lake to Big Arbor Vitae Lake (20%). Most misassignments in the Turtle Flambeau

Flowage were to Kawaguesaga Lake (16%). Populations with high misassignment rates also

tended to have short branch lengths in the dendrogram and were often located near the root of a

clade (Fig. 2b). Furthermore, the two populations from the upper Chippewa basin (Manitowish

Lake and Turtle Flambeau Flowage) had lower pairwise $F_{ST}$ values, on average, relative to

populations from the upper Wisconsin basin than they did with other populations from the upper

Chippewa basin.

The Diversity$_{600}$ panel had the highest accuracy for assigning kin relationships, the

Composite$_{600}$ panel showed intermediate performance and the $F_{ST\_600}$ panel had the lowest

accuracy rate (Fig. 3b, Table 3). For all panels, FPRs were $< 10^{-20}$ for PO and FS relationships,

indicating all panels would perform adequately for reconstructing most relationships in most

study systems. Inter-panel performance did, however, range widely, from an FPR of $4.68 \times 10^{-34}$

for $F_{ST}\_600$ to $2.74 \times 10^{-80}$ for Diversity$_{600}$ panel at an FNR of 0.01. Within panels, FPR was

inversely related to FNR.

Primers were designed using a modified Composite_600 panel, with 250 loci chosen

based on $H_{E\_mhap}$ and 350 chosen based on $F_{ST}$, as this panel delivered the best joint accuracy for

GSI and kinship analyses (Fig. 3, Table 3). Of the initial 600 loci initially selected for primer

design, 100 were not suitable for primer design, and thus, iterative selection of loci meeting

primer design requirements was continued until the targeted number of $F_{ST}$ and diversity markers

was met.

*GT-seq optimization*

Initial amplification and MiSeq sequencing of all 600 loci yielded 4,655,071 reads

containing intact i7 barcode sequences, with 4,150,910 reads (89%) matching *in-silico* primer

sequences. Locus specificity was considered via the proportion of total reads that were on-target,

which was 1,031,707 (24.9%) (Table 4). In terms of amplification uniformity among loci,

prop_reads_T10 accounted for 3,526,201 (85.0%) of the 4,150,910 total reads. A cutoff of 3,000

reads per locus was then visually identified (Fig. 4a); loci producing more than 3,000 reads (n =

123) were deemed overamplifiers and discarded prior to further optimization.

For the second round of optimization, the remaining 477 primers pairs produced

12,653,262 reads containing intact i7 barcode sequences, and 9,347,591 (74%) matched *in-silico*

primer sequences. Locus specificity improved, with 3,268,293 (35.0%) of the total reads

successfully aligning to *in-silico* probe sequences (Table 4). Improvement was also observed in

the uniformity of amplification across loci, with prop_reads_T10 equating to 72.5% (6,776,302)

of total reads. Because locus performance was less variable in this round of testing, the

individual on-target proportion of reads at a locus was also considered while culling undesirable

loci. As such, loci visually identified as overamplifiers were again discarded if they did not

display high on-target read proportions (n = 41, Fig. 4b).

The third GT-seq test was used to determine the functional performance of the panel and

aimed to target 858 SNPs across 436 loci (Fig. 4c). This test produced 7,282,101 reads with

intact i7 barcodes, and 6,827,424 (94%) matched to *in-silico* primers. Locus specificity of primer

pairs improved greatly in this test, as 6,262,523 (91.7%) of the total reads were also on-target

(Table 4). Likewise, the variation in amplification rates across loci decreased as evidenced by

prop_reads_T10 decreasing to 36.6% (2,148,932) of the total reads.

Upon completion of panel optimization, a small but significant difference was observed

between the GC content of primers that were retained (mean = 49.2%) and primers that were

removed (mean = 51.4%, df = 602, t = 5.4, p < 0.001). Similar differences were found when

comparing the GC content of the DNA template; significantly higher GC proportions were

present in templates that were culled from the panel (mean = 47.8%) than templates that were

retained (mean = 45.5%, df = 359, t = 3.8, p < 0.001). Additionally, a total of 88 primer pairs in

the original panel contained at least one degenerate nucleotide, 72 (81%) of which were in the

forward primer. After optimization, 56 of the initial 88 (64%) were retained. In comparison, of

the 512 initial primer pairs that did not have degenerate primers, 380 (74%) were retained. The

average $F_{ST}$ for the most informative SNP at a locus and the average $H_{E\_mhap}$ did not change

appreciably between the initial and fully optimized panels (Table 4).

*Methodological modifications and performance analysis*

Significant differences for on-target read counts and the proportion of total reads that

were on-target were detected among genomic DNA extraction and purification method

combinations. Subsequent analysis using Tukey's HSD revealed that Chelex-extracted DNAs

produced the highest on-target read count, and Qiagen-extracted DNAs with ExoSAP-

purification produced the lowest (Fig. 5, p < 0.001).  While the proportion of on-target reads did

not differ between Chelex with ExoSAP and Qiagen with ExoSAP, both methods produced a

significantly lower proportion of on-target reads than the Chelex-only library (Fig. 5, p < 0.001).

Additionally, when comparing results from the full panel of 436 primer pairs to those obtained

using the same panel divided into two unique multiplexes of 209 and 227 primer pairs (n = 436)

and repooled prior to barcoding, no significant differences were found in total primer reads (df =

860, t = 0.10, p = 0.92), on-target reads (df = 858, t = 0.16, p = 0.87), or the proportion of total

reads that were on target (df = 806, t = 0.66, p = 0.51).

A total of 4,063 genotypes across 406 loci (820 SNPs) could be used in comparisons

between GT-seq data and those obtained from the original RAD study. Of these genotypes,

96.6% of calls were identical between methods, and modeled expectations of genotype

concordance (residual sum of squares = 0.02) indicated that a concordance rate of 99.0% could

be expected at a GT-seq read depth of 31 (Fig. 6).

For a final proof of concept, a new sample of 570 walleye was sequenced using the

current panel of 436 loci. After filtering, 551 individuals and 303 loci were retained with an

average of 32.9 (SD = 29.1) reads per locus; 116 of the 303 loci exhibited an average coverage

greater than the 31× target identified for 99.0% genotyping concordance (Fig. 7). The average

percent of missing data was 6.4% (SD = 13.0%) across individuals and 30.0% (SD = 38.0%)

across loci.

**Discussion**

GT-seq and other amplicon sequencing methods have tremendous potential for

facilitating high-throughput genotyping in non-model organisms (Meek & Larson, 2019). The

general steps for GT-seq panel development: SNP ascertainment, SNP selection, primer design,

and panel optimization have been previously detailed (Baetscher et al., 2018; McKinney et al.,

2019; Schmidt et al., 2019); however, the process of GT-seq panel development is not static.

Here, we leverage our experiences developing a GT-seq panel for walleye with testing various

aspects of the GT-seq methodological process to provide additional guidelines usable by other

researchers to simplify panel construction and validation, particularly in non-model species. Our

walleye panel has the necessary power to conduct GSI in a study system with highly variable

degrees of genetic differentiation and perturbation by historical stocking, while also being

capable of identifying PO and FS relationships within large populations. The robust performance

of our panel was facilitated by exploring the upper limits of how many loci a GT-seq panel can

target and the trade-offs between choosing loci for GSI versus parentage analysis. The

information presented here will aid in the efficient creation of multipurpose GT-seq panels in

organisms with little to no available genomic resources.

*Patterns of population structure: historical stocking influences GSI accuracy*

The largest genetic differentiation in our data was observed between populations from

Wisconsin and Minnesota; this structure was likely the result of recolonization from different

refugia following the Wisconsin glaciation, which ended ~10,000 years ago. A range-wide

analysis of walleye genetic structure using microsatellite loci produced similar patterns, with the

most genetically independent populations found in northern Minnesota and Canada (Stepien et

al., 2009). Additionally, we found that while populations in Minnesota displayed strong isolation

on relatively small spatial scales, broad-scale patterns of isolation were less evident in

Wisconsin. In particular, the Ceded Territory of Wisconsin, which included our Chippewa River

and Wisconsin River reporting groups, displayed patchy and low genetic structure overall. It is

likely that structure in this region has been compromised by stocking. Hammen and Sloss (2019),

for instance, observed that several populations of walleye in the upper Chippewa were more

genetically similar to populations in the upper Wisconsin than to other populations in the upper

Chippewa, while nongame species in the Ceded Territory of Wisconsin displayed patterns of

genetic divergence strictly associated with drainage basin boundaries (Westbrook, 2012). We

also observed that four proximate populations spanning the Chippewa and Wisconsin River

boundaries were nearly indistinguishable (Turtle Flambeau Flowage, Manitowish Lake,

Kawaguesaga Lake, Big Arbor Vitae Lake). These populations are within 50 km of each other

and are located near a state walleye hatchery in Woodruff, Wisconsin, that has historically used

broodstock solely from the Wisconsin River drainage basin. It is therefore highly likely that the

genetic similarity of these four populations is due to stocking. Several of the sampled

populations from Minnesota also had poorly documented stocking histories yet they remained

highly distinct. Genetic structure in Minnesota may have been less eroded if local, genetically

similar sources were used, stocking was into larger, healthier resident populations, or stocking

was less intense or ended a longer time ago.

Despite the challenges posed by low $F_{ST}$ and evidence of supplemental stocking altering

genetic structure in some populations, the SNPs discovered here provide greatly increased

resolution for defining reporting units across the Midwestern, USA. Additionally, simulations

suggested that a panel of several hundred loci would be highly capable of conducting individual-

based GSI for most genetic units in the region. Given the regional complexity, however,

improvements to accuracy could be made by further sampling areas that have shown

heterogeneous signals of genetic structure (e.g., due to stocking). For example, increased

sampling effort directed at the Chippewa and Wisconsin Rivers' drainage basins could prove

especially beneficial as analyzing populations in the lower reaches of each basin may provide a

better understanding of signals of historical recolonization, while populations in the upper

reaches (e.g., Ceded Territory of Wisconsin) could better define the effects stocking may have

had. Additional samples could also serve as a holdout dataset, as suggested by Anderson (2010),

to test the assignment accuracy of our panel.

*Tradeoffs associated with choosing loci based on differentiation versus diversity*

We evaluated the tradeoffs associated with selecting SNPs based on differentiation or

diversity and found that there was relatively little variation in GSI accuracies across panels.

Markers selected based on differentiation have been shown to provide increased resolution for

defining reporting groups in systems with low levels of genetic structure (Larson et al., 2014;

McKinney et al., 2019). This approach has not, however, been applied to systems where stocking

may be a major factor for reduced levels of population structure, such as in upper Midwestern,

USA, walleye. Interestingly, we found that assignment accuracies with our smaller panels was

relatively similar to accuracies obtained using ~30,000 SNPs discovered with RAD-seq (data not

shown). This suggests that assignment accuracy in our system may be limited more by biological

realities associated with human-mediated gene flow than by the power of our genetic markers.

Further increases in assignment accuracy are therefore likely to be realized through sampling of

additional populations and a more refined understanding of population history as opposed to

genotyping additional markers.

Conversely, we found that FPRs for assigning kin relationships were highly variable

among panels, with the microhaplotype diversity-based panel displaying the lowest FPRs by

several orders of magnitude for each kin relationship (Table 3). This contrast in inter-panel

variation between GSI and kinship simulations is reflective of the variation in information

content of each panel (Fig. S1), and supports previous findings that while microhaplotype

information provided added benefit to both applications, the greatest increase in assignment

accuracy will likely be for kinship analysis (Baetscher et al., 2018; McKinney, Seeb, et al.,

2017). When attempting to target microhaplotype loci via GT-seq, attention should be given to

the number of SNPs one aims to genotype within a locus, as attempting to include loci with too

many SNPs may result in targeting repetitive regions that fail to amplify properly in a multiplex.

The expected maximum number of alleles per locus and the degree to which loci with large

numbers of alleles perturbs primer design will likely vary among taxa. We chose a cutoff of 10

alleles per locus as this appeared to be a natural break point in the allele distribution for walleye;

we suggest that researchers investigate this in their system and come up with a logical cutoff

prior to selecting loci. Finally, while our results suggested this panel could facilitate HS

identification in small systems, performing this task in large systems would likely require more

loci. Our tests of panel implementation suggest this could be achievable by combining PCR

products from several panels within individuals prior to barcoding.

*Optimizing primer design and removing overamplifying loci*

The main objective of GT-seq primer development is to produce a single pool of primer

pairs that will amplify uniformly, while retaining as many loci as possible. To achieve this, it is

important to minimize heterogeneity of primer and product characteristics (e.g., primer size,

product size) and to understand that the highly multiplexed PCR required by GT-seq can be

complicated by hairpin- and inter-primer hybridization artifacts. To best control PCR artifacts, it

is important to avoid developing primers with complimentary regions (e.g., complimentary 3'

regions and self-complementarity) and apply conservative thresholds to the upper $T_m$ of primer

design parameters (Rychlik, 1993). Incorporating loci with multiple SNPs can lead to further

difficulties when the ideal priming region also contains a SNP. We found that, while degenerate

primers could be successfully amplified in a multiplex, they were culled during optimization at a

higher rate than non-degenerate primers. Further performance benefits could be gained from

examining DNA template quality beyond just the availability of priming regions, as shown by

Benita et al. (2003) who found regionalized GC content of template DNA to be a predictor of

PCR success. This was supported by our data, as loci removed from the panel during

optimization displayed significantly higher GC content in the amplicon and primer. Finally,

while GT-seq primers can theoretically be designed for a range of amplicon sizes, we suggest

that researchers design panels targeting similarly sized products that can be sequenced using

PE150 technology. Panels containing similarly sized and relatively short amplicons should

reduce variation in amplification rates (Baetscher et al., 2018) and ensure that genotyping is

robust to variation in sample quality. Moreover, PE150 sequencing is common to benchtop and

core facility sequencing platforms, such as Illumina® MiSeq and HiSeq.

In exploring the upper limits of how many loci a GT-seq panel can target, we found that

the number of amplicons reliably genotyped in a single pool is highly dependent on variable

rates of amplification among primer pairs during PCR and, to a lesser extent, the degree of

primer specificity. Despite efforts to limit primer inter-hybridization through diligent primer

design, the presence of overamplifying loci is likely inevitable during early phases of panel

development (see also McKinney et al. 2019). We found it best to focus primarily on the

uniformity of amplification within the primer pool in early optimization steps, by removing

primer pairs found to overamplify. Although achieving perfect uniformity is challenging,

application of strict cutoffs during initial optimization steps likely results in a final panel that is

less influenced by overamplification, thereby increasing the upper limit of GT-seq performance.

The importance of this was illustrated by prop_reads_T10 reducing from 85.0% of all primer

reads to 36.6% after optimization. Likewise, on-target rates were greatly improved by addressing

overamplification, as demonstrated by the on-target proportion of reads increasing from 24.9% to

91.7% by the third test.

*Further optimization of the GT-seq protocol*

Although there may be an upper as-yet-unidentified limit in the number of primers that

can be included in a single primer pool, we found that the total number of loci targeted can be

increased by PCR amplifying multiple primer pools separately on a sample and pooling PCR

products within individuals prior to barcoding. This approach could be used to genotype multiple

complementary or even independent GT-seq panels using the same primer tail systems at a small

cost increase compared to genotyping a single panel, as the most expensive steps in the GT-seq

protocol (e.g., DNA normalization) are only conducted once (Campbell et al., 2015). Combining

multiple panels could facilitate genotyping of > 1,000 loci rather than a few hundred, providing

greatly increased power for kinship analysis and GSI (Baetscher et al., 2018; McKinney, Seeb, et

al., 2017). Additionally, further optimization of individual panels could be conducted by

manipulating the initial concentrations of primer pairs based on observed panel performance,

reducing the concentration of loci that appear to overamplify. While this process would be

cumbersome to perform by hand, a liquid handling robot could enable a researcher to fine-tune

the performance of existing and new panels alike, thereby enhancing efficiency.

DNA extraction can comprise a large portion of the total cost of genetic analysis,

especially for relatively affordable approaches such as GT-seq, in terms of finances and time.

Extractions using chelating beads provided a cost-effective alternative to more expensive salting-

out approaches, such as Qiagen DNeasy kits. Chelating extractions, however, can also produce

lower quality DNA and may include suspended impurities (Singh et al., 2018). Campbell et al.

(2015) did show that GT-seq can be conducted using DNA from chelating extractions but did not

directly compare results using multiple extraction protocols. Here, we directly showed that cost-

effective chelating extractions can produce equally high quality, if not superior, sequence data

compared to more expensive methods. Although consideration should be given to the quality of

tissue samples, the chelating approach appears to be a viable approach for reducing per-sample

costs with GT-seq. It is important to be aware that proper laboratory technique is essential when

using this method, however, as chelating beads will inhibit PCR and greatly reduce library

product yields. This may be especially problematic when using a liquid handling robot that is

unable to visually detect chelating beads. Therefore, we suggest researchers carefully pipette the

DNA-containing supernatant from chelating resin extractions by hand into a secondary container

(e.g., 96-well PCR plate) before aliquoting DNA with a robot. Finally, we found that the

ExoSAP procedure included in the original GT-seq protocol did not produce higher quality data

and was not necessary for our purposes; removing this step from the protocol will further reduce

GT-seq costs and time commitment.

*Suggestions for designing GT-seq studies and conclusions*

A major consideration when designing a GT-seq panel is deciding how large of an

ascertainment dataset is necessary. We constructed a comprehensive ascertainment set with

RAD-seq, which was expensive and resource intensive. Despite this, we found that the panel

chosen based on diversity produced similar results to the panel chosen based on differentiation.

In our case, we believe that a smaller ascertainment set of ~96 individuals sampled from across

the same geographic range may have resulted in a panel of relatively similar quality. Smaller

ascertainment datasets are likely sufficient when the main applications of a given GT-seq panel

are kinship analysis and GSI of highly diverged populations; however, when designing GT-seq

panels to differentiate closely related populations (e.g. Chinook salmon *Oncorhynchus*

*tshawytscha* in western Alaska), accurate characterization of ascertainment populations is vital

(Larson et al., 2014; McKinney et al., 2019).

Another major consideration when conducting GT-seq analysis is deciding how deep to

sequence individuals. We found that a read depth of 31× could be expected to produce genotypes

that were 99% concordant with those derived from RADseq. Read depths were, however, highly

variable across loci; we only retained 303 of the 436 loci in our panel when we genotyped 536

individuals at an average depth of 33×. We also found that a large and variable proportion of

reads can be discarded prior to genotyping. Therefore, we suggest that researchers target an

average depth of at least 100× to ensure that most loci in the panel can be genotyped and that all

acquired genotypes are highly reliable. At this level of coverage, researchers could genotype

~500 individuals with a panel of 500 loci on a single MiSeq lane (~25 million reads) and ~8,000

individuals on a HiSeq lane (~400 million reads). It is possible this level of coverage is not

necessary for some applications, such as GSI, but we strongly suggest obtaining high coverage

for more sensitive applications that require high genotyping accuracy, such as kinship analysis.

Finally, researchers conducting GT-seq must consider trade-offs associated with different

genotyping approaches. The two main approaches we are aware of are: (1) in-silico probe-based

methods that use pattern matching to genotype specific alleles (Campbell et al., 2015; McKinney

et al., 2019) and (2) alignment-based methods that call all polymorphisms in a given amplicon

(Baetscher et al., 2019). A major advantage of probe-based methods is that databases of probes

can be shared among laboratories, facilitating standardization. It is difficult, however, to discover

new variation with these methods, whereas alignment-based methods discover new variation by

default. We suggest a hybrid approach, where researchers periodically use alignment-based

approaches to discover new variation and add this variation to a probe database that forms the

basis of genotyping and standardizing genotyping among laboratories.

GT-seq is a powerful addition to the molecular ecologist's toolkit that facilitates rapid,

accurate, and cost-effective genetic analysis. Yet, creating a GT-seq panel is non-trivial, and

there are many considerations for maximizing the utility of this approach. We found that the

greatest challenge when designing our GT-seq panel was locus-specific overamplification, and

we suggest that researchers remove these loci liberally. We also found that chelating extractions

without an ExoSAP step produce high-quality results, providing a lower-cost alternative to

salting-out extractions. Additionally, we showed that combining multiplex PCR products from

multiple panels prior to barcoding can ensure additional, potentially important, loci can be

genotyped with only a moderate cost increase. Finally, we found that a relatively substantial

proportion of sequencing reads are lost before genotyping, and we suggest researchers target

higher sequencing coverage (100×) than may apparently be necessary to ensure that GT-seq

datasets are robust across loci. The GT-seq approach promises to be a mainstay of population

genetics for the foreseeable future, and the guidelines and suggestions outlined here may help

increase the effective use of this powerful method.

**Acknowledgements**

We thank field crews from the Wisconsin and Minnesota Departments of Natural Resources for

collecting tissue samples. We also thank Keith Turnquist from UW-Stevens Point for assistance

with laboratory analysis and Ana Ramón-Laca for advice regarding primer development.  This

study was funded by the United States Fish and Wildlife Service, Federal Aid in Sportfish

Restoration program and the Wisconsin Department of Natural Resources. Any use of trade,

firm, or product names is for descriptive purposes only and does not imply endorsement by the

U.S. Government.

**References**

Ackerman, M. W., Habicht, C., & Seeb, L. W. (2011). Single-Nucleotide Polymorphisms (SNPs)
under Diversifying Selection Provide Increased Accuracy and Precision in Mixed-Stock
Analyses of Sockeye Salmon from the Copper River, Alaska. *Transactions of the*
*American Fisheries Society, 140*(3), 865-881. doi:10.1080/00028487.2011.588137

Ali, O. A., O'Rourke, S. M., Amish, S. J., Meek, M. H., Luikart, G., Jeffres, C., & Miller, M. R.
(2016). RAD Capture (Rapture): Flexible and Efficient Sequence-Based Genotyping.
*Genetics, 202*(2), 389-400. doi:10.1534/genetics.115.183665

Anderson, E. C. (2010). Assessing the power of informative subsets of loci for population
assignment: standard methods are upwardly biased. *Molecular Ecology Resources, 10*(4),
701-710. doi:10.1111/j.1755-0998.2010.02846.x

Anderson, E. C., & Garza, J. C. (2006). The power of single-nucleotide polymorphisms for
large-scale parentage inference. *Genetics, 172*(4), 2567-2582.
doi:10.1534/genetics.105.048074

Baetscher, D. S., Anderson, E. C., Gilbert-Horvath, E. A., Malone, D. P., Saarman, E. T., Carr,
M. H., & Garza, J. C. (2019). Dispersal of a nearshore marine fish connects marine
reserves and adjacent fished areas along an open coast. *Molecular Ecology, 28*(7), 1611-
1623. doi:10.1111/mec.15044

Baetscher, D. S., Clemento, A. J., Ng, T. C., Anderson, E. C., & Garza, J. C. (2018).
Microhaplotypes provide increased power from short-read DNA sequences for
relationship inference. *Mol Ecol Resour, 18*(2), 296-305. doi:10.1111/1755-0998.12737

Bay, R. A., Taylor, E. B., & Schluter, D. (2019). Parallel introgression and selection on
introduced alleles in a native species. *Molecular Ecology, 28*(11), 2802-2813.
doi:10.1111/mec.15097

Benita, Y., Oosting, R. S., Lok, M. C., Wise, M. J., & Humphery-Smith, I. (2003). Regionalized
GC content of template DNA as a predictor of PCR success. *Nucleic Acids Research,*
*31*(16), e99-e99. doi:10.1093/nar/gng101

Bradbury, I. R., Hubert, S., Higgins, B., Sharen, B., Paterson, I. G., Snelgrove, P. V. R., . . .
Bentzen, P. (2011). Evaluating SNP ascertainment bias and its impact on population
assignment in Atlantic cod, Gadus morhua. *Molecular Ecology Resources, 11*(s1), 218-
225. doi:10.1111/j.1755-0998.2010.02949.x

Brumfield, R. T., Beerli, P., Nickerson, D. A., & Edwards, S. V. (2003). The utility of single
nucleotide polymorphisms in inferences of population history. *Trends in Ecology &*
*Evolution, 18*(5), 249-256. doi:https://doi.org/10.1016/S0169-5347(03)00018-1

Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by
sequencing (GT-seq): A cost effective SNP genotyping method based on custom
amplicon sequencing. *Molecular Ecology Resources, 15*(4), 855-867. doi:10.1111/1755-
0998.12357

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an
analysis tool set for population genomics. *Molecular Ecology, 22*(11), 3124-3140.
doi:10.1111/mec.12354

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., . . . Group, G. P.
A. (2011). The variant call format and VCFtools. *Bioinformatics, 27*(15), 2156-2158.
doi:10.1093/bioinformatics/btr330

Davey, J. W., Hohenlohe, P. A., Etter, P. D., Boone, J. Q., Catchen, J. M., & Blaxter, M. L.
(2011). Genome-wide genetic marker discovery and genotyping using next-generation
sequencing. *Nature Reviews Genetics, 12*(7), 499-510. doi:10.1038/nrg3012

Embke, H. S., Rypel, A. L., Carpenter, S. R., Sass, G. G., Ogle, D., Cichosz, T., . . . Vander
    Zanden, M. J. (2019). Production dynamics reveal hidden overharvest of inland
    recreational fisheries. *Proceedings of the National Academy of Sciences, 116*(49), 24676.
    doi:10.1073/pnas.1913196116

Garza, J. C., & Freimer, N. B. (1996). Homoplasy for size at microsatellite loci in humans and
    chimpanzees. *Genome Research, 6*(3), 211-217. doi:10.1101/gr.6.3.211

Hammen, J. J., & Sloss, B. L. (2019). Walleye Genetic Characterization in the Northern Ceded
    Territory of Wisconsin: Implications for Stocking Using Conservation Strategies. *North
    American Journal of Fisheries Management, 0*(0). doi:10.1002/nafm.10302

Hansen, J., Sass, G., Gaeta, J., Hansen, G., Isermann, D., Lyons, J., . . . Carpenter, S. (2015).
    Largemouth Bass Management in Wisconsin: Intra- and Inter-Specific Implications of
    Abundance Increases.

Hoffman, J. I., Simpson, F., David, P., Rijks, J. M., Kuiken, T., Thorne, M. A. S., . . .
    Dasmahapatra, K. K. (2014). High-throughput sequencing reveals inbreeding depression
    in a natural population. *Proceedings of the National Academy of Sciences, 111*(10), 3775.
    doi:10.1073/pnas.1318945111

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide
    SNP data. *Bioinformatics, 27*(21), 3070-3071. doi:10.1093/bioinformatics/btr521

Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W., & Prodöhl, P. A. (2013). diveRsity: An
    R package for the estimation and exploration of population genetics parameters and their
    associated errors. *Methods in Ecology and Evolution, 4*(8), 782-788. doi:10.1111/2041-
    210X.12067

Larson, W. A., Seeb, J. E., Pascal, C. E., Templin, W. D., Seeb, L. W., & Taylor, E. (2014).
    Single-nucleotide polymorphisms (SNPs) identified through genotyping-by-sequencing
    improve genetic stock identification of Chinook salmon ( Oncorhynchus tshawytscha)
    from western Alaska. *Canadian Journal of Fisheries & Aquatic Sciences, 71*(5), 698-708.
    doi:10.1139/cjfas-2013-0502

Logsdon, D. E., Anderson, C. S., & Miller, L. M. (2016). Contribution and Performance of
    Stocked Walleyes in the Recovery of the Red Lakes, Minnesota, Fishery. *North
    American Journal of Fisheries Management, 36*(4), 828-843.
    doi:10.1080/02755947.2016.1167143

Luikart, G., & England, P. R. (1999). Statistical analysis of microsatellite DNA data. *Trends in
    Ecology & Evolution, 14*(7), 253-256. doi:10.1016/S0169-5347(99)01632-8

McKinney, G. J., Pascal, C. E., Templin, W. D., Gilk-Baumer, S. E., Dann, T. H., Seeb, L. W., &
    Seeb, J. E. (2019). Dense SNP panels resolve closely related Chinook salmon
    populations. *Canadian Journal of Fisheries and Aquatic Sciences*. doi:10.1139/cjfas-
    2019-0067

McKinney, G. J., Seeb, J. E., & Seeb, L. W. (2017). Managing mixed-stock fisheries: genotyping
    multi-SNP haplotypes increases power for genetic stock identification. *Canadian Journal
    of Fisheries and Aquatic Sciences, 74*(4), 429-434. doi:10.1139/cjfas-2016-0443

McKinney, G. J., Waples, R. K., Pascal, C. E., Seeb, L. W., & Seeb, J. E. (2018). Resolving
    allele dosage in duplicated loci using genotyping-by-sequencing data: A path forward for
    population genetic analysis. *Molecular Ecology Resources, 18*(3), 570-579.
    doi:10.1111/1755-0998.12763

McKinney, G. J., Waples, R. K., Seeb, L. W., & Seeb, J. E. (2017). Paralogs are revealed by
    proportion of heterozygotes and deviations in read ratios in genotyping-by-sequencing

data from natural populations. *Mol Ecol Resour, 17*(4), 656-669. doi:10.1111/1755-
0998.12613

Meek, M. H., & Larson, W. A. (2019). The future is now: Amplicon sequencing and sequence
capture usher in the conservation genomics era. *Molecular Ecology Resources, 19*(4),
795-803. doi:10.1111/1755-0998.12998

Moran, B. M., & Anderson, E. C. (2018). Bayesian inference from the conditional genetic stock
identification model. *Canadian Journal of Fisheries & Aquatic Sciences, 76*(4), 551-560.
doi:10.1139/cjfas-2018-0016

Morin, P. A., Luikart, G., Wayne, R. K., & the, S. N. P. w. g. (2004). SNPs in ecology, evolution
and conservation. *Trends in Ecology & Evolution, 19*(4), 208-216.
doi:https://doi.org/10.1016/j.tree.2004.01.009

Narum, S. R., Buerkle, C. A., Davey, J. W., Miller, M. R., & Hohenlohe, P. A. (2013).
Genotyping-by-sequencing in ecological and conservation genomics. *Mol Ecol, 22*(11),
2841-2847. doi:10.1111/mec.12350

Natesh, M., Taylor, R. W., Truelove, N. K., Hadly, E. A., Palumbi, S. R., Petrov, D. A., &
Ramakrishnan, U. (2019). Empowering conservation practice with efficient and
economical genotyping from poor quality samples. *Methods in Ecology and Evolution,*
*10*(6), 853-859. doi:10.1111/2041-210X.13173

Navajas, M. J., Thistlewood, H. M. A., Lagnel, J., & Hughes, C. (1998). Microsatellite
sequences are under-represented in two mite genomes. *Insect Molecular Biology, 7*(3),
249-256. doi:10.1111/j.1365-2583.1998.00066.x

Pavinato, V. A. C., Wijeratne, S., Spacht, D., Denlinger, D. L., Meulia, T., & Michel, A. P.
(2019). Leveraging targeted sequencing for non-model species: a step-by-step guide to
obtain a reduced SNP set and a pipeline to automate data processing in the Antarctic
Midge, <em>Belgica antarctica</em>. *bioRxiv*, 772384. doi:10.1101/772384

Reid, K., Carlos Garza, J., Gephard, S. R., Caccone, A., Post, D. M., & Palkovacs, E. P. (2019).
Restoration-mediated secondary contact leads to introgression of alewife ecotypes
separated by a colonial-era dam. *Evolutionary Applications, n/a*(n/a).
doi:10.1111/eva.12890

Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for
paired-end sequencing improve RADseq-based population genomics. *Molecular Ecology,*
*28*(21), 4737-4754. doi:10.1111/mec.15253

Rychlik, W. (1993). Selection of Primers for Polymerase Chain Reaction. In B. A. White (Ed.),
*PCR Protocols: Current Methods and Applications* (pp. 31-40). Totowa, NJ: Humana
Press.

Rypel, A. L., Goto, D., Sass, G. G., & Zanden, M. J. V. (2018). Eroding productivity of walleye
populations in northern Wisconsin lakes. *Canadian Journal of Fisheries and Aquatic*
*Sciences, 75*, 2291+.

Schmidt, D., Campbell, N. R., Govindarajulu, P., Larsen, K. W., & Russello, M. A. (2019).
Genotyping-in-Thousands by sequencing (GT-seq) panel development and application to
minimally-invasive DNA samples to support studies in molecular ecology. *Molecular*
*Ecology Resources, 0*(ja). doi:10.1111/1755-0998.13090

Seeb, J. E., Carvalho, G., Hauser, L., Naish, K., Roberts, S., & Seeb, L. W. (2011). Single-
nucleotide polymorphism (SNP) discovery and applications of SNP genotyping in
nonmodel organisms. *Molecular Ecology Resources, 11*, 1-8. doi:10.1111/j.1755-
0998.2010.02979.x

Singh, U. A., Kumari, M., & Iyengar, S. (2018). Method for improving the quality of genomic
DNA obtained from minute quantities of tissue and blood samples using Chelex 100
resin. *Biological Procedures Online, 20.*
Smith, C. T., Antonovich, A., Templin, W. D., Elfstrom, C. M., Narum, S. R., & Seeb, L. W.
(2007). Impacts of Marker Class Bias Relative to Locus-Specific Variability on
Population Inferences in Chinook Salmon: A Comparison of Single-Nucleotide
Polymorphisms with Short Tandem Repeats and Allozymes. *Transactions of the*
*American Fisheries Society, 136*(6), 1674-1687. doi:10.1577/T06-227.1
Stepien, C., A., Murphy, D., J., Lohner, R., N., Sepulveda-Villet O, J., & Haponski, A., E.
(2009). Signatures of vicariance, postglacial dispersal and spawning philopatry:
population genetics of the walleye Sander vitreus. *Molecular Ecology, 18*(16), 3411-
3428. doi:10.1111/j.1365-294X.2009.04291.x
Storer, C. G., Pascal, C. E., Roberts, S. B., Templin, W. D., Seeb, L. W., & Seeb, J. E. (2012).
Rank and Order: Evaluating the Performance of SNPs for Individual Assignment in a
Non-Model Organism. *PLOS ONE, 7*(11), 1-13. doi:10.1371/journal.pone.0049018
Wagner, C. E., Keller, I., Wittwer, S., Selz, O. M., Mwaiko, S., Greuter, L., . . . Seehausen, O.
(2013). Genome-wide RAD sequence data provide unprecedented resolution of species
boundaries and relationships in the Lake Victoria cichlid adaptive radiation. *Molecular*
*Ecology, 22*(3), 787-798. doi:10.1111/mec.12023
Westbrook, L. J. (2012). *Genetic structure of rock bass and johnny darters: Implications for*
*gamefish management in Wisconsin.* (Master's), University of Wisconsin, Stevens Point,
Willi, Y., van Kleunen, M., Dietrich, S., & Fischer, M. (2007). Genetic rescue persists beyond
first-generation outbreeding in small populations of a rare plant. *Proceedings of the Royal*
*Society B: Biological Sciences, 274*(1623), 2357-2364. doi:10.1098/rspb.2007.0768

## Data accessibility

876 Raw data for the RADseq and GT-seq data obtained in this study was deposited to the NCBI
877 sequence read archive (SUB####) and VCF files of genotypes are available on DRYAD (DOI:
878 PENDING). Python and R scripts used in the statistical analysis pipeline are available at GIT

## Author contributions

880 WL, GS, KG, and LM designed the study with input from MB. Data analyses were conducted by
881 MB with assistance from GM. Laboratory analysis was conducted by MB, KG, and LS. All
882 authors contributed to the writing of the manuscript.

## Tables

**Table 1.** Information on walleye *Sander vitreus* collections from 23 sites in Wisconsin and Minnesota. Reporting units are aggregations of genetically similar populations grouped for GSI analysis, n past filters is the number of individuals missing genotypes at < 30% of SNPs and retained after quality filtering. Diversity statistics calculated using 20,579 SNPs. The $F_{ST\_600}$, Composite$_{600}$, and Diversity$_{600}$ columns are the percent correct assignment to reporting group for each population with 100% simulations conducted using the corresponding panel.

| Population ID | Reporting Unit | Population | Latitude | Longitude | n sampled | n past filters | $H_E$ | $H_O$ | $F_{IS}$ | AR | $F_{ST\_600}$ | Composite$_{600}$ | Diversity$_{600}$ |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | Rock-Fox | Delavan Lake | 42.58 | -88.63 | 48 | 48 | 0.169 | 0.168 | 0.008 | 1.607 | 1.00 | 1.00 | 1.00 |
| 2 | Wolf River | Lake Winnebago | 44.36 | -88.69 | 47 | 41 | 0.173 | 0.186 | -0.05 | 1.645 | 1.00 | 1.00 | 1.00 |
| 3 | Upper Wisconsin | Lake Wisconsin | 43.38 | -89.58 | 48 | 45 | 0.179 | 0.175 | 0.017 | 1.674 | 1.00 | 1.00 | 1.00 |
| 4 | Upper Wisconsin | Medicine Lake Chain | 45.81 | -89.13 | 47 | 47 | 0.166 | 0.166 | 0.004 | 1.604 | 0.96 | 0.98 | 0.98 |
| 5 | Upper Wisconsin | Willow Flowage | 45.71 | -89.87 | 48 | 48 | 0.176 | 0.174 | 0.013 | 1.657 | 1.00 | 1.00 | 0.99 |
| 6 | Upper Wisconsin | Kawaguesaga Lake | 45.86 | -89.74 | 48 | 42 | 0.17 | 0.167 | 0.013 | 1.638 | 0.96 | 0.94 | 0.94 |
| 7 | Upper Wisconsin | Big Arbor Vitae Lake | 45.93 | -89.65 | 48 | 44 | 0.174 | 0.174 | 0.005 | 1.654 | 0.74 | 0.96 | 0.99 |
| 8 | Upper Chippewa | Escanaba Lake | 46.06 | -89.59 | 48 | 44 | 0.168 | 0.173 | -0.018 | 1.623 | NA | NA | NA |
| 9 | Upper Chippewa | Sanford Lake | 46.18 | -89.69 | 48 | 44 | 0.157 | 0.164 | -0.033 | 1.528 | NA | NA | NA |
| 10 | Upper Chippewa | Manitowish Lake | 46.11 | -89.85 | 47 | 35 | 0.172 | 0.175 | -0.006 | 1.647 | 0.58 | 0.57 | 0.51 |
| 11 | Upper Chippewa | Turtle Flambeau Flowage | 46.06 | -90.13 | 47 | 38 | 0.173 | 0.172 | 0.005 | 1.661 | 0.63 | 0.55 | 0.76 |
| 12 | Upper Chippewa | Chippewa Flowage | 45.90 | -91.09 | 47 | 43 | 0.173 | 0.175 | -0.006 | 1.658 | 0.88 | 0.89 | 0.93 |
| 13 | Upper Chippewa | Eau Claire River | 44.80 | -91.50 | 47 | 47 | 0.161 | 0.162 | -0.001 | 1.583 | 0.98 | 0.98 | 0.98 |
| 14 | Upper Chippewa | Lake Millicent | 46.53 | -91.37 | 48 | 32 | 0.167 | 0.176 | -0.034 | 1.623 | NA | NA | NA |
| 15 | Lake Superior | St. Louis River | 46.65 | -92.21 | 32 | 30 | 0.17 | 0.168 | 0.006 | 1.621 | 0.77 | 0.77 | 0.77 |
| 16 | Vermilion River | Pike River | 47.59 | -92.39 | 32 | 28 | 0.144 | 0.142 | 0.005 | 1.498 | 1.00 | 1.00 | 1.00 |
| 17 | Des Moines River | Lake Sarah | 44.15 | -95.77 | 32 | 30 | 0.164 | 0.166 | -0.006 | 1.597 | 1.00 | 1.00 | 1.00 |
| 18 | North Fork Crow River | Lake Koronis | 45.33 | -94.70 | 32 | 17 | 0.155 | 0.155 | -0.011 | 1.579 | 0.82 | 0.82 | 0.75 |
| 19 | Rum River | Mille Lacs Lake | 46.25 | -93.67 | 32 | 29 | 0.148 | 0.151 | -0.018 | 1.511 | 1.00 | 1.00 | 1.00 |
| 20 | Pine River | Pine River | 46.70 | -94.39 | 32 | 30 | 0.156 | 0.162 | -0.028 | 1.547 | 0.97 | 0.97 | 0.97 |
| 21 | Mississippi River - Headwaters | Cutfoot Sioux Lake | 47.50 | -94.09 | 32 | 25 | 0.147 | 0.148 | -0.011 | 1.517 | 1.00 | 1.00 | 1.00 |
| 22 | Otter Tail River | Ottertail Lake | 46.41 | -95.66 | 32 | 23 | 0.158 | 0.16 | -0.016 | 1.568 | 1.00 | 1.00 | 0.97 |
| 23 | Red Lake | Red Lake | 47.91 | -95.04 | 32 | 29 | 0.149 | 0.153 | -0.025 | 1.514 | 0.90 | 0.86 | 0.83 |

**Table 2.** Summary of pairwise $F_{ST}$ comparisons between walleye *Sander vitreus* populations grouped by state of origin. Abbreviations are Wisconsin (WI) and Minnesota (MN).

|  | WI-WI | MN-MN | WI-MN |
|---|---|---|---|
| Max | 0.106 | 0.142 | 0.142 |
| Mean | 0.032 | 0.068 | 0.072 |
| Min | 0.001 | 0.019 | 0.026 |

898 **Table 3.** Summary statistics by SNP panel tested for walleye *Sander vitreus* in Wisconsin and
899 Minnesota, USA, including: average $F_{ST}$, heterozygosity ($H_{E\_mhap}$), assignment accuracy to
900 population and reporting unit of origin in 100% simulations, and estimated false-positive rates
901 (FPR) for a given kin relationship at a false-negative rate (FNR) of 0.01.

| | $F_{ST\_600}$ | Composite$_{600}$ | Diversity$_{600}$ |
|---|---|---|---|
| Average $F_{ST}$ | 0.117 | 0.076 | 0.047 |
| Average $H_{E\_mhap}$ | 0.389 | 0.569 | 0.633 |
| Average accuracy by reporting unit | 0.937 | 0.937 | 0.929 |
| Average accuracy by population | 0.864 | 0.861 | 0.862 |
| Parent-offspring FPR (FNR = 0.01) | $4.68 \times 10^{-34}$ | $7.92 \times 10^{-62}$ | $2.74 \times 10^{-80}$ |
| Full-sibling FPR (FNR = 0.01) | $3.42 \times 10^{-29}$ | $5.34 \times 10^{-50}$ | $1.16 \times 10^{-64}$ |
| Half-sibling FPR (FNR = 0.01) | $6.44 \times 10^{-6}$ | $2.56 \times 10^{-10}$ | $2.06 \times 10^{-13}$ |

**Table 4.** Summary of GT-seq optimization runs for walleye *Sander vitreus* in Wisconsin and Minnesota, USA. Rows report number of primer pairs targeted, number of reads with intact i-7 barcodes (retained reads), number of retained reads with *in-silico* primer sequences (total reads), number of total reads with *in-silico* probe sequences (on-target reads), percent of total reads on-target, percent of total reads allocated to the 10% of loci tested with highest rank total read counts, average number of SNPs per locus, and average GC content in the forward and reverse primers.

|  | Test 1 | Test 2 | Test 3 |
|---|---|---|---|
| Total primer pairs | 600 | 477 | 436 |
| i7 reads | 4,655,071 | 12,653,262 | 7,282,101 |
| i7 reads w/ primers (total reads) | 4,150,910 | 9,347,591 | 6,827,424 |
| i7 reads w/ primers & probes (on- target) | 1,031,707 | 3,268,293 | 6,262,523 |
| On-target percent of total reads | 24.9% | 35.0% | 91.7% |
| Percent reads in top 10% of loci | 85.0% | 72.5% | 36.6% |
| mean SNPs per locus | 2.06 | 2.00 | 1.97 |
| mean GC percent forward primer | 51.0% | 50.4% | 50.3% |
| mean GC percent reverse primer | 49.0% | 48.3% | 48.2% |
| mean $F_{ST}$ | 0.133 | 0.133 | 0.133 |
| mean $H_{E\_mhap}$ | 0.425 | 0.415 | 0.416 |

914 **Figures**

916 **Figure 1.** Generalized workflow describing major steps inherent to *de novo* construction of a
917 high-density SNP panel for walleye *Sander vitreus* in Wisconsin and Minnesota, USA. Numbers
918 of SNPs or loci present in each phase for this panel shown in parentheses.

**Figure 2.** (a) Map of walleye *Sander vitreus* in Wisconsin (populations 1-14), the St. Louis River (population 15), and Minnesota (populations 16-23), USA, collection locations and (b) dendrogram of sampled populations with bootstrap support (n = 1000) estimates above nodes. Branch lengths correspond to genetic distances estimated using Nei's $D_A$. Figures color coded according to major drainage of origin (Hudson Bay: yellow, Mississippi: green, Great Lakes: blue) and numbered with respect to order in Table 1.

**Figure 3.** (a) Violin plots showing densitity distributions of accuracy estimates from 100%
simulations of 23 populations of walleye *Sander vitreus* in Wisconsin and Minnesota, USA,
performed using 1,000 iterations for each test panel by reporting unit and (b) simulated false-
positive rate (FPR) estimates across a range of false-negative rates (FNR). Figures color coded
according to SNP panel tested: $F_{ST\_600}$ (red, 600 rank $F_{ST}$ loci), Composite$_{600}$ (black, 300 rank
$F_{ST}$ and 300 rank $H_{E\_mhap}$ loci), and Diversity$_{600}$ (purple, 600 rank $H_{E\_mhap}$ loci).

**Figure 4.** Relative $\log_{10}$ total read counts per locus (black) and relative $\log_{10}$ on-target read
counts per locus (green) of the GT-seq panel for walleye *Sander vitreus* in Wisconsin and
Minnesota, USA,  prior to optimization (a, 600 loci), after first optimization (b, 477 loci), and
after second optimization (c, 436 loci). Loci identified for culling during optimization steps
shown in orange. Raw read counts annotated in boxes.

**Figure 5.** Number of on-target reads (green) and proportion of total reads on-target obtained
from GT-seq libraries produced using DNAs extracted via Chelex, Chelex with Exo-SAP, and
Qiagen with Exo-SAP. Significantly different groups denoted by letters on box.

**Figure 6.** Modeled relationship between GT-seq read depth and genotype concordance between GT-seq and RADseq shown in gray (1.00-0.34/GT-seq read depth, rss = 0.02) with 95% confidence intervals in red. GT-seq read depth at which estimated genotype concordance equals 99% (96.2%-100%) represented by blue line. Black points display proportion of genotypes found identical between GT-seq and RADseq for GT-seq read depth bins with > 30 genotypes.

**Figure 7.** Variation in read depth among individuals at loci successfully genotyped after quality filtering (303 loci with < 30% missing data). Average read depth at each locus shown with black points, while gray points denote first and third quartile for each locus. Dotted blue line denotes target read depth of 30×. Data from 551 walleye sequenced using fully optimized panel. Average read depth among all loci is 33×.

**Supplementary materials**

**Table S1.** Pairwise $F_{ST}$ estimates for all sampled walleye *Sander vitreus* populations (sites numbered according to Table 1 and Fig. 1 A). Estimates produced in arlequin v3.5.2.

**Table S2.** Summary statistics for 20,597 SNPs retained through initial filtering based on maximum missingness rates of < 30% and HDplot cutoffs of H > 0.5 and -7 < D < 7. Columns include a locus tag (CHROM), position of SNP within locus (Reid et al.), a unique SNP value (ID), reference (REF) and alternate (Keenan et al.) SNP alleles, global $F_{IS}$ (Willi et al.), single SNP $F_{ST}$ (Smith et al.), expected microhaplotype heterozygosity (mhap_$H_E$), and number of alleles per locus tag (n_alleles). Diversity statistics estimated in diveRsity v1.9.90 (global $F_{IS}$ and single SNP $F_{ST}$) and adegenet v2.1.1 (single locus $H_E$, number of alleles).

**Table S3.** Summary matrix of 100% simulations (reps = 1,000, mixsize = 200) for each sampled population retained through filtering, performed using the $F_{ST\_600}$ panel. Each row represents a simulation for the listed population name. Each column within a row represents the proportion of individuals assigned to the population denoted at the top of the column. Unassigned individuals (< 70% probability of origin from a given population) accounted for in last column.

**Table S4.** Summary matrix of 100% simulations (reps = 1,000, mixsize = 200) for each sampled population retained through filtering steps, performed using the Composite_$_{600}$ panel. Each row represents a simulation for the listed population name. Each column within a row represents the proportion of individuals assigned to the population denoted at the top of the column. Unassigned individuals (< 70% probability of origin from a given population) are accounted for in the last column.

**Table S5.** Summary matrix of 100% simulations (reps = 1,000, mixsize = 200) for each sampled population retained through filtering steps, performed using the Diversity_$_{600}$ panel. Each row represents a simulation for the listed population name. Each column within a row represents the proportion of individuals assigned to the population denoted at the top of the column. Unassigned individuals (< 70% probability of origin from a given population) are accounted for in the last column.

986     **Figure S1.** Frequency distribution of number of alleles among 600 loci tested in each panel.