Title: Deficits in Behavioral and Neuronal Pattern Separation in Temporal Lobe Epilepsy

Abbreviated Title: Pattern Separation and Mnemonic Discrimination in Epilepsy

Authors and Affiliations: A.D. Madar[1,2], J.A. Pfammatter[1], J. Bordenave[3], E.I. Plumley[3], S.

Ravi[1], M. Cowie[1], E.P. Wallace[1,3], B.P. Hermann[3], R.K. Maganti[3], M.V. Jones[1]

[1]Department of Neuroscience, University of Wisconsin-Madison, 53703

[2]Department of Neurobiology, Grossman Institute for Neuroscience, Quantitative Biology and

Human Behavior, University of Chicago, 60637

[3]Department of Neurology, University of Wisconsin-Madison, School of Medicine and Public

Health

Corresponding authors: madar@uchicago.edu, mathewjones@wisc.edu

Number of pages: 59

Number of figures: 9

Number of words: Abstract (158), Introduction (650), Discussion (1499)

Conflicts of interest statement: The authors declare no conflict of interest.

Acknowledgements: We thank Lin Lin who helped start behavioral tests in mice in the Jones lab

and Drs. C.E.L Stark and M.A. Yassa for allowing us to use software and materials they developed.

This work was supported by the US National Institutes of Health (MVJ: RO1 NS075366), the

University of Wisconsin Institute for Clinical and Translational Research (MVJ.; NIH/NCATS

UL1TR000427), the US Department of Defense (RKM: PR161864) and Lily's Fund for Epilepsy

Research (ADM.; 2015 fellow).

Abstract:

In temporal lobe epilepsy, the ability of the dentate gyrus to limit excitatory cortical input to the hippocampus breaks down, leading to seizures. The dentate gyrus is also thought to help discriminate between similar memories by performing pattern separation, but whether epilepsy leads to a breakdown in this neural computation, and thus to mnemonic discrimination impairments, remains unknown. Here we show that temporal lobe epilepsy is characterized by behavioral deficits in mnemonic discrimination tasks, in both humans and mice (C57Bl6 males, systemic low-dose kainate model). Using a recently developed assay in brain slices of the same epileptic mice, we reveal a decreased ability of the dentate gyrus to perform certain forms of pattern separation. This is due to a subset of granule cells with abnormal bursting that can develop independently of early EEG abnormalities. Overall, our results linking physiology, computation and cognition in the same mice advance our understanding of episodic memory mechanisms and their dysfunction in epilepsy.

Keywords:

 Patch-clamp, low-dose systemic kainate model**,** kainic acid, interictal spikes, novelty recognition, firing rate, spiking patterns, similarity metrics, input-output, entorhinal cortex, perforant path, EPSC, excitation/inhibition ratio, dentate gate, Hebb-Marr theory

Introduction:

Temporal lobe epilepsy (TLE) is characterized by recurring focal seizures originating in or near the hippocampus (Toyoda et al., 2013), microcircuit pathologies in brain regions including the hippocampus (Alexander et al., 2016) and memory-related cognitive deficits (Helmstaedter et al., 2003; Zhao et al., 2014). Although the hippocampus is a nexus for episodic memory that is critically affected during TLE, the relationship between TLE and hippocampus-dependent memory is insufficiently understood.

Episodic memory formation is thought to involve storage of neural representations in area CA3 of the hippocampus via Hebbian plasticity at recurrent excitatory synapses of coactivated cells (Rolls, 2010). In this view, a partial cue reactivates a subset of the CA3 ensemble that in turn recruits the other neurons of the original pattern resulting in recall of the original event. However, recurrent excitation is problematic because it theoretically a) predisposes the network to over-excitation that could trigger seizures (Traub and Wong, 1982; Le Duigou et al., 2014) and b) limits the number of patterns that can be stored without overlap (Rolls, 2010). Overlapping memory representations would in turn lead to interference during recall and thus cognitive confusion. To solve these problems, it was proposed that the dentate gyrus (DG) of the hippocampus acts as a) a gate and b) a pattern separator, so that similar cortical representations are transformed into sparse and dissimilar patterns before reaching CA3 (O'Reilly and McClelland, 1994; Hsu, 2007; Dengler and Coulter, 2016).

The function of DG as a gate for cortical activity to prevent seizure generation fails in TLE (Hsu, 2007; Krook-Magnuson et al., 2015; Dengler and Coulter, 2016; Lu et al., 2016). Granule cells (GCs), the output neurons of DG, lose their usual sparseness due to network reorganization (Artinian et al., 2011; Dengler and Coulter, 2016; Dengler et al., 2017) making cortical excitation

easier to propagate to CA3 (Behr et al., 1998; Patrylo et al., 1999; Ouedraogo et al., 2016). Even

in healthy animals, repeated excitation of GCs induces seizures (Krook-Magnuson et al., 2015)

and prolonged stimulation of DG causes TLE (Sloviter, 1983).

An alternate view of the DG function, developed in parallel to the concept of dentate gate,

is that it performs pattern separation: the transformation of similar cortical patterns into dissimilar

hippocampal representations. This process theoretically supports mnemonic discrimination, the

ability to distinguish between similar memories. Indeed, DG lesions impair mnemonic

discrimination both in rodents (Treves et al., 2008; Kesner and Rolls, 2015; Kesner et al., 2016)

and humans (Yassa et al., 2011; Baker et al., 2016; Bennett and Stark, 2016; Dillon et al., 2017).

Moreover, computational models (Chavlis and Poirazi, 2017) and recent experiments (Berron et

al., 2016; Knierim and Neunuebel, 2016; Madar et al., 2019a, 2019b; Woods et al., 2020) suggest

that the DG circuitry supports multiple forms of pattern separation (Santoro, 2013). Whether such

computations underlie mnemonic discrimination remains unclear.

Unsurprisingly, TLE negatively impacts hippocampal-dependent memory in humans

(Coras et al., 2014) and rodents (Gröticke et al., 2008; Chauvière et al., 2009; Müller et al., 2009;

Inostroza et al., 2013; Lenck-Santini and Scott, 2015). However, the effect of TLE specifically on

DG computations and DG-dependent cognition is understudied. It was only recently reported that

patients with TLE are impaired on mnemonic discrimination tasks (Reyes et al., 2018; Poch et al.,

2020) and that TLE causes deficits in DG-dependent object location memory in mice (Bui et al.,

2018). A computational model has also suggested that hippocampal pathologies in TLE would

degrade DG pattern separation (Yim et al., 2015) but the hypotheses that 1) TLE causes a

breakdown in DG neural pattern separation and 2) that such a failure causes mnemonic

discrimination impairments remain experimentally untested.

Here we investigated, in humans and mice, whether TLE is characterized by deficits in

mnemonic discrimination and then recorded, in brain slices from the same mice, the spiking

patterns of single GCs in response to parametrically varied afferent stimulation in order to gauge

neuronal pattern separation.

Materials and Methods:

Human Behavior. A mnemonic similarity task *(Stark et al., 2019)*, also known as a

behavioral pattern separation (BPS) task *(Stark et al., 2013)*, was administered to 15 patients in

the University of Wisconsin-Madison Epilepsy Monitoring Unit. Under an approved Institutional

Review Board protocol and after obtaining informed consent, the task was administered in

conjunction with a standard neuropsychiatric evaluation and during electroencephalographic

(EEG) recording, both of which are part of standard practice for diagnosing the patients' seizures.

Patients were 18-65 years old, male and female. Only patients with a preliminary diagnosis of TLE

were included in our analysis. As controls, 20 subjects without epilepsy, recruited to match the

patients' age and sex distributions (family members of the patients when possible, or other

volunteers), were also tested. Consent documents, medical records and primary data are on file in

a secure location within the Dept. of Neurology. Data were deidentified prior to analysis. Subjects

received no compensation for participation.

The visual, object recognition-based BPS task is described in Yassa et al. (2011) and has

been further validated by demonstrating mnemonic discrimination deficits during normal aging

and multiple neurologic and psychiatric disorders (Stark et al., 2019). It was implemented on a

laptop by trained neuropsychiatric postdoctoral fellows (J.B. and E.I.P.) using software distributed

freely by the Stark lab (http://faculty.sites.uci.edu/starklab/mnemonic-similarity-task-mst/).

Briefly, participants viewed a series of pictures of everyday objects: During Phase 1, 128 different

images were presented and the subject was asked to classify each as "indoor" or "outdoor", simply

to engage the subject's attention. During Phase 2, a new series of 192 images was presented: 1/3

repeated from Phase 1 (*Repeated*), 1/3 new but similar to images from Phase 1 (*Lure*), and 1/3 new

and completely different from Phase 1 (*Novel*). Participants were asked to classify each image of

the Phase 2 set as "Old", "Similar", or "New". BPS was evaluated with a *discrimination index*

computed as the difference between p("Similar"|Lure) and p("Similar"|Novel), as in past research

(Yassa et al., 2011; Stark et al., 2019).

Mouse Experiments. Male mice (C57BL6J, 5-6 weeks old) were received from Envigo

(formerly Harlan, Madison, WI), housed in groups and allowed to acclimate to their new home

environment for one or two weeks. Animals then underwent epilepsy induction with kainic acid

(KA) injections (see below) (J.A.P.). A control group of mice was injected with saline and another

received no injection: both were pooled together and considered as the control group because our

analyses did not reveal any difference. 7-9 weeks after injection, mice started 4 weeks of

behavioral testing on the BPS task. After completion of behavioral testing, animals (~18-20 weeks

old) were transferred to a different building for EEG implantation. Each animal was recorded

continuously for three days before being transferred back to the original building and sacrificed

for slice electrophysiology (~20-24 weeks old). After each building-to-building transfer, mice

were allowed a period of acclimation ranging from two days to two weeks. Behavioral testing

(J.A.P, S.R. and M.C.), EEG recordings (E.P.W.) and slice electrophysiology (A.D.M) were

performed by different researchers, all of whom were blind to KA or Control treatments.

Epilepsy induction. During the induction process, when not being handled, mice were

individually housed in enclosed ~150 cm^3 acrylic cubicles with opaque sides and clear front portals

with holes to allow air exchange, food and bedding. Animals were then randomly assigned to KA

or Control treatment, ear punched for identification and weighed. Animals were induced for

epileptogenesis using the repeated low-dose kainate method *(Hellier et al., 1998; Sharma et al.,*

2018): a series of intraperitoneal injections based on the following schedule. Animals first received

10 mg/kg of kainic acid (Tocris Bioscience, UK) mixed in 1x PBS (8 ml per 10 mg) prepared from

PBS tablets (Dot Scientific, Michigan) with deionized distilled water and filter sterilized

(Millipore, Burlington, MA). Control animals were injected with an equivalent volume of saline

made of 1x PBS. Animals continued to receive injections at 5mg/kg every 20 minutes until status

epilepticus (SE) occurred (same number of saline injections in control mice). Some animals

received alternating 5mg/kg and 2.5 mg/kg injections after the initial 10 mg/kg dose, but we found

this schedule took more time than 5 mg/kg injections and our survival rate (>90% across all

cohorts) was no different between the two schedules. Animals were considered to be in SE when

displaying persistent behavioral seizures of level 4-5 on the Racine Scale (Racine, 1972), less than

5 minutes apart, for a minimum of 30 minutes. Animals received 4-9 injections depending on

tolerance to kainic acid and injection schedule. After the injection schedule, animals were given

fresh apple slices, monitored until SE ceased (assessed from normal posturing, within 1-1.5 hours),

and returned to group housing. They were monitored, weighed and animals weighing less than

their preinjection levels were given 0.4 ml of 1x PBS via intraperitoneal injection on a daily basis

until their weight exceeded preinjection level. No animal injected with saline ever experienced

status epilepticus, lost weight, or required recovery injections of 1x PBS. Mice remained in the

vivarium for 7-9 weeks after injections, allowing time for the development of epilepsy in KA mice

− the latent period in the KA model of TLE has previously been evaluated as 10-30 days before

the first spontaneous electrographic seizure, and ~11±5.4 weeks before the first spontaneous

convulsive seizure *(Lévesque and Avoli, 2013)*. During this time some KA mice became aggressive

and many of the aggressor animals were removed from group housing and caged individually.

Mouse Behavior. At 14-16 weeks of age, mice started behavioral testing on a variant of a

novelty recognition-based BPS task that is well established in rodents to report mnemonic

discrimination abilities *(van Hagen et al., 2015)*. Our protocol lasted 4 weeks, including 1 week

of habituation and 3 weeks of trials. During the first week of habituation, each mouse was gently

handled by the experimenter for five minutes every day. On the second week mice were split into

three groups (A, B, and C). Each group went through a three-day schedule where animals were

habituated on the first and second days and performed the object location task on the third day.

Groups were staggered by one day each, such that group A was scheduled Monday-Wednesday,

B Tuesday-Thursday and C Wednesday-Friday. Animals were housed following a 12:12 light-

dark cycle and always handled or tested between 1 pm and 4pm (light period).

The arena used for the BPS task was a 63.5 x 63.5 x 15 cm open topped Plexiglas box

wrapped on the outside and bottom with black felt cloth. On each side of the arena were different

images serving as proximal cues for orientation. The room also had numerous distinct distal cues

(e.g. the video camera placed over the center of the arena, shelving, etc.), that were kept

unchanged. During behavioral testing, the room was lit with a single bank of overhead fluorescent

lamps.

For the BPS task, animals underwent three phases of exploration in the arena, each lasting

3 min and separated by 1.5 min of single housing in a solitary chamber. In Phase 1 (habituation)

the arena was empty, in Phase 2 (sampling) the arena contained two identical objects in the center,

and in Phase 3 (testing) one object had been randomly selected and displaced by a distance of 7,

14 or 21 cm. The arena and objects were cleaned with 70% ethanol before and after each phase.

During Phases 2 and 3, the experimenter recorded the amount of time that the mouse spent

interacting with each object with two silent stopwatches. The mouse was considered to be

interacting with an object if it was within 1 cm of it and oriented towards, sniffing, scratching, or

on top of the object. These three phases were repeated three times with a different distance each

week, with the order of distances randomized. Objects used for the task were ~6 cm tall block style

plastic colored figures or metal cylinders, with a footprint of ~2 cm in diameter.

All phases of the experiment were recorded with a 1080p webcam positioned above the

arena. During all phases, the experimenter sat in the corner of the room out of the line of sight of

the mouse. The experimenter was consistent in appearance and smell (e.g., same experimenter for

a given cohort, wearing same white lab coat and gloves of the same color each day; use of scented

soap, etc. was minimized).

BPS Analysis. The discrimination ratio for each trial was computed as $(T_{moved} - T_{unmoved})$ /

T_{total} where T_{moved} is the time spent exploring the moved object, $T_{unmoved}$ is the time spent exploring

the unmoved object and T_{total} is the sum of T_{moved} and $T_{unmoved}$. The exploration times used to

calculate the reported discrimination ratio were manually collected by experimenters during the

BPS task. We (M.V.J. and J.A.P.) also developed a home-written program using video recordings

for motion-tracking and trajectory analysis of each mouse during behavioral testing.

Discrimination ratios computed from automated tracking data were well correlated to the ratios

from manually recorded times, and led to similar results ($R^2 = 0.68$, $T(117) = 16.07$, $P < 0.001$).

Automated tracking data were also used to assess a) distance traveled as an estimate of motility

and exploration and b) tendency to stay close to walls (i.e., thigmotaxis) as an estimate of anxiety.

Thigmotaxis *(Simon et al., 1994)* was evaluated as the proportion of time spent within 6.35 mm of

the wall (10% of the arena width).

Mouse electroencephalography (EEG). EEG electrode implantation was performed at

~18-20 weeks of age for all animals, following a previously established protocol *(Wallace et al.,*

2015). Briefly, mice were anesthetized with isoflurane and stainless steel screw electrodes were

implanted in the skull (bregma +1.5 mm and 1 mm right, bregma -3 mm and 1 mm left, and lamda

-1 mm at midline). Two stainless steel braided wires were placed in the nuchal muscles for

electromyography (EMG) recording. After a 72 hour recovery, we transferred mice into individual

tethered EEG acquisition chambers and allowed a >12 hour acclimation period. We acquired EEG

and EMG signals continuously for 3 days. Recordings were digitized with an XLTek amplifier

(XLTEK, USA) sampled at 1024 Hz. Ad libitum access to food and water was ensured. Note that

the resulting EEG dataset was previously reported and analyzed in Pfammatter et al. (2018).

Interictal spike (IIS) analysis. In humans and most animal models of acquired TLE, overt

seizures are relatively rare (often much less than once per day) *(Lévesque et al., 2016).* In contrast,

nonconvulsive and subclinical epileptiform events such as interictal spikes (IISs) can be very

frequent, up to hundreds per day. Therefore, in order to assess epileptiform activity in KA animals,

we used a modification of our previously published principal components (PC)-based method to

quantify IISs *(Pfammatter et al., 2018)* and calculated an "Hourly IIS index" for each animal. All

detected high-amplitude EEG events from 9 Ctrl and 15 KA animals were projected into the space

spanned by their first three PCs. The main modification here is that, instead of using a Gaussian

Mixture Model (GMM) to assign events to 'clusters', we simply gridded the PC space into 'voxels'

and computed relevant quantities within each voxel exactly as we computed those same quantities

within GMM 'clusters' previously: 1) The probability that events within a voxel are characteristic

of an epileptogenic treatment (i.e., KA) was computed as the voxel-wise proportion of events

coming from KA mice, 2) Only the voxels above chance level were considered specific to

epileptogenic treatment and probabilities were scaled accordingly, 3) The Hourly IIS Index of a

given animal was defined as the average frequency of detected events weighted by the scaled

probabilities of each event's voxel. This new voxel-based method has the advantages to make no

assumptions about the structure of the data in PC space or about the number of clusters to fit.

Instead, the main free parameter is now the voxel volume: we selected a size of 10 cubic PC units

following the same optimization procedure described in Pfammatter et al. (2018) in order to avoid

overfitting. Extensive exploration (not shown) revealed that the final results are similar to the

GMM method and are not importantly affected by moderate changes in voxel volume.

Slice electrophysiology. Mice used for electrophysiology were p115-p182 at the time of

experiment (mean ± SEM: p141 +/- 4 days; no difference in age between treatments: KA n = 13,

Ctrl n = 6, U-test: P = 0.9, Z = 0.2, rank sum = 246). Age was also not correlated with any summary

statistics presented in this study.

Adult mice were euthanized by transcardial perfusion with oxygenated PBS under

isoflurane anesthesia, before decapitation and brain extraction. 400 μm horizontal slices of the

ventral and intermediate hippocampus were prepared as detailed in Madar et al. (2019a). After

slicing in a sucrose-based cutting solution *(Yi et al., 2015)*, slices were transferred to an incubation

chamber filled with 50% cutting solution and 50% artificial cerebrospinal fluid (aCSF) at 37°C for

30 minutes, then room temperature. Patch-clamp recordings were done in a chamber submerged

with aCSF containing (in mM) 125 NaCl, 25 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 2 CaCl2, 1

MgCl2, and 25 D-Glucose, flowing at 5 ml/min and saturated with a gas mixture of 95% O_2 and

5% CO_2. Stimulation was applied through a double-barreled "theta" pipette filled with aCSF. Patch

pipettes were filled with an intracellular solution of the following composition (in mM): 135 K-

gluconate, 5 KCl, 0.1 EGTA, 10 HEPES, 20 Na-Phosphocreatine, 2 Mg_2-ATP, 0.3 Na-GTP, 0.25

CaCl$_2$ adjusted to pH 7.3 with KOH and 310 mOsm with H$_2$O, leading to a 2-5 MΩ pipette

resistance in aCSF. Whole-cell patch-clamp recordings of single DG GCs in response to electric

stimulation of the outer molecular layer (perforant path) were performed as detailed in Madar et

al. (2019a), and the stimulation protocols used to test neural pattern separation were the same as

in Madar et al. (2019b). Briefly, input sets used for stimulation were composed of five (type 1) or

ten (type 2 and 3) spiketrains (two seconds long), delivered sequentially (separated by five seconds

of pause) and repeated ten or five times, respectively, in order to yield fifty output spiketrains. The

stimulation pipette was placed >100µm lateral to the recorded GC to avoid direct stimulation of

GC dendrites, with the baseline membrane potential held at -70 mV for current and voltage-clamp

recordings.

Intrinsic electrophysiological properties of recorded GCs were the following (mean ±

SEM for Ctrl / KA): resting membrane potential V$_{rest}$ = -78.0 ± 1.9 / -81.4 ± 1.2 mV; membrane

resistance R$_m$ = 139 ± 17 / 178 ± 16 MΩ; membrane capacitance C$_m$ = 18 ± 1.2 / 17 ± 0.7 pF. There

were no significant differences between control and KA mice (U-tests: P = 0.1, 0.2, 0.7; Z = 1.6, -

1.2, 0.4; rank sums = 294, 198.5, 252.5 respectively).

Neural pattern separation analysis. Similarity between spiketrains was assessed as in

Madar et al. (2019a, b), using three metrics based on dividing spiketrains into time bins of a

specific duration τ$_w$: the Pearson's correlation coefficient (R), the normalized dot product (NDP)

and the scaling factor (SF). NDP and SF consider spiketrains as vectors of spike count per bin and

measure the angle (NDP is the cosine) and the ratio of norms (SF) between the two vectors. R

considers vectors of deviation from mean spike count, which loses the original angle between the

vectors of raw spike counts and leads to considering common periods of silence as correlated,

unlike NDP. Both R and NDP are sensitive to binwise synchrony, whereas SF is more sensitive to

differences in firing rate and burstiness (see Madar et al. 2019b for a detailed discussion on the

different neural codes assumed by each metric). To further test the role of various spiketrain

features in mediating pattern separation, we also considered neural codes purely focused on either

1) the average firing rate (FR), 2) the compactness (1 - proportion of time bins with at least one

spike) and 3) the occupancy (average number of spikes in bins with at least one spike). These were

computed as in Madar et al. (2019b) and the degree of pattern separation (or convergence, for

negative values) was evaluated as the difference between the dispersion of the output spiketrains

minus the dispersion of the input spiketrains. The dispersion was computed as the mean absolute

value of pairwise differences for each spiketrain feature (FR, Compactness or Occupancy) over all

spiketrains of a set, excluding self-comparisons and, in the case of the output sets, all comparisons

between output spiketrains resulting from the same input spiketrain. When using a similarity

metric S (R, NDP or SF), pattern separation was assessed following the same logic. First, the

similarity between each pair of spiketrains of a given set was computed. S_{input} was the average for

all pairs of an input set (excluding self-comparisons), and S_{output} was the average of all pairs in

output sets excluding comparisons between output spiketrains coming from repetitions of the same

input train. This yielded a single output similarity value for a given output set, and the degree of

pattern separation was thus S_{input} - S_{output}. To gain a finer view, we also performed a pairwise

analysis (as in Madar et al. 2019b) where each of the pairwise S_{input} of an input set were

distinguished (10 pairs for input sets of type 1, 45 pairs for input sets of type 2 and 3). The pairwise

S_{output} (pw S_{output}) was thus the average similarity across all pairs of output spiketrains resulting

from a given pair of input spiketrains. Pattern separation was computed as pw S_{input} – pw S_{output}.

Experimental design and statistical analysis. Sample sizes correspond to the standards of

each field (human behavior, mouse behavior, slice electrophysiology) such that large effects could

be detected. Data analysis was performed using MATLAB (Mathworks, Natick, MA, USA). The

Lilliefors test was used to verify the normality of data distributions *(Lilliefors, 1967)*. Parametric

or nonparametric statistical tests were appropriately used to assess significance (p-value < 0.05).

Throughout the Results section, KW ANOVA corresponds to the nonparametric Kruskal-Wallis

analysis of variance, U-test corresponds to the Wilcoxon rank-sum test equivalent to the Mann-

Whitney U-test, and KS test corresponds to the two-sample Kolmogorov-Smirnov test (sidedness

is specified in the legends).

To analyze performance of mice on the BPS task we performed a two-way ANOVA with

the Matlab function *anova* based on a linear mixed-effects model built using *fitlme,* with the

distance of the moved object as a continuous fixed effect, animal treatment (Ctrl vs. KA) as a

categorical fixed effect, and animal identity nested within object distances as a random effect to

account for repeated measurements. The same procedure was used to test for differences in

distances travelled and thigmotaxis.

In the pattern separation analysis, to determine whether (S_{input}, S_{output}) distributions were

significantly different, we performed an analysis of the covariance (ANCOVA) using separate

parabolic or linear regression models, implemented in MATLAB with a custom-written code

following the method described in (Motulsky and Ransnas, 1987), as in Madar et al. (2019a, b)

For the non-linear regression between pBurst and pattern separation, we used the Matlab

function *fitnlm* with a three-parameter model function $f(x) = a/(x + b) + c$.

Code accessibility. Raw data for all figures are available at:

https://www.ebi.ac.uk/biostudies/studies/SBSST353. Code to read and analyze the data is

available at: https://github.com/jessePfammatter/DeficitsBehavNeuronalPatternSep and

https://github.com/antoinemadar/PatSepSpikeTrains

Results:

Behavioral pattern separation deficits in TLE. Mnemonic discrimination is often called behavioral pattern separation (BPS) because it is hypothesized to be the behavioral outcome of neural pattern separation. It is the ability to discriminate between similar experiences (event, environment, object, etc.) that occurred at different times *(Santoro, 2013)*.

We first investigated the effect of TLE on mnemonic discrimination in humans. We used the *Mnemonic Similarity Task* developed by the Stark lab, an established DG-dependent BPS task (Baker et al., 2016; Stark et al., 2019) where participants must distinguish between similar images presented at different times (**Figure 1A**, **Materials and Methods – Human behavior**). Patients previously diagnosed with TLE showed a severe deficit (~50%) compared to nonepileptic subjects in their ability to correctly identify objects as being similar but not identical (**Figure 1**). These results confirmed our hypothesis that TLE impairs DG-dependent mnemonic discrimination, thus warranting further study of the impact of TLE on DG computations that might explain this cognitive deficit.

To investigate the effect of TLE at the behavioral, computational and cellular levels, we turned to the common post-kainate animal model of acquired TLE (KA, **Methods – Mouse experiments**). An automated detection algorithm of interictal spikes (IISs) allowed us to quantify epileptiform activity in each animal even in the absence of seizures (**Methods – IIS analysis** and see Pfammatter et al. 2018). Although we recorded seizures in only two KA animals (Pfammatter et al. 2018), most KA mice developed IISs (**Figure 2**), demonstrating epileptogenesis. The detected IISs belonged to two distinct groups (**Figure 2D**), both having waveform features typical

of epilepsy (**Figure 2E-F**) (White et al., 2010; Chauvière et al., 2012; Huneau et al., 2013;

Pfammatter et al., 2018).

Prior to EEG recordings, mice were subjected to an object-location novelty-recognition

BPS task (**Methods – Mouse Behavior**) where their ability to discriminate between a moved

object versus an identical unmoved object was measured for different object displacement

distances (**Figure 3A**). Our results demonstrate that, on average, KA mice have lower object-

location mnemonic discrimination than control mice over all object distances (**Figure 3B-C**).

Control analyses revealed that KA mice have no locomotor deficits (assessed by the total distance

travelled) and do not display higher anxiety levels (assessed by the time spent near walls) during

the sampling and testing phases (**Figure 3D-E**). Moreover, there was no correlation between

individual mnemonic discrimination ratios and total distance moved during the testing phase (R^2

$= 28\%$, *T(118) = 1.86*, *P* = 0.066) or time spent near walls ($R^2 = 2\%$, *T(118) = 0.50*, *P* = 0.62).

These results suggest that differences between KA and control mice in the BPS task were not due

to differences in motor ability, exploration or anxiety.

TLE is thus accompanied by mnemonic discrimination deficits in both humans and mice.

DG computational impairments in TLE. To test our hypothesis that DG pathologies in

TLE lead to a breakdown of the neural pattern separation function of the DG, which in turn would

translate into the cognitive impairments we observed, we measured pattern separation in

hippocampal slices from the same KA and control mice used above in behavioral and EEG

experiments. Although the slice preparation isolates the DG from its natural inputs, it provides the

advantage that applied inputs and recorded outputs can be known with certainty, a necessary

condition for quantifying circuit-level pattern separation (Santoro, 2013; Madar et al. 2019a).

Briefly, the assay has three steps (**Figure 4A-B** and see Madar et al. 2019a, b): 1) ensembles of

stimulus patterns (simulating afferent input spiketrains) are generated, with known degrees of

similarity to each other. These spiketrains are then fed into the DG by stimulating the lateral

perforant path. 2) The response of a single GC is recorded in whole-cell current-clamp. 3) The

similarity between the output spiketrains is compared to the similarity between the input

spiketrains, revealing the degree of separation or convergence (**Figure 4C** and see **Material and**

Methods – Neural pattern separation analysis).

We tested pattern separation levels in response to three types of input sets. Input sets of

type 1 were constituted of Poisson spiketrains with a 10 Hz mean firing rate designed to have a

prespecified average similarity as measured by the Pearson's correlation coefficient (R) (**Figure**

4B). For an input set with spiketrains correlated by 77% between each other (timescale: $\tau_w = 20$

ms), the output spiketrains of GCs from epileptic mice had a higher average correlation (R_{output})

than GCs from control mice (**Figure 4C**). This demonstrates a decrease in pattern separation in

the DG of epileptic mice. Exploring a wider range of input correlations shows that the pattern

separation function of the DG of KA mice is generally impaired compared to the normal pattern

separation function (**Figure 4D**). Furthermore, a decrease in pattern separation was observed at

multiple timescales (millisecond to second) and using multiple similarity metrics that assume

different neural codes (**Figure 4E**, see **Material and Methods – Neural pattern separation**

analysis). The impairment is most noticeable using the NDP metric, showing that DG output

patterns are less orthogonalized in KA animals at all timescales, whereas differences in pattern

separation via scaling (SF metric) are small but significant at large timescales.

An input set of type 2, made of Poisson spiketrains similar in terms of R but with varying

firing rates (**Figure 5A1**), and an input set of type 3, with 10.5 Hz uncorrelated trains (R = 0) with

varying burstiness (**Figure 5B1**), were designed to explore a wide range of input similarities as

measured by SF and to characterize DG computations on inputs with a variety of statistical

structures (Madar et al., 2019b). As we have shown before, the normal DG exhibits low pattern

separation via scaling for highly similar inputs, and significant pattern convergence for dissimilar

inputs (Madar et al., 2019b). In KA mice, GCs show a decrease in both pattern convergence and

separation via scaling: the DG computation function is steeper, closer to the identity line, meaning

that the DG of KA mice is weaker at transforming the similarity of its inputs in general (**Figure**

5A1 and B1). When similarity is measured with R or NDP (binwise synchrony code) instead of

SF (neural code focused on FR and bursting), these experiments confirmed that pattern separation

is decreased in TLE (**Figure 5A2**) and that this deficit is strongest when inputs are highly similar

(i.e., when pattern separation is theoretically most important) and weaker when inputs are already

dissimilar (**Figure 5B2**).

398   ***Pathological spiking patterns in a subset of GCs from epileptic mice.*** What features of

GC output spiketrains are changed in TLE to explain the difference in DG computations between

KA and control mice? On visual inspection, we noticed that many KA neurons occasionally fired

short bursts of action potentials after a single input pulse (**Figure 6A**), which is unusual for control

GCs (Madar et al., 2019a). To determine whether there was a difference in the spiking patterns of

KA and control GCs, we measured three spiketrain features: 1) the average firing rate (FR) of a

GC across a full recording set (fifty spiketrains), 2) the probability of spiking after a single input

pulse (SP) and 3) the probability of bursting (p(Burst), i.e., more than one spike) after a single

input pulse. Our results show that on average GCs from KA mice fire more faithfully after an input

(**Figure 6B**) and also tend to fire in bursts (**Figure 6C-D**), which together lead to a higher FR

(**Figure 6B, E**).

Closer inspection suggests that the distributions of SP, p(Burst) and FR are different

between KA and controls, with larger variance and upper tails for GCs from KA mice **(Figure 6B-**

E). Although we did not formally test for discrete clusters, visual inspection of Figure 6E shows

that some GCs from KA animals have higher SP, p(Burst) and FR than any control GC. Thus, it

appears that, following an epileptogenic insult, only a subset of GCs displayed pathological

characteristics amidst a background of seemingly normal GCs.

What is the origin of the pathological spiking patterns? We first controlled that stimulation

intensities were comparable between treatment groups and were not causing the abnormal spiking

patterns (**Figure 7A**). We next asked whether the abnormal firing came from changes in the

synaptic drive. To answer this question, we measured the synaptic input-output relation (Ewell and

Jones, 2010) by recording GCs first under current-clamp then under voltage-clamp, in response to

the same pattern separation stimulus protocol (**Figure 7B1-2**). Our data show that the average

excitatory drive per neuron was not different between KA and control groups, and that it does not

predict burstiness (**Figure 7B3**). A finer grained analysis on the individual currents suggests that

although the excitatory drive and the spiking output are correlated, high amplitude EPSCs are

neither sufficient nor necessary to elicit a burst (**Figure 7B4**). Bursting does not seem to result

from other changes in excitatory input-output coupling either (**Figure 7B5**), which suggest that

inhibition might be involved. GCs are indeed subjected to strong tonic, feedforward and feedback

inhibition that controls the sparseness of their activity (Coulter and Carlson, 2007; Ewell and

Jones, 2010; Pardi et al., 2015; Lee et al., 2016) and is altered in TLE (Alexander et al., 2016;

Dengler and Coulter, 2016). We did not record evoked IPSCs directly but, in normal mice, partial

block of inhibition elevates GC firing rates and causes bursts (Madar et al., 2019b) similar to what

we observed here in KA mice. Although indirect, our results thus converge to support the

hypothesis that pathological spiking arises from an excitation/inhibition imbalance in a subset of

GCs, likely driven by a decrease in inhibition.

How do high FR and bursts relate to the DG computational impairment measured in KA

mice? We first tested the hypothesis that higher GC firing rates and burstiness yield less pattern

separation as measured with R (**Figure 8A**). In GCs of young mice, FR and pattern separation are

related, but loosely (Madar et al., 2019a), whereas here in adult mice there is a strong linear

relationship, such that GCs with pathologically high FR exhibit pathologically low pattern

separation. Similarly, abnormal bursting corresponds to abnormally low levels of pattern

separation. This analysis relates cell-wise spiking features (FR or pBurst averaged across all spike

trains of a recorded neuron) to a form of pattern separation that is not theoretically concerned with

such features (Madar et al., 2019b).

Other forms of pattern separation, more directly related to FR or bursting, can in theory

also be performed: for example by increasing spiketrain-to-spiketrain *variability* in FR or

burstiness of the output neuron, even if the average quantities were identical (Madar et al., 2019b).

Because pathological GCs only occasionally fired bursts, we asked whether TLE could affect such

forms of pattern separation. To test this, we measured the spiketrain-to-spiketrain variability in FR

as well as in two complementary measures of burstiness (occupancy and compactness, see

Materials and Methods – Neural pattern separation analysis). For input sets of type 1 (10 Hz

Poisson), GC output variability was slightly lower in terms of burstiness or FR. This weak type of

pattern convergence disappeared in GCs from KA mice (**Figure 8B**). For input sets of type 2 and

3, no significant difference between KA and Ctrl was detected (**Figure 8C-D**). Overall, Figure 8

suggests that TLE reduces DG neural pattern separation mostly by raising average levels of FR

and burstiness, rather than through spiketrain-to-spiketrain variations of those features.

Electrographic, behavioral and computational pathologies in individual mice. We have

shown that, on average, KA mice develop EEG abnormalities (**Figure 2**), suffer mnemonic

discrimination impairments (**Figure 3**) and have a bursting subpopulation of GCs with pattern

separation deficits (**Figures 4-8**). Because we often performed all of the aforementioned

experiments in the same mice, we next asked how those different epilepsy-related pathologies are

linked at the individual level (**Figure 9**). In a simple framework where behavioral impairments are

caused by computational deficits that are due to reorganization of DG network function that also

leads to an increase in epileptiform EEG events, one could expect a simple relationship between

all the variables we measured. Interestingly, our data suggest a more complex view. For example,

animals with few interictal spikes may still harbor some pathological neurons (**Figure 9A**) and be

highly impaired at the BPS task (**Figure 9B**). This shows that both DG network pathologies and

mnemonic impairments can occur independent of, or before, early EEG abnormalities. Conversely,

both normal and pathological GCs were recorded in mice with more advanced EEG pathology,

which confirms that those two subpopulations can coexist in the same KA animal (**Figure 9A**).

Mice with clearly epileptiform EEG activity can also sometimes perform normally on the BPS

task (**Figure 9B**). As expected from theory we found a correlation between pattern separation in

single GCs and mnemonic discrimination, but only for the behavioral test with medium

interference levels (**Figure 9C**). When considering behavioral performance over all interference

levels, the variability in individual behavior and in cell sampling prevents us to definitively

conclude, but we note that KA animals with the largest overall cognitive impairments all had at

least one pathological recorded GC (**Figure 9C**). Finally, in cases where EEG, BPS and patch-

clamp data were all obtained from each animal, the combination of these measurements yielded a

very obvious separation between normal and KA subjects (**Figure 9D**). Note that considering an

EEG index based on individual clusters of IISs identified in Figure 2D did not change any of the

conclusions stated above. Overall, our results suggest that epilepsy-related pathologies do not all

develop in concert and that computational, behavioral and electrographic measures provide

complementary informative dimensions that, together, better assess the epileptic state than any

single dimension or pair of dimensions.

483
Discussion

Cognitive and computational deficits in TLE. We provide the first experimental

evidence that TLE is characterized by both mnemonic discrimination impairments and neuronal

pattern separation deficits.

Our data in human patients (Figure 1) is consistent with recent studies showing that TLE leads to

spatial or object mnemonic discrimination impairments (Reyes et al., 2018; Poch et al., 2020).

Reyes and colleagues employed a spatial memory task whereas Poch and colleagues used an

object-recognition task akin to ours. However, the specific task that we used is known to be DG-

dependent in humans (Baker et al., 2016; Stark et al., 2019), allowing us to conclude that

impairments in TLE patients are likely due to DG alterations. Our behavioral experiment in mice

(Figure 3) also complements a previous study (Bui et al., 2018) by testing short-term (minutes)

rather than long-term (24h) object-location memory and by testing multiple, parameterized levels

of mnemonic interference, as recommended for rigorous testing of mnemonic discrimination

(Hunsaker and Kesner, 2013; Liu et al., 2016). Although not demonstrated, the DG-dependency

of our task in mice is likely because: 1) variants of this task are DG-dependent (Bui et al., 2018;

Pofahl et al., 2020), 2) in rodents, DG supports discrimination for short to long term memory, as

long as the task is not too easy or difficult (Gilbert et al., 2001; Engin et al., 2015; Pofahl et al.,

2020), 3) we tested several degrees of difficulty and found a graded response and an overall

deficit in TLE. Our results and the literature together suggest that mnemonic discrimination

deficits are shared by humans and mice with TLE, occur across different memory domains and

modalities, and point to DG malfunctions.

The impact of TLE-related DG pathologies on neural pattern separation had so far been

investigated only in a computational model (Yim et al., 2015). This model suggested that mossy

fiber sprouting and increased synaptic transmission from the perforant path to GCs can

theoretically lead to a breakdown of pattern separation. We experimentally confirmed that TLE

leads to a deficit in neural pattern separation (Figure 4). However, the deficit we observed is not

as large, possibly because the model 1) considered synaptic transmission as deterministic rather

than probabilistic and plastic, 2) assessed a different form of pattern separation (patterns were

defined as population ensembles of very short spiketrains, thus focusing on a population code

whereas we investigated temporal codes) and 3) considered severe degrees of epilepsy.

The role of sparsity in neuronal pattern separation and DG gating. We found that GCs

from epileptic mice exhibit pathological spiking consistent with a breach of the DG gatekeeper

function (Figure 6) that leads to decreased neuronal pattern separation (Figure 8A). As noted by

others, the gating and pattern separation functions of DG may be related (Dengler and Coulter,

2016) because the filtering of incoming cortical activity into a sparser output can be a

mechanism underlying pattern separation based on population codes (O'Reilly and McClelland,

1994; Dieni et al., 2016; Cayco-Gajic and Silver, 2019; Braganza et al., 2020). Our results

suggest that maintaining a sparse output also supports pattern separation of spiketrains in the

time domain, at the level of single GCs.

Rare investigations of the spiking output of single GCs in epileptic tissue have reported

that single stimulation of the perforant path sometimes leads GCs to fire bursts of spikes (Lynch

et al., 2000; Kobayashi and Buckmaster, 2003). We expand on this by showing that bursts also

emerge in response to complex, naturalistic stimulation patterns (**Figures 6 and 7**). It is important

to note that occasional bursting is observed in normal GCs in vivo (Pernía-Andrade and Jonas,

2014), but this is very unusual in slices (Mongiat et al., 2009; Ewell and Jones, 2010; Zhang et al.,

2012; Dieni et al., 2016; Madar et al., 2019a). In our experiments, bursting GCs therefore

demonstrate a breach of the dentate gate. Because bursts are known to be more efficient at driving

spiking in CA3 pyramidal neurons (Henze et al., 2002), a single GC firing bursts will have a higher

probability of making downstream CA3 pyramidal cells fire. This could have deleterious effects

on memory encoding and promote seizures, as more active CA3 neurons would 1) increase the

chance of overlap between memory representations (Madar et al., 2019a, 2019b; Lee and Lee,

2020) and 2) overexcite a recurrent excitatory circuit, a mechanism of seizure generation (Traub

and Wong, 1982; Santhakumar et al., 2005; de la Prida et al., 2006). Thus, our findings are

consistent with the idea that pathological firing or bursting of even a subset of GCs could

simultaneously contribute to deficits in mnemonic discrimination and epileptiform activity in TLE.

Our study also provides new insight on the mechanisms underlying GCs increased ability

to burst by testing the influence of the excitatory drive (**Figure 7B**). The main source of excitation

to GCs, the perforant path, is sometimes assumed to be augmented in TLE (Yim et al., 2015) given

evidence of increased release probability at the lateral perforant path synapse (Scimemi et al.,

2006). Our results do not directly contradict this, but we did not find a difference in the average

synaptic current evoked in GCs by Poisson input spiketrains applied to the perforant path (**Figure**

7B3). More work is required to carefully test potential TLE-related changes in the coupling

between perforant path and GCs, as well as EPSC kinetics and short-term dynamics (Madar et al.,

2019a), but it is clear that the size of excitatory drive alone is not a good predictor of bursting

propensity in GCs (**Figure 7B3-4**). We also previously showed that partial blockade of inhibition

can produce GC bursting and pattern separation deficits (Madar et al., 2019b). Thus, in accordance

with past research (Kobayashi and Buckmaster, 2003), we suggest that the strong decrease in

synaptic inhibition characteristic of TLE (Dengler and Coulter, 2016; Dengler et al., 2017) is a

likely driver of pathological bursting.

Perhaps the most intriguing insight we bring on the failure of the DG as a gate and pattern

separator in TLE is that this failure occurs only in a subset of GCs, upon a background of

apparently normal GCs in the same epileptic network. This finding invites many more questions:

In the epileptic DG, is there a spectrum of spiking patterns among GCs going from normal to

pathological, or are there two discrete types? What is the ratio of normal to pathological GCs and

does it evolve during epileptogenesis? Do GCs become pathological due to changes in intrinsic

properties or networking? Our results suggest deficits in inhibition, but additional mechanisms

could be in play (Kelly and Beck, 2017). Anatomical or biophysical differences between GCs

could be related to different birthdates of adult-born GCs relative to an epileptogenic brain insult,

resulting in only a subset of GCs integrating abnormally into the DG network (Kron et al., 2010).

Alternatively, pathological GCs could develop from a more susceptible and active subclass of

mature GCs (Erwin et al., 2020). In any case, the discovery that the GC population in the epileptic

DG is not homogeneous in terms of spiking patterns is a crucial step towards developing new

treatments that would specifically target pathological neurons to avoid deleterious side-effects.

The link between mnemonic discrimination, pattern separation and TLE. For thirty

years the DG has been hypothesized to support mnemonic discrimination by performing

neuronal pattern separation (Treves et al., 2008), but despite important efforts (McHugh et al.,

2007; Marrone et al., 2011; Ramirez et al., 2013; Yokoyama and Matsuo, 2016; Woods et al.,

2020), no experiment has yet directly connected DG input-output computations with their role in

memory. Here, we confirmed our previous work showing that the DG itself performs temporal

pattern separation (Madar et al., 2019a, b) and we showed that a deficit in pattern separation is

associated with a deficit in mnemonic discrimination, on average (Figure 3, 4) and possibly at

the individual level (Figure 9C). This is correlative evidence, linking non-simultaneous

experiments, we thus cannot exclude that other DG or non-DG processes (Bernier et al., 2017;

Guise and Shapiro, 2017; Hainmueller and Bartos, 2020) are impaired in KA mice and partially

lead to the observed cognitive deficits. A full demonstration of the causal link between DG

pattern separation and mnemonic discrimination will require measuring and manipulating DG

computations in vivo during episodic memory tasks, but our study is a preliminary step toward

understanding how DG circuit-level computations relate to a high-level cognitive function and

how these processes fail in TLE. A mechanistic understanding of this relation may lead to new

early diagnosis tools (e.g. cognitive tests like in Figure 1) as well as therapies alleviating

memory disorders in TLE. For example, our finding that a subset of GCs with pattern separation

deficits can develop without EEG hallmarks of epilepsy (Figure 9A) suggests a mechanism for

memory impairments that often occur early during epileptogenesis (Jones et al., 2007; Chauvière

et al., 2009; Witt and Helmstaedter, 2015). Similarly, the coexistence of pathological and normal

neurons could explain memory impairments in patients without hippocampal sclerosis (Rayner et

al., 2019).

Figures

Figure 1. Human patients with TLE have mnemonic discrimination deficits

(A) A schematic (adapted with permission from Yassa et al., 2011) of the mnemonic

discrimination task given to patients with TLE and nonepileptic control subjects. The

discrimination index measures the ability of participants to correctly identify images as similar but

not identical to a previously seen image.

(B) Human patients with temporal lobe epilepsy (red, n = 15) display a significant deficit in this

object recognition-based mnemonic discrimination task compared to nonepileptic control subjects

(black, n = 20). Unpaired two-sided T-test: P = 0.0183, T(33) = 2.4832.

(C) The same data from B, presented as cumulative frequency distributions. A non-parametric

one-way KW ANOVA on the indices grouped by treatment confirms a highly significant deficit

in visual pattern separation memory for human subjects with TLE (P = 0.0343, $\chi^2(34)$ = 4.4809).

(D) TLE and Ctrl groups were properly age matched (unpaired two-sided T-test: P = 0.9411, T(31)

= -0.0745).

(E) There were slightly more women than men in the control group, but the gender difference was

not significant between the two treatments (one-way KW ANOVA on gender values, 1 or 2,

grouped by treatment: P = 0.0663, $\chi^2(32)$ = 3.3724).

Figure 2. Kainate-injected mice have epileptiform electrographic events with a range of event frequency

(**A**) The Hourly Interictal Spikes (IIS) index, a proxy for epilepsy severity based on EEG recordings (Pfammatter et al., 2018), was calculated for 9 Ctrl (black) and 15 KA (red) animals. KA animals have a significantly higher Hourly IIS index than Ctrl animals (U-test: $P = 0.009$, $Z = 2.207$, rank sum = 225).

(**B**) To compute the index, high-amplitude events (200 ms around a spike) were identified from left frontal EEG and projected on the first three principal components (PC) of the ensemble of events from all mice. Each dot here is an EEG event, red when from a KA animal and black when from a Ctrl. Notice the 'fingers' of red KA dots extending from the central cluster of mixed Ctrl and KA data points.

(C) Using a voxel gridding of the PC space (voxel size: 10 cubic units), we calculated the

probability *P(KA)* for each event to be specific to KA treatment. The hourly IIS index of a mouse

is the average number of detected events/h, weighted by their probability of being specific to the

KA treatment.

(D) Events with *P(KA) > 0.80*. Two visually identified groups (red ellipses) were separated along

the PC1 axis (group a < -10 and group b > -10).

(E) Each row shows example events from groups identified in D and from the remaining events in

C. EEG amplitude was normalized and reported in terms of standard deviation (SD).

(F) Average waveforms from group a (top), b (middle), and events with *P(KA) < 0.80*. Despite

morphological differences, group a and b both show a wave following the detected spike, which

is a known marker of epilepsy in TLE (Chauvière et al., 2012; Huneau et al., 2013).

Figure 3. Kainate-injected mice have deficits in mnemonic discrimination

(**A**) Object location BPS task arena setup. The experiment was run in three phases of three minutes. In Phase 1, a mouse was allowed to explore the empty square arena. In Phase 2, the same mouse was allowed to explore the same arena but with two identical objects (position indicated by grey circles). In Phase 3, one randomly chosen object was moved either 7, 14 or 21 cm from its original position and the mouse was allowed to explore again. Between phases, the mouse was placed in a solitary chamber for 1.5 minutes. Each mouse performed the task once at each distance with different objects, in randomized order, with a week in between trials.

(**B**) Performance at the BPS task is evaluated with a discrimination ratio computed for each animal at each distance. Healthy mice generally prefer to explore objects with novel characteristics (such as change in relative position) and are thus expected to spend more time with the moved object if they notice it has changed location. Discrimination is thus quantified as the difference between the time exploring the moved object and the time exploring the unmoved object. A two-way ANOVA with repeated measures shows that mice discriminate better when objects are moved farther apart, and that Ctrl mice (n = 29) discriminate better than KA mice (n = 27) (Distance Object Moved: $P = 0.002$, $F(1,118) = 9.84$; Treatment: $P = 0.027$, $F(1,118) = 5.02$).

(**C**) Subset of the data in B plotted in the space of discrimination at each distance, allowing a clearer picture of individual performance. Each animal is represented by a single dot in this 3D space (42 animals with records at the three distances: 24 KA and 18 Ctrl). For each data point, the Mahalanobis distance (De Maesschalck et al., 2000) from the Ctrl centroid is represented by the dot size, and is also displayed in the inset (mean +/- SEM). Mahalanobis distances are larger for KA than Ctrl animals, confirming that the KA population displays mnemonic discrimination deficits (two-sided T-test: $P = 0.003$, $T(40) = -3.15$).

(D,E) Control analysis to determine whether there were motor impairments or anxiety/curiosity

differences between KA and Ctrl mice, across phase 1 (habituation), 2 and 3 of the BPS task. In

each panel, individual mice are represented by 1-3 data points, as in B, depending on how many

conditions (Distance Object Moved) they were tested on. Mean +/- SEM. **(D)** Distance traveled

was used as a proxy for motor ability. During Phase 1 and Phase 2, control and KA mice moved

similar distances ($P = 0.507$, $F(1, 120) = 0.443$ and $P = 0.070$, $F(1, 122) = 3.35$). During Phase 3

KA mice moved farther than Ctrl mice ($P = 0.023$, $F(1, 124) = 5.34$). **(E)** Percent of time near wall

was used as a proxy for anxiety. During Phase 1, KA animals spent more time near the maze walls

compared to Ctrl ($P = 0.03$ $F(1,120) = 4.82$). During Phases 2 and 3, Ctrl and KA animals spent a

similar amount of time near walls ($P = 0.68$, $F(1,122) = 0.173$ and $P = 0.35$, $F(1, 124) = 0.881$).

F-tests correspond to one-way ANOVAs with repeated measures, with Treatment as a categorical

fixed effect.

Figure 4. Epileptic mice have neural pattern separation deficits

(A) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 μm),

overlaid with a schematic of the experimental setup: a theta pipette in the ML is used to focally

stimulate the outer molecular layer (input) while a responding GC is recorded via whole-cell patch-

clamp (output). GCL: granule cell layer, H: hilus, ML: molecular layer. Solid lines represent

dendrites and dashed lines axons.

(B) Current-clamp recordings of the membrane potential of two different GCs (Ctrl and KA) in

response to the same set of input trains. An input set is constituted of five different trains of

electrical pulses following a Poisson distribution with an average rate of 10 Hz. The Pearson's

correlation coefficient (R) between two input trains is computed with a binning window (τ_w) of 20

ms (*Left*). R_{input} is the average of the ten pairwise coefficients, diagonal excluded. After converting

the GC recordings to vectors of binned spike counts, the pairwise coefficients and their average

(R_{output}) can be computed the same way (Madar et al, 2019b). Note that the input set was repeated

10 times to yield output sets of fifty sweeps (only one repeat is shown).

(C) The R_{output} of GCs from kainate-injected animals is higher, on average, than in GCs from

controls. Data points correspond to single output sets (KA: 27 recordings from 24 GCs; Ctrl: 12

recordings from 11 GCs), all being responses to the same input set as in B. Black crosses with

error bars are means +/- SEM. The asterisk signals significance: U-test comparing the medians: P

= 0.011, Z = -2.541, rank sum = 156; two-sided two-sample KS test comparing the distribution

shapes: P = 0.0037, D = 0.583. The dashed line corresponds to R_{input} (0.77): any value below the

line implies effective pattern separation. Thus, GCs from kainate animals exhibit less pattern

separation than controls.

(D) Pattern separation was investigated over a wide range of input correlations by using four

different input sets of five 10 Hz Poisson trains (at τ_w = 20 ms: R_{input} = 0.24, 0.45, 0.77, 0.95).

Crosses and error bars correspond to mean +/- SEM (pw R_{input}, pw R_{output}) across multiple

recordings (KA: 6-24 GCs per input set; Ctrl: 4-11 GCs). Data points below the identity line

(dashed) correspond to pattern separation. The distributions between KA and Ctrl are significantly

different (ANCOVA with separate parabolic models fitting the data points and not the means: P <

0.0001, F(3,714) = 15.485. Solid curves are the parabolic models used for the ANCOVA.

(E) Levels of pattern separation measured using different similarity metrics (S: R, NDP and SF,

see Methods) and timescales. Cumulative frequency distributions of the distance of (pw S_{input}, pw

S_{output}) data points to the identity line in pattern separation graphs like in E. Positive values of the

x-axis correspond to pattern separation, and negative values to pattern convergence. Insets show

medians. For R and NDP, distributions are significantly shifted to the left, showing that GCs from

KA exhibit less decorrelation and less orthogonalization (ANCOVA as in D, but using separate

linear models: $P < 0.01$ for $\tau_w = 5$ to 1000 ms. For R/NDP and $\tau_w = 10, 20, 100, 250$ ms, $P =$

<0.0001/<0.0001, <0.0001/<0.0001, 0.001/<0.0001, 0.011/0.0001, $F_{(2, 716)} = 15.2/18.2$,

20.3/23.4, 6.87/15.97, 4.5/9.4). For SF, scaling levels are weakly but significantly shifted to less

separation at large τ_w (SF: $P > 0.2$ except at 250 ms and 500 ms, with $P = 0.026, 0.046, F_{(2, 716)}$

$= 3.65, 3.09$ respectively).

Figure 5. Multiple DG computations are affected by epilepsy

Our standard input sets (type 1, Figure 4) consisted of 10 Hz Poisson trains. Two other input sets (type 2 and 3) were designed to explore single GCs responses to inputs with diverse structures and statistics, both with a wide range of pairwise similarity as measured by SF (in contrast to input sets of type 1).

(A1) *Left*: Input set type 2 was constituted of spiketrains following a Poisson distribution, each with a different firing rate (FR), but with an average R_{input} constrained around 0.75 ($\tau_w = 10$ ms). *Middle and Right*: Pattern separation graphs showing the pairwise output spiketrain similarity as a function of the pairwise input similarity, as measured by SF with two different timescales, averaged across multiple GCs (KA: 16, Ctrl: 5). Distributions are significantly different, suggesting in particular that epilepsy causes a decrease in pattern convergence (via scaling) for low input similarities (ANCOVA with separate linear models, for $\tau_w = 20$ and 250 ms respectively : P = 0.0016 and < 0.0001, F(2, 941) = 6.5 and 10.4). **(A2)** Levels of pattern separation for type 2 inputs, measured using different similarity metrics and timescales as in Figure 4. It confirms that epilepsy decreases the separation of similar Poisson input spiketrains as measured by R and NDP, consistent with Figure 4, and shows that differences in terms of SF, although small, are significant (ANCOVA as in A1: R: P < 0.0001 for $\tau_w = 20$ ms up to 1000 ms, P = 0.02 and 0.18 for 5 and 10 ms; NDP and SF: P < 0.025 for τ_w up to 1000 ms. Detailed statistics for $\tau_w = 5, 10, 20, 50, 100, 250, 500$ and 1000 ms respectively: R, P = 0.0243, 0.1820, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, F(2, 941) = 3.7, 1.7, 10.2, 15.1, 24.1, 36.0, 50.2, 27.0; NDP, P = 0.0022, 0.0240, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, F(2, 941) = 6.2, 3.7, 13.8, 17.4, 21.4, 26.0, 21.0, 16.8; SF, P = 0.0002, 0.0012, 0.0016, 0.0240, 0.0073, <0.0001, <0.0001, 0.0002, F(2, 941) = 8.7, 6.7, 6.5, 3.7, 4.9, 10.9, 10.4, 8.7).

(B1) *Left:* Input set type 3 was constituted of spiketrains with 21 spikes (FR = 10.5Hz) that were distributed among bins to produce trains with varying burstiness (Madar et al., 2019b). R is close to 0 for all pairs. *Middle and Right*: same analysis as in A1 (KA: 8 GCs, Ctrl: 3 GCs). Distributions are significantly different, suggesting again that epilepsy causes a decrease in pattern convergence via scaling, for low input similarities (ANCOVA for τ_w = 20 and 100 ms: P < 0.0001 and F(2, 491) = 10.3 and 13.8 respectively). **(B2)** Same analysis as in A2. The directions of impairments are the same as in A2, showing that epilepsy decreases pattern separation in terms of R and NDP but slightly improves it in terms of SF. The shift via scaling is weak but significant at all timescales, and larger at the longest τ_w. ANCOVA: R, P < 0.05 for τ_w = 5, 10 and 500 ms; NDP and SF, P < 0.007 for τ_w up to 500 ms. Detailed statistics for τ_w = 5, 10, 20, 50, 100, 250, 500 and 1000 ms respectively: R, P = 0.0451, 0.0004, 0.3727, 0.7357, 0.9340, 0.4271, 0.0012, 0.2889, F(2, 491) = 3.1, 8.1, 1.0, 0.3, 0.1, 0.8, 6.8, 1.2; NDP, P = <0.0001, <0.0001, 0.0003, 0.0012, <0.0001, 0.0069, <0.0001, 0.4154, F(2, 491) = 11.1, 24.3, 8.4, 6.8, 11.2, 5.0, 14.6, 0.9, SF, P = <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, 0.0065, 0.0002, 0.0003, F(2, 941) = 42.1, 20.9, 10.3, 13.6, 13.8, 5.1, 8.5, 8.1.

Figure 6. In epileptic mice, a subpopulation of GCs shows pathological spiking patterns

(A) Example of current-clamp recordings in GCs from a kainate-injected (KA) vs a control animal in response to the same Poisson input train, illustrating that some GCs from KA spiked more, and had a tendency to fire short bursts (2-4 spikes riding a single EPSP, see inset).

(B-E) Some GCs from KA exhibited larger firing rates due to a higher probability of spiking at least once after an input spike (spiking probability), and sometimes a higher probability of spiking more than once between two input spikes (p(Burst)) than GCs from controls. The firing rate, spiking probability and p(Burst) of a neuron were all computed as the average over the fifty sweeps of an output set from a pattern separation experiment (input set type 1, see Figure 4B).

(B) *Left*: data points correspond to the same recordings as in Figure 4C (KA: 27; Ctrl: 12). Black dash and error bars are mean +/- SEM. A U-test was used to compare the medians, showing that on average FR and SP are higher in KA (FR: P = 0.0415, Z = -2.04, rank sum = 172.5; SP: P =

0.0613, $Z = -1.87$, rank sum $= 168.5$). *Right*: Frequency distributions of the same data. A one-sided

two-sample KS test shows that the KA distribution has a larger tail in both cases (FR and SP: $P =$

0.0465, $D = 0.4074$), indicating that a subset of KA GCs are pathological.

(C) Same as in B for p(Burst). Note that the left graph has a $\log_{10}$ scale showing all data points,

whereas the middle graph has a linear scale zoomed in (i.e. not showing the 5 largest KA values).

The graph on the right has the same scale as the middle graph. Together, they suggest that in KA,

there might be a healthy population of GCs coexisting with a different population of pathologically

bursty GCs (U-test: $P = 0.0153$, $Z = -2.16$, rank sum $= 168.5$; KS test: $P = 0.0259$, $D = 0.4444$).

(D) Some bursts were detected by our algorithm in GCs from Ctrl mice, but the vast majority of

those were due to the temporal summation of two EPSPs resulting from input spikes occurring

close in time. In contrast, a large number of bursts in GCs from KA mice come after an isolated

input spike. *Top*: Time interval between the two input spikes (inter-spike-interval, ISI) preceding

a given burst. Data points correspond to all detected bursts from the recordings in C. U-test: $P =$

0.0003, $Z = -3.58$, rank sum $= 5906.2$. *Bottom*: Cumulative frequency distributions of the same

data. A one-sided two-sample KS test demonstrates that the distributions are different ($P < 0.0001$,

$D = 0.2699$).

(E) Same data as in B and C: the elbow in the distribution of SP, p(Burst) and FR values visually

defines two clusters of neurons. One cluster contains GCs from both KA and Ctrl mice (with GCs

from both groups spanning a large range of SP values but low p(Burst)) and can be considered

"normal". In contrast, the other cluster, with highest SP combined with high p(Burst), contains

only GCs from KA animals and can thus be considered "pathological".

Figure 7. Pathological bursting is not explained by aberrant excitation

(**A**) Bursting and high firing rates were not an artefact of stimulation differences: The current intensity of the electric stimulation did not significantly differ between KA and Ctrl (same recordings as in Figure 4C and 6B-E, U-test: $P = 0.2991$, $Z = 1.04$, rank sum = 274.5) and stimulus intensity was a poor predictor of pathological firing patterns (as indicated by the low R^2 values of linear regressions. For FR/SP/pBurst: $R^2 = 12.75\%/11.69\%/4.35\%$, $F(2,37) = 5.4/4.9/1.7$, $P = 0.0257/0.0332/0.2027$).

(**B**) For a subset of neurons in A (KA: 12; Ctrl: 6), we followed the current-clamp recording with a voltage-clamp recording (V_{hold} = -70mV) in response to the same pattern separation protocol in

order to assess the excitatory synaptic drive and its relationship with spiking behavior. The

stimulus electrode location and current intensity was unchanged. **(B1)** Example of current-clamp

(top) and voltage-clamp (bottom) recording in the same GC from a Ctrl animal. Only the first 500

ms of the ten responses to repetitions of the first input train shown in Figure 4B are displayed. *Top*:

action potentials are truncated at 0 mV. *Bottom*: stimulation artefacts and occasional unclamped

spikes were blanked. **(B2)** The ten EPSCs (grey) associated to first pulse of input trains in B1 and

the mean EPSC (black). We assessed the excitatory drive as the maximum inward current in the

interval between each input pulse minus the current baseline (i.e., mode of the current over the full

sweep). P(Spike) was defined as the probability of spiking during this interval across the ten

sweeps of the current-clamp recording. **(B3)** *Top*: The mean EPSC density for each neuron was

not significantly different between KA and Ctrl (U-test: $P = 0.5532$, rank sum $= 50$; two-sided

two-sample KS test: $P = 0.1935$, $D = 0.4167$), suggesting that pathological spiking cannot be

explained by larger EPSCs in KA mice. *Bottom*: Indeed, no clear relationship exists between

p(Burst) and the mean current density, as some GCs with a great synaptic drive do not exhibit

much bursting, and vice versa. **(B4)** Consistently, the peak current density associated to each inter-

input-interval in a recording is not well correlated to the corresponding number of spikes in the

current-clamp trace, for both Ctrl (top) and KA (bottom). For example, excitatory currents of the

same magnitude can be associated to 0, 1, 2, 3 or 4 spikes. EPSC amplitude is thus not sufficient

to explain the spiking output, which suggests that other factors, like reduced inhibition, are likely

implicated in causing pathological bursts. **(B5)** *Top*: The average EPSC density between two input

spikes is plotted against the corresponding probability of spiking p(Spike), as in Ewell and Jones

(2010). The resulting input-output (IO) distribution is fitted with a sigmoid (two GCs

representative of KA and Ctrl are shown). *Bottom*: In a subset of GCs, mostly from KA, The IO

distribution appears disorganized and sometimes difficult to fit. A proxy of this disorganization is

the standard deviation (SD) of the current density, averaged across each level of p(Spike)

excluding p(Spike) = 0 and 1. The average SD for a given GC was plotted against the propensity

of that neuron to fire bursts. There is no clear relationship between these two quantities, suggesting

that: 1) decoupling between the excitatory drive and spiking probability is not directly related to

bursting and 2) a subset of GCs from KA with apparently healthy spiking patterns could already

exhibit a more subtle but pathological IO disorganization.

Figure 8. Pathological spiking patterns explain pattern separation differences in epilepsy

(A) Neurons with high average firing rates and burstiness exhibit lower pattern separation

(computed as $R_{input} - R_{output}$). Same data as in Figure 4C and 6B-C. There is a strong linear

relationship between firing rates and decorrelation (Grey line, $R^2 = 63.68\%$, $F(2,37) = 64.9$, $P <$

0.0001). The relationship between p(Burst) and decorrelation is better described by an inverse

function with a horizontal asymptote (see Methods – Software and statistics): $R^2 = 53.1\%$, R^2

adjusted for 3 parameters = 50.5%, $F(3,36) = 20.4$, $P < 0.0001$.

(B-D) Mean +/- SEM pattern separation levels across recordings when considering burstiness

codes (Compactness or Occupancy) or a rate code (spiking frequency across a 2s sweep) instead

of the binwise synchrony code assumed by R. Compactness is the proportion of time bins with at

least one spike, whereas occupancy is the average number of spikes in a bin. Pattern separation

corresponds to more dispersion in compactness, occupancy or firing rate in the output spiketrains

than the input spiketrains. Negative values mean there is pattern convergence. U-tests comparing

medians of GCs from KA and Ctrl groups were performed (p-values in panel).

(B) In response to input sets of type 1 (10 Hz Poisson trains, same recordings as in A and Figure

4), GCs exhibit low levels of convergence, if any, in terms of compactness, occupancy and rate

codes, and these levels are slightly but significantly shifted to less convergence in GCs from KA

for $\tau_w = 10$ ms. This means that the relationships between firing rate or burstiness and pattern

decorrelation observed in A is not due to variations of the firing rate or burstiness across sweeps.

Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: $P = 0.0482$, .1892, .1628, .2925, .7584,

$Z = -1.97, -1.97, -1.31, -1.39, 0.31$, rank sum = 745, 745, 801, 794, 939 ; Occupancy: $P = 0.0177$,

0.2586, 0.3624, 0.1425, 0.0211, $Z = -2.37, -1.13, -0.91, -1.47, -2.31$, rank sum = 712.5, 816.5, 835,

788, 717; FR: $P = 0.0553$, $Z = -1.92$, rank sum = 750.

(C) In response to an input set of type 2 (Poisson trains with different firing rates, same recordings

as in Figure 5A), GCs from KA and Ctrl do not show significant differences (P > 0.2 for all

timescales and codes). Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: P = 0.7726,

0.7726, 0.8365, 0.9014, 0.2312, Z = -0.29, -0.29, -0.21, 0.12, 1.20, rank sum = 51, 51, 52, 57, 70;

Occupancy: P = 1, 1, 0.9671, 0.5915, 0.2006, Z = 0, 0, 0.04, -0.54, -1.28, rank sum = 55, 55, 56,

48, 39; FR: P = 0.9014, Z = -0.12, rank sum = 53.

(D) In response to an input set of type 3 (10 Hz trains with varying compactness and occupancy,

same recordings as in Figure 5B), GCs exhibit convergence in terms of compactness and

occupancy and separation in terms of rate codes. Only computations in terms of compactness and

firing rate are mildly shifted in GCs from KA for τ_w = 10 ms. Detailed statistics at 10, 20, 50, 100,

250 ms, Compactness: P = 0.0848, rank sum = 9; Occupancy: P = 0.8, 0.7758, 0.9212, 0.3758, 1,

rank sum = 19.5, 20, 17, 23, 18; FR: P = 0.0848, rank sum = 9.

Figure 9. EEG, behavioral and computational measures provide complementary insights about epileptic pathology in individual animals.

(A-D) Scatter plots relating the electrographic measure of IISs developed in Figure 2 (black axis labels), measures from patch-clamp recordings of individual GCs from the same animals (purple axis labels) and behavioral discrimination measures described in Figure 3 (green axis labels). Not all three types of records are available for every animal, which leads to differences in the samples shown in each panel.

(A) Data points correspond to individual GCs from animals in which both EEG and patch-clamp data were recorded. Each animal has a unique average hourly IIS index value. Only a subset of GCs are outside of the Ctrl range (i.e. pathological). For example, in the mouse with the highest IIS index (~12), only one of the three recorded GCs exhibited a pathological FR (~8 Hz, left panel)

and all had a similar R_{output} slightly above the Ctrl range (~0.5, right panel). In contrast, some GCs

with abnormally high FR (~6-12 Hz)_and R_{output} (~0.5-0.6) came from animals with low IIS index

(~0-2).

(B) Data points correspond to individual animals in which both EEG and behavior were measured.

Discrimination Impairment is the Mahalanobis distance computed in Figure 3C. In KA animals,

low IIS index can correspond to either normal or abnormal mnemonic discrimination, and despite

having a high IIS index, some animals have normal discrimination.

(C) Data points correspond to individual GCs from animals in which both behavior and patch-

clamp data were recorded. *Left*: Significant linear correlation (grey line) between behavioral

discrimination after a 14 cm object displacement (Figure 3B) and pattern separation in DG (R^2 =

0.0355, F(1,21) = 11.02, P = 0.0034). No obvious relationship for the other conditions (7 cm and

21 cm). *Right*: Performance on all displacement distances are taken in account. The relationship is

unclear, but all KA mice with large impairments had at least one pathological GC.

(D) Data points correspond to individual GCs from animals in which EEG, behavior and patch-

clamp data were all collected. Stems show which GCs were recorded from each animal. The

lower right corner would correspond to the highest pathology in electrographic, behavioral and

computational categories. Despite any heterogeneity or apparent ambiguity when only one or

two of these measures are considered (above), when all three measures are available for each

animal, the KA group is very clearly separable from the Ctrl group.

References

Alexander A, Maroso M, Soltesz I (2016) Organization and control of epileptic circuits in temporal lobe

epilepsy. Prog Brain Res 226:127–154.

Artinian J, Peret A, Marti G, Epsztein J, Crépel V (2011) Synaptic kainate receptors in interplay with INaP

shift the sparse firing of dentate granule cells to a sustained rhythmic mode in temporal lobe

epilepsy. J Neurosci Off J Soc Neurosci 31:10811–10818.

Baker S, Vieweg P, Gao F, Gilboa A, Wolbers T, Black SE, Rosenbaum RS (2016) The Human Dentate

Gyrus Plays a Necessary Role in Discriminating New Memories. Curr Biol CB 26:2629–2634.

Behr J, Lyson KJ, Mody I (1998) Enhanced propagation of epileptiform activity through the kindled

dentate gyrus. J Neurophysiol 79:1726–1732.

Bennett IJ, Stark CEL (2016) Mnemonic discrimination relates to perforant path integrity: An ultra-high

resolution diffusion tensor imaging study. Neurobiol Learn Mem 129:107–112.

Bernier BE, Lacagnina AF, Ayoub A, Shue F, Zemelman BV, Krasne FB, Drew MR (2017) Dentate Gyrus

Contributes to Retrieval as well as Encoding: Evidence from Context Fear Conditioning, Recall, and

Extinction. J Neurosci Off J Soc Neurosci 37:6359–6371.

Berron D, Schütze H, Maass A, Cardenas-Blanco A, Kuijf HJ, Kumaran D, Düzel E (2016) Strong Evidence

for Pattern Separation in Human Dentate Gyrus. J Neurosci Off J Soc Neurosci 36:7569–7579.

Bielefeld P, van Vliet EA, Gorter JA, Lucassen PJ, Fitzsimons CP (2014) Different subsets of newborn

granule cells: a possible role in epileptogenesis? Eur J Neurosci 39:1–11.

Braganza O, Mueller-Komorowska D, Kelly T, Beck H (2020) Quantitative properties of a feedback circuit

predict frequency-dependent pattern separation. eLife 9.

Bui AD, Nguyen TM, Limouse C, Kim HK, Szabo GG, Felong S, Maroso M, Soltesz I (2018) Dentate gyrus

mossy cells control spontaneous convulsive seizures and spatial memory. Science 359:787–790.

Cayco-Gajic NA, Silver RA (2019) Re-evaluating Circuit Mechanisms Underlying Pattern Separation.

Neuron 101:584–602.

Chauvière L, Doublet T, Ghestem A, Siyoucef SS, Wendling F, Huys R, Jirsa V, Bartolomei F, Bernard C

(2012) Changes in interictal spike features precede the onset of temporal lobe epilepsy. Ann Neurol

71:805–814.

Chauvière L, Rafrafi N, Thinus-Blanc C, Bartolomei F, Esclapez M, Bernard C (2009) Early deficits in spatial

memory and theta rhythm in experimental temporal lobe epilepsy. J Neurosci Off J Soc Neurosci

29:5402–5410.

Chavlis S, Poirazi P (2017) Pattern separation in the hippocampus through the eyes of computational

modeling. Synap N Y N 71.

Coras R, Pauli E, Li J, Schwarz M, Rössler K, Buchfelder M, Hamer H, Stefan H, Blumcke I (2014)

Differential influence of hippocampal subfields to memory formation: insights from patients with

temporal lobe epilepsy. Brain J Neurol 137:1945–1957.

Coulter DA, Carlson GC (2007) Functional regulation of the dentate gyrus by GABA-mediated inhibition.

Prog Brain Res 163:235–243.

de la Prida LM, Huberfeld G, Cohen I, Miles R (2006) Threshold behavior in the initiation of hippocampal

population bursts. Neuron 49:131–142.

De Maesschalck R, Jouan-Rimbaud D, Massart DL (2000) The Mahalanobis distance. Chemom Intell Lab

Syst 50:1–18.

Dengler CG, Coulter DA (2016) Normal and epilepsy-associated pathologic function of the dentate gyrus.

Prog Brain Res 226:155–178.

Dengler CG, Yue C, Takano H, Coulter DA (2017) Massively augmented hippocampal dentate granule cell

activation accompanies epilepsy development. Sci Rep 7:42090.

Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L (2016) Low excitatory

innervation balances high intrinsic excitability of immature dentate neurons. Nat Commun 7:11313.

Dillon SE, Tsivos D, Knight M, McCann B, Pennington C, Shiel AI, Conway ME, Newson MA, Kauppinen

RA, Coulthard EJ (2017) The impact of ageing reveals distinct roles for human dentate gyrus and CA3

in pattern separation and object recognition memory. Sci Rep 7:14069.

Engin E, Zarnowska ED, Benke D, Tsvetkov E, Sigal M, Keist R, Bolshakov VY, Pearce RA, Rudolph U (2015)

Tonic Inhibitory Control of Dentate Gyrus Granule Cells by α5-Containing GABAA Receptors Reduces

Memory Interference. J Neurosci Off J Soc Neurosci 35:13698–13712.

Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2020) A Sparse, Spatially Biased

Subtype of Mature Granule Cell Dominates Recruitment in Hippocampal-Associated Behaviors. Cell

Rep 31:107551.

Ewell LA, Jones MV (2010) Frequency-tuned distribution of inhibition in the dentate gyrus. J Neurosci Off

J Soc Neurosci 30:12597–12607.

Gilbert PE, Kesner RP, Lee I (2001) Dissociating hippocampal subregions: double dissociation between

dentate gyrus and CA1. Hippocampus 11:626–636.

Gröticke I, Hoffmann K, Löscher W (2008) Behavioral alterations in a mouse model of temporal lobe

epilepsy induced by intrahippocampal injection of kainate. Exp Neurol 213:71–83.

Guise KG, Shapiro ML (2017) Medial Prefrontal Cortex Reduces Memory Interference by Modifying

Hippocampal Encoding. Neuron 94:183-192.e8.

Hainmueller T, Bartos M (2020) Dentate gyrus circuits for encoding, retrieval and discrimination of

episodic memories. Nat Rev Neurosci 21:153–168.

Hellier JL, Patrylo PR, Buckmaster PS, Dudek FE (1998) Recurrent spontaneous motor seizures after

repeated low-dose systemic treatment with kainate: assessment of a rat model of temporal lobe

epilepsy. Epilepsy Res 31:73–84.

Helmstaedter C, Kurthen M, Lux S, Reuber M, Elger CE (2003) Chronic epilepsy and cognition: a

longitudinal study in temporal lobe epilepsy. Ann Neurol 54:425–432.

Henze DA, Wittner L, Buzsáki G (2002) Single granule cells reliably discharge targets in the hippocampal

CA3 network in vivo. Nat Neurosci 5:790–795.

Hsu D (2007) The dentate gyrus as a filter or gate: a look back and a look ahead. Prog Brain Res 163:601–

613.

Huneau C, Benquet P, Dieuset G, Biraben A, Martin B, Wendling F (2013) Shape features of epileptic

spikes are a marker of epileptogenesis in mice. Epilepsia 54:2219–2227.

Hunsaker MR, Kesner RP (2013) The operation of pattern separation and pattern completion processes

associated with different attributes or domains of memory. Neurosci Biobehav Rev 37:36–58.

Inostroza M, Brotons-Mas JR, Laurent F, Cid E, de la Prida LM (2013) Specific impairment of "what-

where-when" episodic-like memory in experimental models of temporal lobe epilepsy. J Neurosci

Off J Soc Neurosci 33:17749–17762.

Jones JE, Watson R, Sheth R, Caplan R, Koehn M, Seidenberg M, Hermann B (2007) Psychiatric

comorbidity in children with new onset epilepsy. Dev Med Child Neurol 49:493–497.

Kelly T, Beck H (2017) Functional properties of granule cells with hilar basal dendrites in the epileptic

dentate gyrus. Epilepsia 58:160–171.

Kesner RP, Kirk RA, Yu Z, Polansky C, Musso ND (2016) Dentate gyrus supports slope recognition

memory, shades of grey-context pattern separation and recognition memory, and CA3 supports

pattern completion for object memory. Neurobiol Learn Mem 129:29–37.

Kesner RP, Rolls ET (2015) A computational theory of hippocampal function, and tests of the theory:

new developments. Neurosci Biobehav Rev 48:92–147.

Knierim JJ, Neunuebel JP (2016) Tracking the flow of hippocampal computation: Pattern separation,

pattern completion, and attractor dynamics. Neurobiol Learn Mem 129:38–49.

Kobayashi M, Buckmaster PS (2003) Reduced inhibition of dentate granule cells in a model of temporal

lobe epilepsy. J Neurosci Off J Soc Neurosci 23:2440–2452.

Kron MM, Zhang H, Parent JM (2010) The developmental stage of dentate granule cells dictates their

contribution to seizure-induced plasticity. J Neurosci Off J Soc Neurosci 30:2051–2059.

Krook-Magnuson E, Armstrong C, Bui A, Lew S, Oijala M, Soltesz I (2015) In vivo evaluation of the

dentate gate theory in epilepsy. J Physiol 593:2379–2388.

Le Duigou C, Simonnet J, Teleñczuk MT, Fricker D, Miles R (2014) Recurrent synapses and circuits in the

CA3 region of the hippocampus: an associative network. Front Cell Neurosci 7:262.

Lee C-H, Lee I (2020) Impairment of Pattern Separation of Ambiguous Scenes by Single Units in the CA3

in the Absence of the Dentate Gyrus. J Neurosci Off J Soc Neurosci 40:3576–3590.

Lee C-T, Kao M-H, Hou W-H, Wei Y-T, Chen C-L, Lien C-C (2016) Causal Evidence for the Role of Specific

GABAergic Interneuron Types in Entorhinal Recruitment of Dentate Granule Cells. Sci Rep 6:36885.

Lenck-Santini P-P, Scott RC (2015) Mechanisms Responsible for Cognitive Impairment in Epilepsy. Cold

Spring Harb Perspect Med 5.

Lévesque M, Avoli M (2013) The kainic acid model of temporal lobe epilepsy. Neurosci Biobehav Rev

37:2887–2899.

Lévesque M, Avoli M, Bernard C (2016) Animal models of temporal lobe epilepsy following systemic

chemoconvulsant administration. J Neurosci Methods 260:45–52.

Lilliefors HW (1967) On the Kolmogorov-Smirnov Test for Normality with Mean and Variance Unknown. J

Am Stat Assoc 62:399–402.

Liu KY, Gould RL, Coulson MC, Ward EV, Howard RJ (2016) Tests of pattern separation and pattern

completion in humans-A systematic review. Hippocampus 26:705–717.

Lu Y, Zhong C, Wang L, Wei P, He W, Huang K, Zhang Y, Zhan Y, Feng G, Wang L (2016) Optogenetic

dissection of ictal propagation in the hippocampal-entorhinal cortex structures. Nat Commun

7:10962.

Lynch M, Sayin U, Golarai G, Sutula T (2000) NMDA receptor-dependent plasticity of granule cell spiking

in the dentate gyrus of normal and epileptic rats. J Neurophysiol 84:2868–2879.

Madar AD, Ewell LA, Jones MV (2019a) Pattern separation of spiketrains in hippocampal neurons. Sci

Rep 9:5282.

Madar AD, Ewell LA, Jones MV (2019b) Temporal pattern separation in hippocampal neurons through

multiplexed neural codes. PLoS Comput Biol 15:e1006932.

Marrone DF, Adams AA, Satvat E (2011) Increased pattern separation in the aged fascia dentata.

Neurobiol Aging 32:2317.e23-32.

McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB, Fanselow MS, Wilson

MA, Tonegawa S (2007) Dentate gyrus NMDA receptors mediate rapid pattern separation in the

hippocampal network. Science 317:94–99.

Mongiat LA, Espósito MS, Lombardi G, Schinder AF (2009) Reliable activation of immature neurons in the

adult hippocampus. PloS One 4:e5320.

Motulsky HJ, Ransnas LA (1987) Fitting curves to data using nonlinear regression: a practical and

nonmathematical review. FASEB J Off Publ Fed Am Soc Exp Biol 1:365–374.

Müller CJ, Gröticke I, Bankstahl M, Löscher W (2009) Behavioral and cognitive alterations, spontaneous

seizures, and neuropathology developing after a pilocarpine-induced status epilepticus in C57BL/6

mice. Exp Neurol 219:284–297.

O'Reilly RC, McClelland JL (1994) Hippocampal conjunctive encoding, storage, and recall: avoiding a

trade-off. Hippocampus 4:661–682.

Ouedraogo DW, Lenck-Santini P-P, Marti G, Robbe D, Crépel V, Epsztein J (2016) Abnormal UP/DOWN

Membrane Potential Dynamics Coupled with the Neocortical Slow Oscillation in Dentate Granule

Cells during the Latent Phase of Temporal Lobe Epilepsy. eNeuro 3.

Pardi MB, Ogando MB, Schinder AF, Marin-Burgin A (2015) Differential inhibition onto developing and

mature granule cells generates high-frequency filters with variable gain. eLife 4:e08764.

Patrylo PR, Schweitzer JS, Dudek FE (1999) Abnormal responses to perforant path stimulation in the

dentate gyrus of slices from rats with kainate-induced epilepsy and mossy fiber reorganization.

Epilepsy Res 36:31–42.

Pernía-Andrade AJ, Jonas P (2014) Theta-gamma-modulated synaptic currents in hippocampal granule

cells in vivo define a mechanism for network oscillations. Neuron 81:140–152.

Pfammatter JA, Bergstrom RA, Wallace EP, Maganti RK, Jones MV (2018) A predictive epilepsy index

based on probabilistic classification of interictal spike waveforms. PLOS ONE 13:e0207158.

Poch C, Toledano R, García-Morales I, Prieto A, García-Barragán N, Aledo-Serrano Á, Gil-Nagel A, Campo

P (2020) Mnemonic discrimination in patients with unilateral mesial temporal lobe epilepsy relates

to similarity and number of events stored in memory. Neurobiol Learn Mem 169:107177.

Pofahl M, Nikbakht N, Haubrich AN, Nguyen T, Masala N, Braganza O, Macke JH, Ewell LA, Golcuk K, Beck

H (2020) Dentate gyrus population activity during immobility drives formation of precise memories.

bioRxiv:2020.03.05.978320.

Ramirez S, Liu X, Lin P-A, Suh J, Pignatelli M, Redondo RL, Ryan TJ, Tonegawa S (2013) Creating a false

memory in the hippocampus. Science 341:387–391.

Rayner G, Tailby C, Jackson G, Wilson S (2019) Looking beyond lesions for causes of neuropsychological

impairment in epilepsy. Neurology 92:e680–e689.

Reyes A, Holden HM, Chang Y-HA, Uttarwar VS, Sheppard DP, DeFord NE, DeJesus SY, Kansal L, Gilbert

PE, McDonald CR (2018) Impaired spatial pattern separation performance in temporal lobe epilepsy

is associated with visuospatial memory deficits and hippocampal volume loss. Neuropsychologia

111:209–215.

Rolls ET (2010) A computational theory of episodic memory formation in the hippocampus. Behav Brain

Res 215:180–196.

Santhakumar V, Aradi I, Soltesz I (2005) Role of mossy fiber sprouting and mossy cell loss in

hyperexcitability: a network model of the dentate gyrus incorporating cell types and axonal

topography. J Neurophysiol 93:437–453.

Santoro A (2013) Reassessing pattern separation in the dentate gyrus. Front Behav Neurosci 7:96.

Scimemi A, Schorge S, Kullmann DM, Walker MC (2006) Epileptogenesis is associated with enhanced

glutamatergic transmission in the perforant path. J Neurophysiol 95:1213–1220.

Sharma S, Puttachary S, Thippeswamy A, Kanthasamy AG, Thippeswamy T (2018) Status Epilepticus:

Behavioral and Electroencephalography Seizure Correlates in Kainate Experimental Models. Front

Neurol 9:7.

Simon P, Dupuis R, Costentin J (1994) Thigmotaxis as an index of anxiety in mice. Influence of

dopaminergic transmissions. Behav Brain Res 61:59–64.

Sloviter RS (1983) "Epileptic" brain damage in rats induced by sustained electrical stimulation of the

perforant path. I. Acute electrophysiological and light microscopic studies. Brain Res Bull 10:675–

697.

Stark SM, Kirwan CB, Stark CEL (2019) Mnemonic Similarity Task: A Tool for Assessing Hippocampal

Integrity. Trends Cogn Sci 23:938–951.

Stark SM, Yassa MA, Lacy JW, Stark CEL (2013) A task to assess behavioral pattern separation (BPS) in

humans: Data from healthy aging and mild cognitive impairment. Neuropsychologia 51:2442–2449.

Toyoda I, Bower MR, Leyva F, Buckmaster PS (2013) Early activation of ventral hippocampus and

subiculum during spontaneous seizures in a rat model of temporal lobe epilepsy. J Neurosci Off J Soc

Neurosci 33:11100–11115.

Traub RD, Wong RK (1982) Cellular mechanism of neuronal synchronization in epilepsy. Science

216:745–747.

Treves A, Tashiro A, Witter MP, Moser EI (2008) What is the mammalian dentate gyrus good for?

Neuroscience 154:1155–1172.

van Hagen BTJ, van Goethem NP, Lagatta DC, Prickaerts J (2015) The object pattern separation (OPS)

task: a behavioral paradigm derived from the object recognition task. Behav Brain Res 285:44–52.

Wallace E, Kim DY, Kim K-M, Chen S, Blair Braden B, Williams J, Jasso K, Garcia A, Rho JM, Bimonte-

Nelson H, Maganti R (2015) Differential effects of duration of sleep fragmentation on spatial

learning and synaptic plasticity in pubertal mice. Brain Res 1615:116–128.

White A, Williams PA, Hellier JL, Clark S, Dudek FE, Staley KJ (2010) EEG spike activity precedes epilepsy

after kainate-induced status epilepticus. Epilepsia 51:371–383.

Witt J-A, Helmstaedter C (2015) Cognition in the early stages of adult epilepsy. Seizure 26:65–68.

Woods NI, Stefanini F, Apodaca-Montano DL, Tan IMC, Biane JS, Kheirbek MA (2020) The Dentate Gyrus

Classifies Cortical Representations of Learned Stimuli. Neuron.

Yassa MA, Mattfeld AT, Stark SM, Stark CEL (2011) Age-related memory deficits linked to circuit-specific

disruptions in the hippocampus. Proc Natl Acad Sci U S A 108:8873–8878.

Yi F, Catudio-Garrett E, Gábriel R, Wilhelm M, Erdelyi F, Szabo G, Deisseroth K, Lawrence J (2015)

Hippocampal "cholinergic interneurons" visualized with the choline acetyltransferase promoter:

anatomical distribution, intrinsic membrane properties, neurochemical characteristics, and capacity

for cholinergic modulation. Front Synaptic Neurosci 7:4.

Yim MY, Hanuschkin A, Wolfart J (2015) Intrinsic rescaling of granule cells restores pattern separation

ability of a dentate gyrus network model during epileptic hyperexcitability. Hippocampus 25:297–

308.

Yokoyama M, Matsuo N (2016) Loss of Ensemble Segregation in Dentate Gyrus, but not in

Somatosensory Cortex, during Contextual Fear Memory Generalization. Front Behav Neurosci

10:218.

Zhang W, Huguenard JR, Buckmaster PS (2012) Increased excitatory synaptic input to granule cells from

hilar and CA3 regions in a rat model of temporal lobe epilepsy. J Neurosci Off J Soc Neurosci

32:1183–1196.

Zhao F, Kang H, You L, Rastogi P, Venkatesh D, Chandra M (2014) Neuropsychological deficits in

temporal lobe epilepsy: A comprehensive review. Ann Indian Acad Neurol 17:374–382.

1111