Title: Deficits in Behavioral and Neuronal Pattern Separation in Temporal Lobe Epilepsy

Abbreviated Title: Pattern Separation and Mnemonic Discrimination in Epilepsy

Authors and Affiliations: A.D. Madar[1,2], J.A. Pfammatter[1], J. Bordenave[3], E.I. Plumley[3], S. Ravi[1], M. Cowie[1], E.P. Wallace[1,3], B.P. Hermann[3], R.K. Maganti[3], M.V. Jones[1]

[1]Department of Neuroscience, University of Wisconsin-Madison, 53703

[2]Department of Neurobiology, Grossman Institute for Neuroscience, Quantitative Biology and Human Behavior, University of Chicago, 60637

[3]Department of Neurology, University of Wisconsin-Madison, School of Medicine and Public Health

Acknowledgement: We thank Lin Lin who helped start behavioral tests in mice in the Jones lab and Drs. C.E.L Stark and M.A. Yassa for allowing us to use software and materials they developed.

Author contributions: Conceptualization: MVJ, ADM. Data curation: ADM, JAP. Formal analysis: ADM, JAP, MVJ. Funding acquisition: MVJ, ADM, RKM, JAP. Investigation: ADM, JAP, JB, EIP, SR, MC, EPW. Methodology: ADM, JAP, MVJ. Project administration: JAP, ADM, MVJ. Resources: MVJ, BPH, RKM. Software: ADM, JAP, MVJ. Supervision: MVJ. Validation: JAP, ADM. Visualization: ADM, JAP, MVJ. Writing – original draft: ADM. Writing – review & editing: all authors

Funding: This work was supported by the US National Institutes of Health (MVJ: RO1 NS075366), the University of Wisconsin Institute for Clinical and Translational Research (MVJ.; NIH/NCATS UL1TR000427), the US Department of Defense (RKM: PR161864) and Lily's Fund for Epilepsy Research (ADM.; 2015 fellow).

Conflicts of Interest: The authors declare no conflict of interest.

Abstract:

In temporal lobe epilepsy, the ability of the dentate gyrus to limit excitatory cortical input to the hippocampus breaks down, leading to seizures. The dentate gyrus is also thought to help discriminate between similar memories by performing pattern separation, but whether epilepsy leads to a breakdown in this neural computation, and thus to mnemonic discrimination impairments, remains unknown. Here we show that temporal lobe epilepsy is characterized by behavioral deficits in mnemonic discrimination tasks, in both humans and mice. Using a recently developed assay in brain slices of the same epileptic mice, we reveal a decreased ability of the dentate gyrus to perform certain forms of pattern separation. This is due to a subset of granule cells with abnormal bursting that can develop independently of early EEG abnormalities. Overall, our results linking physiology, computation and cognition in the same mice, advance our understanding of episodic memory mechanisms and their dysfunction in epilepsy.

Keywords:

 Patch-clamp, low-dose systemic kainate model**,** kainic acid, interictal spikes, novelty recognition, firing rate, spiking patterns, similarity metrics, input-output, entorhinal cortex, perforant path, EPSC, excitation/inhibition ratio, dentate gate, Hebb-Marr theory

Introduction:

Temporal lobe epilepsy (TLE) represents about 60% of all epilepsy cases, a third of which are refractory to medication (Tellez-Zenteno and Hernandez-Ronquillo, 2012). TLE is characterized by recurring focal seizures originating in or near the hippocampus (Toyoda et al., 2013), microcircuit pathologies in brain regions including the hippocampus (Alexander et al., 2016) and memory-related cognitive deficits (Helmstaedter et al., 2003; Zhao et al., 2014). Although the hippocampus is a nexus for episodic memory that is critically affected during TLE, the relationship between TLE and hippocampus-dependent memory is insufficiently understood.

Episodic memory formation is thought to involve storage of neural representations in area CA3 of the hippocampus via Hebbian plasticity at recurrent excitatory synapses of coactivated cells (Rolls, 2010). In this view, a partial cue reactivates a subset of the CA3 ensemble that in turn recruits the other neurons of the original pattern resulting in recall of the original event (Rolls, 2010). However, recurrent excitation is problematic because it theoretically a) predisposes the network to over-excitation that could trigger seizures (Le Duigou et al., 2014) and b) limits the number of patterns that can be stored without overlap (Rolls, 2010). Overlapping memory representations would in turn lead to interference during recall and thus cognitive confusion. To solve these problems, it was proposed that the dentate gyrus (DG) of the hippocampus acts as a) a gate and b) a pattern separator, so that similar cortical representations are transformed into sparse and dissimilar patterns before reaching CA3 (O'Reilly and McClelland, 1994; Hsu, 2007; Treves et al., 2008; Dengler and Coulter, 2016).

The function of DG as a gate for cortical activity to prevent seizure generation fails in TLE (Hsu, 2007; Krook-Magnuson et al., 2015; Dengler and Coulter, 2016; Lu et al., 2016): granule cells (GCs), the output neurons of DG, lose their usual sparseness due to network

reorganization (Artinian et al., 2011; Dengler and Coulter, 2016; Dengler et al., 2017) making

cortical excitation easier to propagate to CA3 (Behr et al., 1998; Patrylo et al., 1999; Ouedraogo

et al., 2016). Even in healthy animals, repeated excitation of GCs induces seizures (Krook-

Magnuson et al., 2015) and prolonged stimulation of DG causes TLE (Sloviter, 1983).

An alternate view of the DG function, developed in parallel to the concept of dentate

gate, is that it performs pattern separation: the transformation of similar cortical patterns into

dissimilar hippocampal representations. This process theoretically supports mnemonic

discrimination, the ability to distinguish between similar memories. Indeed, DG lesions impair

mnemonic discrimination both in rodents (Treves et al., 2008; Kesner and Rolls, 2015; Kesner et

al., 2016) and humans (Yassa et al., 2011; Baker et al., 2016; Bennett and Stark, 2016; Dillon et

al., 2017). Moreover, computational models (Chavlis and Poirazi, 2017) and recent experiments

(Knierim and Neunuebel, 2016; Berron et al., 2016; Madar et al., 2019a, b) suggest that the DG

circuitry supports multiple forms of pattern separation (Santoro, 2013). Whether such

computations underlie mnemonic discrimination remains unknown.

Unsurprisingly, TLE negatively impacts hippocampal-dependent memory in humans

(Coras et al., 2014) and rodents (Groticke et al., 2008; Muller et al., 2009; Inostroza et al., 2013;

Lenck-Santini and Scott, 2015). However, the effect of TLE specifically on DG computations

and DG-dependent cognition remains understudied. It was only recently reported that patients

with TLE are impaired at spatial mnemonic discrimination (Reyes et al., 2018) and that TLE

causes deficits in DG-dependent object location memory in mice (Bui et al., 2018). A

computational model has also suggested that hippocampal pathologies in TLE would degrade

DG pattern separation (Yim et al., 2014) but the hypotheses that 1) TLE causes a breakdown in

DG neural pattern separation and 2) that such a failure causes mnemonic discrimination

impairments remain experimentally untested.

Here we tested, in humans and mice, whether TLE is characterized by deficits in

mnemonic discrimination and then recorded, in brain slices from the same mice, the spiking

patterns of single GCs in response to parametrically varied afferent stimulation in order to gauge

neuronal pattern separation.

Materials and Methods:

Human Behavior. A mnemonic similarity task (Stark et al., 2019), also known as a

behavioral pattern separation (BPS) task (Stark et al., 2013), was administered to 15 patients in

the University of Wisconsin-Madison Epilepsy Monitoring Unit. Under an approved Institutional

Review Board protocol and after obtaining informed consent, the task was administered in

conjunction with a standard neuropsychiatric evaluation and during electroencephalographic

(EEG) recording, both of which are part of standard practice for diagnosing the patients' seizures.

Patients were 18-65 years old, male and female. Only patients with a preliminary diagnosis of

TLE were included in our analysis. As controls, 20 subjects without epilepsy, recruited to match

the patients' age and sex distributions (family members of the patients when possible, or other

volunteers), were also tested. Consent documents, medical records and primary data are on file in

a secure location within the Dept. of Neurology. Data were deidentified prior to analysis.

Subjects received no compensation for participation.

The visual, object recognition-based BPS task is described in Yassa et al. (2011) and has

been further validated by demonstrating mnemonic discrimination deficits during normal aging

and multiple neurologic and psychiatric disorders (Stark et al., 2019). It was implemented on a

laptop by trained neuropsychiatric postdoctoral fellows (J.B. and E.I.P.) using software

distributed freely by the Stark lab (http://faculty.sites.uci.edu/starklab/mnemonic-similarity-task-

mst/). Briefly, participants viewed a series of pictures of everyday objects (**Figure 1A**). During

Phase 1, 128 different images were presented and the subject was asked to classify each as

"indoor" or "outdoor", simply to engage the subject's attention. During Phase 2, a new series of

192 images was presented: 1/3 repeated from Phase 1 (*Repeated*), 1/3 new but similar to images

from Phase 1 (*Lure*), and 1/3 new and completely different from Phase 1 (*Novel*). Participants

were asked to classify each image of the Phase 2 set as "Old", "Similar", or "New". BPS was

evaluated with a *discrimination index* computed as the difference between p("Similar"|Lure) and

p("Similar"|Novel), as in past research (Yassa et al., 2011; Stark et al., 2019).

Mouse Experiments. Male mice (C57BL6J, 5-6 weeks old) were received from Envigo

(formerly Harlan, Madison, WI), housed in groups and allowed to acclimate to their new home

environment for one or two weeks. Animals then underwent epilepsy induction with kainic acid

(KA) injections (see below) (J.A.P.). A control group of mice was injected with saline and

another received no injection: both were pooled together and considered as the control group

because our analyses did not reveal any difference. 7-9 weeks after injection, mice started 4

weeks of behavioral testing on the BPS task. After completion of behavioral testing, animals

(~18-20 weeks old) were transferred to a different building for EEG implantation. Each animal

was recorded continuously for three days before being transferred back to the original building

and sacrificed for slice electrophysiology (~20-24 weeks old). After each building-to-building

transfer, mice were allowed a period of acclimation ranging from two days to two weeks.

Behavioral testing (J.A.P, S.R. and M.C.), EEG recordings (E.P.W.) and slice electrophysiology

(A.D.M) were performed by different researchers, all of whom were blind to KA or Control

treatments.

Epilepsy induction. During the induction process, when not being handled, mice were

individually housed in enclosed ~150 cm^3 acrylic cubicles with opaque sides and clear front

portals with holes to allow air exchange, food and bedding. Animals were then randomly

assigned to KA or Control treatment, ear punched for identification and weighed. Animals were

induced for epileptogenesis using the repeated low-dose kainate method (Hellier et al., 1998;

Sharma et al., 2018): a series of intraperitoneal injections based on the following schedule.

Animals first received 10 mg/kg of kainic acid (Tocris Bioscience, UK) mixed in 1x PBS (8 ml

per 10 mg) prepared from PBS tablets (Dot Scientific, Michigan) with deionized distilled water

and filter sterilized (Millipore, Burlington, MA). Control animals were injected with an

equivalent volume of saline made of 1x PBS. Animals continued to receive injections at 5mg/kg

every 20 minutes until status epilepticus (SE) occurred (same number of saline injections in

control mice). Some animals received alternating 5mg/kg and 2.5 mg/kg injections after the

initial 10 mg/kg dose, but we found this schedule took more time than 5 mg/kg injections and our

survival rate (>90% across all cohorts) was no different between the two schedules. Animals

were considered to be in SE when displaying persistent behavioral seizures of level 4-5 on the

Racine Scale (Racine, 1972), less than 5 minutes apart, for a minimum of 30 minutes. Animals

received 4-9 injections depending on tolerance to kainic acid and injection schedule. After the

injection schedule, animals were given fresh apple slices, monitored until SE ceased (assessed

from normal posturing, within 1-1.5 hours), and returned to group housing. They were

monitored, weighed and animals weighing less than their preinjection levels were given 0.4 ml of

1x PBS via intraperitoneal injection on a daily basis until their weight exceeded preinjection

level. No animal injected with saline ever experienced status epilepticus, lost weight, or required

recovery injections of 1x PBS. Mice remained in the vivarium for 7-9 weeks after injections,

allowing time for the development of epilepsy in KA mice (the latent period in the KA model of

TLE has previously been evaluated as 10-30 days before the first spontaneous electrographic

seizure, and ~11±5.4 weeks before the first spontaneous convulsive seizure (Levesque and Avoli,

2013). During this time some KA mice became aggressive and many of the aggressor animals

were removed from group housing and caged individually.

Mouse Behavior. At 14-16 weeks of age, mice started behavioral testing on a variant of a

novelty recognition-based BPS task that is well established in rodents to report mnemonic

discrimination abilities (van Hagen et al., 2015). Our protocol lasted 4 weeks, including 1 week

of habituation and 3 weeks of trials. During the first week of habituation, each mouse was gently

handled by the experimenter for five minutes every day. On the second week mice were split into

three groups (A, B, and C). Each group went through a three-day schedule where animals were

habituated on the first and second days and performed the object location task on the third day.

Groups were staggered by one day each, such that group A was scheduled Monday-Wednesday,

B Tuesday-Thursday and C Wednesday-Friday. Animals were housed following a 12:12 light-

dark cycle and always handled or tested between 1 pm and 4pm (light period).

The arena used for the BPS task (see **Figure 3A**) was a 63.5 x 63.5 x 15 cm open topped

Plexiglas box wrapped on the outside and bottom with black felt cloth. On each side of the arena

were different images serving as proximal cues for orientation. The room also had numerous

distinct distal cues (e.g. the video camera placed over the center of the arena, shelving, etc.), that

were kept unchanged. During behavioral testing, the room was lit with a single bank of overhead

fluorescent lamps.

For the BPS task, animals underwent three phases of exploration in the arena, each

lasting 3 min and separated by 1.5 min of single housing in a solitary chamber. In Phase 1

(habituation) the arena was empty, in Phase 2 (sampling) the arena contained two identical

objects in the center, and in Phase 3 (testing) one object had been randomly selected and

displaced by a distance of 7, 14 or 21 cm. The arena and objects were cleaned with 70% ethanol

before and after each phase. During Phases 2 and 3, the experimenter recorded the amount of

time that the mouse spent interacting with each object with two silent stopwatches. The mouse

was considered to be interacting with an object if it was within 1 cm of it and oriented towards,

sniffing, scratching, or on top of the object. These three phases were repeated three times with a

different distance each week, with the order of distances randomized. Objects used for the task

were ~6 cm tall block style plastic colored figures or metal cylinders, with a footprint of ~2 cm

in diameter.

All phases of the experiment were recorded with a 1080p webcam positioned above the

arena. During all phases, the experimenter sat in the corner of the room out of the line of sight of

the mouse. The experimenter was consistent in appearance and smell (e.g., same experimenter

for a given cohort, wearing same white lab coat and gloves of the same color each day; use of

scented soap, etc. was minimized).

BPS Analysis. The discrimination ratio for each trial was computed as $(T_{moved} - T_{unmoved})$

/ T_{total} where T_{moved} is the time spent exploring the moved object, $T_{unmoved}$ is the time spent

exploring the unmoved object and T_{total} is the sum of T_{moved} and $T_{unmoved}$. The exploration times

used to calculate the reported discrimination ratio were manually collected by experimenters

during the BPS task. We (M.V.J. and J.A.P.) also developed a home-written program using video

recordings for motion-tracking and trajectory analysis of each mouse during behavioral testing.

Discrimination ratios computed from automated tracking data were well correlated to the ratios

from manually recorded times, and led to similar results ($R^2 = 0.68$, $T(117) = 16.07$, $P < 0.001$).

Automated tracking data were also used to assess a) distance traveled as an estimate of motility

and exploration and b) tendency to stay close to walls (i.e., thigmotaxis) as an estimate of

anxiety. Thigmotaxis (Simon et al., 1994) was evaluated as the proportion of time spent within

6.35 mm of the wall (10% of the arena width).

Mouse electroencephalography (EEG). EEG electrode implantation was performed at

~18-20 weeks of age for all animals, following a previously established protocol (Wallace et al.,

2015). Briefly, mice were anesthetized with isoflurane and stainless steel screw electrodes were

implanted in the skull (bregma +1.5 mm and 1 mm right, bregma -3 mm and 1 mm left, and

lamda -1 mm at midline). Two stainless steel braided wires were placed in the nuchal muscles

for electromyography (EMG) recording. After a 72 hour recovery, we transferred mice into

individual tethered EEG acquisition chambers and allowed a >12 hour acclimation period. We

acquired EEG and EMG signals continuously for 3 days. Recordings were digitized with an

XLTek amplifier (XLTEK, USA) sampled at 1024 Hz. Ad libitum access to food and water was

ensured.

Interictal spike (IIS) analysis. In humans and most animal models of acquired TLE,

overt seizures are relatively rare (often much less than once per day) (Levesque et al., 2016). In

contrast, nonconvulsive and subclinical epileptiform events such as interictal spikes (IISs) can be

very frequent, as much as many hundreds per day. Therefore, in order to assess epileptiform

activity in KA animals, we used a modification of our previously published principal

components (PC)-based method to quantify IISs (Pfammatter et al., 2018) and calculated an

"Hourly IIS index" for each animal. All detected high-amplitude EEG events from 9 Ctrl and 15

KA animals were projected into the space spanned by their first three PCs. The main

modification here is that, instead of using a Gaussian Mixture Model (GMM) to assign events to

'clusters', we simply gridded the PC space into 'voxels' and computed relevant quantities within

each voxel exactly as we computed those same quantities within GMM 'clusters' previously: 1)

The probability that events within a voxel are characteristic of an epileptogenic treatment (i.e.,

KA) was computed as the voxel-wise proportion of events coming from KA mice, 2) Only the

voxels above chance level were considered specific to epileptogenic treatment and probabilities

were scaled accordingly, 3) The Hourly IIS Index of a given animal was defined as the average

frequency of detected events weighted by the scaled probabilities of each event's voxel. This new

voxel-based method has the advantages to make no assumptions about the structure of the data in

PC space or about the number of clusters to fit. Instead, the main free parameter is now the voxel

volume: we selected a size of 10 cubic PC units following the same optimization procedure

described in Pfammatter et al. (2018) in order to avoid overfitting. Extensive exploration (not

shown) revealed that the final results are similar to the GMM method and are not importantly

affected by moderate changes in voxel volume.

Slice electrophysiology. Mice used for electrophysiology were p115-p182 at the time of

experiment (mean ± SEM: p141 +/- 4 days; no difference in age between treatments: KA n = 13,

Ctrl n = 6, U-test: P = 0.9, Z = 0.2, rank sum = 246). Age was also not correlated with any

summary statistics presented in this study.

Adult mice were euthanized by transcardial perfusion with oxygenated PBS under

isoflurane anesthesia, before decapitation and brain extraction. 400 μm horizontal slices of the

ventral and intermediate hippocampus were prepared as detailed in Madar et al. (2019a). After

slicing in a sucrose-based cutting solution (Yi et al., 2015), slices were transferred to an

incubation chamber filled with 50% cutting solution and 50% artificial cerebrospinal fluid

(aCSF) at 37°C for 30 minutes, then room temperature. Patch-clamp recordings were done in a

chamber submerged with aCSF containing (in mM) 125 NaCl, 25 NaHCO3, 2.5 KCl, 1.25

NaH2PO4, 2 CaCl2, 1 MgCl2, and 25 D-Glucose, flowing at 5 ml/min and saturated with a gas

mixture of 95% O_2 and 5% CO_2. Stimulation was applied through a double-barreled "theta"

pipette filled with aCSF. Patch pipettes were filled with an intracellular solution of the following

composition (in mM): 135 K-gluconate, 5 KCl, 0.1 EGTA, 10 HEPES, 20 Na-Phosphocreatine,

2 Mg_2-ATP, 0.3 Na-GTP, 0.25 $CaCl_2$ adjusted to pH 7.3 with KOH and 310 mOsm with H_2O,

leading to a 2-5 MΩ pipette resistance in aCSF. Whole-cell patch-clamp recordings of single DG

GCs in response to electric stimulation of the outer molecular layer (perforant path) were

performed as detailed in Madar et al. (2019a) (see **Figure 4A**), and the stimulation protocols

used to test neural pattern separation were the same as in Madar et al.(2019b) (see **Figure 4B** and

5). Briefly, input sets used for stimulation were composed of five (type 1) or ten (type 2 and 3)

spiketrains (two seconds long), delivered sequentially (separated by five seconds of pause) and

repeated ten or five times, respectively, in order to yield fifty output spiketrains. The stimulation

pipette was placed >100µm lateral to the recorded GC to avoid direct stimulation of GC

dendrites, with the baseline membrane potential held at -70 mV for current and voltage-clamp

recordings (see **Figure 4A-B, 7B**).

Intrinsic electrophysiological properties of recorded GCs were the following (mean ±

SEM for Ctrl / KA): resting membrane potential V_{rest} = -78.0 ± 1.9 / -81.4 ± 1.2 mV; membrane

resistance R_m = 139 ± 17 / 178 ± 16 MΩ; membrane capacitance C_m = 18 ± 1.2 / 17 ± 0.7 pF.

There were no significant differences between control and KA mice (U-tests: P = 0.1, 0.2, 0.7; Z

= 1.6, -1.2, 0.4; rank sums = 294, 198.5, 252.5 respectively).

Neural pattern separation analysis. Similarity between spiketrains was assessed as in

Madar et al. (2019a, b), using three metrics based on dividing spiketrains into time bins of a

specific duration τ_w: the Pearson's correlation coefficient (R), the normalized dot product (NDP)

and the scaling factor (SF). NDP and SF consider spiketrains as vectors of spike count per bin

and measure the angle (NDP is the cosine) and the ratio of norms (SF) between the two vectors.

R considers vectors of deviation from mean spike count, which loses the original angle between

the vectors of raw spike counts and leads to considering common periods of silence as correlated,

unlike NDP. Both R and NDP are sensitive to binwise synchrony, whereas SF is more sensitive

to differences in firing rate and burstiness (see Madar et al. 2019b for a detailed discussion on the

different neural codes assumed by each metric). To further test the role of various spiketrain

features in mediating pattern separation, we also considered neural codes purely focused on

either 1) the average firing rate (FR), 2) the compactness (1 - proportion of time bins with at least

one spike) and 3) the occupancy (average number of spikes in bins with at least one spike).

These were computed as in Madar et al. (2019b) and the degree of pattern separation (or

convergence, for negative values) was evaluated as the difference between the dispersion of the

output spiketrains minus the dispersion of the input spiketrains. The dispersion was computed as

the mean absolute value of pairwise differences for each spiketrain feature (FR, Compactness or

Occupancy) over all spiketrains of a set, excluding self-comparisons and, in the case of the

output sets, all comparisons between output spiketrains resulting from the same input spiketrain.

When using a similarity metric S (R, NDP or SF), pattern separation was assessed following the

same logic. First, the similarity between each pair of spiketrains of a given set was computed.

S_{input} was the average for all pairs of an input set (excluding self-comparisons), and S_{output} was

the average of all pairs in output sets excluding comparisons between output spiketrains coming

from repetitions of the same input train. This yielded a single output similarity value for a given

output set, and the degree of pattern separation was thus S_{input} - S_{output}. To gain a finer view, we

also performed a pairwise analysis (as in Madar et al. 2019b) where each of the pairwise S_{input} of

an input set were distinguished (10 pairs for input sets of type 1, 45 pairs for input sets of type 2

and 3). The pairwise S_{output} (pw S_{output}) was thus the average similarity across all pairs of output

spiketrains resulting from a given pair of input spiketrains. Pattern separation was computed as

pw S_{input} – pw S_{output}.

Software and statistics. Data analysis was performed using MATLAB (Mathworks,

Natick, MA, USA). The Lilliefors test was used to verify the normality of data distributions

(Lilliefors, 1967). Parametric or nonparametric statistical tests were appropriately used to assess

significance (p-value < 0.05). Throughout the Results section, KW ANOVA corresponds to the

nonparametric Kruskal-Wallis analysis of variance, U-test corresponds to the Wilcoxon rank-

sum test equivalent to the Mann-Whitney U-test, and KS test corresponds to the two-sample

Kolmogorov-Smirnov test (sidedness is specified in the legends).

To analyze performance of mice on the BPS task (**Figure 3B**) we performed a two-way

ANOVA with the Matlab function *anova* based on a linear mixed-effects model built using

fitlme, with the distance of the moved object as a continuous fixed effect, animal treatment (Ctrl

vs. KA) as a categorical fixed effect, and animal identity nested within object distances as a

random effect to account for repeated measurements. The same procedure was used for each

panel in **Figure 3D-E.**

To determine whether (S_{input}, S_{output}) distributions were significantly different (**Figures 4-**

5), we performed an analysis of the covariance (ANCOVA) using separate parabolic or linear

regression models, implemented in MATLAB with a custom-written code following the method

described in (Motulsky and Ransnas, 1987), as in Madar et al. (2019a, b)

In **Figure 8A**, the non-linear regression was performed using the Matlab function *fitnlm*

with a three-parameter model function $f(x) = a/(x+b)+c$

Results:

Behavioral pattern separation deficits in TLE. Mnemonic discrimination is often called

behavioral pattern separation (BPS) because it is hypothesized to be the behavioral outcome of

neural pattern separation. It is the ability to discriminate between similar experiences (event,

environment, object, etc.) that occurred at different times (Santoro, 2013). Multiple BPS tasks

have been demonstrated to be DG-dependent in humans (Baker et al., 2016) and rodents (Kesner

and Rolls, 2015; Kesner et al., 2016; Bui et al., 2018). We thus chose some of these established

tasks to test whether TLE impacts DG-dependent cognition, both in humans and mice.

In humans, we used the object recognition-based *Mnemonic Similarity Task* developed by

the Stark lab (Stark et al., 2019), where participants must distinguish between similar images

presented at different times (**Figure 1A**, **Materials and Methods – Human behavior**). Patients

previously diagnosed with TLE showed a severe deficit (~50%) compared to nonepileptic

subjects in their ability to correctly identify objects as being similar but not identical (**Figure 1**).

These results confirmed our hypothesis that TLE impairs DG-dependent mnemonic

discrimination, thus warranting further study of the impact of TLE on DG computations that

might explain this cognitive deficit.

Figure 1. Human patients with TLE have mnemonic discrimination deficits

(**A**) A schematic (adapted with permission from Yassa et al., 2011) of the mnemonic discrimination task given to patients with TLE and nonepileptic control subjects. The discrimination index measures the ability of participants to correctly identify images as similar but not identical to a previously seen image.

(**B**) Human patients with temporal lobe epilepsy (red, n = 15) display a significant deficit in this object recognition-based mnemonic discrimination task compared to nonepileptic control subjects (black, n = 20). Unpaired two-sided T-test: P = 0.0183, T(33) = 2.4832.

(**C**) The same data from B, presented as cumulative frequency distributions. A non-parametric one-way KW ANOVA on the indices grouped by treatment confirms a highly significant deficit in visual pattern separation memory for human subjects with TLE (P = 0.0343, χ^2(34) = 4.4809).

(**D**) TLE and Ctrl groups were properly age matched (unpaired two-sided T-test: P = 0.9411, T(31) = -0.0745).

(**E**) There were slightly more women than men in the control group, but the gender difference was not significant between the two treatments (one-way KW ANOVA on gender values, 1 or 2, grouped by treatment: P = 0.0663, χ^2(32) = 3.3724).

To investigate the effect of TLE at the behavioral, computational and cellular levels, we

turned to the common post-kainate animal model of acquired TLE (KA, **Methods – Mouse**

experiments). An automated detection algorithm of interictal spikes (IISs) allowed us to

quantify epileptiform activity in each animal even in the absence of seizures (**Methods –**

Automated EEG analysis and see Pfammatter et al. 2018). Most KA animals developed IISs

(**Figure 2**).

361
Figure 2. Kainate-injected mice have epileptiform electrographic events with a range of
event frequency
(**A**) The Hourly Interictal Spikes (IIS) index, a proxy for epilepsy severity based on EEG
recordings (Pfammatter et al., 2018), was calculated for 9 Ctrl (black) and 15 KA (red) animals.
KA animals have a significantly higher Hourly IIS index than Ctrl animals (U-test: $P = 0.009$, Z
= 2.207, rank sum = 225).
(**B**) To compute the index, high-amplitude events (200 ms duration) were identified from left
frontal EEG and projected on the first three principal components (PC) of the ensemble of events
from all mice. Each dot here is an EEG event, red when from a KA animal and black when from
a Ctrl. Notice the 'fingers' of red KA dots extending from the central cluster of mixed Ctrl and
KA data points.
(**C**) Using a voxel gridding of the PC space (voxel size: 10 cubic units), we calculated the
probability for each event to be specific to KA treatment (see scale bar). The hourly IIS index of
a mouse is the average number of detected events/hour, weighted by their probability of being
specific to the KA treatment.
377

Prior to EEG recordings, mice were subjected to an object-location novelty-recognition BPS task (**Methods – Mouse Behavior**) where their ability to discriminate between a moved object versus an identical unmoved object was measured for different object displacement distances (**Figure 3A**). Our results demonstrate that, on average, KA mice have lower object-location mnemonic discrimination than control mice at all object distances (**Figure 3B-C**). Control analyses of motility (assessed by the total distance moved) and anxiety levels (assessed by the time spent near walls) revealed that KA mice have no obvious locomotor deficits nor do they spend more time near walls during the testing phases (**Figure 3D-E**). Moreover, there was no correlation between individual mnemonic discrimination ratios and total distance moved (R^2 = 0.028, $T(118)$ = 1.86, P = 0.066) or time spent near walls (R^2 = 0.002, $T(118)$ = 0.50, P = 0.62) during phase 3. These results suggest that differences between KA and control mice in the BPS task were not due to differences in motor ability, exploration or anxiety.

TLE is thus accompanied by mnemonic discrimination deficits in both humans and mice.

Figure 3. Kainate-injected mice have deficits in mnemonic discrimination

(A) Object location BPS task arena setup. The experiment was run in three phases of three minutes. In Phase 1, a mouse was allowed to explore the empty square arena. In Phase 2, the same mouse was allowed to explore the same arena but with two identical objects (position indicated by grey circles). In Phase 3, one randomly chosen object was moved either 7, 14 or 21 cm from its original position and the mouse was allowed to explore again. Between phases, the mouse was placed in a solitary chamber for 1.5 minutes. Each mouse performed the task once at each distance with different objects, in randomized order, with a week in between trials.

(B) Performance at the BPS task is evaluated with a discrimination ratio computed for each animal at each distance. Healthy mice generally prefer to explore objects with novel characteristics (such as change in relative position) and are thus expected to spend more time with the moved object if they notice it has changed location. Discrimination is thus quantified as the difference between the time exploring the moved object and the time exploring the unmoved object. A two-way ANOVA with repeated measures shows that mice discriminate better when objects are moved farther apart, and that Ctrl mice (n = 29) discriminate better than KA mice (n = 27) (Distance Object Moved: $P = 0.002$, $F_{(1,118)} = 9.84$; Treatment: $P = 0.027$, $F_{(1,118)} = 5.02$).

(C) Subset of the data in B plotted in the space of discrimination at each distance, allowing a clearer picture of individual performance. Each animal is represented by a single dot in this 3D space (42 animals with records at the three distances: 24 KA and 18 Ctrl). For each data point, the Mahalanobis distance (De Maesschalck et al., 2000) from the Ctrl centroid is represented by the dot size, and is also displayed in the inset (mean +/- SEM). Mahalanobis distances are larger for KA than Ctrl animals, confirming that the KA population displays mnemonic discrimination deficits (two-sided T-test: $P = 0.003$, $T_{(40)} = -3.15$).

(D,E) Control analysis to determine whether there were motor impairments or anxiety/curiosity differences between KA and Ctrl mice, across phase 1, 2 and 3 of the BPS task. In each panel, individual mice are represented by one to three data points, as in B, depending on how many conditions (Distance Object Moved) they were tested on. Mean +/- SEM.

(D) Distance traveled was used as a proxy for motor ability. During Phase 1 and Phase 2, control and KA mice moved similar distances ($P = 0.507$, $F_{(1, 120)} = 0.443$ and $P = 0.070$, $F_{(1, 122)} = 3.35$). During Phase 3 KA mice moved farther than Ctrl mice ($P = 0.023$, $F_{(1, 124)} = 5.34$).

(E) Percent of time near wall was used as a proxy for anxiety. During Phase 1 (habituation), KA animals spent more time near the maze walls compared to Ctrl ($P = 0.03$ $F_{(1,120)} = 4.82$). During Phases 2 and 3, Ctrl and KA animals spent a similar amount of time near walls ($P = 0.68$, $F_{(1,122)} = 0.173$ and $P = 0.35$, $F_{(1, 124)} = 0.881$). F-tests correspond to one-way ANOVAs with repeated measures, with Treatment as a categorical fixed effect.

DG computational impairments in TLE. To test our hypothesis that DG pathologies in TLE lead to a breakdown of the neural pattern separation function of the DG, which in turn would translate into the cognitive impairments we observed, we measured pattern separation in hippocampal slices from the same KA and control mice used above in behavioral and EEG

experiments. Briefly, the assay has three steps (**Figure 4A-B** and see Madar et al. 2019a, b): 1)

ensembles of stimulus patterns (simulating afferent input spiketrains) are generated, with known

degrees of similarity to each other. These spiketrains are then fed into the DG by stimulating the

lateral perforant path. 2) The response of a single GC is recorded in whole-cell current-clamp. 3)

The similarity between the output spiketrains is compared to the similarity between the input

spiketrains, revealing the degree of separation or convergence (**Figure 4C** and see **Material and**

Methods – Neural pattern separation analysis).

We tested pattern separation levels in response to three types of input sets. Input sets of

type 1 were constituted of Poisson spiketrains with a 10 Hz mean firing rate designed to have a

prespecified average similarity as measured by the Pearson's correlation coefficient (R) (**Figure**

4B). For an input set with spiketrains correlated by 77% between each other (timescale: $\tau_w = 20$

ms), the output spiketrains of GCs from epileptic mice had a higher average correlation (R_{output})

than GCs from control mice (**Figure 4C**). This demonstrates a decrease in pattern separation in

the DG of epileptic mice. Exploring a wider range of input correlations shows that the pattern

separation function of the DG of KA mice is generally impaired compared to the normal pattern

separation function (**Figure 4D**). Furthermore, a decrease in pattern separation was observed at

multiple timescales (millisecond to second) and using multiple similarity metrics that assume

different neural codes (**Figure 4E**, see **Material and Methods – Neural pattern separation**

analysis). The impairment is most noticeable using the NDP metric, showing that DG output

patterns are less orthogonalized in KA animals at all timescales, whereas differences in pattern

separation via scaling (SF metric) are small but significant at large timescales.

457
Figure 4. Epileptic mice have neural pattern separation deficits
(A) Histology of the DG in a horizontal slice (Cresyl violet/Nissl staining; scale bar: 250 μm),
overlaid with a schematic of the experimental setup: a theta pipette in the ML is used to focally
stimulate the outer molecular layer (input) while a responding GC is recorded via whole-cell
patch-clamp (output). GCL: granule cell layer, H: hilus, ML: molecular layer. Solid lines
represent dendrites and dashed lines axons.
(B) Current-clamp recordings of the membrane potential of two different GCs (Ctrl and KA) in
response to the same set of input trains. An input set is constituted of five different trains of
electrical pulses following a Poisson distribution with an average rate of 10 Hz. The Pearson's
correlation coefficient (R) between two input trains is computed with a binning window (τ_w) of
20 ms (*Left*). R_{input} is the average of the ten pairwise coefficients, diagonal excluded. After
converting the GC recordings to vectors of binned spike counts, the pairwise coefficients and
their average (R_{output}) can be computed the same way (Madar et al, 2019b). Note that the input set
was repeated 10 times to yield output sets of fifty sweeps (only one repeat is shown).

(C) The R_{output} of GCs from kainate-injected animals is higher, on average, than in GCs from
controls. Data points correspond to single output sets (KA: 27 recordings from 24 GCs; Ctrl: 12
recordings from 11 GCs), all being responses to the same input set as in B. Black crosses with
error bars are means +/- SEM. The asterisk signals significance: U-test comparing the medians: P
= 0.011, Z = -2.541, rank sum = 156; two-sided two-sample KS test comparing the distribution
shapes: P = 0.0037, D = 0.583. The dashed line corresponds to R_{input} (0.77): any value below the
line implies effective pattern separation. Thus, GCs from kainate animals exhibit less pattern
separation than controls.
(D) Pattern separation was investigated over a wide range of input correlations by using four
different input sets of five 10 Hz Poisson trains (at τ_w = 20 ms: R_{input} = 0.24, 0.45, 0.77, 0.95).
Crosses and error bars correspond to mean +/- SEM (pw R_{input}, pw R_{output}) across multiple
recordings (KA: 6-24 GCs per input set; Ctrl: 4-11 GCs). Data points below the identity line
(dashed) correspond to pattern separation. The distributions between KA and Ctrl are
significantly different (ANCOVA with separate parabolic models fitting the data points and not
the means: P < 0.0001, F(3,714) = 15.485. Solid curves are the parabolic models used for the
ANCOVA.
(E) Levels of pattern separation measured using different similarity metrics (S: R, NDP and SF,
see Methods) and timescales. Cumulative frequency distributions of the distance of (pw S_{input},
pw S_{output}) data points to the identity line in pattern separation graphs like in E. Positive values of
the x-axis correspond to pattern separation, and negative values to pattern convergence. Insets
show medians. For R and NDP, distributions are significantly shifted to the left, showing that
GCs from KA exhibit less decorrelation and less orthogonalization (ANCOVA as in D, but using
separate linear models: P < 0.01 for τ_w = 5 to 1000 ms. For R/NDP and τ_w = 10, 20, 100, 250 ms,
P = <0.0001/<0.0001, <0.0001/<0.0001, 0.001/<0.0001, 0.011/0.0001, F(2, 716) = 15.2/18.2,
20.3/23.4, 6.87/15.97, 4.5/9.4). For SF, scaling levels are weakly but significantly shifted to less
separation at large τ_w (SF: P > 0.2 except at 250 ms and 500 ms, with P = 0.026, 0.046, F(2, 716)
= 3.65, 3.09 respectively).
499
An input set of type 2, made of Poisson spiketrains similar in terms of R but with varying

firing rates (**Figure 5A1**), and an input set of type 3, with 10.5 Hz uncorrelated trains (R = 0)

with varying burstiness (**Figure 5B1**), were designed to explore a wide range of input similarities

as measured by SF and to characterize DG computations on inputs with a variety of statistical

structures (Madar et al., 2019b). As we have shown before, the normal DG exhibits low pattern

separation via scaling for highly similar inputs, and significant pattern convergence for dissimilar

inputs (Madar et al., 2019b). In KA mice, GCs show a decrease in both pattern convergence and

separation via scaling: the DG computation function is steeper, closer to the identity line,

meaning that the DG of KA mice is weaker at transforming the similarity of its inputs in general

(**Figure 5A1 and B1**). When similarity is measured with R or NDP (binwise synchrony code)

instead of SF (neural code focused on FR and bursting), these experiments confirmed that pattern

separation is decreased in TLE (**Figure 5A2**) and that this deficit is strongest when inputs are

highly similar (i.e., when pattern separation is theoretically most important) and weaker when

inputs are already dissimilar (**Figure 5B2**).

515
Figure 5. Multiple DG computations are affected by epilepsy
Our standard input sets (type 1, Figure 4) consisted of 10 Hz Poisson trains. Two other input sets
(type 2 and 3) were designed to explore single GCs responses to inputs with diverse structures

and statistics, both with a wide range of pairwise similarity as measured by SF (in contrast to
input sets of type 1).
(A1) *Left*: Input set type 2 was constituted of spiketrains following a Poisson distribution, each
with a different firing rate (FR), but with an average R_{input} constrained around 0.75 (τ_w = 10 ms).
Middle and Right: Pattern separation graphs showing the pairwise output spiketrain similarity as
a function of the pairwise input similarity, as measured by SF with two different timescales,
averaged across multiple GCs (KA: 16, Ctrl: 5). Distributions are significantly different,
suggesting in particular that epilepsy causes a decrease in pattern convergence (via scaling) for
low input similarities (ANCOVA with separate linear models, for τ_w = 20 and 250 ms
respectively : P = 0.0016 and < 0.0001, F(2, 941) = 6.5 and 10.4). **(A2)** Levels of pattern
separation for type 2 inputs, measured using different similarity metrics and timescales as in
Figure 4. It confirms that epilepsy decreases the separation of similar Poisson input spiketrains as
measured by R and NDP, consistent with Figure 4, and shows that differences in terms of SF,
although small, are significant (ANCOVA as in A1: R: P < 0.0001 for τ_w = 20 ms up to 1000 ms,
P = 0.02 and 0.18 for 5 and 10 ms; NDP and SF: P < 0.025 for τ_w up to 1000 ms. Detailed
statistics for τ_w = 5, 10, 20, 50, 100, 250, 500 and 1000 ms respectively: R, P = 0.0243, 0.1820,
<0.0001, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001, F(2, 941) = 3.7, 1.7, 10.2, 15.1, 24.1,
36.0, 50.2, 27.0; NDP, P = 0.0022, 0.0240, <0.0001, <0.0001, <0.0001, <0.0001, <0.0001,
<0.0001, F(2, 941) = 6.2, 3.7, 13.8, 17.4, 21.4, 26.0, 21.0, 16.8; SF, P = 0.0002, 0.0012, 0.0016,
0.0240, 0.0073, <0.0001, <0.0001, 0.0002, F(2, 941) = 8.7, 6.7, 6.5, 3.7, 4.9, 10.9, 10.4, 8.7).
(B1) *Left:* Input set type 3 was constituted of spiketrains with 21 spikes (FR = 10.5Hz) that were
distributed among bins to produce trains with varying burstiness (Madar et al., 2019b). R is close
to 0 for all pairs. *Middle and Right*: same analysis as in A1 (KA: 8 GCs, Ctrl: 3 GCs).
Distributions are significantly different, suggesting again that epilepsy causes a decrease in
pattern convergence via scaling, for low input similarities (ANCOVA for τ_w = 20 and 100 ms: P
< 0.0001 and F(2, 491) = 10.3 and 13.8 respectively).
(B2) Same analysis as in A2. The directions of impairments are the same as in A2, showing that
epilepsy decreases pattern separation in terms of R and NDP but slightly improves it in terms of
SF. The shift via scaling is weak but significant at all timescales, and larger at the longest τ_w.
ANCOVA: R, P < 0.05 for τ_w = 5, 10 and 500 ms; NDP and SF, P < 0.007 for τ_w up to 500 ms.
Detailed statistics for τ_w = 5, 10, 20, 50, 100, 250, 500 and 1000 ms respectively: R, P = 0.0451,
0.0004, 0.3727, 0.7357, 0.9340, 0.4271, 0.0012, 0.2889, F(2, 491) = 3.1, 8.1, 1.0, 0.3, 0.1, 0.8,
6.8, 1.2; NDP, P = <0.0001, <0.0001, 0.0003, 0.0012, <0.0001, 0.0069, <0.0001, 0.4154, F(2,
491) = 11.1, 24.3, 8.4, 6.8, 11.2, 5.0, 14.6, 0.9, SF, P = <0.0001, <0.0001, <0.0001, <0.0001,
<0.0001, 0.0065, 0.0002, 0.0003, F(2, 941) = 42.1, 20.9, 10.3, 13.6, 13.8, 5.1, 8.5, 8.1.
554

Pathological spiking patterns in a subset of GCs from epileptic mice. What features of

GC output spiketrains are changed in TLE to explain the difference in DG computations between

KA and control mice? On visual inspection, we noticed that many KA neurons occasionally fired

short bursts of action potentials after a single input pulse (**Figure 6A**), which is quite unusual for

control GCs (Madar et al., 2019a). To determine whether there was a difference in the spiking

patterns of KA and control GCs, we measured three spiketrain features: 1) the average firing rate

(FR) of a GC across a full recording set (fifty spiketrains), 2) the probability of spiking after a

single input pulse (SP) and 3) the probability of bursting (p(Burst), i.e., more than one spike)

after a single input pulse. Our results show that on average GCs from KA mice fire more

faithfully after an input (**Figure 6B**) and also tend to fire in bursts (**Figure 6C-D**), which

together lead to a higher FR (**Figure 6B, E**).

Closer inspection suggests that the distributions of SP, p(Burst) and FR are different

between KA and controls, with larger variance and upper tails for GCs from KA mice **(Figure**

6B-E). Although we did not formally test for discrete clusters, visual inspection of Figure 6E

shows that a subset of GCs from KA animals have higher SP, p(Burst) and FR than any control

GC. Thus, it appears that, following an epileptogenic insult, only a subset of GCs displayed

pathological characteristics amidst a background of seemingly normal GCs.

573
Figure 6. In epileptic mice, a subpopulation of GCs shows pathological spiking patterns
(**A**) Example of current-clamp recordings in GCs from a kainate-injected (KA) vs a control
animal in response to the same Poisson input train, illustrating that some GCs from KA spiked
more, and had a tendency to fire short bursts (2-4 spikes riding a single EPSP).
(**B-E**) Some GCs from KA exhibited larger firing rates due to a higher probability of spiking at
least once after an input spike (spiking probability), and sometimes a higher probability of
spiking more than once between two input spikes (p(Burst)) than GCs from controls. The firing
rate, spiking probability and p(Burst) of a neuron were all computed as the average over the fifty
sweeps of an output set from a pattern separation experiment (input set type 1, see Figure 4B).
(**B**) *Left*: data points correspond to the same recordings as in Figure 4C (KA: 27; Ctrl: 12). Black
dash and error bars are mean +/- SEM. A U-test was used to compare the medians, showing that
on average FR and SP are higher in KA (FR: P = 0.0415, Z = -2.04, rank sum = 172.5; SP: P =
0.0613, Z = -1.87, rank sum = 168.5). *Right*: Frequency distributions of the same data. A one-
sided two-sample KS test shows that the KA distribution has a larger tail in both cases (FR and
SP: P = 0.0465, D = 0.4074), indicating that a subset of KA GCs are pathological.
(**C**) Same as in B for p(Burst). Note that the left graph has a $\log_{10}$ scale showing all data points,
whereas the middle graph has a linear scale zoomed in (i.e. not showing the 5 largest KA values).
The graph on the right has the same scale as the middle graph. Together, they suggest that in KA,
there might be a healthy population of GCs coexisting with a different population of
pathologically bursty GCs (U-test: P = 0.0153, Z = -2.16, rank sum = 168.5; KS test: P = 0.0259,
D = 0.4444).
(**D**) Some bursts were detected by our algorithm in GCs from Ctrl mice, but the vast majority of
those were due to the temporal summation of two EPSPs resulting from input spikes occurring
close in time. In contrast, a large number of bursts in GCs from KA mice come after an isolated

input spike. *Top*: Time interval between the two input spikes (inter-spike-interval, ISI) preceding
a given burst. Data points correspond to all detected bursts from the recordings in C. U-test: P =
0.0003, Z = -3.58, rank sum = 5906.2. *Bottom*: Cumulative frequency distributions of the same
data. A one-sided two-sample KS test demonstrates that the distributions are different (P <
0.0001, D = 0.2699).
(E) Same data as in B and C: the elbow in the distribution of SP, p(Burst) and FR values visually
defines two clusters of neurons. One cluster contains GCs from both KA and Ctrl mice (with
GCs from both groups spanning a large range of SP values but low p(Burst)) and can be
considered "normal". In contrast, the other cluster, with highest SP combined with high p(Burst),
contains only GCs from KA animals and can thus be considered "pathological".
608

What is the origin of the pathological spiking patterns? We first controlled that

stimulation intensities were comparable between treatment groups and were not causative of

abnormal spiking patterns (**Figure 7A**). We next asked whether the abnormal firing came from

changes in the synaptic drive. To answer this question, we measured the synaptic input-output

relation (Ewell and Jones, 2010) by recording GCs first under current-clamp then under voltage-

clamp, in response to the same pattern separation stimulus protocol (**Figure 7B1-2**). Our data

show that the average excitatory drive per neuron was not different between KA and control

groups, and that it does not predict burstiness (**Figure 7B3**). A finer grained analysis on the

individual currents suggests that although the excitatory drive and the spiking output are

correlated, high amplitude EPSCs are neither sufficient nor necessary to elicit a burst (**Figure**

7B4). Bursting does not seem to result from other changes in excitatory input-output coupling

either (**Figure 7B5**), which suggest that inhibition might be involved. GCs are indeed subjected

to strong tonic, feedforward and feedback inhibition that controls the sparseness of their activity

(Coulter and Carlson, 2007; Ewell and Jones, 2010; Pardi et al., 2015; Lee et al., 2016) and is

altered in TLE (Alexander et al., 2016; Dengler and Coulter, 2016). We did not record evoked

IPSCs directly but, in normal mice, partial block of inhibition elevates GC firing rates and causes

bursts (Madar et al., 2019b) similar to what we observed here in KA mice. All our results thus

 converge to support the hypothesis that pathological spiking arises from an excitation/inhibition

627 imbalance in a subset of GCs, likely driven by a decrease in inhibition.

629
Figure 7. Pathological bursting arises from an excitation/inhibition imbalance
(A) Bursting and high firing rates were not an artefact of stimulation differences: The current intensity of the electric stimulation did not significantly differ between KA and Ctrl (same recordings as in Figure 4C and 6B-E, U-test: P = 0.2991, Z = 1.04, rank sum = 274.5) and stimulus intensity was a poor predictor of pathological firing patterns (as indicated by the low R^2 values of linear regressions. For FR/SP/pBurst: R^2 = 12.75%/11.69%/4.35%, F(2,37) = 5.4/4.9/1.7, P = 0.0257/0.0332/0.2027).
(B) For a subset of neurons in A (KA: 12; Ctrl: 6), we followed the current-clamp recording with a voltage-clamp recording (V_{hold} = -70mV) in response to the same pattern separation protocol in order to assess the excitatory synaptic drive and its relationship with spiking behavior. The stimulus electrode location and current intensity was unchanged. **(B1)** Example of current-clamp (top) and voltage-clamp (bottom) recording in the same GC from a Ctrl animal. Only the first

500 ms of the ten responses to repetitions of the first input train shown in Figure 4B are
displayed. *Top*: action potentials are truncated at 0 mV. *Bottom*: stimulation artefacts and
occasional unclamped spikes were blanked. **(B2)** The ten EPSCs (grey) associated to first pulse
of input trains in B1 and the mean EPSC (black). We assessed the excitatory drive as the
maximum inward current in the interval between each input pulse minus the current baseline
(i.e., mode of the current over the full sweep). P(Spike) was defined as the probability of spiking
during this interval across the ten sweeps of the current-clamp recording. **(B3)** *Top*: The mean
EPSC density for each neuron was not significantly different between KA and Ctrl (U-test: $P = $
0.5532, rank sum = 50; two-sided two-sample KS test: $P = 0.1935$, $D = 0.4167$), suggesting that
pathological spiking cannot be explained by larger EPSCs in KA mice. *Bottom*: Indeed, no clear
relationship exists between p(Burst) and the mean current density, as some GCs with a great
synaptic drive do not exhibit much bursting, and vice versa. **(B4)** Consistently, the peak current
density associated to each inter-input-interval in a recording is not well correlated to the
corresponding number of spikes in the current-clamp trace, for both Ctrl (top) and KA (bottom).
For example, excitatory currents of the same magnitude can be associated to 0, 1, 2, 3 or 4
spikes. EPSC amplitude is thus not sufficient to explain the spiking output, which suggests that
other factors, like reduced inhibition, are likely implicated in causing pathological bursts. **(B5)**
Top: The average EPSC density between two input spikes is plotted against the corresponding
probability of spiking p(Spike), as in Ewell and Jones (2010). The resulting input-output (IO)
distribution is fitted with a sigmoid (two GCs representative of KA and Ctrl are shown). *Bottom*:
In a subset of GCs, mostly from KA, The IO distribution appears disorganized and sometimes
difficult to fit. A nonparametric proxy of this disorganization is the standard deviation (SD) of
the current density, averaged across each level of p(Spike) excluding p(Spike) = 0 and 1. The
average SD for a given GC was plotted against the propensity of that neuron to fire bursts. There
is no clear relationship between these two quantities, suggesting that: 1) decoupling between the
excitatory drive and spiking probability is not directly related to bursting and 2) a subset of GCs
from KA with apparently healthy spiking patterns could already exhibit a more subtle but
pathological IO disorganization.
670

How do high FR and bursts relate to the DG computational impairment measured in KA

mice? We first tested the hypothesis that higher GC firing rates and burstiness yield less pattern

separation as measured with R (**Figure 8A**). In GCs of young mice, FR and pattern separation

are related, but loosely (Madar et al., 2019a), whereas here in adult mice there is a strong linear

relationship, such that GCs with pathologically high FR exhibit pathologically low pattern

separation. Similarly, abnormal bursting corresponds to abnormally low levels of pattern

separation. This analysis relates cell-wise spiking features (FR or pBurst averaged across all

spike trains of a recorded neuron) to a form of pattern separation that is not theoretically

concerned with such features (Madar et al., 2019b).

Other forms of pattern separation, more directly related to FR or bursting, can in theory

also be performed: for example by increasing spiketrain-to-spiketrain *variability* in FR or

burstiness of the output neuron, even if the average quantities were identical (Madar et al.,

2019b). Because pathological GCs only occasionally fired bursts, we asked whether TLE could

affect such forms of pattern separation. To test this, we measured the spiketrain-to-spiketrain

variability in FR as well as in two complementary measures of burstiness (occupancy and

compactness, see **Materials and Methods – Neural pattern separation analysis**). For input

sets of type 1 (10 Hz Poisson), GC output variability was slightly lower in terms of burstiness or

FR. This weak type of pattern convergence disappeared in GCs from KA mice (**Figure 8B**). For

input sets of type 2 and 3, no significant difference between KA and Ctrl was detected (**Figure**

8C-D). Overall, Figure 8 suggests that TLE reduces DG neural pattern separation mostly by

raising average levels of FR and burstiness, rather than through spiketrain-to-spiketrain

variations of those features.

Figure 8. Pathological spiking patterns explain pattern separation differences in epilepsy

(A) Neurons with high average firing rates and burstiness exhibit lower pattern separation (computed as R_{input} – R_{output}). Same data as in Figure 4C and 6B-C. There is a strong linear relationship between firing rates and decorrelation (Grey line, R^2 = 63.68%, F(2,37) = 64.9, P < 0.0001). The relationship between p(Burst) and decorrelation is better described by an inverse function with a horizontal asymptote (see Methods – Software and statistics): R^2 = 53.1%, R^2 adjusted for 3 parameters = 50.5%, F(3,36) = 20.4, P < 0.0001.

(B-D) Mean +/- SEM pattern separation levels across recordings when considering burstiness codes (Compactness or Occupancy) or a rate code (spiking frequency across a 2s sweep) instead of the binwise synchrony code assumed by R. Compactness is the proportion of time bins with at least one spike, whereas occupancy is the average number of spikes in a bin. Pattern separation corresponds to more dispersion in compactness, occupancy or firing rate in the output spiketrains than the input spiketrains. Negative values mean there is pattern convergence. U-tests comparing medians of GCs from KA and Ctrl groups were performed (p-values in panel).

(B) In response to input sets of type 1 (10 Hz Poisson trains, same recordings as in A and Figure 4), GCs exhibit low levels of convergence, if any, in terms of compactness, occupancy and rate codes, and these levels are slightly but significantly shifted to less convergence in GCs from KA for τ_w = 10 ms. This means that the relationships between firing rate or burstiness and pattern decorrelation observed in A is not due to variations of the firing rate or burstiness across sweeps. Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: P = 0.0482, .1892, .1628, .2925, .7584, Z = -1.97, -1.97, -1.31, -1.39, 0.31, rank sum = 745, 745, 801, 794, 939 ; Occupancy: P = 0.0177, 0.2586, 0.3624, 0.1425, 0.0211, Z = -2.37, -1.13, -0.91, -1.47, -2.31, rank sum = 712.5, 816.5, 835, 788, 717; FR: P = 0.0553, Z = -1.92, rank sum = 750.

(C) In response to an input set of type 2 (Poisson trains with different firing rates, same recordings as in Figure 5A), GCs from KA and Ctrl do not show significant differences (P > 0.2 for all timescales and codes). Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: P = 0.7726, 0.7726, 0.8365, 0.9014, 0.2312, Z = -0.29, -0.29, -0.21, 0.12, 1.20, rank sum = 51, 51, 52, 57, 70; Occupancy: P = 1, 1, 0.9671, 0.5915, 0.2006, Z = 0, 0, 0.04, -0.54, -1.28, rank sum = 55, 55, 56, 48, 39; FR: P = 0.9014, Z = -0.12, rank sum = 53.

(D) In response to an input set of type 3 (10 Hz trains with varying compactness and occupancy, same recordings as in Figure 5B), GCs exhibit convergence in terms of compactness and occupancy and separation in terms of rate codes. Only computations in terms of compactness and firing rate are mildly shifted in GCs from KA for τ_w = 10 ms. Detailed statistics at 10, 20, 50, 100, 250 ms, Compactness: P = 0.0848, rank sum = 9; Occupancy: P = 0.8, 0.7758, 0.9212, 0.3758, 1, rank sum = 19.5, 20, 17, 23, 18; FR: P = 0.0848, rank sum = 9.

Electrographic, behavioral and computational pathologies in individual mice. We have shown that, on average, KA mice develop EEG abnormalities (**Figure 2**), suffer mnemonic discrimination impairments (**Figure 3**) and have a bursting subpopulation of GCs with pattern separation deficits (**Figures 4-8**). Because we often performed all of the aforementioned experiments in the same mice, we next asked how those different epilepsy-related pathologies

are linked at the individual level (**Figure 9**). In a simple framework where behavioral

impairments are caused by computational deficits that are due to reorganization of DG network

function that also leads to an increase in epileptiform EEG events, one could expect a simple

relationship between all the variables we measured. Interestingly, our data suggest a more

complex view. For example, animals with few interictal spikes may still harbor some

pathological neurons (**Figure 9A**) and be highly impaired at the BPS task (**Figure 9B**). This

shows that both DG network pathologies and mnemonic impairments can occur independent of,

or before, early EEG abnormalities. Inversely, both normal and pathological GCs were recorded

in mice with more advanced EEG pathology, which confirms that those two subpopulations can

coexist in the same KA animal (**Figure 9A**). Mice with clearly epileptiform EEG activity can

also sometimes perform normally on the BPS task (**Figure 9B**). The variability in individual

behavior and in cell sampling prevents us to definitively conclude on the relationship between

single-cell pattern separation and mnemonic discrimination but, importantly, KA animals with

the largest cognitive impairments all had at least one pathological recorded GC (**Figure 9C**).

Finally, in cases where EEG, BPS and patch-clamp data were all obtained from each animal, the

combination of these measurements yielded a very obvious separation between normal and KA

subjects. Overall, our results suggest that epilepsy-related pathologies do not all develop in

concert and that computational, behavioral and electrographic measures provide complementary

informative dimensions that, together, better assess the epileptic state than any single dimension

or pair of dimensions.

Figure 9. EEG, behavioral and computational measures provide complementary insights about epileptic pathology in individual animals.

(A-D) Scatter plots relating the electrographic measure of IISs developed in Figure 2 (black axis labels), measures from patch-clamp recordings of individual GCs from the same animals (purple axis labels) and a summary score of behavioral discrimination impairment described in Figure 3C (green axis labels). Not all three types of records are available for every animal, which leads to differences in the samples shown in each panel.

(A) Data points correspond to individual GCs from animals in which both IISs and patch-clamp data were recorded. Each animal has a unique average hourly IIS index value. Only a subset of GCs are outside of the Ctrl range (i.e. pathological). For example, in the mouse with the highest IIS index (~12), only one of the three recorded GCs exhibited a pathological FR (~8 Hz, left panel) and all had a similar R_{output} slightly above the Ctrl range (~0.5, right panel). In contrast, some GCs with abnormally high FR (~6-12 Hz)_and R_{output} (~0.5-0.6) came from animals with low IIS index (~0-2).

(B) Data points correspond to individual animals in which both IISs and BPS were measured. In KA animals, low IIS index can correspond to either normal or abnormal mnemonic discrimination, and despite having a high IIS index, some animals have normal discrimination.

(C) Data points correspond to individual GCs from animals in which both BPS and patch-clamp data were recorded. In some KA mice with large behavioral impairments and multiple recorded GCs, some GCs exhibited normal pattern separation whereas others showed pathologically low levels. The relationship is unclear, but all KA mice with large impairments had at least one pathological GC.

(D) Data points correspond to individual GCs from animals in which IISs, BPS and patch-clamp data were all collected. Stems show which GCs were recorded from each animal. The lower right corner would correspond to the highest pathology in electrographic, behavioral and computational categories. Despite any heterogeneity or apparent ambiguity when only one or two of these measures are considered (above), when all three measures are available for each animal, the KA group is very clearly separable from the Ctrl group.

Discussion

We provide the first experimental evidence that TLE is characterized by both mnemonic discrimination impairments and neuronal pattern separation deficits, associated with pathological spiking behavior in a subset of GCs.

Cognitive and computational deficits in TLE. Our findings are consistent with the few previous studies that focused on similar questions. First, our data in human patients confirm the findings of Reyes et al. (2018) that TLE leads to mnemonic discrimination impairments. However, they used a different behavioral paradigm: a short-term memory object-location discrimination task akin to the task we used for mice. Although animal studies suggest that DG is necessary for such spatial mnemonic discrimination (Hunsaker and Kesner, 2013; Bui et al., 2018), it remains unclear whether this is the case in humans. In contrast, we used a non-spatial memory discrimination task that is known to be DG-dependent in humans (Baker et al., 2016; Stark et al., 2019), allowing us to conclude that impairments in TLE patients are likely due to DG alterations.

Our behavioral experiment in mice (**Figure 3**) also complements a previous study (Bui et al., 2018) by testing short-term (minutes) rather than long-term (24h) object-location memory and by testing multiple, parameterized levels of mnemonic interference, as recommended for rigorous testing of mnemonic discrimination (Hunsaker and Kesner, 2013; Liu et al., 2016). As in humans, the BPS task we used in mice is likely dependent on DG activity (Bui et al., 2018).

Taken together, our results and those studies suggest that mnemonic discrimination deficits

observed in TLE are shared between humans and mice, occur for different types of memory and

modalities, and point to DG malfunctions.

Indeed, DG normally functions as a pattern separator (Madar et al., 2019a) which is

hypothesized to support mnemonic discrimination. However, only one modelling study has so far

investigated the impact of TLE-related DG pathologies on neural pattern separation (Yim et al.,

2014). This model suggested that mossy fiber sprouting and increased synaptic transmission

from the perforant path to GCs, both hallmarks of TLE, can theoretically lead to a breakdown of

pattern separation. Our results experimentally confirm that TLE leads to a deficit in neural

pattern separation (**Figure 4**). However, the deficit we observed is not as large, possibly because

the model 1) considered synaptic transmission as deterministic rather than probabilistic and

plastic, 2) assessed a different form of pattern separation (patterns were defined as population

ensembles of very short spiketrains, thus focusing on a population code whereas we investigated

temporal codes) and 3) considered more severe degrees of epilepsy than was perhaps the case for

our mice.

Note that by using several similarity metrics, timescales and input statistical structures,

and thus exploring various potential neural codes, we also discovered that other normal DG

computations performed through rate and burstiness codes are perturbed in TLE (**Figure 5, 8**).

The role of sparsity in neuronal pattern separation and DG gating. We found that GCs

from epileptic mice exhibit pathological spiking consistent with a breach of the DG gatekeeper

function (**Figure 6**) that leads to decreased neuronal pattern separation (**Figure 8A**). As noted by

others, the gating and pattern separation functions of DG may be related (Dengler and Coulter,

2016) because the filtering of incoming cortical activity into a sparser output can be a simple

mechanism underlying a form of pattern separation based on a pure population code (O'Reilly

and McClelland, 1994; Severa et al., 2016; Dieni et al., 2016; Cayco-Gajic and Silver, 2019).

Our results suggest that maintaining a sparse output also supports pattern separation of

spiketrains in the time domain, at the level of single DGs.

Rare investigations of the spiking output of single GCs in epileptic tissue have reported

that single stimulation of the perforant path sometimes leads GCs to fire bursts of spikes (Lynch

et al., 2000; Kobayashi and Buckmaster, 2003). We expand on this by showing that bursts also

emerge in response to complex, naturalistic stimulation patterns (**Figures 6 and 7**). It is

important to note that occasional bursting is observed in normal GCs in vivo (Pernía-Andrade

and Jonas, 2014), but this is very unusual in slices (Mongiat et al., 2009; Ewell and Jones, 2010;

Zhang et al., 2012; Dieni et al., 2016). In our experiments, bursting GCs therefore demonstrate a

breach of the dentate gate. Because bursts are known to be more efficient at driving spiking in

CA3 pyramidal neurons (Henze et al., 2002), a single GC firing bursts will have a higher

probability of making downstream CA3 pyramidal cells fire. This could have deleterious effects

on memory encoding and promote seizures, as more active CA3 neurons would 1) increase the

chance of overlap between memory representations and 2) overexcite a recurrent excitatory

circuit, a mechanism of seizure generation. Thus, our findings are consistent with the idea that

pathological firing or bursting of even a subset of GCs could simultaneously contribute to

deficits in mnemonic discrimination and epileptiform activity in TLE.

Our study also provides new insight on the mechanisms underlying GCs increased ability

to burst by testing the influence of the excitatory drive (**Figure 7B**). The main source of

excitation to GCs, the perforant path, is sometimes assumed to be augmented in TLE (Yim et al.,

2014) given evidence of increased release probability at the lateral perforant path synapse

(Scimemi et al., 2006). Our results do not directly contradict this, but we did not find a difference

in the average synaptic current evoked in GCs by Poisson input spiketrains applied to the

perforant path (**Figure 7B3**). More work is required to carefully test potential TLE-related

changes in the coupling between perforant path and GCs, as well as EPSC kinetics and short-

term dynamics (Madar et al., 2019a), but it is clear that the size of excitatory drive alone is not a

good predictor of bursting propensity in GCs (**Figure 7B3-4**). We also previously showed that

partial blockade of inhibition can produce GC bursting and pattern separation deficits (Madar et

al., 2019b). Thus, in accordance with past research (Kobayashi and Buckmaster, 2003), we

conclude that the strong decrease in synaptic inhibition characteristic of TLE (Dengler and

Coulter, 2016; Dengler et al., 2017) is likely the main driver of pathological bursting.

Perhaps the most intriguing insight we bring on the failure of the DG as a gate and

pattern separator in TLE is that this failure occurs only in a subset of GCs, upon a background of

apparently normal GCs in the same the epileptic network. This finding invites many more

questions: In the epileptic DG, is there a spectrum of spiking patterns among GCs going from

normal to pathological, or are there two discrete types? What is the ratio of normal to

pathological GCs? Does this ratio evolve during epileptogenesis? What makes individual GCs

become pathological? Is that due to differences in intrinsic properties or networking? As

explained above, our results suggest deficits in inhibition, but other anatomical or biophysical

changes could be in play. For example, in individual mice with TLE, some GCs display

pathological hilar basal dendrites that have been associated with increased intrinsic excitability

that might allow them to fire bursts more easily (Kelly and Beck, 2017). It is also possible that

there exist heterogeneity in GC mossy fiber sprouting, with some GCs receiving recurrent

excitatory connections and some not. Some of these differences could be related to different

birthdates of adult-born GCs relative to an epileptogenic brain insult, resulting in only a subset of

GCs integrating abnormally into the DG network (Kron et al., 2010). Alternatively, pathological

GCs could develop from a more susceptible and active subclass of mature GCs (Erwin et al.,

2019). In any case, the discovery that the GC population in the epileptic DG is not homogeneous

in terms of spiking patterns is a crucial step towards developing new treatments that would

specifically target pathological neurons to avoid deleterious side-effects (Bielefeld et al., 2014).

The link between mnemonic discrimination, pattern separation and TLE. For thirty

years the DG has been hypothesized to support mnemonic discrimination by performing

neuronal pattern separation (Treves et al., 2008), but no experiment has yet directly connected

DG input-output computations with memory. Important reports have shown that 1) abnormal

neurotransmission in the DG can lead to overlapping spatial representations in CA3 as well as

impairments in context discrimination learning (McHugh et al., 2007) and 2) that overlapping

engrams in DG are a cause of confusion between the associated memories (Ramirez et al., 2013).

However, those reports did not investigate whether pattern separation was performed in the DG

per se, as opposed to upstream areas. Here, we confirm our previous work showing that the DG

itself performs temporal pattern separation (Madar et al., 2019a, b) and we show that, on

average, a deficit in pattern separation is associated with a deficit in mnemonic discrimination. A

full demonstration of the causal link will require measuring and manipulating DG computations

in vivo during episodic memory tasks, but our study is a preliminary step toward understanding

how DG circuit-level computations relate to a high-level cognitive function and how these

processes fail in TLE. A mechanistic understanding of this relation may lead to new early

diagnosis tools (e.g. cognitive tests like in Figure 1) as well as therapies alleviating memory

disorders in TLE. For example, our finding that a subset of GCs with pattern separation deficits

can develop without EEG hallmarks of epilepsy (**Figure 9A**) suggests a mechanism for memory

impairments that often occur in early stages of human epileptogenesis (Jones et al., 2007; Witt

and Helmstaedter, 2015) and the coexistence of pathological and normal neurons could explain

memory impairments without hippocampal sclerosis (Rayner et al., 2019).

References

Alexander A, Maroso M, Soltesz I (2016) Organization and control of epileptic circuits in temporal lobe epilepsy. Progress in brain research 226:127-154.

Artinian J, Peret A, Marti G, Epsztein J, Crepel V (2011) Synaptic kainate receptors in interplay with INaP shift the sparse firing of dentate granule cells to a sustained rhythmic mode in temporal lobe epilepsy. The Journal of neuroscience : the official journal of the Society for Neuroscience 31:10811-10818.

Baker S, Vieweg P, Gao F, Gilboa A, Wolbers T, Black SE, Rosenbaum RS (2016) The Human Dentate Gyrus Plays a Necessary Role in Discriminating New Memories. Current biology : CB.

Behr J, Lyson KJ, Mody I (1998) Enhanced propagation of epileptiform activity through the kindled dentate gyrus. Journal of neurophysiology 79:1726-1732.

Bennett IJ, Stark CE (2016) Mnemonic discrimination relates to perforant path integrity: An ultra-high resolution diffusion tensor imaging study. Neurobiology of learning and memory 129:107-112.

Berron D, Schutze H, Maass A, Cardenas-Blanco A, Kuijf HJ, Kumaran D, Duzel E (2016) Strong Evidence for Pattern Separation in Human Dentate Gyrus. The Journal of neuroscience : the official journal of the Society for Neuroscience 36:7569-7579.

Bielefeld P, van Vliet EA, Gorter JA, Lucassen PJ, Fitzsimons CP (2014) Different subsets of newborn granule cells: a possible role in epileptogenesis? The European journal of neuroscience 39:1-11.

Bui AD, Nguyen TM, Limouse C, Kim HK, Szabo GG, Felong S, Maroso M, Soltesz I (2018) Dentate gyrus mossy cells control spontaneous convulsive seizures and spatial memory. Science 359:787-790.

Cayco-Gajic NA, Silver RA (2019) Re-evaluating Circuit Mechanisms Underlying Pattern Separation. Neuron 101:584-602.

Chavlis S, Poirazi P (2017) Pattern separation in the hippocampus through the eyes of computational modeling. Synapse (New York, NY).

Coras R, Pauli E, Li J, Schwarz M, Rossler K, Buchfelder M, Hamer H, Stefan H, Blumcke I (2014) Differential influence of hippocampal subfields to memory formation: insights from patients with temporal lobe epilepsy. Brain : a journal of neurology 137:1945-1957.

Coulter DA, Carlson GC (2007) Functional regulation of the dentate gyrus by GABA-mediated inhibition. 163:235-812.

De Maesschalck R, Jouan-Rimbaud D, Massart DL (2000) The mahalanobis distance. Chemometrics and intelligent laboratory systems 50:1-18.

Dengler CG, Coulter DA (2016) Normal and epilepsy-associated pathologic function of the dentate gyrus. Progress in brain research 226:155-178.

Dengler CG, Yue C, Takano H, Coulter DA (2017) Massively augmented hippocampal dentate granule cell activation accompanies epilepsy development. Scientific reports 7:42090.

Dieni CV, Panichi R, Aimone JB, Kuo CT, Wadiche JI, Overstreet-Wadiche L (2016) Low excitatory innervation balances high intrinsic excitability of immature dentate neurons. Nature communications 7:11313.

Dillon SE, Tsivos D, Knight M, McCann B, Pennington C, Shiel AI, Conway ME, Newson MA, Kauppinen RA, Coulthard EJ (2017) The impact of ageing reveals distinct roles for human dentate gyrus and CA3 in pattern separation and object recognition memory. Scientific reports 7:14069.

Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2019) A sparse, spatially biased subtype of mature granule cell is preferentially recruited in hippocampal-associated behaviors. bioRxiv:804393.

Ewell LA, Jones MV (2010) Frequency-tuned distribution of inhibition in the dentate gyrus. The Journal of neuroscience : the official journal of the Society for Neuroscience 30:12597-12607.

Groticke I, Hoffmann K, Loscher W (2008) Behavioral alterations in a mouse model of temporal lobe epilepsy induced by intrahippocampal injection of kainate. Experimental neurology 213:71-83.

948   Hellier JL, Patrylo PR, Buckmaster PS, Dudek FE (1998) Recurrent spontaneous motor seizures after repeated low-
949   dose systemic treatment with kainate: assessment of a rat model of temporal lobe epilepsy. Epilepsy
950   research 31:73-84.
951   Helmstaedter C, Kurthen M, Lux S, Reuber M, Elger CE (2003) Chronic epilepsy and cognition: a longitudinal
952   study in temporal lobe epilepsy. Annals of neurology 54:425-432.
953   Henze DA, Wittner L, Buzsaki G (2002) Single granule cells reliably discharge targets in the hippocampal CA3
954   network in vivo. Nature neuroscience 5:790-795.
955   Hsu D (2007) The dentate gyrus as a filter or gate: a look back and a look ahead. Progress in brain research 163:601-
956   613.
957   Hunsaker MR, Kesner RP (2013) The operation of pattern separation and pattern completion processes associated
958   with different attributes or domains of memory. Neuroscience and biobehavioral reviews 37:36-58.
959   Inostroza M, Brotons-Mas JR, Laurent F, Cid E, de la Prida LM (2013) Specific impairment of "what-where-when"
960   episodic-like memory in experimental models of temporal lobe epilepsy. Journal of Neuroscience
961   33:17749-17762.
962   Jones JE, Watson R, Sheth R, Caplan R, Koehn M, Seidenberg M, Hermann B (2007) Psychiatric comorbidity in
963   children with new onset epilepsy. Developmental medicine and child neurology 49:493-497.
964   Kelly T, Beck H (2017) Functional properties of granule cells with hilar basal dendrites in the epileptic dentate
965   gyrus. Epilepsia 58:160-171.
966   Kesner RP, Rolls ET (2015) A computational theory of hippocampal function, and tests of the theory: new
967   developments. Neuroscience and biobehavioral reviews 48:92-147.
968   Kesner RP, Kirk RA, Yu Z, Polansky C, Musso ND (2016) Dentate gyrus supports slope recognition memory,
969   shades of grey-context pattern separation and recognition memory, and CA3 supports pattern completion
970   for object memory. Neurobiology of learning and memory 129:29-37.
971   Knierim JJ, Neunuebel JP (2016) Tracking the flow of hippocampal computation: Pattern separation, pattern
972   completion, and attractor dynamics. Neurobiology of learning and memory 129:38-49.
973   Kobayashi M, Buckmaster PS (2003) Reduced inhibition of dentate granule cells in a model of temporal lobe
974   epilepsy. The Journal of neuroscience : the official journal of the Society for Neuroscience 23:2440-2452.
975   Kron MM, Zhang H, Parent JM (2010) The developmental stage of dentate granule cells dictates their contribution
976   to seizure-induced plasticity. The Journal of neuroscience : the official journal of the Society for
977   Neuroscience 30:2051-2059.
978   Krook-Magnuson E, Armstrong C, Bui A, Lew S, Oijala M, Soltesz I (2015) In vivo evaluation of the dentate gate
979   theory in epilepsy. The Journal of physiology 593:2379-2388.
980   Le Duigou C, Simonnet J, Telenczuk MT, Fricker D, Miles R (2014) Recurrent synapses and circuits in the CA3
981   region of the hippocampus: an associative network. Frontiers in cellular neuroscience 7:262.
982   Lee CT, Kao MH, Hou WH, Wei YT, Chen CL, Lien CC (2016) Causal Evidence for the Role of Specific
983   GABAergic Interneuron Types in Entorhinal Recruitment of Dentate Granule Cells. Scientific reports
984   6:36885.
985   Lenck-Santini PP, Scott RC (2015) Mechanisms Responsible for Cognitive Impairment in Epilepsy. Cold Spring
986   Harbor perspectives in medicine 5.
987   Levesque M, Avoli M (2013) The kainic acid model of temporal lobe epilepsy. Neuroscience and biobehavioral
988   reviews 37:2887-2899.
989   Levesque M, Avoli M, Bernard C (2016) Animal models of temporal lobe epilepsy following systemic
990   chemoconvulsant administration. Journal of neuroscience methods 260:45-52.
991   Lilliefors HW (1967) On the Kolmogorov-Smirnov test for normality with mean and variance unknown. Journal of
992   the American statistical Association 62:399-402.
993   Liu KY, Gould RL, Coulson MC, Ward EV, Howard RJ (2016) Tests of pattern separation and pattern completion
994   in humans-A systematic review. Hippocampus 26:705-717.
995   Lu Y, Zhong C, Wang L, Wei P, He W, Huang K, Zhang Y, Zhan Y, Feng G, Wang L (2016) Optogenetic
996   dissection of ictal propagation in the hippocampal–entorhinal cortex structures. Nature communications
997   7:10962.
998   Lynch M, Sayin U, Golarai G, Sutula T (2000) NMDA receptor-dependent plasticity of granule cell spiking in the
999   dentate gyrus of normal and epileptic rats. Journal of neurophysiology 84:2868-2879.
Madar AD, Ewell LA, Jones MV (2019a) Pattern separation of spiketrains in hippocampal neurons. Scientific
reports 9:5282.
Madar AD, Ewell LA, Jones MV (2019b) Temporal pattern separation in hippocampal neurons through multiplexed
neural codes. PLoS computational biology 15:e1006932.

McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB, Fanselow MS, Wilson MA,
Tonegawa S (2007) Dentate gyrus NMDA receptors mediate rapid pattern separation in the hippocampal
network. Science 317:94-99.
Mongiat LA, Esposito MS, Lombardi G, Schinder AF (2009) Reliable activation of immature neurons in the adult
hippocampus. PloS one 4:e5320.
Motulsky HJ, Ransnas LA (1987) Fitting curves to data using nonlinear regression: a practical and nonmathematical
review. FASEB journal : official publication of the Federation of American Societies for Experimental
Biology 1:365-374.
Muller CJ, Groticke I, Bankstahl M, Loscher W (2009) Behavioral and cognitive alterations, spontaneous seizures,
and neuropathology developing after a pilocarpine-induced status epilepticus in C57BL/6 mice.
Experimental neurology 219:284-297.
O'Reilly RC, McClelland JL (1994) Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off.
Hippocampus 4:661-682.
Ouedraogo DW, Lenck-Santini PP, Marti G, Robbe D, Crepel V, Epsztein J (2016) Abnormal UP/DOWN
Membrane Potential Dynamics Coupled with the Neocortical Slow Oscillation in Dentate Granule Cells
during the Latent Phase of Temporal Lobe Epilepsy. eNeuro 3.
Pardi MB, Ogando MB, Schinder AF, Marin-Burgin A (2015) Differential inhibition onto developing and mature
granule cells generates high-frequency filters with variable gain. eLife 4:e08764.
Patrylo PR, Schweitzer JS, Dudek FE (1999) Abnormal responses to perforant path stimulation in the dentate gyrus
of slices from rats with kainate-induced epilepsy and mossy fiber reorganization. Epilepsy research 36:31-
42.
Pernía-Andrade Alejandro J, Jonas P (2014) Theta-Gamma-Modulated Synaptic Currents in Hippocampal Granule
Cells In Vivo Define a Mechanism for Network Oscillations. Neuron 81:140-152.
Pfammatter JA, Bergstrom RA, Wallace EP, Maganti RK, Jones MV (2018) A predictive epilepsy index based on
probabilistic classification of interictal spike waveforms. PloS one 13:e0207158.
Racine RJ (1972) Modification of seizure activity by electrical stimulation: II. Motor seizure.
Electroencephalography and clinical neurophysiology 32:281-294.
Ramirez S, Liu X, Lin PA, Suh J, Pignatelli M, Redondo RL, Ryan TJ, Tonegawa S (2013) Creating a false memory
in the hippocampus. Science 341:387-391.
Rayner G, Tailby C, Jackson G, Wilson S (2019) Looking beyond lesions for causes of neuropsychological
impairment in epilepsy. Neurology 92:e680-e689.
Reyes A, Holden HM, Chang YA, Uttarwar VS, Sheppard DP, DeFord NE, DeJesus SY, Kansal L, Gilbert PE,
McDonald CR (2018) Impaired spatial pattern separation performance in temporal lobe epilepsy is
associated with visuospatial memory deficits and hippocampal volume loss. Neuropsychologia 111:209-
215.
Rolls ET (2010) A computational theory of episodic memory formation in the hippocampus. Behavioural brain
research 215:180-196.
Santoro A (2013) Reassessing pattern separation in the dentate gyrus. Frontiers in behavioral neuroscience 7:96.
Scimemi A, Schorge S, Kullmann DM, Walker MC (2006) Epileptogenesis is associated with enhanced
glutamatergic transmission in the perforant path. Journal of neurophysiology 95:1213-1220.
Severa W, Parekh O, James CD, Aimone JB (2016) A Combinatorial Model for Dentate Gyrus Sparse Coding.
Neural computation:1-24.
Sharma S, Puttachary S, Thippeswamy A, Kanthasamy AG, Thippeswamy T (2018) Status epilepticus: Behavioral
and electroencephalography Seizure Correlates in Kainate experimental Models. Frontiers in neurology 9:7.
Simon P, Dupuis R, Costentin J (1994) Thigmotaxis as an index of anxiety in mice. Influence of dopaminergic
transmissions. Behavioural brain research 61:59-64.
Sloviter RS (1983) "Epileptic" brain damage in rats induced by sustained electrical stimulation of the perforant path.
I. Acute electrophysiological and light microscopic studies. Brain research bulletin 10:675-697.
Stark SM, Kirwan CB, Stark CEL (2019) Mnemonic Similarity Task: A Tool for Assessing Hippocampal Integrity.
Trends in cognitive sciences.
Stark SM, Yassa MA, Lacy JW, Stark CE (2013) A task to assess behavioral pattern separation (BPS) in humans:
Data from healthy aging and mild cognitive impairment. Neuropsychologia 51:2442-2449.
Tellez-Zenteno JF, Hernandez-Ronquillo L (2012) A review of the epidemiology of temporal lobe epilepsy.
Epilepsy research and treatment 2012:630853.

Toyoda I, Bower MR, Leyva F, Buckmaster PS (2013) Early activation of ventral hippocampus and subiculum
during spontaneous seizures in a rat model of temporal lobe epilepsy. The Journal of neuroscience : the
official journal of the Society for Neuroscience 33:11100-11115.
Treves A, Tashiro A, Witter MP, Moser EI (2008) What is the mammalian dentate gyrus good for? Neuroscience
154:1155-1172.
van Hagen BT, van Goethem NP, Lagatta DC, Prickaerts J (2015) The object pattern separation (OPS) task: a
behavioral paradigm derived from the object recognition task. Behavioural brain research 285:44-52.
Wallace E, Kim DY, Kim KM, Chen S, Blair Braden B, Williams J, Jasso K, Garcia A, Rho JM, Bimonte-Nelson H,
Maganti R (2015) Differential effects of duration of sleep fragmentation on spatial learning and synaptic
plasticity in pubertal mice. Brain research 1615:116-128.
Witt JA, Helmstaedter C (2015) Cognition in the early stages of adult epilepsy. Seizure 26:65-68.
Yassa MA, Mattfeld AT, Stark SM, Stark CE (2011) Age-related memory deficits linked to circuit-specific
disruptions in the hippocampus. Proceedings of the National Academy of Sciences of the United States of
America 108:8873-8878.
Yi F, Catudio-Garrett E, Gabriel R, Wilhelm M, Erdelyi F, Szabo G, Deisseroth K, Lawrence J (2015) Hippocampal
"cholinergic interneurons" visualized with the choline acetyltransferase promoter: anatomical distribution,
intrinsic membrane properties, neurochemical characteristics, and capacity for cholinergic modulation.
Frontiers in synaptic neuroscience 7:4.
Yim MY, Hanuschkin A, Wolfart J (2014) Intrinsic rescaling of granule cells restores pattern separation ability of a
dentate gyrus network model during epileptic hyperexcitability. Hippocampus.
Zhang W, Huguenard JR, Buckmaster PS (2012) Increased excitatory synaptic input to granule cells from hilar and
CA3 regions in a rat model of temporal lobe epilepsy. The Journal of neuroscience : the official journal of
the Society for Neuroscience 32:1183-1196.
Zhao F, Kang H, You L, Rastogi P, Venkatesh D, Chandra M (2014) Neuropsychological deficits in temporal lobe
epilepsy: A comprehensive review. Annals of Indian Academy of Neurology 17:374-382.
1083
1084

1085