Title: High Sucrose Diets Contribute to Brain Angiopathy with Impaired Glucose

Uptake, and Psychosis-related Higher Brain Dysfunctions in Mice

Short title: Angiopathy in mice and humans with psychosis

Authors: Shinobu Hirai[1,†], Hideki Miwa[1,2], Tomoko Tanaka[1], Kazuya Toriumi[3],

Yasuto Kunii[4], Hiroko Shimbo[1], Takuya Sakamoto[5], Mizuki Hino[4], Ryuta Izumi[4],

Atsuko Nagaoka[4], Hirooki Yabe[4], Tomoya Nakamachi[6], Seiji Shioda[7], Takashi Dan[8],

Toshio Miyata[8], Yasumasa Nishito[9], Kazuhiro Suzuki[3], Mitsuhiro Miyashita[3],

Toshifumi Tomoda[10], Takatoshi Hikida[11], Junjiro Horiuchi[12], Masanari Itokawa[3],

Makoto Arai[3], Haruo Okado[1,†]

Affiliations: [1]Neural Development Project, Department of Brain Development and Neural

Regeneration, Tokyo Metropolitan Institute of Medical Science, Tokyo 156-8506, Japan.

[2]Molecular Neuropsychopharmacology Section, Department of Neuropsychopharmacology,

National Institute of Mental Health, National Center of Neurology and Psychiatry, Tokyo

187-8553, Japan.

[3]Schizophrenia Research Project, Department of Psychiatry and Behavioral Sciences, Tokyo

Metropolitan Institute of Medical Science, Tokyo 156-8506, Japan.

[4]Department of Neuropsychiatry, School of Medicine, Fukushima Medical University,

Fukushima 960-1295, Japan.

[5]Department of Applied Biological Science, Faculty of Science and Technology, Tokyo

University of Science, 2641 Yamazaki, Noda, Chiba 278-8510, Japan.

[6]Laboratory of Regulatory Biology, Graduate School of Science and Engineering,

University of Toyama, Toyama 930-8555, Japan.

[7] Department of Clinical Pharmacy, Shonan University of Medical Sciences, Yokohama

244-0806, Japan

[8]Division of Molecular Medicine and Therapy, Tohoku University Graduate School of

Medicine, Miyagi 980-8575, Japan.

[9]Center for Basic Technology Research, Tokyo Metropolitan Institute of Medical Science,

Tokyo 156-8506, Japan.

[10]Centre for Addiction and Mental Health, Campbell Family Mental Health Research

Institute, University of Toronto, Toronto, Ontario M5T 1R8, Canada.

[11]Department of Laboratory for Advanced Brain Functions, Institute for Protein

Research, *Osaka* University, Osaka 565-0871, Japan.

[12]Tokyo Metropolitan Institute of Medical Science, Tokyo 156-8506, Japan.

[ここに入力]

†These authors contributed equally to this work.

Correspondence should be addressed to Shinobu Hirai and Haruo Okado, Neural

Development Project, Department of Brain Development and Neural Regeneration, Tokyo

Metropolitan Institute of Medical Science, Tokyo 156-8506, Japan.

E-mail: hirai-sn@igakuken.or.jp (S.H.) and okado-hr@igakuken.or.jp (H.O.)

Abstract: Metabolic dysfunction is thought to contribute to the severity of psychiatric

disorders; however, it has been unclear whether current high-simple-sugar diets contribute

to pathogenesis of these diseases. Here we demonstrate that a high-sucrose diet during

adolescence induces psychosis-related behavioral endophenotypes, including hyperactivity,

poor working memory, impaired sensory gating, and disrupted interneuron function in mice

deficient for glyoxalase-1 (Glo1), an enzyme involved in detoxification of sucrose

metabolites. Further, the high-sucrose diet induced microcapillary impairments and reduced

brain glucose uptake in brains of *Glo1* deficient mice. Aspirin protected against this

angiopathy, enhancing brain glucose uptake, and preventing abnormal behavioral

phenotypes. Similar vascular damage to our model mice was found in the brains of

randomly collected schizophrenia and bipolar disorder patients, suggesting that psychiatric

[ここに入力]

disorders are associated with angiopathy in the brain caused by various environmental

stresses, including metabolic stress.

Introduction:

Considering the global increase in consumption of simple sugars, the World Health

Organization published guidelines that addressed concerns regarding body weight gain and

dental caries development in 1995 (*1*). High-sugar intake alone increases the risks of

numerous chronic diseases, including diabetes, hypertension, and kidney disease. However,

there are few studies on the effects of high-sugar intake during adolescence on future mental

health. The daily caloric intake from simple sugar by teenagers is higher than that observed

for other age groups (~20% of total daily caloric intake) (*2*). Most chronic psychiatric

disorders, including schizophrenia (SZ) and bipolar disorder (BD) discussed in the present

study, develop before the age of 30 via complex interactions among multiple genetic and

environmental risk factors. Interestingly, patients with SZ and BD consume approximately

2-fold more sugar than age-matched healthy individuals, and patients with SZ who consume

more sucrose exhibit more severe symptoms (*3-5*). Moreover, the odds ratios for mental

distress, hyperactivity, and behavioral disorders were the highest among adolescents who

self-reported the highest consumption of soft drinks (*6*).

[ここに入力]

Advanced glycation end products (AGEs) and reactive carbonyl compounds are

produced from irreversible non-enzymatic crosslinking reactions between

carbonyl-containing molecules (e.g. sugars, reactive carbonyl compounds) and other

molecules (e.g. amino groups of proteins), and AGEs can lead to the induction of

inflammation and oxidative stress in various tissues, leading to a positive feedback loop that

further increases AGEs (*7*). There is substantial evidence that patients with psychiatric

disorders have elevated proinflammatory cytokines (e.g. interleukin (IL)-1β, IL-6) and

increased oxidative stress (*8-11*).   Glyoxalase I (GLO1) is a zinc metalloenzyme that

protects cells from AGE toxicity by catalyzing the reaction between the reactive carbonyl

compound methylglyoxal to glutathione to form S-lactoyl-glutathione (*12*). Patients with

depressive-state BD and major depressive disorder have reduced amounts of GLO1,

suggesting that this reduction, in combination with an increase in sugar consumption, may be

a cause of the increased inflammation and oxidative stress observed in these diseases (*13*).

Furthermore, a SZ patient exhibiting poor convalescence is found to harbor a frameshift

mutation in *GLO1* leading to reduced enzyme activity (*14, 15*).

Despite accumulating evidence, it is still unproven that excessive sugar intake

contributes to the pathogenesis of psychiatric disorders among susceptible individuals. We

addressed this question by generating mice with psychosis-related phenotypes associated

[ここに入力]

the gene × environment interactions (G × E) to determine whether excessive sucrose

consumption during adolescence is a novel environmental risk factor for SZ and BD. We

further identified angiopathy in mice with high-sucrose intake and genetic venerability, and

confirmed similar vascular damage in randomly collected brain samples from BD and SZ

patients. This suggests that a variety of genetic and environmental factors can adversely

affect brain capillaries. In addition, we found that chronic anti-inflammatory drug treatment

prevented core phenotypes of psychiatric disorders as well as vascular damage, suggesting

that angiopathy may have an important influence on the development and/or condition of

psychiatric disorders.

Results

Psychosis-related behavioral phenotypes in mice on a high-sucrose diet

We fed mice one of two diets containing the same total calories and caloric proportions

of carbohydrates, fat, and proteins, but with either starch or sucrose as the main

carbohydrate (Fig. 1A). We investigated four groups of mice fed these diets for 50 days

immediately after weaning (from postnatal day 21, corresponding to the juvenile/adolescent

stage): starch-fed wild-type (WT) mice (control, CTL), sucrose-fed WT mice

(environmental stressor, Env), starch-fed *Glo1* heterozygous knockout (*Glo1/+*) mice

[ここに入力]

(genetic factor, Gen), and sucrose-fed *Glo1/+* mice (G × E) (Fig. 1A). Reduced Glo1

expression in *Glo1/+* mice was confirmed in the cerebral cortices, including the

hippocampus, using western blotting (fig. S1A, B). The body weight trajectories of these

mice were similar to those of control mice for at least 11 weeks of age, indicative of normal

structural development (fig. S1C) and removing obesity as a possible cause of the observed

group differences described below. We tested eight different behaviors in mice, including

open-field locomotor activity (Fig. 1B), acoustic startle response (Fig. 1C), pre-pulse

inhibition (PPI; a measure of sensory-motor gating) (Fig. 1D), object location performance

(used as a test of working memory) (Fig. 1E), self-grooming (Fig. 1F), nest building (Fig.

1G), elevated plus maze activity (fig. S2A), and social interaction (fig. S2B). These

behaviors could be classified into those that are not affected by either diet or genotype,

those that are affected by genotype but not diet, those affected by diet but not genotype, and

those affected by the combination of diet and genotype. Thus, neither diet nor genotype, nor

a combination of diet and genotype had any significant effect on social interactions (fig.

S2B), while acoustic startle responses were reduced in *Glo1/+* mice but not significantly

affected by feeding type (Fig. 1C). Self-grooming (Fig. 1F), nest building (Fig. 1G), and

elevated plus maze activity (fig. S2A) were affected by diet, but not genotype, while open

field locomotor activity, PPI, and object location performance required a combination of

[ここに入力]

sucrose feeding and a *Glo1/+* genotype to be affected. In the open field test (Fig. 1B),

neither starch-fed *Glo1/+* mice, nor sucrose-fed WT mice, showed significant differences in

distance traveled compared to starch-fed WT mice, while sucrose-fed *Glo1/+* mice showed

significant increases. Similarly, starch-fed *Glo1/+* mice and sucrose-fed WT mice had

similar PPI scores compared to starch-fed WT mice, while sucrose-fed *Glo1/+* mice showed

significant decreases (Fig. 1D). Sucrose-fed WT mice had a non-significant reduction in

object location test scores, and this reduction became significant in sucrose-fed *Glo1/+*

mice (Fig. 1E). Altogether, our results suggest that a combination of sucrose feeding and

Glo1/+ mutation can generate novel phenotypes reminiscent of psychiatric disease

phenotypes that are not found by altering diet or genotype alone.

To further support our idea that synthetic G × E interactions may induce or enhance the

severity of psychosis-like symptoms, we examined the effects of the high sucrose diet on an

additional major psychosis-associated mutation in *Disrupted-in-schizophrenia 1* (*DISC1*)

locus (*16*) (fig. S3). DISC1 plays important roles in neurodevelopment and synaptic activity,

but a loss-of-function mutation in *DISC1* gene alone is not sufficient to cause psychiatric

disorders (*17*). Similar to the results for *Glo1/+* mice, *Disc1/+* mice fed a high sucrose diet (G

× E mice) showed severely impaired object location performance (working memory) and PPI

(fig. S3C, D), whereas WT mice fed a high sucrose diet or *Disc1/+* mice fed control diet did

[ここに入力]

not,   again suggesting that a combination of genetic and environmental influences can

induce psychosis-like behaviors.

Schizophrenia (SZ) is associated with increased amphetamine-induced dopamine release

in the striatum (*18, 19*) and unbalanced dopamine regulation between the medial prefrontal

cortex (mPFC) and striatum (*20*). To examine the effects of high sucrose diet on dopamine

release in *Glo1/+* mice, we next measured dopamine levels in the nucleus accumbens (NAc)

and mPFC of these mice using *in vivo* microdialysis. We found that the high sucrose diet

increased both basal and amphetamine-induced dopamine release in the NAc specifically in

Glo1/+ mice (Fig. 1H), while reducing basal dopamine release in the mPFC in either genetic

background (fig. S2C). To assess whether enhanced dopamine release in the NAc could

induce the observed behavioral phenotypes, we examined the effects of aripiprazole, a D2

receptor partial agonist and clinical antipsychotic (*21*), administered during the last 7 days (0.5

mg/kg/day) of sucrose feeding (Fig. 1A, I, J and fig. S2D–I). We found that the

hyper-locomotion and increased striatal dopamine release in G × E mice were completely

reversed by aripiprazole treatment (Fig. 1I, J), while other behavioral defects were not

improved (fig. S2D–I). Thus, aripiprazole treatment selectively improves a subset of

behavioral abnormalities in G × E mice that are likely caused by dysregulated dopamine

signaling.

[ここに入力]

Dysfunction of parvalbumin-positive inhibitory interneurons in G × E mice

A reduction in density of parvalbumin (PV)-positive GABAergic interneurons is

reported for schizophrenia (*22-25*),   and the precisely coordinated activity of these neurons

is crucial for the regulation of PPI and working memory (*26*). Thus, we next investigated

PV expression using immunohistochemistry and western blotting to determine whether

alterations in PV signaling may contribute to our observed psychiatric disease-associated

phenotypes. A high sucrose diet caused a reduction in PV-positive cells in both *Glo1/+* and

WT mice (Fig. 2A, B) and significantly reduced PV expression in sucrose-fed *Glo1/+* mice

(Fig. 2C, D), suggesting that the sucrose diet and the *Glo1*/+ mutation had additive or

synergistic effects on PV signaling. Since gamma oscillations are produced via the

synchronous activation of PV neurons, we measured gamma oscillations (30–45 Hz) using

surface electroencephalography (EEG) to determine whether downregulation of PV protein

was accompanied by functional abnormalities in neural activity. Interestingly, sucrose-fed

mice (WT and *Glo1*/+ mice) exhibited elevated baseline gamma oscillation power

compared with control mice (starch-fed WT) in the home cage (Fig. 2E). Further, only

sucrose-fed *Glo1*/+ mice failed to exhibit an increase in the gamma oscillation power when

approaching a novel object (Fig. 2F). These results are consistent with findings in patients

[ここに入力]

with SZ and BD as well as other mouse models of psychosis demonstrating increased

baseline gamma oscillations and decreased sensory stimulus-evoked synchronized gamma

power (*27, 28*). Thus, our results suggest that G × E mice mimic the pathophysiological

changes of PV neurons observed in psychiatric disorders.

To summarize, the administration of a high-sucrose diet to *Glo1/+* mice induces

behavioral, histological, and pathophysiological phenotypes reminiscent of phenotypes

observed in psychiatric disorders. This suggests that excessive sucrose intake during

adolescence is a potential environmental risk factor for these diseases.

AGE accumulation and impaired astrocyte function in G × E mice

How does a decrease in GLO1 expression and an increase in sucrose consumption

lead to defects in activity of PV and dopaminergic neurons? We first examined GLO1

expression and detected high expression in astrocytes (fig. S4A–G), especially those

surrounding capillaries (fig. S4B, C), moderate expression in neurons (fig. S4H, I), and low

expression (below detection limits) in vascular endothelial cells (fig. S4A–G) and

microglial cells (fig. S4H, J). Expression was not detected in the brains of *Glo1*

homozygous mice (fig. S3K) indicating that the antibody we used was specific for GLO1.

[ここに入力]

Since both GLO1 and a high sucrose diet are associated with glycation toxicity, we

next examined AGE immunoreactivity to detect levels of AGE products of glucose

metabolism and/or chemical reaction of glucose. We found stronger AGE immunoreactive

signals in the vascular endothelial cells of G × E mice compared with control mice (Fig.

3A–H). Moreover, we detected AGE products of methylglyoxal metabolism and/or

chemical reaction of methylglyoxal using AGE-4 antibody, in the microglia of sucrose-fed

mice (fig. S5A–E). Compared to controls, AGE accumulation in sucrose-fed *Glo1/+* mice

was accompanied by elevated IBA1 immunofluorescence intensity and an enlarged

CD68-positive area (fig. S5F, G, H), both of which are phenotypes identified in activated

microglia (*29, 30*). Thus, sucrose feeding in *Glo1/+* mice results in microglial and

endothelial AGE accumulation and microglial activation.

Since astrocytes express Glo1 most strongly (fig. S4A–G), we next used mice

expressing green fluorescent protein (GFP) under control of the glial fibrillary acid protein

(GFAP) promoter (*31, 32*) to evaluate how astrocytes are affected in our G x E condition.

Typically, cellular damage from AGEs is caused by inflammatory responses induced by

Receptor of AGEs (RAGE) activation or by a loss of normal protein function following

AGE-forming reactions (*33*). Astrocytes exhibit a well-described reactive phenotype in

response to pathogenic conditions including neuroinflammation characterized by enhanced

[ここに入力]

GFAP expression (*34, 35*). Strongly enhanced *GFAP* promoter function was observed in G

× E mice, without changes in the number of GFAP-positive astrocytes (Fig. 3I–K),

indicating that the astrocytes in G × E mice are in the reactive pre-condition during the

high-sucrose feeding. Taken together, these results suggest that a high sucrose diet enhances

AGE production in endothelial cells and microglia (Fig. 3A–H and fig. S5A–E), and

converts astrocytes into a pre-inflammatory state in *Glo1*-deficient mice (Fig. 3I–K).

Microcapillary angiopathy and impaired glucose intake in G × E mice

Endothelial cells and astrocytes are functional and physical components of the

blood-brain barrier (BBB) which tightly controls the parenchymal environment by

modulating the selective passage of nutrients and various factors (*36*). Thus, endothelial

AGE accumulation and astrocyte reactivity may impair BBB function. To examine changes

in endothelial function, we first conducted a transcriptome analysis of the prefrontal cortex

(PFC), a region strongly implicated in psychiatric impairments, using microarrays (Fig. 4A,

B). The coagulation factor V, essential for the production of fibrin from fibrinogen, ranked

seventh on a list of transcripts exhibiting more than doubled expression in G × E mice

compared with the other three groups (Fig. 4A, Supplementary Table 1, 2). Fibrin controls

hemostasis via polymerization with platelets to form blood clots, and deposits of this protein

[ここに入力]

are indicative of endothelial abnormalities (*37*). In early stages of endothelial cell

impairment, fibrin accumulates in capillaries. Therefore, we investigated vascular fibrin

accumulation using immunohistochemistry and confirmed the presence of significant fibrin

accumulation on the vascular lumen side of endothelial cells in brain capillaries of G × E

mice (Fig. 4C–G). Fibrin leakage and deposition in the brain parenchyma, as observed in

Alzheimer's disease, was not detected in G × E mice, indicating that physical BBB

disruption did not occur (*38*).

We next speculated that the abnormal vascular endothelial cells and astrocytes

observed in G × E mice could alter glucose uptake from the plasma into the brain

parenchyma. Extracellular glucose concentrations in the brain parenchyma were measured

under three conditions: 1) fasting, 2) 1 h after feeding, and 3) 2 h after feeding. We first

measured a baseline glucose concentration in the mPFC in starch-fed WT mice after

starvation for 16 hrs (Fig. 4H). This concentration increased significantly within an hour

after starch feeding, and decreased close to baseline 2 hrs after feeding. Sucrose-fed WT

mice and starch-fed *Glo1/+* mice had similar baseline glucose concentrations, similar

increases 1 hr after starch or sucrose feeding, and similar decreases back to baseline 2 hrs

after feeding. In contrast, sucrose-fed *Glo1/+* mice had both significantly decreased basal

glucose concentrations in the mPFC, and no apparent increase in glucose concentrations

[ここに入力]

after sucrose feeding. We measured vascular diameter and expression of glucose transporter

1, the major glucose transporter expressed in vascular endothelial cells and astrocytes, and

observed no differences among the four groups (fig. S6). Further, no differences were

detected in plasma glucose after starvation and feeding, or in fasting plasma insulin levels

among the four groups (Fig. 4I, J), indicating that the lower parenchymal glucose

concentrations in G × E mice is due to reduced uptake across the BBB rather than

dysregulation of plasma glucose or insulin signaling.

Protective effects of chronic low-dose aspirin against behavioral abnormalities and

angiopathy

Previous reports have shown that adjunct non-steroidal anti-inflammatory drug (NSAID)

treatment can improve psychiatric disorder scores (*39, 40*). Aspirin, an NSAID, is routinely

used for the prevention and alleviation of vascular-related adverse events associated with

high blood pressure, ischemia, and cardiovascular diseases (*41, 42*). Here, we examined

whether aspirin treatment can protect against the development of psychiatric phenotypes in

G × E mice (Fig. 1A, Fig. 5A–D and fig. S7A–I). Low-dose aspirin (1 mg/kg/day) fully

prevented hyperlocomotor activity, deficits in PPI, working memory, grooming duration

(Fig. 5A–D), and partially prevented abnormal enhancement of DA release after

[ここに入力]

methamphetamine administration and reduced PV expression among G × E mice (fig.

S7D–F), but did not improve acoustic startle responses, nest building and elevated plus

maze scores and abnormal astrocyte activation (fig. S7A–C, G–I). These improvements

were accompanied by a decrease in endothelial fibrin accumulation (Fig. 5E, F) and a

partial restoration of glucose intake into the brain parenchyma (Fig. 5G). Collectively, these

results suggest that aspirin treatment in G × E mice significantly improves angiopathy and

brain glucose availability, contributing to prevention of several behavioral abnormalities.

Angiopathy in postmortem brains of patients with psychiatric disorders

Finally, we compared immunostains of brain slices from healthy controls and patients with

SZ or BD to examine whether these patients exhibit angiopathic fibrin accumulation in

vascular endothelial cells. Similar to our findings in G × E mice, we found significantly

elevated fibrin accumulation in the vascular endothelium of brain slices from randomly

collected SZ and BD patients (Fig. 6A–C). Therefore, fibrin-related angiopathy and

vascular damage may be a novel and common phenotype of psychiatric illness contributing

to disease progression.

Discussion

[ここに入力]

Using a mouse model, we demonstrated that high dietary sucrose consumption during

adolescence is a potential risk factor for the development of behavioral phenotypes

associated with psychiatric illnesses, such as SZ and BD. These behavioral phenotypes

include impaired sensory gating, dysfunctions in PV interneurons and working memory,

hyperactivity, and increased basal and stimulus-evoked striatal dopamine release. Second,

we identified endothelial fibrin accumulation ("angiopathy") in both our mouse model and

in randomly collected postmortem brains of patients with SZ or BD. We also observed that

glucose intake from the plasma into the brain parenchyma was impaired in $G \times E$ mouse

model. Chronic low-dose aspirin treatment prevented fibrin deposition in the capillaries,

improved glucose transport, and reversed several behavioral phenotypes in $G \times E$ mice,

suggesting angiopathy as a seminal pathogenic event in mental illness.

In Fig. 6D we propose a possible model summarizing our results. A high sucrose diet

causes AGE accumulation in microglia and endothelial cells which express low amounts of

GLO1 (Fig. 3A–H and fig. S5A–E). Astrocytes are less sensitive to this diet in WT animals

because they express high amounts of GLO1 to relieve glycation stress (fig.

S4A–G).   However, the reduced GLO1 in *Glo1*/+ mice causes astrocytes to become

activated by a high sucrose diet (Fig. 3I–K). This activation may be caused directly by

glycation stress in astrocytes or may occur as a consequence of increased microglial and

[ここに入力]

endothelial AGEs. Astrocytic activation further increases microglial and endothelial AGEs

in a positive feedback loop (*7*). Increased microglial and endothelial AGEs and astrocytic

activation may induce fibrin accumulation and damage to the brain vasculature (*33*) (Fig.

4C–G). This results in decreased glucose import into the brain and dopaminergic and PV

neuron dysfunction which in turn cause psychosis-associated behavioral defects (Fig. 1B–H,

Fig. 2 and Fig. 4H).

Strikingly, administration of the NSAID aspirin, which inhibits inflammation and

oxidative stress, protected against the emergence of angiopathy as evidenced by reduced

fibrin accumulation, partially restored parenchymal glucose concentrations, and prevention

of several psychiatric-related phenotypes (Fig. 5 and fig. S7D–F). The increase in

coagulation factor V and the accumulation of fibrin in endothelial cells in G x E mice

indicate the presence of some vascular damage (*43*) (Fig.4A–G). Aspirin may prevent this

damage through its antiplatelet coagulation and anti-inflammatory properties. Since

oxidative stress and chronic inflammation are also common features of psychiatric disorders,

aspirin may provide a protective effect against the etiology of these diseases as well (*8, 11,*

44, 45). We have tried to quantify ROSs in our experimental mice, but thus far have not

been able to see differences among the four groups (fig. S8), so a more sensitive detection

method such as FRET probe for ROS detection may be required.

[ここに入力]

Sucrose consists of glucose and fructose, when these sugars are ingested at the same time,

we find that they generate AGEs in different cell types. Glucose-metabolism-derived AGEs,

which can be detected using α-AGE polyclonal antibody, accumulated primarily

in vascular endothelial cells (Fig. 3A–H), while methylglyoxsal-derived AGEs, recognized

by the α-AGE-4 monoclonal antibody accumulated specifically in microglia (fig. S5A–E).

Our results suggest that these two cell types may import and metabolize these two sugars

differently, but more detailed validation, such as one-cell metabolomic analysis, is required

to validate this idea.

Robust PV neuron function is required for PPI, working memory,

amphetamine-induced hyper-locomotion, dopamine (or 3,4-dihydroxyphenylacetic acid)

release regulation, and gamma oscillation generation, all of which are considered core

symptoms of psychiatric disorders (26). The PV neuron dysfunction in $G \times E$ mice may be

caused by the unique properties of these cells. PV-expressing interneurons are the

fast-spiking neuron subtype exhibiting a lower input resistance and higher-amplitude rapid

after-hyperpolarization than several projection neurons, and the combination of these

properties generates higher frequency action potentials compared with those by other

neuron types. This frequency of high action potentials in PV neurons can be exploited by

their fast spike phenotype to produce fast oscillations (30-100 Hz) in the gamma band of

[ここに入力]

neural oscillations (*46*). To maintain this rapid spiking property, PV neurons require high

energy expenditure as evidenced by mitochondrial and cytochrome c oxidase enrichment

(*47*). Therefore, reduced glucose within the brain parenchyma may preferentially disturb PV

neuron function, resulting in reduced PV expression. Indeed, reduced PV neuron number

has been repeatedly reported in postmortem brains of patients with psychiatric disorders

such as SZ and BP, and this reduction is proposed to be caused by a decrease in PV mRNA

or protein expression in these cells, rather than by a loss of PV neurons themselves (*22-25*).

Since PV is a Ca^{2+} binding protein, proper regulation of the expression of this protein is

considered to be important for maintaining the plasticity of PV neurons. For example, in PV

neurons of the hippocampus and mPFC, the expression of PV protein changes after

contextual fear conditioning, exposure to rich environments, and pharmacogenetic

activation of PV neurons, suggesting PV expression and plasticity are correlated (*25, 48*).

Further, it is known that the inhibitory activity of PV neurons is critical under

environmental stress to prevent a sequela of excessive excitatory activity such as oxidative

stress and inflammation (*47*). Pathogenic processes might more readily occur during the

critical adolescent prodromal period, wherein PV neurons may have increased vulnerability

to environmental stresses because of delayed maturation, which is another feature of PV

neurons (*49*).

[ここに入力]

In the present study, we identified capillary angiopathy in both G × E mice and postmortem brains of patients with SZ and BD (Fig. 4C–G and Fig. 6A–C). In our G × E mice, angiopathy was caused by the high AGE production capacity of sucrose combined with a GLO1 deficiency. Since high sugar consumption has been associated with SZ and BD, and decreased GLO1 expression and function have been seen in SZ and BD, similar interactions may be associated with some cases of psychiatric diseases. However, various other environmental stresses and genetic conditions may also converge to induce angiopathy. Several studies have reported that stressors, such as social defeat, isolation, and viral infection, induce vascular defects (*50-52*). These stressors are also risk factors for SZ and BD and induce PV neuron hypofunction (*53*), suggesting that angiopathy may be a common trigger for psychiatric phenotypes.

Acknowledgments: We thank Sayaka Ogikubo, Yoshie Matsumoto, Haimei Zhang, Minami Murata, Izumi Nohara, Yukiko Shimada, Emiko Hama, Nanako Obata, Mai Hatakenaka, and Chikako Ishida for their contribution to experiments related to this research. We also thank Dr. Tohru Kodama for explaining the procedures for microdialysis and encephalogram recording. We are grateful to Dr. Kenji Tanaka for reviewing the study. We acknowledge Chiaki Watanabe and Hiromi Onuma for coordinating the donations of

[ここに入力]

postmortem brains. Further, we thank Prof. Hideki Chiba for the preparation of postmortem

brain samples. We express our gratitude to the families of the deceased individuals for the

donations of brain tissue and their time and effort devoted to the consent process and

interviews.

Funding: This work was supported by the Japan Society for the Promotion of Science

(KAKENHI grants 18K14832 to S.H., 17K18395 and 19K08033 to H.M., 17K16408 to

T.T., and 18H02537 and 18K19383 to H.O.); the Ichiro Kanehara Foundation, Japan Prize

Foundation, and Takeda Science Foundation (to S.H.); the Naito Foundation (to H.M.); the

Strategic Research Program for Brain Sciences from AMED (grant JP19dm0107107 to

H.Y.); and a Grant-in-Aid for Scientific Research on Innovative Areas from the MEXT

(JP16H06277 to H.Y.). This research was also supported by KAKENHI Grant Numbers

16H05380 (to M.A.), 18K06977 (to K.T.), 19H03589 (to M.I.), and 18K15354 (to K.S.) as

well as AMED Grant Number JP19dm0107088 (to M.I.). This study was also supported by

The Kanae Foundation for the Promotion of Medical Science (to K.T.) and The Uehara

Memorial Foundation (to M.A.), and the Collaborative Research Program of Institute for

Protein Research, Osaka University, ICR-21-3.

[ここに入力]

Author contributions: S.H. and H.O. designed the study. S.H. performed and analyzed all

experiments. H.M. assisted with the experimental design of EEG recording and analysis of

the results and edited the manuscript. T.T. coordinated the EEG recording. Y.K., M.H., R.I.,

A.N., and H.Y. helped with the design of experiments using human specimens and provided

fixed human brain sections. T.S. performed data and image analysis. T.N. and S.S. provided

GFAP-GFP mice. T.D. and T.M. generated and provided *Glo1* knockout (KO) mice. K.T.

and K.S. backcrossed *Glo1* KO mice to B6J mice for all experiments. Y.N. performed

cDNA microarray analysis of gene expression. H.S. assisted with behavioral experiments.

M.I., M.A., K.T., K.S., and M.M. provided important suggestions to this study. T. T. and T.

H provided critical input on *Disc1-LI* mouse experiments. J.H. edited the manuscript. S.H.

generated all figures, tables, and wrote the manuscript. H.O. edited the manuscript and

supervised this study.

Competing interests: The authors declare no competing interests.

Data and materials availability: All materials used in this paper are available upon request.

The accession number for transcriptome analysis data is provided in the Materials and

Methods.

[ここに入力]

Figure Legends

Fig. 1 Generation of G × E model mice and psychosis-related phenotype analyses.

(**A**) Experimental Timeline. See Materials and Methods for a complete description. (**B–G**)

Behavioral analyses in the four groups of mice (n = 18–23 mice per group). (**B**)

Spontaneous locomotor activity in the open-field test. (**C**) Acoustic startle responses. (**D**)

Pre-pulse inhibition (PPI) using a 70 dB pre-pulse. (**E**) Object location test to evaluate

working memory. (**F**) Duration of self-grooming in the home cage. (**G**) Quantification of

nest-building skills over 8 h. Post hoc Dunnett's multiple comparisons test among groups at

each time point indicated ###$p < 0.001$ for Suc *glo1*/+ vs. Ctrl (Starch +/+), $$$p < 0.01$ for

Suc +/+ vs. Ctrl (Starch +/+). (**H, J**) Extracellular dopamine (DA) concentration in the

nucleus accumbens measured at 20 min intervals using an *in vivo* microdialysis system in

the presence or absence of aripiprazole (Aripi). Methamphetamine (1.25 mg/kg) was

administered via intraperitoneal (i.p.) injection at time 0 (arrow) (n = 6–11 mice per group).

See Materials and Methods for detailed statistical analyses. (**I**) Effects of Aripi treatment on

locomotor activity (n = 12–18 mice per group). (**B–H**) Starch +/+ group was used as a

control for post hoc Dunnett's test. All data are presented as mean ± SEM. ***$p < 0.001$,

**$p < 0.01$, *$p < 0.05$

[ここに入力]

Fig. 2 Parvalbumin-positive interneuron dysfunction in G × E mice.

(**A**) Parvalbumin (PV) immunohistochemistry in the medial prefrontal cortex (mPFC). (**B**)

Number of PV-positive cells in the mPFC (n = 4 mice per group). (**C**) Western blot analysis of

PV protein expression using tubulin as internal control. (**D**) Densitometric analysis of PV

protein expression: PV band intensities were divided by the corresponding tubulin band

intensities (n = 3 mice per group). (**E**) Average gamma band power in the home cage and

during novel object recognition in the open field (n = 7–8 mice per group). (**F**) Changes in

gamma band power from the home cage to the novel object phase in individual mice (n = 7–8

mice per group). Each *p* value indicates the result of repeated measures analysis of variance

(ANOVA). The effect of changes in gamma power at Starch +/+ ($F_{1, 5}$ = 8.29, $p = 0.035$), at

Starch *Glo1*/+ ($F_{1, 6}$ = 29.75, $p = 0.0016$), at Sucrose +/+ ($F_{1, 6}$ = 12.04, $p = 0.013$), at

Sucrose *Glo1*/+ ($F_{1, 7}$ = 4.038, $p = 0.084$). (B, D, H) Starch +/+ group was used as control for

post hoc Dunnett's test. All data are presented as mean ± SEM. ***p < 0.001, **p < 0.01, *p

< 0.05

Fig. 3 AGE accumulation in the neurovascular endothelium and pre-inflammatory

status of astrocytes in G × E mice.

[ここに入力]

(**A**) AGE immunohistochemistry in the medial prefrontal cortex. (**B**) Measurement of AGE

immunoreactive area. The mean intensity of the AGE-immunopositive area in the entire

image was measured for each section (n = 3 mice per group). (**C–E**) Colocalization of the

endothelial cell marker tomato lectin with the astrocytic marker ALDH1L1 or AGEs. (**F**)

Enlarged version of images in (**C**) presenting the region used for colocalization analysis.

(**G**) 2D intensity histogram of the two indicated channels for identifying the colocalization

of AGE with ALDH1L1 (Astrocyte) or tomato lectin (Endothelial cell). (**H**) Average

R-value of colocalization data including (**G**) three different locations of three mice. (**I**)

Immunohistological images of GFP-positive astrocytes and ALDH1L1 in the mPFC region

and merged GFP/ALDH1L1 image. (**J**) Percentage of GFP-positive cells per total

ALDH1L1-positive cells in each image in (**I**). (**K**) Mean fluorescent GFP intensities of 10

randomly selected cells per image in (**I**) (from four independent mice). (**B, J, K**) The Starch

+/+ group was used as control for post hoc Dunnett's test. All data are presented as mean ±

SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Fig. 4 Angiopathy and impaired glucose transport in G × E mice.

(**A, B**) Venn graph showing overlap in prefrontal cortex genes exhibiting a >2-fold (**A**) or

<0.5-fold (**B**) expression change compared with the control (CTL) group (n = 3 mice per

[ここに入力]

group). (**C**) Immunohistochemical images of fibrin and the endothelial cell marker tomato

lectin. (**D**) Measurement of the area of fibrin immunoexpression in (**C**). The mean intensity

of the fibrin-immunopositive area of the entire image was measured for each section (n = 4

mice per group). (**E**) Magnified immunohistochemical images of fibrin and the endothelial

cell marker tomato lectin. (**F**) 2D intensity histogram of the two indicated channels using

the left and middle panels in (**E**) for identifying fibrin colocalization with tomato lectin. (**G**)

Average R-value of colocalization data in (**F**) from three different locations of three mice.

(**H**) Extracellular glucose concentrations in the dialysis buffer from mPFC samples at each

time point (n = 5–6 mice per group). See the Materials and Methods for detailed statistical

analyses. (**I**) Plasma glucose concentrations (n = 6–7 mice per group). The first

measurement was performed after 16 h of fasting, and the second blood collection 30 min

after eating 0.05 g of carbohydrates. (**J**) Fasting plasma insulin levels (n = 5–6 mice per

group). (**D, H–J**) Starch +/+ group was used as a control for post hoc Dunnett's test. All

data are presented as mean ± SEM. *$p < 0.05$

Fig. 5 Protective effects of low-dose aspirin in G × E mice.

(**A–D**) Results of behavioral tests performed to evaluate the effects of aspirin treatment (n =

12–21 mice per group). (**A**) Quantification of spontaneous locomotor activity in the open

[ここに入力]

field. (**B**) Pre-pulse inhibition at 70 dB. (**C**) Object location test of working memory. (**D**)

Quantification of nest-building skills over 8 h. (**E**) Immunohistochemical images of fibrin

and the endothelial cell marker tomato lectin. (**F**) Measurement of the area of fibrin

immunoexpression in (**E**). The mean intensity of the fibrin-immunopositive area of the

entire image was measured for each section (n = 3 mice per group). (**G**) Extracellular

glucose concentrations in the dialysis buffer at each time point (n = 4–6 mice per group).

Post hoc Tukey's multiple comparisons test of groups at each time point, $^{\#\#\#}p < 0.001$, $^{\#\#}p <$

0.01, $^{\#}p < 0.05$ for Starch +/+ vs. Suc *Glo1*/+, $^{\$\$\$}p < 0.001$, $^{\$}p < 0.05$ for Starch +/+ Aspi vs.

Suc *Glo1*/+, $^{\&}p < 0.05$ for Starch +/+ vs. Suc *Glo1*/+ Asp. All data are presented as mean ±

SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Fig. 6 Angiopathy in postmortem brains from individuals with psychiatric disorders.

(**A**) Representative immunohistochemical fibrin images in the BA9 region of postmortem

brains from controls and patients with schizophrenia (SZ) or bipolar disorder. (**B**)

Measurement of the area of fibrin immunoexpression in (**A**). The mean intensity of the

fibrin-immunopositive area of the entire image was measured for each section. (**C**)

Representative immunohistochemical images of fibrin (magenta) and the endothelial cell

marker tomato lectin (green) in postmortem brains from a patient with SZ. Fibrin-positive

[ここに入力]

areas are merged with areas of vascular endothelial cell marker expression. (**D**) Diagrams

describing the hypothesis proposed to explain functional and behavioral abnormalities in

control (CTL) mice (left) and G × E mice (right) (see the Discussion for details). All data

are presented as mean ± SEM. **$p < 0.01$

Materials and Methods

Experimental design (related to Fig.1A)

After weaning (P21), wild-type (WT) and *Glo1* heterozygous mutant mice were fed either a

starch diet (control) or a sucrose diet (experimental). Diets were equal in total calories and

proportions of calories from carbohydrates, lipids, and proteins. The behavioral test battery

was administered starting at 2.5 months of age (upper panel). Middle panel: Macronutrient

composition of the two diets. We used *Glo1* heterozygous mice (or *Disc1* heterozygous

mice) to mimic patients with psychiatric disorders who exhibit decreased GLO1 activity or

expression (or decreased *Disc1* expression), whereas high-sucrose intake was used as an

environmental risk factor (bottom panel). We investigated four groups of mice: WT,

starch-fed mice (Starch +/+); *Glo1* (or *Disc1*) heterozygous, starch-fed mice (Starch *Glo1*/+

(or Starch *Disc1*/+)); WT, sucrose-fed mice (Suc +/+); and *Glo1* (or *Disc1*) heterozygous,

sucrose-fed mice (Suc *Glo1*/+ (or Suc *Disc1*/+)).

[ここに入力]

Animals

All experimental procedures were approved by the Animal Experimentation Ethics

Committee of the Tokyo Metropolitan Institute of Medical Science (49040). All mice were

maintained under a 12:12 h light:dark cycle (lights on at 8:00 AM) with free access to the

indicated diet. All efforts were made to minimize the number of animals used and their

suffering. *Glo1* knockout mice were generated as described previously (*54, 55*). Briefly,

Glo1-trapped ES cell lines from the International Gene Trap Consortium were used for the

generation of 3 founder mice, which were then backcrossed to C57BL/6 mice. In the

manuscript, trapped *Glo1* is referred to as *Glo1*. *Disc1-LI* (locus-impairment) heterozygous

mutant mice were generated as previously described (*16*). In the manuscript, *Disc1-LI* is

referred to as *Disc1*. Alternatively, mice were backcrossed to GFAP-GFP mice to monitor

astrocyte activation (*31, 32*). Male mice were exclusively used in the behavioral tests,

whereas mice of both sexes were used in histological, biochemical, and physiological

experiments.

Diet preparation

[ここに入力]

The two diets used in the present study were newly created in collaboration with Oriental

Yeast Co. Ltd. (Tokyo, Japan). We named the sucrose diet HSD-70 (# OYC 2405100) and

the starch diet HCD-70 (# OYC 2405000) (Supplementary Table 4). They contained the

same caloric proportions of carbohydrate, fat, and protein; however, all carbohydrate

calories are derived from either starch or sucrose.

Drug preparation

Aripiprazole was dissolved in acetic acid and diluted to 3.5 mg/L in water for

administration at 0.5 mg/kg/day. The final acetic acid concentration in the drinking water

was 0.7%. Aspirin was dissolved in ethanol and diluted to 70 mg/L in water for

administration at 1 mg/kg/day. The final ethanol concentration in the drinking water was

0.15%. The daily dose was based on the measurement of mean water consumption daily in

all sucrose-fed *Glo1* heterozygous mice and starch-fed wild-type (WT) mice (fig. S1D, E).

There were no between-group differences in terms of aripiprazole consumption (Student's

t-test, Starch +/+ and Suc *Glo1*/+: $p = 0.86$) or aspirin-containing water (Student's *t*-test,

Starch +/+ and Suc *Glo1*/+: $p = 0.61$).

Behavioral tests

[ここに入力]

Mice were habituated in the behavioral room for >30 min before each test. Behavioral tests

were performed in the following sequence of increasing stress: elevated plus maze,

grooming, nest building, open field, object location, social interaction, and pre-pulse

inhibition (PPI). All test apparatuses were cleaned using 70% ethanol and water between

trials, and the subsequent test session was started only after the ethanol vapor odors had

disappeared and the apparatuses had dried.

The elevated plus maze (EPM-04M, Muromachi, Japan) consisted of two opposing

open arms (297×54 mm) and two closed arms ($300 \times 60 \times 150$ mm) extending from a

central platform (60×60 mm). The entire apparatus was elevated 400 mm above the floor.

Each mouse was placed on the central platform facing a closed arm and allowed to freely

explore the maze for 10 min. Arm entry was defined as the entry of all four paws into the

arm. The time spent in the open arms over 10 min was recorded as an index of state anxiety.

For the self-grooming test, all mice housed in the same home cage were moved into a

new cage for 10 min. Each mouse was then individually placed in a standard mouse home

cage ($31 \times 16.5 \times 14$ cm) illuminated at ~200 lux. After a 10-min habituation period, each

mouse was scored for cumulative time spent grooming all body regions (*56*) over 10 min

using a stopwatch. Self-grooming behavior is conserved across species and is indicative of

[ここに入力]

certain pathological conditions or factors. In humans, for example, self-grooming increases

during stressful conditions and in certain psychiatric disorders (*56*).

For the nest building test, 200 g of corncob was spread across the bottom of each cage

for bedding, and a square-shaped piece of cotton was placed in the cage center as raw

material for the nest. Each mouse was individually placed in the cage for 8 h. Photos of the

constructed nest were acquired every 2 h, and the nest building process was evaluated by

measuring the proportion of loose cotton as follows: 1 point for 25% weight (Wt%)

loosened, 2 points for 50 Wt% loosened, 3 points for 75 Wt% loosened, and 4 points for

100 Wt% loosened. After 8 h, we checked the shape of the nest and added 1 point if the

mice had completed a nest with a bird's nest-like shape. The temperature of the room was

maintained at 25°C and illumination at 150–180 lux during nest building. Nest-building

behavior is an indicator of general well-being in mice (*57*), whereas poor nest building is an

indicator of psychological or physiological abnormalities (*58, 59*).

For the open field (OF) test, each mouse was placed in the center of the apparatus (40 ×

40 × 40 cm; 150–180 lux illumination) and allowed to move freely for 10 min. The

behavior of each mouse was monitored using a charge-coupled device camera mounted on

the ceiling above the OF. The total distance traveled (cm) was measured using CompACT

VAS software (Muromachi).

[ここに入力]

For the object location test (OLT) of working memory (*60*), mice first explored the

empty OF box, and then, two identical objects A and B (two 500-mL PET bottles filled with

blue-colored water) were placed in two corners 5 cm from the wall. After a 10-min

exploration/learning period, the mice were returned to their home cage for 5 min, and

Object A was moved to a new corner (Object A′). The animals were then placed back in the

OF box and allowed to explore for 5 min. The time spent exploring A′ and B were

measured to calculate a discrimination index representing working memory according to the

following equation: Discrimination Index = (Novel Object A′ exploration time − Familiar

Object B exploration time) / (Novel Object A′ + Familiar Object B exploration times). The

OLT was performed under illumination at 10–15 lux.

The social interaction test was conducted as previously described (*61*) using a

specialized Sociability Apparatus (SC-03M, Muromachi). The time spent sniffing a novel

stimulus mouse or object was manually scored from videos recorded using an overhead

color USB camera (aa000080a02, Iroiro House). Stimulus mice (129Sv/SvJcl strain) age-

and sex-matched to test mice were habituated to the apparatus and to the enclosure cup for

30 min per day for 2 days prior to testing. The location (left or right) of the novel object and

novel mouse within an enclosure were alternated across test subjects. The test mouse was

allowed to acclimate to the apparatus for a total of 20 min before the sociability test—the

[ここに入力]

first 10 min in the central chamber with the doors closed and then 10 min in the empty arena

with the doors open. The test subject was briefly confined to the center chamber while a

novel stimulus mouse in an enclosure cup was placed on one of the side chamber and

another empty enclosure cup (novel object) was placed on the other side of the chamber.

The test subject was allowed to approach the novel object or mouse freely for 10 min. The

time spent interacting with the stimulus mouse versus the novel object was calculated as an

index of sociability.

The SR-LAB-Startle Response System (San Diego Instruments) was used to detect

acoustic startle reflexes and PPI. Startle responses were measured using 5 stimulus

intensities (80, 90, 100, 110, and 120 dB) delivered 10 times each for 40 ms over a white

noise background (65 dB). The stimuli were presented in quasi-random order at random

inter-trial intervals (10–20 s). In the PPI session, mice were exposed to 2 stimulus patterns:

1) a startle stimulus alone (120 dB, 40 ms) with no pre-pulse stimulus and 2) a startle

stimulus (120 dB, 40 ms) following a pre-pulse stimulus (70 dB for 20 ms; lead time, 100

ms). Each trial was repeated 10 times in quasi-random order at random inter-trial intervals

(10–20 s). PPI was defined as the percent decline in startle response because of pre-pulse

stimuli according to the following equation: 100 − [(120 dB startle amplitude after any

[ここに入力]

pre-pulse) / (120 dB startle amplitude only)] × 100. A new accelerometer was used for the

experiments with *Disc1* knockout mice.

Immunohistochemistry

Following transcardial perfusion with PBS and 4% paraformaldehyde, entire brains were

collected, post-fixed at 4°C overnight, and cryoprotected in 20% sucrose at 4°C overnight.

Serial coronal sections (50 μm) were then cut using a cryostat (CM3050 S; Leica

Microsystems). The antigens in the tissues were reactivated by heating in HistoVT One

solution (Nacalai Tesque) for 30 min at 70°C using a water bath. Sections were

permeabilized with 0.2% Triton X-100 and 1% Block Ace (DS Pharma Biomedical) in PBS

for 30 min at room temperature, following which they were incubated overnight with the

indicated primary antibodies at room temperature. For immunohistochemistry of

postmortem human brain tissues, paraffin blocks including BA9 (a region of frontal cortex)

were sliced into 10-μm sections, deparaffinized with xylene, and rehydrated with decreasing

concentrations of ethanol in water. Antigens were reactivated by heating in HistoVT One

solution for 30 min at 90°C using a water bath. Sections were treated with TrueBlack

Lipofuscin Autofluorescence Quencher (Biotium Inc.) for 30 s at room temperature and

blocked with 1% Block Ace (DS Pharma Biomedical) in PBS for 30 min at room

[ここに入力]

temperature. Thereafter, the mouse and human brain sections were subjected to the same

immunostaining procedures. The following primary antibodies were diluted in PBS

containing 0.4% Block Ace: goat anti-PV (Frontier Institute, PV-Go-Af860; 1:2000), mouse

anti-ALDH1L1 (Rockland, 600-101-HB6S; 1:200), FITC-conjugated tomato lectin

(VECTOR, FL-1171; 1:200), chick anti-GFP (Abcam, ab13970; 1:500), goat anti-IBA1

(Abcam, ab48004; 1:100), mouse anti-NeuN (Millipore, MAB377; 1:500), rabbit anti-AGE

(Abcam, ab23722; 1:2000), rabbit anti-IBA1 (Wako, WDJ3047; 1:300), rat anti-CD68

(Abcam, ab53444; 1:500), and rabbit anti-fibrin (Dako, A0080, 1:500). Thereafter, sections

were washed three times with PBS containing 0.05% Tween-20, incubated for 2 h with

fluorochrome-conjugated secondary antibodies in PBS containing 0.4% Block Ace, and

washed an additional three times in PBS containing 0.4% Block Ace. For enhanced

horseradish peroxidase (HRP) immunostaining, samples were treated with 3% H_2O_2 in PBS

for 20 min following the reactivation step to quench endogenous peroxidase activity and

washed in PBS. Sections were incubated with rabbit anti-GLO1 (Novusbio, NBP2-75514,

1:1500) and/or mouse anti-AGE4 (Trans Genic Inc, 14B5, 1:400), followed by incubation

with anti-IgG antibodies conjugated to biotin (Vector, 1:200). After washing as described

for other secondary antibodies, sections were incubated with streptavidin-conjugated HRP

(Jackson ImmunoResearch, 1:200) for 120 min and washed three times with PBS

[ここに入力]

containing 0.05% Tween-20. The TSA Plus Fluorescence System (PerkinElmer) was used

to detect HRP activity. All preparations were counterstained with DAPI (Nacalai Tesque) to

reveal cell nuclei, washed three additional times, mounted in Permaflow (Thermo

Scientific), and observed using a FluoView® FV3000 Confocal Laser Scanning Microscope

(Olympus).

Image analysis

Unless otherwise noted, all image analyses were performed using ImageJ version

2.0.0-rc-59/1.51n, and images were first binarized. PV-positive cells with a threshold

exceeding 30 were counted.

To measure the areas immunopositive for AGE, fibrin, and AGE-4, the threshold

settings were applied to ensure that only the correct immunopositive area was properly

selected, and the stained area of the entire screen was measured. For measuring the

fluorescence intensity of GFP and IBA1, after setting the ROI to select only target

immunoreactive cells, the intensity of >5 cells in each image were measured. To count

microglial cells containing a CD68-positive area exceeding 20 μm^2, the threshold was set to

select only the CD68-positive area. Colocalization analysis of two different fluorescence

markers was performed using the Fiji plug-in Coloc 2 with the default settings. We adopted

[ここに入力]

Pearson's R value below the threshold to judge the strength of colocalization: R ≥ 0.7,

strong; 0.7 > R ≥ 0.4, moderate; 0.4 > R ≥ 0.2, weak; and R < 0.2, none or very weak.

Immunoblotting

Extracts from mouse hippocampi were homogenized in lysis buffer containing 40 mM Tris

base, 0.4% sodium dodecyl sulfate (SDS), 0.01 M EDTA (pH 8.0), 8 M urea, and 1 mM

phenylmethylsulfonyl fluoride. The total lysate protein content was quantified using a DC

Protein Assay Kit (Bio-Rad). Total protein (30 µg per gel lane) was separated using

SDS-PAGE and transferred to PVDF membranes (Millipore). Membranes were blocked

with TBST buffer (137m M NaCl, 2.7 mM KCl, and 25mM Tris, pH 8.0) including 0.2%

Triton X-100 and 5% bovine serum albumin (BSA) for 30 min at room temperature with

slow shaking, followed by incubation overnight with primary antibodies in TBST including

2% BSA at 4°C. The primary antibodies used were rabbit anti-GLO1 (Santa Cruz,

sc-67351; 1:1000), mouse-anti-PV (Swant, PV-235; 1:1000), rabbit anti-glucose transporter

1 (Glut1, Frontier Institute, Af1020; 1:1000) and mouse anti-tubulin (Santa Cruz, sc-32293;

1:10000). After washing three times with TBST, membranes were incubated with the

secondary antibody (HRP-conjugated anti-mouse or anti-rabbit IgG antibody, GE

Healthcare; 1:2000) in TBST plus 2% BSA. After washing three times with TBST, blots

[ここに入力]

were processed for chemiluminescence using standard protocols (ECL Prime Western

Blotting Detection Regent #RPN2236, GE Healthcare), and signals were detected using a

LAS 4000 Imager (Fuji Film).

Microdialysis

We used an *in vivo* microdialysis system for the measurement of extracellular dopamine

concentration (*62*) and collection of brain parenchymal dialysate (Fig.1H, J and fig. S7D).

After anesthesia by intraperitoneal injection of ketamine (80 mg/kg)/xylazine (16 mg/kg),

mice were fixed in a stereotaxic apparatus (Narishige) and a microdialysis guide cannula

(CXG-8, Eicom) was implanted in the medial prefrontal cortex (mPFC; antero-posterior

(AP), +1.8 mm; medio-lateral (ML), ±0.15 mm; dorso-ventral (DV), −1.5 mm from

bregma), or nucleus accumbens (NAc) (AP, +1.5 mm; ML, ±0.6 mm; DV, −3.5 mm from

bregma). After recovery for at least 10 days, a microdialysis probe (CX-I-8-01 for the

mPFC and CX-I-8-02 for NAc; Eicom) was inserted via the guide cannula. Following

insertion, the probe was connected to a syringe pump and perfusion was performed at 2 μ

L/min for NAc and 0.5 μL/min for mPFC using Ringer's solution (147 mM NaCl, 4 mM

KCl, and 2.3 mM $CaCl_2$). Dialysate samples were collected every 10 min and automatically

loaded onto an HTEC-500EPS HPLC unit (Eicom). Constant 5-HT concentration in three

[ここに入力]

consecutive collection periods was first confirmed to rule out blood contamination before

initiating the measurements of dopamine concentration or collection of parenchymal

dialysates. Analytes were then separated on an affinity column (PP-ODS III, Eicom), and

compounds were subjected to redox reactions within an electrochemical detection unit

(amperometric DC mode; applied potential range, 450 mV). The resulting chromatograms

were analyzed using an EPC-500 data processor (Eicom), and actual sample concentrations

were computed based on the peak heights obtained using 0.01, 0.1, and 1 pg dopamine in

standard solution (Sigma). The locations of the microdialysis probes were histologically

confirmed.

For glucose measurements in Fig. 4H and Fig. 5G, we collected dialysates from the

mPFC for 1 h after 16 h of fasting, followed by 0–1 h after eating 0.05 g of carbohydrates

(starch or sucrose), and 1–2 h after eating 0.05 g of carbohydrates (starch or sucrose).

EEG recordings

For behavioral and video/EEG monitoring, mice were anesthetized by an intraperitoneal

injection of ketamine (80 mg/kg)/xylazine (16 mg/kg), fixed in a stereotaxic apparatus

(Narishige, Japan), and implanted with EEG and electromyography (EMG) electrodes. The

EEG electrodes were gold-coated stainless steel screws (SUS303) soldered with lead wires

[ここに入力]

(ANE-0190, Adler's Nest, Japan) implanted epidurally over the left frontal cortex (AP, 1

mm; ML, 1 mm) and the bilateral parietal cortex (AP, −2 mm; ML, ±2 mm). All wires

were soldered to a multichannel electrical connector (R833-83-006-001, TOKIWA SHOKO,

Japan). The left parietal cortex electrode was used as a reference for monitoring the frontal

cortex EEG. The EMG electrodes were lead wires placed bilaterally into the trapezius

muscle. Following recovery for at least 10 days, EEG/EMG signals were amplified and

band-pass filtered (EEG: 1.5–1000 Hz; EGM: 15–3000 Hz) using a MEG-6116 system

(NIHON KOHDEN), digitized at a sampling rate of 200 Hz, recorded using a data

acquisition system (PowerLab 8/30, ADInstruments), and analyzed using LabChart

Software (ADInstruments). Behavioral activities were recorded using a USB camera

(aa000080a02, Iroiro House, Japan). Behavioral and electrophysiological responses to a

novel object (an empty 100 mL DURAN bin) were recorded in an OF chamber (20 × 20 ×

26 cm). The novel object was placed in one corner of the OF chamber to induce exploration.

The 30 s preceding the first contact with the novel object was analyzed for object

recognition ("object activity"). For EEG monitoring in the home cage, mice were first

habituated for 8 h. Home cage EEG data were then acquired for 2 min after awaking as

confirmed by clear EMG signals and movement images from an offline video camera

analysis ("home cage activity"). All recordings were converted into power spectra using a

[ここに入力]

fast Fourier transform (FFT) algorithm with a 5-s Hann cosine-bell window and 50%

overlap between successive window measurements. All FFTs were maintained at 1024

points to obtain 0.512 Hz resolution. The total signal amplitude or power (V^2) in each 5-s

period was measured as the power magnitude at each frequency. The mean of the grouped

power spectra was calculated over the following frequency ranges: 1–4 Hz (delta), 5–10 Hz

(theta), 30–45 Hz (low gamma), and 55–80 Hz (high gamma). The power values detected at

each frequency range for 30 s were further averaged over 30 s of total EEG power using the

mean values to remove potential noise. These analyses were performed using custom

software written in MATLAB (R2019b; MathWorks).

Transcriptome analysis

Three independent total RNA samples from each group were mixed and purified using a

RNeasy Mini Kit (Qiagen). RNA quality was assessed using a 2100 bioanalyzer (Agilent

Technologies). Cy3-labeled cRNA was prepared using a Low Input Quick Amp Labeling

Kit (Agilent Technologies), in accordance with the manufacturer's protocol. Samples were

hybridized to the SurePrint G3 Mouse Gene Expression v2 Microarray (G4852B; Agilent

Technologies). Thereafter, the array was washed and scanned using the SureScan

Microarray Scanner (Agilent Technologies). Microarray images were analyzed using the

[ここに入力]

Feature Extraction software with default settings for all parameters (Agilent Technologies).

Data from each microarray analysis were normalized by shift to the 75th percentile without

baseline transformation. Microarray results were deposited in the Gene Expression

Omnibus database under the accession number GSE141829.

Insulin and glucose measurements

Blood plasma was collected from the mouse cheek as described by Golde (*63*). Plasma

glucose concentration was measured using a Precision-Neo blood glucose meter (#71386-80,

Abbott Japan), plasma insulin concentration using an ELISA kit (#M1102, MORINAGA),

and glucose concentration in the dialysate samples using a different ELISA kit (#ab65333,

Abcam), all according to the manufacturers' instructions. Data were collected on a

microplate reader (Varioskan, Thermo Fisher Scientific).

Human postmortem brain tissue collection

Postmortem brain tissues from patients with SZ and BD were obtained from the Fukushima

Brain Bank at the Department of Neuropsychiatry, Fukushima Medical University.

Postmortem brain tissues from control individuals were obtained from the Section of

Pathology, Fukushima Medical University Hospital. The use of postmortem human brain

[ここに入力]

tissues in the present study was approved by the Ethics Committee of Fukushima Medical

University (No.1685) and Tokyo Metropolitan Institute of Medical Science (No. 18-20) and

complied with the Declaration of Helsinki and its later amendments. All procedures were

conducted with the informed written consent of the next of kin. Detailed demographic

information of the 10 patients with SZ, 9 patients with BD, and the 12 age- and sex-matched

control individuals is provided in Supplementary Table 3. No between-group differences

were observed in terms of sex (Fisher's exact test, Ctrl and SZ: $p = 1.00$, Ctrl and BD: $p =$

0.40), age (Student's t-test, Ctrl and SZ: $p = 0.69$, Welch's t-test, Ctrl and BD: $p = 0.66$),

postmortem interval (Student's t-test, Ctrl and SZ: $p = 0.89$, Ctrl and BD: $p = 0.98$), or a

history of diabetes mellitus (Fisher's exact test, Ctrl and SZ: $p = 0.59$, Ctrl and BD: $p =$

0.59). Each patient with SZ and BD fulfilled the diagnostic criteria established by the

American Psychiatric Association (Diagnostic and Statistical Manual of Mental Disorders,

DSM-IV) and did not have a past history of other neurological disorders or substance abuse.

Moreover, none of the control individuals had any record of mental disorders, neurological

disorders, or substance abuse.

ROS detection

[ここに入力]

DHE is oxidized by superoxide and other reactive species to produce a precipitate that emits red fluorescence, whose intensity can be used to measure the degree of cellular oxidative stress. Animals were quickly perfused with ice-cold phosphate-buffered salts (PBS), and the decapitated brains were embedded in OCT compound (Sakura Finetek Japan), and 10 μm thick coronary sections were prepared in a cryostat. These sections were incubated with phosphate buffer containing 100 μM diethylene triamine pentaacetic acid (PB/DTPA) for 10 min, and then incubated with PB/DTPA containing 10 μM DHE (Invitrogen) for 20 min. After washing with PB/DTPA twice, sections were fixed with 4% PFA for 15 min. Once washed with PBS, PBS containing 1 μM TO-PRO 3 (Thermo Scientific) was applied to each section followed by 3 washes with PBS. Then, sections were sealed with Permaflow (Thermo Scientific) and observed using a FluoView® FV3000 Confocal Laser Scanning Microscope (Olympus). For DHE detection, we used the following filter set: excitation 405 nm, emission 600–650 nm.

Statistical analyses

Statistical differences among ≥4 groups were determined using one-way analysis of variance (ANOVA), two-way ANOVA, three-way ANOVA, or repeated measures ANOVA, followed by the Bonferroni multiple comparison test or Tukey-Kramer test as a post hoc test, as

[ここに入力]

summarized in Supplementary Table 5. We used the same animals for all results coming from

Starch +/+ and Suc *Glo1*/+ groups according to the 3R rule. Data were collected at the same

time regardless of whether the drug was administered or not. The number of animals

employed for each experiment/analysis can be found in each figure legend. Detailed

descriptions of statistical analyses in Fig.1H, J, and 4H are shown below. In Figure 1H, post

hoc Dunnett's multiple comparisons test for groups was performed at specific time points (0

min and 40 min), $^{\#\#\#}p < 0.001$ for Suc *Glo1*/+ vs. Ctrl (Starch +/+), $^{\$}p < 0.05$ for Starch *Glo1*/+

vs. Ctrl (Starch +/+). In Fig. 1J, post hoc Bonferroni's multiple comparisons test for groups

was performed at specific time points (0 min and 40 min), $^{\#\#\#}p < 0.001$ for Suc *Glo1*/+ vs.

Starch +/+, $^{\$\$\$}p < 0.001$ for Suc *Glo1*/+ vs. Starch +/+ Aripi, $^{\&\&\&}p < 0.001$ for Suc *Glo1*/+ vs.

Suc *Glo1*/+ Aripi. In Fig. 4H, post hoc Dunnett's multiple comparisons test for groups was

performed at each time point, $^{\#\#\#}p < 0.001$, $^{\#\#}p < 0.01$ for Suc *Glo1*/+ vs. Ctrl (Starch +/+).

Supplementary Figure Legends

Supplementary Figure 1. Characterization of regional GLO1 expression in WT and

heterozygous *Glo1* mutant mice fed starch or sucrose.

(**A**) Western blot analysis of GLO1 protein expression using tubulin as internal control. The

cerebral cortex, including the hippocampus, was used as loading sample. (**B**) Densitometric

[ここに入力]

analysis of GLO1 protein expression (n = 3 or 4 mice per group). To quantify expression,

GLO1 band intensities in (**A**) were divided by the intensities of the corresponding tubulin

bands. (**C**) Body weight trajectories. No significant differences were observed among

groups (n = 6–10 mice per group). (**D, E**) Measurement of aripiprazole- or

aspirin-containing water consumption per day to estimate drug intake (n = 4 mice per

group). Student's *t*-test was used for statistical analysis. All data are presented as mean ±

SEM. **p < 0.01, *p < 0.05

Supplementary Figure 2. Analyses of psychiatric disease-related phenotypes in *Glo1*

heterozygotic mice.

(**A, B**) Behavioral analyses in the four groups of mice (n = 18–23 mice per group). (**A**)

Elevated plus maze test to evaluate anxiety. (**B**) Preference for novel mouse in the

three-chamber test, calculated as [(time spent exploring novel mouse)/(total time spent

exploring novel mouse and novel object)]□×□100%. (**C**) Extracellular dopamine

concentration in the medial prefrontal cortex measured at 20-min intervals using an *in vivo*

microdialysis system (n = 7–10 mice per group). (**D–I**) Effect of aripiprazole treatment on

behavioral phenotypes (n = 12–18 mice per group). (**D**) Effects of aripiprazole treatment on

the acoustic startle response, (**E**) pre-pulse inhibition (PPI) using a 70-dB pre-pulse, (**F**) object

[ここに入力]

location test, (**G**) self-grooming, (**H**) elevated plus maze performance, and (**I**) nest-building

skill. In (**I**), post hoc Tukey's multiple comparisons test of groups at specific time points, $^{###}p$

< 0.001, $^{#}p < 0.05$ for Starch +/+ vs. Suc *Glo1*/+ Aripi, $^{\$\$}p < 0.01$ for Starch +/+ Aripi vs. Suc

Glo1/+ Aripi, $^{\$\$\$}p < 0.001$ for Starch +/+ vs. Suc *Glo1*/+, $^{*}p < 0.05$ for Starch +/+ Aripi vs.

Suc *Glo1*/+. All data are presented as mean ± SEM. $^{***}p < 0.001$, $^{**}p < 0.01$, $^{*}p < 0.05$

Supplementary Figure 3. Analyses of psychiatric disease-related phenotypes in *Disc1*

heterozygotic mice.

(**A-E**) Behavioral analyses in the four groups of mice (n = 8–11 mice per group). (**A**)

Spontaneous locomotor activity in the open-field test. (**B**) Acoustic startle responses. (**C**)

Pre-pulse inhibition (PPI) using a 70 dB pre-pulse. (**D**) Object location test. (**E**) Preference for

novel mouse in the three-chamber test, calculated as fig. S2B. All data are presented as mean

± SEM. $^{***}p < 0.001$

Supplementary Figure 4. GLO1 cellular localization in the prefrontal cortex of

sucrose-fed wild-type mice.

(**A–F**) GLO1 localization in astrocytes. (**A**) GLO1 co-immunostaining with an astrocyte

marker (ALDH1L1) or an endothelial cell marker (tomato lectin). (**B**) Merged

[ここに入力]

GLO1/ALDH1L1 image from (**A**). Yellow arrows in (**B**) indicate cells with

GLO1/ALDH1L1 colocalization. (**C**) Merged GLO1/lectin image from (**A**). White arrows

denote representative GLO1-positive cells close to endothelial cells. (**D, E**) High

magnification images of GLO1 co-immunostaining with ALDH1L1 or with tomato lectin.

(**E**) Merged GLO1/ALDH1L1 image from (**D**). (**F**) 2D intensity histogram of the two

indicated channels to identify GLO1 colocalization with ALDH1L1 or tomato lectin,

interpreting the strength of a relationship based on R as follows: $R \geq 0.7$, strong; $0.7 > R \geq$

0.4, moderate; $0.4 > R \geq 0.2$, weak; and $R < 0.2$, none or very weak. (**G**) Average R-value

of colocalization data including (**F**) from three different locations of three mice. (**H**) GLO1

co-immunostaining with the neuronal marker NeuN and the microglial marker IBA1. (**I**)

Merged GLO1/NeuN image from (**H**). Yellow arrows indicate neurons with mild GLO1

immunoreactivity. (**J**) Merged GLO1/IBA1 image from (**H**). (**K**) GLO1 immunostaining

and DAPI staining in *Glo1* homozygous mice.

Supplementary Figure 5. Fructose-derived AGE4 accumulation in microglia of

sucrose-fed mice.

(**A, B**) Immunohistochemical images of the microglial marker IBA1 and AGE4 in the

medial prefrontal cortex region (mPFC). (**C**) Merged AGE4/IBA1 image. (**D**) High

[ここに入力]

magnification images of AGE4 co-immunostaining with IBA1. (**E**) Measurement of the

area of AGE-4 immunoexpression in (**B**). The mean AGE4-immunopositive area intensity

of the entire image was measured in each section (n = 4 mice per group). (**F**) Mean IBA1

fluorescence intensities of five randomly selected cells per image in four independent mice.

(**G**) Immunohistological images of IBA1- and CD68-positive microglia in the mPFC region.

White arrows in (**G**) indicate microglial cells containing CD68-positive areas >20 μm^2. (**H**)

Percentage of CD68-positive cells per total microglial cells in each image in (**G**). All data

are presented as mean ± SEM. ***$p < 0.001$, **$p < 0.01$

Supplementary Figure 6. Blood vessels Characterization in the four groups.

(**A**) Western blot analysis of glucose transporter 1 (Glut1) protein expression using tubulin

as internal control. The cerebral cortex, including the hippocampus, was used as loading

sample. (**B**) Densitometric analysis of Glut1 protein expression (n = 3 mice per group). To

quantify expression, Glut1 band intensities in (**A**) were divided by the intensities of the

corresponding tubulin bands. (**C**) Measurement of capillary diameter. Vessels <10 μm in

diameter were defined as capillaries, and the mean diameter of 10 randomly selected

capillaries per image measured in Fig. 4C (n = 3 mice per group). All data are presented as

mean ± SEM.

[ここに入力]

Supplementary Figure 7. Protective effects of aspirin against the development of

psychiatric phenotypes in G × E mice.

(**A–C**) Effects of aspirin treatment on the acoustic startle response (**A**), nest-building skill

(**B**), elevated plus maze test performance (**C**) and dopamine concentration (**D**). (**E**)

Parvalbumin (PV) Immunohistochemistry in the medial prefrontal cortex (mPFC). (**F**)

Number of PV-positive cells in the mPFC (n = 4 mice per group). (**G**) Immunohistological

images of GFP-positive astrocytes and ALDH1L1in the mPFC region. (**H**) Percentage of

GFP-positive cells per total ALDH1L1-positive cells in each image in (**G**). (**I**) Mean

fluorescence GFP intensities of 10 randomly selected cells per image in (**G**) (from four

independent mice). In (**B**), post hoc Tukey's multiple comparisons test of groups at specific

time points, $^{###}p < 0.001$, $^{##}p < 0.01$ for Starch +/+ vs. Suc *Glo1*/+ Aspi, $^{\$\$}p < 0.01$, $^{\$}p <$

0.05 for Starch +/+ Aspi vs. Suc *Glo1*/+ Aspi. In (**D**), post hoc Bonferroni's multiple

comparisons test of groups at specific time points (0 min and 40 min), $^{###}p < 0.001$ for Suc

Glo1/+ vs. Starch +/+, $^{\$\$\$}p < 0.001$ for Suc *Glo1*/+ vs. Starch +/+ Aspi, $^{\&\&\&}p < 0.001$ for

Suc *Glo1*/+ Aspi vs. Starch +/+, $^{\dagger\dagger}p < 0.01$ for Suc *Glo1*/+ Aspi vs. SucG*glo1*/+, $^{\ddagger\ddagger}p < 0.01$

for Suc *Glo1*/+ Aspi vs. Starch +/+ Aspi. All data are presented as mean ± SEM. $^{***}p <$

0.001, $^{**}p < 0.01$, $^{*}p < 0.05$

[ここに入力]

Supplementary Figure 8. ROS detection in the four groups.

(**A**) Reactive oxygen species (ROS) detection with *dihydroethidium* (*DHE*). (**B**) Merged

DHE/Nucleus (TO-PRO 3) image. All cells are DHE positive, according to nucleus images.

(**C**) ROS detection images without *DHE*. (**D**) Fluorescent intensity of DHE positive cells in

(**A**). The mean DHE fluorescent intensity of each cell was measured in each section (n = 4

mice per group). All data are presented as mean ± SEM.

References

1. *Sugar Intake for adults and children* (2015).
2. N. Fidler Mis, C. Braegger, J. Bronsky, C. Campoy, M. Domellöf, N. D. Embleton, I. Hojsak, J. Hulst, F. Indrio, A. Lapillonne, W. Mihatsch, C. Molgaard, R. Vora, M. Fewtrell, Sugar in Infants, Children and Adolescents: A Position Paper of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* **65**, 681-696 (2017).
3. M. Peet, International variations in the outcome of schizophrenia and the prevalence of depression in relation to national dietary practices: an ecological analysis. *Br J Psychiatry* **184**, 404-408 (2004).
4. J. C. Ratliff, L. B. Palmese, E. L. Reutenauer, E. Liskov, C. M. Grilo, C. Tek, The effect of dietary and physical activity pattern on metabolic profile in individuals with schizophrenia: a cross-sectional study. *Compr Psychiatry* **53**, 1028-1033 (2012).
5. J. L. Elmslie, J. I. Mann, J. T. Silverstone, S. M. Williams, S. E. Romans, Determinants of overweight and obesity in patients with bipolar disorder. *J Clin Psychiatry* **62**, 486-491; quiz 492-483 (2001).

6.   L. Lien, N. Lien, S. Heyerdahl, M. Thoresen, E. Bjertness, Consumption of soft drinks and hyperactivity, mental distress, and conduct problems among adolescents in Oslo, Norway. *Am J Public Health* **96**, 1815-1820 (2006).

7.   K. Byun, Y. Yoo, M. Son, J. Lee, G. B. Jeong, Y. M. Park, G. H. Salekdeh, B. Lee, Advanced glycation end-products produced systemically and by macrophages: A common contributor to inflammation and degenerative diseases. *Pharmacol Ther* **177**, 44-55 (2017).

8.   B. K. Bitanihirwe, T. U. Woo, Oxidative stress in schizophrenia: an integrated approach. *Neurosci Biobehav Rev* **35**, 878-893 (2011).

9.   L. G. Nucifora, T. Tanaka, L. N. Hayes, M. Kim, B. J. Lee, T. Matsuda, F. C. Nucifora, Jr., T. Sedlak, R. Mojtabai, W. Eaton, A. Sawa, Reduction of plasma glutathione in psychosis associated with schizophrenia and bipolar disorder in translational psychiatry. *Transl Psychiatry* **7**, e1215 (2017).

10.   D. W. Volk, A. Chitrapu, J. R. Edelson, K. M. Roman, A. E. Moroco, D. A. Lewis, Molecular mechanisms and timing of cortical immune activation in schizophrenia. *Am J Psychiatry* **172**, 1112-1121 (2015).

11.   H. Q. Cai, V. S. Catts, M. J. Webster, C. Galletly, D. Liu, M. O'Donnell, T. W. Weickert, C. S. Weickert, Increased macrophages and changed brain endothelial cell gene expression in the frontal cortex of people with schizophrenia displaying inflammation. *Mol Psychiatry* **25**, 761-775 (2020).

12.   P. J. Thornalley, Glutathione-dependent detoxification of alpha-oxoaldehydes by the glyoxalase system: involvement in disease mechanisms and antiproliferative activity of glyoxalase I inhibitors. *Chem Biol Interact* **111-112**, 137-151 (1998).

13.   M. Fujimoto, S. Uchida, T. Watanuki, Y. Wakabayashi, K. Otsuki, T. Matsubara, M. Suetsugi, H. Funato, Y. Watanabe, Reduced expression of glyoxalase-1 mRNA in mood disorder patients. *Neurosci Lett* **438**, 196-199 (2008).

14.   M. Miyashita, M. Arai, A. Kobori, T. Ichikawa, K. Toriumi, K. Niizato, K. Oshima, Y. Okazaki, T. Yoshikawa, N. Amano, T. Miyata, M. Itokawa, Clinical features of schizophrenia with enhanced carbonyl stress. *Schizophr Bull* **40**, 1040-1046 (2014).

[ここに入力]

15. M. Toyosima, M. Maekawa, T. Toyota, Y. Iwayama, M. Arai, T. Ichikawa, M. Miyashita, T. Arinami, M. Itokawa, T. Yoshikawa, Schizophrenia with the 22q11.2 deletion and additional genetic defects: case history. *Br J Psychiatry* **199**, 245-246 (2011).

16. N. Shahani, S. Seshadri, H. Jaaro-Peled, K. Ishizuka, Y. Hirota-Tsuyada, Q. Wang, M. Koga, T. W. Sedlak, C. Korth, N. J. Brandon, A. Kamiya, S. Subramaniam, T. Tomoda, A. Sawa, DISC1 regulates trafficking and processing of APP and Aß generation. *Mol Psychiatry* **20**, 874-879 (2015).

17. T. Tomoda, A. Sumitomo, H. Jaaro-Peled, A. Sawa, Utility and validity of DISC1 mouse models in biological psychiatry. *Neuroscience* **321**, 99-107 (2016).

18. O. D. Howes, A. J. Montgomery, M. C. Asselin, R. M. Murray, P. M. Grasby, P. K. McGuire, Molecular imaging studies of the striatal dopaminergic system in psychosis and predictions for the prodromal phase of psychosis. *Br J Psychiatry Suppl* **51**, s13-18 (2007).

19. M. Laruelle, A. Abi-Dargham, C. H. van Dyck, R. Gil, C. D. D'Souza, J. Erdos, E. McCance, W. Rosenblatt, C. Fingado, S. S. Zoghbi, R. M. Baldwin, J. P. Seibyl, J. H. Krystal, D. S. Charney, R. B. Innis, Single photon emission computerized tomography imaging of amphetamine-induced dopamine release in drug-free schizophrenic subjects. *Proc Natl Acad Sci U S A* **93**, 9235-9240 (1996).

20. G. Winterer, D. R. Weinberger, Genes, dopamine and cortical signal-to-noise ratio in schizophrenia. *Trends Neurosci* **27**, 683-690 (2004).

21. A. de Bartolomeis, C. Tomasetti, F. Iasevoli, Update on the Mechanism of Action of Aripiprazole: Translational Insights into Antipsychotic Strategies Beyond Dopamine Receptor Antagonism. *CNS Drugs* **29**, 773-799 (2015).

22. C. L. Beasley, G. P. Reynolds, Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophr Res* **24**, 349-355 (1997).

23. T. Hashimoto, D. W. Volk, S. M. Eggan, K. Mirnics, J. N. Pierri, Z. Sun, A. R. Sampson, D. A. Lewis, Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia. *J Neurosci* **23**, 6315-6326 (2003).

[ここに入力]

24. M. Thompson, C. S. Weickert, E. Wyatt, M. J. Webster, Decreased glutamic acid decarboxylase(67) mRNA expression in multiple brain areas of patients with schizophrenia and mood disorders. *J Psychiatr Res* 43, 970-977 (2009).

25. A. Mukherjee, F. Carvalho, S. Eliez, P. Caroni, Long-Lasting Rescue of Network and Cognitive Dysfunction in a Genetic Schizophrenia Model. *Cell* 178, 1387-1402.e1314 (2019).

26. R. Nguyen, Investigating the Roles of Parvalbumin and Cholecystokinin Interneurons of the Ventral Hippocampus and Medial Prefrontal Cortex in Schizophrenia-Related Behaviours. (2018).

27. M. I. Atagun, Brain oscillations in bipolar disorder and lithium-induced changes. *Neuropsychiatr Dis Treat* 12, 589-601 (2016).

28. P. J. Uhlhaas, W. Singer, Abnormal neural oscillations and synchrony in schizophrenia. *Nat Rev Neurosci* 11, 100-113 (2010).

29. A. R. Bialas, J. Presumey, A. Das, C. E. van der Poel, P. H. Lapchak, L. Mesin, G. Victora, G. C. Tsokos, C. Mawrin, R. Herbst, M. C. Carroll, Microglia-dependent synapse loss in type I interferon-mediated lupus. *Nature* 546, 539-543 (2017).

30. K. E. Linker, M. G. Elabd, P. Tawadrous, M. Cano, K. N. Green, M. A. Wood, F. M. Leslie, Microglial activation increases cocaine self-administration following adolescent nicotine exposure. *Nat Commun* 11, 306 (2020).

31. R. Suzuki, S. Arata, S. Nakajo, K. Ikenaka, S. Kikuyama, S. Shioda, Expression of the receptor for pituitary adenylate cyclase-activating polypeptide (PAC1-R) in reactive astrocytes. *Brain Res Mol Brain Res* 115, 10-20 (2003).

32. R. Suzuki, J. Watanabe, S. Arata, H. Funahashi, S. Kikuyama, S. Shioda, A transgenic mouse model for the detailed morphological study of astrocytes. *Neurosci Res* 47, 451-454 (2003).

33. M. P. Wautier, O. Chappey, S. Corda, D. M. Stern, A. M. Schmidt, J. L. Wautier, Activation of NADPH oxidase by AGE links oxidant stress to altered gene expression via RAGE. *Am J Physiol Endocrinol Metab* 280, E685-694 (2001).

34. A. Das, G. C. t. Wallace, C. Holmes, M. L. McDowell, J. A. Smith, J. D. Marshall, L. Bonilha, J. C. Edwards, S. S. Glazier, S. K. Ray, N. L. Banik, Hippocampal tissue of patients with refractory temporal lobe epilepsy is

associated with astrocyte activation, inflammation, and altered expression of channels and receptors. *Neuroscience* **220**, 237-246 (2012).

35.	D. D. Mao, W. Y. Yang, Y. Li, J. W. Lin, S. Y. Gao, Y. R. Wang, H. Y. Hu, Effect of Qingxin Kaiqiao Fang on Hippocampus mRNA Expression of the Inflammation-Related Genes IL-1beta, GFAP, and Abeta in an Alzheimer's Disease Rat Model. *Evid Based Complement Alternat Med* **2018**, 9267653 (2018).

36.	C. García-Cáceres, C. Quarta, L. Varela, Y. Gao, T. Gruber, B. Legutko, M. Jastroch, P. Johansson, J. Ninkovic, C. X. Yi, O. Le Thuc, K. Szigeti-Buck, W. Cai, C. W. Meyer, P. T. Pfluger, A. M. Fernandez, S. Luquet, S. C. Woods, I. Torres-Alemán, C. R. Kahn, M. Götz, T. L. Horvath, M. H. Tschöp, Astrocytic Insulin Signaling Couples Brain Glucose Uptake with Nutrient Availability. *Cell* **166**, 867-880 (2016).

37.	J. P. Luyendyk, J. G. Schoenecker, M. J. Flick, The multifaceted role of fibrinogen in tissue injury and inflammation. *Blood* **133**, 511-520 (2019).

38.	J. Paul, S. Strickland, J. P. Melchor, Fibrin deposition accelerates neurovascular damage and neuroinflammation in mouse models of Alzheimer's disease. *J Exp Med* **204**, 1999-2008 (2007).

39.	M. Nitta, T. Kishimoto, N. Muller, M. Weiser, M. Davidson, J. M. Kane, C. U. Correll, Adjunctive use of nonsteroidal anti-inflammatory drugs for schizophrenia: a meta-analytic investigation of randomized controlled trials. *Schizophr Bull* **39**, 1230-1241 (2013).

40.	J. Savitz, S. Preskorn, T. K. Teague, D. Drevets, W. Yates, W. Drevets, Minocycline and aspirin in the treatment of bipolar depression: a protocol for a proof-of-concept, randomised, double-blind, placebo-controlled, 2x2 clinical trial. *BMJ Open* **2**, e000643 (2012).

41.	G. de Gaetano, Low-dose aspirin and vitamin E in people at cardiovascular risk: a randomised trial in general practice. Collaborative Group of the Primary Prevention Project. *Lancet* **357**, 89-95 (2001).

42.	E. Magen, J. R. Viskoper, J. Mishal, R. Priluk, D. London, C. Yosefy, Effects of low-dose aspirin on blood pressure and endothelial function of treated hypertensive hypercholesterolaemic subjects. *J Hum Hypertens* **19**, 667-673 (2005).

43.	M. Cortes-Canteli, D. Zamolodchikov, H. J. Ahn, S. Strickland, E. H. Norris, Fibrinogen and altered hemostasis in Alzheimer's disease. *J Alzheimers Dis* **32**, 599-608 (2012).

[ここに入力]

44. A. Ciobica, M. Padurariu, I. Dobrin, C. Stefanescu, R. Dobrin, Oxidative stress in schizophrenia - focusing on the main markers. *Psychiatr Danub* **23**, 237-245 (2011).

45. E. Kim, Z. Keskey, M. Kang, C. Kitchen, W. E. Bentley, S. Chen, D. L. Kelly, G. F. Payne, Validation of oxidative stress assay for schizophrenia. *Schizophr Res* **212**, 126-133 (2019).

46. B. Rudy, C. J. McBain, Kv3 channels: voltage-gated K+ channels designed for high-frequency repetitive firing. *Trends Neurosci* **24**, 517-526 (2001).

47. O. Kann, The interneuron energy hypothesis: Implications for brain disease. *Neurobiol Dis* **90**, 75-85 (2016).

48. F. Donato, S. B. Rompani, P. Caroni, Parvalbumin-expressing basket-cell network plasticity induced by experience regulates adult learning. *Nature* **504**, 272-276 (2013).

49. T. R. Insel, Rethinking schizophrenia. *Nature* **468**, 187-193 (2010).

50. C. Menard, M. L. Pfau, G. E. Hodes, V. Kana, V. X. Wang, S. Bouchard, A. Takahashi, M. E. Flanigan, H. Aleyasin, K. B. LeClair, W. G. Janssen, B. Labonté, E. M. Parise, Z. S. Lorsch, S. A. Golden, M. Heshmati, C. Tamminga, G. Turecki, M. Campbell, Z. A. Fayad, C. Y. Tang, M. Merad, S. J. Russo, Social stress induces neurovascular pathology promoting depression. *Nat Neurosci* **20**, 1752-1760 (2017).

51. W. A. Banks, A. M. Gray, M. A. Erickson, T. S. Salameh, M. Damodarasamy, N. Sheibani, J. S. Meabon, E. E. Wing, Y. Morofuji, D. G. Cook, M. J. Reed, Lipopolysaccharide-induced blood-brain barrier disruption: roles of cyclooxygenase, oxidative stress, neuroinflammation, and elements of the neurovascular unit. *J Neuroinflammation* **12**, 223 (2015).

52. D. Ben-Nathan, S. Lustig, H. D. Danenberg, Stress-induced neuroinvasiveness of a neurovirulent noninvasive Sindbis virus in cold or isolation subjected mice. *Life Sci* **48**, 1493-1500 (1991).

53. D. Koshiyama, M. Fukunaga, N. Okada, K. Morita, K. Nemoto, K. Usui, H. Yamamori, Y. Yasuda, M. Fujimoto, N. Kudo, H. Azechi, Y. Watanabe, N. Hashimoto, H. Narita, I. Kusumi, K. Ohi, T. Shimada, Y. Kataoka, M. Yamamoto, N. Ozaki, G. Okada, Y. Okamoto, K. Harada, K. Matsuo, H. Yamasue, O. Abe, R. Hashimoto, T. Takahashi, T. Hori, M. Nakataki, T. Onitsuka, L. Holleran, N. Jahanshad, T. G. M. van Erp, J. Turner, G. Donohoe, P. M. Thompson, K. Kasai, R. Hashimoto, White matter

microstructural alterations across four major psychiatric disorders: mega-analysis study in 2937 individuals. *Mol Psychiatry*, (2019).

54. S. Koike, K. Toriumi, S. Kasahara, Y. Kibune, Y. I. Ishida, T. Dan, T. Miyata, M. Arai, Y. Ogasawara, Accumulation of Carbonyl Proteins in the Brain of Mouse Model for Methylglyoxal Detoxification Deficits. *Antioxidants (Basel)* 10, (2021).

55. K. Toriumi, S. Berto, S. Koike, N. Usui, T. Dan, K. Suzuki, M. Miyashita, Y. Horiuchi, A. Yoshikawa, M. Asakura, K. Nagahama, H.-C. Lin, Y. Sugaya, T. Watanabe, M. Kano, Y. Ogasawara, T. Miyata, M. Itokawa, G. Konopka, M. Arai, Combined glyoxalase 1 dysfunction and vitamin B6 deficiency in a schizophrenia model system causes mitochondrial dysfunction in the prefrontal cortex. *Redox Biology* 45, 102057 (2021).

56. A. V. Kalueff, A. M. Stewart, C. Song, K. C. Berridge, A. M. Graybiel, J. C. Fentress, Neurobiology of rodent self-grooming and its value for translational neuroscience. *Nat Rev Neurosci* 17, 45-59 (2016).

57. P. Jirkof, Burrowing and nest building behavior as indicators of well-being in mice. *J Neurosci Methods* 234, 139-146 (2014).

58. C. S. Pedersen, D. B. Sorensen, A. I. Parachikova, N. Plath, PCP-induced deficits in murine nest building activity: employment of an ethological rodent behavior to mimic negative-like symptoms of schizophrenia. *Behav Brain Res* 273, 63-72 (2014).

59. A. Forsingdal, K. Fejgin, V. Nielsen, T. Werge, J. Nielsen, 15q13.3 homozygous knockout mouse model display epilepsy-, autism- and schizophrenia-related phenotypes. *Transl Psychiatry* 6, e860 (2016).

60. A. Ennaceur, J. Delacour, A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data. *Behav Brain Res* 31, 47-59 (1988).

61. J. L. Silverman, S. S. Tolu, C. L. Barkan, J. N. Crawley, Repetitive self-grooming behavior in the BTBR mouse model of autism is blocked by the mGluR5 antagonist MPEP. *Neuropsychopharmacology* 35, 976-989 (2010).

62. L. A. Tellez, W. Han, X. Zhang, T. L. Ferreira, I. O. Perez, S. J. Shammah-Lagnado, A. N. van den Pol, I. E. de Araujo, Separate circuitries encode the hedonic and nutritional values of sugar. *Nat Neurosci* 19, 465-470 (2016).

[ここに入力]

63. W. T. Golde, P. Gollobin, L. L. Rodriguez, A rapid, simple, and humane method for submandibular bleeding of mice using a lancet. *Lab Anim (NY)* **34**, 39-43 (2005).

Figure 1

Fig. 1 Generation of G × E model mice and psychosis-related phenotype analyses.
(**A**) Experimental Timeline. See Materials and Methods for a complete description. (**B–G**) Behavioral analyses in the four groups of mice (n = 18–23 mice per group). (**B**) Spontaneous locomotor activity in the open-field test. (**C**) Acoustic startle responses. (**D**) Pre-pulse inhibition (PPI) using a 70 dB pre-pulse. (**E**) Object location test to evaluate working memory. (**F**) Duration of self-grooming in the home cage. (**G**) Quantification of nest-building skills over 8 h. Post hoc Dunnett's multiple comparisons test among groups at each time point indicated ###$p < 0.001$ for Suc *glo1*/+ vs. Ctrl (Starch +/+), \$\$$p < 0.01$ for Suc +/+ vs. Ctrl (Starch +/+). (**H, J**) Extracellular dopamine (DA) concentration in the nucleus accumbens measured at 20 min intervals using an *in vivo* microdialysis system in the presence or absence of aripiprazole (Aripi). Methamphetamine (1.25 mg/kg) was administered via intraperitoneal (i.p.) injection at time 0 (arrow) (n = 6–11 mice per group). See Materials and Methods for detailed statistical analyses. (**I**) Effects of Aripi treatment on locomotor activity (n = 12–18 mice per group). (**B–H**) Starch +/+ group was used as a control for post hoc Dunnett's test. All data are presented as mean ± SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Figure 2

Fig. 2 Parvalbumin-positive interneuron dysfunction in G × E mice.
(**A**) Parvalbumin (PV) immunohistochemistry in the medial prefrontal cortex (mPFC). (**B**) Number of PV-positive cells in the mPFC (n = 4 mice per group). (**C**) Western blot analysis of PV protein expression using tubulin as internal control. (**D**) Densitometric analysis of PV protein expression: PV band intensities were divided by the corresponding tubulin band intensities (n = 3 mice per group). (**E**) Average gamma band power in the home cage and during novel object recognition in the open field (n = 7–8 mice per group). (**F**) Changes in gamma band power from the home cage to the novel object phase in individual mice (n = 7–8 mice per group). Each *p* value indicates the result of repeated measures analysis of variance (ANOVA). The effect of changes in gamma power at Starch +/+ (F1, 5 = 8.29, *p* = 0.035), at Starch *glo1*/+ (F1, 6 = 29.75, *p* = 0.0016), at Sucrose +/+ (F1, 6 = 12.04, *p* = 0.013), at Sucrose *glo1*/+ (F1, 7 = 4.038, *p* = 0.084). (B, D, H) Starch +/+ group was used as control for post hoc Dunnett's test. All data are presented as mean ± SEM. ***p* < 0.001, ***p* < 0.01, **p* < 0.05

Figure 3

Fig. 3 AGE accumulation in the neurovascular endothelium and pre-inflammatory status of astrocytes in G × E mice.
(**A**) AGE immunohistochemistry in the medial prefrontal cortex. (**B**) Measurement of AGE immunoreactive area. The mean intensity of the AGE-immunopositive area in the entire image was measured for each section (n = 3 mice per group). (**C–E**) Colocalization of the endothelial cell marker tomato lectin with the astrocytic marker ALDH1L1 or AGEs. (**F**) Enlarged version of images in (**C**) presenting the region used for colocalization analysis. (**G**) 2D intensity histogram of the two indicated channels for identifying the colocalization of AGE with ALDH1L1 (Astrocyte) or tomato lectin (Endothelial cell). (**H**) Average R-value of colocalization data including (**G**) three different locations of three mice. (**I**) Immunohistological images of GFP-positive astrocytes and ALDH1L1in the mPFC region and merged GFP/ALDH1L1 image. (**J**) Percentage of GFP-positive cells per total ALDH1L1-positive cells in each image in (**I**). (**K**) Mean fluorescent GFP intensities of 10 randomly selected cells per image in (**I**) (from four independent mice). (**B, J, K**) The Starch +/+ group was used as control for post hoc Dunnett's test. All data are presented as mean ± SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Figure 4

Fig. 4 Angiopathy and impaired glucose transport in G × E mice.
(**A, B**) Venn graph showing overlap in prefrontal cortex genes exhibiting a >2-fold (**A**) or <0.5-fold (**B**) expression change compared with the control (CTL) group (n = 3 mice per group). (**C**) Immunohistochemical images of fibrin and the endothelial cell marker tomato lectin. (**D**) Measurement of the area of fibrin immunoexpression in (**C**). The mean intensity of the fibrin-immunopositive area of the entire image was measured for each section (n = 4 mice per group). (**E**) Magnified immunohistochemical images of fibrin and the endothelial cell marker tomato lectin. (**F**) 2D intensity histogram of the two indicated channels using the left and middle panels in (**E**) for identifying fibrin colocalization with tomato lectin. (**G**) Average R-value of colocalization data in (**F**) from three different locations of three mice. (**H**) Extracellular glucose concentrations in the dialysis buffer from mPFC samples at each time point (n = 5–6 mice per group). See the Materials and Methods for detailed statistical analyses. (**I**) Plasma glucose concentrations (n = 6–7 mice per group). The first measurement was performed after 16 h of fasting, and the second blood collection 30 min after eating 0.05 g of carbohydrates. (**J**) Fasting plasma insulin levels (n = 5–6 mice per group). (**D, H-J**) Starch +/+ group was used as a control for post hoc Dunnett's test. All data are presented as mean ± SEM. *$p < 0.05$
Remark: Please clarify the difference with B) in this same figure.

Figure 5

Fig. 5 Protective effects of low-dose aspirin in G × E mice.
(A–D) Results of behavioral tests performed to evaluate the effects of aspirin treatment (n = 12–21 mice per group). (A) Quantification of spontaneous locomotor activity in the open field. (B) Pre-pulse inhibition at 70 dB. (C) Object location test of working memory. (D) Quantification of nest-building skills over 8 h. (E) Immunohistochemical images of fibrin and the endothelial cell marker tomato lectin. (F) Measurement of the area of fibrin immunoexpression in (E). The mean intensity of the fibrin-immunopositive area of the entire image was measured for each section (n = 3 mice per group). (G) Extracellular glucose concentrations in the dialysis buffer at each time point (n = 4–6 mice per group). Post hoc Tukey's multiple comparisons test of groups at each time point, $^{###}p < 0.001$, $^{##}p < 0.01$, $^{#}p < 0.05$ for Starch +/+ vs. Suc *glo1/+*, $^{$$$}p < 0.001$, $^{$}p < 0.05$ for Starch +/+ Aspi vs. Suc *glo1/+*, $^{&}p < 0.05$ for Starch +/+ vs. Suc *glo1/+* Asp . All data are presented as mean ± SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Figure 6

Fig. 6 Angiopathy in postmortem brains from individuals with psychiatric disorders.
(**A**) Representative immunohistochemical fibrin images in the BA9 region of postmortem brains from controls and patients with
schizophrenia (SZ) or bipolar disorder. (**B**) Measurement of the area of fibrin immunoexpression in (**A**). The mean intensity of the fibrin-
immunopositive area of the entire image was measured for each section. (**C**) Representative immunohistochemical images of fibrin
(magenta) and the endothelial cell marker tomato lectin (green) in postmortem brains from a patient with SZ. Fibrin-positive areas are
merged with areas of vascular endothelial cell marker expression. (**D**) Diagrams describing the hypothesis proposed to explain functional
and behavioral abnormalities in control (CTL) mice (left) and G × E mice (right) (see the Discussion for details). All data are presented
as mean ± SEM. **$p < 0.01$

Supplementary Figure 1 Characterization of regional GLO1 expression in WT and heterozygous *Glo1* mutant mice fed starch or sucrose.

(**A**) Western blot analysis of GLO1 protein expression using tubulin as internal control. The cerebral cortex, including the hippocampus, was used as loading sample. (**B**) Densitometric analysis of GLO1 protein expression (n = 3 or 4 mice per group). To quantify expression, GLO1 band intensities in (**A**) were divided by the intensities of the corresponding tubulin bands. (**C**) Body weight trajectories. No significant differences were observed among groups (n = 6–10 mice per group). (**D, E**) Measurement of aripiprazole- or aspirin-containing water consumption per day to estimate drug intake (n = 4 mice per group). Student's *t*-test was used for statistical analysis. All data are presented as mean ± SEM. **$p < 0.01$, *$p < 0.05$

Supplementary Figure 2 Analyses of psychiatric disease-related phenotypes in *Glo1* heterozygotic mice.

(**A, B**) Behavioral analyses in the four groups of mice (n = 18–23 mice per group). (**A**) Elevated plus maze test to evaluate anxiety. (**B**) Preference for novel mouse in the three-chamber test, calculated as [(time spent exploring novel mouse)/(total time spent exploring novel mouse and novel object)] × 100%. (**C**) Extracellular dopamine concentration in the medial prefrontal cortex measured at 20-min intervals using an *in vivo* microdialysis system (n = 7–10 mice per group). (**D–I**) Effect of aripiprazole treatment on behavioral phenotypes (n = 12–18 mice per group). (**D**) Effects of aripiprazole treatment on the acoustic startle response, (**E**) pre-pulse inhibition (PPI) using a 70-dB pre-pulse, (**F**) object location test, (**G**) self-grooming, (**H**) elevated plus maze performance, and (**I**) nest-building skill. In (**I**), post hoc Tukey's multiple comparisons test of groups at specific time points, ###$p < 0.001$, #$p < 0.05$ for Starch +/+ vs. Suc *glo1*/+ Aripi, $$$p < 0.01$ for Starch +/+ Aripi vs. Suc *glo1*/+ Aripi, $$$$p < 0.001$ for Starch +/+ vs. Suc *glo1*/+, *$p < 0.05$ for Starch +/+ Aripi vs. Suc *glo1*/+. All data are presented as mean ± SEM. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$

Supplementary Figure 3 Analyses of psychiatric disease-related phenotypes in *Disc1* heterozygotic mice.
(**A-E**) Behavioral analyses in the four groups of mice (n = 8–11 mice per group). (**A**) Spontaneous locomotor activity in the open-field test. (**B**) Acoustic startle responses. (**C**) Pre-pulse inhibition (PPI) using a 70 dB pre-pulse. (**D**) Object location test. (**E**) Preference for novel mouse in the three-chamber test, calculated as Fig. S2B. All data are presented as mean ± SEM. ***p < 0.001

Supplementary Figure 4 GLO1 cellular localization in the prefrontal cortex of sucrose-fed wild-type mice.
(**A–F**) GLO1 localization in astrocytes. (**A**) GLO1 co-immunostaining with an astrocyte marker (ALDH1L1) or an endothelial cell marker (tomato lectin). (**B**) Merged GLO1/ALDH1L1 image from (**A**). Yellow arrows in (**B**) indicate cells with GLO1/ALDH1L1 colocalization. (**C**) Merged GLO1/lectin image from (**A**). White arrows denote representative GLO1-positive cells close to endothelial cells. (**D, E**) High magnification images of GLO1 co-immunostaining with ALDH1L1 or with tomato lectin. (**E**) Merged GLO1/ALDH1L1 image from (**D**). (**F**) 2D intensity histogram of the two indicated channels to identify GLO1 colocalization with ALDH1L1 or tomato lectin, interpreting the strength of a relationship based on R as follows: R ≥ 0.7, strong; 0.7 > R ≥ 0.4, moderate; 0.4 > R ≥ 0.2, weak; and R < 0.2, none or very weak. (**G**) Average R-value of colocalization data including (**F**) from three different locations of three mice. (**H**) GLO1 co-immunostaining with the neuronal marker NeuN and the microglial marker IBA1. (**I**) Merged GLO1/NeuN image from (**H**). Yellow arrows indicate neurons with mild GLO1 immunoreactivity. (**J**) Merged GLO1/IBA1 image from (**H**). (**K**) GLO1 immunostaining and DAPI staining in *Glo1* homozygous mice.

Supplementary Figure 5 Fructose-derived AGE4 accumulation in microglia of sucrose-fed mice.
(**A, B**) Immunohistochemical images of the microglial marker IBA1 and AGE4 in the medial prefrontal cortex region (mPFC). (**C**) Merged AGE4/IBA1 image. (**D**) High magnification images of AGE4 co-immunostaining with IBA1. (**E**) Measurement of the area of AGE4 immunoexpression in (**B**). The mean AGE4-immunopositive area intensity of the entire image was measured in each section (n = 4 mice per group). (**F**) Mean IBA1 fluorescence intensities of five randomly selected cells per image in four independent mice. (**G**) Immunohistological images of IBA1- and CD68-positive microglia in the mPFC region. White arrows in (**G**) indicate microglial cells containing CD68-positive areas >20 μm². (**H**) Percentage of CD68-positive cells per total microglial cells in each image in (**G**). All data are presented as mean $\pm$ SEM. ***$p < 0.001$, **$p < 0.01$

Supplementary Figure 6 Blood vessels Characterization in the four groups.
(**A**) Western blot analysis of glucose transporter 1 (Glut1) protein expression using tubulin as internal control. The cerebral cortex, including the hippocampus, was used as loading sample. (**B**) Densitometric analysis of Glut1 protein expression (n = 3 mice per group). To quantify expression, Glut1 band intensities in (**A**) were divided by the intensities of the corresponding tubulin bands. (**C**) Measurement of capillary diameter. Vessels <10 μm in diameter were defined as capillaries, and the mean diameter of 10 randomly selected capillaries per image measured in Fig. 4C (n = 3 mice per group). All data are presented as mean ± SEM.

Supplementary Figure 7 Protective effects of aspirin against the development of psychiatric phenotypes in G × E mice.
(**A–C**) Effects of aspirin treatment on the acoustic startle response (**A**), nest-building skill (**B**), elevated plus maze test performance (**C**) and dopamine concentration (**D**). (**E**) Parvalbumin (PV) Immunohistochemistry in the medial prefrontal cortex (mPFC). (**F**) Number of PV-positive cells in the mPFC (n = 4 mice per group). (**G**) Immunohistological images of GFP-positive astrocytes and ALDH1L1 in the mPFC region. (**H**) Percentage of GFP-positive cells per total ALDH1L1-positive cells in each image in (**G**). (**I**) Mean fluorescence GFP intensities of 10 randomly selected cells per image in (**G**) (from four independent mice). In (**B**), post hoc Tukey's multiple comparisons test of groups at specific time points, $^{###}p < 0.001$, $^{##}p < 0.01$ for Starch +/+ vs. Suc *glo1*/+ Aspi, $^{\$\$}p < 0.01$, $^{\$}p < 0.05$ for Starch +/+ Aspi vs. Suc *glo1*/+ Aspi. In (**D**), post hoc Bonferroni's multiple comparisons test of groups at specific time points (0 min and 40 min), $^{###}p < 0.001$ for Suc *glo1*/+ vs. Starch +/+, $^{\$\$\$}p < 0.001$ for Suc *glo1*/+ vs. Starch +/+ Aspi, $^{\&\&\&}p < 0.001$ for Suc *glo1*/+ Aspi vs. Starch +/+, $^{††}p < 0.01$ for Suc *glo1*/+ Aspi vs. Suc *glo1*/+, $^{‡‡}p < 0.01$ for Suc *glo1*/+ Aspi vs. Starch +/+ Aspi. All data are presented as mean ± SEM. $^{***}p < 0.001$, $^{**}p < 0.01$, $^{*}p < 0.05$

Supplementary Figure 8 ROS detection in the four groups.
(**A**) Reactive oxygen species (ROS) detection with *dihydroethidium* (*DHE*). (**B**) Merged DHE/Nucleus (TO-PRO 3) image. All cells are DHE positive, according to nucleus images. (**C**) ROS detection images without *DHE*. (**D**) Fluorescent intensity of DHE positive cells in (**A**). The mean DHE fluorescent intensity of each cell was measured in each section (n = 4 mice per group). All data are presented as mean ± SEM.