Calcium homeostasis plays important roles in the internalisation and activities of the small synthetic antifungal peptide PAF26

Akira JT Alexander[1], Alberto Munoz[2], Jose F. Marcos[3] and Nick D. Read[4].

1 Institute of Infection, Immunity & Inflammation, The University of Glasgow, 464 Bearsden Rd, Bearsden, Glasgow, G61 1QH
2 Medical Research Council Centre for Medical Mycology at the University of Exeter, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter, EX4 4QD
3 Instituto de Agroquímica y Tecnología de Alimentos (IATA), Consejo Superior de Investigaciones Científicas (CSIC), Avda Agustín Escardino 7, 46980 Paterna, Valencia, Spain
4 Manchester Fungal Infection Group, Infection, Immunity & Respiratory Medicine, University of Manchester, Core Technology Facility, 46 Grafton St, Manchester, M13 9WU

ABSTRACT

Fungal diseases are responsible for the deaths of over 1.5 million people worldwide annually. Antifungal peptides represent a useful source of antifungals with novel mechanisms-of-action, and potentially provide new methods of overcoming resistance. Here we investigate the mode-of-action of the small, rationally designed synthetic antifungal peptide PAF26 using the model fungus *Neurospora crassa*. Here we show that the cell killing activity of PAF26 is dependent on extracellular Ca^{2+} and the presence of fully functioning fungal Ca^{2+} homeostatic/signalling machinery. In a screen of mutants with deletions in Ca^{2+}-signalling machinery, we identified three mutants more tolerant to PAF26. The Ca^{2+} ATPase NCA-2 was found to be involved in the initial interaction of PAF26 with the cell envelope. The vacuolar Ca^{2+} channel YVC-1 was shown to be essential for its accumulation and concentration within the vacuolar system. The Ca^{2+} channel CCH-1 was found to be required to prevent the translocation of PAF26 across the plasma membrane. In the wild type, Ca^{2+} removal from the medium resulted in the peptide remaining trapped in small vesicles as in the *Δyvc-1* mutant. It is therefore apparent that cell killing by PAF26 is complex and unusually dependent on extracellular Ca^{2+} and components of the Ca^{2+}-regulatory machinery.

AUTHOR SUMMARY

Life threatening diseases can be caused when fungi invade human tissues. These invasions often occur when a person's immune defences are down, often due to treatments for cancer or transplantation. These infections are commonly buried deep within the body and as such are difficult to access and treat. Current medications are often highly toxic to the patient.

There is also a worrying rise in drug resistance seen in fungi sampled from patients, with

infections effectively untreatable – a death sentence. Antifungal peptides such as PAF26

provide a possible solution by offering a cheap and rapidly produced alternative to

conventional drugs. However, unlike antibacterial peptides, little is known about how these

small molecules mostly exert their effects and cause death. Using live-cell imaging and

deletion mutants, this study provides an analysis of the important roles that Ca^{2+}-

homeostasis and Ca^{2+}-signalling, and possible accompanying vacuolar fusion, play during

the dynamic internalization and interaction with and within the fungal cell following PAF26

treatment.

Introduction

Fungal infections today are among the most difficult diseases to manage in humans(Kohler, Casadevall *et al.* 2014). Fungi collectively kill over 1.5 million people annually which is more than malaria and similar to the death toll from tuberculosis (Brown, Denning *et al.* 2012, Bongomin, Gago *et al.* 2017). Increasing resistance to the limited arsenal of antifungal drugs is a serious concern, especially for Candida and Aspergillus infections, for which the therapeutic options are limited. Overall, there is an urgent need to develop new antifungal strategies to tackle fungal infections (Denning and Bromley 2015, Nicola, Albuquerque *et al.* 2019).

Antifungal peptides (AFPs) and peptide-related molecules are being intensively studied as alternatives for the therapeutic control of pathogenic fungi (Matejuk, Leng *et al.* 2010, Duncan and O'Neil 2013, Rautenbach, Troskie *et al.* 2016, Nicola, Albuquerque *et al.* 2019). A detailed understanding of their antimicrobial mechanisms is of high priority if peptides are to be considered as useful antifungal agents. These studies may also aid the identification of novel targets for antifungal therapy (Muñoz, Gandía *et al.* 2013, Rautenbach, Troskie *et al.* 2016). Furthermore, this mechanistic understanding is guiding the de novo design and modification of natural peptides in order to circumvent their limitations (e.g. instability, toxicity, interactions with other drugs, poor kinetics, resistance mechanisms etc) and thus improve their antimicrobial efficacy (Nicola, Albuquerque *et al.* 2019). Overall, AFPs o er promising alternatives to standard therapies as anti-infectives and immunomodulatory agents with mechanisms-of-action which are less prone to resistance induction compared to conventional antibiotics (Mahlapuu, Håkansson *et al.* 2016).

PAF26 is a synthetic hexapeptide that has been shown to be highly effective at killing filamentous fungi whilst showing low toxicity against human and bacterial cells (Munoz, Lopez-Garcia *et al.* 2006). Unlike membrane permeabilising antimicrobials, initial investigations into the mode-of-action of PAF26 indicated that it did not directly permeabilise the plasma membrane. Instead, at low fungicidal concentrations, PAF26 has a dynamic antifungal mechanism-of-action that involves at least three stages: peptide interaction with the fungal cell envelope (cell wall and/or plasma membrane), endocytic internalisation and accumulation in the vacuole followed by vacuolar expansion. At a certain point, PAF26 is actively transported out of the vacuole into the cytoplasm, followed by a series of complex and specific intracellular effects whose relationship with the eventual death of the target fungus is still unclear (Muñoz, Marcos *et al.* 2012). Two functional and separate motifs (cationic and hydrophobic domains) in the peptide amino acid sequence have been identified as playing important roles in the antimicrobial mode-of-action (Muñoz, Harries *et*

al. 2013). As a result of these studies, PAF26 has been proposed as a simple peptide model

for the characterization and study of cationic, cell-penetrating antifungal peptides (Muñoz,

Marcos *et al.* 2012, Muñoz, Gandía *et al.* 2013).

From average measurements of fungal cell populations expressing the genetically

encoded Ca^{2+}-reporter aequorin, cytosolic free Ca^{2+} concentrations ($[Ca^{2+}]_{cyt}$) within

Neurospora crassa spore germlings treated with a low inhibitory concentration of PAF26

(2.5-5 µM) have been shown to exhibit a biphasic increase in response to PAF26, and this

biphasic $[Ca^{2+}]_{cyt}$ increase is completely dependent on the PAF26 and extracellular Ca^{2+}

concentrations. The second phase of the biphasic $[Ca^{2+}]_{cyt}$ increase was found to be energy

dependent because it was blocked by treatment with the metabolic inhibitor, sodium azide

(NaN_3). This may link PAF26 internalization with actin-dependent endocytosis. Consistent

with endocytic internalization playing a key role in PAF26 internalization, is the inhibition by

the F-actin inhibitor, Latrucunulin A and a reduced uptake rate in endocytosis by the

endocytic mutants *Δrvs-167*, *Δrvs-161* and *Δrab-5* (Muñoz, Marcos *et al.* 2012, Muñoz,

Gandía *et al.* 2013). At high fungicidal concentrations (15 µM), PAF26 killed cells even after

NaN_3 treatment, although in a different manner. Under these conditions, the peptide first

bound to the cell envelope as before, but was then observed to directly enter the cytoplasm,

indicating passive transport across the plasma membrane at high concentrations (Muñoz,

Marcos *et al.* 2012).

In the current research, evidence for Ca^{2+}-signalling having a significant role in the

PAF26 mode-of-action was analysed by testing and analysing the PAF26 sensitivity of

deletion mutants defective in different components of their Ca^{2+}-signalling machinery. The

pattern and kinetics of peptide interaction, internalization and distribution within the cells of

PAF26-resistant mutants were compared with the parental wild type by using fluorescently

labelled PAF26 combined with live-cell imaging. The role of external Ca^{2+} in these processes

and vacuolar fusion was assessed.

RESULTS

Sensitivity of fungal cells to killing by PAF26 is dependent on extracellular Ca^{2+}

To confirm that extracellular Ca^{2+} plays a role in determining the sensitivity of *N.*

crassa to PAF26 (Muñoz, Marcos *et al.* 2012), and thus Ca^{2+} homeostasis being involved in

the mode-of-action of PAF26; the wild type strain was grown in: (1) standard Vögels medium

(VM) (0.64 mM Ca^{2+}); (2) Ca^{2+} free VM (in which Ca^{2+} had been omitted); and (3) VM with

twice the normal concentration of Ca^{2+} (1.28 mM). The IC_{50} values for the wild type under

these different media are shown in Fig. 1A. The IC_{50} in standard VM was 3.7 ± 0.6 µM and in

VM lacking Ca^{2+} was 6.4 ± 0.9 µM, a significant increase in resistance against PAF26 ($P \leq$

0.001). VM with 2 X Ca^{2+} had an IC_{50} value of 2.5 ± 0.4 µM, a significant increase in

sensitivity to PAF26 ($P \leq$ 0.01) compared with standard VM. These results are consistent

with a Ca^{2+}-dependent mechanism playing a role in determining the sensitivity of *N. crassa*

to PAF26 involving uptake of Ca^{2+} from the external medium.

PAF26 requires a functional $[Ca^{2+}]_{cyt}$ homeostatic machinery in order to kill fungal cells

To assess whether Ca^{2+}-homeostasis and signalling play a role in the mode-of-action

of PAF26; 24 homokaryotic deletion mutants of genes encoding proteins predicted to be

components of the *N. crassa* Ca^{2+}-homeostatic machinery were screened for increased

PAF26 tolerance / sensitivity. The predicted locations of these proteins in the fungal cell are

shown in Fig. 1C; The results of the screen are in Table 1.

The first group of mutants screened were defective in the following Ca^{2+}-channel

proteins: CCH-1 and MID-1, which are components of the high affinity Ca^{2+}-influx system

(HACS); FIG-1, which forms the low-affinity Ca^{2+}-influx system (LACS); and, the vacuolar

Ca^{2+} channel, YVC-1. Two of these mutants (∆*cch-1* and ∆*yvc-1*) exhibited significantly

increased tolerance against PAF26 ($P \leq$ 0.001) when compared with the wild type (Fig. 1B).

The IC_{50} values of ∆*cch-1* and ∆*yvc-1* were both above the level (~ 10 µM) at which PAF26

directly permeabilizes the plasma membrane of *N. crassa* wild type cells (Muñoz, Marcos *et*

al. 2012).

N. crassa has eight Ca^{2+} ATPases (Zelter *et al.*, 2004), for which seven deletion

mutants were available as homokaryons for screening: Δ*nca-1*, Δ*nca-2*, Δ*nca-3*, Δ*pmr-1*,

Δ*ph-7*, Δ*cta-3* and Δ*cta-4. Δnca-2,* Δ*cta-3* and Δ*cta-4* exhibited significantly increased

tolerance against PAF26 ($P \leq$ 0.001) when compared with the wild type (Fig. 1B); with *Δnca-*

2 approaching inhibitory levels of PAF26 that directly permeabilise the membrane in the wild

type.

Homokaryotic deletion mutants of six of the eight predicted *N. crassa* antiporters

involved in Ca^{2+} transport (Zelter *et al.*, 2004) were available for screening: Δ*cax*, Δ*trm-14*,

Δ*trm-15,* Δ*trm-16*, Δ*trm-18*, and Δ*trm-20* (Fig. 1B, Table 1). Δ*trm-20* showed significantly

greater tolerance ($P \leq$ 0.01) to PAF26 compared with the wild type although this was less

than a two-fold increase (Fig. 1B).

Fig. 1. Ca²⁺ plays a significant role in the mode-of-action of PAF26. (A) shows the effects
of both the removal and increase of free Ca^{2+} in the media on the IC_{50} of PAF26 by fungal cells.
Increasing the level lowers the concentration at which PAF26 is effective and removing Ca^{2+}
increases the concentration. Both of these results are significant; One way ANOVA with Dunnet's
comparison test: $F(2,12) = 58.85$, $P = \leq 0.001$, $R^2 = 90.75$, $R^2(adj) = 89.21$. The Low Ca^{2+} VM (M =
6.410 ± 0.941) was significantly different from the Normal VM (M = 3.868 ± 0.288) at $P = \leq 0.001$ and
the 2X Ca^{2+} (M = 2.342 ± 0.331) was significantly different from Normal VM at $P = \leq 0.01$ **(B)** shows
the effects of deleting components of the Ca^{2+} homeostasis machinery on susceptibility to PAF26. 6
were significantly more tolerant than the wildtype; One way ANOVA with Dunnet's comparison test:
$F(17,106) = 60.13$, $P = \leq 0.001$, $R^2 = 90.61$, $R^2(adj) = 89.10$. $\Delta cch-1$ (M = 15.595 ± 2.649), $\Delta yvc-1$ (M
= 11.375 ± 1.419), $\Delta nca-2$ (M = 8.442 ± 0.838), $\Delta cta-3$ (M = 9.099 ± 1.677) and $\Delta cta-4$ (M = 8.212 ±
0.594) were significantly more tolerant than the wildtype (M = 3.756 ± 0.193) at $P = \leq 0.001$, and

Δtrm-20 (M = 6.637 ± 1.234) was significant at *P = ≤ 0.01*. Each assay consisted of 8 technical and 3
biological replicates, figures are representative. The predicted localisations of these is shown in **(C)**
based on the localization of their protein orthologs in *S. cerevisiae* yeast cells and vegetative hyphae
of *N. crassa* (Wada, Ohsumi *et al.* 1987, Yang and Sachs 1989, Bertl and Slayman 1990, Diamond,
Zasloff *et al.* 1991, Antebi and Fink 1992, Iida, Nakamura *et al.* 1994, Lapinskas, Cunningham *et al.*
1995, Levina, Lew *et al.* 1995, Paidhungat and Garrett 1997, Erdman, Lin *et al.* 1998, Kanzaki 1999,
Benito, Garciadeblas *et al.* 2000, Locke, Bonilla *et al.* 2000, Muller, Locke *et al.* 2001, Palmer, Zhou
et al. 2001, Courchesne 2002, Courchesne and Ozturk 2002, Denis and Cyert 2002, Gupta, Ton *et*
al. 2003, Kaiserer, Oberparleiter *et al.* 2003, Muller, Mackin *et al.* 2003, Zhou, Batiza *et al.* 2003,
Zelter, Bencina *et al.* 2004, Brand, Shanks *et al.* 2007, Hallen and Trail 2008, Lew, Abbas *et al.*
2008, Benito, Garciadeblás *et al.* 2009, Bormann and Tudzynski 2009, Bowman, Draskovic *et al.*
2009, Binder, Chu *et al.* 2010, Bowman, Abreu *et al.* 2011, Cavinder, Hamam *et al.* 2011).

Table 1: The half maximal inhibitory concentration (IC$_{50}$) of PAF26 in Ca^{2+}
signalling/homeostasis mutants.

	Strain	Genotype	IC$_{50}$ PAF 26 µM	±	Source	*P*
	WT	74-OR23-1V	3.76	0.19	FGSC2489	
	Δ*cch-1*	**NCU02762**	**13.94**	**1.26**	**Chu, 2013**	**≤0.001**
	Δ*mid-1*	NCU06703	5.74	1.33	FGSC11708	
	Δ*yvc-1*	**NCU07605**	**11.37**	**1.42**	**FGSC11253**	**≤0.001**
	Δ*fig-1*	NCU02219	5.89	0.74	FGSC17273	
	Δ*nca-1*	NCU03305	5.96	1.07	FGSC13287	
	Δ*nca-2*	**NCU04736**	**12.73**	**2.37**	**FGSC13071**	**≤0.001**
	Δ*nca-3*	NCU05154	5.12	0.42	FGSC13037	
	Δ*pmr-1*	NCU03292	4.36	0.24	FGSC11616	
	Δ*ph-7*	NCU08147	5.08	0.59	FGSC11256	
Ca^{2+} signalling / homeostasis	Δ*cta-3*	NCU07966	9.10	1.68	FGSC11409	≤0.001
	Δ*cta-4*	NCU04898	8.22	0.59	FGSC13040	≤0.001
	Δ*ena-5*	NCU10143	3.27	0.29	FGSC12645	
	Δ*cax*	NCU07075	4.11	0.50	FGSC11249	
	Δ*trm-15*	NCU00916	5.27	0.62	FGSC11686	
	Δ*trm-14*	NCU00795	6.19	0.38	FGSC12375	
	Δ*trm-18*	NCU06366	4.29	0.56	FGSC11408	
	Δ*trm-16*	NCU02826	6.06	1.00	FGSC11529	
	Δ*trm-20*	NCU08490	6.64	1.23	FGSC12468	≤0.01
	Δ*plc-1*	NCU06245	3.27	0.46	FGSC11411	
	Δ*plc-2*	NCU01266	6.24	1.60	FGSC12022	
	Δ*plc-3*	NCU02175	3.59	0.46	FGSC12023	
	Δ*plc-4*	NCU09655	4.17	0.30	FGSC11271	
	Δ*CamK-1*	NCU09123	6.99	0.26	FGSC12547	
	Δ*CamK-2*	NCU02283	5.09	0.36	FGSC12449	
	Δ*CamK-3*	NCU06177	3.84	0.39	FGSC11536	

Mutants highlighted in bold were tolerant to PAF26 at levels approaching passive membrane permeabilization and were selected for further study.

PAF26 is taken up by wild type macroconidia into vacuoles which fuse and then lyse before releasing the peptide into the cytoplasm

In order to compare the dynamic pattern of peptide-cell interactions between the wild

type and the mutants, PAF26 labelled with the fluorophore TAMRA (TMR-PAF26) was

imaged. Fluorescent labelling has been previously employed to monitor, using confocal live-

cell imaging, the dynamic localization of PAF26 or related peptides in a range of fungi. We

are confident that the dynamic pattern of TMR-PAF26 staining provides a faithful localization

of the unlabelled peptide in fungal cells and labelled/unlabelled peptides have similar IC$_{50}$

values (Muñoz, Marcos *et al.* 2012, Muñoz, Gandía *et al.* 2013, Muñoz, Harries *et al.* 2013).

Previously it has been shown that wild type macroconidia treated with a low fungicidal

concentration of FITC-PAF26 (2.5 μM) exhibit a similar time-dependent staining pattern as

germlings treated with the same concentration of PAF26 (Muñoz, Marcos *et al.* 2012). A

similar localization pattern of TMR-PAF26 interaction, internalization and distribution was

observed with ungerminated macroconidia to that of FITC-PAF26. The use of macroconidia

allowed a basic monitoring of cell volume, surface area and developmental state in relation

to possible variations in morphogenesis and staining patterns.

Fig. 2. Graphical representation showing that the interaction between PAF26 and
macroconidia is altered through deletion of Ca^{2+}-channels and pumps. (**A**) shows maximum
intensity projections from a Z series captured using confocal microscopy. Initially, the peptide
appears at the cell envelope and within small vesicles (a). The peptide is then accumulated in larger
vesicles and vacuoles (b) until there appears a lower number of larger vacuoles (c) which will

eventually fuse into a single vacuole which then releases the peptide into the cytoplasm (d). Scale
bar is 5 µm. **(B)** shows quantification of this interaction over time for the deletion mutants compared
to the wild type; there are marked changes in the localisation of the peptide during the time course.
(C) shows the interaction between PAF26 and macroconidia is Ca^{2+}-dependent and energy
dependent. This is demonstrated by the significant delay in the uptake of PAF26 in the wild type
when extracellular free Ca^{2+} is removed using BAPTA, at top. The peptide remains trapped in small
vesicles and there is little transport into the vacuolar system and consequently a reduction in internal
membrane permeabilisation. The lower figure shows the effect of the removal of cellular free energy
on the *Δcch-1* mutant using the metabolic inhibitor NaN_3. There is no uptake of PAF26 into the cell
and therefore the direct translocation across the plasma membrane must be an energy dependent
process. N = 100 macroconidia minimum per field of view, 10 fields of view per timepoint, repeated
at least twice.

208 The dynamic pattern of TMR-PAF26 in Ca^{2+} homeostasis mutant
209 staining provides insights into the localization and roles of the
210 defective mutant proteins

In order to gain more insight into which stages of the PAF26 interaction, uptake and

distribution process the CCH-1, YVC-1 and NCA-2 proteins influence, the time-dependent

staining by TMR-PAF26 in each of the *Δcch-1, Δyvc-1,* and *Δnca-2* mutants were compared

with that in the wild type. As a baseline for the comparison, four staining patterns were

identified in the wild type macroconidia that could be readily quantified at 30 min intervals

following treatment with 3.5 µM TMR-PAF26. These staining patterns related to different

events in the time-dependent uptake and distribution to different organelles which precedes

eventual cell death. They were: (a) Accumulation in multiple small vesicles. These were

predicted to be mostly endosomes typically up to ~ 1 µm in width. (b) Accumulation in 1-3

large vacuole(s) typically greater than ~ 1.5 µm in width. (c) Accumulation in the cytoplasm

and excluded from vacuoles. This is the stage just after the peptide has been released out

from vacuoles. (d) Accumulation in both the cytoplasm and vacuoles. This is the stage after

which the intracellular membranes have become permeabilized and thus represents

commitment to cell death, as previously reported (Muñoz, Marcos *et al.* 2012).

In the wild type macroconidia (Figs. 2B and 3A), this quantitative analysis showed that after

treatment for 30 min with TMR-PAF26, all the macroconidia had interacted with TMR-PAF26

and many were beginning to internalise the peptide into small vesicles. After 60 min, many of

the conidia had already undergone vacuolar expansion and some had exported the peptide

into the cytoplasm, and this correlated with a transient reduction in the percentage of cells

with peptide only localized within small vesicles. After 60 min, 40% of the cells had actively

transported the peptide out of their vacuoles and into the surrounding cytoplasm. In another

40% of the cells at the 60 min time point, vacuolar membrane permeabilisation had occurred

and the peptide had equilibrated throughout the cells. By 120 min, 57% of the cells were

permeabilised (Figs. 2B and 3A).

Fig. 3. The interaction with PAF26 is altered through deletion of Ca^{2+}-channels and -pumps as shown by visualisation of wild type and mutant conidia stained with 3.5 µM TAMRA labelled PAF26. These figures show maximum intensity projections of Z stacks false coloured with the 'physics' lookup table to highlight the different fluorescent intensities measured. (**A**) The wild type conidia undergo a sequence of events which ultimately results in cell death (Muñoz, Marcos *et al.*, 2012). (**B**) The *Δyvc-1* mutant retains the peptide in small vesicles, without the formation of a large central vacuole, as in the wild type. (**C**) The *Δnca-2* mutant appears much reduced in peptide interaction, with very few conidia showing the peptide bound to the envelope. (**D**) The process is reversed in the *Δcch-1* mutant, where the peptide first appears within the cytoplasm and is pumped into small vesicles which then appear to fragment. No conidia were seen to completely lose loaded peptide. The internalisation process is Ca^{2+} dependent and energy-dependent, as chelation of Ca^{2+} with BAPTA (**E**), and treatment with Sodium azide (NaN$_3$) (**F**), significantly reduced uptake of TAMRA-PAF26. 1000 individual macroconidia per timepoint per strain. Assay was repeated twice and images are representative. Scale bar is 20 µm.

YVC-1 is required for transport of PAF26 from vesicles to vacuoles

When the *Δyvc-1* mutant was treated with the labelled peptide (Fig. 2B, 3B), it rapidly appeared at the cell envelope. Unlike with the wild type, the peptide was internalised into small vesicles at a more-or-less linear rate with much remaining trapped at the cell surface. Accumulation of the peptide in vacuoles was much slower and reached a maximum of 6% of the cells with this localization pattern after 120 min. Virtually no cells were observed with peptide exclusively in the cytoplasm at the three time points of measurement over the 120 min by which time only 5% of cells were internally permeabilised. After 120 min, 79% of the *Δyvc-1* cells had taken up TMR-PAF26 compared with 100% in the wild type. Thus, less TMR-PAF26 was taken up by *Δyvc-1* cells, most accumulated in small vesicles and few cells had internal membrane permeabilisation after 120 min (5% compared with 57% in the wild type). These results are consistent with YVC-1 being required for the transport of PAF26 from vesicles to vacuoles.

NCA-2 is required for the interaction of PAF26 and the cell envelope

When the *Δnca-2* mutant was treated with TMR-PAF26 (Fig. 2B and 3C) the cell envelopes of the macroconidia were visibly less fluorescent than that of the wild type. Thus, the affinity of TMR-PAF26 for the cell envelope of this mutant seemed to be reduced and its staining was delayed compared with that of the wild type. The *Δnca-2* macroconidia also internalized TMR-PAF26 at a far reduced rate compared with the other mutants and the peptide mostly became trapped within the small vesicles, resulting in virtually no TMR-PAF26 being taken up by vacuoles and a correspondingly extremely small percentage (~ 2%) of the cells having permeabilised internal membranes after 120 min (Fig. 2B).

CCH-1 plays a key role in the energy- and Ca^{2+}-dependent internalisation of PAF26 by fungal cells

The pattern of TMR-PAF26 internalization and intracellular transport within

macroconidia of the *Δcch-1* mutant appeared to be opposite to that in the wild type (Fig. 3D).
After treatment for 30 min the peptide was primarily localized in the cytoplasm (in 67% of
cells) and to a much lesser extent in small vesicles (~ 18% of cells) and to a very low level in
large vacuoles (~ 5% of cells). Between 30 and 90 min, TMR-PAF26 was removed from the
cytoplasm and increased in amount in the small vesicles and large vacuoles. Between 90
and 120 min the large vacuoles containing TMR-PAF26 are fragmenting into smaller
vesicles. This was reflected by a dramatic increase in the number of stained small vesicles
(Fig. 2B).

The overall rate of internalisation of TMR-PAF26 by *Δcch-1* macroconidia was faster
than in the wild type because in the former it was initially taken up directly into the cytoplasm
whilst in the latter it first appeared intracellularly in small vesicles that are presumed to be
mostly endosomes. It seems unlikely that TMR-PAF26 is taken up by means of non-specific
permeabilization of the plasma membrane because this would likely have resulted in rapid
cell death. Furthermore, 30 min after the addition of TMR-PAF26 it was clear that the
vacuolar membrane had not become permeabilized because the peptide was excluded from
the vacuoles of cells in which the cytoplasm was fluorescent.

In order to clarify whether the passage of PAF26 across the plasma membrane into
the cytoplasm of the *Δcch-1* mutant was a result of passive translocation or active uptake,
macroconidia were pre-treated with the metabolic inhibitor NaN_3 (Muñoz, Marcos *et al.* 2012)
at a concentration of 5 µM for 15 min before the addition of TMR-PAF26 and subsequent
imaging and quantification of localization over 120 min (Figs. 2C and 3F).

In the presence of NaN_3, over the whole 120 min period of incubation with TMR-
PAF26, the peptide remained bound to the cell envelope and was not internalized by most of
the macroconidia. Clearly the metabolic inhibitor NaN_3 had almost completely abolished the
movement of TMR-PAF26 across the plasma membrane indicating that the uptake of the
peptide into the cells of the *Δcch-1* mutant is an ATP-dependent process. The results also
showed that the rate of internal membrane permeabilisation, whilst faster over the first 30
min, remained similar to that of the non-azide treated *Δcch-1* cells; after 120 min ~ 25% of
the cells were fluorescent throughout the cell. These results are consistent with the uptake of
TMR-PAF26 by this subpopulation of *Δcch-1* macroconidia being energy independent as a
result of a passive process.

These results are consistent with the normal endocytic internalization of PAF26 being
dependent on the Ca^{2+} channel protein, CCH-1. As CCH-1 appears to initiate PAF26
internalization by endocytosis (see previous section), this suggests that the uptake of Ca^{2+}
from the external medium may be mediated by this Ca^{2+} channel, which has previously been

reported to be immunolocalized to the plasma membrane (Locke *et al.*, 2000). Furthermore,
we had previously shown that removal of Ca^{2+} from VM made macroconidia more resistant
to being killed by PAF26. To attempt to mimic the effect of *cch-1* deletion in the wild type,
Ca^{2+} was removed from the external medium by the addition of the Ca^{2+} chelator BAPTA (5
mM) 30 min prior to the TMR-PAF26 treatment (Figs. 2C and 3E). Rather than preventing
the entry of the peptide, the BAPTA treatment had the unexpected effect of trapping the
peptide in small vesicles. Over the 120 min period very little TMR-PAF26 had localized in
large vacuoles, in the cytoplasm and only ~ 11% of the macroconidia had fluorescence
throughout the cell. The phenotype of the *Δcch-1* mutant is consistent with the CCH-1
protein playing a key role in the energy- and Ca^{2+}-dependent internalisation of PAF26 by
fungal cells.

The $[Ca^{2+}]_{cyt}$ response during PAF26 treatment is disrupted in the $[Ca^{2+}]_{ext}$ tolerant mutants

The wild type had previously been shown to undergo a dose dependent biphasic rise
in $[Ca^{2+}]_{cyt}$ upon addition of PAF26 at concentrations between 0.8 and 2.0 µM (Muñoz,
Marcos *et al.* 2012). The experiment was repeated here but with over a concentration range
of 1.25-10 µM PAF26. For all measurements, the unstimulated $[Ca^{2+}]_{cyt}$ resting level was
measured for 50 sec, in the wild type this was calculated to be 0.05 ± 0.02 µM (Fig. 4A),
before the peptide was added at time 0. After treatment with a low final concentration of
PAF26 (1.25 µM), an immediate increase in $[Ca^{2+}]_{cyt}$ to 0.19 ± 0.01 µM was observed and
this was sustained throughout the measurement period (1,200 sec) and slightly increased
towards the end of this period (Fig. 4B). When the PAF26 added was at a final concentration
of 2.5 µM, which is close to its IC_{50} value, the initial increase in $[Ca^{2+}]_{cyt}$ was slightly greater
(to 0.22 ± 0.03 µM) (Fig. 4B). Again, this was followed by a period of sustained $[Ca^{2+}]_{cyt}$
increase, but there was also a more pronounced exponential increase in $[Ca^{2+}]_{cyt}$ which
began at ~ 900 sec after treatment. With the higher dose, there was also an increase in the
standard deviations of the measurements with time. This is due to aequorin consumption
during the course of the experiment, resulting in less sensitivity after prolonged exposure to
high $[Ca^{2+}]$ (note that luminescence measurements from 6 wells are averaged per time
point). When the wild type was treated with 5 µM PAF26, which was above its IC_{50} value, the
$[Ca^{2+}]_{cyt}$ response followed the same general pattern but with a much larger initial $[Ca^{2+}]_{cyt}$
increase to 0.32 ± 0.02 µM and a shorter period of sustained increase before the $[Ca^{2+}]_{cyt}$
increase became exponential at ~ 430 sec (Fig. 4B). Furthermore, after 590 sec following
peptide treatment, $[Ca^{2+}]_{cyt}$ spiking was observed. The $[Ca^{2+}]_{cyt}$ response was further
accentuated after treatment with a very high dose (10 µM) of PAF26 resulting in a rapid
initial rise in $[Ca^{2+}]_{cyt}$ to 0.54 ± 0.03 µM, followed by an immediate exponential increase in

$[Ca^{2+}]_{cyt}$. This in turn was followed by the $[Ca^{2+}]_{cyt}$ spiking and then a sudden drop in $[Ca^{2+}]_{cyt}$

after ~ 220 sec, which then exponentially increased and subsequently underwent spiking

again (Fig. 4B).

The unstimulated resting level of $[Ca^{2+}]_{cyt}$ in the $\Delta yvc\text{-}1$ mutant was measured as 0.02

± 0.01 µM (Fig 4A), significantly different at P >0.01. The $[Ca^{2+}]_{cyt}$ response of the $\Delta yvc\text{-}1$

strain to PAF26 was markedly different to that of the wild type. Whilst there was a similar

initial immediate, dose dependent increase in $[Ca^{2+}]_{cyt}$, there was no second exponential

increase in $[Ca^{2+}]_{cyt}$, which was particularly evident after treating the wild type with PAF26 at

2.5 µM or above (Fig. 4B). Instead the $\Delta yvc\text{-}1$ mutant underwent a slight reduction in $[Ca^{2+}]_{cyt}$

followed by a sustained, more-or-less constant $[Ca^{2+}]_{cyt}$ level that was significantly raised

compared with the unstimulated resting levels recorded prior to treatment. Thus the $\Delta yvc\text{-}1$

mutant lacked the second $[Ca^{2+}]_{cyt}$ increase of the biphasic $[Ca^{2+}]_{cyt}$ response of the wild

type, which is consistent with this second $[Ca^{2+}]_{cyt}$ increase resulting from the release of Ca^{2+}

from the vacuole. The resting level of $[Ca^{2+}]_{cyt}$ in the $\Delta nca\text{-}2$ mutant was measured at 0.05

µM ± 0.02, Fig 4A, and was not significantly different from the wild type. In general terms,

the $\Delta nca\text{-}2$ mutant showed a similar response to PAF26 as did the $\Delta yvc\text{-}1$ mutant. However,

the Ca^{2+} signatures of the $\Delta nca\text{-}2$ and $\Delta yvc\text{-}1$ mutants were clearly but not dramatically

different from each other; there was a more pronounced dose dependent $[Ca^{2+}]_{cyt}$ increase

in the $\Delta nca\text{-}2$ strain compared with that in $\Delta yvc\text{-}1$ in response to PAF26 (Fig. 4B). The

resting level of $[Ca^{2+}]_{cyt}$ in the $\Delta cch\text{-}1$ mutant was 0.03 ± 0.00 µM Fig. 4A, significantly

different to the wild type at P > 0.01. Interestingly, deletion of $cch\text{-}1$ does not inhibit the initial

increase in $[Ca^{2+}]_{cyt}$ after PAF26 addition, surprising given the lack of evidence for another

Ca^{2+} channel at the cell surface. The Ca^{2+} signatures of the $\Delta cch\text{-}1$ mutant in response to

different concentrations of PAF26 were broadly similar to those of the $\Delta yvc\text{-}1$ and $\Delta nca\text{-}2$

mutants (Figs. 4B). Thus the $\Delta cch\text{-}1$ mutant also showed an initial $[Ca^{2+}]_{cyt}$ increase but

lacked the second $[Ca^{2+}]_{cyt}$ increase of the typical biphasic response of the wild type to

PAF26. These results suggest there may be an unidentified L-type channel within the

plasma membrane, not CCH-1, being responsible for the initial $[Ca^{2+}]_{cyt}$ increase in response

to PAF26.

The influence on the $[Ca^{2+}]_{cyt}$ response to 5.0 µM PAF26 of the wild type following

pre-treatment with 5 mM BAPTA (to chelate extracellular Ca^{2+}) or with the L-type Ca^{2+}

channel blockers diltiazem or verapamil (both at 5 mM), was analysed (Fig. 4C). A 15 min

pre-treatment with ddH_2O (the control) caused a biphasic increase in $[Ca^{2+}]_{cyt}$ whilst all of the

other treatments prevented any significant $[Ca^{2+}]_{cyt}$ increase (Fig. 2). These results are

consistent with the biphasic $[Ca^{2+}]_{cyt}$ increase being dependent on extracellular Ca^{2+} as

reported before (Muñoz, Marcos *et al.* 2012) and on L-type Ca^{2+} channel activity. The only

known L-type Ca^{2+} channel in *N. crassa* is CCH-1(Zelter, Bencina *et al.* 2004). These results
also suggest that the second component of the biphasic [Ca^{2+}]$_{cyt}$ increase is dependent on
the first [Ca^{2+}]$_{cyt}$ increase.

Fig. 4. The [Ca^{2+}]$_{cyt}$ response to PAF26 is altered by deletion of Ca^{2+} channels and pumps. (**A**)
shows the unstimulated resting level of the wildtype and the Ca^{2+} homeostatic mutants: the resting
level in *Δyvc-1* and *Δcch-1* were significantly different from the wild type. One way ANOVA with
Dunnet's comparison test: $F(3,108) = 32.83$, $P = \le 0.001$, $R^2 = 47.70$, R^2(adj) = 46.24. (**B**) shows the
[Ca^{2+}]$_{cyt}$ response to PAF26 by population wide measurements using the genetically encoded reporter
aequorin supplied with an excess of coelenterazine. The [PAF26] μM is shown above each
measurement series and was added at time 0, baseline measurements were recorded for 50s before
the addition of the peptide. (**C**) shows that this response in the wild type is both dependent on external
Ca^{2+} and on the presence of an external Ca^{2+} channel, the biphasic response to 5μM PAF26 (seen in
the ddH$_2$O control) is completely abolished following chelation of Ca^{2+} with BAPTA or treatment with
L-Type Ca^{2+} channel blockers. All measurements are averages of 6 individual wells in a 96 well plate,
with [Ca^{2+}]$_{cyt}$ calculated following quenching of the remaining aequorin using EtOH and Ca^{2+}. Error
bars are standard deviations. All experiments were repeated a minimum of three times and the above
figures are representative.

Imaging using the fluorescent reporter GCaMP6 shows PAF26 does not cause a dose dependent rise in $[Ca^{2+}]_{cyt}$

In order to investigate whether PAF26 was indeed causing a dose dependent rise in $[Ca^{2+}]_{cyt}$, conidia expressing GCaMP6s were imaged during PAF26 treatment using widefield fluorescence microscopy. Individual macroconidia were then isolated from the background in FIJI and fluorescence measured over time. As Figure 5A shows, the individual responses are far more varied than the measurements using aequorin suggested. In the middle of the timecourse, the moment the PAF26 reaches the field of view and interacts with the cells can be clearly seen in a sudden increase in Ca^{2+} spiking in all the individuals. Given that the aequorin measurements are across a whole population, it is not hard to see how the cumulative effects of an increase in Ca^{2+} spikes and waves could be interpreted as a dose dependent rise in $[Ca^{2+}]_{cyt}$.In order to determine whether there was a predictable response to PAF26 addition, several repeated experiments were run in which the peptide was added and fluorescence intensity recorded. No distinct pattern or regularity was found within any of the data sets, except for an increase in Ca^{2+} signalling after PAF26 addition. As the only noticeable response, the amplitude and frequency of these Ca^{2+} spikes was quantified. Conidia were treated either with ddH_2O or ddH_2O containing 3.5 µM PAF26. A second set were pre-treated with 5 mM BAPTA before ddH_2O or PAF26 treatment. The conidia treated with ddH_2O have a relatively low frequency of Ca^{2+} spiking with the mean being 4 spikes over the course of the measurement. This is consistent with our findings as to the rate of Ca^{2+} spiking in germinating and fusing conidia (Read, unpublished). When the cells are treated with PAF26 however, there is a marked increase in both frequency and amplitude. The mean frequency of the Ca^{2+} spikes increases to 19 and the amplitude increases to around 6 times the resting level from 4 (Figure 5B). Both of these results are significant at $P < 0.01$. When the conidia are pre-treated with BAPTA however, all Ca^{2+} spiking ceases completely. Whilst there were no apparent recurring patterns in the spiking events, observations appeared to show that they correspond to vesicles and vacuoles coming into close proximity. A Ca^{2+} trace is shown in Fig. 5C and images are shown in Fig. 5D. The images shown in Fig. 5D are taken every five seconds from the timeframes indicated by the black arrows in Fig. 5C. The first Ca^{2+} spike marked by *, appears to correspond with the meeting of the two vacuoles at the lower part of the image, whether they fuse or not is unclear. In the images corresponding to the second spike marked with *, the fusion of two vacuoles is clear at the top of the image.

Figure 5. PAF26 does not cause a dose dependent rise in [Ca^{2+}]$_{cyt}$. (**A**) shows the relative fluorescence intensity of several macroconidia expressing GCaMP6s during PAF26 treatment. Fluorescence values are normalised and given as arbitrary values. After the addition of PAF26 there is a marked increase in the amplitude and frequency of Ca^{2+} spiking. (**B**) Quantification of this revealed an actual increase in spiking from an average of 4 spikes per 30 min to 19 spikes per 30 min (i). The spiking was entirely dependent on the presence of external Ca^{2+} as chelation with BAPTA reduced the frequency of spiking to 0. The amplitude of the individual spikes also increased from 4 to 6 arbitrary units (ii). (**C**) these spikes often appeared to correspond to the meeting of vesicles and vacuoles within the cell; the images captured between the two arrows are shown in (**D**) which shows widefield fluorescence images captured every 5 s. The two spikes marked by * in C appear to correspond to vacuolar fusion events. Scale bar = 2µm.

YVC-1 is localized in vacuoles and CCH-1 is both plasma membrane and vacuolar localized

In *N. crassa*, CCH-1 and YVC-1 have not yet been definitively localised while NCA-2 has previously been localised to the vacuole and plasma membrane in mature hyphae (Bowman, Draskovic *et al.* 2009). We conducted experiments to localize NCA-2 tagged with RFP in macroconidia of *N. crassa* and the data showed labelling of the vacuoles, but no

obvious signal at the plasma membrane (Fig. 6).).

In order to localise the CCH-1 and YVC-1 proteins, plasmids were constructed based
on FJ457002 for N-terminal tagging and FJ457006 for C-terminal tagging (Honda and Selker
2009).

Both the C- and N- terminal YVC-1:GFP fusion proteins fluoresced, although the N-
tagged protein showed a stronger more stable fluorescence. Fluorescence mainly appeared
within vacuoles in conidia, germ tubes and mature hyphae (Fig. 6). The CCH-1:GFP fusion
products produced very little fluorescence, both from N and C tagged versions. This
appeared to be localised to vacuoles within conidia and intracellular membranes within older
hyphae. Single punctate spots of GFP fluorescence could however be seen at the plasma
membrane of mature hyphae, Fig.6.

Fig. 6. Localization of CCH-1 and YVC-1 labelled with GFP in macroconidia, germlings and
vegetative hyphae. Representative widefield fluorescence images of the different cell types of GFP
tagged CCH-1 and YVC-1 are shown. As NCA-2 has been previously localised in hyphae, only

macroconida re shown for this RFP expressing strain(Bowman, Draskovic *et al.* 2009). CCH-1
appears to be localised in the vacuolar system in macroconidia and germlings, but also appears as
distinct points in the plasma membrane of mature hyphae. YVC-1 appears to localise to the vacuolar
system in all cell types. RFP tagged NCA-2 localises to the vacuole in macroconidia. Bar = 5 μm.

DISCUSSION

Ca^{2+} plays important roles in the mode-of-action of PAF26

In this research, evidence for Ca^{2+} signaling and homeostasis having a significant

role in the PAF26 mode-of-action was obtained initially by testing the PAF26 sensitivity of

conidia grown in the presence or absence of calcium. When the level of calcium in the media

is kept at a minimal level, tolerance increases. Conversely when the level is raised, tolerance

decreases. This was confirmed with deletion mutants defective in different components of

their cellular transport machinery of Ca^{2+}. Deletion of most of the Ca^{2+} channels significantly

increased the concentrations at which PAF26 inhibits fungal growth. The elimination of the

high affinity Ca^{2+} plasma membrane channel CCH-1 and the vacuolar channel YVC-1 as well

as the Ca^{2+} ATPase resulted in inhibitory concentrations of the peptide approaching the

point at which passive membrane permeabilisation occurs. Ca^{2+} has been shown to be an

important factor in the mechanistics of PMAP-23 against *Candida albicans*, the disruption

results in reactive oxygen species (ROS) accumulation which triggers apoptosis (Kim and

Lee 2019). This effect is also seen in the peptide CGA-N9 against *C. tropicalis* (Li, Chen *et*

al. 2019). High levels of cytosolic free Ca^{2+} have also been shown to confer protection to *C.*

albicans against the antimicrobial MUC7 12mer, thought to be by a resulting change in the

cell membrane properties preventing peptide entry (Lis, Liu *et al.* 2010). It appears in this

case that the elevated levels of cytosolic free Ca^{2+} are a stress response designed to

maximise survival by initiating the endocytosis of PAF26. There are significant changes to

the pattern of peptide cell interaction in the mutants and to the quantity of peptide the

mutants take up. The mutations in the Ca^{2+} distribution machinery appear to disrupt the

internalisation and accumulation of the peptide which in turn increases tolerance to PAF26.

Deletion of *nca-2* appears to influence the binding of PAF26 to the cell envelope, the *Δnca-2*

mutant has been shown to accumulate up to 10 fold more Ca^{2+} than wild type cells,

suggesting that NCA-2 serves to remove Ca^{2+} from the cell (Bowman, Abreu *et al.* 2011).

We found no significant difference in the resting level of [Ca^{2+}[$_{cyt}$ from the wild type however

under unstimulated conditions. Interestingly however, the *Δnca-2* mutant has a membrane

potential reversed from the wild type due to a lack of cell surface H$^+$ATPase function

(Hamam and Lew 2012). In *S. cerevisiae* the H$^+$ATPase is one of the most abundant cell

surface proteins (Bagnat, Chang *et al.* 2001) and makes up a significant amount (up to 10%)

of the cell surface in *N. crassa* (Bowman, Blasco *et al.* 1981). PAF26 has been shown to

cause a rapid depolarization of the membrane in wild type cells in an energy independent
manner (Muñoz, Marcos *et al.* 2012). Given that PAF26 has a net positive charge, the
membrane potential reversed *Δnca-2* should technically not be inhibited in PAF26
membrane binding from an electrostatic view. It is therefore possible to propose that PAF26
directly inhibits H^+-ATPase action, possibly by direct binding; misfunctioning H^+--ATPase at
the plasma membrane is sent to the vacuole for degradation in yeast, a possible mechanistic
for the accumulation of PAF26 (Bagnat, Chang *et al.* 2001, Liu, Sitaraman *et al.* 2006).
When the peptide is internalised in the *Δyvc-1* mutant, it remains in small vesicles with very
little sign of the peptide entering the vacuolar system; adding support for the hypothesis that
YVC-1 and a threshold amount of PAF26 is required to initiate vacuolar fusion. Vacuolar
fusion occurs through conformational change of the docking SNARE proteins, triggered by
the release of Ca^{2+} from the vacuole in *S. cerevisiae* (Bayer, Reese *et al.* 2003, Merz and
Wickner 2004, Coonrod, Graham *et al.* 2013). In *S. cerevisiae*, isolated vacuoles are able to
catalyse their own fusion through the release of Ca^{2+} from yvc1p present in the vacuolar
membrane (Peters and Mayer 1998). This does not appear to be the case in *N. crassa*
however, as both deletion of YVC-1 and removal of extracellular Ca^{2+} resulted in the trapping
of the peptide in small vesicles. Therefore extracellular Ca^{2+} is required to initiate the release
of Ca^{2+} from the vacuoles and trigger fusion. This raises questions as to why internal
membrane fusion is reliant on external stimuli. In the *Δcch-1* mutant the peptide was directly
translocated across the plasma membrane into the cytoplasm in an energy dependent
manner, before accumulation in vacuoles, meaning PAF26 does not kill by being present in
the cytosolic space. CCH-1 is therefore required to initiate the endocytic pathway. Energy
dependent peptide uptake into the vacuolar system is also seen in the *Penicillium*
chrysogenum Penicillium antifungal protein B (PAFB), where the fungal cells show no signs
of cell death as long as the peptide remains in the vacuole (Huber, Hajdu *et al.* 2018). This is
similarly seen in the *Neosartorya (Aspergillus) fischeri* antifungal protein (NFAP), which is
also accumulated in the vacuole in an energy dependent maner. Again, this peptide also
does not exert its antifungal effects until it is present within the cytoplasm (Hajdu, Huber *et*
al. 2019). This contrasts the findings here of PAF26 being present within the cytoplasmic
space without killing the cells. The importance of the vacuole in the mode of action of many
antifungal peptides is becoming increasingly apparent. A genetic screen of *S. cerevisiae*
revealed tolerance to the plant antifungal defensins NbD6 and SBI6 could be increased by
deletion of genes involved in vacuolar transport (Parisi, Doyle *et al.* 2019).

Ca^{2+} measurement at the population level revealed that the Ca^{2+} responses to PAF26
were markedly different in the PAF26 tolerant mutants. Whilst all underwent a similar initial
increase in $[Ca^{2+}]_{cyt}$ the second biphasic increase seen in the wildtype was abolished. This

initial increase was independent of CCH-1, as the signal in the $\Delta cch-1$ mutant increased, but

was dependent on external Ca^{2+} and L-type Ca^{2+} channels. The fact that L-type channel

blockers completely stop the $[Ca^{2+}]_{cyt}$ increase as does the chelation of external Ca^{2+} using

BAPTA would suggest that there is another, as yet unknown cell surface Ca^{2+} channel.

These findings mirror those of Binder *et al.,* 2010, who found that deletion of *cch-1* did not

prevent the increase in $[Ca^{2+}]_{cyt}$ in response to *P. chrysogenum Penicillium* antifungal protein

(PAF). When these measurements were taken at the level of the individual, rather than the

population, it became apparent that PAF26 does not in fact cause a dose dependent rise in

$[Ca^{2+}]_{cyt}$ but rather increases both the amplitude and frequency of Ca^{2+} spiking, again

dependent on the presence of external Ca^{2+}. These findings have allowed us to propose the

model shown in Fig. 7 for the role of Ca^{2+} signalling and homeostasis in the mode of action

of PAF26. Now the challenge is to identify the trigger of vacuolar release of PAF26 and the

initiator of cell death.

Fig 7. Proposed role of Ca^{2+} signalling in the mode of action of PAF26. 1 initially the cell is unstimulated and maintains a low [Ca^{2+}]$_{cyt}$ and stable membrane potential. **2** PAF26 inhibits the cell surface H$^+$ATPase initiating membrane depolarisation. **3** Depolarisation of the plasma membrane activates the voltage gated CCH-1, causing [Ca^{2+}]$_{cyt}$ to rise. **4** This rise in [Ca^{2+}]$_{cyt}$ initiates the turnover of the H$^+$ATPase including bound PAF26 by endocytosis. **5** further rises in [Ca^{2+}]$_{cyt}$ caused by external Ca^{2+} entering through CCH-1 initiate Ca^{2+} release through YVC-1 **6** driving the fusion of membranes. **7** PAF26 is released from the H$^+$ATPase as it is degraded and accumulates in the vacuolar system.

Materials and methods

Neurospora crassa stocks

N. crassa strains were obtained from the Fungal Genetics Stock Centre (FGSC, 563   University of Missouri, Kansas City, USA), unless otherwise specified. FGSC2489 (*matA*,74-

OR23-1VA) was used as the wild type control. *N. crassa* strains generated in this study were
derived from FGSC9717 (*matA,Δmus-51::bar⁺; his-3⁻*), FGSC9720 (*matA,Δmus-52::bar⁺; his-*
3⁻) and FGSC6103 (*matA,his-3⁻*) *N. crassa* was cultured and maintained according to FGSC
protocols (www.fgsc.net/Neurospora/NeurosporaProtocolGuide.htm) unless otherwise
stated.

For microscopy, conidia were cultured in liquid VM for up to 4 hours at 30°C using 8
well Nunc™ Lab-Tek™ II Chambered Coverglass.

N. crassa transformation

572 Electroporation of macroconidia was carried out using the following settings:
573 resistance 600 Ω, voltage 1.5 kV/cm and capacitance 25 μFd.

Heterokaryon purification

575 *N. crassa* strains that were obtained as heterokaryons, were purified to homokaryons
576 by standard procedure (www.fgsc.net/Neurospora/NeurosporaProtocolGuide.htm).

Plasmids

In order to create pTef1AeqS, aequorin was amplified from pAB19 using the primers
AeqS-F and AeqS-R. The vector pCC019 was digested with PacI and EcoRI and the vector
gel purified. Aequorin was then amplified with AeqS-IF-Fw and AeqS-IF-Rv to add overlaps
homologous to the backbone whilst maintaining digestion sites and the vector was
assembled using Gibson assembly. To generate pYVC1CGFP and pYVC1NGFP, the *yvc-1*
ORF was amplified from gDNA using the primers NC-YVC1-AMP-FW and NC-YVC-1-AMP-
RV. The purified product was then amplified using NC-YVC1-N-FW & NC-YVC1-N-RV for N
terminal tagging and NC-YVC1-C-FW & NC-YVC1-C-RV for C terminal tagging. The
FJ457002 vector was digested using PacI and XbaI and the FJ457006 vector digested with
AscI and NotI. Following gel extraction both vectors were assembled using Gibson
assembly.

Tagging of CCH-1 was not straightforward due to the size of the ORF - 6.5kb. The
CCH-1 ORF was amplified first using the primer sets NC-CCH1 AMP-FW and NC-CCH1-
3.5Rv and NC-CCH1-3.5Fw and NC-CCH1-AMP-RV to amplify the gene in two segments, 5'
and 3'. The complete constructs were built using fusion PCR; primer NC-CCH1-C-FW, 5'
segment, 3' segment and primer NC-CCH1-C-RV, the N terminal tagging vector used the
primers NC-CCH1-N-FW and NC-CCH1-N-RV. Following successful fusion the vectors were
assembled as for YVC-1. Primer sequences are given in supplementary material.

IC$_{50}$ values

IC$_{50}$ values were calculated using clear, U bottom polystyrene microtitre plates. PAF 26 was prepared at two concentrations, 35 µM and 30 µM, and diluted two-fold appropriately in to reach the final experimental concentrations . Conidia were diluted to a concentration of 1 X 10^6 in 20% liquid VM and 50 µl dispensed into each well. Experimental PAF26 concentrations (µM) were: [17.50][15.00][8.75][7.50][4.38][3.75][2.19][1.88][1.09][0.94][0.55][0]. Final conidial concentration was of 5 X 10^5 in 10% VM. Absorbance measurements were obtained at 610 nm. This wavelength is close to the 595nm at which the relationship between optical density and fungal biomass is linear (Broekaert, Terras *et al.* 1990).

Calcium measurements

Coelenterazine (#C-7001, www.biosynth.com)was prepared by dissolving in ice cold methanol (MetOH) under a protective atmosphere in complete darkness to a concentration of 10 µg/µl. Stocks were diluted to the working concentration of 3.175 µg in 10 µl MetOH, 74.78 µM.

Conidia were diluted to a concentration of 1 X 10^6 in 10% VM, coelenterazine was added to a final concentration of 2.5 µM. 100 µl of conidial suspension was used per well of a white microtitre plate and the plates incubated at 25°C in the dark for 6 hours. The light output of aequorin was measured by counting the photons emitted by a single well over one second, each well in a row of six being measured once every cycle of 7 seconds. After recording baseline luminescence, a single row was treated with PAF26 and the light output measured over time. Following this the second row was run with 100 µl 3M CaCl$_2$:20% ethanol (EtOH) injected on cycle 8 to immediately discharge all the aequorin.

[Ca^{2+}] calculation

The raw relative light units (RLU) were converted to [Ca^{2+}] using the formula:(Bonora, Giorgi *et al.* 2013)

$$\left[\text{Ca}^{2+}\right]\text{M} = \left(\frac{\left(\frac{L}{L_{\max}} \times \lambda\right)^{\frac{1}{n}} + \left(\left(\frac{L}{L_{max}} \times \lambda\right)^{\frac{1}{n}} \times K_{TR}\right) - 1}{K_R - \left(\left(\frac{L}{L_{max}} \times \lambda\right)^{\frac{1}{n}} \times K_R\right)} \right)$$

Where:
- *L* is the light intensity at sampling time
- *L$_{max}$* is the total light emitted at sampling time

• K_R is the constant for calcium bound state **(7,230,000)**

• K_{TR} is the constant for calcium unbound state **(120)**

• λ is the rate constant for aequorin consumption at saturating $[Ca^{2+}]$ **(1)**

• **n**   is the number of Ca^{2+} binding sites **(2.99)**

In order to calculate L_{max} the total RLU (RLU_{total}) available was calculated from the discharge

of aequorin (RLUd) using the trapezoid rule formula, where the total RLUd available at time

point t_n equals:

$$\mathbf{RLUd}_{t_{n-1}} + \frac{(\mathbf{t_n} - \mathbf{t_{n-1}}) \times (\mathbf{RLUd_{t_n}} + \mathbf{RLUd_{t_{n-1}}})}{\mathbf{2}}$$

632 and $RLUd_{total}$ is the final value reached. Switching to the experimental data (RLUe), L_{max} was

633 calculated for each time point as:

$$\mathbf{RLUd_{total}} - \left(\mathbf{RLUe}_{t_{n-1}} + \left(\frac{(\mathbf{t_n} - \mathbf{t_{n-1}}) \times \left(\mathbf{RLUe_{t_n}} + \mathbf{RLUe_{t_{n-1}}}\right)}{\mathbf{2}} \right) \right)$$

Peptides and peptide handling

Peptides were ordered from GenScript (www.genscript.com) at > 98 % purity. Stock

solutions were prepared in 50:50 dimethyl sulfoxide (DMSO):H_2O buffer. Working peptide

solutions were prepared in ddH_2O.

Microscopy

All microscopy was carried out on a Nikon Eclipse® TE2000-E inverted microscope,

using a Nikon Plan Apo 60 X 1.2 N.A. DIC H water immersion objective and Nikon G-2A and

B-2A filter sets, excitation was provided by a CoolLED illumination system set at 550 nm for

use with the G-2A filter or 470 nm for use with the B2-A filter.

Confocal microscopy was carried out using a Leica TCS SP8 equipped with two

hybrid GaAsP detectors (HyD) and two photomultiplier tubes (PMT). Excitation was provided

using either the Leica tuneable white light laser (450 -750 nm), an argon laser (458nm,

476nm, 488nm and 496nm) or UV laser (405nm). Images were captured using LAS X

software and the Leica 63 X water immersion objective. All image handling and analysis was

carried out using Fiji (fiji.sc/Fiji).

Quantification of Ca^{2+} signalling

Fluorescence values were exported from Fiji and analysed in Excel. Data was

normalised using feature scaling. This scales the data to removes irregularities in GCaMP6

expression and photon yield. The function used was: x^{l} = MIN + (x-min_x)(MAX-MIN)/(max_x-

min_x). Where MIN and MAX are the required scale range, x is the original value and x^{l} is the

normalised value. min_x is the minimum value for that data set and max_x is the maximum,
max_x was defined as the maximum value from all experiments plus 10 to avoid artificial
amplification of noise. The noise was removed using an IF function: =IF($x>y,x,0$) where x is
the data point and y is the noise threshold limit. Finally, an IF(AND($x>x_{-1}$, $x>x_{+1}$),1,0)
function, where x_{-1} and x_{+1} are the surrounding measurements, quantified each peak in
GCaMP6 fluorescence

Data availability

661   Data used in this study is available at: http://dx.doi.org/10.17632/sb2r2zsxdz.1

Funding Statement

663 Funding for this project was provided by a Biotechnology and Biological Sciences Research
664 Council Research Studentship to AJTA. The funders had no role in study design, data
665 collection and analysis, decision to publish, or preparation of the manuscript.

References

Antebi, A. and G. R. Fink (1992). "The yeast Ca(2+)-ATPase homologue, PMR1, is required for normal
Golgi function and localizes in a novel Golgi-like distribution." Molecular Biology of the Cell **3**(6): 633-
654.
Bagnat, M., A. Chang and K. Simons (2001). "Plasma Membrane Proton ATPase Pma1p Requires Raft
Association for Surface Delivery in Yeast." Molecular Biology of the Cell **12**(12): 4129-4138.
Bayer, M. J., C. Reese, S. Buhler, C. Peters and A. Mayer (2003). "Vacuole membrane fusion: V0
functions after trans-SNARE pairing and is coupled to the Ca2+-releasing channel." The Journal of cell
biology **162**(2): 211-222.
Benito, B., B. Garciadeblás, J. Pérez-Martín and A. Rodríguez-Navarro (2009). "Growth at High pH
and Sodium and Potassium Tolerance in Media above the Cytoplasmic pH Depend on ENA ATPases in
Ustilago maydis." Eukaryotic Cell **8**(6): 821-829.
Benito, B., B. Garciadeblas and A. Rodriguez-Navarro (2000). "Molecular cloning of the calcium and
sodium ATPases in Neurospora crassa." Molecular Microbiology **35**(5): 1079-1088.
Bertl, A. and C. L. Slayman (1990). "Cation-selective channels in the vacuolar membrane of
Saccharomyces: dependence on calcium, redox state, and voltage." Proceedings of the National
Academy of Sciences **87**(20): 7824-7828.
Binder, U., M. Chu, N. D. Read and F. Marx (2010). "The Antifungal Activity of the Penicillium
chrysogenum Protein PAF Disrupts Calcium Homeostasis in Neurospora crassa." Eukaryotic Cell **9**(9):
1374-1382.
Bongomin, F., S. Gago, O. R. Oladele and W. D. Denning (2017). "Global and Multi-National
Prevalence of Fungal Diseases—Estimate Precision." Journal of Fungi **3**(4).
Bonora, M., C. Giorgi, A. Bononi, S. Marchi, S. Patergnani, A. Rimessi, R. Rizzuto and P. Pinton (2013).
"Subcellular calcium measurements in mammalian cells using jellyfish photoprotein aequorin-based
probes." Nature Protocols **8**(11): 2105-2118.

Bormann, J. and P. Tudzynski (2009). "Deletion of Mid1, a putative stretch-activated calcium channel
in Claviceps purpurea, affects vegetative growth, cell wall synthesis and virulence." <u>Microbiology</u>
155(12): 3922-3933.
Bowman, B. J., S. Abreu, E. Margolles-Clark, M. Draskovic and E. J. Bowman (2011). "Role of Four
Calcium Transport Proteins, Encoded by *nca-1* , *nca-2* , *nca-3* , and *cax* , in Maintaining Intracellular
Calcium Levels in Neurospora crassa." <u>Eukaryotic Cell</u> **10**(5): 654-661.
Bowman, B. J., F. Blasco and C. W. Slayman (1981). "Purification and characterization of the plasma
membrane ATPase of Neurospora crassa." <u>J Biol Chem</u> **256**(23): 12343-12349.
Bowman, B. J., M. Draskovic, M. Freitag and E. J. Bowman (2009). "Structure and Distribution of
Organelles and Cellular Location of Calcium Transporters in *Neurospora crassa*." <u>Eukaryotic Cell</u>
8(12): 1845-1855.
Brand, A., S. Shanks, V. M. S. Duncan, M. Yang, K. Mackenzie and N. A. R. Gow (2007). "Hyphal
Orientation of Candida albicans Is Regulated by a Calcium-Dependent Mechanism." <u>Current Biology</u>
17(4): 347-352.
Broekaert, W. F., F. R. Terras, B. P. Cammue and J. Vanderleyden (1990). "An automated quantitative
assay for fungal growth inhibition." <u>FEMS Microbiology Letters</u> **69**(1-2): 55-59.
Brown, G. D., D. W. Denning, N. A. R. Gow, S. M. Levitz, M. G. Netea and T. C. White (2012). "Hidden
Killers: Human Fungal Infections." <u>Science Translational Medicine</u> **4**(165): 165rv113-165rv113.
Cavinder, B., A. Hamam, R. R. Lew and F. Trail (2011). "Mid1, a Mechanosensitive Calcium Ion
Channel, Affects Growth, Development, and Ascospore Discharge in the Filamentous Fungus
Gibberella zeae." <u>Eukaryotic Cell</u> **10**(6): 832-841.
Coonrod, E. M., L. A. Graham, L. N. Carpp, T. M. Carr, L. Stirrat, K. Bowers, N. J. Bryant and T. H.
Stevens (2013). "Homotypic vacuole fusion in yeast requires organelle acidification and not the V-
ATPase membrane domain." <u>Developmental cell</u> **27**(4): 462-468.
Courchesne, W. E. (2002). "Characterization of a Novel, Broad-Based Fungicidal Activity for the
Antiarrhythmic Drug Amiodarone." <u>Journal of Pharmacology and Experimental Therapeutics</u> **300**(1):
195-199.
Courchesne, W. E. and S. Ozturk (2002). "Amiodarone induces a caffeine-inhibited, MID1-depedent
rise in free cytoplasmic calcium in Saccharomyces cerevisiae: Amiodarone induces a rise in
cytoplasmic calcium in yeast." <u>Molecular Microbiology</u> **47**(1): 223-234.
Denis, V. and M. S. Cyert (2002). "Internal Ca $^{2+}$ release in yeast is triggered by hypertonic shock and
mediated by a TRP channel homologue." <u>The Journal of Cell Biology</u> **156**(1): 29-34.
Denning, D. W. and M. J. Bromley (2015). "How to bolster the antifungal pipeline." <u>Science</u>
347(6229): 1414.
Diamond, G., M. Zasloff, H. Eck, M. Brasseur, W. L. Maloy and C. L. Bevins (1991). "Tracheal
antimicrobial peptide, a cysteine-rich peptide from mammalian tracheal mucosa: peptide isolation
and cloning of a cDNA." <u>Proceedings of the National Academy of Sciences</u> **88**(9): 3952-3956.
Duncan, V. M. S. and D. A. O'Neil (2013). "Commercialization of antifungal peptides." <u>Fungal Biology</u>
<u>Reviews</u> **26**(4): 156-165.
Erdman, S., L. Lin, M. Malczynski and M. Snyder (1998). "Pheromone-regulated Genes Required for
Yeast Mating Differentiation." <u>The Journal of Cell Biology</u> **140**(3): 461-483.
Gupta, S. S., V.-K. Ton, V. Beaudry, S. Rulli, K. Cunningham and R. Rao (2003). "Antifungal Activity of
Amiodarone Is Mediated by Disruption of Calcium Homeostasis." <u>Journal of Biological Chemistry</u>
278(31): 28831-28839.
Hajdu, D., A. Huber, A. Czajlik, L. Tóth, Z. Kele, S. Kocsubé, Á. Fizil, F. Marx, L. Galgóczy and G. Batta
(2019). "Solution structure and novel insights into phylogeny and mode of action of the Neosartorya
(Aspergillus) fischeri antifungal protein (NFAP)." <u>International Journal of Biological Macromolecules</u>
129: 511-522.
Hallen, H. E. and F. Trail (2008). "The L-Type Calcium Ion Channel Cch1 Affects Ascospore Discharge
and Mycelial Growth in the Filamentous Fungus *Gibberella zeae* (Anamorph *Fusarium graminearum*
)." <u>Eukaryotic Cell</u> **7**(2): 415-424.

Hamam, A. and R. R. Lew (2012). "Electrical phenotypes of calcium transport mutant strains of a
filamentous fungus, Neurospora crassa." Eukaryotic cell **11**(5): 694-702.
Honda, S. and E. U. Selker (2009). "Tools for fungal proteomics: multifunctional neurospora vectors
for gene replacement, protein expression and protein purification." Genetics **182**(1): 11-23.
Huber, A., D. Hajdu, D. Bratschun-Khan, Z. Gáspári, M. Varbanov, S. Philippot, Á. Fizil, A. Czajlik, Z.
Kele, C. Sonderegger, L. Galgóczy, A. Bodor, F. Marx and G. Batta (2018). "New Antimicrobial
Potential and Structural Properties of PAFB: A Cationic, Cysteine-Rich Protein from Penicillium
chrysogenum Q176." Scientific Reports **8**(1): 1751.
Iida, H., H. Nakamura, T. Ono, M. S. Okumura and Y. Anraku (1994). "MID1, a novel Saccharomyces
cerevisiae gene encoding a plasma membrane protein, is required for Ca2+ influx and mating."
Molecular and Cellular Biology **14**(12): 8259-8271.
Kaiserer, L., C. Oberparleiter, R. Weiler-G□rz, W. Burgstaller, E. Leiter and F. Marx (2003).
"Characterization of the Penicillium chrysogenum antifungal protein PAF." Archives of Microbiology
180(3): 204-210.
Kanzaki, M. (1999). "Molecular Identification of a Eukaryotic, Stretch-Activated Nonselective Cation
Channel." Science **285**(5429): 882-886.
Kim, S. and D. G. Lee (2019). "Role of calcium in reactive oxygen species-induced apoptosis in
Candida albicans: an antifungal mechanism of antimicrobial peptide, PMAP-23." Free Radic Res
53(1): 8-17.
Kohler, J. R., A. Casadevall and J. Perfect (2014). "The spectrum of fungi that infects humans." Cold
Spring Harb Perspect Med **5**(1): a019273.
Lapinskas, P. J., K. W. Cunningham, X. F. Liu, G. R. Fink and V. C. Culotta (1995). "Mutations in PMR1
suppress oxidative damage in yeast cells lacking superoxide dismutase." Molecular and Cellular
Biology **15**(3): 1382-1388.
Levina, N. N., R. R. Lew, G. J. Hyde and I. B. Heath (1995). "The roles of Ca2+ and plasma membrane
ion channels in hyphal tip growth of Neurospora crassa." J Cell Sci **108 (Pt 11)**: 3405-3417.
Lew, R. R., Z. Abbas, M. I. Anderca and S. J. Free (2008). "Phenotype of a Mechanosensitive Channel
Mutant, mid - 1 , in a Filamentous Fungus, Neurospora crassa." Eukaryotic Cell **7**(4): 647-655.
Li, R., C. Chen, B. Zhang, H. Jing, Z. Wang, C. Wu, P. Hao, Y. Kuang and M. Yang (2019). "The
chromogranin A-derived antifungal peptide CGA-N9 induces apoptosis in Candida tropicalis."
Biochemical Journal **476**(20): 3069-3080.
Lis, M., T. T. Liu, K. S. Barker, P. D. Rogers and L. A. Bobek (2010). "Antimicrobial peptide MUC7 12-
mer activates the calcium/calcineurin pathway in Candida albicans." FEMS Yeast Research **10**(5):
579-586.
Liu, Y., S. Sitaraman and A. Chang (2006). "Multiple Degradation Pathways for Misfolded Mutants of
the Yeast Plasma Membrane ATPase, PMA1." Journal of Biological Chemistry **281**(42): 31457-31466.
Locke, E. G., M. Bonilla, L. Liang, Y. Takita and K. W. Cunningham (2000). "A Homolog of Voltage-
Gated Ca2+ Channels Stimulated by Depletion of Secretory Ca2+ in Yeast." Molecular and Cellular
Biology **20**(18): 6686-6694.
Mahlapuu, M., J. Håkansson, L. Ringstad and C. Björn (2016). "Antimicrobial Peptides: An Emerging
Category of Therapeutic Agents." Frontiers in Cellular and Infection Microbiology **6**.
Matejuk, A., Q. Leng, M. D. Begum, M. C. Woodle, P. Scaria, S. T. Chou and A. J. Mixson (2010).
"Peptide-based Antifungal Therapies against Emerging Infections." Drugs Future **35**(3): 197.
Merz , A. J. and W. T. Wickner (2004). "Trans-SNARE interactions elicit Ca2+ efflux from the yeast
vacuole lumen." The Journal of Cell Biology **164**(2): 195-206.
Muller, E. M., E. G. Locke and K. W. Cunningham (2001). "Differential regulation of two Ca(2+) influx
systems by pheromone signaling in Saccharomyces cerevisiae." Genetics **159**(4): 1527-1538.
Muller, E. M., N. A. Mackin, S. E. Erdman and K. W. Cunningham (2003). "Fig1p facilitates Ca2+ influx
and cell fusion during mating of Saccharomyces cerevisiae." J Biol Chem **278**(40): 38461-38469.

Muñoz, A., M. Gandía, E. Harries, L. Carmona, N. D. Read and J. F. Marcos (2013). "Understanding
the mechanism of action of cell-penetrating antifungal peptides using the rationally designed
hexapeptide PAF26 as a model." Fungal Biology Reviews **26**(4): 146-155.
Muñoz, A., M. Gandía, E. Harries, L. Carmona López, N. Read and J. F. Marcos (2013). Understanding
the mechanism of action of cell-penetrating antifungal peptides using the rationally designed
hexapeptide PAF26 as a model.
Muñoz, A., E. Harries, A. Contreras-Valenzuela, L. Carmona, N. D. Read and J. F. Marcos (2013). "Two
Functional Motifs Define the Interaction, Internalization and Toxicity of the Cell-Penetrating
Antifungal Peptide PAF26 on Fungal Cells." PLOS ONE **8**(1): e54813.
Munoz, A., B. Lopez-Garcia and J. F. Marcos (2006). "Studies on the Mode of Action of the Antifungal
Hexapeptide PAF26." Antimicrobial Agents and Chemotherapy **50**(11): 3847-3855.
Muñoz, A., J. F. Marcos and N. D. Read (2012). "Concentration-dependent mechanisms of cell
penetration and killing by the de novo designed antifungal hexapeptide PAF26: Uptake mechanisms
of the antifungal peptide PAF26." Molecular Microbiology **85**(1): 89-106.
Nicola, A. M., P. Albuquerque, H. C. Paes, L. Fernandes, F. F. Costa, E. S. Kioshima, A. K. R. Abadio, A.
L. Bocca and M. S. Felipe (2019). "Antifungal drugs: New insights in research & development."
Pharmacol Ther **195**: 21-38.
Paidhungat, M. and S. Garrett (1997). "A homolog of mammalian, voltage-gated calcium channels
mediates yeast pheromone-stimulated Ca2+ uptake and exacerbates the cdc1(Ts) growth defect."
Mol Cell Biol **17**(11): 6339-6347.
Palmer, C. P., X. L. Zhou, J. Lin, S. H. Loukin, C. Kung and Y. Saimi (2001). "A TRP homolog in
Saccharomyces cerevisiae forms an intracellular Ca(2+)-permeable channel in the yeast vacuolar
membrane." Proc Natl Acad Sci U S A **98**(14): 7801-7805.
Parisi, K., S. R. Doyle, E. Lee, R. G. T. Lowe, N. L. van der Weerden, M. A. Anderson and M. R.
Bleackley (2019). "Screening the Saccharomyces cerevisiae Nonessential
Gene Deletion Library Reveals Diverse Mechanisms of Action for Antifungal Plant Defensins."
Antimicrobial Agents and Chemotherapy **63**(11): e01097-01019.
Peters, C. and A. Mayer (1998). "Ca2+/calmodulin signals the completion of docking and triggers a
late step of vacuole fusion." Nature **396**(6711): 575-580.
Rautenbach, M., A. M. Troskie and J. A. Vosloo (2016). "Antifungal peptides: To be or not to be
membrane active." Biochimie **130**: 132-145.
Wada, Y., Y. Ohsumi, M. Tanifuji, M. Kasai and Y. Anraku (1987). "Vacuolar ion channel of the yeast,
Saccharomyces cerevisiae." J Biol Chem **262**(36): 17260-17263.
Yang, X. and F. Sachs (1989). "Block of stretch-activated ion channels in Xenopus oocytes by
gadolinium and calcium ions." Science **243**(4894): 1068-1071.
Zelter, A., M. Bencina, B. J. Bowman, O. Yarden and N. D. Read (2004). "A comparative genomic
analysis of the calcium signaling machinery in Neurospora crassa, Magnaporthe grisea, and
Saccharomyces cerevisiae." Fungal Genetics and Biology **41**(9): 827-841.
Zhou, X. L., A. F. Batiza, S. H. Loukin, C. P. Palmer, C. Kung and Y. Saimi (2003). "The transient
receptor potential channel on the yeast vacuole is mechanosensitive." Proceedings of the National
Academy of Sciences **100**(12): 7105-7110.

Supplementary

836 Primers

Name	Sequence
NC-cch1-AMP-FW	ATCTATGGCTGTGACGCTAG
NC-cch1-AMP-RV	GTCATCTTCGCATCTCCA
NC-yvc1-AMP-FW	AGCATGGCAAGGTTCGG
NC-yvc1-AMP-RV	TCACTTCCCATCACTACTGCTACTG
NC-cch1-C-FW	aaccctcacatcaaccaaatctagATGAGCGATCATCCAGCAC
NC-cch1-C-RV	ctccgccgcctccgcccttaattaaGTAAGCCCCCGAGTATCCTC
NC-cch1-N-FW	cggaggcggcggaggcggcgcgccaAGCGATCATCCAGCACCT
NC-cch1-N-RV	ggagctccaccgcggtggcggccgcTTAGTAAGCCCCCGAGTATCCT
NC-yvc1-C-FW	aaccctcacatcaaccaaatctagATGGCAAGGTTCGGGGC
NC-yvc1-C-RV	ctccgccgcctccgcccttaattaaCTTCCCATCACTACTGCTACTGCTACTA
NC-yvc1-N-FW	cggaggcggcggaggcggcgcgccaGCAAGGTTCGGGGCCTT
NC-yvc1-N-RV	ggagctccaccgcggtggcggccgcTCACTTCCCATCACTACTGCTACTG
NC-cch1-3.5Rv	TAGAAAGGGTTGGGTTCACG
NC-cch1-3.5Fw	CGTGAACCCAACCCTTTCTA

839 Plasmids

Plasmid	Backbone	Genotype	Source
FJ457002		*Pccg-1::c-gly::gfp*	Honda & Selker, 2009
FJ457006		*Pccg-1::n-gfp*	Honda & Selker, 2009
pAB19	pAZ6	*Pccg-1-AeqS-PtrpC-bar$^+$*	Marris, 2007
pCC019	FJ457002	*his-3-Ptef-1-gcamp6s*	Chang, 2015
pTef1AeqS	pCC019	*his-3-Ptef-1-AeqS*	This study
pYVC1CGFP	FJ457002	*his-3-Pccg-1-yvc-1-gfp*	This study
pYVC1NGFP	FJ457006	*his-3-Pccg-1-gfp-yvc-1*	This study
pCCH1CGFP	FJ457002	*his-3-Pccg-1-yvc-1-gfp*	This study
pCCH1NGFP	FJ457006	*his-3-Pccg-1-gfp-yvc-1*	This study