Calcium homeostasis plays important roles in the internalisation and activities of the small synthetic antifungal peptide PAF26

Akira JT Alexander[1], Alberto Munoz[2], Jose F. Marcos[3] and Nick D. Read[4*].

1   Institute of Infection, Immunity & Inflammation, The University of Glasgow, 464 Bearsden Rd, Bearsden, Glasgow, G61 1QH
2   Medical Research Council Centre for Medical Mycology at the University of Exeter, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter, EX4 4QD
3   Instituto de Agroquímica y Tecnología de Alimentos (IATA), Consejo Superior de Investigaciones Científicas (CSIC), Avda Agustín Escardino 7, 46980 Paterna, Valencia, Spain
4   Manchester Fungal Infection Group, Infection, Immunity & Respiratory Medicine, University of Manchester, Core Technology Facility, 46 Grafton St, Manchester, M13 9WU
* Correspondence: nick.read@manchester.ac.uk

ABSTRACT

Fungal diseases are responsible for the deaths of over 1.5 million people annually. The range of treatments in current use is limited and, is hampered by drug resistance, which is increasingly found in clinical isolates. Antifungal peptides represent a useful source of antifungals with novel mechanisms-of-action, and potentially provide new methods of overcoming resistance. Here we investigate the mode-of-action of the small, rationally designed synthetic antifungal peptide PAF26. This peptide is a hexamer, which had previously been shown to exhibit highly specific antifungal activity against a range of fungal species. Here we show that the cell killing activity of PAF26 is dependent on extracellular Ca^{2+} and the presence of fully functioning fungal Ca^{2+} homeostatic/signalling machinery. Disruption of this machinery influenced the internalization and vacuolar accumulation of fluorescently labelled peptide and reduced its cell killing effects. In a screen of mutants with deletions in components of the Ca^{2+}-signalling machinery, we identified three mutants more tolerant to PAF26 that approach levels at which the peptide is able to directly permeabilise the plasma membrane. The Ca^{2+} ATPase NCA-2 was found to be involved in the initial interaction of PAF26 with the cell envelope. The vacuolar Ca^{2+} channel YVC-1 was shown to be essential for the initial uptake of the peptide and its accumulation and concentration within the vacuolar system. The Ca^{2+} channel CCH-1 was found to be required to prevent the translocation of PAF26 across the plasma membrane. In the wild type, Ca^{2+} removal from the medium resulted in the peptide remaining trapped in small vesicles (as in the *Δyvc-1* mutant). It is therefore apparent that cell killing by PAF26 is complex and unusually dependent on extracellular Ca^{2+} and different components of the Ca^{2+}-regulatory machinery.

AUTHOR SUMMARY

Life threatening diseases can be caused when fungi invade human tissues. These invasions often occur when a person's immune defences are down, often due to treatments for cancer or transplantation. These infections are commonly buried deep within the body and as such are difficult to access and treat. Current medications are often highly toxic to the patient. There is also a worrying rise in drug resistance seen in fungi sampled from patients, with infections effectively untreatable – a death sentence. Antifungal peptides such as PAF26 provide a possible solution by offering a cheap and rapidly produced alternative to conventional drugs. However, unlike antibacterial peptides, little is known about how these small molecules mostly exert their effects and cause death. Using live-cell imaging and deletion mutants, this study provides an analysis of the important roles that Ca^{2+}-homeostasis and Ca^{2+}-signalling, and possible accompanying vacuolar fusion, play during the dynamic internalization and interaction with and within the fungal cell following PAF26 treatment.

Introduction

Fungal infections today are among the most difficult diseases to manage in humans(Kohler, Casadevall et al. 2014). Fungi collectively kill over 1.5 million people annually which is more than malaria and similar to the death toll from tuberculosis (Brown, Denning et al. 2012, Bongomin, Gago et al. 2017). Increasing resistance to the limited arsenal of antifungal drugs is a serious concern, especially for Candida and Aspergillus infections, for which the therapeutic options are limited. Overall, there is an urgent need to develop new antifungal strategies to tackle fungal infections (Denning and Bromley 2015, Nicola, Albuquerque et al. 2019).

Antifungal peptides (AFPs) and peptide-related molecules are being intensively studied as alternatives for the therapeutic control of pathogenic fungi (Matejuk, Leng et al. 2010, Duncan and O'Neil 2013, Muñoz, Gandía et al. 2013, Rautenbach, Troskie et al. 2016, Nicola, Albuquerque et al. 2019). A detailed understanding of their antimicrobial mechanisms is of high priority if peptides are to be considered as useful antifungal agents. These studies may also aid the identification of novel targets for antifungal therapy. Furthermore, this mechanistic understanding is guiding the de novo design and modification of natural peptides in order to circumvent their limitations (e.g. instability, toxicity, interactions with other drugs, poor kinetics, resistance mechanisms etc) and thus improve their antimicrobial efficacy (Nicola, Albuquerque et al. 2019). Overall, AFPs o□er promising alternatives to standard therapies as anti-infectives and immunomodulatory agents with mechanisms-of-action which are less prone to resistance induction compared to conventional antibiotics (Mahlapuu, Håkansson et al. 2016).

PAF26 is a synthetic hexapeptide that has been shown to be highly effective at killing filamentous fungi whilst showing low toxicity against human and bacterial cells (Munoz, Lopez-Garcia et al. 2006). Unlike membrane permeabilising antimicrobials, initial investigations into the mode-of-action of PAF26 indicated that it did not directly permeabilise the plasma membrane. Instead, at low fungicidal concentrations, PAF26 has a dynamic antifungal mechanism-of-action that involves at least three stages: peptide interaction with the fungal cell envelope (cell wall and/or plasma membrane), endocytic internalisation and accumulation in the vacuole followed by vacuolar expansion. At a certain point, PAF26 is actively transported out of the vacuole into the cytoplasm, followed by a series of complex and specific intracellular effects whose relationship with the eventual death of the target fungus is still unclear. Two functional and separate motifs (cationic and hydrophobic domains) in the peptide amino acid sequence have been identified as playing important roles in the antimicrobial mode-of-action. As a result of these studies, PAF26 has been

proposed as a simple peptide model for the characterization and study of cationic, cell-
penetrating antifungal peptides (Muñoz, Marcos et al. 2012, Muñoz, Gandía et al. 2013).

From average measurements of fungal cell populations expressing the genetically
encoded Ca2+-reporter aequorin, cytosolic free Ca2+ concentrations ([Ca2+]c) within
Neurospora spore germlings treated with a low inhibitory concentration of PAF26 (2.5-5 µM)
have been shown to exhibit a biphasic increase in response to PAF26, and this biphasic
[Ca2+]c increase is completely dependent on the PAF26 and extracellular Ca2+
concentrations. The second phase of the biphasic [Ca2+]c increase was found to be energy
dependent because it was blocked by treatment with the metabolic inhibitor, sodium azide
(NaN3). This may link PAF26 internalization with actin-dependent endocytosis. Consistent
with endocytic internalization playing a key role in PAF26 internalization, is the inhibition by
the F-actin inhibitor, Latrucunulin A and a reduced uptake rate in endocytosis by the
endocytic mutants Δrvs-167, Δrvs-161 and Δrab-5 (Muñoz, Marcos et al. 2012, Muñoz,
Gandía et al. 2013). At high fungicidal concentrations (15 µM), PAF26 killed cells even after
NaN3 treatment, although in a different manner. Under these conditions, the peptide first
bound to the cell envelope as before, but was then observed to directly enter the cytoplasm,
indicating passive transport across the plasma membrane at high concentrations.

In the current research, evidence for Ca2+-signalling having a significant role in the
PAF26 mode-of-action was analysed by testing and analysing the PAF26 sensitivity of
deletion mutants defective in different components of their Ca2+-signalling machinery,
vacuolar fusion and possible programmed cell death. The pattern and kinetics of peptide
interaction, internalization and distribution within the cells of PAF26-resistant mutants were
compared with the parental wild type by using fluorescently labelled PAF26 combined with
live-cell imaging. The role of external Ca2+ in these processes and vacuolar fusion was
assessed.

RESULTS

Sensitivity of fungal cells to killing by PAF26 is dependent on extracellular Ca^{2+}

To confirm that extracellular Ca^{2+} plays a role in determining the sensitivity of *N.*
crassa to PAF26 (Muñoz, Marcos et al. 2012), and thus possibly Ca^{2+} homeostasis being
involved in the mode-of-action of PAF26, the wild type strain was grown in: (1) standard VM
medium (which contains 0.64 mM Ca^{2+}); (2) Ca^{2+} free VM medium (in which Ca^{2+} had been
omitted from the recipe); and (3) VM medium with twice the normal concentration of Ca^{2+}
(1.28 mM). The IC_{50} values for the wild type under these different media were: standard VM

= 3.7 ± 0.6 µM; VM lacking Ca^{2+} = 6.4 ± 0.9 µM which represents a significant increase in
resistance against PAF26 ($P < 0.001$) compared with that in standard VM; VM with 2 X Ca^{2+}
= 2.5 ± 0.4 µM which represented a significant increase in sensitivity to PAF26 ($P < 0.05$)
compared with standard VM (Fig. 1A). These results are consistent with a Ca^{2+}-dependent
mechanism playing a role in determining the sensitivity of *N. crassa* to PAF26 involving
uptake of Ca^{2+} from the external medium.

PAF26 requires a functional $[Ca^{2+}]_{cyt}$ homeostatic machinery in order to kill fungal cells

To assess whether Ca^{2+}-homeostasis and signalling play a role in the mode-of-action
of PAF26, 24 homokaryotic deletion mutants of genes encoding proteins predicted to be
components of the *N. crassa* Ca^{2+}-homeostatic machinery were screened for increased
PAF26 tolerance / sensitivity in comparison with the wild type. The predicted locations of
these proteins in the fungal cell are shown in Fig. 1C; The results of the screen are in Table
1.

The first group of mutants screened were defective in the following Ca^{2+}-channel
proteins: CCH-1 and MID-1, which are components of the high affinity Ca^{2+}-influx system
(HACS); FIG-1, which forms the low-affinity Ca^{2+}-influx system (LACS); and, the vacuolar
Ca^{2+} channel, YVC-1 Two of these mutants (∆*cch-1* and ∆*yvc-1*) exhibited significantly
increased tolerance against PAF26 ($P < 0.001$) when compared with the wild type (Fig. 1B).
The IC_{50} values of ∆*cch-1* and ∆*yvc-1* were both above the level (~ 10 µM) at which PAF26
directly permeabilizes the plasma membrane of *Neurospora* wild type cells (Muñoz, Marcos
et al. 2012).

N. crassa has eight Ca^{2+} ATPases (Zelter *et al.*, 2004), for which seven deletion
mutants were available as homokaryons for screening: ∆*nca-1*, ∆*nca-2*, ∆*nca-3*, ∆*pmr-1*,
∆*ph-7*, ∆*cta-3* and ∆*cta-4* (Fig 1C). These seven mutants exhibited significantly increased
tolerance against PAF26 ($P < 0.001$) when compared with the wild type (Fig. 1B), and *∆nca-
2* and ∆*cta-4* stood out over the rest with *∆nca-2* again approaching inhibitory levels of
PAF26 that directly permeabilise the membrane in the wild type.

Homokaryotic deletion mutants of six of the eight predicted *N. crassa* antiporters
involved in Ca^{2+} transport (Zelter *et al.*, 2004) were available for screening: ∆*cax*, ∆*trm-14*,
∆*trm-15,* ∆*trm-16*, ∆*trm-18*, and ∆*trm-20* (Table 1). Four of the mutants showed significantly
greater tolerance ($P < 0.001$) against PAF26 compared with the wild type although this was
less than a two-fold increase (Fig. 1B).

Fig. 1. Ca^{2+} plays a significant role in the mode-of-action of PAF26. (A) shows the effects of both the removal and increase of free Ca2+ in the media on the IC50 of PAF26 by fungal cells. Increasing the level lowers the concentration at which PAF26 is effective and removing Ca2+ increases the concentration. Four of the mutants (highlighted in bold in Table 1) showed a significantly greater resistance (P < 0.001) against PAF26 compared with the wild type although this was less than a two-fold increase in resistance (Fig. 1B). (B) shows the effects of deleting components of the Ca2+ homeostasis machinery on susceptibility to PAF26. The predicted localisations of these is shown in (C) based on the localization of their protein orthologs in S. cerevisiae yeast cells and vegetative hyphae of N. crassa (Wada, Ohsumi et al. 1987, Yang and Sachs 1989, Bertl and Slayman 1990, Diamond, Zasloff et al. 1991, Antebi and Fink 1992, Iida, Nakamura et al. 1994, Lapinskas, Cunningham et al. 1995, Levina, Lew et al. 1995, Paidhungat and Garrett 1997, Erdman, Lin et al. 1998, Kanzaki 1999, Benito, Garciadeblas et al. 2000, Locke, Bonilla et al. 2000, Muller, Locke et al. 2001, Palmer, Zhou et al. 2001, Courchesne 2002, Courchesne and Ozturk 2002, Denis and Cyert 2002, Gupta, Ton et al. 2003, Kaiserer, Oberparleiter et al. 2003, Muller, Mackin et al. 2003, Zhou, Batiza et al. 2003, Zelter, Bencina et al. 2004, Brand, Shanks et al. 2007, Hallen and Trail 2008, Lew, Abbas et al. 2008, Benito, Garciadeblás et al. 2009, Bormann and Tudzynski 2009, Bowman, Draskovic et al. 2009, Binder, Chu et al. 2010, Bowman, Abreu et al. 2011, Cavinder, Hamam et al. 2011). ** indicates significantly different from wild type at P < 0.001. * indicates significantly different from wildtype at P < 0.05. n = 6. Error bars show standard deviations.

175 **Table 1: The half maximal inhibitory concentration (IC50) of PAF26 in Ca2+**
176 **signalling/homeostasis mutants.**

	Strain	Genotype	IC$_{50}$ PAF 26 µM	±	Source
	WT	74-OR23-1V	3.76	0.19	FGSC2489
Δcch-1	**NCU02762**		**13.94**	**1.26**	**Chu, 2013**
Δmid-1	NCU06703		5.74	1.33	FGSC11708
Δyvc-1	**NCU07605**		**11.37**	**1.42**	**FGSC11253**
Δfig-1	NCU02219		5.89	0.74	FGSC17273
Δnca-1	NCU03305		5.96	1.07	FGSC13287
Δnca-2	**NCU04736**		**12.73**	**2.37**	**FGSC13071**
Δnca-3	NCU05154		5.12	0.42	FGSC13037
Δpmr-1	NCU03292		4.36	0.24	FGSC11616
Δph-7	NCU08147		5.08	0.59	FGSC11256
Δcta-3	NCU07966		9.10	1.68	FGSC11409
Δcta-4	NCU04898		8.22	0.59	FGSC13040
Δena-5	NCU10143		3.27	0.29	FGSC12645
Δcax	NCU07075		4.11	0.50	FGSC11249
Δtrm-15	NCU00916		5.27	0.62	FGSC11686
Δtrm-14	NCU00795		6.19	0.38	FGSC12375
Δtrm-18	NCU06366		4.29	0.56	FGSC11408
Δtrm-16	NCU02826		6.06	1.00	FGSC11529
Δtrm-20	NCU08490		6.64	1.23	FGSC12468
Δplc-1	NCU06245		3.27	0.46	FGSC11411
Δplc-2	NCU01266		6.24	1.60	FGSC12022
Δplc-3	NCU02175		3.59	0.46	FGSC12023
Δplc-4	NCU09655		4.17	0.30	FGSC11271
ΔCamK-1	NCU09123		6.99	0.26	FGSC12547
ΔCamK-2	NCU02283		5.09	0.36	FGSC12449
ΔCamK-3	NCU06177		3.84	0.39	FGSC11536

(Row label at left of table, rotated: Ca^{2+} signalling / homeostasis)

PAF26 is taken up by wild type macroconidia into vacuoles which fuse and then lyse before releasing the peptide into the cytoplasm

In order to compare the dynamic pattern of peptide-cell interactions between the wild
type mutants, PAF26 labelled with the fluorophore TAMRA (TMR-PAF26) was imaged.
Fluorescent labelling has been previously employed to monitor, using confocal live-cell
imaging, the dynamic localization of PAF26 or related peptides in a range of fungi and we
are confident that the dynamic pattern of TMR-PAF26 staining provides a faithful localization
of the unlabelled peptide in fungal cells (Muñoz, Marcos et al. 2012, Muñoz, Gandía et al.
2013, Muñoz, Harries et al. 2013). Previously it has been shown that wild type macroconidia
treated with a low fungicidal concentration of FITC-PAF26 (2.5 µM) exhibit a similar time-
dependent staining pattern as germlings treated with the same concentration of PAF26
(Muñoz, Marcos et al. 2012).

A similar localization pattern of TMR-PAF26 interaction, internalization and
distribution was observed with ungerminated macroconidia to that of FITC-PAF26 (Fig. 2A).
The use of macroconidia allowed a basic monitoring of cell volume, surface area and
developmental state in relation to possible variations in morphogenesis and staining
patterns. Although higher resolution can be achieved with confocal microscopy, widefield
fluorescence microscopy was predominately used for time lapse imaging because of
reduced problems with laser beam irradiation.

Fig. 2. Graphical representation showing that the interaction between PAF26 and
macroconidia is altered through deletion of Ca^{2+}-channels and pumps. **A** shows maximum
intensity projections from a Z series captured using confocal microscopy. Initially, the peptide
appears at the cell envelope and within small vesicles **(a).** The peptide is then accumulated in
larger vesicles and vacuoles **(b)** until there appears a lower number of larger vacuoles **(c)** which
will eventually fuse into a single vacuole which then releases the peptide into the cytoplasm **(d)**.
Scale bar is 10 µm. **B** shows quantification of this interaction over time for the deletion mutants
compared to the wild type ; there are marked changes in the localisation of the peptide during the
time course. **C** shows the interaction between PAF26 and macroconidia is Ca^{2+}-dependent and
energy-independent. This is demonstrated by the significant delay in the uptake of PAF26 in the
wild type when extracellular free Ca^{2+} is removed using BAPTA, at top. The peptide remains
trapped in small vesicles and there is little transport into the vacuolar system and consequently a
reduction in cell death. The lower figure shows the effect of the removal of cellular free energy on
the *Δcch-1* mutant using the metabolic inhibitor NaN$_3$. There is no uptake of PAF26 into the cell
and therefore the direct translocation across the plasma membrane must be an energy
dependent process. N = 100 macroconidia minimum per field of view, three fields of view per
timepoint, repeated at least twice.

The dynamic pattern of TMR-PAF26 in Ca^{2+} homeostasis mutant staining provides insights into the localization and roles of the defective mutant proteins

In order to gain more insight into which stages of the PAF26 interaction, uptake and

distribution process the CCH-1, YVC-1 and NCA-2 proteins influence, the time-dependent

staining by TMR-PAF26 in each of the *Δcch-1, Δyvc-1,* and *Δnca-2* mutants were compared

with that in the wild type. As a baseline for the comparison, four staining patterns were

identified in the wild type macroconidia that could be readily quantified at 30 min intervals

following treatment with 3.5 µM TMR-PAF26. These staining patterns related to different

events in the time-dependent uptake and distribution to different organelles culminating in

cell death. They were: (a) Accumulation in multiple small vesicles. These were predicted to

be mostly endosomes typically up to ~ 1 µm in width. (b) Accumulation in 1-3 large

vacuole(s) typically greater than ~ 1.5 µm in width. (c) Accumulation in the cytoplasm and

excluded from vacuoles. This is the stage just after the peptide has been released out from

vacuoles. (d) Accumulation in both the cytoplasm and vacuoles. This is the stage after which

the intracellular membranes have become permeabilized and thus represents cell death, as

previously reported (Muñoz, Marcos et al. 2012).

In the wild type macroconidia (Figs. 2B and 3A), this quantitative analysis showed that after

treatment for 30 min with TMR-PAF26, all the macroconidia had interacted with TMR-PAF26

and many were beginning to internalise the peptide into small vesicles. After 60 min, many

of the conidia had already undergone vacuolar expansion and some had exported the

peptide into the cytoplasm, and this correlated with a transient reduction in the percentage of

cells with peptide only localized within small vesicles. After 60 min, 40% of the cells had

actively transported the peptide out of their vacuoles and into the surrounding cytoplasm. In

another 40% of the cells at the 60 min time point, vacuolar membrane permeabilization had

occurred and the peptide had equilibrated throughout the cells, which were thus dead. By

120 min, 57% of the cells were dead (Figs. 2B and 3A).

Fig. 3. The interaction with PAF26 is altered through deletion of Ca2+-channels and -pumps as shown by visualisation of wild type and mutant conidia stained with 3.5 μM TAMRA labelled PAF26. These figures show maximum intensity projections of Z stacks false coloured with the 'physics' lookup table to highlight the different fluorescent intensities measured. The wild type conidia undergo a sequence of events which ultimately results in cell death. (4A) (also see 5A) and (4B) (also see 5A) This process is reversed in the Δcch-1 mutant, where the peptide first appears within the cytoplasm and is pumped into small vesicles which then appear to fragment. No conidia were seen to completely lose loaded peptide. (4C) (also see 5A) The Δyvc-1 mutant retains the peptide in small vesicles, without the formation of a large central vacuole, as in the wild type. (4D) (also see 5A) The Δnca-2 mutant appears much reduced in peptide interaction, with very few conidia showing the peptide bound to the envelope. The internalisation process is Ca2+ dependent and energy-dependent, as chelation of Ca2+ with BAPTA. (4E) (also see 6), and treatment with azide (NaN3) (4F) (also see 6), significantly reduced uptake of TAMRA-PAF26. Scale bar is 20 μm.

YVC-1 is required for transport of PAF26 from vesicles to vacuoles

When the Δ*yvc-1* mutant was treated with the labelled peptide (Fig. 2B, 3B), it rapidly appeared at the cell envelope. Unlike with the wild type, the peptide was internalised into small vesicles at a more-or-less linear rate with much remaining trapped at the cell surface. Accumulation of the peptide in vacuoles was much slower and reached a maximum of 6% of the cells with this localization pattern after 120 min. Virtually no cells were observed with peptide exclusively in the cytoplasm at the three time points of measurement over the 120 min by which time only 5% of cells were clearly dead. After 120 min, 79% of the Δ*yvc-1* cells had taken up TMR-PAF26 compared with 100% in the wild type. Thus, less TMR-PAF26 was taken up by Δ*yvc-1* cells, most accumulated in small vesicles and few cells had been killed after 120 min (5% compared with 57% in the wild type). These results are consistent with YVC-1 being required for the transport of PAF26 from vesicles to vacuoles.

NCA-2 is required for the interaction of PAF26 and the cell envelope

When the Δ*nca-2* mutant was treated with TMR-PAF26 (Fig. 2B and 3C) the cell envelopes of the macroconidia were visibly less fluorescent than that of the wild type. Thus the affinity of TMR-PAF26 for the cell envelope of this mutant seemed to be reduced and its staining was delayed compared with that of the wild type. The Δ*nca-2* macroconidia also internalized TMR-PAF26 at a far reduced rate compared with the other mutants and the peptide mostly became trapped within the small vesicles, resulting in virtually no TMR-PAF26 being taken up by vacuoles and a correspondingly extremely small percentage (~ 2%) of the cells being killed after 120 min (Fig. 2B).

CCH-1 plays a key role in the energy- and Ca^{2+}-dependent internalisation of PAF26 by fungal cells

The pattern of TMR-PAF26 internalization and intracellular transport within macroconidia of the Δ*cch-1* mutant appeared to be opposite to that in the wild type (Fig. 3D). The cell envelope was not obviously labelled by the fluorescent peptide in the mutant even after being incubated for 120 min with it (Fig. 2B). After treatment for 30 min the peptide was

primarily localized in the cytoplasm (in 67% of cells) and to a much lesser extent in small
vesicles (~ 18% of cells) and to a very low level in large vacuoles (~ 5% of cells). Between
30 and 90 min, TMR-PAF26 was removed from the cytoplasm and increased in amount in
the small vesicles and large vacuoles. Between 90 and 120 min the large vacuoles
containing TMR-PAF26 are fragmenting into smaller vesicles. This was reflected by a
dramatic increase in the number of stained small vesicles (Fig. 2B).

The overall rate of internalisation of TMR-PAF26 by Δcch-1 macroconidia was faster
than in the wild type because in the former it was initially taken up directly into the cytoplasm
whilst in the latter it first appeared intracellularly in small vesicles that are presumed to be
mostly endosomes. It seems unlikely that TMR-PAF26 is taken up by means of non-specific
permeabilization of the plasma membrane because this would have resulted in more rapid
cell death than in the wild type. Furthermore, 30 min after the addition of TMR-PAF26 it was
clear that the vacuolar membrane had not become permeabilized because the peptide was
excluded from the vacuoles of cells in which the cytoplasm was fluorescent. However,
between 60 and 90 min TMR-PAF26 was transported into the cytoplasm.

In order to clarify whether the passage of PAF26 across the plasma membrane into
the cytoplasm of the Δcch-1 mutant was a result of passive translocation or active uptake,
macroconidia were pre-treated with the metabolic inhibitor NaN_3 (Muñoz, Marcos et al. 2012)
at a concentration of 5 μM for 15 min before the addition of TMR-PAF26 and subsequent
imaging and quantification of localization over 120 min (Figs. 2C and 3F).

In the presence of NaN_3, over the whole 120 min period of incubation with TMR-
PAF26, the peptide remained bound to the cell envelope and was not internalized by most of
the macroconidia. Clearly the metabolic inhibitor NaN_3 had almost completely abolished the
movement of TMR-PAF26 across the plasma membrane indicating that the uptake of the
peptide into these cells of the Δcch-1 mutant is an ATP-dependent process. The results also
showed that the rate of cell death, whilst faster over the first 30 min, remained similar to that
of the non-azide treated Δcch-1 cells; after 120 min ~ 25% of the cells had been killed.
These results are consistent with the uptake of TMR-PAF26 by this subpopulation of Δcch-1
macroconidia being energy independent as a result of a passive process. The kinetics of the
passive uptake of TMR-PAF26 into the cytoplasm and concomitant cell death in the Δcch-1
macroconidia pre-treated with NaN_3 was very similar to the kinetics of the WT macroconidia
similarly pre-treated with NaN_3 over the same time period. These results are consistent with
the normal endocytic internalization of PAF26 being dependent on the Ca^{2+} channel protein,
CCH-1. As CCH-1 appears to initiate PAF26 internalization by endocytosis (see previous
section), this suggests that the uptake of Ca^{2+} from the external medium may be mediated

by this Ca^{2+} channel, which has previously been reported to be immunolocalized to the

plasma membrane (Locke *et al.*, 2000). Furthermore, we had previously shown that removal

of Ca^{2+} from VM made macroconidia more resistant to being killed by PAF26. To attempt to

mimic the effect of *cch-1* deletion in the wild type, Ca^{2+} was removed from the external

medium by the addition of the Ca^{2+} chelator BAPTA (5 mM) 30 min prior to the TMR-PAF26

treatment (Figs. 2C and 3E). Rather than preventing the entry of the peptide, the BAPTA

treatment had the unexpected effect of trapping the peptide in small vesicles. Over the 120

min period very little TMR-PAF26 had localized in large vacuoles, in the cytoplasm and only

~ 11% of the macroconidia were killed. The phenotype of the $\Delta cch\text{-}1$ mutant is consistent

with the CCH-1 protein playing a key role in the energy- and Ca^{2+}-dependent internalisation

of PAF26 by fungal cells.

328 The $[Ca^{2+}]_{cyt}$ response during PAF26 treatment is disrupted in the 329 $[Ca^{2+}]_{ext}$ tolerant mutants

The wild type had previously been shown to undergo a dose dependent biphasic rise

in $[Ca^{2+}]_{cyt}$ upon addition of PAF26 at concentrations between 0.8 and 2.0 µM (Muñoz,

Marcos et al. 2012). The experiment was repeated here but with over a concentration range

of 1.25-10 µM PAF26. For all measurements, the unstimulated $[Ca^{2+}]_{cyt}$ resting level was

measured for 50 sec, in the wild type this was calculated to be 0.05 ± 0.02 µM (Fig. 4A),

before the peptide was added at time 0. After treatment with a low final concentration of

PAF26 (1.25 µM), an immediate increase in $[Ca^{2+}]_{cyt}$ to 0.19 ± 0.01 µM was observed and

this was sustained throughout the measurement period (1,200 sec) and slightly increased

towards the end of this period (Fig. 4B). When the PAF26 added was at a final concentration

of 2.5 µM, which is close to its IC_{50} value, the initial increase in $[Ca^{2+}]_{cyt}$ was slightly greater

(to 0.22 ± 0.03 µM) (Fig. 4B). Again, this was followed by a period of sustained $[Ca^{2+}]_{cyt}$

increase, but there was also a more pronounced exponential increase in $[Ca^{2+}]_{cyt}$ which

began at ~ 900 sec after treatment. With the higher dose, there was also an increase in the

standard deviations of the measurements with time. This is due to aequorin consumption

during the course of the experiment, resulting in less sensitivity after prolonged exposure to

high $[Ca^{2+}]$ (note that luminescence measurements from 6 wells are averaged per time

point). When the wild type was treated with 5 µM PAF26, which was above its IC_{50} value, the

$[Ca^{2+}]_{cyt}$ response followed the same general pattern but with a much larger initial $[Ca^{2+}]_{cyt}$

increase to 0.32 ± 0.02 µM and a shorter period of sustained increase before the $[Ca^{2+}]_{cyt}$

increase became exponential at ~ 430 sec (Fig. 4B). Furthermore, after 590 sec following

peptide treatment, $[Ca^{2+}]_{cyt}$ spiking was observed. The $[Ca^{2+}]_{cyt}$ response was further

accentuated after treatment with a very high dose (10 µM) of PAF26 resulting in a rapid

initial rise in $[Ca^{2+}]_{cyt}$ to 0.54 ± 0.03 µM, followed by an immediate exponential increase in

$[Ca^{2+}]_{cyt}$. This in turn was followed by the $[Ca^{2+}]_{cyt}$ spiking and then a sudden drop in $[Ca^{2+}]_{cyt}$

after ~ 220 sec, which then exponentially increased and subsequently underwent spiking

again (Fig. 4B).

The unstimulated resting level of $[Ca^{2+}]_{cyt}$ in the Δyvc-1 mutant was measured as 0.02

± 0.01 µM (Fig 4A), significantly different at P >0.01. The $[Ca^{2+}]_{cyt}$ response of the Δyvc-1

strain to PAF26 was markedly different to that of the wild type. Whilst there was a similar

initial immediate, dose dependent increase in $[Ca^{2+}]_{cyt}$, there was no second exponential

increase in $[Ca^{2+}]_{cyt}$, which was particularly evident after treating the wild type with PAF26 at

2.5 µM or above (Fig. 4B). Instead the Δyvc-1 mutant underwent a slight reduction in $[Ca^{2+}]_{cyt}$

followed by a sustained, more-or-less constant $[Ca^{2+}]_{cyt}$ level that was significantly raised

compared with the unstimulated resting levels recorded prior to treatment. Thus the Δyvc-1

mutant lacked the second $[Ca^{2+}]_{cyt}$ increase of the biphasic $[Ca^{2+}]_{cyt}$ response of the wild

type, which is consistent with this second $[Ca^{2+}]_{cyt}$ increase resulting from the release of Ca^{2+}

from the vacuole. The resting level of $[Ca^{2+}]_{cyt}$ in the Δnca-2 mutant was measured at 0.05

µM ± 0.02, Fig 4A, and was not significantly different from the wild type. In general terms,

the Δnca-2 mutant showed a similar response to PAF26 as did the Δyvc-1 mutant. However,

the Ca^{2+} signatures of the Δnca-2 and Δyvc-1 mutants were clearly but not dramatically

different from each other; there was a more pronounced dose dependent $[Ca^{2+}]_{cyt}$ increase

in the Δnca-2 strain compared with that in Δyvc-1 in response to PAF26 (Fig. 4B). The

resting level of $[Ca^{2+}]_{cyt}$ in the Δcch-1 mutant was 0.03 ± 0.00 µM Fig. 4A, significantly

different to the wild type at P > 0.01. Interestingly, deletion of cch-1 does not inhibit the

increase in $[Ca^{2+}]_{cyt}$ similar to PAF26 addition, surprising given the lack of evidence for

another Ca^{2+} channel at the cell surface. The Ca^{2+} signatures of the Δcch-1 mutant in

response to different concentrations of PAF26 were broadly similar to those of the Δyvc-1

and Δnca-2 mutants (Figs. 4B). Thus the Δcch-1 mutant also showed an initial $[Ca^{2+}]_{cyt}$

increase but lacked the second $[Ca^{2+}]_{cyt}$ increase of the typical biphasic response of the wild

type to PAF26. These results are consistent with an unidentified L-type channel within the

plasma membrane, which is not CCH-1, being responsible for the initial $[Ca^{2+}]_{cyt}$ increase in

response to PAF26.

The influence on the $[Ca^{2+}]_{cyt}$ response to 5.0 µM PAF26 of the wild type following

pre-treatment with 5 mM BAPTA (to chelate extracellular Ca2+) or with the L-type Ca2+

channel blockers diltiazem or verapamil (both at 5 mM), was analysed (Fig. 4C). A 15 min

pre-treatment with ddH_2O (the control) caused a biphasic increase in $[Ca^{2+}]_{cyt}$ whilst all of the

other treatments prevented any significant $[Ca^{2+}]_{cyt}$ increase (Fig. 2). These results are

consistent with the biphasic $[Ca^{2+}]_{cyt}$ increase being dependent on extracellular Ca2+ as

reported before (Muñoz, Marcos et al. 2012) and on L-type Ca2+ channel activity. The only

known L-type Ca2+ channel in *N. crassa* is CCH-1(Zelter, Bencina et al. 2004). These
results also suggest that the second component of the biphasic $[Ca^{2+}]_{cyt}$ increase is
dependent on the first $[Ca^{2+}]_{cyt}$ increase.

Imaging using the fluorescent reporter GCaMP6 shows PAF26 does not cause a dose dependent rise in $[Ca^{2+}]_{cyt}$

In order to investigate whether PAF26 was indeed causing a dose dependent rise in $[Ca^{2+}]_{cyt}$, conidia expressing GCaMP6s were imaged during PAF26 treatment using widefield fluorescence microscopy. Individual macroconidia were then isolated from the background in FIJI and fluorescence measured over time. As Figure 5A shows, the individual responses are far more varied than the measurements using aequorin suggested. In the middle of the timecourse, the moment the PAF26 reaches the field of view and interacts with the cells can be clearly seen in a sudden increase in Ca^{2+} spiking in all the individuals. Given that the aequorin measurements are across a whole population, it is not hard to see how the cumulative effects of an increase in Ca^{2+} spikes and waves could be interpreted as a dose dependent rise in $[Ca^{2+}]_{cyt}$.In order to determine whether there was a predictable response to PAF26 addition, several repeated experiments were run in which the peptide was added and fluorescence intensity recorded. No distinct pattern or regularity was found within any of the data sets, except for an increase in Ca^{2+} signalling after PAF26 addition. As the only noticeable response, the amplitude and frequency of these Ca^{2+} spikes was quantified. Conidia were treated either with ddH_2O or ddH_2O containing 3.5 µM PAF26. A second set were pre-treated with 5 mM BAPTA before ddH_2O or PAF26 treatment. The conidia treated with ddH_2O have a relatively low frequency of Ca^{2+} spiking with the mean being 4 spikes over the course of the measurement. This is consistent with our findings as to the rate of Ca^{2+} spiking in germinating and fusing conidia (Read, unpublished). When the cells are treated with PAF26 however, there is a marked increase in both frequency and amplitude. The mean frequency of the Ca^{2+} spikes increases to 19 and the amplitude increases to around 6 times the resting level from 4 (Figure 5B). Both of these results are significant at $P < 0.01$. When the conidia are pre-treated with BAPTA however, all Ca^{2+} spiking ceases completely. Whilst there were no apparent recurring patterns in the spiking events, observations appeared to show that they correspond to vesicles and vacuoles coming into close proximity. A Ca^{2+} trace is shown in Fig. 5C and images are shown in Fig. 5D. The images shown in Fig. 5D are taken every five seconds from the timeframes indicated by the black arrows in Fig. 5C. The first Ca^{2+} spike marked by *, appears to correspond with the meeting of the two vacuoles at the lower part of the image, whether they fuse or not is unclear. In the images corresponding to the second spike marked with *, the fusion of two vacuoles is clear at the top of the image.

Figure 5. PAF26 does not cause a dose dependent rise in [Ca^{2+}]$_{cyt}$. (5A) shows the relative fluorescence intensity of several macroconidia expressing GCaMP6s during PAf26 treatment. Fluorescence values are normalised and given as arbitrary values. After the addition of PAF26 there is a marked increase in the amplitude and frequency of Ca2+ spiking. (5B) Quantification of this revealed an actual increase in spiking from an average of 4 spikes per 30 min (left hand panel) to 19 spikes per 30 min (right hand panel). The spiking was entirely dependent on the presence of external Ca2+ as chelation with BAPTA reduced the frequency of spiking to 0. The amplitude of the individual spikes also increased from 4 to 6 arbitrary units. (5C) these spikes often appeared to correspond to the meeting of vesicles and vacuoles within the cell; the images captured from this trace are shown in D which shows images captured every 5 s. The two spikes marked by * in C appear to correspond to vacuolar fusion events. Scale bar = 5µm.

YVC-1 is localized in vacuoles and CCH-1 is both plasma membrane and vacuolar localized

In *N. crassa*, CCH-1 and YVC-1 have not yet been definitively localised while NCA-2 has previously been localised to the vacuole and plasma membrane in mature hyphae (Bowman, Draskovic et al. 2009). We conducted experiments to localize NCA-2 tagged with RFP in macroconidia of *N. crassa* and the data showed labelling of the vacuoles, but no obvious signal at the plasma membrane (Fig. 6).).

In order to localise the CCH-1 and YVC-1 proteins, plasmids were constructed based on FJ457002 for N-terminal tagging and FJ457006 for C-terminal tagging (Honda and Selker 2009).

Both the C- and N- terminal YVC-1:GFP fusion proteins fluoresced, although the N-
tagged protein showed a stronger more stable fluorescence. Fluorescence mainly appeared
within vacuoles in conidia, germ tubes and mature hyphae (Fig. 6). The CCH-1:GFP fusion
products produced very little fluorescence, both from N and C tagged versions. This
appeared to be localised to vacuoles within conidia and intracellular membranes within older
hyphae. Single punctate spots of GFP fluorescence could however be seen at the plasma
membrane of mature hyphae, Fig.6.

Fig. 7. Localization of CCH-1 and YVC-1 labelled with GFP in macroconidia, germlings and vegetative hyphae. Representative widefield fluorescence images of the different cell types of GFP tagged CCH-1 and YVC-1 are shown. As NCA-2 has been previously localised in hyphae, only macroconida re shown for this RFP expressing strain(Bowman, Draskovic et al. 2009). CCH-1 appears to be localised in the vacuolar system in macroconidia and germlings, but also appears as distinct points in the plasma membrane of mature hyphae. YVC-1 appears to localise to the vacuolar system in all cell types. RFP tagged NCA-2 localises to the vacuole in macroconidia. Bar = 10 μm.

DISCUSSION

Ca^{2+} plays important roles in the mode-of-action of PAF26

In this research, evidence for Ca^{2+} signaling and homeostasis having a significant

role in the PAF26 mode-of-action was obtained initially by testing the PAF26 sensitivity of

conidia grown in the presence or absence of calcium. When the level of calcium in the media

is kept at a minimal level, tolerance increases. Conversely when the level is raised,

tolerance decreases. This was confirmed with deletion mutants defective in different

components of their cellular transport machinery of Ca^{2+}. Deletion of most of the Ca^{2+}

channels significantly increased the concentrations at which PAF26 inhibits fungal growth.

The elimination of the high affinity Ca^{2+} plasma membrane channel CCH-1 and the vacuolar

channel YVC-1 as well as the Ca^{2+} ATPase resulted in inhibitory concentrations of the

peptide approaching the point at which passive membrane permeabilisation occurs. Ca^{2+}

has been shown to be an important factor in the mechanistics of PMAP-23 against *Candida*

albicans, the disruption results in reactive oxygen species (ROS) accumulation which

triggers apoptosis (Kim and Lee 2019). This effect is also seen in the peptide CGA-N9

against *C. tropicalis* (Li, Chen et al. 2019). High levels of cytosolic free Ca^{2+} have also been

shown to confer protection to *C. albicans* against the antimicrobial MUC7 12mer, thought to

be by a resulting change in the cell membrane properties preventing peptide entry (Lis, Liu

et al. 2010). It appears in this case that the elevated levels of cytosolic free Ca^{2+} are a stress

response designed to maximise survival by initiating the endocytosis of PAF26. There are

significant changes to the pattern of peptide cell interaction in the mutants and to the

quantity of peptide the mutants take up. The mutations in the Ca^{2+} distribution machinery

appear to disrupt the internalisation and accumulation of the peptide which in turn increases

tolerance to PAF26. Deletion of *nca-2* appears to influence the binding of PAF26 to the cell

envelope, the *Δnca-2* mutant has been shown to accumulate up to 10 fold more Ca^{2+} than

wild type cells, suggesting that NCA-2 serves to remove Ca^{2+} from the cell (Bowman, Abreu

et al. 2011). We found no significant difference in the resting level of [Ca^{2+}[$_{cyt}$ from the wild

type however under unstimulated conditions. Interestingly however, the *Δnca-2* mutant has a

membrane potential reversed from the wild type due to a lack of cell surface H^{+}ATPase

function (Hamam and Lew 2012). In *S. cerevisiae* the H^{+}ATPase is one of the most

abundant cell surface proteins (Bagnat, Chang et al. 2001) and makes up a significant

amount (up to 10%) of the cell surface in *N. crassa* (Bowman, Blasco et al. 1981). PAF26

has been shown to cause a rapid depolarization of the membrane in wild type cells in an

energy independent manner (Muñoz, Marcos et al. 2012). Given that PAF26 has a net

positive charge, the membrane potential reversed *Δnca-2* should technically not be inhibited

in PAF26 membrane binding from an electrostatic view. It is therefore possible to propose

that PAF26 directly inhibits H^+ATPase action, possibly by direct binding; misfunctioning

H^+ATPase at the plasma membrane is sent to the vacuole for degradation in yeast, a

possible mechanistic for the accumulation of PAF26 (Bagnat, Chang et al. 2001, Liu,

Sitaraman et al. 2006). When the peptide is internalised in the *Δyvc-1* mutant, it remains in

small vesicles with very little sign of the peptide entering the vacuolar system; adding

support for the hypothesis that YVC-1 and a threshold amount of PAF26 is required to

initiate vacuolar fusion. Vacuolar fusion occurs through conformational change of the

docking SNARE proteins, triggered by the release of Ca^{2+} from the vacuole in *S. cerevisiae*

(Bayer, Reese et al. 2003, Merz and Wickner 2004, Coonrod, Graham et al. 2013). In *S.*

cerevisiae, isolated vacuoles are able to catalyse their own fusion through the release of

Ca^{2+} from yvc1p present in the vacuolar membrane (Peters and Mayer 1998). This does not

appear to be the case in *N. crassa* however, as both deletion of YVC-1 and removal of

extracellular Ca^{2+} resulted in the trapping of the peptide in small vesicles. Therefore

extracellular Ca^{2+} is required to initiate the release of Ca^{2+} from the vacuoles and trigger

fusion. This raises questions as to why internal membrane fusion is reliant on external

stimuli. In the *Δcch-1* mutant the peptide was directly translocated across the plasma

membrane into the cytoplasm in an energy dependent manner, before accumulation in

vacuoles, meaning PAF26 does not kill by being present in the cytosolic space. CCH-1 is

therefore required to initiate the endocytic pathway.

Ca^{2+} measurement at the population level revealed that the Ca^{2+} responses to PAF26

were markedly different in the PAF26 tolerant mutants. Whilst all underwent a similar initial

increase in $[Ca^{2+}]_{cyt}$ the second biphasic increase seen in the wildtype was abolished. This

initial increase was independent of CCH-1, as the Δ*cch-1* mutant increased, but was

dependent on external Ca^{2+} and L-type Ca^{2+} channels. The fact that L-type channel blockers

completely stop the $[Ca^{2+}]_{cyt}$ increase as does the chelation of external Ca^{2+} using BAPTA

would suggest that there is another, as yet unknown cell surface Ca^{2+} channel. These

findings mirror those of Binder *et al.,* 2010, who found that deletion of *cch-1* did not prevent

the increase in $[Ca^{2+}]_{cyt}$ in response to PAF. When these measurements were taken at the

level of the individual, rather than the population, it became apparent that PAF26 does not in

fact cause a dose dependent rise in $[Ca^{2+}]_{cyt}$ but rather increases both the amplitude and

frequency of Ca^{2+} spiking, again dependent on the presence of external Ca^{2+}. These findings

have allowed us to propose the model shown in Fig. 8 for the role of Ca^{2+} signalling and

homeostasis in the mode of action of PAF26.

Fig 7. Proposed role of Ca²⁺ signalling in the mode of action of PAF26. 1 initially the cell is unstimulated and maintains a low $[Ca^{2+}]_{cyt}$ and stable membrane potential. **2** PAF26 inhibits the cell surface H⁺ATPase initiating membrane depolarisation. **3** Depolarisation of the plasma membrane activates the voltage gated CCH-1, causing $[Ca^{2+}]_{cyt}$ to rise. **4** This rise in $[Ca^{2+}]_{cyt}$ initiates the turnover of the H⁺ATPase including bound PAF26 by endocytosis. **5** further rises in $[Ca^{2+}]_{cyt}$ caused by external Ca²⁺ entering through CCH-1 initiate Ca²⁺ release through YVC-1 **6** driving the fusion of membranes. **7** PAF26 is released from the H⁺ATPase as it is degraded and accumulates in the vacuolar system.

Materials and methods

Neurospora crassa stocks

N. crassa strains were obtained from the Fungal Genetics Stock Centre (FGSC, University of Missouri, Kansas City, USA), unless otherwise specified. FGSC2489 (*matA*,74-OR23-1VA) was used as the wild type control. *N. crassa* strains generated in this study were derived from FGSC9717 (*matA,Δmus-51::bar⁺; his-3⁻*), FGSC9720 (*matA,Δmus-52::bar⁺; his-3⁻*) and FGSC6103 (*matA,his-3⁻*) *N. crassa* was cultured and maintained according to FGSC protocols (www.fgsc.net/Neurospora/NeurosporaProtocolGuide.htm) unless otherwise stated.

564   For microscopy, conidia were cultured in liquid VM for up to 4 hours at 30°C using 8
565   well Nunc™ Lab-Tek™ II Chambered Coverglass.

N. crassa transformation

567   Electroporation of macroconidia was carried out using the following settings:
568   resistance 600 Ω, voltage 1.5 kV/cm and capacitance 25 µFd.

Heterokaryon purification

570   *N. crassa* strains that were obtained as heterokaryons, were purified to homokaryons
571   by standard procedure (www.fgsc.net/Neurospora/NeurosporaProtocolGuide.htm).

Plasmids

In order to create pTef1AeqS, aequorin was amplified from pAB19 using the primers
AeqS-F and AeqS-R.. The vector pCC019 was digested with PacI and EcoRI and the vector
gel purified. Aequorin was then amplified with AeqS-IF-Fw and AeqS-IF-Rv to add overlaps
homologous to the backbone whilst maintaining digestion sites and the vector was
assembled using Gibson assembly. To generate pYVC1CGFP and pYVC1NGFP, the *yvc-1*
ORF was amplified from gDNA using the primers NC-YVC1-AMP-FW and NC-YVC-1-AMP-
RV. The purified product was then amplified using NC-YVC1-N-FW & NC-YVC1-N-RV for N
terminal tagging and NC-YVC1-C-FW & NC-YVC1-C-RV for C terminal tagging. The
FJ457002 vector was digested using PacI and XbaI and the FJ457006 vector digested with
AscI and NotI. Following gel extraction both vectors were assembled using Gibson
assembly.

Tagging of CCH-1 was not straightforward due to the size of the ORF - 6.5kb. The
CCH-1 ORF was amplified first using the primer sets NC-CCH1 AMP-FW and NC-CCH1-
3.5Rv and NC-CCH1-3.5Fw and NC-CCH1-AMP-RV to amplify the gene in two segments, 5'
and 3'. The complete constructs were built using fusion PCR; primer NC-CCH1-C-FW, 5'
segment, 3' segment and primer NC-CCH1-C-RV, the N terminal tagging vector used the
primers NC-CCH1-N-FW and NC-CCH1-N-RV. Following successful fusion the vectors were
assembled as for YVC-1. Primer sequences are given in supplementary material.

IC$_{50}$ values

IC$_{50}$ values were calculated using clear, U bottom polystyrene microtitre
plates. PAF 26 was prepared at two concentrations, 35 µM and 30 µM, and diluted two-fold
appropriately in to reach the final experimental concentrations . Conidia were diluted to a
concentration of 1 X 10^6 in 20% liquid VM and 50 µl dispensed into each well. Experimental

PAF26   concentrations   (µM)   were:
[17.50][15.00][8.75][7.50][4.38][3.75][2.19][1.88][1.09][0.94][0.55][0].   Final   conidial
concentration was of 5 X 10^5 in 10% VM. Absorbance measurements were obtained at 610
nm. This wavelength is close to the 595nm at which the relationship between optical density
and fungal biomass is linear (Broekaert, Terras et al. 1990).

Calcium measurements

Coelenterazine (#C-7001, www.biosynth.com)was prepared by dissolving in ice cold
methanol (MetOH) under a protective atmosphere in complete darkness to a concentration
of 10 µg/µl. Stocks were diluted to the working concentration of 3.175 µg in 10 µl MetOH,
74.78 µM.

Conidia were diluted to a concentration of 1 X 10^6 in 10% VM, coelenterazine was
added to a final concentration of 2.5 µM. 100 µl of conidial suspension was used per well of
a white microtitre plate and the plates incubated at 25°C in the dark for 6 hours. The light
output of aequorin was measured by counting the photons emitted by a single well over one
second, each well in a row of six being measured once every cycle of 7 seconds. After
recording baseline luminescence, a single row was treated with PAF26 and the light output
measured over time. Following this the second row was run with 100 µl 3M $CaCl_2$:20%
ethanol (EtOH) injected on cycle 8 to immediately discharge all the aequorin.

[Ca^{2+}] calculation

The raw relative light units (RLU) were converted to [Ca^{2+}] using the formula:(Bonora,
Giorgi et al. 2013)

$$[\text{Ca}^{2+}]\text{M} = \left(\frac{\left(\frac{L}{L_{\max}} \times \lambda\right)^{\frac{1}{n}} + \left(\left(\frac{L}{L_{max}} \times \lambda\right)^{\frac{1}{n}} \times K_{TR}\right) - 1}{K_R - \left(\left(\frac{L}{L_{max}} \times \lambda\right)^{\frac{1}{n}} \times K_R\right)} \right)$$

Where:
•L is the light intensity at sampling time
•L_{max} is the total light emitted at sampling time
•K_R is the constant for calcium bound state **(7,230,000)**
•K_{TR} is the constant for calcium unbound state **(120)**
•λ is the rate constant for aequorin consumption at saturating [Ca^{2+}] **(1)**
•n is the number of Ca^{2+} binding sites **(2.99)**

In order to calculate L_{max} the total RLU (RLU_{total}) available was calculated from the discharge

of aequorin (RLUd) using the trapezoid rule formula, where the total RLUd available at time

point t_n equals:

$$RLUd_{t_{n-1}} + \frac{(t_n - t_{n-1}) \times (RLUd_{t_n} + RLUd_{t_{n-1}})}{2}$$

627 and $RLUd_{total}$ is the final value reached. Switching to the experimental data (RLUe), L_{max} was

628 calculated for each time point as:

$$RLUd_{total} - \left(RLUe_{t_{n-1}} + \left(\frac{(t_n - t_{n-1}) \times \left(RLUe_{t_n} + RLUe_{t_{n-1}} \right)}{2} \right) \right)$$

Peptides and peptide handling

Peptides were ordered from GenScript (www.genscript.com) at > 98 % purity. Stock

solutions were prepared in 50:50 dimethyl sulfoxide (DMSO):H_2O buffer. Working peptide

solutions were prepared in ddH_2O.

Microscopy

All microscopy was carried out on a Nikon Eclipse® TE2000-E inverted microscope,

using a Nikon Plan Apo 60 X 1.2 N.A. DIC H water immersion objective and Nikon G-2A and

B-2A filter sets, excitation was provided by a CoolLED illumination system set at 550 nm for

use with the G-2A filter or 470 nm for use with the B2-A filter.

Confocal microscopy was carried out using a Leica TCS SP8 equipped with two

hybrid GaAsP detectors (HyD) and two photomultiplier tubes (PMT). Excitation was provided

using either the Leica tuneable white light laser (450 -750 nm), an argon laser (458nm,

476nm, 488nm and 496nm) or UV laser (405nm). Images were captured using LAS X

software and the Leica 63 X water immersion objective. All image handling and analysis was

carried out using Fiji (fiji.sc/Fiji).

Quantification of Ca^{2+} signalling

Fluorescence values were exported from Fiji and analysed in Excel. Data was

normalised using feature scaling. This scales the data to removes irregularities in GCaMP6

expression and photon yield. The function used was: $x' = $ MIN + (x-min_x)(MAX-MIN)/(max_x-

min_x). Where MIN and MAX are the required scale range, x is the original value and x' is the

normalised value. min_x is the minimum value for that data set and max_x is the maximum,

max_x was defined as the maximum value from all experiments plus 10 to avoid artificial

amplification of noise. The noise was removed using an IF function: =IF(x>y,x,0) where x is

the data point and y is the noise threshold limit. Finally, an IF(AND(x>x_{-1}, x>x_{+1}),1,0)

function, where x_{-1} and x_{+1} are the surrounding measurements, quantified each peak in

GCamP6   fluorescence

Funding Statement

Funding for this project was provided by a Biotechnology and Biological Sciences Research

Council Research Studentship to AJTA. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

References

Antebi, A. and G. R. Fink (1992). "The yeast Ca(2+)-ATPase homologue, PMR1, is required
for normal Golgi function and localizes in a novel Golgi-like distribution." Molecular Biology of
the Cell **3**(6): 633-654.
Bagnat, M., A. Chang and K. Simons (2001). "Plasma Membrane Proton ATPase Pma1p
Requires Raft Association for Surface Delivery in Yeast." Molecular Biology of the Cell
12(12): 4129-4138.
Bayer, M. J., C. Reese, S. Buhler, C. Peters and A. Mayer (2003). "Vacuole membrane
fusion: V0 functions after trans-SNARE pairing and is coupled to the Ca2+-releasing
channel." The Journal of cell biology **162**(2): 211-222.
Benito, B., B. Garciadeblás, J. Pérez-Martín and A. Rodríguez-Navarro (2009). "Growth at
High pH and Sodium and Potassium Tolerance in Media above the Cytoplasmic pH Depend
on ENA ATPases in *Ustilago maydis*." Eukaryotic Cell **8**(6): 821-829.
Benito, B., B. Garciadeblas and A. Rodriguez-Navarro (2000). "Molecular cloning of the
calcium and sodium ATPases in Neurospora crassa." Molecular Microbiology **35**(5): 1079-
1088.
Bertl, A. and C. L. Slayman (1990). "Cation-selective channels in the vacuolar membrane of
Saccharomyces: dependence on calcium, redox state, and voltage." Proceedings of the
National Academy of Sciences **87**(20): 7824-7828.
Binder, U., M. Chu, N. D. Read and F. Marx (2010). "The Antifungal Activity of the
Penicillium chrysogenum Protein PAF Disrupts Calcium Homeostasis in Neurospora
crassa." Eukaryotic Cell **9**(9): 1374-1382.
Bongomin, F., S. Gago, O. R. Oladele and W. D. Denning (2017). "Global and Multi-National
Prevalence of Fungal Diseases—Estimate Precision." Journal of Fungi **3**(4).
Bonora, M., C. Giorgi, A. Bononi, S. Marchi, S. Patergnani, A. Rimessi, R. Rizzuto and P.
Pinton (2013). "Subcellular calcium measurements in mammalian cells using jellyfish
photoprotein aequorin-based probes." Nature Protocols **8**(11): 2105-2118.
Bormann, J. and P. Tudzynski (2009). "Deletion of Mid1, a putative stretch-activated calcium
channel in Claviceps purpurea, affects vegetative growth, cell wall synthesis and virulence."
Microbiology **155**(12): 3922-3933.
Bowman, B. J., S. Abreu, E. Margolles-Clark, M. Draskovic and E. J. Bowman (2011). "Role
of Four Calcium Transport Proteins, Encoded by *nca-1*, *nca-2*, *nca-3*, and *cax*, in
Maintaining Intracellular Calcium Levels in Neurospora crassa." Eukaryotic Cell **10**(5): 654-
661.
Bowman, B. J., F. Blasco and C. W. Slayman (1981). "Purification and characterization of
the plasma membrane ATPase of Neurospora crassa." J Biol Chem **256**(23): 12343-12349.
Bowman, B. J., M. Draskovic, M. Freitag and E. J. Bowman (2009). "Structure and
Distribution of Organelles and Cellular Location of Calcium Transporters in *Neurospora
crassa*." Eukaryotic Cell **8**(12): 1845-1855.
Brand, A., S. Shanks, V. M. S. Duncan, M. Yang, K. Mackenzie and N. A. R. Gow (2007).
"Hyphal Orientation of Candida albicans Is Regulated by a Calcium-Dependent Mechanism."
Current Biology **17**(4): 347-352.

Broekaert, W. F., F. R. Terras, B. P. Cammue and J. Vanderleyden (1990). "An automated
quantitative assay for fungal growth inhibition." <u>FEMS Microbiology Letters</u> **69**(1-2): 55-59.
Brown, G. D., D. W. Denning, N. A. R. Gow, S. M. Levitz, M. G. Netea and T. C. White
(2012). "Hidden Killers: Human Fungal Infections." <u>Science Translational Medicine</u> **4**(165):
165rv113-165rv113.
Cavinder, B., A. Hamam, R. R. Lew and F. Trail (2011). "Mid1, a Mechanosensitive Calcium
Ion Channel, Affects Growth, Development, and Ascospore Discharge in the Filamentous
Fungus Gibberella zeae." <u>Eukaryotic Cell</u> **10**(6): 832-841.
Coonrod, E. M., L. A. Graham, L. N. Carpp, T. M. Carr, L. Stirrat, K. Bowers, N. J. Bryant
and T. H. Stevens (2013). "Homotypic vacuole fusion in yeast requires organelle acidification
and not the V-ATPase membrane domain." <u>Developmental cell</u> **27**(4): 462-468.
Courchesne, W. E. (2002). "Characterization of a Novel, Broad-Based Fungicidal Activity for
the Antiarrhythmic Drug Amiodarone." <u>Journal of Pharmacology and Experimental</u>
<u>Therapeutics</u> **300**(1): 195-199.
Courchesne, W. E. and S. Ozturk (2002). "Amiodarone induces a caffeine-inhibited, MID1-
depedent rise in free cytoplasmic calcium in Saccharomyces cerevisiae: Amiodarone
induces a rise in cytoplasmic calcium in yeast." <u>Molecular Microbiology</u> **47**(1): 223-234.
Denis, V. and M. S. Cyert (2002). "Internal Ca $^{2+}$ release in yeast is triggered by hypertonic
shock and mediated by a TRP channel homologue." <u>The Journal of Cell Biology</u> **156**(1): 29-
34.
Denning, D. W. and M. J. Bromley (2015). "How to bolster the antifungal pipeline." <u>Science</u>
347(6229): 1414.
Diamond, G., M. Zasloff, H. Eck, M. Brasseur, W. L. Maloy and C. L. Bevins (1991).
"Tracheal antimicrobial peptide, a cysteine-rich peptide from mammalian tracheal mucosa:
peptide isolation and cloning of a cDNA." <u>Proceedings of the National Academy of Sciences</u>
88(9): 3952-3956.
Duncan, V. M. S. and D. A. O'Neil (2013). "Commercialization of antifungal peptides."
<u>Fungal Biology Reviews</u> **26**(4): 156-165.
Erdman, S., L. Lin, M. Malczynski and M. Snyder (1998). "Pheromone-regulated Genes
Required for Yeast Mating Differentiation." <u>The Journal of Cell Biology</u> **140**(3): 461-483.
Gupta, S. S., V.-K. Ton, V. Beaudry, S. Rulli, K. Cunningham and R. Rao (2003). "Antifungal
Activity of Amiodarone Is Mediated by Disruption of Calcium Homeostasis." <u>Journal of</u>
<u>Biological Chemistry</u> **278**(31): 28831-28839.
Hallen, H. E. and F. Trail (2008). "The L-Type Calcium Ion Channel Cch1 Affects Ascospore
Discharge and Mycelial Growth in the Filamentous Fungus *Gibberella zeae* (Anamorph
Fusarium graminearum)." <u>Eukaryotic Cell</u> **7**(2): 415-424.
Hamam, A. and R. R. Lew (2012). "Electrical phenotypes of calcium transport mutant strains
of a filamentous fungus, Neurospora crassa." <u>Eukaryotic cell</u> **11**(5): 694-702.
Honda, S. and E. U. Selker (2009). "Tools for fungal proteomics: multifunctional neurospora
vectors for gene replacement, protein expression and protein purification." <u>Genetics</u> **182**(1):
11-23.
Iida, H., H. Nakamura, T. Ono, M. S. Okumura and Y. Anraku (1994). "MID1, a novel
Saccharomyces cerevisiae gene encoding a plasma membrane protein, is required for Ca2+
influx and mating." <u>Molecular and Cellular Biology</u> **14**(12): 8259-8271.
Kaiserer, L., C. Oberparleiter, R. Weiler-G□rz, W. Burgstaller, E. Leiter and F. Marx (2003).
"Characterization of the Penicillium chrysogenum antifungal protein PAF." <u>Archives of</u>
<u>Microbiology</u> **180**(3): 204-210.
Kanzaki, M. (1999). "Molecular Identification of a Eukaryotic, Stretch-Activated Nonselective
Cation Channel." <u>Science</u> **285**(5429): 882-886.
Kim, S. and D. G. Lee (2019). "Role of calcium in reactive oxygen species-induced
apoptosis in Candida albicans: an antifungal mechanism of antimicrobial peptide, PMAP-23."
<u>Free Radic Res</u> **53**(1): 8-17.
Kohler, J. R., A. Casadevall and J. Perfect (2014). "The spectrum of fungi that infects
humans." <u>Cold Spring Harb Perspect Med</u> **5**(1): a019273.

Lapinskas, P. J., K. W. Cunningham, X. F. Liu, G. R. Fink and V. C. Culotta (1995). "Mutations in PMR1 suppress oxidative damage in yeast cells lacking superoxide dismutase." <u>Molecular and Cellular Biology</u> **15**(3): 1382-1388.

Levina, N. N., R. R. Lew, G. J. Hyde and I. B. Heath (1995). "The roles of Ca2+ and plasma membrane ion channels in hyphal tip growth of Neurospora crassa." <u>J Cell Sci</u> **108 (Pt 11)**: 3405-3417.

Lew, R. R., Z. Abbas, M. I. Anderca and S. J. Free (2008). "Phenotype of a Mechanosensitive Channel Mutant, *mid - 1*, in a Filamentous Fungus, *Neurospora crassa*." <u>Eukaryotic Cell</u> **7**(4): 647-655.

Li, R., C. Chen, B. Zhang, H. Jing, Z. Wang, C. Wu, P. Hao, Y. Kuang and M. Yang (2019). "The chromogranin A-derived antifungal peptide CGA-N9 induces apoptosis in Candida tropicalis." <u>Biochemical Journal</u> **476**(20): 3069-3080.

Lis, M., T. T. Liu, K. S. Barker, P. D. Rogers and L. A. Bobek (2010). "Antimicrobial peptide MUC7 12-mer activates the calcium/calcineurin pathway in Candida albicans." <u>FEMS Yeast Research</u> **10**(5): 579-586.

Liu, Y., S. Sitaraman and A. Chang (2006). "Multiple Degradation Pathways for Misfolded Mutants of the Yeast Plasma Membrane ATPase, PMA1." <u>Journal of Biological Chemistry</u> **281**(42): 31457-31466.

Locke, E. G., M. Bonilla, L. Liang, Y. Takita and K. W. Cunningham (2000). "A Homolog of Voltage-Gated Ca2+ Channels Stimulated by Depletion of Secretory Ca2+ in Yeast." <u>Molecular and Cellular Biology</u> **20**(18): 6686-6694.

Mahlapuu, M., J. Håkansson, L. Ringstad and C. Björn (2016). "Antimicrobial Peptides: An Emerging Category of Therapeutic Agents." <u>Frontiers in Cellular and Infection Microbiology</u> **6**.

Matejuk, A., Q. Leng, M. D. Begum, M. C. Woodle, P. Scaria, S. T. Chou and A. J. Mixson (2010). "Peptide-based Antifungal Therapies against Emerging Infections." <u>Drugs Future</u> **35**(3): 197.

Merz , A. J. and W. T. Wickner (2004). "Trans-SNARE interactions elicit Ca2+ efflux from the yeast vacuole lumen." <u>The Journal of Cell Biology</u> **164**(2): 195-206.

Muller, E. M., E. G. Locke and K. W. Cunningham (2001). "Differential regulation of two Ca(2+) influx systems by pheromone signaling in Saccharomyces cerevisiae." <u>Genetics</u> **159**(4): 1527-1538.

Muller, E. M., N. A. Mackin, S. E. Erdman and K. W. Cunningham (2003). "Fig1p facilitates Ca2+ influx and cell fusion during mating of Saccharomyces cerevisiae." <u>J Biol Chem</u> **278**(40): 38461-38469.

Muñoz, A., M. Gandía, E. Harries, L. Carmona, N. D. Read and J. F. Marcos (2013). "Understanding the mechanism of action of cell-penetrating antifungal peptides using the rationally designed hexapeptide PAF26 as a model." <u>Fungal Biology Reviews</u> **26**(4): 146-155.

Muñoz, A., M. Gandía, E. Harries, L. Carmona López, N. Read and J. F. Marcos (2013). <u>Understanding the mechanism of action of cell-penetrating antifungal peptides using the rationally designed hexapeptide PAF26 as a model</u>.

Muñoz, A., E. Harries, A. Contreras-Valenzuela, L. Carmona, N. D. Read and J. F. Marcos (2013). "Two Functional Motifs Define the Interaction, Internalization and Toxicity of the Cell-Penetrating Antifungal Peptide PAF26 on Fungal Cells." <u>PLOS ONE</u> **8**(1): e54813.

Munoz, A., B. Lopez-Garcia and J. F. Marcos (2006). "Studies on the Mode of Action of the Antifungal Hexapeptide PAF26." <u>Antimicrobial Agents and Chemotherapy</u> **50**(11): 3847-3855.

Muñoz, A., J. F. Marcos and N. D. Read (2012). "Concentration-dependent mechanisms of cell penetration and killing by the de novo designed antifungal hexapeptide PAF26: Uptake mechanisms of the antifungal peptide PAF26." <u>Molecular Microbiology</u> **85**(1): 89-106.

Nicola, A. M., P. Albuquerque, H. C. Paes, L. Fernandes, F. F. Costa, E. S. Kioshima, A. K. R. Abadio, A. L. Bocca and M. S. Felipe (2019). "Antifungal drugs: New insights in research & development." <u>Pharmacol Ther</u> **195**: 21-38.

Paidhungat, M. and S. Garrett (1997). "A homolog of mammalian, voltage-gated calcium
channels mediates yeast pheromone-stimulated Ca2+ uptake and exacerbates the cdc1(Ts)
growth defect." Mol Cell Biol **17**(11): 6339-6347.
Palmer, C. P., X. L. Zhou, J. Lin, S. H. Loukin, C. Kung and Y. Saimi (2001). "A TRP
homolog in Saccharomyces cerevisiae forms an intracellular Ca(2+)-permeable channel in
the yeast vacuolar membrane." Proc Natl Acad Sci U S A **98**(14): 7801-7805.
Peters, C. and A. Mayer (1998). "Ca2+/calmodulin signals the completion of docking and
triggers a late step of vacuole fusion." Nature **396**(6711): 575-580.
Rautenbach, M., A. M. Troskie and J. A. Vosloo (2016). "Antifungal peptides: To be or not to
be membrane active." Biochimie **130**: 132-145.
Wada, Y., Y. Ohsumi, M. Tanifuji, M. Kasai and Y. Anraku (1987). "Vacuolar ion channel of
the yeast, Saccharomyces cerevisiae." J Biol Chem **262**(36): 17260-17263.
Yang, X. and F. Sachs (1989). "Block of stretch-activated ion channels in Xenopus oocytes
by gadolinium and calcium ions." Science **243**(4894): 1068-1071.
Zelter, A., M. Bencina, B. J. Bowman, O. Yarden and N. D. Read (2004). "A comparative
genomic analysis of the calcium signaling machinery in Neurospora crassa, Magnaporthe
grisea, and Saccharomyces cerevisiae." Fungal Genetics and Biology **41**(9): 827-841.
Zhou, X. L., A. F. Batiza, S. H. Loukin, C. P. Palmer, C. Kung and Y. Saimi (2003). "The
transient receptor potential channel on the yeast vacuole is mechanosensitive." Proceedings
of the National Academy of Sciences **100**(12): 7105-7110.

Supplementary

Name	Sequence
NC-cch1-AMP-FW	ATCTATGGCTGTGACGCTAG
NC-cch1-AMP-RV	GTCATCTTCGCATCTCCA
NC-yvc1-AMP-FW	AGCATGGCAAGGTTCGG
NC-yvc1-AMP-RV	TCACTTCCCATCACTACTGCTACTG
NC-cch1-C-FW	aacccctcacatcaaccaaatctagATGAGCGATCATCCAGCAC
NC-cch1-C-RV	ctccgccgcctccgcccttaattaaGTAAGCCCCCGAGTATCCTC
NC-cch1-N-FW	cggaggcggcggaggcggcgcgccaAGCGATCATCCAGCACCT
NC-cch1-N-RV	ggagctccaccgcggtggcggccgcTTAGTAAGCCCCCGAGTATCCT
NC-yvc1-C-FW	aacccctcacatcaaccaaatctagATGGCAAGGTTCGGGGC
NC-yvc1-C-RV	ctccgccgcctccgcccttaattaaCTTCCCATCACTACTGCTACTGCTACTA
NC-yvc1-N-FW	cggaggcggcggaggcggcgcgccaGCAAGGTTCGGGGCCTT
NC-yvc1-N-RV	ggagctccaccgcggtggcggccgcTCACTTCCCATCACTACTGCTACTG
NC-cch1-3.5Rv	TAGAAAGGGTTGGGTTCACG
NC-cch1-3.5Fw	CGTGAACCCAACCCTTTCTA

Primers

Plasmids

Plasmid	Backbone	Genotype	Source
FJ457002		*Pccg-1::c-gly::gfp*	Honda & Selker, 2009
FJ457006		*Pccg-1::n-gfp*	Honda & Selker, 2009
pAB19	pAZ6	*Pccg-1-AeqS-PtrpC-bar$^+$*	Marris, 2007
pCC019	FJ457002	*his-3-Ptef-1-gcamp6s*	Chang, 2015
pTef1AeqS	pCC019	*his-3-Ptef-1-AeqS*	This study
pYVC1CGFP	FJ457002	*his-3-Pccg-1-yvc-1-gfp*	This study
pYVC1NGFP	FJ457006	*his-3-Pccg-1-gfp-yvc-1*	This study
pCCH1CGFP	FJ457002	*his-3-Pccg-1-yvc-1-gfp*	This study
pCCH1NGFP	FJ457006	*his-3-Pccg-1-gfp-yvc-1*	This study