Motor training strengthens corticospinal suppression during movement preparation

Pierre Vassiliadis[1,2]*, Gerard Derosiere[1], Julien Grandjean[1], Julie Duque[1]

[1] Institute of Neuroscience, Université Catholique de Louvain, 1200, Brussels, Belgium.

[2] Center for Neuroprosthetics (CNP) and Brain Mind Institute (BMI), Swiss Federal Institute of Technology (EPFL), 1202, Geneva, Switzerland

Abbreviated title: Training strengthens preparatory suppression

* CORRESPONDENCE TO:
Pierre Vassiliadis
Institute of Neuroscience
Université catholique de Louvain
53, Avenue Mounier
COSY- B1.53.04
B-1200 BRUSSELS
Belgium
Email: pierre.vassiliadis@uclouvain.be
Web: http://www.coactionslab.com

Key Points

- Motor training modifies neural activity at rest and during movement execution. Training may also influence how movements are prepared but the neural correlates and the potential behavioral relevance of such adjustments in movement preparation remain unclear.

- In humans, studies have shown that movement preparation is accompanied by a suppression of corticospinal excitability (CSE), relative to a resting measure, a phenomenon called preparatory suppression.

- Here, we show that training on a reaction time (RT) task strengthens preparatory suppression and that this effect occurs in parallel with an increase in resting CSE.

- Training also accelerated RTs. This RT improvement was related to the deepening in preparatory suppression (but not to changes in resting CSE): subjects showing a stronger expansion of preparatory suppression were also those exhibiting larger gains in RTs.

- These results suggest that training induces changes in motor preparatory processes that are linked to an enhanced ability to initiate fast movements.

Abstract

Training can improve motor skills and modify neural activity at rest and during movement execution. Learning-related modulations may also concern motor preparation but the neural correlates and the potential behavioral relevance of such adjustments remain unclear. In humans, preparatory processes have been largely investigated using transcranial magnetic stimulation (TMS) with several studies reporting decreased corticospinal excitability (CSE) relative to a baseline measure; a phenomenon called preparatory suppression. Here, we investigated the effect of motor training on preparatory suppression of CSE in humans, as compared to modulatory changes at rest. We trained participants to initiate quick movements in an instructed-delay reaction time (RT) task and used TMS to investigate changes in CSE over the practice blocks. Training on the task speeded up RTs, with no repercussion on error rates. Training also increased baseline CSE at rest. Most interestingly, we found that motor activity during action preparation did not mirror the training-related rise in resting CSE. Rather, the degree of preparatory suppression from the rising baseline strengthened with practice. This training-related change in preparatory suppression predicted RT gains (but not the changes in baseline CSE): subjects showing a stronger expansion of preparatory suppression were also those exhibiting larger gains in RTs. Finally, such relationship between RTs and preparatory suppression was also evident at the single-trial level: RTs were generally faster in trials where preparatory suppression was deeper. These findings suggest that training induces changes in motor preparatory processes that are linked to an enhanced ability to initiate fast movements.

64   **Keywords:** motor training; action preparation; transcranial magnetic stimulation; motor-evoked

65   potentials; primary motor cortex; corticospinal excitability; corticospinal suppression, reaction

66   time.

1. Introduction

Motor training improves the speed and/or accuracy at which movements are selected, initiated and executed. Significant research has been devoted to unveiling the functional changes at the basis of such improvements (Krakauer *et al*., 2019). At the neural level, neuroimaging (*e.g.,* Wiestler & Diedrichsen, 2013; Wenger *et al.*, 2017; Yokoi & Diedrichsen, 2019) and transcranial magnetic stimulation (TMS) studies (*e.g.,* Rosenkranz *et al.*, 2007; Reis *et al.*, 2008; Mawase *et al.*, 2017) have shown that training is accompanied by a plastic reorganization of the motor system, supporting the formation of new motor memories. Specifically, training amplifies resting motor activity (*e.g.,* Pascual-Leone *et al.*, 1995; Butefisch *et al.*, 2000; Duque *et al.*, 2008; Galea & Celnik, 2009; Christiansen *et al.*, 2018) and induces learning-specific changes of motor activity during movement execution (Krakauer *et al.*, 2004; Steele & Penhune, 2010; Shmuelof *et al.*, 2014). Animal studies also show learning-related modulations of motor activity during action preparation (Paz *et al.*, 2003; Makino *et al.*, 2017; Vyas *et al.*, 2018, 2020) that could reflect an optimization of preparatory processes with training (Mawase *et al.*, 2018). Yet, the behavioral relevance of the effects of training on action preparation remain unclear.

In humans, the excitability of the motor system can be assessed by applying TMS over primary motor cortex (M1), eliciting motor-evoked potentials (MEPs), whose amplitude reflects the excitability of the corticospinal pathway (Derosiere & Duque, 2020; Derosiere *et al*., 2020). When applied during reaction time (RT) tasks, TMS elicits MEPs that are used to assess corticospinal excitability (CSE) changes associated with action preparation and initiation. CSE is often suppressed during action preparation when compared to a baseline, measured at rest. The function of this preparatory suppression (or inhibition) remains unclear (*e.g.,*Greenhouse *et al.*, 2015; Duque *et al.*, 2017; Derosiere, 2018; Hannah *et al.*, 2018). A prominent view is that it assists action selection processes, by preventing the release of premature or incorrect responses (Duque *et al.*, 2010; Quoilin *et al.*, 2018). Indeed, the amount of preparatory suppression was shown to scale with the complexity of selection processes (Klein *et al.*, 2014; Duque *et al.*, 2016). Another hypothesis is that preparatory suppression eases action initiation (Greenhouse *et al.*, 2015; Hasegawa *et al.*, 2017). In this line, a study showed a dependence of RTs on the amount of preparatory suppression on a single-trial basis: the stronger the suppression, the faster the initiation of the ensuing movement (Hannah *et al.*, 2018). Importantly, these hypotheses could be both valid

as they focus on different levels of control, which are both known to shape motor activity: while the choice hypothesis suggests that suppression originates from processes that help select accurate actions (*i.e.*, therefore reducing the error rate), the motor hypothesis entails that preparatory suppression is also generated by processes speeding up action initiation (*i.e.*, therefore reducing RTs).

Here, we investigated the impact of motor training on preparatory suppression, while subjects practiced an instructed-delay RT task. The choice aspects were clear-cut, as evident from the low error rates, even before training. Hence, in such task, the selection requirements are so low that there is no room for improvement; subjects can only become more skilled at the motor level, by initiating their action faster. Based on this, we predicted that RTs would decrease over the course of practice but that error rates would remain marginal. In addition, we expected resting CSE to increase with training, in accordance with previous work (Pascual-Leone *et al*., 1995; Butefisch *et al*., 2000; Duque *et al.*, 2008; Galea & Celnik, 2009; Christiansen *et al*., 2018). Based on the motor hypothesis (*i.e.*, that preparatory suppression fastens RTs), we expected that training would deepen the drop in excitability during action preparation with respect to rest, reflecting an increased preparatory suppression. Hence, we predicted that preparatory activity would not follow the training-related rise in resting CSE.

2. Materials and Methods

2.1. Participants

Fifteen right-handed healthy subjects participated in the present study (n=15; 10 women; 22.4±1.63 years old). Handedness was assessed via Edinburgh Handedness inventory (Oldfield, 1971). Participants filled out a TMS safety questionnaire to look for any contra-indication and gave written informed consent in accordance with the Ethics Committee of the Université Catholique de Louvain and the principles of the Declaration of Helsinki. We had to exclude one subject because we encountered a technical problem during the experiment; hence, analyses were run on the fourteen remaining subjects. Part of the data reported here has been exploited in a separate study (Vassiliadis *et al*., 2018). All of the data are expressed as mean±SE.

2.2. Task

Subjects were sited in front of a computer screen with the hands on response devices (**Fig.1A,** (Quoilin *et al*., 2016, 2018, 2019; Grandjean *et al*., 2019). They performed an instructed-delay RT task, which required them to choose between abduction movements of the left or right index finger according to the position of a preparatory cue (*i.e.*, a left- or right-sided ball separated from a goal by a gap). Participants had to prepare their movement once the ball appeared but to withhold responding until the onset of a "Go" signal (*i.e.*, a bridge). When the bridge appeared on the screen, subjects had to respond as fast as possible, allowing the ball to roll on the bridge and to reach the goal. To reduce anticipation of the "Go" signal, the bridge did not appear in some of the trials (5%). Trials always ended with a feedback score reflecting performance (see **Fig.1B**).

2.3 TMS Protocol

Monophasic pulses were delivered through one or two figure-of-eight shaped coils, each connected either to a Magstim 200² magnetic stimulator (Magstim, Whitland, Dyfed, UK) or a Magstim Bistim² magnetic stimulator. The TMS machine used to stimulate each hemisphere was counterbalanced between subjects. Pulses could be triggered in one (*i.e.*, single-coil TMS) or in the two coils (*i.e.*, double-coil TMS, **Fig.1C**) because the dataset was initially acquired for a separate study to establish the reliability of double-coil TMS to probe CSE bilaterally (Vassiliadis *et al*.,

2018; Grandjean *et al.*, 2018). Here, MEPs are considered regardless of the protocol used to elicit
them.

Each TMS coil was placed tangentially over one M1 with the handle pointing backward
and laterally at a 45° angle away from the midline **(Fig.1C)**. TMS was applied over the hotspot of
the first dorsal interosseous muscle (FDI), which was the prime-mover in our task (Duque *et al.*,
2014; Derosiere *et al.*, 2017a, 2017b). The resting Motor Threshold (rMT) was determined for each
M1. It was defined as the minimal intensity required to evoke MEPs of $50\mu V$ at rest in at least 5
out of 10 stimulations. The rMTs equalled 41.7±5.05% and 40.8±6.39% of the maximum
stimulator output for the left and the right FDI, respectively. For each hemisphere, the intensity
used throughout the experiment was set at 115% of the individual rMT (Derosiere *et al.*, 2019).

Figure 1. A, The response device. Index finger responses were recorded using a home-made
device composed of two pairs of metal edges fixed on a wooden platform and positioned under the
left (graphic representation) and right (photographic representation) hands. **B, "Rolling Ball" task.**
Subjects were asked to choose between responding with the left or right index finger according to
the position of a ball (Preparatory cue) appearing on the left or right part of the screen (left in the
current example). They had to wait until the onset of a bridge (Go signal) to release their response
as quickly as possible. The ball then rolled on the bridge (when the subjects answered correctly) to
reach a goal located on the other side of the gap. A feedback reflecting how fast and accurate the
subjects were concluded each trial. On correct trials, scores ranged from 1 to 100 points and were
displayed in green. Participants were informed that the score was inversely proportional to the RT:

the faster the response, the higher the score. In order to homogenize the score across subjects, scores on correct trials were individualized according to RTs measured during a familiarization session just before the main experiment (Vassiliadis *et al.,* 2018; Grandjean *et al*., 2019). Incorrect responses were penalized with negative scores displayed in red. They involved responses occurring too early (RT<100 ms), referred to as "anticipation errors" (-75 points), responses occurring too late (RT>500 ms), referred to as "time-out errors" (-50 points), responses provided with the incorrect hand (-20 points), referred to as "choice errors" and responses provided on catch trials (-12 points), referred as "catch errors". When subjects succeeded not to respond on a catch trial, they were rewarded by +12 points. The total score was displayed at the end of each block. **C, TMS protocol.** Two figure-eight-shaped coils were placed over the subject's M1, eliciting MEPs in the left and/or right FDI. In double-coil trials, a 1-ms interval separated the onset of the two pulses, eliciting MEPs in both hands at a near simultaneous time (Vassil*iadis et al*., 2018; Grandjean *et al*., 2018; Algoet *et al*., 2018; Quoilin *et al*., 2019). This interval was used to avoid direct electromagnetic interference between the two coils (Cincotta *et al.*, 2005), while preventing transcallosal interactions that would occur between motor areas with longer delays (Ferbert *et al.*, 1992; Hanajima *et al.*, 2001). Notably, in double-coil trials, half of the trials involved a pulse over left M1 first whereas the other half involved a pulse over right M1 first (1ms delay). These data were assembled because the a prior analysis reported elsewhere showed that the order of stimulation does not influence the double-coil MEP amplitudes, which were identical to single-coil MEPs (Vassiliadis *et al.*, 2018).

2.4 Experimental procedure

The experiment started with two familiarization blocks. The first block allowed subjects to become acquainted with the task. The second block involved TMS and served to compute the median RT for each subject. The latter was used to individualize the feedback scores on correct trials according to the initial performance (see **Fig.1B**).

Then, subjects performed 400 trials of the task, divided in 10 blocks. Each block involved an equal combination of single- and double-coil stimulations, occurring in a random order (*i.e.*, subjects could not anticipate the type of stimulation they would face). Given that both techniques produce equivalent MEPs (Vassiliadis *et al.,* 2018; Grandjean *et al.,* 2018), these data were considered regardless of the protocol used to elicit them.

TMS could occur in three different settings. First, some TMS pulses were delivered outside the blocks (TMS$_{baseline-out}$), providing MEPs reflecting baseline CSE at complete rest. TMS$_{baseline-out}$ pulses occurred every other block, starting before block 1 and ending after block 8 (5 time points; 30 MEPs per time point; **Fig.2A**). Second, TMS occurred during the intertrial interval, 300 ms before the beginning of the trial (**Fig.2B**). MEPs recorded at this time (12 per block) provided another baseline measure of CSE, with subjects at rest but engaged in the task (TMS$_{baseline-in}$ Labruna *et al.,* 2011). Finally, TMS occurred at 900 or 950 ms after the onset of the preparatory cue (TMS$_{preparation}$). Since no difference was found between MEPs recorded at these two timings in our previous analysis (Vassiliadis *et al.*, 2018), these data were pooled together (48 MEPs per block). Half of these MEPs fell in left response trials, while the other half occurred in right response trials. Hence, MEPs could either fall in a hand that was selected for the forthcoming response (MEP$_{selected}$; *e.g.*, left MEPs preceding a left index finger response) or in a hand that was non-selected (MEP$_{non-selected}$).

Figure 2. A, Time-course of the experiment. After two familiarization blocks (Fam1: 20 trials without TMS and Fam 2: 40 trials with TMS), subjects executed ten blocks of forty trials during

which MEPs were elicited at $TMS_{baseline-in}$ or $TMS_{preparation}$ (see 2A). The effect of training was assessed by comparing five sets of data ($Training_1$ to $Training_5$), each involving MEPs pooled over two consecutive blocks. MEPs were also elicited outside the blocks ($TMS_{baseline-out}$) at five points in time, before block 1 and after blocks 2, 4, 6 and 8, categorized as $Training_1$ to $Training_5$, similar to the MEPs elicited during the blocks. We obtained 15 $TMS_{baseline-out}$ MEPs per hand for each $Training_{STAGE}$. Comparing these data sets allowed us to consider potential training-related changes in resting CSE outside the context of the task. **B, Time course of a trial.** Trials were separated by a blank screen (intertrial interval; 2050 to 2300 ms) and always started with a preparatory cue appearing for a variable delay period (1000 to 1200 ms). Then, a Go signal was presented and remained on the screen until a response was detected, hence for the duration of the reaction time (RT). The feedback was presented at the end of each trial for 500 ms and depended on the RT on correct trials. Importantly, the variable delays in the task were sampled from uniform distributions to induce temporal uncertainty and therefore reduce anticipation that could emerge with the repetition of trials. TMS pulses occurred either during the intertrial interval (300 ms before the beginning of the trial; $TMS_{baseline-in}$), or during the delay period (900 or 950 ms after the preparatory cue onset, timings pooled; $TMS_{preparation}$). In double-coil trials, motor-evoked potentials (MEPs) were elicited in the first dorsal interosseous (FDI) of both hands at a near simultaneous time (1 ms delay); in single-coil trials, MEPs were elicited in the left or right hand. The figure displays a left hand trial with double-coil TMS at $TMS_{preparation}$.

2.5 Data processing and statistical analyses

The purpose of the study was twofold: (1) to characterize changes in CSE at rest and during action preparation occurring along with training in a basic instructed-delay RT task, (2) to assess whether modulations in CSE were correlated to training-related improvements in RTs. To do so, the behavioral and MEP data were evaluated according to the block within which they were elicited and data from two consecutive blocks were pooled together. Given the 10 blocks, we obtained five data sets reflecting five training stages ($Training_{STAGE}$: $Training_1$ to $Training_5$; **Fig.2A**).

Statistical analyses were carried out with Matlab 2018a (the Mathworks, Natick, Massachusetts, USA) and Statistica 10 (StatSoft Inc., Tulsa, Oklahoma, USA). All data were systematically tested for the sphericity assumption using Maunchley's tests. The Greenhouse–

245 Geisser (GG) correction was used for sphericity when necessary. Post-hocs comparisons were

246 always conducted using the Fisher's LSD procedure. The significance level was set at $p \leq 0.05$.

2.5.1 RTs and errors

Left and right hand RTs were computed as the difference between the onset of the Go signal

and movement onset (when the finger quitted the outer metal edge of the device). Trials where

subjects made an error were removed from the data set for the RT analysis. An average of 35 left

and 34 right response trials remained for each subject at each Training$_{STAGE}$. We computed the

mean RT for left and right responses separately and then averaged these data together. Besides, we

also assessed response accuracy over training, by computing, for each Training$_{STAGE}$, the amount

of anticipation, time-out and catch errors as well as the total error rate. For each of these variables,

we expressed the number of incorrect trials in percentage of the total amount of trials, regardless

of the responding hand. Choice errors were not analysed because they were rare (4 choice errors

across all subjects). For the statistical analysis of RTs and errors (*i.e.*, anticipation, time-out, catch

and global errors), we used one-way analyses of variance for repeated measures (ANOVA$_{RM}$) with

the factor Training$_{STAGE}$ (Training$_1$ to Training$_5$).

2.5.2 MEP amplitudes

MEPs were obtained by recording electromyography (EMG) bilaterally from surface

electrodes (Neuroline, Medicotest, Oelstykke, Denmark) placed over the FDI. The signals were

amplified (x1000), bandpass filtered (10-500Hz; NeuroloLog; Digitimer), digitalized at 2000 Hz

and collected with Signal (Signal 3.0, Cambridge, UK) for offline analysis. Trials with background

EMG activity preceding the pulse exceeding 3 SDs above the mean were discarded ($1.46\pm0.32\%$

removal; Duque *et al.,* 2014; Klein *et al.,* 2016).

To assess training-related changes in resting CSE based on MEPs elicited at TMS$_{baseline-out}$

and TMS$_{baseline-in}$, we averaged separately left and right hand MEPs for each Training$_{STAGE}$ before

computing the mean of these averages. These data were analysed using a two-way ANOVA$_{RM}$ with

TMS$_{TIMING}$ (TMS$_{baseline-out}$ or TMS$_{baseline-in}$), and Training$_{STAGE}$ (Training$_1$ to Training$_5$) as within-

subject factors. To assess training-related changes in preparatory suppression based on MEPs at

TMS$_{preparation}$ (expressed in percentage of MEPs at TMS$_{baseline-in}$), we first removed the trials in

which subjects made a mistake (10.78±1.81% removal) and then grouped left and right hand MEPs according to whether they corresponded to a $MEP_{selected}$ or $MEP_{non-selected}$. Within these categories, we averaged the separate means of left and right hand MEPs for each $Training_{STAGE}$. To analyse these data, we first focused on percentage MEPs at $Training_1$, assessing with t-tests (against a constant value of 100%) the significance of preparatory suppression at the beginning of training. Then, we analyzed all training stages using a two-way $ANOVA_{RM}$ with the factors $MEP_{SELECTION}$ ($MEP_{selected}$ or $MEP_{non-selected}$) and $Training_{STAGE}$ ($Training_1$ to $Training_5$). This ANOVA was also run on absolute MEP amplitudes (in mV).

2.5.3 Relationship between training-related changes in RTs and CSE

As described in the Result section, training influenced RTs and CSE. We studied the relationship between changes at these two levels, with CSE considered separately at rest and during action preparation. We computed ratios reflecting training-related changes. Based on the RT data, we realized that the subjects' behavior improved substantially during the first practice stage ($Training_1$ to $Training_3$) but then, RTs remained quite stable (from $Training_3$ to $Training_5$; Result section). For this reason, we considered ratios for these two phases of training separately, providing us with an indication of early ($Training_{ratio-early}$: $Training_3/Training_1$) and late ($Training_{ratio-late}$: $Training_5/Training_3$) training-related changes in RTs and CSE. For the latter, we computed separate ratios for MEPs at $TMS_{baseline-out}$, $TMS_{baseline-in}$ and $TMS_{preparation}$ (expressed in percentage of MEPs at $TMS_{baseline-in}$). We then examined the correlation between the RT and MEP $Training_{ratios}$ by using least squares linear regressions.

Finally, we compared the strength of the RT relationship to training-related changes in MEP amplitudes at $TMS_{baseline-in}$ (reflecting resting CSE) and changes in percentage MEPs at $TMS_{preparation}$ (reflecting preparatory suppression of CSE). To do so, in order to obtain a robust estimate of the absolute Pearson's R, we ran a bootstrap analysis with 10000 resamples and calculated a median R for each correlation (Efron, 1979). These R-values were then compared to each other using Pearson and Fillon's z test (Pearson & Filon, 1898).

2.5.4 Single-trial relationship between RTs and preparatory suppression

The correlation analyses revealed a relationship between RTs and preparatory suppression: the subjects who showed the greatest training-related reduction in RTs were also those who displayed the strongest deepening in preparatory suppression (see Result section). To better understand the dependency of RTs to the strength of preparatory suppression, we investigated whether this relationship was evident on a single-trial basis, as suggested previously (Hann*ah et al.,* 2018). We selected the MEPs elicited at $TMS_{preparation}$ and again, expressed them as a percentage of $TMS_{baseline-in}$. We only used the double-coil trials, to consider a homogeneous set of data, with preparatory MEPs falling in both hands systematically. For each trial, we extracted the RT, as well as the MEPs recorded at $TMS_{preparation}$ in both selected ($MEP_{selected}$) and non-selected ($MEP_{non-selected}$) hands. Hence, for each trial, we obtained one RT measure linked to two different MEPs.

To examine the relationship between RTs and preparatory suppression, we pooled the trials from all 10 blocks together and sorted them according to the amplitude of MEPs within each trial. Given that there were two MEPs in each trial, we repeated this procedure twice, providing us with two different orderings of the trials according to the $MEP_{selected}$ or $MEP_{non-selected}$. Within each arrangement, trials were grouped into 6 consecutive percentile bins (MEP_{BIN}: MEP_{BIN-1} = 0 to 16.7%, MEP_{BIN-2} = 16.7 to 33.3% … MEP_{BIN-6} = 83.3 to 100% of the data). MEP_{BIN-1} contained the trials with the stronger preparatory suppression whereas the MEP_{BIN-6} included the trials with the weaker preparatory suppression. We then computed the mean RT of trials within each MEP_{BIN} (23 trials per condition on average and never less than 19 trials), and then averaged the corresponding RTs. Hence, we obtained six average RT values (i.e. one for each MEP_{BIN}) for each of the trial arrangements based on the two MEP types. These two sets of RT data were analysed using two separate $ANOVA_{RM}$ with the factor MEP_{BIN} (MEP_{BIN-1} to MEP_{BIN-6}).

3. Results

3.1. RTs and errors

Fig.3A shows the evolution of RTs with training. The $ANOVA_{RM}$ revealed a significant influence of $Training_{STAGE}$ on RT ($F_{(4,52)}$=4.31, p=0.0043). Post-hoc tests showed that RTs measured from $Training_3$ to $Training_5$ were shorter than at $Training_1$ (all p<0.004). In contrast, the total error rate remained stable over the blocks ($F_{(4,52)}$=0.82, p=0.52, **Fig.3B**). We did not observe any modification of the percentage of anticipation ($F_{(4,52)}$=1.12, p=0.36), time-out (GG-corrected $F_{(2.50,32.50)}$=0.90, p=0.44) or catch errors ($F_{(4,52)}$=1.73, p=0.16). Hence, training enabled subjects to respond more quickly while maintaining the same accuracy level.

Figure 3. Evolution of reaction times (RTs) and total error rate throughout training. The mean RTs (**A**, in ms) and total error rate (**B,** in % of all trials) are represented for each $Training_{STAGE}$, regardless of the responding hand. Stars denote a significant difference between a given $Training_{STAGE}$ and $Training_1$ (p<0.05). Individual data for $Training_1$ and $Training_5$.

3.2. MEP amplitude

First, we evaluated the effect of training on MEPs acquired at rest. As evident on **Fig.4,** MEPs were larger when assessed in the context of the task ($TMS_{baseline-in}$: 1.79±0.17mV) compared to when subjects were fully at rest ($TMS_{baseline-out}$: 1.34±0.17mV), as supported by the significant factor TMS_{TIMING} ($F_{(1,13)}$=28.43, p<0.001) and consistent with previous studies (Labruna *et al.,*

2011; Derosière *et al.,* 2015). The ANOVA$_{RM}$ also revealed an effect of Training$_{STAGE}$ on baseline

MEPs ($F_{(4,52)}$=6.34, p<0.001). MEPs recorded at Training$_2$ to Training$_5$ were larger than at

Training$_1$ (all p<0.03). This training effect on MEPs occurred independently of the TMS$_{TIMING}$:

there was a parallel increase in the amplitude of MEPs elicited at TMS$_{baseline-out}$ and TMS$_{baseline-in}$

(Training$_{STAGE}$xTMS$_{TIMING}$: $F_{(4,52)}$=0.18, p=0.95).

Figure 4. Evolution of baseline MEPs throughout training. MEP amplitudes (in mV) elicited at

TMS$_{baseline-out}$ (black) and TMS$_{baseline-in}$ (pink) at the different Training$_{STAGES}$. Hash signs indicate a

TMS$_{TIMING}$ effect. Stars denote a significant difference between a given Training$_{STAGE}$ and

Training$_1$ (p<0.05). Individual data for Training$_1$ and Training$_5$ are also displayed.

Second, we analyzed the effect of training on preparatory suppression by considering MEPs

elicited at TMS$_{preparation}$ (expressed in percentage of TMS$_{baseline-in}$). As evident on **Fig.5A**,

percentage FDI MEPs were initially suppressed at Training$_1$ (MEPs smaller than 100%), consistent

with the presence of preparatory suppression in the prime-mover, whether selected for the

forthcoming response (MEP$_{selected}$: 73.98±4.00%; $t_{(13)}$=-6.50, p<0.0001) or not (MEP$_{non-selected}$:

76.26±4.36%; $t_{(13)}$=-5.44, p<0.001). Interestingly, preparatory suppression became more

prominent with training: the ANOVA$_{RM}$ revealed a significant decrease in percentage MEP

amplitudes over the Training$_{STAGES}$ ($F_{(4,52)}$=2.79, p=0.036). This change was marginal at Training$_4$

(*i.e.,* Training$_4$: p=0.058 when compared to Training$_1$) and became significant at Training$_5$

(p=0.006). It concerned MEPs obtained from the selected and non-selected hands

(Training$_{STAGE}$xMEP$_{SELECTION}$: $F_{(4,52)}$=0.56, p=0.70). To further our understanding of training-

related changes of preparatory activity, we ran another set of $ANOVA_{RM}$ on absolute MEP amplitudes (rather than percentages) at $TMS_{preparation}$ (**Fig.5B**). These MEPs did not show any fluctuation over the $Training_{STAGES}$ ($F_{(4,52)}=1.30$, $p=0.28$). Moreover, we did not find any $MEP_{SELECTION}$ effect ($F_{(1,13)}=1.69$, $p=0.22$) or $Training_{STAGE}xMEP_{SELECTION}$ interaction ($F_{(4,52)}=0.85$, $p=0.50$).

In conclusion, our results indicate that training did not produce even modulatory changes in motor activity at rest and during action preparation: while resting CSE increased, preparatory activity remained flat over the blocks, thus revealing an augmenting drop (i.e. preparatory suppression) with respect to the rising baseline excitability state. These changes in CSE occurred in parallel with an acceleration of RTs.

Figure 5. Evolution of preparatory MEPs throughout training. Normalized MEP amplitudes recorded at $TMS_{preparation}$ (in percentage of MEPs elicited at $TMS_{baseline-in}$) muscles at the different $Training_{STAGES}$ **(A)**. Absolute MEP data (in mV) are also represented muscles **(B)**. The star denotes a significant difference between a $Training_5$ and $Training_1$ ($p<0.05$). Note that the change in preparatory suppression was close to significance at $Training_4$ (*i.e.,* $p=0.058$ when compared to $Training_1$). Individual data for $Training_1$ and $Training_5$ are also displayed.

Because RTs became shorter over the blocks, one may argue that MEPs at $TMS_{preparation}$ were not recorded in a comparable preparatory state throughout training; that is, the delay between TMS and movement onset ($Delay_{TMS-TO-MOVE}$) may have decreased over the blocks. Importantly, we shuffled the delay between the pulse and the Go signal in the present study (see Methods), in

order to prevent changes in RT to convert into equivalent changes in the $Delay_{TMS-TO-MOVE}$.
However, because TMS fell on average closer to movement onset at $Training_5$ (399.70 ± 8.48ms)
than $Training_1$ (419.99 ± 9.99ms, $t_{(13)}=-3.10$, $p=0.008$), we performed an additional analysis to
control for a potential bias of the $Delay_{TMS-TO-MOVE}$. We conducted a response-locked analysis
whereby we classified MEP data at $TMS_{preparation}$ (regardless of the $Training_{STAGE}$) according to the
$Delay_{TMS-TO-MOVE}$ in 5 consecutive bins of trials ($Delay_{BIN}=Delay_{BIN-1}=0$ to 20%, $Delay_{BIN-2}=20$
to 40%, …, $Delay_{BIN-5}=80$ to 100% of the $Delay_{TMS-TO-MOVE}$ data). An $ANOVA_{RM}$ ran on these
data did not reveal any effect of $Delay_{BIN}$ ($F_{(4,52)}=1.45$; $p=0.23$), nor was there any significant
$MEP_{SELECTION}$x$Delay_{BIN}$ interaction ($F_{(4,52)}=0.40$; $p=0.81$; **Fig.6**). These results indicate that MEPs
elicited preceding a Go signal remain quite unaffected by the delay separating the $TMS_{preparation}$
pulse and movement onset.

Figure 6. Preparatory suppression according to time before movement onset. MEP amplitudes
recorded at $TMS_{preparation}$ (in percentage of MEPs elicited at $TMS_{baseline-in}$) are represented for each
$Delay_{BIN}$ in a selected (red) or non-selected (blue) muscle.

3.3 Relationship between training-related changes in RTs and CSE
Given that training influenced RTs and CSE, we studied the relationship between changes
at these two levels, with CSE considered separately at rest and during action preparation. To assess
the relationship between RTs and resting CSE, we ran correlations between training-related
changes in RTs and changes in MEPs at $TMS_{baseline-in}$ and $TMS_{baseline-out}$. These analyses did not
reveal any link between variations in resting measures of CSE and changes observed in RTs, neither

at Training$_{ratio\text{-}early}$ **(Fig.7A,** R=-0.27, p=0.36 and R=0.079, p=0.79 for TMS$_{baseline\text{-}in}$ and TMS$_{baseline\text{-}out}$, respectively) nor at Training$_{ratio\text{-}late}$ (R=-0.28, p=0.33 and R=-0.16, p=0.59).

In contrast, changes in RTs at Training$_{ratio\text{-}early}$ were linked to variations in preparatory suppression observed in the selected **(Fig.7B;** R=0.55, p=0.043**)** and non-selected FDI **(Fig.7C;** R=0.74, p=0.0027): subjects showing a greater training-related strengthening of preparatory suppression also showed larger improvements in RTs. This correlation was not significant at Training$_{ratio\text{-}late}$, neither for the selected (R=0.12, p=0.67) nor for the non-selected effectors (R=0.48, p=0.084). Our results suggest that RT improvements were related to early changes in preparatory suppression.

This conclusion is further supported by an additional analysis showing that the strength of the correlation between RTs and CSE at Training$_{ratio\text{-}early}$ was significantly higher when considering percentage MEPs at TMS$_{preparation}$ (i.e., preparatory suppression) in the non-selected FDI (bootstrap estimate of absolute R=0.76), than when MEPs were considered at TMS$_{baseline\text{-}in}$ (R=0.29; z=1.75; p=0.040, **Fig.7D**). This difference was not significant when taking preparatory suppression in the selected FDI (R=0.56; z-score=0.85, p=0.20). Hence, training-related changes in preparatory suppression of the non-selected effector turned out to be the best predictor of RT improvements.

Figure 7. Correlation between early training-related changes in RTs and CSE. Changes in RTs as a function of changes in MEP amplitudes at TMS$_{baseline\text{-}in}$ (reflecting resting CSE, **A**) and changes in percentage MEPs at TMS$_{preparation}$ (reflecting preparatory suppression of CSE) in the selected **(B)** and non-selected FDI muscle **(C)** during the early Training$_{stage}$. For this analysis, changes in RTs and MEPs were assessed by computing percentage ratios between the values obtained at Training$_3$ and Training$_1$. **(D)** Bootstrap estimates of absolute R values are also displayed

440   (± standard deviation of the samples) for each condition. These R values were compared by means

441   of a Pearson and Fillon's z test. One tail p-values were used given our a priori hypothesis

442   concerning the directionality of the effect ($p<0.05$).

3.4 Single-trial relationship between RTs and preparatory suppression

Finally, we asked whether the dependency of RTs to preparatory suppression is also evident

on a single-trial basis. This was the case for MEPs recorded from the non-selected hand: the greater

the preparatory suppression in that hand, the shorter the following RT (**Fig.8**, right panel), as

supported by the ANOVA$_{RM}$ revealing an effect of the factor MEP$_{BIN}$ on RTs ($F_{(5,65)}=2.57$,

$p=0.035$). Post-hoc tests revealed that RTs in MEP$_{BIN-1}$ and MEP$_{BIN-2}$ (i.e. strongest preparatory

suppression) were systematically shorter than those in MEP$_{BIN-6}$ ($p=0.0021$ and $p=0.0090$). We did

not observe any relationship between RTs and MEPs obtained in the selected hand (MEP$_{BIN}$: GG-

corrected $F_{(2.22,28.88)}=0.85$, $p=0.45$; **Fig.8**, left panel**).** Hence, the training-related effects and the

single-trial relationship indicates that preparatory suppression in the non-selected (non-responding)

hand is a predictor of the following RT. The lower this activity, the faster the response.

Figure 8. Single-trial MEP-RT relationship. Averaged RTs as a function of the preceding

preparatory suppression in a selected (left panel) or non-selected muscle (right panel). For this

analysis, the MEP data were divided in 6 MEP$_{BIN}$ of increasing amplitude and the RTs

corresponding to each MEP$_{BIN}$ were averaged. The star denotes a significant difference between

RTs at MEP$_{BIN-1}$ and MEP$_{BIN-2}$ and RTs at MEP$_{BIN-6}$ in the non-selected muscle ($p<0.05$). Note that

there was also a trend for RTs at MEP_{BIN-1} to be shorter than those in MEP_{BIN-4} (p=0.070) and

MEP_{BIN-5} (p=0.066).

4. Discussion

Training accelerated RTs while errors remained low. CSE became larger at rest and preparatory suppression of CSE was stronger after training. Interestingly, subjects who showed the strongest RT improvements at the early Training$_{STAGES}$ were also those displaying the largest initial strengthening in preparatory suppression, especially when probed in the non-selected hand. Such a relationship between RTs and preparatory suppression was also evident at a single-trial level: RTs were generally faster in trials where preparatory suppression was deeper.

Subjects responded faster with training. RTs reflect the sum of the time required for processing the imperative cue, preparing the motor command and initiating the action (Haith *et al.,* 2016; Derosiere *et al.,* 2019) and, theoretically, training may impact any of these sensory-motor components. Previous studies have shown that RT improvements can result from both faster sensory processing (Clark *et al.*, 2015) and more efficient motor preparation (Mawase *et al.*, 2018). Yet, in an instructed-delay task, the time required for sensory processing and motor preparation is strongly constrained and most of the RT is assumed to reflect the time needed for action initiation (Haith *et al.*, 2016). Hence, the RT gains reported here are likely to reflect a reduction in initiation time. Our findings thus yield an extension of former work, suggesting that, in addition to accelerating sensory processing and motor preparation, training can also boost action initiation.

Resting CSE was higher when assessed in the context of the task (i.e., at TMS$_{baseline-in}$) than between the blocks (i.e., at TMS$_{baseline-out}$), consistent with previous data (Labruna *et al.,* 2011; Vassiliadis *et al.,* 2018) and with the observation that task-driven increases in attention amplifies cortical excitability (Kastner *et al.*, 1998, 1999). As expected based on prior observations (*e.g.*, Pascual-Leone *et al.*, 1995; Butefisch *et al.*, 2000; Duque *et al.*, 2008; Galea & Celnik, 2009; Christiansen *et al.*, 2018), practicing the task led to an increase in resting CSE. Interestingly, this increase was not exclusive to the task and was in fact strongly similar at TMS$_{baseline-in}$ and TMS$_{baseline-out}$, ruling out the possibility that it resulted from a change in task-related attention over practice (Derosière *et al.*, 2015). Rather, our findings support the idea of a plastic reorganization of the motor system, measurable when engaged in the task as well as at rest.

CSE was reduced during action preparation when compared to baseline (during the task), reflecting the well-known preparatory suppression effect (Duque *et al.,* 2017), which was evident

in the selected and non-selected hands from the beginning of the training. Contrary to rest, the amplitude of MEPs at TMS$_{preparation}$ did not increase with practice (they remained unchanged), reflecting a strengthening drop in CSE from the rising baseline state. Notably, although at the group level this reinforcement of preparatory suppression appeared late (**Fig.5A**), at the individual level, a majority of subjects already exhibited a strengthening of preparatory suppression at early training stages (**Fig.6**).

Based on these findings, one could propose that changes in resting excitability are key to RT improvements, as suggested by the inverse relationship between baseline CSE and RTs described recently (Greenhouse *et al.,* 2017). Yet, we did not find a relationship between training-related changes in baseline excitability and improvements in performance. This is in line with the idea that increased resting CSE is not crucial for immediate performance (Bologna *et al.*, 2015), but may be involved in the long-term retention of the motor behavior (Cantarero *et al.*, 2013). Rather, what was predictive of RT gains in the present study was the change in relative CSE, as measured during action preparation: subjects showing the strongest reinforcement of preparatory suppression at the early Training$_{STAGES}$ were those who became fastest. These results are consistent with animal studies showing that behavioral improvements in motor learning tasks are associated with changes in relative preparatory activity (Paz *et al.*, 2003; Mandelblat-cerf *et al.*, 2009; Vyas *et al.*, 2018). Similarly, a recent study using paired-pulse TMS showed that changes in preparatory activity of M1 intra-cortical circuits are correlated to training-related behavioral gains, contrary to changes observed at rest (Dupont-Hadwen *et al.*, 2018). More generally, our findings agree with the idea that efficient action preparation relies on dynamical shifts of neural activity from a baseline state to a preparatory state (Churchland *et al.*, 2012). From this point of view, training may allow tuning the dynamics of preparatory activity, bringing it closer to an optimal state for action initiation (Vyas *et al.*, 2018). In this line, strengthening of preparatory suppression would facilitate action initiation by allowing excitatory inputs targeting the selected motor representation to better stand out against a quiescent background (mostly reflected in the excitability of non-selected effector), ultimately speeding up RTs (Hasbroucq *et al.*, 1997; Greenhouse *et al.*, 2015; Hasegawa *et al.*, 2017).

This interpretation was reinforced by our single-trial analysis showing that RTs depended on the foregoing amount of preparatory suppression. That is, stronger levels of suppression were related to faster initiation times in the very same trials, in agreement with previous results

525 (Hasegawa *et al.,* 2017; Hannah *et al.,* 2018). Interestingly, we found such relationship when

526 considering the non-selected prime-mover but not the selected one. This was also the case for

527 training-related effects, with preparatory suppression in the non-selected effector appearing as the

528 best predictor of RT changes, possibly because the selected effector is targeted by too many

529 overlapping inputs to supply as meaningful MEP amplitudes (Duque & Ivry, 2009).

Conclusion

This study shows that a simple training paradigm can lead to improvements in action initiation that are accompanied by an increase in resting CSE and a strengthening of corticospinal suppression from the rising baseline state. Moreover, contrary to changes in resting CSE, such strengthening of preparatory suppression was linked to RTs improvements. These findings could have implications for the rehabilitation of patients suffering from impaired action initiation such as in cerebellar ataxia (Battaglia *et al.,* 2006) or Parkinson's disease (Mure *et al.*, 2012).

References

Algoet M, Duque J, Iannetti GD & Mouraux A (2018). Temporal Profile and Limb-specificity of Phasic Pain-Evoked Changes in Motor Excitability. *Neuroscience* **386,** 240–255.

Battaglia F, Quartarone A, Ghilardi MF, Dattola R, Bagnato S, Rizzo V, Morgante L & Girlanda P (2006). Unilateral cerebellar stroke disrupts movement preparation and motor imagery. *Clin Neurophysiol* **117,** 1009–1016.

Bologna M, Rocchi L, Paparella G, Nardella A, Li P, Conte A, Kojovic M, Rothwell JC & Berardelli A (2015). Brain Stimulation Reversal of Practice-related Effects on Corticospinal Excitability has no Immediate Effect on Behavioral Outcome. *Brain Stimul* **8,** 603–612.

Butefisch CM, Davis BC, Wise SP, Sawaki L, Kopylev L, Classen J & Cohen LG (2000). Mechanisms of use-dependent plasticity in the human motor cortex. *Proc Natl Acad Sci* **97,** 3661–3665.

Cantarero G, Lloyd A & Celnik P (2013). Reversal of Long-Term Potentiation-Like Plasticity Processes after Motor Learning Disrupts Skill Retention. **33,** 12862–12869.

Christiansen L, Madsen MJ, Bojsen-møller E, Thomas R & Nielsen JB (2018). Brain Stimulation Progressive practice promotes motor learning and repeated transient increases in corticospinal excitability across multiple days. *Brain Stimul* **11,** 346–357.

Churchland MM, Cunningham JP, Kaufman MT, Foster JD, Nuyujukian P, Ryu SI, Shenoy K V. & Shenoy K V. (2012). Neural population dynamics during reaching. *Nature* **487,** 51–56.

Cincotta M, Borgheresi A, Jung P, Balestrieri F, Giovannelli F, Zaccara G & Ziemann U (2005). Physical interactions between induced electrical fields can have substantial effects on neuronal excitation during simultaneous TMS of two brain areas. *Clin Neurophysiol* **116,** 1733–1742.

Clark K, Gregory Appelbaum L, van den Berg B, Mitroff SR & Woldorff MG (2015). Improvement in visual search with practice: Mapping learning-related changes in neurocognitive stages of processing. *J Neurosci* **35,** 5351–5359.

Derosiere G (2018). A Dynamical System Framework for Theorizing Preparatory Inhibition. *J Neurosci* **38,** 3391–3393.

Derosière G, Billot M, Ward ET & Perrey S (2015). Adaptations of motor neural structures'

activity to lapses in attention. *Cereb Cortex* **25,** 66–74.

Derosiere G & Duque J (2020). Tuning the Corticospinal System: How Distributed Brain Circuits
Shape Human Actions. *Neurosci*107385841989675.

Derosiere G, Thura D, Cisek P & Duque J (2019). Motor cortex disruption delays motor
processes but not deliberation about action choices. *J Neurophysiol* **122,** 1566–1577.

Derosiere G, Vassiliadis P, Demaret S, Zénon A & Duque J (2017*a*). Learning stage-dependent
effect of M1 disruption on value-based motor decisions. *Neuroimage*; DOI:
10.1016/j.neuroimage.2017.08.075.

Derosiere G, Vassiliadis P & Duque J (2020). Advanced TMS approaches to probe corticospinal
excitability during action preparation. *Neuroimage* **213,** 116746.

Derosiere G, Zénon A, Alamia A & Duque J (2017*b*). Primary motor cortex contributes to the
implementation of implicit value-based rules during motor decisions. *Neuroimage* **146,**
1115–1127.

Dupont-Hadwen J, Bestmann S & Stagg CJ (2018). Motor training modulates intracortical
inhibitory dynamics in motor cortex during movement preparation. *Brain Stimul*; DOI:
10.1016/j.brs.2018.11.002.

Duque J, Greenhouse I, Labruna L & Ivry RB (2017). Physiological Markers of Motor Inhibition
during Human Behavior. *Trends Neurosci* **40,** 219–236.

Duque J & Ivry RB (2009). Role of corticospinal suppression during motor preparation. *Cereb*
Cortex **19,** 2013–2024.

Duque J, Labruna L, Cazares C & Ivry RB (2014). Dissociating the influence of response
selection and task anticipation on corticospinal suppression during response preparation.
Neuropsychologia **65,** 287–296.

Duque J, Lew D, Mazzocchio R, Olivier E & Richard B (2010). Evidence for two concurrent
inhibitory mechanisms during response preparation. *J Neurosci* **30,** 3793–3802.

Duque J, Mazzocchio R, Stefan K, Hummel F, Olivier E & Cohen LG (2008). Memory formation
in the motor cortex ipsilateral to a training hand. *Cereb Cortex* **18,** 1395–1406.

Duque J, Petitjean C & Swinnen SP (2016). Effect of aging on motor inhibition during action
preparation under sensory conflict. *Front Aging Neurosci* **8,** 1–14.

Efron B (1979). Bootstrap Methods: Another Look at the Jackknife. *Ann Stat* **7,** 1–26.

Ferbert A, Priori A, Rothwell JC, Day BL, Colebatch JG & Marsden CD (1992).

Interhemispheric inhibition of the human motor cortex. *J Physiol* **453,** 525–546.

Galea JM & Celnik P (2009). Brain Polarization Enhances the Formation and Retention of Motor

Memories. *J Neurophysiol* **102,** 294–301.

Grandjean J, Derosiere G, Vassiliadis P, Quemener L, Wilde Y de & Duque J (2018). Towards

assessing corticospinal excitability bilaterally: Validation of a double-coil TMS method. *J*

Neurosci Methods **293,** 162–168.

Grandjean J, Quoilin C & Duque J (2019). Investigating the effect of anticipating a startling

acoustic stimulus on preparatory inhibition. *Neurophysiol Clin* **49,** 137–147.

Greenhouse I, Sias a., Labruna L & Ivry RB (2015). Nonspecific Inhibition of the Motor System

during Response Preparation. *J Neurosci* **35,** 10675–10684.

Greenhouse XI, King XM, Noah XS, Maddock XRJ & Ivry RB (2017). Individual Differences in

Resting Corticospinal Excitability Are Correlated with Reaction Time and GABA Content

in Motor Cortex. **37,** 2686–2696.

Haith AM, Pakpoor J & Krakauer JW (2016). Independence of Movement Preparation and

Movement Initiation. *J Neurosci* **36,** 3007–3015.

Hanajima R, Ugawa Y, Machii K, Mochizuki H, Terao Y, Enomoto H, Furubayashi T, Shiio Y,

Uesugi H & Kanazawa I (2001). Interhemispheric facilitation of the hand motor area in

humans. *J Physiol* **531,** 849–859.

Hannah R, Cavanagh SE, Tremblay S, Simeoni S & Rothwell JC (2018). Selective suppression of

local interneuron circuits in human motor cortex contributes to movement preparation. *J*

Neurosci **38,** 2869–17.

Hasbroucq T, Kaneko H, Akamatsu M & Possamaï CA (1997). Preparatory inhibition of cortico-

spinal excitability: A transcranial magnetic stimulation study in man. *Cogn Brain Res* **5,**

185–192.

Hasegawa M, Majima K, Itokazu T, Maki T, Albrecht UR, Castner N, Izumo M, Sohya K, Sato

TK, Kamitani Y & Sato TR (2017). Selective Suppression of Local Circuits during

Movement Preparation in the Mouse Motor Cortex. *Cell Rep* **18,** 2676–2686.

Kastner S, Pinsk MA, De Weerd P, Desimone R & Ungerleider LG (1999). Increased Activity in
Human Visual Cortex during Directed Attention in the Absence of Visual Stimulation.
Neuron **22,** 751–761.

Kastner S, De Weerd P, Desimone R & Ungerleider LG (1998). Mechanisms of Directed
Attention in the Human Extrastriate Cortex as Revealed by Functional MRI. *Science (80-)*
282, 108–111.

Klein PA, Duque J, Labruna L & Ivry RB (2016). Comparison of the two cerebral hemispheres in
inhibitory processes operative during movement preparation. *Neuroimage* **125,** 220–232.

Klein PA, Petitjean C, Olivier E & Duque J (2014). Top-down suppression of incompatible motor
activations during response selection under conflict. *Neuroimage* **86,** 138–149.

Krakauer JW, Ghilardi MF, Mentis M, Barnes A, Veytsman M, Eidelberg D & Ghez C (2004).
Differential Cortical and Subcortical Activations in Learning Rotations and Gains for
Reaching: A PET Study. *J Neurophysiol* **91,** 924–933.

Krakauer JW, Hadjiosif AM, Xu J, Wong AL & Haith AM (2019). Motor Learning. **9,** 613–663.

Labruna L, Fernández-Del-Olmo M & Ivry RB (2011). Comparison of different baseline
conditions in evaluating factors that influence motor cortex excitability. *Brain Stimul* **4,**
152–155.

Makino H, Ren C, Liu H, Kim AN, Kondapaneni N, Liu X, Kuzum D & Komiyama T (2017).
Transformation of Cortex-wide Emergent Properties during Motor Learning. *Neuron* **94,**
880-890.e8.

Mandelblat-cerf Y, Paz R & Vaadia E (2009). Trial-to-Trial Variability of Single Cells in Motor
Cortices Is Dynamically Modified during Visuomotor Adaptation. **29,** 15053–15062.

Mawase F, Lopez D, Celnik PA & Haith AM (2018). Movement Repetition Facilitates Response
Preparation. *Cell Rep* **24,** 801–808.

Mawase F, Uehara S, Bastian AJ & Celnik P (2017). Motor Learning Enhances Use-Dependent
Plasticity. *J Neurosci* **37,** 2673–2685.

Mure H, Tang CC, Argyelan M, Ghilardi MF, Kaplitt MG, Dhawan V & Eidelberg D (2012).
Improved sequence learning with subthalamic nucleus deep brain stimulation: Evidence for
treatment-specific network modulation. *J Neurosci* **32,** 2804–2813.

Oldfield RC (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia **9,** 97–113.

Pascual-Leone A et al. (1995). Modulation of muscle responses evoked by transcranial magnetic
stimulation during the acquisition of new fine motor skills. *J Neurophysiol* **74,** 1037–1045.

Paz R, Boraud T, Natan C, Bergman H & Vaadia E (2003). Preparatory activity in motor cortex
reflects learning of local visuomotor skills. **6,** 882–890.

Pearson K & Filon L (1898). VII. Mathematical contributions to the theory of evolution.— IV.
On the probable errors of frequency constants and on the influence of random selection on
variation and correlation. *Philos Trans R Soc London Ser A, Contain Pap a Math or Phys*
Character **191,** 229–311.

Quoilin C, Fievez F & Duque J (2019). Preparatory inhibition: Impact of choice in reaction time
tasks. *Neuropsychologia* **129,** 212–222.

Quoilin C, Lambert J, Jacob B, Klein PA & Duque J (2016). Comparison of motor inhibition in
variants of the instructed-delay choice reaction time task. *PLoS One* **11,** 1–16.

Quoilin C, Wilhelm E, Maurage P, Timary P De & Duque J (2018). Deficient inhibition in
alcohol-dependence : let's consider the role of the motor system !
*Neuropsychopharmacology*0–8.

Reis J, Swayne OB, Vandermeeren Y, Camus M, Dimyan MA, Harris-Love M, Perez MA,
Ragert P, Rothwell JC & Cohen LG (2008). Contribution of transcranial magnetic
stimulation to the understanding of cortical mechanisms involved in motor control. *J Physiol*
586, 325–351.

Rosenkranz K, Kacar A & Rothwell JC (2007). Differential modulation of motor cortical
plasticity and excitability in early and late phases of human motor learning. *J Neurosci* **27,**
12058–12066.

Shmuelof L, Yang J, Caffo B, Mazzoni P & Krakauer JW (2014). The neural correlates of
learned motor acuity. 971–980.

Steele CJ & Penhune VB (2010). Specific increases within global decreases: A functional
magnetic resonance imaging investigation of five days of motor sequence learning. *J*
Neurosci **30,** 8332–8341.

Vassiliadis P, Grandjean J, Derosiere G & Wilde Y De (2018). Using a Double-Coil TMS
Protocol to Assess Preparatory Inhibition Bilaterally. **12,** 1–14.

Vyas S, Even-Chen N, Stavisky SD, Ryu SI, Nuyujukian P & Shenoy K V. (2018). Neural
Population Dynamics Underlying Motor Learning Transfer. *Neuron* **97,** 1177-1186.e3.

Vyas S, O'Shea DJ, Ryu SI & Shenoy K (2020). Causal role of motor preparation during error-
driven learning. *Neuron* 1–11.

Wenger E, Kühn S, Verrel J, Mårtensson J, Bodammer NC, Lindenberger U & Lövdén M (2017).
Repeated Structural Imaging Reveals Nonlinear Progression of Experience-Dependent
Volume Changes in Human Motor Cortex. *Cereb Cortex* **27,** 2911–2925.

Wiestler T & Diedrichsen J (2013). Skill learning strengthens cortical representations of motor
sequences. *Elife* **2013,** 1–20.

Yokoi A & Diedrichsen J (2019). Neural Organization of Hierarchical Motor Sequence
Representations in the Human Neocortex. *Neuron* **103,** 1178-1190.e7.