The prospective sense of agency is rooted in local and global properties of intrinsic functional brain networks

Simone Di Plinio[1*], Mauro Gianni Perrucci[1,2], Sjoerd J.H. Ebisch[1,2]

[1] Department of Neuroscience Imaging and Clinical Sciences, "G. D'Annunzio" University of Chieti-Pescara, Chieti 66100, Italy;

[2] Institute for Advanced Biomedical Technologies (ITAB), G D'Annunzio University of Chieti-Pescara, Chieti, Italy;

[*] **Corresponding author**. Address: Department of Neuroscience Imaging and Clinical Sciences, "G. D'Annunzio" University of Chieti-Pescara, Chieti 66100, Italy. E-mail: simone.diplinio@unich.it

Conflict of Interest: The authors declare no competing financial interests.

Keywords: Intentional binding, Motor control; Resting-state; Modularity; Sense of agency.

Running Title: Agency is embedded in local and global brain parameters.

Acknowledgments: This work was supported by the BIAL Foundation grant number 195/16 to SE, and by the "Departments of Excellence 2018-2022" initiative of the Italian Ministry of Education, University and Research for the Department of Neuroscience, Imaging and Clinical Sciences (DNISC) of the University of Chieti-Pescara.

1 ABSTRACT

The sense of agency (SoA) refers to a constitutional aspect of the self that is based on the

extent to which individuals feel in control over their actions and consequences thereof.

Although the SoA has been associated with mental health and well-being, it is still

unknown how interindividual variability in the SoA is embedded in the intrinsic brain

organization. We hypothesized that the prospective component of an implicit SoA is

associated with individual intrinsic brain connectivity on multiple spatial scales. Firstly,

we replicated previous findings by showing a significant prospective SoA as indicated by

intentional binding effects in healthy individuals. Secondly, using task-free functional

magnetic resonance imaging (fMRI) and graph analysis, we analyzed associations

between intentional binding effects and the intrinsic brain organization at regional,

modular, and whole-brain scales. The results showed that inter-modular connections of a

motor attention module, and particularly of the left supramarginal gyrus, are associated

with individual differences in prospective intentional binding. Notably, prospective

intentional binding effects were also related to global brain modularity within a specific

structural resolution range. These findings suggest that an implicit SoA generated through

sensorimotor predictions relies on the intrinsic organization of the brain connectome on

both local and global scales.

1. INTRODUCTION

The sense of agency (SoA) concerns the experience of oneself as the source of one's actions and their consequences. This phenomenon is often investigated as a state- or event-related subjective experience from which researchers can extrapolate the extent to which an individual feels in control over his or her behavior, thoughts, and the environment. Acquiring a sense of agency is considered a fundamental step in cognitive development (Ruvolo et al., 2015) as well as in human evolutionary adaptation (Taylor et al., 2014). The experience of a lower sense of agency can impair behavioral performances on cognitive tasks (Schooler et al., 2014), cause a loss of awareness (Berberian et al., 2017), and negatively impact the quality of everyday life (Bandura, 2006; Renes and Aarts, 2017) and mental health (Moore and Fletcher, 2012; de Bézenac et al., 2018).

Intentional binding, defined as the shift of the subjectively perceived timing of a performed action toward the consequences of that action, allows the measurement of an implicit sense of agency (Haggard et al., 2002; Moore and Obhi, 2012). Intentional binding can rely on two mechanisms (Moore and Haggard, 2008): firstly, a prospective intentional binding is based on the probabilistic and context-dependent coding of action consequences supported by internal predictive models; secondly, outcome-dependent, a posteriori inferences about action consequences support a retrospective intentional binding.

Studies on intentional binding have highlighted many substantial interindividual differences in the sense of agency and their implications for mental health. For instance, while healthy individuals show a significant prospective intentional binding effect (Moore and Haggard, 2008), increased schizotypal traits and schizophrenia have been associated with weakened prospective and increased retrospective intentional binding effects (Voss et al., 2010). It has also been shown that psychosis-related and positive social personality traits may predict decreases or increases in the prospective intentional binding depending on individual environmental control (Di Plinio et al., 2019a). Furthermore, intentional binding differs between high- and low-hypnotizable individuals (Lush et al., 2019) and correlates with narcissistic personality traits (Hascalovitz and Sukhvinder, 2015). These studies consistently demonstrate interindividual differences in the tendency to experience an enhanced or reduced sense of agency.

Increasing evidence from analyses based on graph theory (Bullmore and Sporns, 2009; Rubinov and Sporns, 2010) suggests that the intrinsic organization of brain networks contributes to the predisposition of an individual's typical behavioral patterns (Gallen and D'Esposito, 2019). Intriguingly, many brain regions implicated in agency, including the supplementary motor area (SMA), the inferior parietal lobule (IPL), the anterior insula, the cerebellum, and the precuneus

(Farrer et al., 2003; Desmurget et al., 2009; Chambon et al., 2013; Rae et al., 2014; Chambon et al.,

2015; Króliczak et al., 2016; Haggard, 2017), have been described as information hubs in the

human brain (Buckner et al., 2009; Power et al., 2013) that act as multimodal integrators of

sensorimotor information (van den Heuvel and Sporns, 2013) within motor control networks.

Whether the intrinsic functional connectivity patterns of these regions predispose

individual propensities regarding the sense of agency is still unknown. Moreover, the potential link

between agency and the intrinsic brain architecture may concern not only local elements of the

brain (i.e., specific regions/nodes or networks/modules), but also whole-brain parameters. For

example, it has been proposed that complex neurocognitive processes may also arise from the

brain's global modular structure (Ito et al., 2019). This view is supported by recent studies that

linked the brain's intrinsic organization to the predisposition of individual behavioral patterns

(Godwin et al., 2015; Hilger et al., 2017; Gupta et al., 2018). Specifically, the prospective sense of

agency reflects a constitutional aspect of self-awareness (Gallagher, 2000; Prinz, 2012) that emerges

from sensorimotor prediction models (Wolpert et al., 2001; Clark, 2013; Haggard, 2017). Predictive

coding, putatively grounded in the integration of actions and their outcomes (Kilner et al., 2007),

is a general property of the brain that is fundamental for both its development (Wolpert, 1997)

and its adaptability (Sato and Yasuda, 2005). Indeed, better contextual predictions mean smaller

prediction errors (Friston and Kiebel, 2009) and increased adaptability of the brain to external

demands. Therefore, the quality of these predictive mechanisms likely depends on global features

of the brain system, and a higher efficiency of the brain in generating accurate predictive schemes

may be necessary for a healthy sense of agency.

The purpose of this functional magnetic resonance imaging (fMRI) study was to investigate

if and how an implicit sense of agency, reflected by intentional binding, is associated with the

brain's intrinsic functional organization at multiple spatial scales. To that end, we first assessed the

prospective and retrospective components of the sense of agency in accordance with previous

studies (Haggard et al., 2002; Voss et al., 2010; Di Plinio et al., 2019a) in a sample of healthy

individuals (N = 39). Then, we studied the associations between implicit measures of the sense of

agency (prospective and retrospective intentional binding) and intrinsic functional network

features during a task-free (i.e., resting) state on three different, complementary spatial scales: nodal

(brain regions), modular (brain networks), and global (whole brain).

We hypothesized that the neural processing of the individual sense of agency may be

specifically modulated according to individual patterns of intrinsic brain connectivity. Given that

predictive mechanisms are more likely to be embedded in the intrinsic brain organization (Körding

et al., 2007; Kannape and Blanke, 2012; Apps and Tsakiris, 2014), while retrospective mechanisms are more likely related to task-evoked signals, we expected the prospective agency to be related with task-free fMRI measures. Specifically, regarding multi-scale properties, we expected that prospective intentional binding is associated with enhanced participation measures on the nodal and modular scales, indicating increased inter-network interactions of agency-related brain regions and modules, including the IPL, the precuneus, the insula, the cerebellum, and the preSMA. Moreover, we hypothesized that the sense of agency may also be entailed in general whole-brain parameters like global modularity, with higher modularity favoring increased prospective intentional binding.

2. MATERIALS and METHODS

2.1. Participants

Thirty-nine healthy Italian adults (19 females and 20 males, aged 23 ± 2) without a history of psychiatric or neurological disease and without contraindications for MRI scanning participated in the experiment. The experiment was approved by the local ethics committee. All participants had a normal or corrected-to-normal vision and provided written informed consent before taking part in the study in accordance with the Declaration of Helsinki (2013). Each participant performed the behavioral sense of agency task one to three days before the MRI acquisition.

2.2. Behavioral procedure

All participants performed a well-established sense of agency paradigm (Haggard et al., 2002; Voss et al, 2010; Di Plinio et al., 2019a). On each trial, the participants performed voluntary, self-initiated keypresses with the right index finger, while watching a clock hand rotating on a screen. The clock was labelled using a circular scale with numbers positioned at 5-minutes intervals. The participants' task was to judge the timing of the keypress, i.e., the clock time indicated by the clock hand when they pressed the key. The participants started each trial by voluntarily pressing a button with the left index finger. The clock hand rotation began at a random position in each trial and performed a full rotation every 2560 ms. The inter-trial interval was jittered between two and three seconds.

Two experimental conditions of action timing were used in the sense of agency task. In the 50% probability condition, participants heard a tone following the keypress in 50% of the trials. Instead, in the 75% probability condition, the keypress was randomly followed by a tone

1 presentation in 75% of the trials. The tone consisted of a single pulse at 1000 Hz and lasting 200

2 ms. The tone was presented binaurally 250 ms after the keypress. The task was performed using

Figure 1. Behavioral task. Participants performed voluntary, self-paced keypresses with the right index finger while watching a clock hand rotating on a screen. Participants started each trial by themselves pressing a button with the left index finger, and they were instructed to judge the clock 'time' indicated by the clock hand when they pressed the key. After the keypress, depending on the trial and on the experimental condition, an auditory feedback (or nothing, contingent on the current trial) was binaurally presented through headphones.

the E-Prime software (Schneider et al., 2012), while participants were sitting approximately 60 cm

from the 20-inches monitor. Two blocks for each condition (50%, 75%) of the sense of agency

task were performed by each participant. Each block consisted of 32 trials. The participants also

performed a fifth, baseline block composed of 32 trials in which no tones occurred. Block order

was pseudo-randomized across participants. The whole procedure lasted approximately 45

minutes for each participant.

The variable measured in each trial of the sense of agency task was the perceptual shift (Δ).

The perceptual shift is defined as the time difference between the real keypress and the perceived

keypress indicated by the subject in each trial, and it is expressed in milliseconds. To note, although

the perceptual shift and the phenomenon of the intentional binding are related, they should not

be confused: while the former represents the divergence between real and perceived keypress, the

second is a phenomenon which concerns the shift of the perceived timing of action toward the

action consequences (sound).

The participants were instructed (i) to avoid the planning of their keypresses, (ii) to avoid

answering in a stereotyped way, and (iii) to avoid answering before the end of the first full clock

rotation. A brief training session was performed by each participant before the experiment. The

task is illustrated in Figure 1.

2.3. Data acquisition

Each participant performed two consecutive task-free fMRI runs, each consisting of 376 volumes. The participants were instructed to watch a white fixation cross on a black screen without performing a cognitive task. Each run lasted approximately 7.5 minutes. Functional images were acquired using a Philips Achieva 3T scanner installed at the Institute for Advanced Biomedical Technologies (Gabriele D'Annunzio University, Chieti-Pescara, Italy). Whole-brain functional images were acquired with a gradient echo-planar sequence using the following parameters: repetition time (TR) = 1.2 s, echo time (TE) = 30 ms, field of view = 240x240x142.5 mm, flip angle = 65°, in-plane voxel size = 2.5 mm^2, slice thickness = 2.5 mm. A high-resolution T1-weighted whole-brain image was also acquired after functional sessions using the following parameters: TR = 8 ms, TE = 3.7, FoV = 256x256x180 mm, flip angle = 8°, in-plane voxel size = 1 mm^2, slice thickness = 1 mm.

2.4. Preprocessing of MRI data

The preprocessing steps and the analysis of functional images were implemented in AFNI (Cox, 1996). A slice-timing correction was applied to remove differences in acquisition times between the slices. The obtained functional images were deobliqued, despiked, corrected for time-shifted acquisition, and motion-corrected using a six-parameter rigid body realignment before being aligned to the Montreal Neurological Institute (MNI) standard brain using non-linear warping. During the preprocessing steps, motion parameters for each participant were collected. The functional images were scaled to have voxels with an average value of 100. A Gaussian filter of 5 mm full-width at half-maximum was applied to spatially smooth the functional images.

Finally, the time series were censored (volumes with 10% or more outliers across voxels and volumes with Euclidean norm of the motion derivative exceeding 0.3 mm were excluded), and band-pass filtered (frequency interval: 0.01-0.10 Hz) in a single regression step (Caballero-Gaudes and Reynolds, 2017) in which the motion parameters were also included as noise regressors together with white matter and cerebrospinal fluid signals. We did not regress out the global signal because it is a controversial approach (Saad et al., 2012) and potentially introduces spurious negative correlations (Weissenbacher et al., 2009).

2.5. Behavioral data analysis

The average baseline perceptual shift (Δ_B) was subtracted from the average perceptual shift in each

experimental condition and in each participant (Voss et al., 2010; Di Plinio et al., 2019a). Linear

mixed-effects analyses were implemented using *tone probability* (levels: 50% and 75%) and *trial type*

(levels: Action & tone, action only) as fixed effects, while a random intercept was added at the

subject level. The dependent variable was the average perceptual shift minus the baseline

perceptual shift (Δ-Δ_B). To identify the significance of prospective/retrospective components, we

performed multiple comparisons using Tukey's honest significant difference test. The

homogeneity of residuals was assessed using the D'Agostino-Pearson test.

We also implemented a bootstrap procedure to estimate the distributions of the effect sizes

and to investigate the modulation of the prospective and retrospective components (Kirby and

Gerlanc, 2013). The prospective component was estimated contrasting the perceptual shifts during

'action only' trials' in the 75% probability condition versus perceptual shifts during 'action only'

trials' in the 50% probability condition, thus, reflecting contextual effect of tone probability on

perception. The retrospective component was estimated contrasting the perceptual shifts between

'action+tone' and 'action only' trials within the 50% probability condition, thus, reflecting the

outcome-dependent effect on perception. The effect size (Hedge's g) was estimated in 10,000

bootstrap cycles with replacement for both the prospective and retrospective components.

19   2.6. Connectomics

The two task-free runs were concatenated and whole-brain functional connectomes were formed

using a set of 418 nodes derived from the cortical (346 parcels) and subcortical (40 grey nuclei)

atlases from Joliot and colleagues (2015) plus the cerebellar (32 nodes) atlas from Diedrichsen and

colleagues (2009). Functional connectivity was calculated using the z-Fisher transform of the

Pearson correlation among average time series extracted from the voxels within each node.

Connectomes were used to build binary undirected graphs after thresholding (the 10% strongest

connections were retained). Graph analyses were performed within Matlab using the Brain

Connectivity Toolbox (Rubinov and Sporns, 2010). Modular subdivisions of the brain were

visualized using the software BrainNet Viewer (https://www.nitrc.org/projects/bnv/; Xia et al.,

2013) implemented in MatLab. Formal definitions of the metrics used in graph analysis are

reported in Table I.

To find the optimal community structure of the networks, we implemented modularity

maximization (Porter et al., 2009) through the application of the widely used and robust Louvain

algorithm (Lancichinetti and Fortunato, 2009). An iterative fine-tuning process was used to find subject-specific optimal modular structures (Sun et al., 2009) and to handle the stochastic nature of the Louvain algorithm (Bassett et al., 2011). In the first step, each node was assigned to a distinct module. Then, the optimal community was detected using the Louvain algorithm and the modularity Q was estimated. This procedure was repeated using the optimal community found in the previous cycle as the starting community affiliation vector, and the process was repeated until Q could not be increased anymore. The group-level modular structure was achieved from the agreement matrix using a community detection algorithm developed for the analysis of complex networks (Lancichinetti and Fortunato, 2012), with the number of repetitions set to 1,000.

During the detection of optimal communities of nodes in the brain, the structural resolution parameter (γ) plays an important role since it weights the null model in the modularity estimation. In the present study, γ was varied in the interval [0.3-5.0] with steps of 0.1 to avoid biasing the subsequent analyses (Betzel et al., 2016). Then, the similarity between the consensus structure and the community structure across participants was calculated using the adjusted Rand coefficient (Traud et al., 2011) for each γ. Finally, an automated maximization algorithm was implemented in Matlab to detect one or more γ_{MAX} associated with local maxima of the Rand coefficient among the 48 tested gammas. The Newman-Grivan procedure was employed to detect significant modules in the consensus structure(s) (Newman and Girvan, 2004), as follows. Null models (10,000) were generated using a random permutation of module assignments while maintaining the original size and number of modules. Within each null model, the modularity contribution Q_C of each module was calculated as the summation of the real minus the expected number of connections between node pairs. The expected number of connections between each node pair was weighted with node degree (Newman and Girvan, 2004; Betzel et al., 2016). Only modules in which the real modularity contribution was greater than 99.9% of the simulated modularity contributions ($p < .001$) were considered significant.

The percentage of each predefined network (Di Plinio and Ebisch, 2018) covered by each module was estimated to characterize the modules' functional fingerprints.

2.7. Brain-behavior relationship analysis

Global statistics of modularity (Q) and nodal statistics of participation (Guimerà and Nunes Ameral, 2005) were extracted and analyzed with respect to the prospective component of the sense of agency.

A two-step procedure was applied to analyze the participation coefficients excluding poor
fitting and overfitting. In the first step, the model was fitted using all the module's nodes as levels
of a random grouping variable, and the prospective component was used as a continuous
predictor. In the second step, the analyses were repeated after removing nodes violating model
assumptions, that is, with inhomogeneous residuals within the corresponding random grouping as
indicated by Anderson-Darling normality tests. Regressions were performed separately for each
module to detect module-specific associations between participation measures and the sense of
agency. Results were corrected for multiple comparisons after model diagnostics and outlier
removal. Best linear unbiased predictors (BLUPs) were extracted to estimate the effect at the nodal
level and highlight nodes with the highest contributions whereas significant effects were detected
(Liu et al., 2008). Individual Conditional Expectation (ICE) plots were generated to visualize
significant effects at both the nodal and modular level (Goldstein et al., 2013). The impact of
significant results was tested also considering different graph density thresholds (10, 15, 20, 25,
30%) through the cost integration approach (Ginestet et al., 2011; Hilger et al., 2017), conforming
to current neuroscientific standards (Nichols et al., 2017; Van den Heuvel et al., 2017).

Starting from the hypothesis that the individual intrinsic brain signatures may be associated
with the implicit sense of agency depending on the structural resolution of the modular
architecture, the modularity Q was analyzed using all the γ values in the interval [0.3-5.0]. Since
this approach represents a methodological novelty, three different analyses were implemented to
test these associations. Firstly, a simple linear model was fitted for each value of γ using the
modularity Q as the dependent variable and the prospective component as a continuous predictor.
Secondly, the link between modularity and the sense of agency was tested in a multivariate model
including all the structural resolutions in a single analysis, with levels of γ being used as repeated
measures. Thirdly, a penalized regression (elastic net, Zou and Hastie, 2005) was adopted to
account for the many potentially correlated variables, i.e., modularity values at different levels of
γ. The optimal regularization coefficient of shrinkage (λ) for the penalized regression was chosen
as the λ minimizing the mean squared error in 100 Monte Carlo cycles of two-folded cross-
validations (Hastie, et al., 2009). Since the LASSO-penalty in the regression constraints irrelevant
variables (irrelevant $γ_s$) to be zero, the stability of feature selection was estimated calculating the
frequency of non-zero coefficients in 1,000 bootstrap cycles (threshold: $p < .05$). It can be pointed
out that functional connectivity may depend on the level of musical expertise or fitness of the
individuals (Herting and Nagel, 2013; Kühnis et al., 2014). To this aim, the analyses described
above were also performed adding years of sport training and music expertise as covariates.

3. RESULTS

3.1. Behavioral Results: confirming a significant prospective sense of agency in healthy subjects

The average perceptual shifts are reported in Table II and illustrated in Figure 2a. As expected, we found a strong prospective component, while the retrospective component was nearly zero. This effect was confirmed by the results of the mixed-effects model. Significant effects were found for the factor *tone probability* (p = .02, $F_{(1,160)}$ = 5.7) and for the interaction between *tone probability* and *trial type* (p = .01, $F_{(1,160)}$ = 6.1). The factor *trial type* by itself was not significant (p = .38, $F_{(1,160)}$ = 0.8). The statistics observed for the intercept (p = .06, $F_{(1,160)}$ = 3.4) showed that the average perceptual shift (Δ) in the experimental conditions was different from the baseline perceptual shift (Δ_B), though only a trend was observed. The prospective component was confirmed by contrasting perceptual shifts between the 'action only' trials of the two probability conditions (p < .001, Tukey's HSD test).

The results of the bootstrap procedure to estimate the effect sizes are shown in Figure 2b and indicated a medium-to-large effect size regarding the prospective component (average Hedges' g = 0.39, bootstrapped 95% CI: [0.19 ~ 0.61]). Effect sizes related to the retrospective component were rather irrelevant (g = 0.10, 95% CI: [-0.04 ~ 0.23]). Finally, the difference from the baseline was medium-to-large (g = 0.38, 95% CI: [0.08 ~ 0.67]).

Figure 2. Behavioral results. (a) Average values and standard errors of perceptual shifts relative to the baseline condition (Δ-ΔB), which represent measures of intentional binding. As expected, participants showed a significant effect relative to the prospective component. (b) Bootstrapped effect sizes (Hedges' g) for the prospective component, retrospective component, and intercept. Null distributions were created scrambling data between experimental conditions in 10,000 bootstrap cycles. While a medium-to-large effect was observed for the prospective component (upper panel) and for the intentional binding (lower panel), the retrospective component showed a close to zero effect size (middle panel). These results replicate findings from Di Plinio et al. (2019a).

3.2. Graph Analyses: the modular brain structure

Three resolution parameters, corresponding to γ_1=0.7, γ_2=1.2, and γ_3=2.1 were found to maximize

the adjusted Rand coefficients (Figure 3a). We report results related to the modular structure

associated with γ_3 since significant relationships with the sense of agency were found within this

modular configuration. Group average matrices of functional connectivity are reported in Figure

3b and show the eight significant modules found with γ_3 (M1-8, p < .001 using the Newman-

Girvan procedure).

Anatomical fingerprints of the eight modules with respect to predefined brain networks
(Di Plinio and Ebisch, 2018) are depicted in Figure 3c. The consensus community detection (with
γ_3) detected four modules encompassing associative cortices in the frontal, parietal, and temporal
lobes (M1-4). Among these modules, M1 was mainly related to default mode and language systems,
M2 encompassed high-order executive and attentional regions, M3 included networks associated
with motor attention and social interactions, and M4 incorporated frontal, insular, and cingulate
regions often ascribed to salience and control networks. Furthermore, two sensory modules were
detected, M5 and M6, that were related to sensorimotor/auditory and visual networks,
respectively. Among the last two modules, M7 included hippocampal and amygdala regions, while
M8 included subcortical and cerebellar structures. Module topographies are illustrated in Figure
3d.

Figure 3. Consensus modularity. (a) Local maxima of the Rand coefficient were detected for three gamma values: γ1 = 0.7; γ2 = 1.2; γ3 = 2.1. (b) Group average functional connectivity matrix (γ3 = 2.1). Each significant module (p < .001, comparison with 10,000 null models following the Newman-Girvan procedure) is enclosed in a square, distinguishing within-modules (inside the squares) and between-modules connections (outside the squares). (c) Functional fingerprints of the eight modules identified with γ3 = 2.1 as compared with predefined brain networks (Di Plinio and Ebisch, 2018). The width of each circle represents the amount of overlap between each network and each module, calculated as the normalized percentage of each network covered by each module. (d) Modular structural topographies with γ3 = 2.1. A full list of regions included in each module is available upon request to the corresponding author.

3.3. Modular and nodal associations with a prospective sense of agency

A significant association between the participation coefficient and the predictive sense of agency was found at the modular level with γ_3. The network participation of M3 (motor attention module)

was positively associated with the prospective component of the sense of agency ($t_{(310)} = 2.7$, p =

.007, β = 4.5e-4 CI: [1.2e-4 ~ 7.8e-4]; adjusted $R^2 = 0.42$). Figure 4a represents the nodal ICE plot

and shows the effect both at the modular (thick line) and nodal level (shaded lines). The nodes

which showed stronger effects are represented with a higher color intensity and a bigger size in

Figure 4b. Scatter plots and effect sizes ($β_s$) for the two nodes in M3 showing the largest effects,

that were located in the left supramarginal gyrus (average t = 3.8, p < .005), are reported in Figure

4c. The impact of significant results was significant also using different graph density thresholds

as indicated by the cost integration approach (Figure 4d). Summarizing, since the participation

coefficient reflects the extent of nodal interactions with other modules, these results indicate the

presence of a fronto-parietal module of which the inter-modular connectivity covaried with the

individual implicit sense of agency. Within this module, the strongest contributors to such effect

were two nodes located in the left supramarginal gyrus. No significant effects were found with

respect to the retrospective component.

Figure 4. Associations between agency and nodal/modular participation. (a) Individual Conditional Expectation (ICE) plot for the module M3 which illustrates the relationships between prospective component and network measures of participation. Each slope represents the zero-centered predictions of a node's participation coefficient (Y axis) for a given individual score in the prospective component (X axis). Color intensity indicates the effect strength (beta value, estimated using Best Linear Unbiased Predictors, BLUPs). The average effect at the module level is indicated by a thick black line (t(310) = 2.7; p = .007; β = 4.5e-4). (b) Module M3. The size and the color intensity of the nodes are regulated according to the strength of the associations (βs) between the prospective component and participation coefficients. (c) Nodes in M3 in which the effect was statistically stronger than the average slope for the module. These nodes were in the left supramarginal gyrus (L-SMG). For each node, the MNI coordinates and the effect sizes are indicated. (d) Estimates and relative standard errors for the interaction between prospective component of the sense of agency and participation coefficient (module M3) are reported for each graph threshold analyzed. To note, results were significant using all the threshold range considered (from 0.10 to 0.30).

3.4. The prospective sense of agency is associated with whole-brain modularity

The association between the prospective sense of agency and multi-scale global modularity was investigated in multiple ways. Using linear regression models, significant positive relationships between the two measures were found for γ ranging in the interval [1.4-3.0] (Figure 5a). The t-statistics for the effect of the prospective component on modularity Q with increasing γ are reported in Figure 5b. In particular, the strongest association was found for γ = 2.2 ($t_{(37)}$ = 2.9, p = .0058., β = 9.4e-5).

Figure 5. Associations between agency and global modularity: Linear regression. (a) Scatter plots representing values of the prospective component (X axis) against global modularity (Y axis) for each value of γ tested. Significant (p < .05) interactions are indicated with bold axes and orange colors. (b) The t-statistic for the association between global modularity (Q) and the prospective component is represented for each γ and is overlaid on the Rand coefficient. Darker colors in the interval γ = [1.4-3.0] indicate significant associations (p < .05).

The same effect was observed using a multivariate model. A higher prospective sense of agency was associated with a higher modularity Q ($F_{(47)}$ = 3.4, p < .001). The effect was still significant after applying the Huynh-Feldt correction for sphericity (p = .04). This relationship was

especially strong for γ_s in the interval [1.5-3.0]. Figure 6a reports the predicted Q for different

values of the prospective component. Although modularity is acknowledged as a measure of the

internal cohesion of modules, increasing the structural resolution parameter automatically results

in a lower Q. This happens because, by using additional smaller modules, more connections

become cross-modular, that is, more connections are excluded in the estimation of Q. To

characterize the relationship between a γ-standardized measure of the modularity and the

prospective sense of agency, we also estimated Q_{ADJ}, that is, a modularity adjusted for the value of

γ ($Q_{ADJ} = \gamma Q$). Figure 6b reports the predicted Q_{ADJ} against increasing values of the prospective

sense of agency using the same multivariate model. The association between Q_{ADJ} and the sense

of agency was still significant ($F_{(47)} = 3.9$, $p < .001$), and remained significant also after Huynh-

Feldt correction for sphericity ($p = .02$).

The cross-validated mean square errors (MSE) regarding the elastic net penalized model

are shown in Figure 6c. The λ associated with the minimum MSE ($\lambda = 7.6$) produced a structure

of beta estimates with 16 non-zero significant values (Figure 6d). Remarkably, as shown in Figure

6e, the elastic net regression returned non-zero estimates in correspondence of almost all (16 out

of 17) the significant γ values from the linear regression models (the standardized βs were in the

range [2.1 ~ 21.9]), with the highest effects associated with $\gamma = 2.1$ ($\beta = 21.1$) and $\gamma = 2.4$ ($\beta =

21.9$). Modularity values with $\gamma = 2.2$, $\gamma = 2.3$, $\gamma = 2.4$, and $\gamma = 2.5$ were significant following the

bootstrap procedure used to control for the stability in feature selection (darker bars in Figure 6e).

Summarizing, the results from linear, multivariate, and penalized regression models

showed a significant positive association between the prospective (but not the retrospective) sense

of agency and the global modular organization of the brain with medium structural resolutions

(with γ in the interval [1.5-3.0]). Importantly, even if it may be considered a sub-optimal model,

results from linear regression would allow possible comparisons with studies focalizing on single

γ values. The same effects were significant when adding years of sport training and music expertise

as covariates, showing the independence of the effects from these features.

Figure 6. Associations between agency and global modularity: Multivariate and penalized regressions. (a) Multivariate model. Predicted values of modularity (Q) with increasing values of γ. The color indicates the value of the prospective component (red = low prospective component, ≈ -35 ms; green = high prospective component, ≈ 65 ms). (b) Multivariate measures model. Predicted values of adjusted modularity (QADJ) with increasing values of gamma. The color indicates the value the prospective component (red = low prospective component, ≈ -35 ms; green = high prospective component, ≈ 65 ms). The comparison of Q plot versus QADJ plots illustrates the dependency of Q on the structural resolution of the network. (c) Penalized regression. Cross-validation for the penalized regression using elastic net (α = 0.5). The minimization of the mean squared error (MSE) occurred with λ = 7.6. (d) Penalized regression. The MSE minimization implied a structure with 16 non-zero beta estimates (degrees of freedom: df = 16). (e) Penalized regression. Standardized effect sizes associated with each value of γ. The 99% confidence intervals in the figure were obtained through 1,000 cycles of penalized regression using bootstrap with replacement. As shown, elastic net regression returned non-zero values of beta in the γ interval [1.5-3.0]. Darker green colors in the interval [2.2-2.5] indicate gammas associated with non-zero betas following the bootstrap procedure (p < .05).

4. DISCUSSION

Combining behavioral and task-free fMRI neurophysiological acquisitions, we showed that signatures of intrinsic brain activity are associated with the individual sense of agency in healthy adults. On the behavioral level, we focused on intentional binding as a measure of predictive and retrospective components of an implicit sense of agency (Voss et al., 2010; Haggard, 2017). On the neurophysiological level, graph-theoretical metrics were used to describe the intrinsic organization of functional brain networks based on a multi-scale, cross-resolution approach (Bullmore and Sporns, 2009; Rubinov and Sporns, 2010). To the best of our knowledge, this is the first study that traces the experimental measure of the sense of agency to individual differences in the intrinsic functional brain architecture. Moreover, we showed that this relationship exists on multiple spatial scales. The detected associations on nodal, modular, and global whole-brain scales, as well as their implications for cognitive neuroscience and psychopathology, are discussed below.

At the regional (nodal) and network (modular) levels, participation coefficients were studied as measures of cross-network integration. Our results revealed an association between the prospective sense of agency and participation coefficients in a fronto-parietal module, labeled M3, which consists of regions related to motor attention, such as the inferior parietal lobule (IPL), the dorsal precuneus, and the premotor cortex (PMC). These regions have previously been found to be interconnected both structurally and functionally in a task-free state, as well as to be connected with other regions related to sensorimotor processes, such as the supplementary motor cortex and the anterior insula (Caspers et al., 2011; Wang et al., 2012). The cross-network connectivity of two nodes in the left supramarginal gyrus (SMG), which is part of the IPL, showed the strongest associations with the prospective sense of agency.

Previous literature has outlined a network of brain regions supporting the sense of agency. Specific frontal and parietal brain regions have been associated with the sense of control over the environment to predispose the body to motor learning (Wolpert et al., 2011) and to generate internal predictions, which are then compared with perceived sensory stimuli (Gallese, 2000; Desmurget and Sirigu, 2009; Haggard, 2017). For example, the IPL is a multimodal integration region (Caspers et al., 2011) involved in the translation of motor intentions into meaningful actions (Farrer et al., 2003; Rae et al., 2014) and in distinguishing the self from other agents to generate a prospective sense of agency (Chambon et al., 2015; Ticini et al., 2018). Some studies reported that within the IPL, the left supramarginal gyrus (SMG) plays a prominent role in motor control and motor attention (Rushworth et al., 2003; Króliczak et al., 2016). On the other hand, the PMC is involved in monitoring conscious movements (Desmurget et al., 2009) and in higher-order

cognitive processes like overt task control (Chambon et al., 2013), while the dorsal precuneus has
been associated with the perceived experience of agency (Cavanna and Trimble, 2006).
Accordingly, certain sensorimotor disorders, such as alien hand syndrome (Assal et al., 2007;
Hassan and Josephs, 2016) and impaired movement initiation (Desmurget et al., 2009), have been
associated with aberrant functioning of the IPL, and others, such as anosognosia for hemiplegia
(Vocat et al., 2019), with impaired functioning of the PMC. These impairments often compromise
the online adjustment of motor actions in voluntary versus externally driven movements (Fried et
al., 2017).

Taken together, these task-related studies have highlighted the role of the SMG, the PMC,
and the dorsal precuneus in sensorimotor processes that support the experience of control over
actions and their intended consequences. In this context, this study's results on the local and
modular scales make two major contributions. Firstly, behavioral measures of an implicit,
prospective sense of agency are related to the cross-module integration of a fronto-parietal
subsystem (module M3) encompassing regions related to motor control and sense of agency,
namely, the SMG, the PMC, and the dorsal precuneus. Secondly, within this subsystem, the most
prominent relationships between intrinsic functional inter-network connections and the
prospective sense of agency exist in the left SMG. Hence, our findings show that the prospective
sense of agency may be rooted in the intrinsic functional connectivity patterns of the same regions
that are modulated during task execution. Additionally, they suggest that altered inter-network
connections in the brain may shift the individual predisposition to contingent effects conveyed by
the contextual sensory feedback, which consequently leads to diversified behavioral responses
across the population. Conceivably, stronger inter-network connections mediated by this fronto-
parietal subsystem, and especially by the left SMG (Wang et al., 2012), may be necessary for
integrating action consequences with ongoing motor behavior to efficiently acquire a feeling of
control over actions and their consequences.

Another intriguing finding in our study concerns the relationship between the prospective
sense of agency and global modularity. Global modularity parameters obtained during a task-free
state may be interpreted as describing intrinsic organizational principles that quantify how
efficiently a brain is represented by subnetworks, or modules (Gallen and D'Esposito, 2019).
Previous studies have reported various associations between whole-brain parameters and behavior,
such as between the modular organization and intelligence (Hilger et al., 2017), between inter-
modular connections and awareness (Godwin et al., 2015), and between global efficiency and
short-term memory skills (Gupta et al., 2018). Expanding on these studies, our findings show that
brain-behavior associations may exist at multiple hierarchical levels of the brain architecture.

It should be noted here that modularity measures may suffer from structural constraints (Betzel et al., 2016). This is a direct consequence of the way in which the modularity itself is estimated: the observed intra-modular connections are compared with the expected intra-modular connections, which are estimated using null models (Newman and Girvan, 2004; Bassett et al., 2011). The constraint in this procedure arises because the weight of the null model is usually assumed to be unitary ($\gamma = 1$), essentially constricting the structural resolution of the network (i.e., the size of the modules). In this study, we overcame this limitation by exploring multiple resolution parameters. The results showed that increased values of modularity within medium-high structural resolutions were associated with increased prospective SoA. This interaction was significant with all the regression models that we tested and using both classic and γ-adjusted modularity.

These findings indicate that whole-brain features like modularity may explain high-order mechanisms like contextual predictions elicited by SoA when the brain subsystems (modules) fall between a medium and a small size. Thus, specific resolutions of the modular structure may support specific cognitive functions.

We suggest that the relationship between the sense of agency and global modularity might be explained by the reliance of prospective intentional binding effects on multimodal predictive mechanisms, which constitute a putative general principle of brain function (Wolpert, 1997; Körding et al., 2007; Clark, 2013). In line with this hypothesis, several researchers have proposed a link between predictive coding and the self (Friston, 2012; Kannape and Blanke, 2012; Apps and Tsakiris, 2014). For instance, an essential feature of the self is the ability to generate probabilistic inferences as efferent copies (Feinberg, 1978; Synofzik et al., 2008; Friston and Kiebel, 2009), which are then compared with sensory perceptions (Wolpert et al., 2001; Apps and Tsakiris, 2014). According to internal prediction models, such comparisons are needed to generate prediction errors (Sato and Yasuda, 2005) that favor the adjustment of the motor response (Bédard and Sanes, 2014). The relevance of such considerations is underlined by the crucial importance of the sense of agency for self-awareness (Gallagher, 2000; Taylor, 2014; Ruvolo, 2015; Haggard, 2017). Further research is needed to directly explore the interplay between global brain properties and predictive coding.

The relation between the sense of agency and the integrative functioning of brain networks may also potentially impact clinical contexts. We argue that a suboptimal prospective sense of agency, commonly observed in neuropathological conditions related to distributed cortical abnormalities such as psychosis (Voss et al., 2010; Moore and Fletcher, 2012; de Bézenac et al., 2018) and in altered awareness (Schooler et al., 2014; Berberian et al., 2017), may arise through

two non-exclusive neurophysiological processes. On the one hand, failure of the brain to adapt its
structural resolution to the ongoing behavioral context may favor polluted probabilistic estimates
of action consequences. On the other hand, inaccurate sensorimotor predictions may result from
aberrant intrinsic patterns of nodal functional connectivity in specific brain modules involved in
agency tasks. Our report provides a first neurophysiological whole-brain backbone to
psychological models of a prediction-based sense of agency that can be evaluated in clinical
samples of psychopathology.

Some limitations of this study need to be mentioned. The study focused on implicit
(prospective and retrospective) measures of the sense of agency and their link with network
metrics, ignoring explicit measures of agency. This choice was made on the grounds that explicit
judgments of agency may be affected by cognitive biases (Wegner, 2002; Synofzik et al., 2008),
implicit beliefs (Desantis et al., 2011; Hoogeveen et al., 2018), and attentional biases (Wen et al.,
2016). Consequently, we contend that implicit behavioral measurements, rather than self-report
measures, could be more suitable to start exploring the link between the sense of agency and
intrinsic features of the brain's physiology. Moreover, this study replicates intentional binding
effects from previous investigations of the implicit predictive and retrospective sense of agency
(Voss et al., 2010; Di Plinio et al., 2019a), supporting the robustness of the prospective intentional
binding effect that putatively reflects the implicit sense of agency (Moore and Haggard, 2008;
Moore and Obhi, 2012). This cross-studies robustness should also dispel potential concerns about
the imbalance of the number of trials across conditions. Finally, some general problems related to
studies based on individual differences need to be mentioned, like low experimental control. In
fact, neural features of agency may covary with the other variables like the level of physical fitness
and musical expertise, as suggested by relationships between these features and brain-related
measures (Herting and Nagel, 2013; Kühnis et al., 2014). To limit such potential confounds,
control analyses were performed using both the years of sport training and music expertise as
covariates. These controls showed that the effects reported in the present study were independent
of these factors.

In conclusion, we showed that both whole-brain intra-modular connections and local
(supramarginal) intrinsic inter-modular connections support the prospective sense of agency.
Comparing predictions and intentions with action feedback is an essential feature of human brain
function. This process may be favored by a neural substrate showing, on the one hand, a modular
architecture represented by highly interconnected, adaptably rewiring subsystems and, on the other
hand, efficient exchange of information mediated by motor-control brain systems. The balance of
these two components may predispose to more efficient processing of extrinsic self-information,

while aberrant network behavior may be a potential marker of increased psychosis-like traits (Di

Plinio et al., 2019b). Future studies may consider combining implicit and explicit measures of

agency to distinguish between contextual effects and personal biases and their relationships with

interindividual differences in brain activity and connectivity.

1 REFERENCES

Apps, MA, Tsakiris, M. 2014. The free-energy self: a predictive coding account for self-recognition. Neurosci Biobehav Rev, 41:85-97.

Assal, F, Schwartz, S, Villeumier, P. 2007. Moving with or without will: functional neural correlates of alien hand syndrome. Ann Neurol, 62(3):301-306.

Bandura, A. 2006. Toward a psychology of human agency. Perspect Psychol Sci, 1(2):164-180.

Bassett, DS, Wymbs, NF, Porter, MA, Mucha, PJ, Carlson, JM, Grafton ST. 2011. Dynamic reconfiguration of human brain networks during learning. Proc Natl Acad Sci USA, 108(18):7641-46.

Bédard, P, Sanes, JN. 2014. Brain representations for acquiring and recalling visual-motor adaptations. NeuroImage, 101:225-235.

Berberian, B, Sarrazin, JC, Le Blaye, P, Haggard, P. 2012. Automation technology and sense of control: a window on human agency. PLoS One, 7(3):e34075.

Betzel, R, Medaglia, JD, Papadopoulos, L, Baum, GL, Gur, R, Gur, R, Roalf, D, Satterthwaite, TD, Bassett, DS. 2016. The modular organization of human anatomical brain networks: accounting for the cost of wiring. Net Neurosci, 1(1):42-68.

Buckner, R, Sepulcre, J, Talukdar, T, Krienen, FM, Liu, H, Hedden, T, Andrews-Hanna, JR, Sperling, RA, Johnson, KA. 2009. Cortical hubs revealed by intrinsic functional connectivity: Mapping, assessment of stability, and relation to Alzheimer's disease. J Neurosci, 29(6):1860-1873.

Bullmore, E, Sporns, O. 2009. Complex brain networks: Graph theoretical analysis of structural and functional systems. Nat Rev Neurosci, 10(3):186-188.

Caballero-Gaudes, C, Reynolds, RC. 2017. Methods for cleaning the BOLD fMRI signal. NeuroImage, 154, 128-47.

Caspers, S, Eickoff, SB, Rick, T, von Kapri, A, Kuhlen, T, Huang, R, Shah, NJ, Zilles, K. 2011. Probabilistic fibre tract analysis of cytoarchitectonically defined human inferior parietal lobule areas reveals similarities to macaques. NeuroImage, 58(2):362-380.

Cavanna, AE, Trimble, MR. 2006. The precuneus: A review of its functional anatomy and behavioural correlates. Brain, 129:564-583.

Chambon, V, Wenke, D, Fleming, SM, Prinz, W, Haggard, P. 2013. An online neural substrate for sense of agency. Cereb Cortex, 26(11):1031-1037.

Chambon, V, Moore, JW, Haggard, P. 2015. TMS stimulation over the inferior parietal cortex disrupts prospective sense of agency. Brain Struct Funct, 220(6):3627-3639.

Clark, A. 2013. Whatever next? Predictive brains, situated agents, and the future of cognitive science. Behav Brain Sci, 36(3):181-204.

Cox, RW. 1996. Software for analysis and visualization of functional magnetic resonance neuroimages. Comput Biomed Res, 29(3):162-73.

Desantis, A, Roussel, C, Waszak, F. 2011. On the influence of causal beliefs on the feeling of agency. Conscious Cogn, 20(4):1211-1220.

Desmurget, M, Sirigu, A. 2009. A parietal-premotor network for movement intention and motor awareness. Trends Cogn Sci, 13(10):411-419.

Desmurget, M, Reilly, KT, Richard, N, Szathmari, A, Mottolese, C, Sirigu, A. 2009. Movement intention after parietal cortex stimulation in humans. Science, 324(5928):811-813.

de Bézenac, CE, Swindells, RA, Corcoran, R. 2018. The necessity of ambiguity in self-other processing: a psychosocial perspective with implications for mental health. Front Psychol, 9(2114):1-15.

Di Plinio, S, Ebisch, SJH. 2018. Brain Network Profiling defines functionally specialized cortical networks. Hum Brain Mapp, 39(18):1-18.

Di Plinio, S, Arnò, S, Perrucci, M G, Ebisch, SJH. 2019a. Environmental control and psychosis-relevant traits modulate the prospective sense of agency in non-clinical individuals. Conscious Cogn, 73.

Di Plinio, S, Perrucci, M G, Ebicsch, S J H. 2019b. I am Me: Brain systems integrate and segregate to establish a multidimensional sense of self. Submitted.

Diedrichsen, J, Balster, JH, Flavell, J, Ramnani, N. 2009. A probabilistic MR atlas of the human cerebellum.
NeuroImage, 46(1):39-46.

Farrer, C, Franck, N, Georgieff, N, Frith, CD, Decety, J, Jeannerod, M. 2003. Modulating the experience of agency:
a positron emission tomography study. Neuroimage, 18:324-333.

Feinberg, I. Efference copy and corollary discharge: implications for thinking and its disorders. Schizop Bull,
4:636:640.

Fried, I, Haggard, P, He, BJ, Schurger, A. 2017. Volition and action in the human brain: processes, pathologies, and
reasons. J Neurosci, 37(45):10842-10847.

Friston, K, Kiebel, S. 2009. Predictive coding under the free-energy principle. Philos Trans Royal Soc B, 364:1211-
1221.

Friston, K. 2012. Prediction, perception and agency. Int J Neurophysiol, 83(2):248-252.

Gallagher, S. 2000. Philosophical conceptions of the self: Implications for cognitive science. Trends Cogn Sci,
4(1):14-21.

Gallen, CL, D'Esposito, M. 2019. Brain modularity: A biomarker of intervention-related plasticity. Trends Cogn Sci,
23(4):293-304.

Gallese, V. 2000. The inner sense of action. Agency and motor representations. J Conscious Stud, 7(10):23-40.

Ginestet, CE, Nichols, TE, Bullmore, ET, Simmons, A. 2011. Brain network analysis: Separating cost from topology
using cost-integration. PLoS ONE, 6(7), p.e21570.

Godwin, D, Barry, RL, Marois, R. 2015. Brakdown of the brain's functional network modularity with awareness.
Proc Natl Acad Sci USA, 112(12):3799-3804.

Goldstein, A, Kapelner, A, Bleich, J, Pitkin, E. 2013. Peeking Inside the Black Box: Visualizing Statistical Learning
with Plots of Individual Conditional Expectation. J Comput Graph Statist, 24(1):44-65.

Guimerà, R, Nunes Ameral, LA. 2005. Functional cartography of complex metabolic networks. Nature,
433(7028):895-900.

Gupta, A, Bhushan, B, Behera, L. 2018. Short-term enhancement of cognitive functions and music: A three-channel
model. Sci Rep, 8(1):1-12.

Haggard, P, Clark, S, Kalogeras, J. 2002. Voluntary action and conscious awareness. Nat Neurosci, 5(4):382.

Haggard, P. 2017. Sense of agency in the human brain. Nat Rev Neursoci, 18(4):196.

Hassan, A, Josephs, KA. 2016. Alien hand syndrome. Curr Neurol Neurosci Rep, 16(8):73.

Hascalovitz, A, Sukhvinder, SO. 2015. Personality and intentional binding: An exploratory study using the
narcissistic personality inventory. Front Hum Neurosci, 9(13):1-8.

Hastie, T, Tibshirani, R, Friedman, J. 2009. The Elements of Statistical Learning; Data mining, Inference and
Prediction. New York: Springer.

Herting, MM, Nagel, BJ. 2013. Differences in brain activity during a verbal associative memory encoding task in
high- and low-fit adolescents. J Cogn Neurosci, 25(4):595-612.

Hilger, K, Ekman, M, Fiebach, CJ, Basten, U. 2017. Intelligence is associated with the modular structure of intrinsic
brain networks. Sci Rep, 7(16088):1-12.

Hoogeveen, S, Schjoedt, U, van Elk, M. 2018. Did I do that? Expectancy effects of brain stimulation on error -
related negativity and sense of agency. J Cogn Neurosci, 30(11):1720-1733.

Ito, T, Hearne, L, Mill, R, Cocuzza, C, Cole, MW. 2019. Brain network organization as the computational
architecture of cognition. Preprint available on BioArxiv: https://arxiv.org/abs/1907.03612.

Joliot, M, Jobard, G, Naveau, M, Delcroix, N, Petit, L, Zago, L, Crivello, F, Mellet, E, Mazoyer, B, Tzourio-
Mazoyer, N. AICHA: An atlas of intrinsic connectivity of homotopic areas. 254:46-59.

Kannape, O A, Blanke, O. 2012. Agency, gait and self-consciousness. Int J Psychophysiol, 83(2):191-199.

Kilner, JM, Friston, KJ, Frith, CD. 2007. Predictive coding: An account of the mirror neuron system. Cogn Process,
8(3):159-166.

Kirby, KN, Gerlanc, D. 2013. BootES: an R package for bootstrap confidence intervals on effect sizes. Behav Res
Met, 45(4):905-927.

Körding, KP, Tenenbaum, JB, Shadmehr, R, 2007. The dynamics of memory as a consequence of optimal adaptation to a changing body. Nat Neurosci, 10(6):779-786.

Króliczak, G, Piper, BJ, Frey, SH. 2016. Specialization of the left supramarginal gyrus for hand-independent praxis representation is not related to hand dominance. Neuropsychologia, 93(B):501-512.

Kühnis, J, Elmer, S, Jäncke, L. 2014. Auditory evoked responses in musicians during passive vowel listening are modulated by functional connectivity between bilateral auditory-related brain regions. J Cogn Neurosci, 26(12):2750-2761.

Lancichinetti, A, Fortunato, S. 2009. Community detection algorithms: A comparative analysis. Phys Rev E, 80(056117).

Lancichinetti, A, Fortunato, S. 2012. Consensus clustering in complex networks. Sci Rep, 2(336):1-7.

Lush, P, Roseboom, W, Cleermans, A, Scott, RB, Seth, AK, Dienes, Z. 2019. Intentional binding as Bayesian cue combination: Testing predictions with trait individual differences. J Exp Psychol Hum Percept Perform, 45(9):1206-1217.

Moore, JW, Haggard, P. 2008. Awareness of action: inference and prediction. Conscious Cogn, 17(1):136-144.

Moore, JW, Fletcher, PC. 2012. Sense of agency in health and disease: a review of cue integration approaches. Conscious Cogn, 21, 59-68.

Moore, JW, Obhi, SS. 2012. Intentional binding and the sense of agency: a review. Conscious Cogn, 21(1):546-561.

Newman, MEJ, Girvan, M. 2004. Finding and evaluating community structures in networks. Phys Rev E, 69(2).

Nichols, TE, Das, S, Eickoff, SB, Evans, AC, Glatard, T, Hanke, M, Kriegeskorte, N, Milham, MP, Poldrack, RA, Poline, JB, Proal, E, Thirion, B, Van Essen, DC, White, T, Yeo, BT. 2017. Best practices in data analysis and sharing in neuroimaging using MRI. Nat Neurosci, 20(3):299-303.

Porter, MA, Onnela, J, Mucha, PJ. 2009. Communities in networks. Not Amer Math Soc, 56(9):1082-1097.

Prinz, W. 2012. Open minds: The social making of agency and intentionality. Cambridge: MIT Press.

Rae, CL, Hughes, LE, Weaver, C, Anderson, MC, Rowe, JB. 2014. Selection and stopping in voluntary action: a meta-analysis and combined fMRI study. NeuroImage 86:381-391.

Renes, RA, Aarts, H. 2018. The sense of agency in health and well-being: Understanding the role of the minimal self in action-control. In D de Ridder, M Adriaanse, K. Fujita (Eds.), Routledge international handbooks. The Routledge international handbook of self-control in health and well-being. New York, NY, US: Routledge/Taylor & Francis Group.

Rubinov, M, Sporns, O. 2010. Complex network measures of brain connectivity: uses and interpretations. NeuroImage, 52(3):1050-1060.

Rushworth, MF, Johansen-Berg, H, Göbel, SM, Devlin, JT. 2003. The left parietal and premotor cortices: motor attention and selection. NeuroImage, 20(S1):S89-S100.

Ruvolo, P, Messinger, D, Movellan, J. 2015. Infants time their smiles to make their moms smile. PLoS One 10:e0136492.

Saad, ZS, Gotts, SJ, Murphy, K, Chen, G, Jo, HJ, Martin, A, Cox, RW. 2012. Trouble at rest: how correlation patterns and group differences become distorted after global signal regression. Brain Connect, 2:25-32.

Sato, A, Yasuda, A. 2005. Illusion of sense of self-agency: discrepancy between the predicted and actual sensory consequences of actions modulates the sense of self-agency, but not the sense of self-ownership. Cognition, 94(3):241-255.

Schneider, W, Eschman, A, Zuccolotto, A. 2012. E-Prime 2.0 Reference Guide Manual. Pittsburgh: Psychology Software Tools Inc.

Schooler, JW, Mrazek, MD, Franklin, MS, Baird, B, Mooneyham, BW, Zedelius, C, Broadway, JM. 2014. The middle way, in Psychology of Learning and Motivation, Vol 60. Burlington: Elsevier.

Sun, Y, Danila, B, Josić, K, Bassler, KE. 2009. Improved community structure detection using a modified fine-tuning strategy. Europhys Letters, 86(2):28004.

Synofzik, M, Vosgerau, G, Newen, A. 2008. Beyond the comparator model: A multifactorial two-step account of agency. Conscius Cogn, 17(1):219-239.

Taylor, AH, Cheke, LG, Waismeyer, A, Meltzoff, AN, Miller, R, Gopnik, A, Clayton, NS, Gray, RD. 2014. Of babies and birds: complex tool behaviors are not sufficient for the evolution of the ability to create a novel causal intervention. Proc Royal Soc B, 281:20140837.

Ticini, LF, Dolk, T, Waszak, F, Schütz-Bosbach, S. 2018. IPL-M1 interaction shapes pre-reflective social differentiation in the human action system: new insights from TBS and TMS combined. Sci Rep, 8(12001):1-10.

Traud, AL, Kelsic, ED, Mucha, PJ, Porter, MA. 2011. Comparing community structure to characteristics in online collegiate social networks. SIAM Rev, 53(3):526-543.

van den Heuvel, MP, Sporns, O. 2013. Network hubs in the human brain. Trends Cogn Sci, 17(12):683-696.

van den Heuvel, MP, de Lange, SC, Zalesky, A, Seguin, C, Yeo, BT, Schmidt, R. 2017. Proportional thresholding in resting-state fMRI functional connectivity networks and consequences for patient-control connectome studies: Issues and recommendations. NeuroImage, 152:437-449.

Vocat, R, Staub, F, Stroppini, T, Vuilleumier, P. 2010. Anosognosia for hemiplegia: a clinical-anatomical perspective study. Brain, 133(12):3578-3597.

Voss, M, Moore, J, Hauser, M, Gallinat, J, Heinz, A, Haggard, P. 2010. Altered awareness of action in schizophrenia: a specific deficit in predicting action consequences. Brain, 133(10):3104-3112.

Wang, J, Fan, L, Zhang, Y, Liu, Y, Jiang, D, hang, Y, yu, C, Jiang, T. 2012. Tractography-based parcellation of the human left inferior parietal lobule. NeuroImage, 63(2):641-652.

Wegner, D M. 2002. The illusion of conscious will. Cambridge: MIT press.

Weissenbacher, A, Kasess, C, Gersti, F, Lanzenberger, R, Moser, E, Windischberger, G. 2009. Correlations and anticorrelations in resting-state functional connectivity MRI: a quantitative comparison of preprocessing strategies. NeuroImage, 47:1408-1416.

Wen, W, Yamashita, A, Asama, H. 2016. Divided attention and processes underlying sense of agency. Front Psychol, 7(35):1-8.

Wolpert, DM. 1997. Computational approaches to motor control. Trends Cogn Sci, 1(6):209-216.

Wolpert, DM, Diedrichsen, J, Flanagan, J R. 2011. Principles of sensorimotor learning. Nat Rev Neurosci, 12(12):739-751.

Xia, M, Wang, J, He, Y. 2013. BrainNet Viewer: A network visualization tool for human brain connectomics. PLoS One, 8(7):e68910.

Zou, H, Hastie, T. 2005. Regularization and variable selection via the elastic net. J Royal Stat Soc B, 67(2):310-320.