FACTORS AFFECTING RESPONSE TO RECURRENT GENOMIC SELECTION IN SOYBEANS

1) Vishnu Ramasubramanian

1525, Agronomy Hall, ISU

Ames, IA 50011

Phone: 515 598 6376

Email: vishnur@iastate.edu

2) William Beavis[*]

1208, Agronomy Hall, ISU

Ames, IA 50011

Phone: 800 450 4854 (4412)

Fax: 515 294 730

E-mail: wdbeavis@iastate.edu

Vishnu Ramasubramanian[*][†] **and William D Beavis**[*][†]

[*] Department of Agronomy

[†]Bioinformatics and Computational Biology Graduate Program

Iowa State University, Ames, Iowa, USA - 50010

Factors affecting Response to Recurrent Genomic Selection in Soybeans

Keywords: Genomic Selection, Genetic Response, Recurrent Selection, Soybean, Nested Association Mapping

Corresponding Author:

Vishnu Ramasubramanian

1525, Agronomy Hall, ISU

Ames, IA 50011

Phone: 515 598 6376

Email: vishnur@iastate.edu

Abstract

Herein we report the impacts of applying five selection methods across 40 cycles of recurrent selection and identify interactions among factors that affect genetic responses in sets of simulated families of recombinant inbred lines derived from 21 homozygous soybean lines. Our use of recurrence equation to model response from recurrent selection allowed us to estimate the half-lives, asymptotic limits to recurrent selection for purposes of assessing the rates of response and future genetic potential of populations under selection. The simulated factors include selection methods, training sets, and selection intensity that are under the control of the plant breeder as well as genetic architecture and heritability. A factorial design to examine and analyze the main and interaction effects of these factors showed that both the rates of genetic improvement in the early cycles and limits to genetic improvement in the later cycles are significantly affected by interactions among all factors. Some consistent trends are that genomic selection methods provide greater initial rates of genetic improvement (per cycle) than phenotypic selection, but phenotypic selection provides the greatest long term responses in these closed genotypic systems. Model updating with training sets consisting of data from prior cycles of selection significantly improved prediction accuracy and genetic response with three parametric genomic prediction models. Ridge Regression, if updated with training sets consisting of data from prior cycles, achieved better rates of response than BayesB and Bayes LASSO models. A Support Vector Machine method, with a radial basis kernel, had the worst estimated prediction accuracies and the least long term genetic response. Application of genomic selection in a closed breeding population of a self-pollinated crop such as soybean will need to consider the impact of these factors on trade-offs between short term gains and conserving useful genetic diversity in the context of the goals for the breeding program.

Introduction

Genetic improvement in commercial and public soybean breeding programs include variety development as the evaluation phase of recurrent selection (Orf 2008). Each cycle consists of making crosses among selected lines, creating replicable homozygous lines by self-pollinating progeny for several generations, evaluating the lines in replicated field trials for several years during which poor performing lines are discarded and superior performing lines are retained to cross and begin a new cycle . Pedigrees of modern soybean varieties confirm that genetic improvement of soybeans in Maturity Zones (MZs) II, III and IV has been largely through intra-population recurrent selection (Byrum, personal communication; Hyten et al. 2006; Mikel et al. 2010; Langewisch et al. 2017; Achard et al. 2020). Prior to the Plant Variety Protection Act in 1970 (https://www.ams.usda.gov/rules-regulations/pvpa), a cycle of genetic improvement would require from 10 to 14 years, whereas in the last 40 years commercial organizations have invested in development of continuous nurseries including software that streamlines inventories and logistics of seed transfer, resulting in the capacity to complete a cycle of recurrent selection in five years (Byrum et al. 2017; Anderson et al 2019). Thus, using current best practices, a soybean breeder might experience three to five cycles of genetic improvement.

Despite reducing cycle times to less than ½ of the time required 40 years ago, yield improvements for soybean have not doubled in the corn-soybean agricultural system of the United States (Mikel et al. 2010; Specht et al. 2014). Recognizing the relatively slow yield improvements, farmer-members of the North Central Soybean Research Program (NCSRP) provided funding to public soybean breeders to identify novel QTL in germplasm adapted to MZ's II, III and IV and subsequently to identify methods to increase soybean yields (https://www.ncsrp.com/NCSRP_research.html#yield). The SoyNAM project provided evidence

of novel useful genetic variability for yield among the SoyNAM lines (Song et al. 2017; Diers et

al. 2018) and genomic selection (GS) is being investigated for improving yield in pubic

germplasm.

After the seminal work by Bernardo and Yu (2007), crop breeders have experimentally

demonstrated the relative advantage of GS to phenotypic selection (PS) for a few cycles of

recurrent selection in barley, maize, oats, rice and wheat (Bernardo 2008, 2014; Asoro et al.

2011; Heslot et al. 2012, 2015; Nakaya and Isobe 2012; Emily and Bernardo 2013; Crossa et al.

2014; Beyene et al. 2015; Bassi et al. 2016; Marulanda et al. 2016; Jonas and de Koning 2013,

2016; Hickey et al. 2017; Goiffon et al. 2017). Genomic prediction (GP) has been integrated in

commercial soybean variety development projects as a tool to identify crosses that are most

likely to produce superior progeny and to a lesser extent as a means to reduce the costs of

conducting preliminary field trials (Kurek 2018). However, implementation of GS in soybeans

has not been based on an understanding of how various factors and their interactions can affect

responses to recurrent GS (RGS) in a crop with population structures and genome organizations

such as found in soybean.

Simulations have been used to demonstrate that genome organization, population structure,

genetic architecture, heritability, selection intensity, GP models and composition of training sets

will affect responses to selection (Robertson 1960; Dempfle 1974; Kang 1979; Cockerham &

Burrows 1980; Kang and Namkoong 1980; Kang 1987; Kang and Nienstaedt 1987; Podlich and

Cooper 1998; Cooper et al. 2002; Habier et al. 2007; Goddard 2009; Zhong et al. 2009; Jannink

2010; Bastiaansen et al. 2012; Bijma 2012; de los Campos et al. 2013; Howard et al. 2014; Liu et

al. 2015; Michel et al. 2016). Most of these studies evaluated impacts of subsets of factors on

responses across a few cycles of recurrent selection, although a couple of studies investigated

limits of responses to RGS across more than 20 cycles (Jannink 2010; Liu et al. 2015).

Comparisons of genomic prediction (GP) models have revealed that simulated genetic

architecture and proportion of phenotypic variance due to the genotypic variance affected

prediction accuracies and simulated responses to selection (de los Campos et al. 2010, 2013;

Wimmer et al. 2013; Howard et al. 2014). Parametric methods such as Ridge Regression (RR)

and Bayesian regression in the mixed effects modeling framework provide more accurate

predictions and greater gains for traits with additive genetic architecture, whereas non-parametric

machine-learning methods such as Neural Networks and Support Vector Machines provide more

accurate predictions for traits with epistatic genetic architectures (Howard et al. 2014). Prediction

accuracies are essentially the same for all GP models applied to data with additive genetic

architectures (Long et al. 2010, 2011; Guo et al. 2012; Howard et al. 2014). Given additive

genetic architectures, RR (Endelman 2011) and Bayesian models (Pérez and de los Campos,

2014) provide similar short-term genetic gains, but different responses after ~ 15 cycles of RGS.

Marker densities impact prediction accuracies and short term genetic gains, with a dense marker

set performing better than a sparse set, although there are limits to improvements from increased

marker densities that depend on linkage disequilibrium (LD) and structure of the breeding

population (de Roos et al. 2009; Schulz-Streeck et al. 2012; Hickey et al. 2014; Xavier et al,

2016; Norman et al. 2018). Decreased prediction accuracies of GP models in later cycles of

recurrent GS are associated with decay of LD between marker loci (ML) and quantitative trait

loci (QTL), loss of relationships between lines in early and later cycles of selection or a

combination of both (Habier et al. 2007; Zhong et al. 2009; Hickey et al. 2014; Liu et al. 2015;

Müller and Melchinger 2017, 2018). It is possible to offset the loss of relationships and LD, by

updating the training sets with each cycle of selection (Jannink 2010; Liu et al. 2015; Müller et

al. 2017, 2018). If training data are from only the current cycle of selection, then predictions do

not take into account relationships between the current population and the founder population or

possible loss of LD. At the other extreme, data from all prior cycles of selection can be included

with data from the current cycle in the training set, but practical computation limits will be

encountered with large training sets.

With the development of male sterile and insect pollination systems for soybean, it will become

possible to conduct one to three cycles of RGS per year (Ortiz-Perez 2008; Davis, personal

communication). This will enable increased selection intensities and greater immediate genetic

gains from GS because much larger numbers of progeny can be evaluated without the

commensurate added expense of increased numbers of field plots (Heslot et al. 2015). However,

it is well-established that increased selection intensities will reduce the genetic potential of a

breeding population through loss of useful genetic variability and ultimately will limit responses

to selection (Robertson 1960; Hill and Robertson 1968; Bulmer 1970).

Emerging technologies, such as genome editing, promise to provide novel useful genetic

variability for crops that do not have natural genetic variability for traits of interest (Brooks et al.

2014; Jenko et al. 2015; Hickey et al. 2017; Rodríguez-Leal et al. 2017; Lemmon et al. 2018).

However, before assuming that genome editing will enable genetic improvement after loss of

useful genetic variance from selection, it is important to recognize that expression of novel

genetic modifications within genetic networks depend on the genomic background, even in

simple model organisms such as yeast (Peccoud et al. 2004; Forsberg et al. 2017; Hou et al.

2018). Since RGS with one to multiple cycles per year (Gaynor et al. 2017) will become feasible

for soybean and because genome editing will need genetic diversity for success, soybean

breeders need to determine appropriate trade-offs between short term genetic gains and retention

of useful genetic variability in the germplasm that will be inherited by future plant breeders

(Rodríguez-Leal et al. 2017; Lemmon et al. 2018).

In anticipation of emerging technologies that will enable soybean breeders to implement a two

part strategy for genetic improvement (Gaynor et al. 2017), we investigated the impacts of

selection methods (PS and GS), training sets (TS), number of QTL, heritability (H), selection

intensity (SI) and their interactions on responses to RGS. We employed a factorial treatment

design, where each combination of factor levels was replicated ten times in simulated RGS and

RPS of recombinant inbred lines (RILs) across 40 cycles. Outcomes reported herein provide a

foundation for comparisons with strategies that use other types of progeny for selection and

crossing as well as practical guidelines for understanding trade-offs between genetic gains in

early cycles and retention of useful genetic potential for future genetic improvement of soybeans.

Methods

Simulations and Treatment Design. The impact of nQTL, SI, h, TS and SM on response to

selection across 40 cycles of recurrent selection were evaluated with a factorial design consisting

of 306 combinations of factors. In contrast to evaluating one or two variables at a time, the

factorial design is widely used to comprehensively determine factors that will optimize processes

(Myers 1976; Collins et al. 2014; Dunn 2020). Specifically the treatments consisted of three

values for number of simulated QTL, three selection intensities, two values for non-genetic

variance, five selection methods and four types of TS used to update four GP models (Table 1).

In summary the treatment design consists of 18 combinations of factors for PS plus 288

combinations of factors for GS methods for a total of 306 combinations of factors. Note that TS

are irrelevant for PS. Each set of factor combinations was replicated with ten simulated recurrent

selection projects across 40 cycles resulting in 3060 simulations with 122400 outcomes.

Simulated soybean RILs were generated by crossing *in silico* 20 homozygous SoyNAM founder

lines with IA3023 to generate 20 distinct F_1 progeny. The F_1 progeny from each of the 20 crosses

were self-pollinated *in silico* for five generations to generate 100 RILs per family.   The resulting

2000 RILS from 20 families were assessed for segregating genotypic information at 4289 SNP

loci (Song et al. 2017; Xavier A et al. 2017; Diers B et al. 2018). On average alleles from the

common founder occurred at a frequency of 0.9 and alleles at the loci from all other founder

lines occurred at a frequency of 0.1. Soybeans has been subjected to phenotypic selection and

genetic improvement for several thousand years (Anderson et al. 2019). Thus, the SoyNAM

founders represent a realized population structure adapted to maturity zones II, III and IV that is

akin to a single coalescent process used in many simulation studies (Daetwyler et al. 2013;

Bandillo et al. 2017).

Of the 4289 SNP markers with genotypic scores for the SoyNAM population 3818 were

polymorphic among the 20 families used as founders for the simulations. On average, 773 were

polymorphic for a family with a standard deviation among families of 34. In the initial founding

set of RILs, the average heterozygosity per SNP locus across 20 families was 0.09. 'Gst' is a

measure of sub-population differentiation estimated as ratio of difference between expected

heterozygosity of sub- populations to total expected heterozygosity. The average estimated G_{st}

value across the genome for the initial founding set of RILs was 0.32, as determined by the

'diff_stats' function in the mmod R package (Jombart 2008; Ryman and Leimar 2009; Jombart

and Ahmed 2011). Relative to previous reported founders that used a coalescent process

(Woolliams and Corbin 2012; Hickey and Gorjanc 2012; Daetwyler HD et al. 2013), our

simulations began with a structure more likely to be found in actual soybean breeding

populations and with less, albeit more realistic, genetic diversity of soybeans adapted to maturity

zones II, III and IV. We've also estimated Pairwise 'Fst' using 'pairwise.fst' in 'hierfstat' R

package (Goudet 2005), which is a measure of population differentiation among pairs of

populations. It is estimated as the ratio of difference between the average of the expected

heterozygosity of the two populations and total expected heterozygosity of the pooled

populations to total expected heterozygosity of the pooled populations. Average Fst among the

20 families in simulated SoyNAM data is 0.20. Whereas the average Fst using genotypic data

from SoyNAM project among 40 families is 0.09 with a maximum pairwise Fst of 0.15 and a

minimum Fst of 0.007. Average Fst among the clusters in USDA soybean germplasm collection

is 0.22 -0.23 (Song et al 2015; Xavier 2018).

Subsets of 40, 400, and 4289 SNP loci were designated as QTL. The QTL were distributed

evenly throughout the genome, and each contributed equal additive effects of 5/-5, 0.5/-0.5, or

0.05/-0.05 units respectively to the total genotypic value of the simulated RILs. Thus, all three

genetic architectures had the same potential to create genotypic values ranging from +200 to -

200 in the initial founder sets of RILs. Positive and negative allelic effects were simulated to

alternate sequentially at QTL that are uniformly distributed across the Soybean genetic map.

Because all marker alleles are QTL alleles when there are 4289 QTL, LD between marker alleles

and QTL will not deteriorate across cycles of selection and recombination. Phenotypic values

were simulated by adding non-genetic variance sampled from an N $(0, \sigma)$ distribution to the

simulated genotypic values, where σ was determined by the heritability on an entry mean basis

among the initial sets of founder sets of RILs. Broad sense heritability on an entry mean basis

(H) values of 0.7 and 0.3 were simulated for each of the three sets of QTL. After the phenotypic

values were simulated in the initial founding RILs, the non-genetic variance was held constant

across subsequent cycles of selection.

For each cycle of recurrent selection, 1%, 2.5% or 10% of the most positive phenotypic or

predicted phenotypic values among 2000 simulated RILs, corresponding to selection intensities

of 2.67, 2.34 and 1.75, were selected as lines to inter-mate for the next cycle. Matings among

pairs of selected RILs used a design with greater contributions from the best selected lines

(Figure 1) also known as networked families designs (Guo et al. 2013; Guo et al. 2014).

Based on previous results from Howard et al (2014), four GS methods were evaluated. Ridge

Regression using restricted maximum likelihood (RR-REML) represented a frequentist

parametric model. Bayes-B (BB) and Bayesian LASSO (BL) represented parametric bayesian

models and Support Vector Machine with Radial Basis Kernel (SVM-RBF) represented a non-

parametric method of machine learning.

Ridge regression was implemented with a method that employs expectation maximization to

obtain Restricted Maximum Likelihood estimates of marker effects (Xavier et al. 2019). This

computational method is faster than the popular implementation of ridge regression in rrBLUP

package (Endelman 2011) and generates values that are highly correlated with the predictions

based on the rrBLUP package (Figure S1). The BGLR package (Perez and de los Campos 2014)

provided implementations of BB and BL models. The 'Rgtsvm' package in R was used for its

implementation of the SVM with RBF kernel method (Wang et al. 2017). 'Rgtsvm' implements

SVM training on GPUs with computing time several hundred times less than that required for the

implementation in the 'caret' package on high performance computing clusters, and produces

similar prediction accuracies and estimates of mean squared errors (Figure S2). The parameters

used to train GP models with R packages are provided in Table 2.

For purposes of this manuscript we use the phrase 'model updating' to refer to retraining GP

models with up to 14 previous cycles of training data (Figure S3). A preliminary analysis of TS

on genotypic values and prediction accuracies was conducted using RR-REML models trained

with data from the current cycle as well as 3, 5, 8, 10, 12, and 14 prior cycles. The results were

compared with responses from the RR-REML model updated with TS's comprised of data from

all prior cycles and a RR-REML model with no updating. Training sets for each cycle were

obtained by randomly sampling 1600 RILs from the set of 2000 simulated RILs in each cycle.

The most accurate prediction and maximum genetic response was obtained with training data

that is cumulatively added every cycle (Figure S4 and S5). The results indicate that 3-5 prior

cycles of training data did not significantly improve prediction accuracies and responses relative

to models that were not updated. Also, the standardized genotypic values and prediction

accuracies, obtained using 10 to 14 prior cycles of data in the TS's, were not significantly

different than results based on TS's consisting of all prior cycles. Based on the results of this

preliminary study, we investigated responses to recurrent selection using TS's consisting of up to

14 prior cycles of selection as well as data from the current cycle. After the 14th cycle, training

data consisted of 14 prior cycles of recurrent selection.

Modeled response to recurrent selection. The averaged genotypic value for each cycle, c, of

recurrent selection was modeled with a linear first order recurrence equation:

$$f_0(\text{c})\,y_{(\text{c}+1)} + f_1(\text{c})\,y_{(\text{c})} = g(\text{c}) \qquad \text{(Eqn 1)}$$

226   Where c is a sequence of integers from 0 to 39 representing each cycle of recurrent selection

227   from cycle 1 to 40 and f_0, f_1 and g are constant functions of c. By rearranging the equation we

228   note that the response in cycle c+1 can be represented as

229
$$y_{(c+1)} = -\frac{f_1(c)}{f_0(c)} y_{(c)} + \frac{g(c)}{f_0(c)} \qquad \text{(Eqn 2)}$$

230   Since the ratios $f_1(c)/f_0(c)$ and $g(c)/f_0(c)$ are constants, we can represent the response in cycle c+1

231   as

232
$$y_{(c+1)} = \alpha y_{(c)} + \beta \qquad \text{(Eqn 3)}$$

233   If y_0 specifies the average genotypic value of the first cycle of RILs derived from the founders,

234   then (3) has a unique solution (Goldberg 1958):

235
$$y_c = \alpha^c y_0 + \beta \frac{1-\alpha^c}{1-\alpha} \text{ if } \alpha \neq 1 \qquad \text{(Eqn 4)}$$
$$y_c = \alpha^c y_0 + \beta c \qquad \text{if } \alpha = 1$$

236   An alternative representation of (4) for the situation of $\alpha \neq 1$ is

237
$$y_c = \alpha^c (y_0 - y') + y'$$
$$with \ y' = \frac{\beta}{1-\alpha},$$

, where α is less than 1 for genotypic response to recurrent selection and y' represents the

asymptotic limit to selection (Goldberg 1958). To illustrate, values of the sequence of c=0 to 39,

with $y_0 = 0$, for the range of α (0.6- 0.9) and β (1.4-38) values are plotted in Figure 2. The curves

can be interpreted as response to selection as a function of the frequencies of alleles with additive

selective advantage, selection intensity, time and effective population size (Robertson 1960).

The parameters, y_0, α, and β, were estimated with a non-linear mixed effects method

implemented in 'nlme' functions in the 'nlme' and 'nlshelper' packages (Pinheiro and Bates

2000; Baty et al. 2015; Pinheiro et al. 2019).

Since the limits of responses are asymptotic, the number of cycles before there is no longer

response to selection is referred to as the half-life (Robertson 1960; Dempfle 1974; Kang 1979;

Cockerham & Burrows 1980; Kang and Namkoong 1980; Kang 1987; Kang and Nienstaedt

1987). From the first order recurrence equation, the half-life is estimated as $\ln(0.5)/\ln(\alpha)$, when

y_0 is '0' and the asymptotic limit is estimated as y'.

Analyses of variance (ANOVA) *of modeled response to recurrent selection.* The purpose of the

ANOVA is to evaluate the impact of factors and their interactions on the modeled responses to

recurrent PS and GS methods. The analyses of variance used single and multi-level nlme models

with modeled (eqn 4) responses grouped by treatment factors. The influence of multiple factor

treatment combinations on estimated non-linear mixed effect models have not been implemented

in standard statistical software packages that report the analysis of variance in terms of sums of

squares and traditional 'F-tests'. For discussions on the challenges of using standard F-test for

non-linear mixed effects (nlme) models see (Pinheiro et al. 2000; Baty et al. 2015; Pinheiro et al.

2019). Consequently, we analyzed the variance among modeled responses using AIC, BIC and

Likelihood metrics that were grouped based on combinations of factors consisting of selection

methods, TS, SI, nQTL and simulated H.

In order to provide a balanced data table for analyses by the non-linear mixed effect model,

responses that included PS, which has no TS's, were assumed constant resulting in a balanced

full factorial set of responses for 360 combinations of factors. The process of fitting, selecting

and refining mixed effects models closely followed the steps described in Pinheiro et al. 2000;

Zuur 2009 and Oddi et al. 2019). The complete process used in the study is provided in File S1.

In the first phase of model fitting, estimates of modeled parameters from nlsList models were

retained as starting values for fixed effects. Both alpha and beta were fit only for intercept and

deviations from estimated means conditioned on grouping variables were modeled as random

effects using the 'nlme' R package. Multiple ANOVA of 'nlme' objects representing the models

were used to identify combinations of factors with significant effects on the non-linear response

model. The model with the lowest AIC score was selected as the best model. The best random

intercept model in the first phase of model fitting process M31 in Table S1 was further refined

by modelling the correlation structure.

Evaluation of Simulated Response to Recurrent Selection

While the modeled genotypic values are evaluated using half-life and asymptotic limits, we have

evaluated the simulated outcomes from recurrent selection using a set of metrics to assess

responses, population characteristics, and GP model performance every cycle of selection.

The standardized genotypic value, R_s (eqn 5), was estimated every cycle as the change in

genotypic value from the average genotypic value of 2000 RILs derived from the initial founders

and standardized to the maximum genotypic potential (200 units) among the founders

(Meuwissen et al. 2001; Liu et al. 2015).

283
$$R_s = \frac{R_c}{(R_m - R_0)} \qquad \text{(Eqn 5)}$$

284
R_s - Standardized genotypic value
R_0 - Average genotypic value of RILs produced by founders
R_c - Average genotypic value in cycle c – R_0
R_m - Maximum possible genotypic value (=200)

The most positive genotypic value (M_{gv}) among RIL's selected in cycle c is a metric used to

evaluate the best RIL produced each cycle, while the standardized genotypic variance (SV_g)

defined as the estimated genotypic variance divided by the estimated genotypic variance of the

initial population, was used to evaluate the loss of genotypic variance. Note that values for the

SV_g range from zero to one.

Response standardized to change in standard deviation of genotypic values captures genetic gain

with respect to loss of genetic variance (eqn 6). The numerator term represents change in

genotypic values of a population in cycle 'c' from cycle '0' founder population normalized to

standard deviation of genotypic values in cycle '0'. The denominator term represents change in

standard deviation of genotypic values from cycle '0' to cycle 'c' as a fraction of standard

deviation of genotypic values in cycle 0. This metric is similar to the metric used to refer to

efficiency of converting loss of genetic diversity to genetic gain of a selection method in

recurrent selection (Gorjanc et al. 2018). While efficiency is estimated as slope in linear

regression model with numerator as 'y' term and denominator as 'x' term in the linear part of

response curve, with Rs_Var it is possible to visualize both linear and non-linear sections of the

response curve.

$$Rs\_var = \frac{G_c - G_0}{SdG_0 - SdG_c} \qquad (Eqn\ 6)$$

303
G_c -average genotypic value of the set of RILs evaluated in cycle c
G_0 -average genotypic value of the founding set of RILs
SdG_0 - estimated standard deviation from genotypic values of founding set of RILs
SdG_c - estimated standard deviation from genotypic values of RILS in cycle c

Estimated Linkage disequilibrium (*LD*) among pairs of marker loci on all 20 chromosomes was

evaluated as the deviation of observed gametic frequency of alleles at a pair of loci from the

product of the individual allele frequencies, assuming independence (Weir 1996). The R function

'get_PG_LD_StatsGSMethods' used to estimate pairwise LD between markers is provided in the

R package 'SoyNAMPredictionMethods'. GP models were assessed using the estimated

prediction accuracies (r_{ps}), defined as the estimated linear correlation (Pearson) between

predicted and simulated phenotypic values and the estimated Mean Squared Error (MSE), and

defined as the mean of the squared deviations of the predicted phenotypic values from the

simulated values.

Analyses and Data Availability

More information on the analyses can be found in the R package 'SoyNAMPredictionMethods'.

Also, simulated data and code are available as part of the package (File S1 found at

http://gfspopgen.agron.iastate.edu/SoyNAMPredictionMethods_v2_2020.html). Supplemental

material including the R package can be found at https://figshare.com/s/f7d4e515e563beef550b

and http://gfspopgen.agron.iastate.edu/SoyNAMPredictionMethods_v2_2020.html describes

how to use the package. The complete process for fitting NLME models and ANOVA can be

found at http://gfspopgen.agron.iastate.edu/Vishnu_MS1/NLME_Models_Part_I.html .SoyNAM

genotypic and phenotypic data are available in SoyBase (Grant et al. 2010).

Results

Prediction accuracies in the founding sets of RILs: Estimates of prediction accuracies, r_{ps}, of GP

models trained with the initial set of 2000 F_5-derived RILs ranged from 0.75-0.82 for H of 0.7

and ranged from 0.38 - 0.49 for h of 0.3 (Figure 3). The initial r_{ps} for both H values was best

with BB and poorest with the SVM-RBF. The nQTL had little effect on r_{ps} within either value of

0.7 or 0.3 for H. RR-REML and BL had smaller magnitude MSE values than BB and SVM RBF

for all numbers of simulated QTL and both values for H (Figure 3). Accuracies are lower when

GP models are trained without QTL in the training set, but follow a similar pattern as the models

with QTL. MSE are greater or comparable for models trained without QTL than for models with

QTL (Figure S6- S7). Average within family prediction accuracies are lesser than prediction

accuracies from a combined TS comprising of RILs from all the families (Figure S8-S9).

However, a combined TS will have (n*population size of family) for 'n' families and estimated

accuracies will have confounding effects from training set size. Estimated accuracies for models

trained with TS generated by random sampling from a combined population to keep the TS size

same as family size are lower than average within family accuracies. MSE are lesser for

combined TS than for models trained using within family TS and sampled TS (Figure S8-S9).

Influence of factors on response metrics. The averaged responses Rs were modeled (eqn 4) and

the results are consistent with theory (Figure 2). Averaged across all simulations there was rapid

increase of Rs across the first five cycles of selection followed by slower responses from cycles 5

to 10 and no response after cycle 20.

While there are observable general trends for each of the individual factors, response metrics are

unique for each combination of all factors (Figure 4-7, S10 - S14). The most parsimonious

model requires unique estimates of α, and β (eqn 4) for each of the combinations of factors

indicating that interactions among all factors have significant influences on the responses (Table

S1). Also analyses of variances on subsets of 10 and 20 cycles of selection demonstrate that the

interactions were important from the earliest cycles (File S2: Table S1, S2). Further, the relative

importance of factors on interaction effects were consistent (nQTL>SM>SI>H>TS) and

significant in analyses of variance conducted on subsets of 10 and 20 selection cycles (Table S1).

Among the three factors SM, TS and SI that are under the control of the plant breeder, we

consider SM as the primary factor of interest, whereas TS and SI are considered secondary

factors of interest that significantly modulate the effect of SM on response. In addition, nQTL

and H that are not under the control of the plant breeder are noted for their large and significant

impacts on response metrics (Table S1).

Relative to PS, the three parametric GP models provided greater initial rates of response, reduced

population half-life and faster loss of genetic variance. Whereas, selection using the SVMRBF

consistently produced the least effective responses (Figure 4-7). Training sets consisting of data

from up to 14 prior cycles of selection, compared to TS's consisting of data from only the current

cycle of selection are observably distinctive for both the short term and long term response

metrics (Figure 4-7). Importantly, use of the TS's significantly improved responses to selection

in both the short term and long term for all other combinations of factors (Table S1). Selection

intensity is the third most important factor to affect the response metrics (Table S1). Consistent

with theory (Robertson 1960; Hill and Robertson 1968; Bulmer 1971,1976), greater SI's

(associated with retention of smaller numbers of RILs to initiate subsequent cycles, were

associated with more rapid response rates, shorter half-lives, faster loss of genetic variance and

significantly lower Rs values as the population approached it's limit to respond to selection.

The modeled responses were highly dependent on each of the factors, with nQTL showing the

greatest deviations from the average response (Table S1). From the modeled responses the half-

life for a responsive population varied from 1.6 to 4.9 cycles for 40 QTL, whereas for 400 and

4289 QTL it ranged from 2.2 to 8.8 and 3 to 9.5 cycles respectively. The modeled responses

were highly correlated with the simulated responses (Pearson correlation coefficient: 0.94-0.97).

To illustrate the impact of nQTL, consider Rs values plotted across forty cycles of recurrent

selection (Figure 4 and S15). If the genetic architecture of the trait consists of 40 and 400 QTL,

responses to selection were limited after 10-15 cycles of selection, whereas for 4289 QTL, limits

to selection responses were not realized until 30 to 40 cycles of selection.

Next consider that the maximum realized response for 40 simulated QTL was from 0.32 to 0.78

of maximum genotypic potential among the founders, whereas for 400 and 4289 QTL, the

realized response was 0.12 - 0.42 and 0.04-0.17 of the maximum (Figure 4 and S15). Also if

there are 40 simulated QTL, the maximum attained values are as high as 80% of the maximum

value of 200 in less than ten cycles of recurrent selection (Figure 8 and S16). In contrast, Rs

values are no greater than 40% of the maximum value and stop responding to selection in 10-15

cycles if there are 400 simulated QTL. Rs values were never greater than 15% of the maximum

value and only begin to approach a limit after 20 cycles if there are 4289 simulated QTL.

As expected, responses to selection reflect declining genetic variances (Figure 9 and S17). The

loss of Sgv's across cycles is much faster with fewer simulated QTL than larger numbers of

QTL. Likewise the estimated prediction accuracies (Figure 10 and S18) approach zero as the

genotypic variance approaches zero. Average MSE for the GP models increase across cycles of

selection (Figure 11 and S19) LD among markers approach zero as the genotypic variance

approaches zero although the covariance among these response metrics depend on the other

simulated factors (Figure S20-S24).

To interpret the role of nQTL as a factor, it is important to recall that: 1) positive and negative

allelic effects were simulated to alternate sequentially at QTL (marker loci) that were distributed

across the genome according to the Soybean genetic linkage map. 2) Crossing nearly

homozygous lines and subsequently self-pollinating progeny for five generation before

genotyping and phenotyping within each cycle creates a limited number of large linkage blocks.

Analyses of the number of linkage blocks each generation reveals that regardless of the number

of QTL, the number of linkage blocks per cycle ranges from 70-90. It was not the same blocks

each cycle, but if the nQTL equals 40, then each linkage block included all segregating QTL

each cycle. For 400 and 4289 simulated QTL, each linkage block had a net genetic effect of zero

or +/- 0.5 or +/- 0.05 multiplied by the number of QTL in the block respectively. Thus, the nQTL

might be better understood as the magnitude of genetic effects associated with segregating

linkage blocks.

The observable general trend for H, or perhaps more accurately understood as contributions of

non-genetic effects to the phenotypes, was that H values of 0.7 in the initial phenotypic variance

resulted in Rs values that were greater than H values of 0.3 of the initial phenotypic variance

(Figure 4 and S15). The trend in Rs values is correlated with the other response metrics, in

particular prediction accuracies of the GP models. The loss of estimated prediction accuracies

are greater with H values of 0.3 than 0.7 with relaxed selection intensities (Figure 10 and S18).

Other combinations of SI and H require model updating to provide reasonable GP model

prediction accuracies and achieve greater responses across more cycles of selection. As we

would expect, for all combinations of SI and nQTL, loss of genotypic variance are greater with H

values of 0.7 than 0.3 (Figure 9 and S17).

Some specific outcomes of interest: For purposes of illustrating interaction effects on SM's

across all 40 cycles of selection, consider the most relaxed SI of 1.75, associated with selecting

10% of the RILs per cycle. When GP models are not updated, BB produced greater R_s values

than PS in the early cycles for all nQTL and both levels of H, whereas PS resulted in greater

responses than all GS methods after the 10^{th} cycle (Figure 6, S15 and S25). SVMRBF did not

demonstrate any better responses than PS in either early or late cycles for any nQTL or level of

H (Figure 6, S15, and S25; File S3 and S4).

If the parametric GP models are updated with training sets consisting of data from up to 14 prior

cycles of recurrent selection, responses to RR-REML demonstrated the greatest responses (Rs)

for 40, 400 and 4289 QTL across both levels of H (Figure 7, S15 and S26; File S3 and S4). If

the RR-REML model is updated with up to 14 prior cycles of training sets, responses are larger

than PS for up to 10-40 cycles depending on the number of QTL, H and SI (Figure 4 and 7; File

S3 and S4). When BB and BL GP models are updated, responses are larger than PS for up to 5,

20 and 40 cycles for 40, 400, and 4289 QTL respectively. Similar, albeit distinctive, comparisons

among outcomes from GP models with model updating for genetic architectures responsible for

0.3 of the phenotypic variance in the initial sets of RILs (Figure S26) are described in File S4.

Relative to responses without model updating application of RR-REML and Bayesian methods

with model updating resulted in greater responses. Model updating with Bayesian methods also

resulted in less favorable responses than the RR-REML (Figure S27 and S28; File S4).

SVMRBF when updated with TS's demonstrated no significant improvement relative to

SVMRBF without updating for all genetic architectures, levels of H and SI's (Figure S27 and

S28; File S4). If the genetic architecture explains only 30% of the phenotypic variability in the

initial sets of RILs, the relative improvements in Rs values across cycles using updated TS's are

better than simulated QTL that explain 70% of the phenotypic variance (Figure S28). Percentage

gain in responses for GS with model updating relative to response from GS without model

updating are provided in File S5.

In terms of lost genetic potential, every cycle of selection reduced the maximum possible genetic

value. When GP models are not updated, the genetic potential is lost at a rapid rate beginning in

the early cycles, whereas when GP models are updated, genetic potential is retained in the

population and genetic variance decreases at a slower rate. PS had the least loss in genetic

potential relative to all four GP models without updating. However, with model updating, the

loss of genetic potential using parametric GP models was almost the same as PS. Among the

parametric GP models, RR-REML and Bayesian methods showed similar slow losses of genetic

potential with and without model updating. SVMRBF GS had the greatest loss of genetic

potential beginning with the early cycles (Figure 5).

The loss of genetic potential in early cycles determines the limits to selection response in later

cycles. For example, with 400 simulated QTL responsible for 70% of the phenotypic variance,

the maximum potential was only 50% of the maximum potential (100 units) for PS and

parametric GS methods with and without model updating. When parametric GP models are

updated, 81% and 75% of the limits of maximum available potential are realized with RR-REML

and Bayesian methods respectively. If GP models are not updated, only 62% of the limits of the

maximum available potential are realized with RR-REML and Bayesian methods. With PS, 78%

of maximum available potential is realized by the cycles in which the population no longer

responds to selection. With SVMRBF, only 35 % of the potential is realized with and without

updating by the cycles in which the population no longer responds to selection (Figure 5).

If GP models are not updated with data from up to 14 prior cycles, the Mgv's were consistently

greater with PS than the four GS methods. Among GP models without updating, BB provided

the best Mgv, while SVM-RBF had the smallest Mgv (Figure 8 and S16). If GP models are

updated, the pattern depends mostly on the number of QTL. For initial H values of either 0.7 and

0.3 and 40 simulated QTL, Mgv's are similar for RR-REML, Bayesian GP models and PS,

whereas for 400 QTL, RR-REML produces greater Mgv's than PS and Bayesian methods. For

4289 QTL, RR-REML and Bayesian methods produce greater Mgv's than PS. Recurrent GS

with SVMRBF produced the least desirable Mgv's for 40, 400 and 4289 QTL.

If GP models are updated, the standardized genotypic variance (Sgv) decreases at a rate similar

to GP models that are not updated (Figure 9 and S17). There is no difference among GS methods

in terms of rate at which Sgv decreases. Also, model updating significantly improved estimated

prediction accuracies, $r_{ps,}$ for all GP models except SVMRBF. Among RR-REML and Bayesian

GP models, model updating has a slightly larger impact on estimated accuracies and MSE using

RR-REML than with Bayesian GP models (Figure 10, 11, S18 and S19). MSE were orders of

magnitude lesser for RR-REML than bayesian GP models with updates after the first 10-15

cycles of selection (Figure 11 and S19).

If models are updated using data from up to 14 prior cycles, the changes to genetic variance

among the RILs selected to be crossed, their average heterozygosity, average rate of inbreeding,

and loss of favorable alleles are similar among GS methods (Figure S29- S32). Model updating

resulted in faster loss of genotypic variance among genotypes selected to be parental lines for the

next cycle of inter-mating. The loss of genotypic variance is similar among parametric GS

methods. (Figure S29). When there are 400 simulated QTL responsible for 70% of the

phenotypic variance, the average number of favorable alleles that are lost across 40 cycles due to

selection and drift are similar among PS and GS methods, but the rate at which they are lost

differs among selection methods for the first 20 cycles until they converge at the same limit.

SVMRBF GS showed the greatest rate of loss and PS had the least rate of loss while the

parametric GS methods had intermediate rates of loss. By the time 40 cycles of selection have

completed, model updating didn't result in any significant difference in rates of loss and the total

number of favorable alleles that are lost (Figure S30).

SVMRBF GS showed the greatest loss of average heterozygosity and PS lost heterozygosity at

the slowest rate, while RR-REML and bayesian GS methods lost heterozygosity at intermediate

rates. Model updating didn't result in significant changes to rates at which heterozygosity was

lost over cycles (Figure S31). PS showed slower rates of inbreeding than GS methods as we

would expect from the decay of standardized genotypic variance. Average rates of inbreeding

were similar among parametric GP models in the early cycles of recurrent selection, whereas the

patterns varied after there is no response to selection (Figure S32).

For all selection methods pairwise LD among markers on the same chromosome decreased

across cycles of recurrent selection (Figure S20-S24). LD decreased slowest with PS (Figure

S20). Loss of LD in early and late cycles of selection are similar among parametric GP models

and SVMRBF with the relaxed selection intensity. By the 20^{th} cycle of recurrent selection, LD

approached zero for all selection methods and there was no evidence that selection methods

affected linkage disequilibrium (LD) differentially in the earlier cycles. The rates at which LD

decays are lower when GP models are updated with training sets compared to GP models

without updating (Figure S20-S24).

The limiting values for RsVar (Response standardized to change in genotypic variance) when PS

is used to select the best 10% of RILS with genetic architectures consisting of 400 and 4289

QTL are greater than the limiting values using GS methods without model updates (Figure S33

and S34). The parametric GP models, without model updating, resulted in similar changes of

RsVar for 40, 400 and 4289 simulated QTL responsible for both 70% and 30% of phenotypic

variability in the initial population. Also, if the GP models are not updated, the rates and limits to

loss of RsVar are similar among the GS methods for all nQTL and SI.

If GP models are updated with the TS's, the patterns of RsVar are significantly different among

GS methods and are dependent on nQTL, SI and H (Figure S35 and S36). With 0.7 heritability,

there are no significant difference in RsVar among GS methods for 40 simulated QTL. If the

genetic architecture consists of 400 and 4289 QTL and weaker selection intensities are practiced,

the RR-REML GS method maintained genetic variance and RsVar for more cycles than PS and

the Bayesian GS methods. Relative Gain in RsVar with RR-REML GS is even larger for 0.3 H

treatment with relaxed selection intensities (Figure S35 and S36). SVMRBF demonstrated the

least limits of RsVar for treatment combinations with and without model updating (Figure S33 –

S36). The plots for all the evaluation metrics discussed above for selection intensities 2.67 and

2.34 are provided in Figure S37 –S60 and discussed in File S6).

Discussion

Previous publications of *in silico* investigations of factors affecting outcomes from RGS have

been conducted using a few factors applied to arbitrary diploid genomes, and an expected

population structure from an assumed coalescent process. To our knowledge, the research

reported herein is the first designed to reveal interactions among five factors that previously had

been shown to affect responses to selection. Our ability to detect and characterize interaction

effects is enabled by use of a first order recurrence equation (eqn 4). Hopefully, our explanation

of how to implement recurrence equation models in available R packages will encourage others

to investigate recurrent genetic improvement designs. Even though our motivation for the use of

non-linear modeling in this study was restricted to the systematic investigation of significance of

variation in response due to the factors and the relative order of magnitude of interaction effects

of factors, non-linear mixed effects models have predictive power that is still relatively

unexploited in the study of limits due to recurrent selection. We plan to investigate the use of

non-linear recurrence modelling in mixed model framework to estimate the magnitude of

interaction effects and optimize the use of genomic information at prediction and selection levels

to optimize response in recurrent selection schemes in the future.

Most *in silico* investigations of GS have simulated diploid genomes consisting of ten or fewer

linkage groups with evenly spaced molecular markers depending on the organism (Jannink 2010;

Liu et al. 2015; Akdemir and Sánchez 2016; Yabe et al. 2016). The organization of the soybean

genome is the result of ancient duplication events (Shoemaker et al. 1996; Grant et al. 2000;

Cannon and Shoemaker 2012) and consists of many copies of functional genes that are

distributed unevenly with high concentrations near the telomeric ends of 20 diploid

chromosomes. Also, most prior simulation studies created population structures using an

assumed coalescent process (Woolliams and Corbin 2012; Hickey and Gorjanc 2012; Daetwyler

et al. 2013). We used a realized structure representing adapted and selected lines by sampling the

SoyNAM founders.

Structures of plant populations are highly dependent on the reproductive biology. Indeed, plant

breeders design genetic improvement projects based on reproductive biology. Similar to many

cereal and pulse crops, the reproductive biology of soybean is primarily through self-pollination

(Wilcox et al. 1979; Fehr 1980, 1991). The frequency of natural cross pollination in soybean is

only about 0.025 (Garber et al. 1925; Caviness 1966; Carlson and Lersten 1987; Ahrent and

Caviness 1994). Because crossing soybean lines is labor intensive and expensive

(https://www.youtube.com/watch?v=VnjGijF4KQI) soybean breeder's use mating designs in

which only one or two elite varieties are crossed with a few dozen recently selected RIL's (Guo

et al. 2013; Guo et al. 2014). We refer to this as a hub network design. Soybean breeders

subsequently take advantage of natural self-pollination to create RILs with sufficient seed for

replicated evaluations across many environments.

Our simulations attempted to emulate cycles of selection, crossing, and self-pollination currently

conducted by commercial and academic soybean breeders. Relative to outcrossing species

selecting homozygotes to participate in a hub network within each cycle will have smaller

effective populations and retain LD for more cycles of recurrent selection. Future studies are

needed to determine if the significant interaction effects that we found are due to population

structure and genome organization.

Implications for application of GS for genetic improvement with current soybean breeding.

There has been considerable concern expressed about the limited genetic variability among

soybean genotypes adapted to specific maturity zones and recurrently selected over 75 years in

North America (Carter et al. 2004; Hyten et al. 2006; Mikel et al. 2010; McCouch et al. 2013).

The SoyNAM founders represent a sample of current improved breeding germplasm adapted to

and used for agricultural production in maturity zones II - IV (Song et al. 2017; Diers et al.

2018).

Despite concerns about limited genetic variability, our assessment of the half-lives of

populations suggest that even if only 40 QTL remain in the founders of the SoyNAM population,

there is still genetic potential for response to selection for at least five cycles. If there are larger

numbers of QTL with smaller additive effects distributed among linkage blocks of SoyNAM

founders, then we can expect reasonable half-lives to RGS for 10 cycles (File S2). Because

current soybean breeding methods require four to five years before RIL's are selected to

participate in crossing nurseries, soybean breeders can expect reasonable responses to selection

using samples from the SoyNAM population for 20 to 60 years. It is possible to obtain

reasonable estimates of half-lives and asymptotic limits using the parameters in eqn 4 with

sufficient number of simulated cycles of selection. The estimated half-lives and asymptotic limits

can be used to assess the impact of selection on genetic potential of populations in the long term.

In our simulations, the estimates of half-lives using 10 or 20 simulated cycles, are significantly

less accurate than estimates obtained with 40 simulated cycles of recurrent selection (Table S2).

Among the five factors we investigated, a soybean breeder can choose SM's, SI's and TS's.

Currently, plant breeders have little control on nQTL and H, because these parameters are

determined by the sample of germplasm and environments. While it is possible to adjust the

value of H on an entry mean basis by increasing/decreasing the number of replicates (Fehr 1991),

estimating the number of segregating QTL and the magnitude of their effects is difficult and

usually extremely expensive (Beavis 1994; Goring et al. 2001; Xu 2003). For a fixed budget, the

breeder will be faced with a trade-off between numbers of replicates and numbers of RIL's that

can be evaluated. In other words, H on an entry mean basis can be estimated, but not adjusted

without adding field plot resources, and nQTL and their contributions to genetic variance can be

estimated, but the estimates will be biased.

Although modulated by nQTL and H, PS consistently retained useful genetic variability across

many cycles of genetic improvement, Bayesian methods provided the fastest genetic gains in the

short term and RR-REML provided a compromise between PS and Bayesian methods. SVM-

RBF should not be considered for genetic improvement for the additive genetic architectures in

soybean population structures. Previously others have reported that long-term response using

GS methods will be more limited than PS in closed populations (Goddard 2009; Zhong et al.

2009; Jannink 2010). We also observed that GS without updated TS's result in rapid loss of

genetic variance in the initial cycles, which results in lower Rs values as responses to selection

approach an asymptotic limit. When models were updated with TS's composed of data from

prior cycles of selection, loss of prediction accuracy slowed for all values of nQTL and H

(Figure 10).

Given similar rates of decreasing genetic variance among parametric GS methods, the different

limits to selection response is possibly due to greater efficiency of retaining genetic diversity

with RR-REML for later cycles, although for many combinations of factors the limits of

response with RR-REML and Bayesian methods are about the same. Unlike previous studies,

we noted in the initial cycles of RGS genetic gains and estimates of accuracy were similar using

BL and RR-REML, whereas after 15 cycles genetic responses were not as limited with RR-

REML, probably because additive genetic variance was retained for later cycles of selection (Liu

et al. 2015).

Replicated responses to high values of SI quickly reach a limit in five to ten cycles of recurrent

selection. Replicated responses to lower values of selection intensity consistently result in greater

gains over more cycles, indicating that genetic drift is the most likely mechanism for loss of

genotypic variance. These constraints on plant breeding programs are well characterized

(Brisbane and Gibson 1995; Hayes et al. 2009; Jannink 2010; Hung et al. 2012; Liu et al. 2015;

Akdemir and Sánchez 2016; Yabe et al. 2016).

Model updating with TS's from prior cycles improves the relationship between training sets and

validation sets and thus improves responses to GS without updated training sets. While model

updating doesn't significantly change the estimated half-life, model updating did produce greater

responses standardized to the rate at which genotypic variance is lost in selected populations with

updated RR-REML GP models (File S2; Table S3).

The choice of which combinations of SM, SI and TS depend on the objectives of the breeding

program. If the objective is to enter and capture market share in a short time, then maintenance

of genetic diversity is not important. Averaged over nQTL and H, the greatest changes in Rs

values, i.e., rate of genetic gain, were attained in early cycles using BayesB and SI = 2.84,

without model updating. Thus, if a soybean breeding project wants to maximize the rate of

genetic gain for a single quantitative trait in a population derived from a sample of the SoyNAM

founders, then application of the BayesB GS method and large SI is the best combination to meet

the breeding objective. If the breeding project has a long term breeding objective to improve

germplasm while maintaining useful genetic diversity for purposes such as providing useful

germplasm for future generations or evaluating genome editing, then PS or GS with RR-REML

with relaxed SI will be the best combination to meet the breeding objective. The greatest values

for Rs were attained using the RR-REML model with model updating and SI = 1.75.   Last if the

breeding project has multiple objectives for both immediate and longer term goals, then pareto

optima among tradeoffs involving responses/cycle and retention of useful genetic variance for

multiple traits need to be identified (Akdemir et al. 2019).

Lessons for future simulation studies. In our simulations we assigned the same amount of time

to develop and evaluate RILs for GS as for PS. Application of PS usually requires field trials for

three years before lines are selected for use in a crossing nursery. In practice, one of the

advantages of GS relative to PS is that only subsets of RILs need to be phenotyped. Commercial

soybean breeding projects have used genotypic values obtained from GP models to cull lines in

as they are being self-pollinated and prior to the first stage of field trials (Kurek 2018). Also it is

possible to train or update GP models with lines derived in earlier filial generations thereby

requiring less time per cycle (Bassi et al. 2016). Even if both GS and PS require the same amount

of time to develop lines before phenotypic evaluation, GS methods can be conducted as soon as

phenotypic information is available from first year field trials (Heffner et al. 2009), while PS

usually requires phenotypic evaluations for three years before lines are selected for crossing

nurseries to begin a new cycle. Herein we did not investigate the trade-offs between less

accurate predictions from models trained with less extensive phenotyping or phenotyping with

lines derived in earlier filial generations. These practical adjustments from application of GS

methods to current soybean genetic improvement projects still need to be investigated.

Accuracies of GP models in the founding set of RILs were similar to that reported in previous

studies (Long et al. 2010, 2011; Guo et al. 2012; Howard et al. 2014). However, our estimates of

accuracies are larger as we've included QTL in our training sets. Relationship among selected

RILs and LD between marker loci (ML) and QTL are considered the two most important sources

of prediction accuracy. In previous studies, relationship among RILs had a greater effect on

prediction accuracies for RR-REML than BL, whereas accuracy of BL was more dependent on

LD and both components showed similar contribution to the accuracy of BayesB GP models

(Habier et al. 2007; Zhong et al. 2009; Liu et al. 2015). Similar to our results without model

updating, Bayesian GS methods resulted in greater responses as the populations approached

asymptotic limits (Meuwissen 1997; Li et al. 2008; Akdemir and Sánchez 2016). However, GP

model updating reduced the difference in rate of decrease of prediction accuracies among RR-

REML and Bayesian GP models but there was no consistent pattern in relationship among

selected RILs and rate of loss for LD to explain the observed estimates of prediction accuracies.

However, GP model updating consistently resulted in lesser MSE for RR-REML than BB and

BL GP models across all levels of SI, nQTL and H. This pattern is consistent with greater

efficiency of converting loss of variance into gain with updated RR-REML GS method. In order

to estimate the contribution of LD and linkage blocks to prediction accuracy of GP models will

require a design similar to that employed by Müller et al. (2017, 2018) for synthetic populations.

Populations with unrelated training and prediction sets with LD and SNP based relationship

estimates showed low prediction accuracy and low genetic response in recurrent GS similar to

GS without updating in this study. Whereas populations with relationship between training and

prediction sets with LD and SNP based relationship estimates showed greater prediction

accuracy and greater genetic response similar to GS with model updating (Müller et al. 2017,

2018).

While TS's had relatively small impacts on response metrics, they were highly significant. Since

relatedness of TS's and validation sets affect estimates of prediction accuracy it has been

suggested that model updating needs to be evaluated for accuracies of prediction within and

across populations (Crossa et al. 2014; Juliana et al. 2018; Stewart-Brown et al. 2019). Herein,

the TS's were comprised of a combined population of RIL's derived from all families across

cycles of selection. Given a constant number of RILs in the TS's, within or across family

prediction accuracies will depend on genetic differentiation among families (de Roos et al. 2009;

Schulz-Streeck et al. 2012; Stewart-Brown et al. 2019). However, actual soybean breeding

projects evaluate a few to many dozen RIL's per family and future simulation studies, especially

of two part systems, should consider whether relationships between evaluation units, RIL's and

selection units (Holland et al. 2003), possibly individuals, need to be used in design of the TS's.

We allowed the size of TS's to increase every cycle by adding data from prior cycles. Increasing

the number of RIL's per TS requires more computational resources. An alternative strategy is to

randomly sample subsets of data from each of the prior cycles to maintain a constant cumulative

training population size. It is also possible to assign weights to the samples from prior cycles to

place more weight on data from more recent cycles. These possibilities suggest determination of

an optimal combination of numbers of RIL's and weights that will provide maximum prediction

accuracy with minimal computational requirements. Some aspects of this optimization problem

have been addressed (Lorenz et al. 2013; Hickey et al. 2014; Akdemir et al. 2015; Xavier et al.

2017). For example, Akdemir et al (2015) devised a genetic algorithm for selecting optimal

training populations to minimize prediction error variance and Xavier et al (2017) developed

sampling methods for training Bayesian GP models. Another consideration is whether TS's need

to be updated every cycle. Instead of updating the model with data from every cycle would it be

more effective to retrain GP models every second third, fourth… cycle while maintaining a

constant training population size? Modifications to design of TS's will need to be addressed with

simulations before implementation.

Like most simulation studies we fixed levels of SI as constant for all combinations of factors

across all cycles of recurrent selection. This is not consistent with actual soybean breeding

projects. Just as sampling families result in some with exceptional genotypes and some with

poor genotypes, in an actual genetic improvement project there is variability among cycles. The

effects of applying a dynamic selection strategy is an alternative and interesting question. We

hypothesize that a strategy consisting of applying different SI's, optimized for each cycle, will

achieve improved long-term genetic response by differentially emphasizing genetic variance and

genetic gain across multiple cycles of selection.

Unlike actual soybean genetic improvement projects we simulated a closed breeding population

derived from a sample of SoyNAM founders in which culled lines were not resampled for

discarded favorable alleles, nor did we simulate exchange of lines among breeding projects. In

MZ's II, III and IV there are six public breeding projects and about a dozen commercial soybean

breeding projects. Thus, there is potential, depending on material transfer agreements, to

exchange genotypes among breeding populations. In our next set of investigations migration

among multiple breeding projects will be evaluated for response to recurrent selection within and

among breeding projects using island model evolutionary algorithms (Hagan et al. 2012; Yabe et

al. 2016). Results reported herein will provide comparators for assessing impacts of migration

policies relative to the other factors that affect responses to RGS.

Also, as with prior simulation studies, we simulated truncation selection, but unlike previous

investigations we did not randomly mate selected lines each cycle. Rather, we used the hub

network design (Guo et al. 2013; Guo et al. 2014). We did not consider relationships among

selected RILs nor the trade-offs between genetic gain, genetic variance (inbreeding) when

selecting RILs to cross. There exist multi-objective optimization methods such as genomic

mating and optimal cross selection (Rutten et al. 2002; Woolliams et al. 2015; Akdemir and

Sánchez 2016; Gorjanc et al. 2018; Akdemir et al. 2019) that have been demonstrated to provide

both greater rates of genetic gain and assure maintenance of population genetic potential across

cycles.

In the future, it is possible that development of male sterile and insect pollination systems for

soybean (Ortiz-Perez 2008; Davis 2020) will enable a cycle of RGS to be conducted using two or

three mating generations per year. This will enable an alternative genetic improvement system

based on decoupling genetic improvement from variety development (Gaynor et al. 2017). By

separating the two types of breeding projects, GS can be applied continuously. In the two part

system TS's will need to be composed of genotypic and phenotypic data obtained from annual

field trials of RIL's, although the TS's will be several selection cycles removed from the cycle

used to create the RILs.

Results reported herein suggest that RGS in a two part system will rapidly produce genetic gains

and loss of useful genetic variance in very short periods of time. Indeed, 40 cycles of RGS could

be completed in 12 to 20 years. To offset the shorter cycle times, application of GP models to

select and cross individual progeny instead of RIL's could result in a larger effective population

size per cycle by creating more opportunities for recombination and slow the unintended loss of

valuable alleles in discarded linkage blocks. However, before investments in development of

male sterile and insect pollination systems for soybean (Ortiz-Perez 2008; Davis, personal

communication) research using simulations are needed to understand the trade-offs and whether

the investments are justified using approaches from Operations Research (Xu et al. 2011;

Cameron et al. 2017, Han et al. 2017).

Acknowledgements

Funding for this research was provided by the Department of Agronomy, Iowa State University,

the North Central Soybean Research Program and an NSF grant (1830478). Supplemental

funding for large scale computing was enabled by the Extreme Science and Engineering

Discovery Environment (XSEDE), which is supported by National Science Foundation. XSEDE

resources consisted of research allocations (DMS190015 & DMS190018) on PSC-Bridges Large

Memory nodes for the simulations involving updating of parametric GP models. The Iowa State

University -Pronto GPU cluster enabled computation of SVM model update simulations. We

want to acknowledge Matheus de Krause for discussions on fitting non-linear models using

'nlme' package, Dr. Lizhi Wang for efficient programs to simulate meiosis and Dr. Alencar

Xavier for sharing an efficient expectation maximization method for fitting ridge regression GP

models. Last, we want to thank anonymous critical reviewers of earlier versions of this

manuscript for providing useful suggestions on how to present such a large volume of

information from such a comprehensive investigation of expected short and long term outcomes

of GS applied to soybeans adapted to MZ's II to IV.

Reference

Achard F, Butruille M, Madjarac S, Nelson PT, Duesing J, Laffont JL, Nelson B, Xiong J, Mikel
MA, Smith JSC: **Single nucleotide polymorphisms facilitate dus testing of soybean cultivars**
for plant variety protection. *Crop Science* 2020, **n/a**.

Ahrent DK, Caviness CE: **Natural Cross-Pollination of Twelve Soybean Cultivars in**
Arkansas. *Crop Science* 1994, **34:**376-378.

Akdemir D, Beavis W, Fritsche-Neto R, Singh AK, Isidro-Sánchez J: **Multi-objective**
optimized genomic breeding strategies for sustainable food improvement. *Heredity* 2019,
122:672-683.

Akdemir D, Sánchez JI: **Efficient Breeding by Genomic Mating.** *Frontiers in Genetics* 2016, **7**.

Akdemir D, Sanchez JI, Jannink JL: **Optimization of genomic selection training populations**
with a genetic algorithm. *Genet Sel Evol* 2015, **47:**38.

Anderson EJ, Ali ML, Beavis WD, Chen P, Clemente TE, Diers BW, Graef GL, Grassini P,
Hyten DL, McHale LK, et al: **Soybean [Glycine max (L.) Merr.] Breeding: History,**
Improvement, Production and Future Opportunities. In *Advances in Plant Breeding*

Strategies: Legumes: Volume 7. Edited by Al-Khayri JM, Jain SM, Johnson DV. Cham: Springer International Publishing; 2019: 431-516

Asoro FG, Newell MA, Beavis WD, Scott MP, Jannink J-L: **Accuracy and Training Population Design for Genomic Selection on Quantitative Traits in Elite North American Oats.** Iowa State University Digital Repository; 2011.

Bassi FM, Bentley AR, Charmet G, Ortiz R, Crossa J: **Breeding schemes for the implementation of genomic selection in wheat (Triticum spp.).** *Plant Science* 2016, **242:**23-36.

Bastiaansen JWM, Coster A, Calus MPL, van Arendonk JAM, Bovenhuis H: **Long-term response to genomic selection: effects of estimation method and reference population structure for different genetic architectures.** *Genetics Selection Evolution* 2012, **44:**3.

Baty F, Ritz C, Charles S, Brutsche M, Flandrois J-P, Delignette-Muller M-L: **A Toolbox for Nonlinear Regression in R: The Package nlstools.** *Journal of Statistical Software; Vol 1, Issue 5 (2015)* 2015.

Beavis WD: **The power and deceit of QTL experiments: lessons from comparative QTL studies.** In *Proceedings of the forty-ninth annual corn and sorghum industry research conference.* Chicago, IL; 1994: 250-266.

Bernardo R: **Molecular Markers and Selection for Complex Traits in Plants: Learning from the Last 20 Years.** *Crop Science* 2008, **48:**1649.

Bernardo R: **Genomewide Selection of Parental Inbreds: Classes of Loci and Virtual Biparental Populations.** *Crop Science* 2014, **54:**2586.

Bernardo R, Yu J: **Prospects for Genomewide Selection for Quantitative Traits in Maize.** *Crop Science* 2007, **47:**1082-1090.

Beyene Y, Semagn K, Mugo S, Tarekegne A, Babu R, Meisel B, Sehabiague P, Makumbi D, Magorokosho C, Oikeh S, et al: **Genetic gains in grain yield through genomic selection in eight bi-parental maize populations under drought stress.(RESEARCH)(Author abstract).** 2015, **55:**154.

Bijma P: **Long-term genomic improvement – new challenges for population genetics.** *Journal of Animal Breeding and Genetics* 2012, **129:**1-2.

Brisbane J, Gibson J: **Balancing selection response and rate of inbreeding by including genetic relationships in selection decisions.** *International Journal of Plant Breeding Research* 1995, **91:**421-431.

Brooks C, Nekrasov V, Lippman ZB, Van Eck J: **Efficient gene editing in tomato in the first generation using the clustered regularly interspaced short palindromic repeats/CRISPR-associated9 system.** *Plant Physiol* 2014, **166:**1292-1297.

Bulmer MG: **The Effect of Selection on Genetic Variability.** *The American Naturalist* 1971,
105:201-211.

Bulmer MG: **The effect of selection on genetic variability: a simulation study.** *Genet Res*
1976, **28:**101-117.

Byrum J, Beavis B, Davis C, Doonan G, Doubler T, Kaster V, Mowers R, Parry S: **Genetic Gain**
Performance Metric Accelerates Agricultural Productivity. *Interfaces* 2017, **47:**442-453.

Cameron JN, Han Y, Wang L, Beavis WD: **Systematic design for trait introgression projects.**
Theoretical and Applied Genetics 2017, **130:**1993-2004.

Cannon SB, Shoemaker RC: **Evolutionary and comparative analyses of the soybean genome.**
Breeding science 2012, **61:**437-444.

Carlson JB, Lersten NR: **Reproductive Morphology. In Soybeans: Improvement,**
Production, and Uses. 2004**:**59-95.

Carter Jr TE, Nelson RL, Sneller CH, Cui Z: **Genetic diversity in soybean.** *Soybeans:*
Improvement, production, and uses 2004, **16:**303-416.

Caviness CE: **Estimates of Natural Crosspollination in Jackson Soybeans in Arkansas1.**
Crop Science 1966, **6:**cropsci1966.0011183X000600020034x.

Cockerham CC, Burrows PM: **Selection limits and strategies.** *Proceedings of the National*
Academy of Sciences of the United States of America 1980, **77:**546.

Collins LM, Dziak JJ, Kugler KC, Trail JB: **Factorial Experiments: Efficient Tools for**
Evaluation of Intervention Components: Efficient Tools for Evaluation of Intervention
Components. *American Journal of Preventive Medicine* 2014, **47:**498-504.

Cooper M, Podlich D, Micallef K, Smith O, Jensen N, Chapman S, Kruger N: **Complexity,**
quantitative traits and plant breeding: a role for simulation modelling in the genetic
improvement of crops. *Quantitative genetics, genomics and plant breeding'(Ed MS Kang) pp*
2002**:**143-166.

Crossa J, Pérez P, Hickey J, Burgueño J, Ornella L, Cerón-Rojas J, Zhang X, Dreisigacker S,
Babu R, Li Y, et al: **Genomic prediction in CIMMYT maize and wheat breeding programs.**
Heredity 2014, **112:**48-60.

Daetwyler HD, Calus MPL, Pong-Wong R, de Los Campos G, Hickey JM: **Genomic prediction**
in animals and plants: simulation of data, validation, reporting, and benchmarking.
Genetics 2013, **193:**347-365.

De Los Campos G, Gianola D, Rosa GJM, Weigel KA, Crossa J: **Semi-parametric genomic-**
enabled prediction of genetic values using reproducing kernel Hilbert spaces methods.
Genetics Research 2010, **92:**295-308.

852 de los Campos G, Hickey JM, Pong-Wong R, Daetwyler HD, Calus MPL: **Whole-Genome**
853 **Regression and Prediction Methods Applied to Plant and Animal Breeding.** *Genetics* 2013,
854 **193:**327-345.

855 de los Campos G, Naya H, Gianola D, Crossa J, Legarra A, Manfredi E, Weigel K, Cotes JM:
856 **Predicting Quantitative Traits With Regression Models for Dense Molecular Markers and**
857 **Pedigree.** *Genetics* 2009, **182:**375-385.

858 de Los Campos G, Vazquez AI, Fernando R, Klimentidis YC, Sorensen D: **Prediction of**
859 **complex human traits using the genomic best linear unbiased predictor.** *PLoS Genet* 2013,
860 **9:**e1003608.

861 de Roos APW, Hayes BJ, Goddard ME: **Reliability of genomic predictions across multiple**
862 **populations.** *Genetics* 2009, **183:**1545-1553.

863 Dempfle L: **A note on increasing the limit of selection through selection within families.**
864 *Genet Res* 1974, **24:**127-135.

865 Diers BW, Specht J, Rainey KM, Cregan P, Song Q, Ramasubramanian V, Graef G, Nelson R,
866 Schapaugh W, Wang D, et al: **Genetic Architecture of Soybean Yield and Agronomic Traits.**
867 *G3: Genes|Genomes|Genetics* 2018, **8:**3367.

868 Dunn K: **Process Improvement using Data.** In.; 2015

869 Emily C, Rex B: **Accuracy of Genomewide Selection for Different Traits with Constant**
870 **Population Size, Heritability, and Number of Markers.** *The Plant Genome* 2013, **6**.

871 Endelman JB: **Ridge Regression and Other Kernels for Genomic Selection with R Package**
872 **rrBLUP.** *The Plant Genome Journal* 2011, **4:**250.

873 Fehr W: *Principles of cultivar development: theory and technique.* Macmillian Publishing
874 Company; 1991.

875 Fehr WR, Hadley HH: *Hybridization of crop plants.* American Society of Agronomy and Crop
876 Science Society of America; 1980.

877 Forsberg SK, Bloom JS, Sadhu MJ, Kruglyak L, Carlborg Ö: **Accounting for genetic**
878 **interactions improves modeling of individual quantitative trait phenotypes in yeast.** *Nat*
879 *Genet* 2017, **49:**497-503.

880 Garber RJ, Odland T, McIlvaine T, Quisenberry K: *Varietal Experiments with Soybeans.*
881 Agricultural Experiment Station, College of Agriculture, West Virginia …; 1925.

882 Gaynor RC, Gorjanc G, Bentley AR, Ober ES, Howell P, Jackson R, Mackay IJ, Hickey JM: **A**
883 **Two-Part Strategy for Using Genomic Selection to Develop Inbred Lines.** *Crop Science*
884 2017, **57:**2372-2386.

885 Goddard M: **Genomic selection: prediction of accuracy and maximisation of long term**

response. *Genetica* 2009, **136:**245-257.

Goiffon M, Kusmec A, Wang L, Hu G, Schnable PS: **Improving Response in Genomic**
Selection with a Population-Based Selection Strategy: Optimal Population Value Selection.
Genetics 2017, **206:**1675.

Goldberg S: *Introduction to difference equations, with illustrative examples from economics,*
psychology, and sociology. New York: New York, Wiley; 1958.

Göring HH, Terwilliger JD, Blangero J: **Large upward bias in estimation of locus-specific**
effects from genomewide scans. *The American Journal of Human Genetics* 2001, **69:**1357-
1369.

Gorjanc G, Gaynor RC, Hickey JM: **Optimal cross selection for long-term genetic gain in**
two-part programs with rapid recurrent genomic selection. *Theoretical and Applied Genetics*
2018, **131:**1953-1966.

GOUDET J: **hierfstat, a package for r to compute and test hierarchical F-statistics.**
Molecular Ecology Notes 2005, **5:**184-186.

Grant D, Cregan P, Shoemaker RC: **Genome organization in dicots: Genome duplication in**
Arabidopsis and synteny between soybean and Arabidopsis. *Proceedings of the National*
Academy of Sciences 2000, **97:**4168-4173.

Grant D, Nelson RT, Cannon SB, Shoemaker RC: **SoyBase, the USDA-ARS soybean genetics**
and genomics database. *Nucleic acids research* 2010, **38:**D843.

Guo B, Sleper DA, Beavis WD: **Nested Association Mapping for Identification of Functional**
Markers. *Genetics* 2010, **186:**373-383.

Guo B, Wang D, Guo Z, Beavis WD: **Family-based association mapping in crop species.**
Theoretical and Applied Genetics 2013, **126:**1419-1430.

Guo Z, Tucker DM, Basten CJ, Gandhi H, Ersoz E, Guo B, Xu Z, Wang D, Gay G: **The impact**
of population structure on genomic prediction in stratified populations. *Theoretical and*
Applied Genetics 2014, **127:**749-762.

Guo Z, Tucker DM, Lu J, Kishore V, Gay G: **Evaluation of genome-wide selection efficiency**
in maize nested association mapping populations. *Theoretical and Applied Genetics* 2012,
124:261-275.

Guzman C, Peña RJ, Singh R, Autrique E, Dreisigacker S, Crossa J, Rutkoski J, Poland J,
Battenfield S: **Wheat quality improvement at CIMMYT and the use of genomic selection on**
it. *Applied & Translational Genomics* 2016, **11:**3-8.

Habier D, Fernando RL, Dekkers JCM: **The Impact of Genetic Relationship Information on**
Genome-Assisted Breeding Values. *Genetics* 2007, **177:**2389.

Hagan S, Knowles J, Kell DB: **Exploiting Genomic Knowledge in Optimising Molecular**
Breeding Programmes: Algorithms from Evolutionary Computing (Evolutionary
Computing for Molecular Breeding). 2012, **7:**e48862.

Han Y, Cameron JN, Wang L, Beavis WD: **The Predicted Cross Value for Genetic**
Introgression of Multiple Alleles. *Genetics* 2017, **205:**1409.

Hayes BJ, Visscher PM, Goddard ME: **Increased accuracy of artificial selection by using the**
realized relationship matrix. *Genetics Research* 2009, **91:**47-60.

Heffner EL, Sorrells ME, Jannink J-L: **Genomic Selection for Crop Improvement.** *Crop*
Science 2009, **49:**1.

Heslot N, Akdemir D, Sorrells ME, Jannink J-L: **Integrating environmental covariates and**
crop modeling into the genomic selection framework to predict genotype by environment
interactions. *Theoretical and Applied Genetics* 2014, **127:**463-480.

Heslot N, Jannink J-L, Sorrells ME: **Perspectives for Genomic Selection Applications and**
Research in Plants. *Crop Science* 2015, **55:**1-12.

Heslot N, Yang H-P, Sorrells ME, Jannink J-L: **Genomic Selection in Plant Breeding: A**
Comparison of Models. *Crop Science* 2012, **52:**146-160.

Hickey JM, Chiurugwi T, Mackay I, Powell W: **Genomic prediction unifies animal and plant**
breeding programs to form platforms for biological discovery. *Nature genetics* 2017,
49:1297.

Hickey JM, Dreisigacker S, Crossa J, Hearne S, Babu R, Prasanna BM, Grondona M, Zambelli
A, Windhausen VS, Mathews K, Gorjanc G: **Evaluation of Genomic Selection Training**
Population Designs and Genotyping Strategies in Plant Breeding Programs Using
Simulation. *Crop Science* 2014, **54:**1476-1488.

Hickey JM, Gorjanc G: **Simulated data for genomic selection and genome-wide association**
studies using a combination of coalescent and gene drop methods. *G3 (Bethesda, Md)* 2012,
2:425.

Hill WG, Robertson A: **The effect of linkage on limits to artificial selection.** *Genetics*
Research 2008, **89:**311-336. (First published in 1968)

Holland J, Nyquist WE, Cervantes-Martinez CT: **Estimating and interpreting heritability for**
plant breeding: Anupdate. *Plant breeding reviews* 2003, **22:**9-111.

Hou J, van Leeuwen J, Andrews BJ, Boone C: **Genetic Network Complexity Shapes**
Background-Dependent Phenotypic Expression. *Trends in genetics : TIG* 2018, **34:**578-586.

Howard R, Carriquiry AL, Beavis WD: **Parametric and nonparametric statistical methods**
for genomic selection of traits with additive and epistatic genetic architectures. *G3*
(Bethesda) 2014, **4:**1027-1046.

Howard R, Carriquiry AL, Beavis WD: **Application of Response Surface Methods To**
Determine Conditions for Optimal Genomic Prediction. *G3: Genes|Genomes|Genetics* 2017,
7:3103.

Hung HY, Browne C, Guill K, Coles N, Eller M, Garcia A, Lepak N, Melia-Hancock S,
Oropeza-Rosas M, Salvo S, et al: **The relationship between parental genetic or phenotypic**
divergence and progeny variation in the maize nested association mapping population.
Heredity (Edinb) 2012, **108:**490-499.

Hyten DL, Song Q, Zhu Y, Choi IY, Nelson RL, Costa JM, Specht JE, Shoemaker RC, Cregan
PB: **Impacts of genetic bottlenecks on soybean genome diversity.** *Proc Natl Acad Sci U S A*
2006, **103:**16666-16671.

Jannink J-L: **Dynamics of long-term genomic selection.** *Genetics Selection Evolution* 2010,
42:35.

Jannink JL, Lorenz AJ, Iwata H: **Genomic selection in plant breeding: from theory to**
practice. *Brief Funct Genomics* 2010, **9:**166-177.

Jenko J, Gorjanc G, Cleveland MA, Varshney RK, Whitelaw CBA, Woolliams JA, Hickey JM:
Potential of promotion of alleles by genome editing to improve quantitative traits in
livestock breeding programs. *Genetics Selection Evolution* 2015, **47:**55.

Jombart T: **adegenet: a R package for the multivariate analysis of genetic markers.**
Bioinformatics 2008, **24:**1403-1405.

Jombart T, Ahmed I: **adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.**
Bioinformatics 2011, **27:**3070-3071.

Jonas E, de Koning D-J: **Does genomic selection have a future in plant breeding?** *Trends in*
Biotechnology 2013, **31:**497-504.

Jonas E, de Koning DJ: **Goals and hurdles for a successful implementation of genomic**
selection in breeding programme for selected annual and perennial crops. *Biotechnology*
& genetic engineering reviews 2016, **32:**18.

Juliana P, Singh RP, Poland J, Mondal S, Crossa J, Montesinos-López OA, Dreisigacker S,
Pérez-Rodríguez P, Huerta-Espino J, Crespo-Herrera L, Govindan V: **Prospects and Challenges**
of Applied Genomic Selection—A New Paradigm in Breeding for Grain Yield in Bread
Wheat. *The Plant Genome* 2018, **11:**180017.

Kang H: **Limits of artificial selection under balanced mating systems.** 1979.

Kang H: **Limits of artificial selection under balanced mating systems with family selection.**
Silvae genetica 1983, **32:**188-195.

Kang H, Namkoong G: **Limits of artificial selection under unbalanced mating systems.**
Theoretical and Applied Genetics 1980, **58:**181-191.

Kang H, Nienstaedt H: **Managing long-term tree breeding stock.** *Silvae genetica* 1987, **36:**30-
39.

Kurek A: **"Phenotypic and genomic selection for multi-trait improvement in soybean line**
and variety development" 2018.

Langewisch T, Lenis J, Jiang G-L, Wang D, Pantalone V, Bilyeu K: **The development and use**
of a molecular model for soybean maturity groups. *BMC Plant Biology* 2017, **17:**91.

Lemmon ZH, Reem NT, Dalrymple J, Soyk S, Swartwood KE, Rodriguez-Leal D, Van Eck J,
Lippman ZB: **Rapid improvement of domestication traits in an orphan crop by genome**
editing. *Nat Plants* 2018, **4:**766-770.

Li Y, Kadarmideen HN, Dekkers JCM: **Selection on multiple QTL with control of gene**
diversity and inbreeding for long-term benefit. *Journal of Animal Breeding and Genetics*
2008, **125:**320-329.

Liu H, Meuwissen TH, Sorensen AC, Berg P: **Upweighting rare favourable alleles increases**
long-term genetic gain in genomic selection programs. *Genet Sel Evol* 2015, **47:**19.

Long N, Gianola D, Rosa GJM, Weigel KA: **Application of support vector regression to**
genome-assisted prediction of quantitative traits. *Theoretical and Applied Genetics* 2011,
123:1065.

Long N, Gianola D, Rosa GJM, Weigel KA, Kranis A, GonzÁLez-Recio O: **Radial basis**
function regression methods for predicting quantitative traits using SNP markers. *Genetics*
Research 2010, **92:**209-225.

Lorenz AJ: **Resource Allocation for Maximizing Prediction Accuracy and Genetic Gain of**
Genomic Selection in Plant Breeding: A Simulation Experiment. *G3:*
Genes|Genomes|Genetics 2013, **3:**481.

Marulanda J, Mi X, Melchinger A, Xu J-L, Würschum T, Longin C: **Optimum breeding**
strategies using genomic selection for hybrid breeding in wheat, maize, rye, barley, rice and
triticale. *Theor Appl Genet* 2016, **129:**1901-1913.

McCouch S, Baute GJ, Bradeen J, Bramel P, Bretting PK, Buckler E, Burke JM, Charest D,
Cloutier S, Cole G: **Feeding the future.** *Nature* 2013, **499:**23-24.

Meuwissen T, Hayes B, Goddard M: **Prediction of total genetic value using genome-wide**
dense marker maps. *Genetics* 2001, **157:**1819-1829.

Meuwissen T, Hayes B, Goddard M: **Accelerating improvement of livestock with genomic**
selection. *Annu Rev Anim Biosci* 2013, **1:**221-237.

Meuwissen TH: **Maximizing the response of selection with a predefined rate of inbreeding.**
Journal of animal science 1997, **75:**934-940.

Meuwissen TH, Goddard ME: **Mapping multiple QTL using linkage disequilibrium and**
linkage analysis information and multitrait data. *Genetics Selection Evolution* 2004, **36:**261.

Michel S, Ametz C, Gungor H, Epure D, Grausgruber H, Loschenberger F, Buerstmayr H:
Genomic selection across multiple breeding cycles in applied bread wheat
breeding.(Original Article)(Report). *Theoretical and Applied Genetics* 2016, **129:**1179.

Mikel MA, Diers BW, Nelson RL, Smith HH: **Genetic Diversity and Agronomic**
Improvement of North American Soybean Germplasm. *Crop Science* 2010, **50:**1219-1229.

Montesinos-López OA, Martín-Vallejo J, Crossa J, Gianola D, Hernández-Suárez CM,
Montesinos-López A, Juliana P, Singh R: **A Benchmarking Between Deep Learning, Support**
Vector Machine and Bayesian Threshold Best Linear Unbiased Prediction for Predicting
Ordinal Traits in Plant Breeding. *G3: Genes/Genomes/Genetics* 2019, **9:**601.

Müller D, Schopp P, Melchinger AE: **Persistency of Prediction Accuracy and Genetic Gain in**
Synthetic Populations Under Recurrent Genomic Selection. *G3: Genes/Genomes/Genetics*
2017, **7:**801.

Müller D, Schopp P, Melchinger AE: **Selection on Expected Maximum Haploid Breeding**
Values Can Increase Genetic Gain in Recurrent Genomic Selection. *G3:*
Genes/Genomes/Genetics 2018, **8:**1173.

Myers RH: *Response surface methodology / Raymond H. Myers.* s.l. : [Ann Arbor: s.l. : s.n. Ann
Arbor : available from Edwards Brothers; 1976.

Nakaya A, Isobe SN: **Will genomic selection be a practical method for plant breeding?**
Annals of Botany 2012, **110:**1303-1316.

Norman A, Taylor J, Edwards J, Kuchel H: **Optimising Genomic Selection in Wheat: Effect of**
Marker Density, Population Size and Population Structure on Prediction Accuracy. *G3:*
Genes/Genomes/Genetics 2018.

Oddi FJ, Miguez FE, Ghermandi L, Bianchi LO, Garibaldi LA: **A nonlinear mixed-effects**
modeling approach for ecological data: Using temporal dynamics of vegetation moisture as
an example. *Ecology and evolution* 2019, **9:**10225-10240.

Orf JH: **2 - Breeding, Genetics, and Production of Soybeans.** In *Soybeans.* Edited by Johnson
LA, White PJ, Galloway R: AOCS Press; 2008: 33-65

Ortiz-Perez E, Mian RMA, Cooper RL, Mendiola T, Tew J, Horner HT, Hanlin SJ, Palmer RG:
Seed-set evaluation of four male-sterile, female-fertile soybean lines using alfalfa
leafcutting bees and honey bees as pollinators. *The Journal of Agricultural Science* 2008,
146:461-469.

Peccoud J, Velden KV, Podlich D, Winkler C, Arthur L, Cooper M: **The selective values of**
alleles in a molecular network model are context dependent. *Genetics* 2004, **166:**1715-1725.

Pérez P, de los Campos G: **Genome-Wide Regression and Prediction with the BGLR**
Statistical Package. *Genetics* 2014, **198:**483-495.

Pinheiro JC: *Mixed-effects models in S and S-PLUS / José C. Pinheiro, Douglas M. Bates.* New
York: New York : Springer; 2000.

Podlich DW, Cooper M: **Modelling Plant Breeding Programs as Search Strategies on a**
Complex Response Surface. In *Simulated Evolution and Learning: Second Asia-Pacific*
Conference on Simulated Evolution and Learning, SEAL'98 Canberra, Australia, November 24–
27, 1998 Selected Papers. Edited by McKay B, Yao X, Newton CS, Kim J-H, Furuhashi T.
Berlin, Heidelberg: Springer Berlin Heidelberg; 1999: 171-178

Robertson A: **A Theory of Limits in Artificial Selection.** *Proceedings of the Royal Society of*
London Series B Biological Sciences 1960, **153:**234.

Rodríguez-Leal D, Lemmon ZH, Man J, Bartlett ME, Lippman ZB: **Engineering Quantitative**
Trait Variation for Crop Improvement by Genome Editing. *Cell* 2017, **171:**470-480.e478.

Rutten MJM, Bijma P, Woolliams JA, van Arendonk JAM: **SelAction: Software to Predict**
Selection Response and Rate of Inbreeding in Livestock Breeding Programs. *Journal of*
Heredity 2002, **93:**456-458.

Ryman N, Leimar O: **GST is still a useful measure of genetic differentiation — a comment**
on Jost's D. *Molecular Ecology* 2009, **18:**2084-2087.

Schulz-Streeck T, Ogutu JO, Karaman Z, Knaak C, Piepho HP: **Genomic Selection using**
Multiple Populations. *Crop Science* 2012, **52:**2453-2461.

Shoemaker R, Polzin K, Labate J, Specht J, Brummer E, Olson T, Young N, Concibido V,
Wilcox J, Tamulonis J: **Genome duplication in soybean (Glycine subgenus soja).** *Genetics*
1996, **144:**329-338.

Song Q, Hyten DL, Jia G, Quigley CV, Fickus EW, Nelson RL, Cregan PB: **Fingerprinting**
Soybean Germplasm and Its Utility in Genomic Research. *G3: Genes|Genomes|Genetics*
2015, **5:**1999.

Song Q, Yan L, Quigley C, Jordan BD, Fickus E, Schroeder S, Song B-H, Charles An Y-Q,
Hyten D, Nelson R, et al: **Genetic Characterization of the Soybean Nested Association**
Mapping Population. *The Plant Genome* 2017, **10**.

Specht JE, Diers BW, Nelson RL, de Toledo JFF, Torrion JA, Grassini P: **Soybean.** *Yield gains*
in major US field crops 2014, **33:**311-355.

Stewart-Brown BB, Song Q, Vaughn JN, Li Z: **Genomic Selection for Yield and Seed**
Composition Traits Within an Applied Soybean Breeding Program. *G3:*
Genes|Genomes|Genetics 2019, **9:**2253.

Wang Z, Chu T, A Choate L, Danko C: *Rgtsvm: Support Vector Machines on a GPU in R.* 2017.

Weir BS: **Disequilibrium.** *methods for discrete population genetic data* 1996**:**91-139.

Wilcox JR, Schapaugh Jr. WT, Bernard RL, Cooper RL, Fehr WR, Niehaus MH: **Genetic**
Improvement of Soybeans in the Midwest1. *Crop Science* 1979,
19:cropsci1979.0011183X001900060014x.

Wimmer V, Lehermeier C, Albrecht T, Auinger H-J, Wang Y, Schön C-C: **Genome-Wide**
Prediction of Traits with Different Genetic Architecture Through Efficient Variable
Selection. *Genetics* 2013, **195:**573.

Woolliams J, Corbin L: **Coalescence theory in livestock breeding.** *Journal of Animal Breeding*
and Genetics 2012, **129:**255-256.

Woolliams JA, Berg P, Dagnachew BS, Meuwissen TH: **Genetic contributions and their**
optimization. *J Anim Breed Genet* 2015, **132:**89-99.

Xavier A: **Efficient Estimation of Marker Effects in Plant Breeding.** *G3:*
Genes|Genomes|Genetics 2019, **9:**3855.

Xavier A, Jarquin D, Howard R, Ramasubramanian V, Specht JE, Graef GL, Beavis WD, Diers
BW, Song Q, Cregan P, et al: **Genome-Wide Analysis of Grain Yield Stability and**
Environmental Interactions in a Multiparental Soybean Population. *G3:*
Genes|Genomes|Genetics 2017.

Xavier A, Xu S, Muir W, Rainey KM: **Genomic prediction using subsampling.** *BMC*
Bioinformatics 2017, **18:**191.

Xavier A, Thapa R, Muir WM, Rainey KM: **Population and quantitative genomic properties**
of the USDA soybean germplasm collection. *Plant Genetic Resources* 2018**:**1-11.

Xavier A, Muir WM, Rainey KM: **Assessing Predictive Properties of Genome-Wide Selection**
in Soybeans. *G3: Genes|Genomes|Genetics* 2016, **6:**2611.

Xu S: **Theoretical basis of the Beavis effect.** *Genetics* 2003, **165:**2259-2268.

Xu P, Wang L, Beavis WD: **An optimization approach to gene stacking.** *European Journal of*
Operational Research 2011, **214:**168-178.

Yabe S, Yamasaki M, Ebana K, Hayashi T, Iwata H: **Island-Model Genomic Selection for**
Long-Term Genetic Improvement of Autogamous Crops. *PLoS One* 2016, **11:**e0153945.

Yu J, Holland JB, McMullen MD, Buckler ES: **Genetic Design and Statistical Power of**
Nested Association Mapping in Maize. *Genetics* 2008, **178:**539.

Zhong S, Dekkers JCM, Fernando RL, Jannink J-L: **Factors Affecting Accuracy From**
Genomic Selection in Populations Derived From Multiple Inbred Lines: A Barley Case
Study. *Genetics* 2009, **182:**355-364.

Zuur A: ***Mixed Effects Models and Extensions in Ecology with R*** *by Alain Zuur, Elena N. Ieno,*
Neil Walker, Anatoly A. Saveliev, Graham M. Smith. 1st ed. 2009.. edn: New York, NY :
Springer New York : Imprint: Springer; 2009.

1130
1131

1132

1133

Figure 1 Flow Chart for Simulations of Recurrent Genomic Selection. The upper half panel represents the steps involved in generating the base population of 2000 F_5 RILs derived from 20 NAM founder lines crossed, *in silico*, to IA3023. It includes the model training step for genomic prediction models. The lower half panel represents recurrent steps of prediction, sorting, truncation selection, crossing, and generation of F_5 RILs for each cycle as well as the decision steps to check if the training set should be updated and if the recurrent process is to be continued for another cycle.

Figure 2 Theoretical Genotypic Values: Theoretical Genotypic Values from 40 cycles of recurrent selection modeled with the recurrence equation, where y_c represents the genotypic value in cycle c, with c= 1, 2… 40 and values of α and β range from 0.6-0.9 and 1.32-38.06 respectively for 360 combinations of factors across all selection methods, training sets, selection intensities, number of simulated QTL and simulated heritabilities. The bold lines represent curves with the smallest and largest beta values and their corresponding alpha values. The curves are colored corresponding to 40 (pink), 400 (blue) and 4289 (gray) simulated QTL.

Figure 3 Estimated prediction accuracies and MSE in Founding Set of RILs: Estimated prediction accuracies (left panel) and mean squared errors (right panel) for four genomic prediction (GP) models: BayesB, BL (Bayes LASSO), RRREML (Ridge Regression with REML) and SVMRBF (Support Vector Machines with Radial Basis Function Kernel) trained with F_5 RILs derived from crosses of 20 homozygous founder lines with IA3023. Phenotypes used to train the GP models consisted of genetic architectures comprised of 40, 400 and 4289 simulated QTL (top, middle and bottom) that were responsible for 70% (blue) and 30% (red) of phenotypic variability in the initial populations.

Figure 4 Standardized Responses for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction Forty cycles of standardized responses to selection of 10% of 2000 soybean RILs per cycle. Standardized responses are plotted by selection methods without (left panels) and with (right panels) model updating using prior cycles as training sets for the four genotypic prediction models. Phenotypic selection (PS) is not updated and hence is the same in the left and right panels. The top panels consist of responses for genetic architectures consisting of 40 simulated QTL. Middle panels consist of responses for genetic architectures consisting of 400 simulated QTL and the bottom panels consist of responses for genetic architectures consisting of 4289 simulated QTL. All 40, 400, and 4289 simulated QTL are responsible for 70% of phenotypic variability in the initial population. PS - Phenotypic Selection, RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL – Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Figure 5 Average Genotypic Value and Maximum Genetic Potential for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction Average genotypic value and maximum possible genotypic value in recurrent selection of 10% of 2000 soybean RILs per cycle. The values are plotted by selection methods without (left panels) and with (right panels) model updating using prior cycles as training sets for the four genotypic prediction models. Plots demonstrate decrease in maximum possible genotypic value due to loss of favorable alleles and increase in average genotypic value for 10 cycles (upper panel) and 40 cycles (lower panel) of selection. Phenotypic selection (PS) is not updated and hence is the same in the left and right panels. The top and bottom panels represent genotypic values for genetic architectures consisting of 400 simulated QTL responsible for 70% of phenotypic variability in the initial population. PS - Phenotypic Selection, RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL - Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Method	QTL	heritability	TH5	TH10	TH15	TH20	TH25	TH30	TH35	TH40
RR-REML	40	0.7	-2.36	-12.71	-9.63	-9.61	-9.61	-9.61	-9.61	-9.61
BayesB	40	0.7	10.96	-4.18	-5.49	-5.49	-5.49	-5.49	-5.49	-5.49
BL	40	0.7	3.46	-9.96	-8.77	-8.24	-8.24	-8.24	-8.24	-8.24
SVMRBF	40	0.7	-39.67	-49.09	-46.90	-46.67	-46.67	-46.67	-46.67	-46.67
RR-REML	400	0.7	2.42	-7.37	-14.57	-17.25	-19.61	-20.96	-20.93	-21.11
BayesB	400	0.7	0.10	-9.53	-14.89	-17.58	-19.94	-21.26	-21.05	-21.11
BL	400	0.7	-3.35	-11.95	-15.15	-18.26	-19.62	-21.21	-20.80	-21.37
SVMRBF	400	0.7	-21.91	-43.06	-49.93	-52.95	-54.26	-55.48	-55.46	-55.42
RR-REML	4289	0.7	-6.26	-20.12	-25.33	-27.60	-29.16	-30.98	-32.60	-33.57
BayesB	4289	0.7	-9.07	-19.79	-26.28	-29.14	-31.30	-33.37	-34.90	-35.84
BL	4289	0.7	-5.85	-17.94	-23.50	-25.70	-27.24	-29.66	-31.36	-32.35
SVMRBF	4289	0.7	-34.49	-55.74	-63.38	-66.01	-66.85	-67.79	-68.60	-69.15

Figure 6 Heat Map for Percent Gain in Rs Relative to PS for 0.7 H for GP Models without Updating: Heat map indicating standardized response relative to PS as percentage gain after 40 cycles of recurrent selection using genomic prediction models without updated training sets for 40, 400, 4289 simulated QTL responsible for 70% of phenotypic variability in the initial population. Blue to red shaded cells represent increasing gain in response relative to PS. RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL – Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Method	QTL	heritability	No. Prior Cycles	TH5	TH10	TH15	TH20	TH25	TH30	TH35	TH40
RR-REML	40	0.7	14	9.04	1.55	0.00	0.00	0.00	0.00	0.00	0.00
BayesB	40	0.7	14	14.41	-1.05	-2.06	-2.06	-2.06	-2.06	-2.06	-2.06
BL	40	0.7	14	11.89	-4.70	-5.49	-5.49	-5.49	-5.49	-5.49	-5.49
SVMRBF	40	0.7	14	-39.85	-48.58	-47.63	-47.36	-47.36	-47.36	-47.36	-47.36
RR-REML	400	0.7	14	11.34	14.84	12.30	8.71	5.73	3.29	3.50	3.22
BayesB	400	0.7	14	13.05	12.00	8.38	1.75	-1.11	-2.96	-3.02	-3.91
BL	400	0.7	14	13.85	10.44	6.45	1.02	-1.89	-4.24	-3.92	-4.55
SVMRBF	400	0.7	14	-21.66	-42.81	-50.39	-53.52	-54.86	-55.38	-55.48	-55.40
RR-REML	4289	0.7	14	10.01	10.51	11.79	13.39	14.52	13.67	11.70	10.13
BayesB	4289	0.7	14	10.10	10.41	9.09	8.31	7.20	4.56	2.06	0.60
BL	4289	0.7	14	10.32	9.22	8.87	8.78	8.22	6.11	3.59	2.10
SVMRBF	4289	0.7	14	-36.73	-56.79	-64.13	-66.87	-68.08	-68.97	-69.49	-69.96

Figure 7 Heat Map for Percent Gain in Rs Relative to PS for 0.7 H for GP Models with Updating: Heat map indicating standardized response relative to PS as percentage gain after 40 cycles of recurrent selection using genomic prediction models with updated training sets from up to 14 prior cycles of selection for 40, 400, and 4289 simulated QTL responsible for 70% of phenotypic variability in the initial population. Blue to red shaded cells represent increasing gain in response relative to PS. RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL – Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Figure 8 Maximal Genotypic Value for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction: Maximum attained genotypic values (Mgvs) in recurrent genomic selection and phenotypic selection (PS) without updating the training sets in the left panels and with training set updates from up to 14 prior cycles in the right panels. PS has no training sets and hence does not change between the left and right panels. a) 40 QTL (top), b) 400 QTL (middle) and c) 4289 QTL (bottom) responsible for 70% of phenotypic variability in the initial population and selection of 10% of the RILs in each cycle. PS – Phenotypic Selection, RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL – Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Figure 9 Standardized Genotypic Variance (Sgv) for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction Standardized genotypic variance without training set updating (left panels) and with training set updating using prior cycle training data (right panels) for the four GP models. PS has no updating and hence is the same in both left and right panels. A) Training data from up to 14 prior cycles for 40 simulated QTL (top), 400 simulated QTL (middle) and 4289 simulated QTL (bottom) responsible for 70% of phenotypic variability in the initial population and top 10% of RILs with the greatest predicted values. PS – Phenotypic Selection, RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL – Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Figure 10 Estimated Prediction Accuracies for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction: Estimated prediction accuracies with updates to the training sets used in genomic prediction (GP) models. Training data from up to 14 prior selection cycles were used to update all four GP models for 40 QTL (top), 400 QTL (middle) and 4289 QTL (bottom) responsible for 70% of phenotypic variability in the initial population and top 10% of RILs with the greatest predicted values. RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL - Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

Figure 11 Standardized Mean Squared Error for Comparison of GS methods with and without Updating for 0.7 H and Top 10% Selected Fraction: Mean Squared Error of GP models with updates to the training sets used in genomic prediction (GP) models. Standardized MSE (>=1) is estimated as the ratio of MSE for GP models in cycle 'c' to MSE for GP models trained with founder population of RILs. While MSE for RRREML model were lesser with model updating, MSE for bayesian methods increased orders of magnitude in late cycles of selection with model updating. SVMEBF showed relatively constant MSE with and without updating. Training data from up to 14 prior selection cycles were used to update all four GP models for 40 QTL (top), 400 QTL (middle) and 4289 QTL (bottom) responsible for 70% of phenotypic variability in the initial population and top 10% of RILs with the greatest predicted values. RR-REML- Ridge Regression with Restricted Maximum Likelihood, BL - Bayes LASSO, and SVMRBF- Support Vector Machine with Radial Basis Kernel.

TABLE 1: FACTORIAL DESIGN

Factors	Number of Levels	Values for Levels
Number of QTL	3	40, 400, 4289
Heritability	2	0.7, 0.3
Selection Intensity	3	2.67, 2.34, 1.75
Selection Method	5	i) PS- Phenotypic value ii) GS – GP(RR-REML) iii) GS - GP(Bayes B) iv) GS - GP(Bayes LASSO) v) GS – GP(SVM, Radial basis Function Kernel)
Model Update: number of prior cycles used in training sets	4	i) 0 previous cycles ii) 10 previous cycles iii) 12 previous cycles iv) 14 previous cycles
Total Number of unique combinations	360	
Total Number of Simulations	3600 with 10 reps /condition	

TABLE 2 PACKAGES IN R FOR PARAMETRIC AND NON-PARAMETRIC MODELS WITH TUNING PARAMETERS

GS Model	Package (R)	Model Tuning Parameters (BGLR package)
Ridge Regression	REML-EM (custom R script Xavier 2019)	EM algorithm for estimation of parameters with REML method without using matrix inversion
Bayesian LASSO	BGLR (Perez et al. 2014)	Priors for varE (df=3,S=0.25); varU (df=3,S=0.63); lambda(shape=0.53,rate=5e-5) type='random',value=30), nIter=20000, burnIn=2000, thin=1
Bayes B	BGLR (Perez et al. 2014)	nIter=41000, burnIn =1000, df0=4, R2=0.7
SVM	Rgtsvm (Wang et al. 2017)	SVM with Radial basis function kernel on GPU