Diversity of culturable actinobacteria producing protease inhibitors from the

intertidal zones of Maharashtra, India

Neha Shintre[1], Ulfat Baig[2], Anagha Pund[2], Rajashree Patwardhan[3], Vaijayanti

Tamhane[4], Neelima Deshpande[1]

[1]Department of Microbiology, M.E.S. Abasaheb Garware College, Karve Road, Pune 411004, Maharashtra, India.

[2]Indian Institute of Science Education and Research, Pune (IISER-P), Dr. Homi Bhabha Road, Pashan, Pune 411008, Maharashtra, India.

[3]Department of Microbiology, Haribhai V. Desai College of Commerce, Arts and Science, Pune – 411002, Maharashtra, India H. V. Desai College, Pune

[4]Institute of Bioinformatics and Biotechnology, Savitribai Phule Pune University, Pune – 411007, Maharashtra, India

Corresponding authors: V. Tamhane (email: vatamhane@gmail.com,) and

N. Deshpande (email: neelimapune@gmail.com)

Keywords: Marine actinobacteria; Molecular phylogeny; Protease inhibitors; chemical ecology

ABSTRACT

Phylogenetic diversity of culturable actinobacteria isolated from the intertidal regions of west

coast of Maharashtra, India was studied using 16S rRNA gene sequencing. Total of 140

actinobacterial isolates were obtained, which belonged to 14 genera, 10 families and 65 putative

species with *Streptomyces* being the most dominant genus followed by *Nocardiopsis* and

Micromonospora. They were screened for production of extracellular protease inhibitors (PI)

against three pure proteases viz. chymotrypsin, trypsin, subtilisin and one crude extracellular

protease from *Pseudomonas sp.* 80% of the isolates showed protease inhibition activity against

at least one of the proteases, majority of them belonged to genus *Streptomyes.* Actinobacterial

diversity from two sites Ade and Harnai with varying degrees of anthropological pressures

showed that more species diversity was obtained from site with minimal human disturbance i.e

Ade, however, site with higher human disturbance, i.e. Harnai contained higher percentage of

PI producing strains. Understanding association of PI production and abundance of

extracellular proteases due to organic load in the marine environments can give interesting

insights into the chemical ecology of microbial community in the intertidal region.

INTRODUCTION

Proteases are enzymes that catalyse proteolytic reactions and are involved in variety of biological processes like digestion, cell signalling or tumour formation (Groll et al., 2002; Koblinski et al., 2000; Overall and Blobel, 2007; Roberts et al., 2012). They are ubiquitously present across different plant, animal and microbial taxa. Two of the many roles of these enzymes in bacteria is to aid in bacterial pathogenesis (Ingmer and Brøndsted, 2009; Maeda, 1996) and bacterial predation (Martin, 2002) where they secrete hydrolytic enzymes in order to degrade prey species in the vicinity (Martin, 2002). Protease inhibitors bind reversibly or irreversibly with the enzymes, rendering them inactive. Hence, protease inhibitors (PIs) play a crucial role as defence molecules in intracellular or extracellular environment. They are classified based on either the type of protease they inhibit (e.g. serine protease inhibitor, aspartic protease inhibitor) or based on the mechanism of action (e.g. reversible or irreversible enzyme inhibitors). Various plants, animals and microorganisms produce PIs in order to deter pests and predators or secrete them as toxins for self-defence (Habib and Fazili, 2007; Hartl et al., 2011; Mourão and Schwartz, 2013). Also, PIs have high commercial potential since they are used as drugs against viral, bacterial and other parasitic infections (Karthik et al., 2014; Liu et al., 2012; Sreedharan and Bhaskara Rao, 2017; Umezawa, 1976).

Marine intercostal community harbours a variety of sedentary invertebrates and microorganisms. These organisms rely primarily on chemicals for self-defence (Engel et al., 2002; Pawlik, 1993). Additionally, as stated by Hay *et al.* the "biggest challenge for marine organisms is to obtain lunch without becoming lunch" (Hay, 2009). Protease inhibitors can play a major role in these situations since they can inhibit the action of digestive proteases present in the surroundings. Literature reports suggest that PIs are indeed produced by marine intercostal communities (Covaleda et al., 2012; Karthik and Kirthi, 2015; Karthik et al., 2014;

Mourão and Schwartz, 2013) Considering these facts, it might be possible to detect PIs

synthesised by marine microorganisms for their defence.

Intertidal zones are exposed to air at low tides and are submerged at high tides. Thus they

experience varying dry and wet periods, fluctuating temperatures and atmospheric pressure.

Intertidal life forms adapt to these constantly changing environment by developing unique

features. Their adaptations for survival in dynamic environments might also increase the

chances of finding unique set of metabolites from them. . Moreover, it is expected that marine

microorganisms will have different set of defence molecules than their terrestrial counterparts

(Mourão and Schwartz, 2013; Xie et al., 2018).

Actinobacteria is one of the largest phyla in domain bacteria. These are gram positive organisms

with high guanine to cytosine ratio (Trujillo, 2016). They at large are known to produce

commercially important diverse metabolites. Some reports show that the marine environment

has become a prime source for discovery of novel actinobacteria and novel natural products

produced by them (Maldonado et al., 2005; Ward and Bora, 2006). There are few recent reports

of PI production by marine actinobacteria (Karthik et al., 2014; Sreedharan and Bhaskara Rao,

2017; Sun et al., 2014) yet, to our knowledge, there are limited reports on PIs from west coast

of India. Thus, in the current study, marine actinobacteria from intertidal zone were screened

for presence of PIs.

It has been reported that sponges and bacteria are two main sources of enzyme inhibitors from

marine environments (Ruocco et al., 2017). Some reports also suggest that many a times,

metabolites obtained from sponges are actually produced by the bacterial symbionts (Haygood

et al., 1999; Mehbub et al., 2014). Therefore, in the current study sponge, sediment and water

samples were collected from the coast of Maharashtra and Goa for isolation of actinobacteria.

Obtained isolates were subjected to molecular identification using 16S rRNA gene sequencing

and isolates were screened for presence of protease inhibitors. Data was analysed to check

correlations between inhibitor molecules and environmental parameters and their probable role

as defence molecules.

MATERIALS AND METHODS

Sampling

Around 60 samples comprising of sponge, sediment and sea water were collected from the

intertidal rock pools from seven different locations along the coast of Ratnagiri district from

Velas, Kelshi, Ade, Anjarle, Harnai, Murud and Aare Ware. (17°57'04.9"N 73°01'43.0"E to

17°04'35.9"N 73°17'17.5"E). From Ade and Harnai, sampling was carried out in different

seasons viz. pre monsoon (Mar, Apr, May), Monsoon (Early Sep), post-monsoon (Oct) and

winter season (Dec, Jan, Feb). Whereas, a single sampling was done from rest of the sites. Sub-

tidal sampling of sediment was also done in a single event at a place off shore to Goa

(15°21'08.4"N 73°46'41.8"E). The depths of collection for this sampling were 9m, 11m, 12m

and 16m. (Figure 1).

Small tissue samples of sponge were collected without damaging the sponge colonies or their

habitat. Samples were rinsed in sterile Poor Ravan Saline (PRS) broth to remove loosely bound

particles and debris and were stored in sterile collection tubes containing the same medium.

Sediment and water samples in the vicinity of sponge were collected in sterile collection tubes.

All the samples were transported to the laboratories in ice box and were processed fresh for

microbial isolations.

Sample Processing, Selective Isolation and Culture Maintenance

All the samples were given heat treatment at 60°C for 15 mins to reduce the load of non-

sporulating bacteria. Sponge samples were homogenised in Poor Ravan Saline (PRS) medium

(Watve et al., 2000) and diluted serially with 10 fold dilutions up to 10^{-5}. Sediment samples

were vigorously shaken for two minutes and diluted up to 10^{-5} dilutions. Sea water samples

were also diluted up to 10^{-5} dilutions. 0.1 ml of undiluted, 10^{-3} and 10^{-5} dilutions were spread

plated in triplicates on different growth media and incubated at room temperature for up to 21

days.

Four different growth media Sponge agar (1% macerated sponge colonies collected from the

site, 50% sea water and 2.5% agar), Sea Water Agar (50% sea water and 2.5% agar), ZOBELL

Marine Agar (ZOBELL, 1941) and Modified poor ravan medium (Watve et al., 2000)) were

used to obtain maximum culture dependent diversity of actinobacteria. Nalidixic acid and

Cyclohexamide were added to the culture medium in 25 µg/ml concentration to inhibit the

growth of Gram-negative bacteria and fungi (Magarvey et al., 2004). Plates were incubated at

30°C and monitored daily for 21 days. Isolated colonies showing resemblance to typical

actinobacterial colony morphology were picked up and subcultured several times for obtaining

pure cultures. All the pure cultures were later stored on modified (1:4 diluted) ZOBELL marine

agar at 4°C for further use.

DNA Sequencing and Analysis

16S rRNA sequences from previous study of (Baig et al., 2020) were used to make maximum

likelihood phylogenetic tree of actinobacteria used in the study. Maximum likelihood tree was

constructed in IQtree (Nguyen et al., 2015). Best nucleotide substitution model was determined

in model finder (Kalyaanamoorthy et al., 2017). Note support was examined using bootstrap

values of 1000 iterations and *Bacillus sp.* were used as out-group.

Screening for Protease Inhibitors (PI) using spot assay

Sample preparation

Detection of extracellular PIs from cell free supernatant was done as follows. Pure cultures of

actinobacteria were inoculated (single colony in 150 ml broth) in ZOBELL Marine broth and

incubated for 5 days at 30°C on incubator shaker with 100 rpm speed. After incubation, the

cultures were centrifuged at 4000 rpm for 20 mins and cell free supernatants (CFS) were

collected. 1 ml of each of these CFSs were kept in hot water bath of 70°C for 15 mins for

denaturation of proteases produced by the bacteria. Heat treated CFSs were used for detection

of PIs. Unprocessed X-ray films coated with gelatine were used to detect PIs. (Gelatine is

degraded by various proteolytic enzymes, so, upon action of the proteases, clear zones are

observed on unprocessed x-ray films at the point of contact of enzymes. On the other hand, if

the gelatine layer remains intact, no clearance is observed. PIs inactivate proteases and thus,

presence of protease inhibitors is marked by no clearance on the x-ray films as shown in Figure

3).

Proteases used for the assay included Trypsin (SRL Pvt. Ltd, Cat no.- 60484) and α-

Chymotrypsin (SRL Pvt. Ltd, Cat no.- 35085) of bovine origin and Subtilisin (Sigma-Aldrich,

Cat no.- P5380) and crude protease obtained in the laboratory from cell free supernatant of

Pseudomonas sp of bacterial origin. were used to detect PIs. Amongst these enzymes, trypsin

and chymotrypsin were of bovine origin. Additionally, Protease inhibitor cocktail (Sigma-

Aldrich, Cat no.- P5380) was used as a positive control and un-inoculated culture broth

incubated with the enzyme was used as a negative control.

Spot assay

Standardization of enzyme concentration was done as described by (Tripathi et al.,

2011).Various dilutions of pure enzyme were spotted on gelatine coated X-ray films. Lowest

dilution that gave complete clearance, in turn indicating complete digestion of gelatine, was

used in the assay.

Spot assay as described by (Cheung et al., 1991) for detection of PIs was carried out as follows.

10 µl of pure enzyme (100 µg/ml) was incubated with 10 µl of heat treated CFS for 10 mins

and then transferred to untreated X-ray-Fuji Medical X-ray, HRU grade-films. The assembly

was kept undisturbed for 15 mins at room temperature to allow the enzyme substrate reaction

to take place. Films were washed with running tap water and allowed to dry before recording

the results.

RESULTS

Sample Collection

A single sampling of sediment and water was carried out from intertidal and sub-tidal regions

of Aare Ware and Goa respectively. Similarly, one time sampling was carried out from

intertidal zones of Velas, Kelshi and Anjarle, Murud however, at the time of sampling, sponge

colonies could not be located on any of these sites and thus only sediment and water samples

were collected. Whereas, at Harnai and Ade four to five distinct morphotypes of sponges were

recorded. Thus, sponge sediment and water samples were collected from these sites. These

sites were also selected for sampling in various seasons throughout the year. Harnai, one of the

busy ports on the western coast is used for various activities like, auction of fishes on a daily

basis, recreational space for the villagers or tourist hang out spot. Certain parts of intertidal rock

patch in the vicinity are used as open defecation sites by the villagers and few other parts of the

rock patch are used for clamp collection and fishing from the rock pools. Conversely Ade, is

hardly used for any of the above purposes except for occasional fishing from the rock pools.

Thus, intertidal regions of the above sites faced varying degree of anthropological disturbance

with Harnai being the most and Ade being the least used site with respect to human activities.

Distribution, Identification and Phylogeny of Actinobacteria from Sponge, Sediment and Sea

Water Samples

We obtained 140 actinobacterial isolates from sponge sediment and sea water samples. They

showed high phylogenetic diversity wherein there were 65 putative species belonging to 14

genera of 10 different families. Most abundant genus amongst the isolates was *Streptomyces*

sp. (~63%) followed by *Nocardiopsis sp.* (~22%) and *Micromonospora sp.* (~4%).

Approximately 11% isolates belonged to other genera (Table 1).

Table 1: Counts of putative species obtained from sponge, sediment and sea water samples

Family	Genus	Putative species identity	Isolates
Actinomycetaceae	*Streptomyces*	*Streptomyces albidoflavus*	6
		Streptomyces albogriseolus	8
		Streptomyces atrovirens	1
		Streptomyces aurantiogriseus	1
		Streptomyces aureofaciens	1
		Streptomyces cellulosae	1
		Streptomyces champavatii	1
		Streptomyces coeruleofuscus	1
		Streptomyces collinus	1
		Streptomyces diastaticus	3
		Streptomyces euryhalinus	4
		Streptomyces fradiae	6
		Streptomyces geysiriensis	1
		Streptomyces graminearus	1
		Streptomyces griseorubens	3
		Streptomyces koyangensis	1
		Streptomyces longispororuber	4
		Streptomyces maritimus	6
		Streptomyces nigra	1
		Streptomyces olivaceus	4
		Streptomyces prasinosporus	1
		Streptomyces pseudogriseolus	4
		Streptomyces redgersensis	1
		Streptomyces rochei	2
		Streptomyces sampsonii	3
		Streptomyces smyrnaeus	1

		Streptomyces sp. 102H11-4	1
		Streptomyces sp. 13650C	8
		Streptomyces sp. CNS-753	1
		Streptomyces sp. FZ42	1
		Streptomyces sp. OAct 12	1
		Streptomyces sp. OAct 89	1
		Streptomyces tempisquensis	2
		Streptomyces tendae	2
		Streptomyces thermocarboxydus	1
		Streptomyces variabilis	1
		Streptomyces viridobrunneus	1
		Streptomyces xylophagus	1
Nocardiopsaceae	*Nocardiopsis*	*Nocardiopsis alba*	21
		Nocardiopsis dasonvelli	2
		Nocardiopsis fildesensis	1
		Nocardiopsis metallicus	2
		Nocardiopsis salina	1
		Nocardiopsis synnematoformis	3
		Nocordiopsis metallicus	1
Micromonosporaceae	*Micromonospora*	*Micromonospora chalcea*	2
		Micromonospora maritima	1
		Micromonospora sp	1
		Micromonospora tulbaghiae	1
Nocardiaceae	*Rhodococcus*	*Rhodococcus aetherivorans*	1
		Rhodococcus cory	1
		Rhodococcus rhodochrous	1
		Rhodococcus zopfii	1
Pseudonocardiaceae	*Actinomycetospora*	*Actinomycetospora chiangmaiensis*	1
		Actinomycetospora straminia	1
	Pseudonocardia	*Pseudonocardia kongjuensis*	1
Micrococcaceae	*Glutamicibacter*	*Arthrobacter mysoreus*	1
	Kocuria	*Kocuria rosea*	1
	Micrococcus	*Micrococcus aloeverae*	1
	Rothia	*Rothia terrae*	1
Brevibacteriaceae	*Brevibacterium*	*Brevibacterium leteolum*	2
Intrasporangiaceae	*Kytococcus*	*Kytococcus sedenteris*	1
Jonesiaceae	*Jonesia*	*Jonesia denitrificans*	1
Microbacteriaceae	*Agrococcus*	*Agrococcus carbonis*	1

191   From sponges, 50 actinobacterial isolates were obtained which belonged to 7 genera and 7

192   families. Whereas, 70 isolates were obtained from sediments that belonged to 7 different genera

193   and 7 families. 19 isolates belonging to 8 genera of 7 families were isolated from sea water

194   (Table 2).

195   **Table 2:** Number of isolates belonging to different families obtained from various sources

Families	Sediment	Water	Sponge	Total
Actinomycetaceae	47	9	32	**88**
Brevibacteriaceae	1	1	-	**2**
Intrasporangiaceae	1	-	-	**1**
Microbacteriaceae	-	1	-	**1**
Micrococcaceae	1	1	2	**4**
Micromonosporaceae	1	-	4	**5**
Nocardiaceae	1	1	2	**4**
Nocardiopsaceae	18	2	11	**31**
Pseudonocardiaceae	-	2	1	**3**
Jonesiaceae	-	-	1	**1**
Grand total	**70**	**17**	**53**	**140**

More number of isolates (67) and more species diversity (10 genera with 43 putative species)

were obtained from anthropologically less disturbed site Ade. Whereas, from Harnai which is

a site with higher disturbance 47 isolates belonging to 25 putative species of 9 genera could be

isolated. (Figure 2).

Production of Protease inhibitors

Results of PI production were recorded using spot assay as shown in Figure 3. It was observed

that, PIs retained their activity even after heat treatment at 70°C for 15 mins. Out of 140 isolates

used for screening of PIs, 113 isolates showed activity against at least one of the three pure

proteases used for the study (viz. chymotrypsin, trypsin and subtilisin), whereas 27 isolates

showed no PI production at all against any of these enzymes. Out of 113 isolates, 37 showed

some degree of activity against all pure enzymes, of which 8 showed strong positive activity.

Total of 17, 7 and 9 isolates showed PI activity exclusively against chymotrypsin, trypsin and

subtilisin respectively (Figure 3). Majority of isolates which showed strong positive activity

against enzyme chymotrypsin, also showed strong positive activity against subtilisin and trypsin. Further, even though 20% of the *Streptomyces* members showed complete absence of protease inhibitors, all the isolates showing strong positive activity against all enzymes belonged entirely to *Streptomyces* sp. Very few isolates of *Nocardiopsis* sp. showed PI activity against subtilisin. Regarding the source, highest number of strong positive isolates (7 out of 8) were obtained from the sediments and only 1 was obtained from the sponge. (Figure 2).

Out of 113 PI producers, strong positive (++) activity was shown by 10, 54, and 27 isolates against chymotrypsin, trypsin and subtilisin respectively (Table 3 A). A subset of 90 isolates belonging to most common genera *viz. Streptomyces sp.*, *Nocardiopsis sp.* and *Micromonospora sp.* were chosen to check the activity against another extracellular bacterial protease which was obtained from cell free supernatant of *Pseudomonas sp*. 36 isolates gave some degree of activity against this crude bacterial protease (Table 3B).

Table 3A: Genus-wise counts of PI producers active against pure proteases

Genus	Chymotrypsin			Trypsin			Subtilisin			Total
	-	+	++	-	+	++	-	+	++	
Actinomycetospora	0	2	0	1	1		2	0	0	**2**
Agrococcus	0	1	0	1	0	0	1	0	0	**1**
Brevibacterium	1	1	0	1	1	0	2	0	0	**2**
Glutamicibacter	0	1	0	0	0	1	1	0	0	**1**
Jonesia	0	1	0	1	0	0	1	0	0	**1**
Kocuria	0	1	0	1	0	0	1	0	0	**1**
Kytococcus	0	1	0	0	1	0	1	0	0	**1**
Micrococcus	0	1	0	0	1	0	1	0	0	**1**
Micromonospora	2	3	0	2	1	2	4	1	0	**5**
Nocardiopsis	6	23	2	17	3	11	27	4	0	**31**
Pseudonocardia	0	1	0	1	0	0	1	0	0	**1**
Rhodococcus	0	4	0	2	1	1	4	0	0	**4**
Rothia	0	1	0	0	0	1	1	0	0	**1**
Streptomyces	43	37	8	31	19	38	33	28	27	**88**
Total	**52**	**78**	**10**	**58**	**28**	**54**	**80**	**33**	**27**	**140**

226 **Table 3B:** Genus-wise counts of PI producers active against crude protease

Genus	Crude *Pseudomonas* Protease			Total
	-	+	++	
Micromonospora	2	1	0	3
Nocardiopsis	3	4	0	7
Streptomyces	49	29	2	80
Grand Total	**54**	**34**	**2**	**90**

Effect of Season and Anthropological Activities on Production of Protease Inhibitors

Least number of actinobacterial isolates were obtained from monsoon and post-monsoon

seasons and the number of isolates obtained from each sampling increased towards the month

of May i.e. the pre-monsoon season. Majority of actinobacterial isolates were obtained from

pre-monsoon season (Figure 4A). From Ade and Harnai, over all, occurrence of inhibitors

against chymotrypsin and trypsin was more as compared to that against subtilisin (Figure 4B).

Proportion of isolates not giving PI activity against any of the enzymes was higher in Ade (14

out of 21 isolates i.e. 67%) compared to that in Harnai (4 isolates out of 47 i.e. 19%) (Figure

4C). Thus to summarize, more number of putative species were obtained from Ade however,

proportion of PI producers from that site was less compared to Harnai.

DISCUSSION:

Actinobacteria are widespread in terrestrial and aquatic environments. Many of them also

produce external spores which are resistant to dehydration (Trujillo, 2016). The reports say that

marine actinobacteria have a high potential for production of biomolecules (Jose and Jha, 2017;

Solanki et al., 2008) however, the number of actinobacterial compounds discovered so far are

limited not by the number of compounds produced by them but by the amount of screening

efforts put in (Watve et al., 2001). Various reports have highlighted the need for studying

marine actinomycete diversity and their potential use for obtaining novel metabolites (Dharmaraj, 2010; Jensen et al., 2005; Lam, 2006; Subramani and Aalbersberg, 2012). Researchers from India have also isolated actinobacteria from various coastal regions of West Bengal (Peela et al., 2005; Ramesh and Mathivanan, 2009), Gujarat (Jose and Jha, 2017), Tamil Nadu (Raja et al., 2010; Valli et al., 2012) and Andaman and Nicobar (Karthik et al., 2014) for bioprospecting studies. However, there are very few reports of isolation of actinobacteria and still fewer reports on PI producing actinobacteria from the coast of Maharashtra. Thus considering these facts, current study focused on isolation of actinobacteria from coast of Maharashtra and their potential as producers of PIs.

Phylogenetics of Actinobacteria from Intertidal Regions of Maharashtra

From 7 different sites including Ade and Harnai used in the study, high diversity of actinobacterial species was obtained. Highest number of isolates were obtained from sediment samples followed by sponge and water samples majority of which belonged to *Streptomyces sp.* followed by *Nocardiopsis sp.* and *Micromonispora sp.* The observations coincided with observations of (Jose and Jha, 2017) who worked on the samples obtained from coast of Gujarat which is situated at north of Maharashtra. These findings might suggest that members of above three genera have adapted well to live in marine environments and show a widespread distribution along the western coast of India.

PI Production by Marine Actinobacteria

Marine actinobacteria are known to produce large number of bioactive molecules like antimicrobial and anticancer compounds intra-as well as extracellularly (Subramani and Aalbersberg, 2012), however, it is difficult to understand the strategy of producing extracellular molecules in the marine ecosystems, since, the molecules secreted outside can be easily diluted in the surroundings. Endopeptidases like chymotrypsin and trypsin are also secreted

extracellularly by bacteria and invertebrates in the marine environments (Holmström and

Kjelleberg, 1999; Thao et al., 2015). Moreover, reports have shown that coastal waters contain

significant amounts of trypsin-type and chymotrypsin-type endopeptidases (Obayashi and

Suzuki, 2005). Interestingly, results of current study demonstrated that high number of

actinobacteria produced PIs against chymotrypsin and trypsin. Further, more than 80% of

cultured actinobacteria produced extracellular inhibitors against at least one of the four

enzymes and almost 20% isolates showed inhibition against all the enzymes used in the study.

The presence of PIs suggests that they are needed for the defence of organisms against exo-

and endoproteases present in the surrounding environment. Moreover, they might be used as

defence molecules against proteases secreted by protists and bacteria.

Micro-ecological Dynamics in PI Producing Actinobacteria

In this study, PI producing actinobacteria were obtained from all locations and from samples

collected in all seasons. However, there was a remarkable difference in proportions of PI

producing actinobacteria obtained from sites with varying human disturbance. Although

comparatively less disturbed site Ade showed more species diversity, majority of non-

producers of PI were reported from this site. On the contrary, almost 80% isolates obtained

from Harnai (which has more human interference and thus chances of containing higher organic

load) were producing PIs. Organic load and presence of extracellular proteases in the

surroundings might be a driving force for production of extracellular PIs. Therefore, these

results can be used as a lead for carrying out studies on a larger expanse to check correlation of

organic load and production of PIs by the marine actinobacteria.

Actinobacteria are involved in bacterial predation at oligophilic conditions (Kumbhar et al.,

2014), they might need to produce proteases for killing prey species. Hence, one of the

possibilities of finding high number of PI producers amongst actinobacterial isolates might be

to achieve self-protection so that their own proteases are rendered ineffective against their own

selves. Hence, it would be interesting to check if there is any significant correlation between

production of PI and bacterial predation, (Baig et al., 2020).

Through this study we show that extracellular protease inhibitor producing actinobacteria are

abundant in the intertidal zones of west coast of Maharashtra. We hypothesize that this might

have a correlation with extracellular proteases and bacterial load in marine environment.

Understanding the need for production of protease inhibitors by actinomycetes can give

interesting insights into marine microbial ecology and clues for production of novel protease

inhibitors.

ACKNOWLEDGEMENT:

Financial support for this project was provided by Rajiv Gandhi Science and Technology

Commission under the Maharashtra Gene Bank Program. We thank Dr. Milind Watve and Dr.

Neelesh Dahanukar for their critical comments and suggestions for this manuscript. We also

thank Mr. Chinmay Kulkarni for assisting in the fieldwork.

REFERENCES:

1. Baig, U., Dahanukar, N., Shintre, N., Holkar, K., Pund, A., Lele, U., Gujarathi, T.,
Patel, K., Jakati, A., Singh, R., et al. Phylogenetic diversity and activity screening of
cultivable actinobacteria isolated from marine sponges and associated environments
from the western coast of India. (2020), BioRxiv 2020.01.09.901108.

2. Cheung, A.L., Ying, P., and Fischetti, V.A. A method to detect proteinase activity
using unprocessed X-ray films. Anal. Biochem. (1991), *193*, 20–23.

3. Covaleda, G., Rivero, M.A. del, Chávez, M.A., Avilés, F.X., and Reverter, D. Crystal
Structure of Novel Metallocarboxypeptidase Inhibitor from Marine Mollusk Nerita
versicolor in Complex with Human Carboxypeptidase A4. J. Biol. Chem. (2012), *287*,
9250–9258.

4. Dharmaraj, S. Marine Streptomyces as a novel source of bioactive substances. World
J. Microbiol. Biotechnol. (2010), *26*, 2123–2139.

5. Engel, S., Jensen, P.R., and Fenical, W. Chemical Ecology of Marine Microbial
Defense. J. Chem. Ecol. (2002), *28*, 1971–1985.

6. Groll, U. von, Berger, D., and Altmann, T. The Subtilisin-Like Serine Protease SDD1
Mediates Cell-to-Cell Signaling during Arabidopsis Stomatal Development. Plant Cell
(2002), *14*, 1527–1539.

7. Habib, H., and Fazili, K.M. Plant protease inhibitors: a defense strategy in plants. p.
(2007).

8. Hartl, M., Giri, A.P., Kaur, H., and Baldwin, I.T. The multiple functions of plant
serine protease inhibitors: defense against herbivores and beyond. Plant Signal.
Behav. (2011), *6*, 1009–1011.

9. Hay, M.E. Marine Chemical Ecology: Chemical Signals and Cues Structure Marine
Populations, Communities, and Ecosystems. Annu. Rev. Mar. Sci. (2009), *1*, 193–
212.

10. Haygood, M.G., Schmidt, E.W., Davidson, S.K., and Faulkner, D.J. Microbial
symbionts of marine invertebrates: opportunities for microbial biotechnology. J. Mol.
Microbiol. Biotechnol. (1999), *1*, 33–43.

11. Holmström, C., and Kjelleberg, S. Marine Pseudoalteromonas species are associated
with higher organisms and produce biologically active extracellular agents. FEMS
Microbiol. Ecol. (1999), *30*, 285–293.

12. Ingmer, H., and Brøndsted, L. Proteases in bacterial pathogenesis. Res. Microbiol.
(2009), *160*, 704–710.

13. Jensen, P.R., Mincer, T.J., Williams, P.G., and Fenical, W. Marine actinomycete
diversity and natural product discovery. Antonie Van Leeuwenhoek (2005), *87*, 43–
48.

14. Jose, P.A., and Jha, B. Intertidal marine sediment harbours Actinobacteria with
promising bioactive and biosynthetic potential. Sci. Rep. (2017), *7*, 1–15.

15. Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., Haeseler, A. von, and Jermiin, L.S.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods
(2017), *14*, 587–589.

16. Karthik, L., and Kirthi, Av. Protease inhibitors from marine organisms. In
Antimicrobials, (CRC Press), pp. (2015), 175–188.

17. Karthik, L., Kumar, G., Keswani, T., Bhattacharyya, A., Chandar, S.S., and Rao,
K.V.B. Protease Inhibitors from Marine Actinobacteria as a Potential Source for
Antimalarial Compound. PLOS ONE (2014), *9*, e90972.

18. Koblinski, J.E., Ahram, M., and Sloane, B.F. Unraveling the role of proteases in
cancer. Clin. Chim. Acta (2000), *291*, 113–135.

19. Kumbhar, C., Mudliar, P., Bhatia, L., Kshirsagar, A., and Watve, M. Widespread
predatory abilities in the genus Streptomyces. Arch. Microbiol. (2014), *196*, 235–248.

20. Lam, K.S. Discovery of novel metabolites from marine actinomycetes. Curr. Opin. Microbiol. (2006), *9*, 245–251.

21. Liu, X., Gan, M., Dong, B., Zhang, T., Li, Y., Zhang, Y., Fan, X., Wu, Y., Bai, S., Chen, M., et al. 4862F, a new inhibitor of HIV-1 protease, from the culture of Streptomyces I03A-04862. Mol. Basel Switz. (2012), *18*, 236–243.

22. Maeda, H. Role of Microbial Proteases in Pathogenesis. Microbiol. Immunol. (1996), *40*, 685–699.

23. Magarvey, N.A., Keller, J.M., Bernan, V., Dworkin, M., and Sherman, D.H. Isolation and characterization of novel marine-derived actinomycete taxa rich in bioactive metabolites. Appl. Environ. Microbiol. (2004), *70*, 7520–7529.

24. Maldonado, L.A., Stach, J.E.M., Pathom-aree, W., Ward, A.C., Bull, A.T., and Goodfellow, M. Diversity of cultivable actinobacteria in geographically widespread marine sediments. Antonie Van Leeuwenhoek (2005), *87*, 11–18.

25. Martin, M.O. Predatory prokaryotes: an emerging research opportunity. J. Mol. Microbiol. Biotechnol. (2002), *4*, 467–477.

26. Mehbub, M.F., Lei, J., Franco, C., and Zhang, W. Marine sponge derived natural products between 2001 and 2010: trends and opportunities for discovery of bioactives. Mar. Drugs (2014), *12*, 4539–4577.

27. Mourão, C.B.F., and Schwartz, E.F. Protease Inhibitors from Marine Venomous Animals and Their Counterparts in Terrestrial Venomous Animals. Mar. Drugs (2013), *11*, 2069–2112.

28. Nakao, Y., Fujita, M., Warabi, K., Matsunaga, S., and Fusetani, N. Miraziridine A, a Novel Cysteine Protease Inhibitor from the Marine Sponge Theonella aff. mirabilis1. J. Am. Chem. Soc. (2000), *122*, 10462–10463.

29. Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., and Minh, B.Q. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol. (2015), *32*, 268–274.

30. Obayashi, Y., and Suzuki, S. Proteolytic enzymes in coastal surface seawater: Significant activity of endopeptidases and exopeptidases. Limnol. Oceanogr. (2005), *50*, 722–726.

31. Overall, C.M., and Blobel, C.P. In search of partners: linking extracellular proteases to substrates. Nat. Rev. Mol. Cell Biol. (2007), *8*, 245–257.

32. Pawlik, J.R. Marine invertebrate chemical defenses. Chem. Rev. (1993), *93*, 1911–1922.

33. Peela, S., Kurada, V.B., and Terli, R. Studies on antagonistic marine actinomycetes from the Bay of Bengal. World J. Microbiol. Biotechnol. (2005), *21*, 583–585.

34. Quintero, D., and Bermudes, D. A Culture-Based Method for Determining the
Production of Secreted Protease Inhibitors. J. Microbiol. Methods (2014), *100*, 105–
110.

35. Raja, S., Ganesan, S., Sivakumar, K., and Thangaradjou, T. Screening of marine
actinobacteria for amylase enzymes inhibitors. Indian J. Microbiol. (2010), *50*, 233–
237.

36. Ramesh, S., and Mathivanan, N. Screening of marine actinomycetes isolated from the
Bay of Bengal, India for antimicrobial activity and industrial enzymes. World J.
Microbiol. Biotechnol. (2009), *25*, 2103–2111.

37. Roberts, I.N., Caputo, C., Criado, M.V., and Funk, C. Senescence-associated
proteases in plants. Physiol. Plant. (2012), *145*, 130–139.

38. Ruocco, N., Costantini, S., Palumbo, F., and Costantini, M. Marine Sponges and
Bacteria as Challenging Sources of Enzyme Inhibitors for Pharmacological
Applications. Mar. Drugs (2017), *15*.

39. Solanki, R., Khanna, M., and Lal, R. Bioactive compounds from marine
actinomycetes. Indian J. Microbiol. (2008), *48*, 410–431.

40. Sreedharan, V., and Bhaskara Rao, K.V. Efficacy of protease inhibitor from marine
Streptomyces sp. VITBVK2 against Leishmania donovani - An in vitro study. Exp.
Parasitol. (2017), *174*, 45–51.

41. Subramani, R., and Aalbersberg, W. Marine actinomycetes: An ongoing source of
novel bioactive metabolites. Microbiol. Res. (2012), *167*, 571–580.

42. Sun, Y., Takada, K., Nogi, Y., Okada, S., and Matsunaga, S. Lower homologues of
ahpatinin, aspartic protease inhibitors, from a marine Streptomyces sp. J. Nat. Prod.
(2014), *77*, 1749–1752.

43. Thao, N.V., Nozawa, A., Obayashi, Y., Kitamura, S.-I., Yokokawa, T., and Suzuki, S.
Extracellular proteases are released by ciliates in defined seawater microcosms. Mar.
Environ. Res. (2015), *109*, 95–102.

44. Tripathi, V.R., Kumar, S., and Garg, S.K. A study on trypsin, Aspergillus flavus and
Bacillus sp. protease inhibitory activity in Cassia tora (L.) syn Senna tora (L.) Roxb.
seed extract. BMC Complement. Altern. Med. (2011), *11*, 56.

45. Trujillo, M.E. Actinobacteria. In ELS, (American Cancer Society), pp. (2016), 1–16.

46. Umezawa, H. [55] Structures and activities of protease inhibitors of microbial origin.
In Methods in Enzymology, (Academic Press), pp. (1976), 678–695.

47. Valli, S., Suvathi, S.S., Aysha, O., Nirmala, P., Vinoth, K.P., and Reena, A.
Antimicrobial potential of Actinomycetes species isolated from marine environment.
Asian Pac. J. Trop. Biomed. (2012), *2*, 469–473.

48. Ward, A.C., and Bora, N. Diversity and biogeography of marine actinobacteria. Curr.
Opin. Microbiol. (2006), *9*, 279–286.

49. Watve, M., Shejval, V., Sonawane, C., Rahalkar, M., Matapurkar, A., Shouche, Y.,
Patole, M., Phadnis, N., Champhenkar, A., Damle, K., et al. The "K" selected
oligophilic bacteria: a key to uncultured diversity? Curr. Sci. (2000), *78*, 1535–1542.

50. Watve, M., Tickoo, R., Jog, M., and Bhole, B. How many antibiotics are produced by
the genus Streptomyces ? Arch. Microbiol. (2001), *176*, 386–390.

51. Xie, C.-L., Xia, J.-M., Wang, J.-S., Lin, D.-H., and Yang, X.-W. Metabolomic
Investigations on Nesterenkonia flava Revealed Significant Differences between
Marine and Terrestrial Actinomycetes. Mar. Drugs (2018), *16*.

52. ZOBELL, C.E. Studies on marine bacteria, I : the cultural requirements of
heterotrophic aerobes. J Mar Res (1941), *4*, 42–75.

Figure 1: Map of west coast of India (A) illustrating sampling sites along the coast of Maharashtra and Goa (B). Corresponding positions are shown in the inset (C & D). The maps were generated using QGIS software (http://qgis.osgeo.org) and Google maps.

Figure 2: Phylogenetic relatedness of isolates in the study along with location, source of isolation and PI profile against different enzymes. Given is a maximum likelihood tree of actinobacteria used for screening of PIs. Outgroup taxa (*Bacillus* sp.) not shown. Percent bootstrap values of 1000 iterations are provided along the nodes. The figure also depicts location and source of isolation of the isolates along with their ability for production of PIs against subtilisin (SI), chymotrypsin (CI) and trypsin (TI) where (- : negative, +: positive and ++ : strong positive)

Figure 3: Spot assay representing PI activity against trypsin (A); and Venn diagram showing overlap of number of isolates producing subtilisin, chymotrypsin and trypsin inhibitors (B).

(A)

(B)

Figure 4: Graphs showing number of actinobacterial isolates obtained from Ade and Harnai in different seasons (A) , number of isolates showing inhibitory activity against subtilisin (SI), chymotrypsin (CI) and trypsin (TI) is shown in (B) and number of isolates negative for PI production are shown in (C).