Antibiotics affect migratory restlessness orientation

Yuval Werber[1], Eviatar Natan[2], Yizhar Lavner[3], Yoni Vortman[4]

1. Hula Research Center, Department of Biotechnology, Tel-Hai College
2. The Aleph Lab, Oxford, England
3. Department of Computer Sciences, Tel Hai College
4. Hula Research Center, Department of Animal Sciences, Tel-Hai College

Abstract

Magnetoreception is a sense that allows the organism to perceive and act according to

different parameters of the magnetic field. This magnetic sense plays a part in many

fundamental processes in various living organisms. Much effort was expended in finding the

'magnetic sensor' in animals. While some experiments show a role of the ophthalmic nerve in

magnetic sensing, others show that effects of light on processes in the retina are involved.

According to these inconclusive and puzzling findings, the scientific community has yet to reach

an agreement concerning the underlying mechanism behind animal magnetic sensing. Recently,

the symbiotic magnetotaxis hypothesis has been forwarded as a mechanistic explanation for

the phenomenon of animal magnetoreception. It suggests a symbiotic relationship between

magnetotactic bacteria (MTB) and the navigating host. Here we show that in contrast to the

control group, antibiotic treatment caused a lack of clear directionality in an Emlen funnel

experiment. Accordingly, the antibiotics treatment group showed a significant increase in

directional variance. This effect of antibiotics on behaviors associated with animal magnetic

sensing is, to the best of our knowledge, the first experimental support of the symbiotic

magnetotactic hypothesis.

Introduction

Actively motile animals need to be able to navigate their environment. Using memory, timing

and external cues, organisms move deliberately and reach their designated destination. Any

non-random environmental feature is a candidate for use in orientation, from chemical cues

[1,2], and visual characteristics of the environment [3,4] to movement of celestial bodies [5]

and seismic signals [6]. In most cases, a combination of such signals is used [7].

Navigation using Earth's magnetic field is ubiquitous throughout the entire tree of life. It occurs

in a great variety of organisms, playing a part in many natural processes, from bacterial

movement to global migrations. Although magnetic sensation in navigating organisms has long

been accepted as fact, its underlying mechanism in multicellular animals is still being debated

[8].

The "magnetite based magnetoreception" hypothesis suggests that biogenic magnetite crystals

serve as magnetic field sensors by arousing mechanosensitive protein structures which

translate mechanic excitation to sensory information [8]. Alternatively, the "radical pair"

hypothesis predicts that the geomagnetic field modulates the outcome of biochemical

reactions by influencing the spin state of light-induced radical pairs in macromolecules on

vertebrate retinas [9]. Both theories fail to provide a complete magnetoreception mechanism

which is functional in a natural setting.

In the absence of convincing proof for a 'magnetotactic sensor', a new hypothesis was

proposed. The hypothesis suggests the existence of a symbiosis between magnetotactic

bacteria and magnetotactile vertebrates [10]. Magnetotactic bacteria (MTB) are gram-negative

aquatic prokaryotes which sense and act upon a magnetic field [11]. MTB mineralize

ferromagnetic crystals in unique organelles called magnetosomes, which are arrayed on the

longitudinal body axis of the bacterium, and respond to the ambient magnetic field much like a

compass needle [12]. Here, we examine the effects of antibiotics on the orientation of a

magnetic-sensing migrating passerine in order to provide first experimental support for the

symbiotic magnetic-sensing hypothesis. Led by a straightforward rationale, we explore whether

exposure to an antibiotic substance affects orientation–related, magnetic sensing behaviors of

migrating passerines, using a well-established protocol to quantify the influence of an antibiotic

substance on navigation-related behaviors.

Materials and Methods

The experiment took place during the spring of 2018 (March-May) at the Hula Research Center

in Israel's Hula Valley (33°06'43.8"N, 35°35'08.1"E), a major stopover site for avian migrants on

the Eurasian-African flyway every autumn and spring.

Study animal: Eurasian reed warblers (*Acrocephalus scirpaceus)* are small, night-migrating

passerines that show wide latitudinal variation in their breeding grounds and migration date

[13]. The location of the Hula Valley with respect to Eurasian reed warblers' migration route

suggests that during autumn migrating individuals will show a southward directional tendency

and during spring most individuals are expected to show a northward tendency. Eurasian reed

warblers were caught early in the morning at the Hula Ringing Station using mist nets. We

chose Eurasian reed warblers with high fat scores (≥ 3), as indicating that they are preparing to

migrate soon [14]. We only took specimens with primary wing feather length of ≥ 66 mm, to

make sure that they belong to populations that breed at higher latitudes and are not intending

to breed in or around the valley [13], which would mean their migration has ended. Suitable

specimens were put into cloth bags and taken to the research station.

Experimental procedure

Birds were housed in 30 cm X 23 cm X 40 cm wooden cages inside an air-conditioned container.

Each cage had a wooden perch, two water-filled bottle caps, and a retractable tray for food

provision. Since onset of capture birds had no view of the sky or the outside environment. The

first 24 hours after capture were set as an adjustment period. Birds that were eating and did

not show signs of stress by the evening of the day of capture were considered adjusted, and

were given food (*Tenebrio molitor)* and water ad libitum. At later stages of the experiment, a

bird that appeared stressed was immediately released. On the second day, birds were randomly

divided into a control group and a treatment group, and the first dose of treatment (or water,

according to group) was orally administered using a pipette. Antibiotic substance, dosage, and

method of administration were chosen according to avian veterinary advice, as would be

administered for treating bacterially induced symptoms in the oculonasal region. Enrofloxacin,

also known as Baytril, is a standard, FDA-approved substance for treating bacterial infections in

vertebrates from all groups. It is a broad-spectrum antibiotic, which we tested against MTB

during June 2017 at the molecular and environmental microbiology lab in CEA, Cadarache,

France, with the aid of Dr. Christopher Lefevre. Enrofloxacin proved lethal to various MTB types

including various unidentified morphotypes from the research area and cultivated

Magnetospirillum magneticum strain AMB-1 and *Magnetovibrio blakemorei* strain MV-1. One

dose comprised 2 µl of the solution (or water), and each bird received four doses: one in the

morning and one before sunset, for two consecutive days. On the third day after capture, half

an hour after sunset, birds were removed from their cages and placed in Emlen funnels (plastic

funnels with a rim diameter of 45 cm, covered by a PVC sheet), which were placed in a mesh

enclosure with no view of the sky. Funnels were filmed from above using HIKvision 2.8 IR

cameras (one for each funnel), connected to an HIK vision HD DVR hard drive. Recording started

exactly one hour after sunset with a shot of the identity number and a compass, to indicate

north for later analysis. Filming continued for 90 minutes, of which the first ten allowed for

recovery, and were not analyzed. At the end of the 90-minute experiment birds were removed

from the funnels and released.

Statistical analysis

Each bird's average hop azimuth was transformed to its projection on both axes of the

trigonometric unit circle according to[15]. We used Rayleigh's Z test to examine whether

individuals and groups displayed significant directionality. To verify that the directional data fit

a unimodal distribution, we assessed the distribution using the AIC criterion, and dedicated

methods for circular data, according to [16]. Comparison of the variance in directional

tendencies between groups was done using Levene's test. Activity levels were compared using

two-tailed t-tests.

<u>**Results and Discussion**</u>

When examining orientation of individuals placed in Emlen funnels, we found significant

difference in orientation between the two experimental groups. Individuals from the control

group showed a highly significant south-westward orientation (n = 14, Z = 8.84, P < 0.0001,

Figure 1A), while individuals from the treatment group were more dispersed and showed no

significant orientation (n = 14, Z = 1.33, P > 0.2, Figure 1B). According to the significant

orientation of controls and lack of significant orientation of the treatment group, using Levene's

tests we show that variance of directionality was significantly lower in the control group in both

the north/south and the east/west axes: (north/south axis: F = 6.5, P = 0.01, sd: control = 0.5,

treatment = 0.78, Figure 2A; east/west axis: F = 7.3, P = 0.01, sd: control = 0.38, treatment = 0.6,

Figure 2B). As seen by standard deviation values, controls exhibit smaller variance in all aspects.

To verify that the lack of significant orientation of individuals from the treatment group is not a

result of reduced activity due to exposure to antibiotics, we compared the number of hops

between groups as a means of negating any non-navigation-related effects of antibiotics.

Groups did not differ in the number of hops per individual (T = 0.1, n_c =14, n_t = 14, P = 0.9).

The results obtained from implementation of the well-established Emlen protocol indicate that

Enrofloxacin affects orientation-related behavior during migration restlessness in a night-

migrating passerine. This is the first documentation of such an effect. As seen in Figure 1,

treatment group graphs are similar to control graphs, with the exception that in the first,

individual azimuths are more dispersed over the diagram. Significant orientation is a function of

low directional variance. The significant difference in directional variance between the

experimental groups is exactly what would be expected if antibiotics are detrimental for

orientation. This difference is significant regardless of the lack of significant orientation of the

treatment group. The size and shape of the funnel do not allow birds to do much more than

take a small skip towards the direction in which they have decided to go, meaning that each

hop should represent the directional decision made prior to jumping. This narrows the window

of antibiotic effects further, to the decision-making process itself. Directional decisions during

migration depend on multiple internal and external factors [17]. Emlen funnels allow us to

examine this process in a highly controlled setting. According to widely used, standard

techniques, in our setup tested birds could only rely on magnetic information for the process of

directional decision making, as they had no access to other celestial cues. This narrows our

window of effect further still. The choice of specimens was such that only migrants at late

stages of preparation to migrate were tested. From these, only adjusted, heavily fueled

individuals were tested. A three-day stay in a cage, including treatment and handling prior to

migration could have affected the resulting patterns in various ways. But in this respect,

individuals from both groups underwent exactly the same procedure. This leaves us with the

potential physical effects of the antibiotic, Enrofloxacin, as the internal cause of the differences

between groups. The amount needed to cause any effect to vertebrate cells is two orders of

magnitude larger than the bactericidal, therapeutic amount [18]. This means that any

disruption of the directional, decision-making process caused by our treatment should have

been mediated by a bacterial factor, assuming, as discussed above, that the only external

directional stimuli to which the birds were exposed was magnetic. Namely, the process of

directional decision making involves (at least in part) bacterial factors. We regard these results

as experimental support for the symbiotic magnetic sensing hypothesis.

Today, many processes in multicellular organisms are found to include bacterial involvement

[19–21]. Thus, the conclusions from this experiment are not surprising. An immediate

conclusion would therefore be that antibiotic pollution should be considered a global concern

not only in the context of pathogen resistance and human health [22] but also for processes

involving magnetic field sensing such as migration in the air, on land, or at sea, pollination,

habitat choice, etc.

We expected the directional tendency in spring to be northwards, according to the migration

route of the species in the region [13]. The resulting south-western tendency could be

explained in several ways. Prior work with the study species has shown it to demonstrate axial

behavior, meaning that part of the population (up to 55%) shows reverse directionality with

regard to migration route [23]. Furthermore, "reverse migration" is a known phenomenon,

which has been related to stress, late night migration [24] and the lack of exposure to celestial

information and other directionally significant environmental factors for a long duration prior

to, and during the test [17,25,26]. The issue of exposure to celestial cues during Emlen tests is

debatable, and there are groups working on both methods [17,5]. We aimed to isolate the

magnetic factor of orientation, so obstruction of celestial cues was important. Regardless of the

reason for the seasonally inappropriate directionality, we show significant directional

tendencies in control groups which were absent in treatment groups. Most importantly, we

show a significant increase in directional variance in the antibiotic treatment group. Considering

all the above, this trend could indicate bacterial involvement in navigation-related processes in

a passerine, specifically magnetic sensing.

Acknowledgments:

We would like to thank Shay Agmon of the Hula ringing station and the Hula research center

staff for their assistance throughout the project.

References

1. Dittman, Quinn. 1996 Homing in Pacific salmon: mechanisms and ecological basis. *J. Exp.*
Biol. **199**, 83–91.

2. Hansson BS. 1995 Olfaction in Lepidoptera. *Experientia* **51**, 1003–1027.
(doi:10.1007/BF01946910)

3. Deutschlander ME, Beason RC. 2014 Avian navigation and geographic positioning. *J. F.*
Ornithol **85**, 111–133. (doi:10.1111/jofo.12055)

4. Wehner R, Michel B, Antonsen PER. 1996 Visual Navigation in Insects: Coupling of
Egocentric and Geocentric Information. **140**, 129–140.

5. Wiltschko, W. and Wiltschko. R. 1990 Magnetic orientation and celestial cues in

migratory orientation. *Experientia.* **46**, 342-352. (doi: 0.1007/BF01952167)

6. Narins PM, Lewis ER, Jarvis JUM, O'riain J. 1997 The use of seismic signals by fossorial
Southern African mammals: A neuroethological gold mine. *Brain Res. Bull.* **44**, 641–646.
(doi:10.1016/S0361-9230(97)00286-4)

7. Mouritsen H. 2018 Long-distance navigation and magnetoreception in migratory animals.
Nature. **558**, 50–59. (doi:10.1038/s41586-018-0176-1)

8. Nordmann GC, Hochstoeger T, Keays DA. 2017 Unsolved mysteries: Magnetoreception—
A sense without a receptor. *PLoS Biol.* **15**, 1–10. (doi:10.1371/journal.pbio.2003234)

9. Hore PJ, Mouritsen H. 2016 The radical pair mechanism of magnetoreception. *Annu. Rev.*
Biophys. **45**, 299-344.

10. Natan E, Vortman Y. 2017 The symbiotic magnetic-sensing hypothesis: do magnetotactic
bacteria underlie the magnetic sensing capability of animals. *Mov. Ecol.* **5**, 22-27. (doi:
10.1186/s40462-017-0113-1)

11. Edwards KJ, Simmons SL, Sievert SM, Frankel RB, Bazylinski D. 2004 Spatiotemporal
distribution of marine magnetotactic bacteria in a seasonally stratified coastal salt pond
spatiotemporal distribution of marine magnetotactic bacteria in a seasonally stratified
coastal salt pond. *Appl. Environ. Microbiol.* **70**, 6230–6239.
(doi:10.1128/AEM.70.10.6230)

12. Franckel RB. 1984 Magnetic guidance of organisms. *Annu Rev Biophys Bioeng.* **13**, 85-
103. (doi: 10.1146/annurev.bb.13.060184.000505)

13.   Pearson DJ, Small BJ, Kennerley PR. 2002 Eurasian Reed Warbler: The characters and
variation associated with the Asian form fuscus. *Br. Birds* **95**, 42–61.

14.   Weber TP, Houston AI, Ens BJ, Weber TP, Houston AI, Ens B. 2018 Optimal Departure Fat
Loads and Stopover Site Use in Avian Migration : An Analytical Model Stable URL :
https://www.jstor.org/stable/49969 Linked references are available on JSTOR for this
article : Optimal departure fat loads arid stopover site use in av. **258**, 29–34.

15.   Mora C V, Walker MM. 2012 Consistent effect of an attached magnet on the initial
orientation of homing pigeons, Columba livia. *Anim. Behav.* **84**, 377–383.
(doi:10.1016/j.anbehav.2012.05.005)

16.   Marr PG. 2014 Directional (Circular) Statistics.

http://webspace.ship.edu/pgmarr/Geo441/Lectures/Lec%2016%20%20Directional%20St
atistics.pdf

17.   Åkesson S, Walinder G, Karlsson L, Ehnbom S. 2001 Reed warbler orientation: Initiation
of nocturnal migratory flights in relation to visibility of celestial cues at dusk. *Anim.*
Behav. **61**, 181–189. (doi:10.1006/anbe.2000.1562)

18.   Mitchell MA. 1988 Enrofloxacin. *Drugs Future*. **13**, 305–307.
(doi:10.1358/dof.1988.013.04.62402)

19.   Gonzalez A, Stombaugh J, Lozupone C, Turnbaugh PJ, Gordon JI, Knight R. 2011 The
mind-body-microbial continuum. *Dialogues Clin. Neurosci.* **13**, 55–62.

20.   Logan AC, Katzman M. 2005 Major depressive disorder: Probiotics may be an adjuvant
therapy. *Med. Hypotheses*. **64**, 533–538. (doi:10.1016/j.mehy.2004.08.019)

21.   Collins S, Berrick P. 2009 The relationship between intestinal microbiota and the central
nervous system in normal gastrointestinal function and disease. *Gastroenterology*. **136**
(6), 2003-2014. (doi:10.1053/j.gastro.2009.01.075)

22.   Inyinbor AA, Bello OS, Fadiji AE, Inyinbor HE. 2018 Threats from antibiotics: A serious
environmental concern. *J. Environ. Chem. Eng.* **6**, 784–793.
(doi:10.1016/j.jece.2017.12.056)

23.   Formella M, Przemyslaw B. 2002 Directional preferences of the reed warbler
(Acrocephalus scirpaceus) and the sedge warbler (*A. schoenobaenus*) on autumn
migration at Lake Drużno (Poland). *The Ring*. **24**, 15-29. (doi 10.2478/v10050-008-0076-
0)

24.   Muheim R, Åkesson S, Bäckman J, Sjöberg S. 2017 Magnetic compass orientation
research with migratory songbirds at Stensoffa Ecological Field Station in southern
Sweden: why is it so difficult to obtain seasonally appropriate orientation? *J. Avian Biol.*
48, 6–18. (doi:10.1111/jav.01303)

25.   Able KP. 1982 Field studies of avian nocturnal migratory orientation I. interaction of sun,

wind and stars as directional cues. *Anim. Behav.* **30**, 761–767. (doi:10.1016/S0003-
3472(82)80148-6)

26.   Engels S, Schneider N-L, Mouritsen H, Heyers D, Zapka M, Weiler S, Kishkinev D, Hein CM,
Wild JM, Dreyer D. 2009 Visual but not trigeminal mediation of magnetic compass
information in a migratory bird. *Nature.* **461**, 1274–1277. (doi:10.1038/nature08528)

Figures

Figure 1: Orientation of treated and untreated reed warblers

Figure 2: Variance of directional preference
A: Variance on east/west axis

B: Variance on north/south axis

Figure Legends

Figure 1: Orientation of treated and untreated reed warblers: Each dot represents the average hop azimuth of an individual in an Emlen funnel. Scales are projections of azimuths on x/y axes in the trigonometric unit circle (see Materials and Methods). On the north/south axis, positive values indicate a northward tendency and negative values indicate a southward tendency. On the east/west, positive values indicate east, while negative values indicate west.

Figure 2: Variance of directional preference: Differences in the variance of directional tendency between treatment and control groups. Scales are projections of azimuths on x/y axes in the trigonometric unit circle (see Materials and Methods).