Widespread genetic effects and sex differences play a crucial role in addiction.

Authors:

Michael C. Saul[1,2], Jared R. Bagley[2,3], Lauren S. Bailey[2,3], Udita Datta[1,2], Price E. Dickson[1,2], Rainy Dodd[1,2], Leona H. Gagnon[1,2], Violet M. Kimble[4], Michael Leonardo[1,2], Sam-Moon Kim[2,5], Ashley Olson[1,2], Tyler Roy[1,2], Sarah A. Schoenrock[2,6], Troy Wilcox[1,2], J. David Jentsch[2,3], Ryan W. Logan[2,5], Colleen A. McClung[2,5], Vivek M. Philip[1,2], Laura G. Reinholdt[1,2], Stacey J. Sukoff Rizzo[1,2,7], Lisa M. Tarantino[2,6,8], and Elissa J. Chesler[1,2,*].

Affiliations: [1]The Jackson Laboratory of Mammalian Genetics, Bar Harbor, ME USA; [2]Center for Systems Neurogenetics of Addiction at The Jackson Laboratory, Bar Harbor, ME USA; [3]Binghamton University Department of Psychology, Binghamton, NY USA; [4]Drew University Department of Neuroscience, Madison, NJ USA; [5]University of Pittsburgh School of Medicine Department of Psychiatry, Pittsburgh, PA USA; [6]University of North Carolina-Chapel Hill Department of Genetics, Chapel Hill, NC USA; [7]University of Pittsburgh School of Medicine Department of Neurobiology, Pittsburgh, PA USA; [8]University of North Carolina-Chapel Hill Eshelman School of Pharmacy Division of Pharmacotherapy and Experimental Therapeutics, Chapel Hill, NC USA.

[*]To whom correspondence should be addressed: elissa.chesler@jax.org

Keywords: addiction, sex differences, genetics, cocaine, genomics.

1 Abstract

Though risk for cocaine use disorder, like most complex disease, is subject to considerable

inter-individual variation, the sources of that variation – including genetics and sex – are

frequently ignored in non-human animal studies. Here, we studied both males and females of

eight different inbred mouse strains whose reproducible genomes capture 90% of the genetic

diversity mice. In this population, individual differences explain a substantial proportion of

variance in important cocaine-related behavioral, physiological, and striatum transcriptional

responses traits. Individual differences thus represent a crucial source of biological information

about addiction mechanisms missing in typical studies.

Introduction

Addictions are a highly prevalent complex disease, leading to a public health crisis associated with substantial morbidity and mortality. Illicit substance use disorders afflict 1 in 14 young adults in the United States[1] and drug overdoses are now the leading cause of accidental death among American adults under 55[2]. Genetic variation and sex differences are both known to influence addiction vulnerability; cocaine use disorder is highly heritable ($H^2 = 0.71$)[3] and substance use behaviors show sex differences in both humans and other animals[4,5]. Consequently, the neurobiology underlying addiction cannot be understood completely without consideration of genetic background and sex.

However, like many complex diseases, behavioral and genomic studies of addiction-related phenotypes often utilize only males of a single inbred mouse strain[6] or outbred rat populations confounded by vendor[7]. While these experimental methodologies certainly produce valuable biological insights, ignoring sex differences and genetic variation limits their generalizability and does not reduce experimental noise[8,9]. Inclusion of genetic variability and both males and females in rodent studies has the power to identify addiction-relevant targets, delineate variants' effects on co-regulation and co-expression networks, and define druggable network nodes. Rodent genetic variation does not capture precise human variants, but it can be exploited to determine underlying mechanisms in addiction-relevant processes. As a further benefit, genotypic and phenotypic precision allows genetics studies in rodents to be performed at orders of magnitude lower cost than human GWAS.

To assess genetics, sex, and their interaction on cocaine-related phenotypes in mice, we undertook a large-scale comprehensive evaluation of behavioral, physiological, and brain transcriptomic measures in both male and female mice from the eight inbred founder strains of the Diversity Outbred (DO) mouse heterogenous stock[10]. The genomes represented by these strains capture approximately 90% of the genetic diversity in *Mus musculus*[11].

Results

In the eight founder strains for the DO, we surveyed behavioral and physiological correlates of future cocaine use – including multiple novelty response behaviors, circadian rhythm

phenotypes observed in cells, and reversal learning as a measure of reward learning and

impulsivity. We directly measured cocaine-related behaviors such as initial locomotor

sensitivity to cocaine and intravenous self-administration (IVSA) of cocaine. Finally, we

measured the striatum transcriptome response to cocaine in all eight founder strains with

RNAseq. Our work represents the first time that many of these behavioral, physiological, and

molecular traits have been studied using methods powered to detect sex and genotype effects

and the extent to which these interact by moderating one another.

Response to novelty predicts psychostimulant addiction-related phenotypes in both humans[12]

and mice[13]. We first assessed differences in behavioral traits related to exploration and

Figure 1: Heritable differences in cocaine-related behavioral and physiological traits. All plots show the mean ± the standard error in both sexes with females on the left. **A)** Hole board total entries, **B)** Novel place preference (% time in novel zone), **C)** Open field (% time in center of open field), **D)** Transitions between light and dark in the light-dark box, **E)** Circadian rhythm fibroblast *Bmal1-dLuc* luminescence amplitude, **F)** Reversal learning premature responses to the hole rewarded during acquisition, **G)** Initial locomotor sensitivity to cocaine (cm moved on Day 3 – Day 2), **H)** Cocaine intravenous self-administration number of infusions at 1.0 mg/kg FR1, and **I)** Sessions to acquisition of cocaine intravenous self-administration.

response to novelty. These behavioral risk measures were moderately to strongly heritable

among these strains (H^2 = 0.16-0.55, **Figure 1A-D**, **Supplementary Table S1**). Further, for the

phenotype of total nose pokes in the hole board, there was a significant sex difference ($F_{1,597}$ =

10.69, p = 0.0014). We detected strain-by-sex interactions in the total number of hole board

nose pokes ($F_{7,597}$ = 2.74, p = 0.0083) and in the proportion of distance traveled in the center of

the open field ($F_{7,684}$ = 2.09, p = 0.043), indicating that sex differences in these novelty response

traits are moderated by genetic background. Total nose pokes in the hole board (**Figure 1A**) and

novel place preference (**Figure 1B**) showed the largest differences in wild-derived strains,

demonstrating the importance of their expanded phenotypic range[14]. Transitions between the

sides of the light-dark box also showed extreme phenotypes in wild-derived mouse strains

(**Figure 1D**).

Circadian rhythm and reward-related behavioral phenotypes are co-inherited and rhythm

disruptions are linked to development and progression of substance use disorders[15]. The

molecular clock system directly influences expression of dopamine receptors in the striatum[16].

A cell-based assay on fibroblasts derived from each of the founder strains in which a *Bmal1-*

dLuc reporter was utilized for circadian measurement of luciferase bioluminescence to assess

differences in circadian rhythmicity[17,18]. We found very high heritability of the amplitude of

these rhythmic patterns (H^2 = 0.59, **Figure 1E**), but no significant sex differences or strain x sex

interactions. These results suggest that genetic differences in the molecular clock are one

potential mechanism for individual differences in addiction-related phenotypes.

Reversal learning tasks evaluate impulsive and compulsive behaviors that predict addiction

liability[19]. Within reversal learning paradigms, one measure of impulsivity is the number of

premature responses produced during the reversal phase of the task. Premature reversal

responses in these showed moderate heritability across this panel of mice (H^2 = 0.29, **Figure**

1F), but no significant sex differences or strain x sex interactions. We observed the lowest rate

of premature responding in the wild-derived PWK/PhJ strain.

We next sought to assess individual differences in initial locomotor response to cocaine, a

behavioral phenotype that predicts subsequent drug use in humans[20]. In these diverse mouse

strains, initial cocaine sensitivity showed strong heritability (H^2 = 0.48 for initial sensitivity,

Figure 1G). Wild-derived strains, particularly WSB/EiJ and PWK/PhJ, exhibited the highest initial

sensitivity to cocaine.

Operant drug self-administration procedures directly quantify reinforced responding for drug[13].

Heritability of cocaine IVSA traits in the founders of the DO was very strong ($H^2 = 0.47$ for

sessions to acquisition of IVSA; $H^2 = 0.60$ for total infusions at FR-1 1.0 mg/kg, **Figure 1H-I**). The

high heritability of infusions earned is quite similar to the observed heritability of human

cocaine use disorder[3]. Of note, 129S1/SvImJ mice do not acquire IVSA and do not take any

cocaine during the acquisition phase. The largest phenotypic range manifests in wild-derived

mice; PWK/PhJ mice acquire IVSA behavior very quickly and self-administer the most cocaine –

about four times as much as C57BL/6J – while WSB/EiJ mice self-administer the least amount of

cocaine relative to the other strains that do acquire IVSA behavior. The broad phenotypic range

of volitional cocaine-taking behaviors in these genetically diverse mouse strains represents a

clear opportunity to study mechanisms underlying variation in the physiology of addiction.

Because addiction-related phenotypes are highly heritable and sometimes exhibit sex

differences in a manner dependent on genetics, we next sought to assess heritable differences

in the molecular response to cocaine. Male and female mice of the eight DO founder strains

were given multiple injections of cocaine or saline over 19 days. Samples for bulk RNAseq were

collected from striatum at least 24 hours after the final injection of repeated administration of

either cocaine or vehicle (sham). First, we documented that in the absence of cocaine, most

transcripts show moderate to high baseline heritability across the founder strains (median $H^2 =$

0.29) and that subspecies of origin explains the greatest amount of variation in expression

(**Figure 2A**). Using a linear modeling approach[21,22], we found few expression differences

attributable solely to the effect of cocaine treatment. Instead, significant effects of cocaine

arose in interaction with individual differences such as genetic background. The greatest

quantity of differentially expressed transcripts was observed in the strain-by-sex-by-drug

treatment three-way interaction (1,282 genes at q < 0.01, **Figure 2B, Supplementary Table S2**).

Some strains responded more strongly than others; 129S1/SvImJ and PWK/PhJ had many

expressed genes influenced by cocaine and sex-by-cocaine interactions (PWK/PhJ: 89 genes at q

< 0.01, **Figure 2C**) while the commonly used C57BL/6J strain had very few (4 genes at q < 0.01,

Figure 2C). Because statistical power

was approximately equal for all strains

involved in this study, these differences

likely reflect real individual differences

in the brain's sex-specific responses to

cocaine.

Many significant effects of cocaine on

gene expression would have been

missed in had only single strain been

analyzed. For instance, we identified

differential expression of the serotonin

receptor 1D gene *Htr1d* in the strain-

by-sex-by-cocaine interaction factor.

This gene is upregulated in males of

two strains but downregulated in males

of another (**Figure 2D**).

Discussion

Some of these heritable transcriptome

differences we observed recapitulate

previous work showing heritable

differences in pharmacokinetics of

cocaine[23] – for example, mice closely

related to the high cocaine taking

PWK/PhJ strain are known to have

some of the highest brain

concentrations of cocaine shortly after

Figure 2: Individual differences in striatum cocaine response transcriptomes. **A)** High heritability of baseline transcriptome partitions the subspecies in a multidimensional scaling plot, **B)** In a three-way analysis of strain, sex, and cocaine, many genes showed significant expression changes in response to cocaine in interaction factors; the strain x sex x cocaine interaction factor identified 1,282 DEGs. In contrast, there were very few significant DEGs in the main effect of cocaine. **C)** Two-way models show the magnitude of transcriptional responses to cocaine differs by strain. B6 has few differentially expressed genes while 129 and PWK have many. The y-axis is log_{10}. **D)** The serotonin 1D receptor gene *Htr1d* is differentially expressed in males of 129S1/SvImJ, PWK/PhJ, and CAST/EiJ strains. CAST/EiJ males show the opposite directionality of differential expression to PWK/PhJ and 129S1/SvImJ males.

injection. These pharmacokinetic differences do not account for the strong differences between

strains such as 129S1/SvImJ and A/J, whose brain cocaine pharmacokinetic profiles are very

similar [23]. Further research on highly diverse mice may resolve the mechanisms driving

individual differences between strains with closely matched pharmacokinetic profiles.

Genetic variation is a valuable resource for the discovery of biological mechanisms of

addiction[24,25]. Similar to humans, individual differences among mice greatly influence

behavioral, physiological, and transcriptomic cocaine-related traits. For many of these traits,

individual differences explain a substantial proportion of the variation. Individual variation in

addiction related traits is a largely untapped resource that would have otherwise remained

unknown had we only examined a single inbred strain. Because the vast majority of the

biomedical literature is built on limited genetic diversity, such individual differences represent a

crucial opportunity to find new, clinically relevant genes and mechanisms.

Methods

Standard Operating Procedures

All methodologies used in this work are documented in depth in the Center for Systems

Neurogenetics of Addiction's Standard Operating Procedures (SOPs)

(https://www.jax.org/research-and-faculty/research-centers/systems-neurogenetics/data-

resources). The brief methods appearing below summarize these SOPs and reference specific

SOPs.

Animals

Mice from the following strains were surveyed in these experiments: A/J (JAX stock #000646),

C57BL/6J (JAX stock #000664), 129S1/SvImJ (JAX stock #002448), NOD/ShiLtJ (JAX stock

#001976), NZO/HlLtJ (JAX stock #002105), CAST/EiJ (JAX stock #000928), PWK/PhJ (JAX stock

#003715), and WSB/EiJ (JAX stock #001145). These strains are the founders of the DO

heterogeneous stock and Collaborative Cross recombinant inbred strains. Surveys of these eight

strains can demonstrate statistical heritability patterns that justify further dissection using the

derived resources[10]. The mice used in the Research Animal Facility at The Jackson Laboratory

came from breeding colonies maintained in the Research Animal Facility. These colonies were

derived from production colonies at The Jackson Laboratory and breeders were replaced with

animals from The Jackson Laboratory's production colony at least every five generations. The

mice used in the Jentsch Lab at Binghamton University were shipped to the Jentsch Lab from

either the Chesler Lab colonies or from production colonies in The Jackson Laboratory. The

studies described utilized a total of 1,085 mice (**Supplementary Table S3**).

Mouse Husbandry and Housing

All procedures were approved by the Jackson Laboratory of Mammalian Genetics institutional

animal care and use committee. Mice were housed according to the CSNA animal housing SOP

(https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-

[neurogenetics/csna-animal-housing.pdf](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/csna-animal-housing.pdf)).

Cocaine

Cocaine hydrochloride was provided by the National Institute on Drug Abuse Drug Supply

Program Division of Therapeutics and Medical Consequences (catalog number: 9041-001).

Cocaine was stored in powder form until it was formulated into 0.9% Saline (100 mg/mL clear

solution) in various concentrations specific to each experiment according to their individual

SOPs.

Novelty Response Behavioral Phenotypes

Open field, light-dark box, hole board, and novel place preference behavioral paradigms were

conducted in order with one test per day on consecutive days during the light phase of the

light:dark cycle. Open field data were collected for 60 minutes according to the SOP

(https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-

[neurogenetics/open-field-](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/open-field-)

[assay.pdf?la=en&hash=32DDAFF2B17B2D4961C136C5616C4982AC23EC3B](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/open-field-assay.pdf?la=en&hash=32DDAFF2B17B2D4961C136C5616C4982AC23EC3B)). Light-dark data

were collected for 20 minutes with the mouse starting in the light side of the chamber facing

the dark side according to the SOP (https://www.jax.org/-/media/jaxweb/files/research-and-

[faculty/tools-and-resources/system-neurogenetics/light-dark-](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/light-dark-)

[assay.pdf?la=en&hash=A63CF8D22EB7936CF6C69A3178373981F4016675](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/light-dark-assay.pdf?la=en&hash=A63CF8D22EB7936CF6C69A3178373981F4016675)). Hole board data

were collected for 20 minutes according to the SOP (https://www.jax.org/-

[/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/hole-](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/hole-)

[board-assay.pdf?la=en&hash=EC343A797D37209CF64D34E6031608A511D8E15D](https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/hole-board-assay.pdf?la=en&hash=EC343A797D37209CF64D34E6031608A511D8E15D)). Novel place

preference included a five minute acclimation period to a center chamber, a 10 minute

exposure period to a randomized exposure side, a five minute acclimation period, then testing

for 20 minutes for preference for novel or exposure side of the novel place preference

apparatus according to the SOP (https://www.jax.org/-/media/jaxweb/files/research-and-

faculty/tools-and-resources/system-neurogenetics/novelty-place-preference-

assay.pdf?la=en&hash=B5D2D0FC9028B408E84729C0C8832C580AB8E039). All mice from the

Center for Systems Neurogenetics of Addiction were tested through this novelty pipeline prior

to any other test and were then randomized and assigned into either reversal learning, cocaine

locomotor sensitization, or cocaine intravenous self-administration. The novelty study

produced observations from a total of 783 mice.

Bmal1-dLuc Circadian Rhythm Data

Data for circadian rhythm were measured in primary fibroblast cultures generated from skin

biopsies in the founders. To isolate fibroblasts, ear biopsies (1 mm in diameter) were digested

in Dulbecco's Modified Eagle's Medium (DMEM, HyClone) containing 2.5 mg/ml collagenase D

(Gibco) and 1.25 mg/ml pronase (Millipore) for 90 mins and then plated in DMEM growth

media containing 10% Fetal Bovine Serum (FBS, HyClone), 292 μg/ml L-glutamine (HyClone),

100 units/ml penicillin (Hyclone) and 100 μg/ml streptomycine (HyClone). *Bmal1-dLuc* reporter

was delivered to fibroblasts by lentiviral-mediated gene delivery (VectorBuilder). Following

synchronization of rhythms by 15 μM forskolin (Sigma) for 2 hrs, the temporal patterns of

Bmal1-dLuc bioluminescence was recorded for ~70 secs at intervals of 10 mins over 6-7 days

from fibroblast cultures in DMEM recording media containing 15 μM forskolin, 25 mM HEPES

(Gibco), 292 μg/ml L-glutamine, 100 units/ml penicillin, 100 μg/ml streptomycine, and 10μM

luciferin (Promega) by an automated 32-channel luminometer (Lumicycle, ActiMetrics) in a

standard tissue culture incubator at 32°C. The amplitude of bioluminescence rhythms was

determined from baseline-subtracted data using the damped sine fit and Levenberg-Marquardt

algorithm[26]. The circadian study produced observations from a total of 56 mice.

Reversal Learning

Data for reversal learning were collected using the SOP (https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/reversal-learning-assay.pdf?la=en&hash=8484E47B170462960E11C1FAEEE6FF3CE6FDFC08). The reversal learning data produced observations from a total of 202 mice.

Initial Locomotor Sensitivity to Cocaine

Data for initial locomotor sensitivity were collected as described previously[27] using data from days 1-3 in the SOP for locomotor behavioral sensitization to cocaine (https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/cocaine-locomotor-sensitization-assay.pdf?la=en&hash=9E5D4C248C3BCCAD947C164AE81663C13A77EB0D). Briefly, mice were placed into the open field arena for 30 minutes, removed, and injected i.p. with either saline (days 1-2) or 10 mg/kg cocaine (day 3) and returned to the open field arena for 60 minutes. Distance moved after injection on day 3 minus day 2 was uses as a measure of initial locomotor sensitivity to cocaine. The sensitization study produced observations from a total of 230 mice.

Cocaine Intravenous Self-Administration

Prior to cocaine intravenous self-administration, mice were implanted with a jugular catheter and allowed 10 days for post-operative recovery. In an operant conditioning paradigm, mice were allowed to acquire cocaine self-administration at 1.0 mg/kg, then evaluated for dose-response effects at eight different doses. After a final stabilizing dose at 1.8 mg/kg, responses during seven days of withdrawal were recorded. Finally, cued reinstatement was recorded for two days. Self-administration in these eight mouse strains was performed according to v1.0 of the CSNA's SOP (https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/intravenous-self-administration-ivsa-paradigm.pdf?la=en&hash=FA64135F219C7DF65937A1CF9270301B0E771836). The intravenous self-administration study produced observations from a total of 217 mice.

Data Deposit

Data for each phenotype will be deposited in the Mouse Phenome Database (MPD)[28] upon publication.

Heritability Calculations

For each trait, heritability was calculated from linear models using the isogenic strain as the independent categorical variable using the following equation:

$$h^2 = \frac{MS_{strain}}{MS_{strain} + (n_{mean} - 1) * MS_{resid}}$$

where MS_{strain} is the mean square of the strain effect, n_{mean} is the mean number of samples within each strain, and MS_{resid} is the mean square of the residuals. For the reversal learning data, an additional additive covariate of site was included in the model to account for inter-lab variation. This term was not utilized in the heritability calculation.

For some traits such as number of infusions at FR-1 1.0 mg/kg cocaine self-administration, a single strain such as 129S1/SvImJ showed little to no variation, which may upwardly bias heritability calculations. For these traits, heritability was calculated both with and without the low variance strain. The data reported in the paper rely upon the inclusive calculation, but results of both methods of calculation are reported for completeness (see **Supplementary Table S1**).

RNAseq

Striatum tissue was collected during the light stage of the light:dark cycle between 24 and 48 hours after the final injection in the cocaine behavioral sensitization protocol according to the SOP (https://www.jax.org/-/media/jaxweb/files/research-and-faculty/tools-and-resources/system-neurogenetics/post-sensitization-tissue-collection.pdf?la=en&hash=9E6CD8DEB39606B791A5D25F6CD0611EF14D96A7). Tissue was collected for both sexes of each founder strain exposed to either sham (saline) or 10 mg/kg IP cocaine.

RNA was isolated from striatum tissue using the MagMAX mirVana Total RNA Isolation Kit (ThermoFisher) and the KingFisher Flex purification system (ThermoFisher). Tissues were lysed and homogenized in TRIzol Reagent (ThermoFisher). After the addition of chloroform, the RNA-containing aqueous layer was removed for RNA isolation according to the manufacturer's protocol, beginning with the RNA bead binding step.

RNA concentration and quality were assessed using the Nanodrop 2000 spectrophotometer (Thermo Scientific) and the RNA Total RNA Nano assay (Agilent Technologies). 2μl of diluted 1:1000 diluted ERCC Spike-in Control Mix 1 (Ambion by Life Technologies) was added to 100ng of each RNA sample prior to library construction. Libraries were prepared by the Genome Technologies core service at The Jackson Laboratory using the KAPA RNA Hyper Prep Kit with RiboErase (HMR) (KAPA Biosystems), according to the manufacturer's instructions. Briefly, the protocol entails depletion of ribosomal RNA (rRNA), RNA fragmentation, first and second strand cDNA synthesis, ligation of Illumina-specific adapters containing a unique barcode sequence for each library, magnetic bead size selection, and PCR amplification. Libraries were checked for quality and concentration using the D5000 ScreenTape assay (Agilent Technologies) and quantitative PCR (KAPA Biosystems), according to the manufacturers' instructions.

RNAseq libraries were pooled and sequenced by Novogene in 150 bp paired-end format on an Illumina NovaSeq 6000 sequencer targeting 90 million read pairs per sample. Sequencing achieved a median read depth of 132 million reads. The resultant reads were determined to be of consistently high quality using fastqc v0.11.3 and MultiQC v1.2.

Reads were generated from raw data and demultiplexed using BCL2Fastq v2.18.0.12, concatenated by sample, and aligned with the STAR aligner v2.6.1[29] to the GRCm38 mouse reference genome with v94 of the Ensembl transcriptome. Transcript-level quantification was estimated using RSEM v1.3.0[30] on a transcriptome BAM file produced as an output of this alignment. The data were imported into R v3.5.1 and summarized to the gene level using tximport v1.10.1[31], TMM-normalized using edgeR v3.24.3[21], and imported into limma v3.38.3[32] using the $\log_2$-transformation function voom. We compared multivariate approaches modeling with interaction factors between edgeR and voom+limma approaches and found that voom+limma performs better than edgeR for controlling false negatives. Upon initial

examination of the findings, we identified intermittent contamination with choroid plexus,

which potentially derives from the ventricular aspect of the dorsal striatum. Correcting for this

contamination necessited an additive covariate for choroid plexus consisting of log-mean CPM

values of *Kl* and *Ttr* expression, unambiguous markers for choroid plexus[33]. These values were

$\log_2$ transformed for work in limma. For An overall model for all strains included this choroid

plexus factor as a nuisance variable plus the main effects of strain, sex, and cocaine injection

and all of their interactions. Individual models included the choroid plexus nuisance variable

plus sex, cocaine injection, and sex:cocaine injection interaction. Correction for local false

discovery rates utilized the qvalue package in R v2.14.1[34]. Because brain transcriptional changes

are subtle[35], all results reported are at q < 0.01 with no fold-change cutoff (**Supplementary**

Table S2).

Raw data and transcript-level expression estimates will be deposited in the Gene Expression

Omnibus [36] upon publication (accession number: GSEXXXXX).

Supplemental Material

Supplementary Table S1: Statistical test results for heritability and genotype-by-sex

interactions.

Supplementary Table S2: Differential expression results for all tests discussed.

Supplementary Table S3: Sample sizes for all strain and sex combinations for studies reported.

Raw behavioral data will be deposited in the Mouse Phenome Database upon publication.

Gene expression data will be deposited in the Gene Expression Omnibus upon publication.

All scripts, code, and metadata used for analysis are deposited in GitHub (repository:

github.com/msaul/csna_founders_survey_2020)

Author Contributions

EJC, PED, LMT, SAS, JDJ, RWL, CAM, LGR, VMP, SJSR, and CSNA conceived the studies. SJSR, PED,

JRB, LSB, SAS, RD, ML, AO, TR, TW, and LHG designed and implemented the behavioral

experiments. SMK designed and implemented the circadian transcriptional experiment. JRB,

LSB, UD, PED, ML, SMK, AO, TR, SAS, TW, LHG, VMP, and MCS analyzed the behavioral data.

SMK, MCS, and VMP analyzed the circadian transcriptional data. MCS, VMK, and VMP analyzed

the RNAseq data. MCS, UD, PED, JRB, LSB, SAS, TW, LHG, VMP, and EJC interpreted the

behavioral results. SMK, MCS, and VMP interpreted the results of the circadian transcriptional

experiment. MCS, VMK, VMP, and EJC interpreted the RNAseq results. MCS and EJC wrote the

manuscript.

Acknowledgements

We gratefully acknowledge Technologies service at The Jackson Laboratory Genome for their

expert assistance on the RNAseq experiment, Surgical Services at The Jackson Laboratory for

their work on the intravenous self-administration paradigm, and the NIDA Drug Supply for

providing cocaine. We would further like to thank Stephen Krasinski for critical reading and

comments during manuscript preparation, Rob Williams for helpful discussion on the analysis,

and C. Herbert Pratt for scientific program management. This work was funded by NIH P50

DA039841 (Center for Systems Neurogenetics of Addiction) to EJC, LMT, JDJ, RWL CAM, VMP,

LGR, and SJSR as well as NIH R01 DA037927 to EJC and K99 DA043573 to PED.

Literature Cited

1. SAMHSA. *The 2017 National Survey on Drug Use and Health (NSDUH)*.

https://www.samhsa.gov/data/nsduh/reports-detailed-tables-2017-NSDUH (2017).

2. Kochanek, K. D., Murphy, S. L., Xu, J. & Arias, E. Deaths: Final Data for 2017. *Natl. Vital*

Stat. Rep. **68**, 1–77.

3. Goldman, D., Oroszi, G. & Ducci, F. The genetics of addictions: uncovering the genes. *Nat.*

Rev. Genet. **6**, 521–532 (2005).

4. Becker, J. B. & Chartoff, E. Sex differences in neural mechanisms mediating reward and

addiction. *Neuropsychopharmacology* **44**, 166–183 (2019).

5. Becker, J. B., Perry, A. N. & Westenbroek, C. Sex differences in the neural mechanisms

mediating addiction: a new synthesis and hypothesis. *Biol. Sex Differ.* **3**, 14 (2012).

6. Walker, D. M. *et al.* Cocaine self-administration alters transcriptome-wide responses in the

brain's reward circuitry. *Biol. Psychiatry* (2018).

7. Fitzpatrick, C. J. *et al.* Variation in the Form of Pavlovian Conditioned Approach Behavior

among Outbred Male Sprague-Dawley Rats from Different Vendors and Colonies: Sign-

Tracking vs. Goal-Tracking. *PLoS ONE* **8**, e75042 (2013).

8. Prendergast, B. J., Onishi, K. G. & Zucker, I. Female mice liberated for inclusion in

neuroscience and biomedical research. *Neurosci. Biobehav. Rev.* **40**, 1–5 (2014).

9. Tuttle, A. H., Philip, V. M., Chesler, E. J. & Mogil, J. S. Comparing phenotypic variation

between inbred and outbred mice. *Nat. Methods* **15**, 994–996 (2018).

10. Saul, M. C., Philip, V. M., Reinholdt, L. G. & Chesler, E. J. High-Diversity Mouse

Populations for Complex Traits. *Trends Genet.* **0**, (2019).

11. Roberts, A., De Villena, F. P.-M., Wang, W., McMillan, L. & Threadgill, D. W. The

polymorphism architecture of mouse genetic resources elucidated using genome-wide

resequencing data: implications for QTL discovery and systems genetics. *Mamm. Genome*

18, 473–481 (2007).

12. Ersche, K. D., Turton, A. J., Pradhan, S., Bullmore, E. T. & Robbins, T. W. Drug addiction

endophenotypes: impulsive versus sensation-seeking personality traits. *Biol. Psychiatry* **68**,

770–773 (2010).

13. Dickson, P. E. *et al.* Association of novelty-related behaviors and intravenous cocaine self-

administration in Diversity Outbred mice. *Psychopharmacology (Berl.)* **232**, 1011–1024

(2015).

14. Wahlsten, D., Metten, P. & Crabbe, J. A rating scale for wildness and ease of handling

laboratory mice: results for 21 inbred strains tested in two laboratories. *Genes Brain Behav.*

2, 71–79 (2003).

15. Logan, R. W., Williams III, W. P. & McClung, C. A. Circadian rhythms and addiction:

Mechanistic insights and future directions. *Behav. Neurosci.* **128**, 387–412 (2014).

16. Ozburn, A. R. *et al.* Direct Regulation of Diurnal Drd3 Expression and Cocaine Reward by

NPAS2. *Biol. Psychiatry* **77**, 425–433 (2015).

17. Ramanathan, C. *et al.* Cell Type-Specific Functions of Period Genes Revealed by Novel

Adipocyte and Hepatocyte Circadian Clock Models. *PLoS Genet.* **10**, e1004244 (2014).

18. Kim, S.-M., Neuendorff, N., Chapkin, R. S. & Earnest, D. J. Role of Inflammatory Signaling

in the Differential Effects of Saturated and Poly-unsaturated Fatty Acids on Peripheral

Circadian Clocks. *EBioMedicine* **7**, 100–111 (2016).

19. Izquierdo, A. & Jentsch, J. D. Reversal learning as a measure of impulsive and compulsive

behavior in addictions. *Psychopharmacology (Berl.)* **219**, 607–620 (2012).

20. de Wit, H. & Phillips, T. J. Do initial responses to drugs predict future use or abuse?

Neurosci. Biobehav. Rev. **36**, 1565–1576 (2012).

21. Chen, Y., Lun, A. T. L. & Smyth, G. K. Differential Expression Analysis of Complex RNA-

seq Experiments Using edgeR. in *Statistical Analysis of Next Generation Sequencing Data*

(eds. Datta, S. & Nettleton, D.) 51–74 (Springer International Publishing, 2014).

doi:10.1007/978-3-319-07212-8_3.

22. Phipson, B., Lee, S., Majewski, I. J., Alexander, W. S. & Smyth, G. K. Robust

Hyperparameter Estimation Protects Against Hypervariable Genes and Improves Power to

Detect Differential Expression. *Ann. Appl. Stat.* **10**, 946–963 (2016).

23. Wiltshire, T. *et al.* Initial locomotor sensitivity to cocaine varies widely among inbred mouse

strains: Initial locomotor sensitivity to cocaine in inbred mice. *Genes Brain Behav.* **14**, 271–

280 (2015).

24. Kumar, V. *et al.* C57BL/6N mutation in cytoplasmic FMRP interacting protein 2 regulates

cocaine response. *Science* **342**, 1508–1512 (2013).

25. Ruan, Q. T. *et al.* A Mutation in Hnrnph1 That Decreases Methamphetamine-Induced

Reinforcement, Reward, and Dopamine Release and Increases Synaptosomal hnRNP H and

Mitochondrial Proteins. *J. Neurosci.* **40**, 107–130 (2020).

26. Izumo, M., Johnson, C. H. & Yamazaki, S. Circadian gene expression in mammalian

fibroblasts revealed by real-time luminescence reporting: Temperature compensation and

damping. *Proc. Natl. Acad. Sci.* **100**, 16089–16094 (2003).

27. Schoenrock, S. A. *et al.* Characterization of genetically complex Collaborative Cross mouse

strains that model divergent locomotor activating and reinforcing properties of cocaine.

Psychopharmacology (Berl.) (2020) doi:10.1007/s00213-019-05429-3.

28. Bogue, M. A. *et al.* Mouse Phenome Database: a data repository and analysis suite for

curated primary mouse phenotype data. *Nucleic Acids Res.* gkz1032 (2019)

doi:10.1093/nar/gkz1032.

29. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

30. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. 16 (2011).

31. Soneson, C., Love, M. & Robinson, M. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. *F1000Research* **4**, (2016).

32. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

33. Sathyanesan, M. *et al.* A molecular characterization of the choroid plexus and stress-induced

gene regulation. *Transl. Psychiatry* **2**, e139 (2012).

34. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc. Natl.*

Acad. Sci. **100**, 9440–9445 (2003).

35. Hitzemann, R. *et al.* Introduction to Sequencing the Brain Transcriptome. in *International*

Review of Neurobiology vol. 116 1–19 (Elsevier, 2014).

36. Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets—update. *Nucleic*

Acids Res. **41**, D991–D995 (2012).

424