1

Client processing is altered by novel myopathy-causing mutations in the HSP40 J domain

Melanie Y. Pullen[1], Conrad C. Weihl[2] and Heather L. True[1*]

[1] Department of Cell Biology and Physiology, Washington University School of Medicine, St

Louis, Missouri, United States of America

[2] Department of Neurology, Washington University School of Medicine, St Louis, Missouri,

United States of America

[*] Corresponding author

E-mail: heather.true@wustl.edu (HLT)

Abstract

The misfolding and aggregation of proteins is often implicated in the development and progression of degenerative diseases. Heat shock proteins (HSPs), such as the ubiquitously expressed Type II Hsp40 molecular chaperone, DNAJB6, assist in protein folding and disaggregation. Historically, mutations within the DNAJB6 G/F domain have been associated with Limb-Girdle Muscular Dystrophy type 1D, now referred to as LGMDD1, a dominantly inherited degenerative disease. Recently, novel mutations within the J domain of DNAJB6 have been reported in patients with LGMDD1. Since novel myopathy-causing mutations in the Hsp40 J domain have yet to be characterized and both the function of DNAJB6 in skeletal muscle and the clients of this chaperone are unknown, we set out to assess the effect of these mutations on chaperone function using the genetically tractable yeast system. The essential yeast Type II Hsp40, Sis1, is homologous to DNAJB6 and is involved in the propagation of yeast prions. Using phenotypic, biochemical, and functional assays we found that homologous mutations in the Sis1 J domain differentially alter the processing of specific yeast prion strains, as well as a non-prion substrate. These data suggest that the newly-identified mutations in the J domain of DNAJB6 cause aberrant chaperone function that leads to the pathogenesis in LGMDD1.

Introduction

Molecular chaperones preserve protein homeostasis (1). A deficient chaperone network may lead to protein misfolding and aggregation often associated with protein conformational disorders such as Alzheimer's Disease, Charcot-Marie-Tooth disease, distal hereditary motor neuropathies, and Limb Girdle Muscular Dystrophy, among others (2,3). Limb Girdle Muscular Dystrophy type 1D (LGMD1D), more recently termed LGMDD1 (4), is a disease characterized

by proximal muscle weakness with moderate progression mediated by defective chaperone

function (5). Historically, dominantly inherited disease-associated mutations in the type II Hsp40

co-chaperone DNAJB6 have been found within a 12 amino acid region known as the G/F

domain (6–11). Recently, three novel pathogenic mutations associated with LGMDD1 have been

identified within the J domain of DNAJB6 (12,13).

Since molecular chaperones are highly conserved from yeast to mammals, we used a yeast

model system to study these disease-associated mutations (14–16). The essential yeast Type II

Hsp40, Sis1, is homologous to DNAJB6 and plays an important role in yeast for the propagation

of two prions, [RNQ^+] and [PSI^+] (17–22). Prions in yeast are epigenetic elements that form

when the prion proteins form amyloid aggregates. These prion protein aggregates are propagated

by fragmentation into propagons, which are then passed on to progeny during mitosis (14).

Somewhat counterintuitively, chaperones not only promote prion propagation in yeast, but are

essential for it (14,23–26). At the heart of this chaperone-mediated process is the Hsp40 Sis1,

with the interaction between Hsp40 and Hsp70 chaperones being crucial for prion propagation in

yeast (24,27–32). When chaperone function is defective, prion propagation is impaired (17,33–

36). Thus, the loss of prion propagation provides a read-out for chaperone dysfunction in yeast.

Homologous LGMDD1-associated mutations of the DNAJB6 G/F domain impair the

propagation of various yeast prion strains when expressed in Sis1 (37). We were inspired by this

model to assess the effect of homologous J domain mutants in Sis1 on chaperone function.

Fortunately, the yeast model system allows for a wide array of phenotypic, biochemical, and

functional assays to assess the effect of these mutations on chaperone function (14,38). As such,

we assessed the ability of the disease-associated mutant chaperone to process two specific clients

of Sis1, Sup35 and Rnq1, which form the prion elements [PSI^+] and [RNQ^+], respectively. In

addition, we performed functional assays with a non-prion client of the yeast chaperone

machinery, firefly luciferase (FFL) (39). In conjunction, these assays elucidated nuances in client

processing due to disease-associated mutations in a manner that would not be possible by

assessing a single client protein or using a limited number of assays.

Here, we present evidence that disease-associated mutations in the J domain cause functional

defects that differentially impair processing of native proteins in a client and conformer specific

manner.　Our work suggests that mutations in the J domain are responsible for altered protein

processing and, potentially, disease pathogenesis.

75　Materials and methods

76　Protein sequence alignment structure modeling

Protein sequences for DNAJB6b orthologs were found on UniProt and subsequently aligned

to homologous disease-causing J domain mutations in Sis1. Sequences were visualized using

BioEdit (77) with multiple sequences aligned through the use of ClustalW2 (77–79). Mutant

DNAJB6b structures were generated through the use of using I-TASSER (80–82) and visualized

using PyMOL (83). Structural comparisons and protein threading of variants to wild-type protein

structures was performed using wild-type Sis1 (PBD: 4RWU) (21).

84　Yeast cultures, transformations and treatments

All experiments were performed in derivates of 74-D694 (*ade1-14 his3-Δ200 leu2-3,112*

trp1-289 ura3-52). Yeast strains are [*PSI*$^+$], [*psi-*], [*RNQ*$^+$], or [*rnq-*], kindly gifted by S.

Liebman (50,84) and J. Weissman (51). Yeast were cultured using YPD (1% yeast extract, 2%

peptone, 2% dextrose), 1/4 YPD (0.25% yeast extract, 2% peptone, 2% dextrose) or synthetic

defined (SD) media (0.67% yeast nitrogen base without amino acids, 2% dextrose) lacking

specific amino acids using standard techniques. In order to study the function of *sis1* mutants in

the absence of wild type *SIS1*, previously described plasmid shuffle strains carrying pRS316-

SIS1 (37) were transformed with pRS314 (85) carrying *sis1* mutants. We selected for colonies

that lost wild type *SIS1* through plasmid shuffle on plates containing 5-fluoroorotic acid (5-

FOA). Plasmid transformations were performed by the PEG/LiOAC method (86). Plasmid

pRS316-Sis1 was a kind gift from E. Craig (17,18). Plasmid 316-GPD-Lux was a kind gift from

J. Weissman (67). Other plasmids are described below and were constructed using standard

molecular techniques. For curing of prion strains, yeast were passaged twice on 3mM guanidine

hydrochloride (GdnHCl) plates, then grown on complete media without GdnHCl for use in

assays.

Plasmid construction

Oligonucleotides used for site-directed mutagenesis are listed in S1 Table. Using pRS314-

SIS1, the J domain LGMDD1 mutations were created by site-directed mutagenesis using the

Agilent QuikChange II XL Site-Directed Mutagenesis Kit, as per manufacturer's instructions

along with the following oligonucleotides: 1890 and 1891 (S49V), 1892 and 1893 (E53A), 1894

and 1895 (N56L). Primer sequences were generated using Agilent's online primer design

program. Mutagenesis was confirmed by sequencing the entire coding region of *SIS1*.

Yeast phenotypic assays

For yeast spottings, cultures were grown overnight in selective media. Cultures were pelleted,

washed, and suspended in water to an optical density of 0.1. The normalized yeast solutions were

pipetted into a 96-well plate, and serial dilutions (1:5) were made using a multichannel pipette.

Yeast were spotted onto plates using an ethanol-sterilized 48-pin replicator. To monitor cell

growth or the [PSI^+] status of yeast cells with the spread plate assay, overnight cultures were

normalized by A600, serially diluted 5-fold, and spotted on the indicated media. [PSI^+] status

was assessed by colony color on large 145mm plates in rich ¼ YPD media. Plates were

incubated for 5 days at 30°C followed by overnight incubation at 4°C for additional color

development.

Protein analysis

For boiled gel assays, yeast strains were cultured overnight. Cells were lysed with glass beads

in buffer (25mM Tris-HCl pH7.5, 100mM NaCl, 1mM EDTA, protease inhibitors) and pre-

cleared at 6,000 rpm for 1 minute at 4°C. Protein concentration of cell lysates was normalized

using a Bradford assay and mixed with SDS-Page sample buffer (200mM Tris-HCl pH 6.8, 4%

SDS, 0.4% bromophenol blue, 40% glycerol). Samples remained un-boiled and were loaded on a

12% polyacrylamide gel and run under constant current of 110V until the dye front migrated

halfway through the resolving gel. The current was then stopped and, with the assembled gel

remaining intact within the glass plates, they were sealed in plastic and boiled upright for 15mins

in a 95-100°C water bath. After boiling, gels were removed from the plastic cover and were

reinserted in the SDS-PAGE apparatus, where voltage was re-applied until the dye migrated to

the bottom of the gel. SDS-PAGE was followed by standard western blotting with Sup35, Sis1,

and Pgk1 antibodies. Sedimentation analysis for Rnq1 was performed as previously described

(37,48). Semi-denaturing agarose gel electrophoresis (SDD-AGE) for [*PSI*⁺] was performed as

previously described (37,87).

Luciferase refolding

Luciferase refolding assays were performed as previously described (68). To monitor the

ability of Sis1 to enhance refolding of luciferase *in vivo*, [*PSI*⁺] or [*rnq-*] yeast strains were

transformed with the plasmid pRS316-GPD-Lux. Cultures were grown overnight in selective

media and back diluted to an optical density of 0.3 in 8mL plastic culture tubes. Subsequently,

cyclohexamide was added to a final concentration of 10ug/mL. Treated samples were then

subjected to heat shock at 42°C for 22 minutes. Meanwhile, control samples were plated in 98-

well clear bottom plates and kept at 30°C. After heat shock, 200ul of each sample was plated in

triplicate on 98-well clear bottom plates. All cultures were allowed to recover at 30°C.

Luminometer readings were taken at 30, 60, 90 and, 120 minutes. For data analysis, each

triplicate was averaged. 2way ANOVA was performed using GraphPad Prism version 8.1.1 for

Windows, GraphPad Software, La Jolla California USA, www.graphpad.com.

Statistical methods

150   Data are reported as the means SEM of at least three independent experiments. Comparisons

151   between multiple groups were performed by one-way analysis of variance test (ANOVA). For

152   comparison between two groups, the paired t-test was used. In all cases, $p < 0.05$ was considered

153   significant. GraphPad Prism 8.1.1 software (GraphPad Software) was used for analysis.

Antibodies and reagents

The following antibodies were used for Western Blots: rabbit anti-Sup35, dilution 1:1000 (True lab); rabbit anti-Sis1, dilution 1:5000 (Cosmo Bio, COP-080051); rabbit anti- Rnq1-11, dilution 1:2500 (True lab); mouse anti-Pgk1, dilution 1:2000 (Abcam, ab113687). For the firefly luciferase refolding assay, D-Luciferin (Sigma, L9504) was used as a substrate.

Results

Homologous mutations in the Sis1 J domain are conserved and potentially alter J domain and G/F domain interaction

Recently, novel mutations located within a 7 amino acid region of the DNAJB6 J domain have been reported to cause dominant distal myopathy (12,13). These variants were identified in five different families presenting with both distal and proximal muscle weakness. Initial studies examined these variants in patient tissue and cell culture studies in an attempt to elucidate whether these mutations lead to functional defects in DNAJB6's anti-aggregation activity. From these studies, the recently identified DNAJB6 variants DNAJB6-A50V, DNAJB6-E54A and DNAJB6-S57L have been categorized as pathogenic, likely pathogenic, and variant of unknown significance, respectively (12).

To further examine the effect of these disease-associated variants on myopathy, we performed protein threading to identify homologous mutations in the yeast Hsp40 Sis1. Our comparative modeling identified homologous J domain mutations in Sis1 to be Sis1-S49V (homologous to DNAJB6-A50V), Sis1-E53A (homologous to DNAJB6-E54A), and Sis1-N56L (homologous to DNAJB6-S57L) (Fig 1A). In addition, our structural model recapitulates the observation that

mutations in the J domain have the potential to disrupt important intramolecular interactions

previously shown to stabilize the interface between the G/F and J domains (Fig 1B) (12,21).

Fig 1. Novel LGMDD1 disease-associated mutations were identified in the J-domain of

DNAJB6. (A) Amino acid sequence comparison of DNAJB6 and orthologs, including the yeast

Hsp40, Sis1, aligned using ClustlW2. (B*)* Protein structure identifying homologous LGMDD1-

associated mutations in the Sis1 J domain.

Homologous mutations in the J domain of Sis1 differentially impair [PSI+] prion propagation

To assess the implications of these disease-causing variants on chaperone function, we turned

to prion propagation models in yeast. Such models have been used extensively to study

chaperones and understand the deleterious effects of disease-associated mutations on protein

folding (40–46). In yeast, prions are naturally occurring, self-propagating protein structures that

require chaperone networks for their efficient propagation (14). Defects in chaperone function

disrupt this process and, therefore, prion propagation (27,33–36,47,48). In order to ask whether

these disease variants disrupt client processing, we studied two known Sis1 substrates: the

translation termination factor Sup35, which can aggregate to form the $[PSI^+]$ prion, and Rnq1,

which can aggregate to form the $[RNQ^+]$ prion (14,29).

In an effort to understand the effect of homologous J domain mutations on $[PSI^+]$ prion

propagation we performed experiments utilizing strains harboring the *ade1-14* allele that has a

premature stop codon in the *ADE1* open reading frame (14). When Sup35 misfolds and

aggregates in the $[PSI^+]$ state, nonsense suppression of the premature stop codon in *ade1-14*

occurs and the adenine biosynthesis pathway is completed. Thus, [*PSI*$^+$] colonies are white in

color and can grow on media lacking adenine. Conversely, if cells are [*psi-*], adenine

biosynthesis is incomplete, resulting in cells that are unable to grow in media lacking adenine

and an accumulation of red pigment in cells grown on rich media. This colorimetric phenotypic

assay has been utilized to demonstrate the effect of LGMDD1-causing G/F domain mutations on

[*PSI*$^+$] propagation (37). Here, we make use of four different [*PSI*$^+$] strains (weak, strong, Sc37,

and Sc4) (49,50). Each prion strain is a different self-propagating conformation of the same

protein sequence and they are differentiated by the strength of the nonsense suppression

phenotype and stability of their mitotic inheritance (49,51–53). Yeast containing *ade1-14* that

propagate stronger [*PSI*$^+$] strains (strong and Sc4) display higher nonsense suppression and are

lighter in color. Yeast propagating weaker [*PSI*$^+$] strains (weak and Sc37) are darker in color due

to decreased nonsense suppression (better translation termination). Sis1 is required for yeast

viability (54). Therefore, to perform these experiments, we used *sis1*Δ yeast (covered by a

plasmid expressing Sis1) propagating different [*PSI*$^+$] strains (weak, strong, Sc37, and Sc4) and

replaced *SIS1* with homologous LGMDD1 mutant *SIS1* constructs or wild-type *SIS1* (as a

control) (Fig 2A). In doing so, we observed viability was not affected in these strains expressing

the mutants.

Fig 2. Homologous LGMDD1-associated mutations in the Hsp40 Sis1 differentially impair

propagation of [PSI$^+$] strains. (A) Yeast 74-D694 *sis1*Δ [*PSI*$^+$] strains expressing wild-type

SIS1 or mutated *SIS1* constructs were serially diluted 5-fold and spotted onto 1/4 YPD media and

SD-Ade to monitor nonsense suppression of the *ade1-14* allele (n=3). (B) [*PSI*$^+$] colonies

expressing the indicated constructs were isolated, grown in liquid YPD overnight, and plated on

large 1/4 YPD spread-plates. An average of 626 colonies were counted and scored for strong

[*PSI+*], weak [*PSI+*] or [*psi-*]/very weak phenotype. Data were collected from three separate

biological replicates. (**C**) Representative western blot of yeast *sis1*Δ [*PSI+*] strains expressing

wild-type *SIS1* or mutated *sis1* constructs. Cells were lysed and subjected to a boiled gel assay in

order to display the amount of aggregated and monomeric Sup35 with antibodies against Sup35.

Pgk1 is shown as a loading control. While we observed some reproducible differences in the

percentage of monomeric Sup35 as compared to wild-type, we found these differences were not

statistically significant. Standard protein markers are inappropriate for this particular assay and

therefore are not shown (n=4-7). All samples were analyzed under the same experimental

conditions.

Interestingly, we found that expression of Sis1-S49V in the weak [*PSI*$^+$] strain, but not the

Sc37 [*PSI*$^+$] strain, altered the weak strain phenotype, as indicated by the lighter color on rich

media and robust growth in selective media as compared to wild-type (Fig 2A). In comparison to

the other J domain mutants, the stronger nonsense suppression phenotype we observed is unique

to expression of Sis1-S49V in the weak [*PSI*$^+$] strain. This suggests altered processing of Sup35

by Sis1-S49V occurs with conformational specificity. We questioned whether expression of

Sis1-S49V could have altered the conformation of Sup35 in weak [*PSI*$^+$] in a manner that leads

to the propagation of a strong [*PSI*$^+$] strain. Intrigued by this result, we introduced pRS316-Sis1

(wildtype) to this strain and re-assessed its [*PSI*$^+$] phenotype (S1 Fig). We found this reverted the

phenotype and was comparable to wild-type controls, suggesting that Sis1-S49V expression does

not alter the [*PSI*$^+$] strain. Previous work demonstrated that a shortage of cytosolic Sis1 may lead

to more efficient transmission of propagons to daughter cells thus "strengthening" the prion

phenotype (55). Thus, we hypothesize this may occur when Sis1-S49V is expressed in a weak

[*PSI*$^+$] strain.

This assay also showed that the expression of either Sis1-E53A or Sis1-N56L altered the

propagation of the weaker [*PSI*$^+$] strains (weak and Sc37), albeit in a different manner and to

varying degrees. This is indicated by the phenotypic change towards a red color (less nonsense

suppression) on rich media and lack of growth on media lacking adenine (Fig 2A). Similarly,

when we introduced a wild type copy of *SIS1* to these phenotypically red strains, we identified a

partial rescue of the weak [*PSI*$^+$] phenotype (S1 Fig) indicating these Sis1 mutants, although

phenotypically [*psi-*], result in propagation of cryptic [*PSI*$^+$] propagons (56).

Complementarily, we used this assay quantitatively by phenotypically scoring large numbers

of colonies on rich media spread plates (Fig 2B). This assay allows for a more detailed

understanding of any changes in [*PSI*$^+$] propagation. We scored colony color as light pink

(strong [*PSI*$^+$]), dark pink (weak [*PSI*$^+$]) or red (very weak [*PSI*$^+$] or [*psi-*]) in yeast expressing

wild type Sis1 or the homologous LGMDD1 mutants. Indeed, through this assay we found that

the expression of these mutants phenotypically altered prion propagation and identified a

phenotypic variation that was not previously appreciated in the spotting assay. We observed a

wider range of phenotypic variation in the weaker strains than in the stronger [*PSI*$^+$] strains,

which rarely convert to a weaker phenotype (57,58). However, an increase in the amount of

darker pink colonies was observed when Sis1-S49V and Sis1-N56L were expressed in the

stronger [*PSI*$^+$] strains. Similar results were observed when Sis1-E53A and Sis1-N56L were

expressed in the Sc4 [*PSI*$^+$] strain (Fig 2B).

To further explore these results with respect to the solubility of the prion protein, we

performed boiled gel assays to examine the relative levels of aggregated and monomeric Sup35.

This assay consists of loading unboiled samples onto a sodium dodecyl sulfate–

polyacrylamide gel electrophoresis (SDS-PAGE) gel. By doing so, non-denatured aggregates are

too large to enter the resolving gel while monomeric Sup35 readily enters the gel due to its size

and lack of aggregation. Halfway through electrophoresis, the gel itself is boiled and then run

again, which allows previously aggregated protein to enter the resolving gel. The boiled gel

assay provides a clear separation of monomeric and aggregated species of Sup35 and is easily

detected through subsequent western blots (59). In order to examine the distribution of Sup35

between the aggregated and monomeric forms, we grew cultures from colonies representative of

the phenotypic distribution observed in our spread plate assay.

Expressing homologous J domain mutations in the strong or Sc4 [PSI^+] strains did not alter

Sup35 distribution nor phenotype significantly (Fig 2C). Interestingly, the distribution of Sup35

between monomer and aggregate does not always correlate to phenotype in the weaker [PSI+]

strains (many replicates of these assays were performed to verify reproducibility). Of note, we

previously described a chaperone mutant that demonstrated differences in the ability to relate

SDS-resistant or small protein aggregates to phenotype (56). Moreover, we found that

expressing Sis1-S49V in the weak [PSI^+] strain led to Sup35 being found mostly in the

aggregated state (Fig 2C), coinciding with our phenotypic assay. With regards to expression of

Sis1-E53A in the weak [PSI^+] strain, we were surprised to find little Sup35 in the monomeric

fraction, while the expression of Sis1-N56L results in most of the Sup35 in the monomeric

fraction. Interestingly, expressing Sis1-S49V, Sis1-E53A or Sis1-N56L in the Sc37 [PSI^+] strain

showed a reproducible yet not statistically significant increase in monomeric Sup35 while

maintaining a modest level of aggregated Sup35. Thus, there are differences with respect to the

interaction between the Sis1 mutants and Sup35 in the weaker [PSI+] strains and how propagons

and large aggregates correlate to phenotype.

Homologous mutations in the J domain of Sis1 differentially impair [RNQ+] prion propagation

We then asked whether mutations in the Sis1 J domain would also affect another known client of Sis1, the [RNQ^+] prion. In a manner similar to that of the previous experiments, we utilized various [RNQ^+] prion strains (*rnq-*, low, and very high) and expressed each of our Sis1 mutants. Here, we observed reduced growth when Sis1-S49V was expressed in [*psi-*] strains that we did not observe in the same yeast strains harboring the [*PSI*+] prion. Thus, we performed spottings (using 5-fold serial dilutions) and confirmed that expression of Sis1-S49V, but not Sis1-E53A or Sis1-N56L, resulted in a growth defect (Fig 3A). This effect was only observed when wild-type Sis1 was removed through plasmid shuffle and occurred when Sis1-S49V was expressed in either [RNQ^+] or [*rnq-*] strains. This indicated the defect in growth was independent of [RNQ^+] and it was possibly impacted by the loss of [PSI^+]. Because Sis1 is essential for yeast growth, mutations in Sis1 have been shown to affect viability (20,54,60,61).

Fig 3. Homologous LGMDD1-associated mutations in the J domain of Sis1 differentially impair propagation of [RNQ$^+$]. (A) Yeast *sis1Δ* cells harboring the indicated constructs and propagating low [RNQ^+], very high [RNQ^+] or [*rnq-*] were serially diluted 5-fold and spotted onto media in order to express the indicated *SIS1* constructs and select for either loss of wild-type Sis1 (-Sis1) or co-expression of wild-type Sis1 (+Sis1). Cells were also spotted on rich media (YPD) as a control for overall growth (n=3). (B) Sedimentation assay to separate Rnq1 into soluble and insoluble fractions with expression of the indicated constructs (n=6). Total (T), soluble (S) and insoluble (I) fractions were subjected to SDS-PAGE and western blot probing for

Rnq1. (C) SDD-AGE assay for the indicated constructs in low [RNQ^+] and very high [RNQ^+] strains (n=4-6). All samples were analyzed under the same experimental conditions.

Unlike the nonsense suppression assay that discerns [PSI^+] strains, strains of the [RNQ^+] prion are characterized by the rate at which they induce the formation of [PSI^+], ranging from low to very high rates (62–65). Therefore, we performed two different biochemical assays to assess [RNQ^+] propagation. First, we performed a sedimentation assay (59) which consists of separating the soluble and insoluble protein fractions using ultracentrifugation to assay the relative solubility of Rnq1. These fractions are subjected to SDS-Page and probed with an antibody specific to Rnq1. We performed this assay using two [RNQ^+] strains, low [RNQ^+] and very high [RNQ^+]. Using this assay, we identified a defect in [RNQ^+] propagation when Sis1-S49V is expressed in both low [RNQ^+] and very high [RNQ^+] strains (Fig 3B). In comparison, there is only a slight change in Rnq1 solubility when Sis1-E53A or Sis1-N56L are expressed, indicating that J domain variants differentially impair processing of client conformers, albeit not completely.

Furthermore, while the sedimentation assay allows us to detect pelletable aggregates, these aggregates differ in their SDS sensitivity. In an effort to elucidate whether Rnq1 aggregate size or SDS resistance was altered by expression of the mutants, we performed a semi-denaturing detergent agarose gel electrophoresis (SDD-AGE) (Fig 3C). This assay allows for the visualization of SDS-resistant aggregates such as aggregated prion conformers (66). Strikingly, we observed a dramatic loss of SDS-resistant Rnq1 aggregates in the low [RNQ^+] and very high [RNQ^+] strains when Sis1-S49V was expressed. We also observed a slight reduction in SDS-resistant Rnq1 aggregates in low [RNQ^+] strains expressing Sis1-E53A, while low [RNQ^+] strains expressing Sis1-N56L appeared unaffected. In addition, very high [RNQ^+] strains expressing

340   Sis1-E53A or Sis1-N56L showed a decreased in SDS-resistant Rnq1 aggregates. Altogether,

341   these biochemical assays present varying degrees of altered [RNQ^+] propagation that occur in a

342   conformer-specific manner when disease-associated J domain mutants are expressed in these

343   strains.

Disruption in prion propagation is due to a defect in substrate

refolding

We observed different degrees of alteration in prion propagation due to the expression of the

homologous LGMDD1 mutations in the J domain, which vary depending on the client and its

conformation. To further understand the effect of these disease-associated mutations, we

assessed how they impact protein folding of a non-prion substrate *in vivo*. Thus, we utilized the

well-established firefly luciferase (FFL) refolding assay (67,68) (Fig 4). FFL is a non-prion

substrate of the Hsp40/Hsp70/Hsp104 chaperone machinery and is denatured in cells by heat

shock. When FFL misfolds and aggregates it requires the Hsp40/Hsp70/Hsp104 chaperone

machinery to efficiently refold. Since Hsp104 is required for efficient refolding of FFL, we used

a *hsp104Δ* strain expressing FFL as a negative control. We observed a significant defect in FFL

refolding when Sis1-S49V was expressed in either a [PSI^+] [$RNQ+$] (Fig 4) or [*psi-*] [*rnq-*] strain

(S2A Fig), but not when other J domain variants are expressed. Of note, we did not observe a

decrease in Sis1-S49V steady-state levels when this construct was expressed in Sc37 [PSI^+]

strains (S3 Fig). We subjected other known G/F domain mutants expressed in a chimeric protein

where the G/F domain of Sis1 is replaced by that of DNAJB6, referred to as SDSS (37), to this

assay and saw no significant differences in luciferase refolding as compared to wild type (S2B

Fig). Hence, these results highlight the importance of substrate recognition – both with the client

itself, as well as conformer specificity. Most importantly, these findings emphasize the

distinction of assessing native chaperone clients as these mutants displayed varying degrees of

altered prion propagation in our assays. Given this specificity, care should be taken when using

certain assays, or a limited number of them, when trying to understand and confirm whether a

potentially pathogenic chaperone variant is disease-causing or not.

Fig 4. A homologous LGMDD1-associated mutant in Sis1 exhibits impaired substrate

refolding. (A) The capability for refolding luciferase was measured in Sc37 [PSI^+] $sis1\Delta$ yeast

strains harboring the indicated construct along with a plasmid expressing firefly luciferase (FFL).

Yeast were normalized, treated with cycloheximide and subjected to heat shock at 42°C for 22

minutes, followed by recovery at 30°C. Luminescence was measured at the indicated timepoints

during recovery and normalized to luminescence of samples without heat shock treatment. The

results represent the amount of luciferase refolding plotted as the percentage of recovery and

represented as mean SEM (n=6). Each triplicate was averaged, (two-way ANOVA was

performed; **p indicates a significant difference in Sis1-S49V FFL refolding relative to WT (p=

0.002) at the indicated timepoint; ***p indicates a significant difference in Sis1-S49V FFL

refolding relative to WT at the indicated timepoint (p<0.0001)). All samples were analyzed

under the same experimental conditions.

Stability of a mutated Hsp40 chaperone is rescued by expression of

the [PSI+] prion

While assessing the effect of the J domain LGMDD1 mutations on [RNQ^+] propagation we

identified a growth defect that only occurred with expression of the Sis1-S49V variant (Fig 3A).

Interestingly, we observed this defect when Sis1-S49V was expressed in [RNQ^+] [psi-] strains,

but not when expressed in [PSI^+] strains (Figs 2A and 3A). Because *SIS1* is essential

(20,54,60,61), we were curious whether this growth defect was due to any change in steady-state

expression levels. We assessed steady state Sis1 levels by western blot (Fig 5A) and found a

significant decrease in the level of Sis1-S49V in various [rnq-] and [RNQ^+] strains (Fig 5B). We

hypothesize that this decrease in steady-state levels is responsible for the growth defect.

Fig 5. The LGMDD1-associated J-domain mutation Sis1-S49V has reduced steady state

levels in the absence of [PSI^+]. (A) Representative western blot showing expression of the

indicated constructs in [rnq-], low [RNQ^+], and very high [RNQ^+] strains. (B) Results represent

the quantified protein levels of Fig 5A displayed as their ratio relative to WT. (C) Representative

western blot of strong [PSI^+] *sis1Δ* yeast expressing the indicated construct (left). The specified

strong [PSI^+] and Sc37 [PSI^+] strains expressing the indicated constructs were passaged twice on

3mM guanidine hydrochloride (GdnHCl) (cured), then grown on complete media without

GdnHCl and lysed for SDS-PAGE followed by western blot (right). Pgk1 served as a loading

control. (D) Results represent the quantified protein levels of Fig 5C displayed as their ratio

relative to WT. For all Western Blot analyses, intensities were quantified and plotted as the mean

SEM (n=3). Independent experiments indicating a significant difference in Sis1 expression levels

relative to WT (paired t-test; n.s. nonsignificant; *$p<0.05$; **$p<0.01$; ***$p<0.001$). All samples

were analyzed under the same experimental conditions.

To confirm this decrease is due to a lack of [PSI^+] presence, we set out to recapitulate this

phenomenon by curing [PSI^+] strains of its prion form. Curing of [PSI^+] was performed by

growth on media containing guanidine hydrochloride (GdnHCl), which inactivates the ATPase

activity of Hsp104 and hinders the replication of [*PSI*$^+$] seeds (69). After curing both a strong

[*PSI*$^+$] and a Sc37 [*PSI*$^+$] strain (each expressing Sis1-S49V) we performed western blots (Fig

5C) and identified a significant decrease in Sis1 steady-state levels (Fig 5D). This decrease was

not observed with expression of wild type Sis1 nor the other LGMDD1 J domain variants. The

decrease in steady-state levels recapitulates what we observed in [*psi-*] strains expressing Sis1-

S49V (Figs 5A and 5B). This striking result suggests the Sis1 client, Sup35, in its aggregated and

[*PSI*$^+$]-propagating form, interacts with this particular Hsp40 mutant in a way that stabilizes the

steady-state level of Sis1-S49V. Of note, this decrease in Sis1 steady-state levels has not been

observed in the context of LGMDD1-associated mutations found in the G/F domain. This

interesting finding is the first known instance wherein a potentially unstable mutant chaperone is

stabilized by the presence of a specific client protein in its aggregated form.

Discussion

LGMDD1 is a myopathy historically characterized by mutations within the G/F domain of the

chaperone DNAJB6 (6–9,11). Here we used the yeast model system, a homologous chaperone,

and native chaperone clients to demonstrate that recently reported mutations in the J domain of

DNAJB6 alter canonical chaperone function. Specifically, expression of homologous disease-

associated mutations in the yeast Hsp40, Sis1, identified defects in prion propagation to varying

degrees depending on the disease-associated variant, the client protein and its conformation, as

well as a significant defect in refolding of a non-prion client protein. We conclude these

mutations in the J domain of Sis1 lead to aberrant chaperone function, altered protein

homeostasis, and potentially drive a variety of defects in chaperone machinery which may

contribute to pathogenesis.

The function and interaction of the various Hsp40 domains has been a topic of interest for

many years (20,21,70–75). Surprisingly, colleagues identified LGMDD1 patients with novel

variants in the J domain of DNAJB6, a disease previously characterized by mutations in the G/F

domain (12,13). As previously mentioned, LGMDD1 disease-associated mutations in the G/F

domain have been found to impair the processing of specific client conformers. These findings

highlighted the G/F domain as having an important role in substrate regulation and conformer

selectivity (37). By contrast, the J domain of Sis1 has a known role in regulating Hsp70 ATPase

activity (76).

Here we demonstrate a novel finding by which disease-associated missense mutations in the J

domain of an Hsp40 result in the chaperone having client and conformer specificity. Strikingly,

this fits with previous structural studies which have highlighted the importance of the

intramolecular interaction between the J and G/F domains (21). Specifically, a particular amino

acid, E50, in the Sis1 J domain has been shown to interact with the EEVD(HSP70) motif, an

interaction required for both chaperones to partner in protein refolding (21). Moreover, this key

amino acid (E50) forms a salt bridge with R73 which is found in the Sis1 G/F domain. We and

others (12,21) believe this interaction is crucial and of functional importance for the stability and

interaction between both domains and, ultimately, the interaction between Hsp40 and Hsp70.

In the context of the LGMDD1-associated mutants found in the J domain, all variants appear

to affect prion propagation to varying degrees depending on client and conformation specificity.

Notably, the homologous Sis1-S49V variant appears to have the most deleterious effect on the

chaperone machinery and its function, probably due to its proximity to the E50:R73 salt bridge.

This would also explain why only Sis1-S49V presented with defective substrate refolding in our

luciferase refolding assay. Overall, we hypothesize that the steric hindrance caused by these

missense mutations may obstruct the intramolecular interaction or communication between the J

and G/F domains. Moreover, these findings support the idea that it may be the interaction

between these two domains that is crucial for the specificity and processing of Hsp40 client

proteins.

By assessing the propagation of various Sis1 clients, we found that there are differences in

prion propagation, but that propagation is not completely abolished by expression of these

mutants. When compared to LGMDD1-associated mutations in the G/F domain, for example, we

observe variants in the G/F domain to have a stronger effect on prion propagation in certain

assays, oftentimes demonstrating complete loss of prion propagation (37). As such, although

these disease-associated mutations in the J domain disrupt prion processing and protein

refolding, they do so in a manner that is slightly different from that of G/F domain mutants.

Given the limited knowledge of the clients of DNAJB6 in skeletal muscle and its role in

disease, we hypothesize, as we did for the G/F domain mutations, that distinct variants may be

associated with different levels of disease severity due to impairment of Hsp40 function to

varying degrees, or in different ways. Nonetheless, we are cautious to imply that a more drastic

loss-of-propagation effect would correlate with or be causative of a more severe disease

pathogenesis, as we have observed different degrees of aberrant chaperone function in the

context of these disease-associated variants that are greatly dependent on client and conformation

specificity. It is not uncommon for LGMDD1-associated mutations to have an effect in some but

not all assays or when assaying specific client proteins (12,37). Moreover, as we continue to

understand and classify disease-associated variants as pathogenic, we should consider how

variable clinical outcomes and disease severity may be influenced by factors such as variable

expressivity and sex influences (11). Given these considerations, there is a need for simple

models, such as the yeast system, by which we can understand if and how identified genetic

variants affect chaperone function in order to better understand variable clinical outcomes within

LGMDD1. Currently, we can only show that there are differences in adequate client processing

and refolding although, undoubtedly, further *in vitro* studies are required to further elucidate the

mechanistic cause of these functional defects and how they translate to pathogenesis.

Lastly, we identified a novel phenomenon by which a mutated Hsp40, Sis1-S49V, appears to

have a decreased steady state level of protein expression. This is, to our knowledge, the first

known instance where steady-state expression of a chaperone is dependent on stabilization by a

known substrate. Although serendipitous, this finding not only highlights an important function

of chaperone-substrate interaction, but also hints at an additional role that specific substrates

might play in chaperone function and protein homeostasis overall. Indeed, there is still much to

be learned about co-chaperones, their function, interactions between themselves and with clients,

and their role in disease pathogenesis.

Acknowledgements

494   We thank S. Liebman, J. Weissman and E. Craig for reagents. We also thank J. Daw for

495   assistance with site-directed mutagenesis. We are appreciative of L. Dublin and A. Bhadra for

496   helpful discussions and comments on the manuscript.

References

1. Jee H. Size dependent classification of heat shock proteins: a mini-review. J Exerc Rehabil. 2016 Aug 26;12(4):255–9.

2. Muranova LK, Ryzhavskaya AS, Sudnitsyna MV, Shatov VM, Gusev NB. Small Heat Shock Proteins and Human Neurodegenerative Diseases. Biochem Biokhimiia. 2019 Nov;84(11):1256–67.

3. Hipp MS, Kasturi P, Hartl FU. The proteostasis network and its decline in ageing. Nat Rev Mol Cell Biol. 2019 Jul;20(7):421–35.

4. Straub V, Murphy A, Udd B, LGMD workshop study group. 229th ENMC international workshop: Limb girdle muscular dystrophies - Nomenclature and reformed classification Naarden, the Netherlands, 17-19 March 2017. Neuromuscul Disord NMD. 2018;28(8):702–10.

5. Ruggieri A, Saredi S, Zanotti S, Pasanisi MB, Maggi L, Mora M. DNAJB6 Myopathies: Focused Review on an Emerging and Expanding Group of Myopathies. Front Mol Biosci. 2016;3:63.

6. Couthouis J, Raphael AR, Siskind C, Findlay AR, Buenrostro JD, Greenleaf WJ, et al. Exome sequencing identifies a DNAJB6 mutation in a family with dominantly-inherited limb-girdle muscular dystrophy. Neuromuscul Disord NMD. 2014 May;24(5):431–5.

7. Harms MB, Sommerville RB, Allred P, Bell S, Ma D, Cooper P, et al. Exome sequencing

reveals DNAJB6 mutations in dominantly-inherited myopathy. Ann Neurol. 2012

Mar;71(3):407–16.

8. Sarparanta J, Jonson PH, Golzio C, Sandell S, Luque H, Screen M, et al. Mutations affecting

the cytoplasmic functions of the co-chaperone DNAJB6 cause limb-girdle muscular

dystrophy. Nat Genet. 2012 Feb 26;44(4):450–5, S1-2.

9. Sato T, Hayashi YK, Oya Y, Kondo T, Sugie K, Kaneda D, et al. DNAJB6 myopathy in an

Asian cohort and cytoplasmic/nuclear inclusions. Neuromuscul Disord NMD. 2013

Mar;23(3):269–76.

10. Bohlega SA, Alfawaz S, Abou-Al-Shaar H, Al-Hindi HN, Murad HN, Bohlega MS, et al.

LGMD1D myopathy with cytoplasmic and nuclear inclusions in a Saudi family due to

DNAJB6 mutation. Acta Myol Myopathies Cardiomyopathies Off J Mediterr Soc Myol.

2018 Sep;37(3):221–6.

11. Zima J, Eaton A, Pál E, Till Á, Ito YA, Warman-Chardon J, et al. Intrafamilial variability of

limb-girdle muscular dystrophy, LGMD1D type. Eur J Med Genet. 2019 Apr 27;

12. Palmio J, Jonson PH, Inoue M, Sarparanta J, Bengoechea R, Savarese M, et al. Mutations in

the J domain of DNAJB6 cause dominant distal myopathy. Neuromuscul Disord [Internet].

2019 Nov 19 [cited 2019 Nov 26]; Available from:

http://www.sciencedirect.com/science/article/pii/S0960896619312015

13. Savarese M, Di Fruscio G, Mutarelli M, Torella A, Magri F, Santorelli FM, et al. MotorPlex

provides accurate variant detection across large muscle genes both in single myopathic

patients and in pools of DNA samples. Acta Neuropathol Commun. 2014 Sep 11;2(1):100.

14. Liebman SW, Chernoff YO. Prions in yeast. Genetics. 2012 Aug;191(4):1041–72.

15. Richter K, Haslbeck M, Buchner J. The heat shock response: life on the verge of death. Mol

Cell. 2010 Oct 22;40(2):253–66.

16. Bukau B, Weissman J, Horwich A. Molecular chaperones and protein quality control. Cell.

2006 May 5;125(3):443–51.

17. Aron R, Higurashi T, Sahi C, Craig EA. J-protein co-chaperone Sis1 required for generation

of [RNQ+] seeds necessary for prion propagation. EMBO J. 2007 Aug 22;26(16):3794–

803.

18. Lopez N, Aron R, Craig EA. Specificity of class II Hsp40 Sis1 in maintenance of yeast prion

[RNQ+]. Mol Biol Cell. 2003 Mar;14(3):1172–81.

19. Sondheimer N, Lopez N, Craig EA, Lindquist S. The role of Sis1 in the maintenance of the

[RNQ+] prion. EMBO J. 2001 May 15;20(10):2435–42.

20. Yan W, Craig EA. The glycine-phenylalanine-rich region determines the specificity of the

yeast Hsp40 Sis1. Mol Cell Biol. 1999 Nov;19(11):7751–8.

21. Yu HY, Ziegelhoffer T, Osipiuk J, Ciesielski SJ, Baranowski M, Zhou M, et al. Roles of

intramolecular and intermolecular interactions in functional regulation of the Hsp70 J-

protein co-chaperone Sis1. J Mol Biol. 2015 Apr 10;427(7):1632–43.

22. Higurashi T, Hines JK, Sahi C, Aron R, Craig EA. Specificity of the J-protein Sis1 in the

556   propagation of 3 yeast prions. Proc Natl Acad Sci U S A. 2008 Oct 28;105(43):16596–601.

557   23. Kryndushkin D, Wickner RB. Nucleotide exchange factors for Hsp70s are required for

558   [URE3] prion propagation in Saccharomyces cerevisiae. Mol Biol Cell. 2007

559   Jun;18(6):2149–54.

560   24. Winkler J, Tyedmers J, Bukau B, Mogk A. Chaperone networks in protein disaggregation

561   and prion propagation. J Struct Biol. 2012 Aug 1;179(2):152–60.

562   25. Tuite MF, Koloteva-Levin N. Propagating prions in fungi and mammals. Mol Cell. 2004 Jun

563   4;14(5):541–52.

564   26. Berger SE, Nolte AM, Kamiya E, Hines JK. Three J-proteins impact Hsp104-mediated

565   variant-specific prion elimination: a new critical role for a low-complexity domain. Curr

566   Genet. 2019 Jun 22;

567   27. Summers DW, Douglas PM, Cyr DM. Prion propagation by Hsp40 molecular chaperones.

568   Prion. 2009;3(2):59–64.

569   28. Xu L, Gong W, Zhang H, Perrett S, Jones GW. The same but different: the role of Hsp70 in

570   heat shock response and prion propagation. Prion. 2018;12(3–4):170–4.

571   29. Killian AN, Miller SC, Hines JK. Impact of Amyloid Polymorphism on Prion-Chaperone

572   Interactions in Yeast. Viruses. 2019 Apr;11(4):349.

573   30. Masison DC, Kirkland PA, Sharma D. Influence of Hsp70s and their regulators on yeast

574   prion propagation. Prion. 2009 Jun;3(2):65–73.

31. Guinan E, Jones GW. Influence of Hsp70 chaperone machinery on yeast prion propagation.

Protein Pept Lett. 2009;16(6):583–6.

32. Sweeny EA, Shorter J. Prion proteostasis: Hsp104 meets its supporting cast. Prion. 2008

Dec;2(4):135–40.

33. Needham PG, Masison DC. Prion-impairing mutations in Hsp70 chaperone Ssa1: effects on

ATPase and chaperone activities. Arch Biochem Biophys. 2008 Oct 15;478(2):167–74.

34. Jung G, Jones G, Wegrzyn RD, Masison DC. A role for cytosolic hsp70 in yeast [PSI(+)]

prion propagation and [PSI(+)] as a cellular stress. Genetics. 2000 Oct;156(2):559–70.

35. Jones GW, Masison DC. Saccharomyces cerevisiae Hsp70 mutations affect [PSI+] prion

propagation and cell growth differently and implicate Hsp40 and tetratricopeptide repeat

cochaperones in impairment of [PSI+]. Genetics. 2003 Feb;163(2):495–506.

36. Loovers HM, Guinan E, Jones GW. Importance of the Hsp70 ATPase domain in yeast prion

propagation. Genetics. 2007 Feb;175(2):621–30.

37. Stein KC, Bengoechea R, Harms MB, Weihl CC, True HL. Myopathy-causing mutations in

an HSP40 chaperone disrupt processing of specific client conformers. J Biol Chem. 2014

Jul 25;289(30):21120–30.

38. Alberti S, Halfmann R, Lindquist S. Chapter 30 - Biochemical, Cell Biological, and Genetic

Assays to Analyze Amyloid and Prion Aggregation in Yeast. In: Methods in Enzymology

[Internet]. Academic Press; 2010 [cited 2019 Dec 16]. p. 709–34. (Guide to Yeast Genetics:

Functional Genomics, Proteomics, and Other Systems Analysis; vol. 470). Available from:

http://www.sciencedirect.com/science/article/pii/S0076687910700306

39. Nathan DF, Vos MH, Lindquist S. In vivo functions of the Saccharomyces cerevisiae Hsp90

chaperone. Proc Natl Acad Sci. 1997 Nov 25;94(24):12949–56.

40. Chernoff YO. Stress and Prions: Lessons from the Yeast Model. FEBS Lett. 2007 Jul

31;581(19):3695–701.

41. Inge-Vechtomov SG. Yeast prions as a model of neurodegenerative infectious amyloidoses

in humans. Russ J Dev Biol. 2011 Oct 6;42(5):293.

42. Dunham MJ, Fowler DM. Contemporary, yeast-based approaches to understanding human

genetic variation. Curr Opin Genet Dev [Internet]. 2013 Dec [cited 2020 Feb 8];23(6).

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3864666/

43. Chernova TA, Chernoff YO, Wilkinson KD. Yeast Models for Amyloids and Prions:

Environmental Modulation and Drug Discovery. Mol Basel Switz. 2019 Sep 18;24(18).

44. Lindström M, Liu B. Yeast as a Model to Unravel Mechanisms Behind FUS Toxicity in

Amyotrophic Lateral Sclerosis. Front Mol Neurosci. 2018;11:218.

45. Monahan ZT, Rhoads SN, Yee DS, Shewmaker FP. Yeast Models of Prion-Like Proteins

That Cause Amyotrophic Lateral Sclerosis Reveal Pathogenic Mechanisms. Front Mol

Neurosci. 2018;11:453.

46. Tenreiro S, Franssens V, Winderickx J, Outeiro TF. Yeast models of Parkinson's disease-

associated molecular pathologies. Curr Opin Genet Dev. 2017 Jun;44:74–83.

47. Sporn ZA, Hines JK. Hsp40 function in yeast prion propagation: Amyloid diversity

necessitates chaperone functional complexity. Prion. 2015;9(2):80–9.

48. Bardill JP, Dulle JE, Fisher JR, True HL. Requirements of Hsp104p activity and Sis1p

binding for propagation of the [RNQ(+)] prion. Prion. 2009 Sep;3(3):151–60.

49. Tanaka M, Chien P, Naber N, Cooke R, Weissman JS. Conformational variations in an

infectious protein determine prion strain differences. Nature. 2004 Mar 18;428(6980):323–

8.

50. Derkatch IL, Chernoff YO, Kushnirov VV, Inge-Vechtomov SG, Liebman SW. Genesis and

variability of [PSI] prion factors in Saccharomyces cerevisiae. Genetics. 1996

Dec;144(4):1375–86.

51. Tanaka M, Collins SR, Toyama BH, Weissman JS. The physical basis of how prion

conformations determine strain phenotypes. Nature. 2006 Aug;442(7102):585–9.

52. Sharma J, Liebman SW. [PSI(+)] prion variant establishment in yeast. Mol Microbiol. 2012

Nov;86(4):866–81.

53. Toyama BH, Kelly MJS, Gross JD, Weissman JS. The structural basis of yeast prion strain

variants. Nature. 2007 Sep 13;449(7159):233–7.

54. Luke MM, Sutton A, Arndt KT. Characterization of SIS1, a Saccharomyces cerevisiae

homologue of bacterial dnaJ proteins. J Cell Biol. 1991 Aug;114(4):623–38.

55. Matveenko AG, Barbitoff YA, Jay-Garcia LM, Chernoff YO, Zhouravleva GA. Differential

effects of chaperones on yeast prions: CURrent view. Curr Genet. 2018 Apr;64(2):317–25.

56. Dulle JE, Bouttenot RE, Underwood LA, True HL. Soluble oligomers are sufficient for

transmission of a yeast prion but do not confer phenotype. J Cell Biol. 2013 Oct

28;203(2):197–204.

57. Crist CG, Kurahashi H, Nakayashiki T, Nakamura Y. Conformation preserved in a weak-to-

strong or strong-to-weak [PSI+] conversion during transmission to Sup35 prion variants.

Biochimie. 2006 May 1;88(5):485–96.

58. Liebman SW, Derkatch IL. The Yeast [PSI +] Prion: Making Sense of Nonsense. J Biol

Chem. 1999 Jan 15;274(3):1181–4.

59. Alberti S, Halfmann R, Lindquist S. Chapter 30 - Biochemical, Cell Biological, and Genetic

Assays to Analyze Amyloid and Prion Aggregation in Yeast. In: Methods in Enzymology

[Internet]. Academic Press; 2010 [cited 2019 Dec 16]. p. 709–34. (Guide to Yeast Genetics:

Functional Genomics, Proteomics, and Other Systems Analysis; vol. 470). Available from:

http://www.sciencedirect.com/science/article/pii/S0076687910700306

60. Lee S, Fan CY, Younger JM, Ren H, Cyr DM. Identification of essential residues in the type

II Hsp40 Sis1 that function in polypeptide binding. J Biol Chem. 2002 Jun

14;277(24):21675–82.

61. Johnson JL, Craig EA. An Essential Role for the Substrate-Binding Region of Hsp40s in

Saccharomyces cerevisiae. J Cell Biol. 2001 Feb 19;152(4):851–6.

62. Westergard L, True HL. Extracellular environment modulates the formation and propagation

of particular amyloid structures. Mol Microbiol. 2014 May;92(4):698–715.

63. Huang VJ, Stein KC, True HL. Spontaneous variants of the [RNQ+] prion in yeast

demonstrate the extensive conformational diversity possible with prion proteins. PloS One.

2013;8(10):e79582.

64. Derkatch IL, Bradley ME, Masse SV, Zadorsky SP, Polozkov GV, Inge-Vechtomov SG, et

al. Dependence and independence of [PSI(+)] and [PIN(+)]: a two-prion system in yeast?

EMBO J. 2000 May 2;19(9):1942–52.

65. Bradley ME, Edskes HK, Hong JY, Wickner RB, Liebman SW. Interactions among prions

and prion "strains" in yeast. Proc Natl Acad Sci. 2002 Dec 10;99(suppl 4):16392–9.

66. Kryndushkin DS, Alexandrov IM, Ter-Avanesyan MD, Kushnirov VV. Yeast [PSI+] Prion

Aggregates Are Formed by Small Sup35 Polymers Fragmented by Hsp104. J Biol Chem.

2003 Dec 5;278(49):49636–43.

67. Tipton KA, Verges KJ, Weissman JS. In vivo monitoring of the prion replication cycle

reveals a critical role for Sis1 in delivering substrates to Hsp104. Mol Cell. 2008 Nov

21;32(4):584–91.

68. Abrams JL, Morano KA. Coupled Assays for Monitoring Protein Refolding in

Saccharomyces cerevisiae. JoVE J Vis Exp. 2013 Jul 9;(77):e50432.

69. Ferreira PC, Ness F, Edwards SR, Cox BS, Tuite MF. The elimination of the yeast [PSI+]

prion by guanidine hydrochloride is the result of Hsp104 inactivation. Mol Microbiol. 2001

Jun;40(6):1357–69.

70. Hennessy F, Nicoll WS, Zimmermann R, Cheetham ME, Blatch GL. Not all J domains are

created equal: Implications for the specificity of Hsp40–Hsp70 interactions. Protein Sci

Publ Protein Soc. 2005 Jul;14(7):1697–709.

71. Ramos CHI, Oliveira CLP, Fan C-Y, Torriani IL, Cyr DM. Conserved central domains

control the quaternary structure of type I and type II Hsp40 molecular chaperones. J Mol

Biol. 2008 Oct 31;383(1):155–66.

72. Pinheiro GMS, Amorim GC, Iqbal A, Almeida FCL, Ramos CHI. Solution NMR

investigation on the structure and function of the isolated J-domain from Sis1: Evidence of

transient inter-domain interactions in the full-length protein. Arch Biochem Biophys. 2019

Jul 15;669:71–9.

73. Kaschner LA, Sharma R, Shrestha OK, Meyer AE, Craig EA. A conserved domain important

for association of eukaryotic J-protein co-chaperones Jjj1 and Zuo1 with the ribosome.

Biochim Biophys Acta. 2015 May;1853(5):1035–45.

74. Ajit Tamadaddi C, Sahi C. J domain independent functions of J proteins. Cell Stress

Chaperones. 2016 Jul 1;21(4):563–70.

75. Schilke BA, Ciesielski SJ, Ziegelhoffer T, Kamiya E, Tonelli M, Lee W, et al. Broadening

the functionality of a J-protein/Hsp70 molecular chaperone system. PLOS Genet. 2017 Oct

30;13(10):e1007084.

76. Bukau B, Horwich AL. The Hsp70 and Hsp60 Chaperone Machines. Cell. 1998 Feb

6;92(3):351–66.

77. Tom A. Hall. BioEdit: a user-friendly biological sequence alignment editor and analysis

program for Windows 95/98/NT. Nucleic acids symposium series. 1999;41(41):1979–2000.

78. Hall T, Biosciences I, Carlsbad C. BioEdit: an important software for molecular biology.

GERF Bull Biosci. 2011;2(1):60–61.

79. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, et al. Multiple sequence

alignment with the Clustal series of programs. Nucleic Acids Res. 2003 Jul 1;31(13):3497–

500.

80. Roy A, Kucukural A, Zhang Y. I-TASSER: a unified platform for automated protein

structure and function prediction. Nat Protoc. 2010 Apr;5(4):725–38.

81. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y. The I-TASSER Suite: protein structure

and function prediction. Nat Methods. 2015 Jan;12(1):7–8.

82. Zhang Y. I-TASSER server for protein 3D structure prediction. BMC Bioinformatics. 2008

Jan 23;9(1):40.

83. The PyMOL Molecular Graphics System. Schrödinger, LLC;

84. Bradley ME, Liebman SW. Destabilizing interactions among [PSI(+)] and [PIN(+)] yeast

prion variants. Genetics. 2003 Dec;165(4):1675–85.

85. A system of shuttle vectors and yeast host strains designed for efficient manipulation of

DNA in Saccharomyces cerevisiae. | Genetics [Internet]. [cited 2019 Dec 4]. Available

from: https://www.genetics.org/content/122/1/19.short

86. Guthrie C, Fink G. Guide to Yeast Genetics and Molecular Biology. San Diego, CA:

Academic Press, Inc; 1991.

87. Dulle JE, Stein KC, True HL. Regulation of the Hsp104 middle domain activity is critical for

yeast prion propagation. PloS One. 2014;9(1):e87521.

FIGURES

Fig 1.

Fig 2.

726
727 **Fig 3.**

728
729   **Fig 4.**

A

B

C

D

731
732 **Fig 5.**

734 **Supporting information**

Oligonucleotide	Description	Sequence
1890	3'-Sis1-S49V	5'-aaatttcaaaggcctctactatctccttaaacttttctgtgtcacctgttgg
1891	5'-Sis1-S49V	5'-ccaacaggtgacacagaaaagtttaaggagatagtagaggcctttgaaattt
1892	3'-Sis1-E53A	5'-ttaaggagatatcagaggcctttgcaattttaaatgatcctcaaaaaagg
1893	5'-Sis1-E53A	5'-cctttttttgaggatcatttaaaattgcaaaggcctctgatatctccttaa
1894	3'-Sis1-N56L	5'-gaaaagtttaaggagatatcagaggcctttgaaattttactagatcctcaaaaaagggaaatat
1895	5'-Sis1-N56L	5'-atatttcccttttttgaggatctagtaaaatttcaaaggcctctgatatctccttaaacttttc
1964	5'-Sis1-ClaI	5'-ccccatcgatatggtcaaggagaca
1965	3'-Sis1-SpeI	5'-gcactagtttaaaaattttcatctatagcacgttt

735
736 **S1 Table. Oligonucleotides used in this study.**

A

[*psi*-]

WT

S49V

S49V +WT

E53A

E53A +WT

N56L

N56L +WT

738
739 **S1 Fig. Novel LGMDD1 J-domain mutations in Sis1 differentially impair propagation of**

740 **[PSI⁺] strains and are rescued by expression of an additional WT-Sis1 copy.** (A) *sis1Δ*

741 [*PSI⁺*] strains expressing a single copy of wild-type or mutated *SIS1,* and strains expressing an

742 additional copy of wild-type *SIS1* (+WT) were spotted onto YPD media (n=3). All spottings,

743 minus the top (labeled [*psi*-]) are weak [*PSI⁺*] strains expressing the indicated constructs.

744
S2 Fig. Sis1-S49V exhibits impaired substrate refolding while G/F domain mutants do not.

(A) the capability for refolding luciferase was measured in [*rnq*-] *sis1Δ* yeast strains harboring

the indicated construct along with a plasmid expressing luciferase. (B) same as in *A*, but in Sc37

[*PSI*+] strains expressing wild-type or the mutated chimeric construct *SDSS* constructs instead of

Sis1. Yeast were normalized, treated with cycloheximide and subjected to heat shock at 42°C for

22 minutes, followed by recovery at 30°C. Luminescence was measured at the indicated

timepoints during recovery and normalized to luminescence of samples without heat shock

treatment. The amount of luciferase refolding is plotted as percentage of recovery and

represented as mean SEM (n=6).

755
756 **S3 Fig. Sis1-S49V, does not show decreased steady-state levels when expressed in Sc37**

757 **[PSI$^+$] strain. (A)** Representative western blot showing expression of Sis1 in Sc37 [*PSI$^+$*] *sis1Δ*

758 yeast strains harboring the indicated construct (n=3). Pgk1 is shown as a loading control. All

759 samples were analyzed under the same experimental conditions.