Client processing is altered by novel myopathy-causing mutations in the HSP40 J domain

Melanie Y. Pullen[§], Conrad C. Weihl[ʆ] and Heather L. True[§1]

From the [§]Departments of Cell Biology and Physiology, and [ʆ]Neurology, Washington University School of Medicine, St Louis, Missouri 63110

Running title: *Novel LGMDD1 mutations in the HSP40 J domain*

[1]To whom correspondence should be addressed: Heather L. True: Tel.: (314) 362-3934, Fax: (314) 362-7463, E-mail: heather.true@wustl.edu; Conrad C. Weihl: Tel.: (314) 747-6394, Fax: (314) 362-4391, E-mail: weihlc@neuro.wustl.edu

Keywords: Genetic disease; Heat shock protein (HSP); Molecular chaperone; Muscular dystrophy; Myopathy; Prion; Protein aggregation; Yeast

---

ABSTRACT

The misfolding and aggregation of proteins is often implicated in the development and progression of degenerative diseases. Heat shock proteins (HSPs), such as the ubiquitously expressed Type II Hsp40 molecular chaperone, DNAJB6, assist in protein folding and disaggregation. Historically, mutations within the DNAJB6 G/F domain have been associated with Limb-Girdle Muscular Dystrophy type 1D, now referred to as LGMDD1, a dominantly inherited degenerative disease. Recently, novel mutations within the J domain of DNAJB6 have been reported in patients with LGMDD1. Since novel myopathy-causing mutations in the Hsp40 J domain have yet to be characterized and both the function of DNAJB6 in skeletal muscle and the clients of this chaperone are unknown, we set out to assess the effect of these mutations on chaperone function using the genetically tractable yeast system. The essential yeast Type II Hsp40, Sis1, is homologous to DNAJB6 and is involved in the propagation of yeast prions. Using phenotypic, biochemical, and functional assays we found that homologous mutations in the Sis1 J domain differentially alter the processing of specific yeast prion strains, as well as a non-prion substrate. These data suggest that the newly-identified mutations in the J domain of DNAJB6 cause aberrant chaperone function that leads to the pathogenesis in LGMDD1.

INTRODUCTION

Molecular chaperones preserve protein homeostasis (1). A deficient chaperone network may lead to protein misfolding and aggregation often associated with protein conformational disorders such as Alzheimer's Disease, Charcot-Marie-Tooth disease, distal hereditary motor neuropathies, and Limb Girdle Muscular Dystrophy, among others (2, 3). Limb Girdle Muscular Dystrophy type 1D (LGMD1D), more recently termed LGMDD1 (4), is a disease characterized by proximal muscle weakness with moderate progression mediated by defective chaperone function (5). Historically, dominantly inherited disease-associated mutations in the type II Hsp40 co-chaperone DNAJB6 have been found within a 12 amino acid region known as the G/F domain (6–11). Recently, three novel pathogenic mutations associated with LGMDD1 have been identified within the J domain of DNAJB6 (12, 13).

Since the heat shock response is highly conserved from yeast to mammals, we used a yeast model system to study these disease-associated mutations (14–16). The essential yeast Type II Hsp40, Sis1, is homologous to DNAJB6 and plays an important role in yeast for the propagation of two prions, [*RNQ*[+]] and [*PSI*[+]] (17–22). Prions in yeast are epigenetic elements that form when the prion proteins form amyloid aggregates. These prion protein aggregates are propagated by fragmentation into propagons, which are then passed on to progeny during mitosis (14). Somewhat counterintuitively, chaperones not only promote prion propagation in yeast, but are essential for it (14, 23–26). At the heart of this chaperone-mediated process is the Hsp40 Sis1, with the interaction between Hsp40

and Hsp70 chaperones being crucial for prion propagation in yeast (24, 27–32). When chaperone function is defective, prion propagation is impaired (17, 33–36). Thus, the loss of prion propagation provides a read-out for chaperone dysfunction in yeast.

Homologous LGMDD1-associated mutations of the DNAJB6 G/F domain impair the propagation of various yeast prion strains when expressed in Sis1 (37). We were inspired by this model to assess the effect of homologous J domain mutants in Sis1 on chaperone function. Fortunately, the yeast model system allows for a wide array of phenotypic, biochemical, and functional assays to assess the effect of these mutations on chaperone function (14, 38). As such, we assessed the ability of the disease-associated mutant chaperone to process two specific clients of Sis1, Sup35 and Rnq1, which form the prion elements [*PSI*+] and [*RNQ*+], respectively. In addition, we performed functional assays with a non-prion client of the yeast chaperone machinery, firefly luciferase (FFL) (39). In conjunction, these assays elucidated nuances in client processing due to disease-associated mutations in a manner that would not be possible by assessing a single client protein or using a limited number of assays.

Here, we present evidence that disease-associated mutations in the J domain cause functional defects that differentially impair processing of native proteins in a client and conformer specific manner. Our work suggests that mutations in the J domain are responsible for altered protein processing and, potentially, disease pathogenesis.

RESULTS

Homologous mutations in the Sis1 J domain are conserved and potentially alter J domain and G/F domain interaction. Recently, novel mutations located within a 7 amino acid region of the DNAJB6 J domain have been reported to cause dominant distal myopathy (12, 13). These variants were identified in five different families presenting with both distal and proximal muscle weakness. Initial studies examined these variants in patient tissue and cell culture studies in an attempt to elucidate whether these mutations lead to functional defects in DNAJB6's anti-aggregation activity. From these studies, the recently identified

DNAJB6 variants DNAJB6-A50V, DNAJB6-E54A and DNAJB6-S57L have been categorized as pathogenic, likely pathogenic, and variant of unknown significance, respectively (12).

To further examine the effect of these disease-associated variants on myopathy, we performed protein threading to identify homologous mutations in the yeast Hsp40 Sis1. Our comparative modeling identified homologous J domain mutations in Sis1 to be Sis1-S49V (homologous to DNAJB6-A50V), Sis1-E53A (homologous to DNAJB6-E54A), and Sis1-N56L (homologous to DNAJB6-S57L) (Figure 1A). In addition, our structural model recapitulates the observation that mutations in the J domain have the potential to disrupt important intramolecular interactions previously shown to stabilize the interface between the G/F and J domains (Figure 1B) (12, 21).

Homologous mutations in the J domain of Sis1 differentially impair [PSI+] prion propagation. To assess the implications of these disease-causing variants on chaperone function, we turned to prion propagation models in yeast. Such models have been used extensively to study chaperones and understand the deleterious effects of disease-associated mutations on protein folding (40–46). In yeast, prions are naturally occurring, self-propagating protein structures that require chaperone networks for their efficient propagation (14). Defects in chaperone function disrupt this process and, therefore, prion propagation (27, 33–36, 47, 48). In order to ask whether these disease variants disrupt client processing, we studied two known Sis1 substrates: the translation termination factor Sup35, which forms the [*PSI*+] prion, and Rnq1, which forms the [*RNQ*+] prion (14, 29).

In an effort to understand the effect of homologous J domain mutations on [*PSI*+] prion propagation we performed experiments utilizing strains harboring the *ade1-14* allele that has a premature stop codon in the *ADE1* open reading frame (14). When Sup35 misfolds and aggregates in the [*PSI*+] state, nonsense suppression of the premature stop codon in *ade1-14* occurs and the adenine biosynthesis pathway is completed. Thus, [*PSI*+] colonies are white in color and can grow on media lacking adenine. Conversely, if cells are [*psi-*], adenine biosynthesis is incomplete, resulting in cells that are unable to grow in media lacking adenine and an accumulation of red

pigment in cells grown on rich media. This colorimetric phenotypic assay has been utilized to demonstrate the effect of LGMDD1-causing G/F domain mutations on [*PSI*+] propagation (37). Here, we make use of four different [*PSI*+] strains (weak, strong, Sc37, and Sc4) (49, 50). Each prion strain is a different self-propagating conformation of the same protein sequence and they are differentiated by the strength of the nonsense suppression phenotype and stability of their mitotic inheritance (49, 51–53). Yeast containing *ade1-14* that propagate stronger [*PSI*+] strains (strong and Sc4) display higher nonsense suppression and are lighter in color. Yeast propagating weaker [*PSI*+] strains (weak and Sc37) are darker in color due to decreased nonsense suppression (better translation termination). Sis1 is required for yeast viability (54). Therefore, to perform these experiments, we used *sis1Δ* yeast (covered by a plasmid expressing Sis1) propagating different [*PSI*+] strains (weak, strong, Sc37, and Sc4) and replaced *SIS1* with homologous LGMDD1 mutant *SIS1* constructs or wild-type *SIS1* (as a control) (Figure 2A). In doing so, we observed viability was not affected in these strains expressing the mutants.

Interestingly, we found that expression of Sis1-S49V in the weak [*PSI*+] strain, but not the Sc37 [*PSI*+] strain, altered the weak strain phenotype, as indicated by the lighter color on rich media and robust growth in selective media as compared to wild-type (Figure 2A). In comparison to the other J domain mutants, the stronger nonsense suppression phenotype we observed is unique to expression of Sis1-S49V in the weak [*PSI*+] strain. This suggests altered processing of Sup35 by Sis1-S49V occurs with conformational specificity. We questioned whether expression of Sis1-S49V could have altered the conformation of Sup35 in weak [*PSI*+] in a manner that leads to the propagation of a strong [*PSI*+] strain. Intrigued by this result, we introduced pRS316-Sis1 (wildtype) to this strain and re-assessed its [*PSI*+] phenotype (Supporting Figure S1). We found this reverted the phenotype and was comparable to wild-type controls, suggesting that Sis1-S49V expression does not alter the [*PSI*+] strain. Previous work demonstrated that a shortage of cytosolic Sis1 may lead to more efficient transmission of propagons to daughter cells thus "strengthening" the prion phenotype (55). Thus, we hypothesize this may

occur when Sis1-S49V is expressed in a weak [*PSI*+] strain.

This assay also showed that the expression of either Sis1-E53A or Sis1-N56L altered the propagation of the weaker [*PSI*+] strains (weak and Sc37), albeit in a different manner and to varying degrees. This is indicated by the phenotypic change towards a red color (less nonsense suppression) on rich media and lack of growth on media lacking adenine (Figure 2A). Similarly, when we introduced a wild type copy of *SIS1* to these phenotypically red strains, we identified a partial rescue of the weak [*PSI*+] phenotype (Supporting Figure S1) indicating these Sis1 mutants, although phenotypically [*psi-*], result in propagation of cryptic [*PSI*+] propagons (56).

Complementarily, we used this assay quantitatively by phenotypically scoring large numbers of colonies on rich media spread plates (Figure 2B). This assay allows for a more detailed understanding of the variability in [*PSI*+] propagation. We scored colony color as light pink (strong [*PSI*+]), dark pink (weak [*PSI*+]) or red (very weak [*PSI*+] or [*psi-*]) in yeast expressing wild type Sis1 or the homologous LGMDD1 mutants. Indeed, through this assay we found that the expression of these mutants phenotypically altered prion propagation and identified a phenotypic variation that was not previously appreciated in the spotting assay. We observed a wider range of phenotypic variation in the weaker strains than in the stronger [*PSI*+] strains, which rarely convert to a weaker phenotype (57, 58). However, an increase in darker pink colonies was observed when Sis1-S49V and Sis1-N56L were expressed in the stronger [*PSI*+] strains. Similar results were observed when Sis1-E53A and Sis1-N56L were expressed in the Sc4 [*PSI*+] strain.

To further explore these results with respect to the solubility of the prion protein, we performed boiled gel assays to examine the relative levels of aggregated and monomeric Sup35. This assay consists of loading unboiled samples onto an SDS-PAGE gel. By doing so, non-denatured aggregates are too large to enter the resolving gel while monomeric Sup35 readily enters the gel due to its size and lack of aggregation. Halfway through electrophoresis, the gel itself is boiled and then run again, which allows previously aggregated protein to enter the resolving gel. The boiled gel assay provides a clear separation of monomeric and

aggregated species of Sup35 and is easily detected through subsequent western blots (59). In order to examine the distribution of Sup35 between the aggregated and monomeric forms, we grew cultures from colonies representative of the phenotypic distribution observed in our spread plate assay.

Expressing homologous J domain mutations in the strong or Sc4 [*PSI*+] strains did not alter Sup35 distribution nor phenotype significantly (Figure 2C). Interestingly, the distribution of Sup35 between monomer and aggregate does not always correlate to phenotype in the weaker [*PSI*+] strains (many replicates of these assays were performed to verify reproducibility). Of note, we previously described a chaperone mutant that demonstrated differences in the ability to relate SDS-resistant or small protein aggregates to phenotype (56). Moreover, we found that expressing Sis1-S49V in the weak [*PSI*+] strain led to Sup35 being found mostly in the aggregated state (Figure 2C), coinciding with our phenotypic assay. With regards to expression of Sis1-E53A in the weak [*PSI*+] strain, we were surprised to find little Sup35 in the monomeric fraction, while the expression of Sis1-N56L results in most of the Sup35 in the monomeric fraction. Interestingly, expressing Sis1-S49V, Sis1-E53A or Sis1-N56L in the Sc37 [*PSI*+] strain showed a marked increase in monomeric Sup35, as compared to wild-type controls, while maintaining a modest level of aggregated Sup35. Thus, there are differences with respect to the interaction between the Sis1 mutants and Sup35 in the weaker [*PSI*+] strains and how propagons and large aggregates correlate to phenotype.

Homologous mutations in the J domain of Sis1 differentially impair [RNQ+] prion propagation. We then asked whether mutations in the Sis1 J domain would also affect another known client of Sis1, the [*RNQ*+] prion. In a manner similar to that of the previous experiments, we utilized various [*RNQ*+] prion strains (*rnq-*, low, and very high) and expressed each of our Sis1 mutants. Here, we observed reduced growth when Sis1-S49V was expressed in [*psi-*] strains that we did not observe in the same yeast strains harboring the [*PSI*+] prion. Thus, we performed spottings (using 5-fold serial dilutions) and confirmed that expression of Sis1-S49V, but not Sis1-E53A or Sis1-N56L, resulted in a growth defect (Figure 3A). This

effect was only observed when wild-type Sis1 was removed through plasmid shuffle and occurred when Sis1-S49V was expressed in either [*RNQ*+] or [*rnq-*] strains. This indicated the defect in growth was independent of [*RNQ*+] and it was possibly impacted by the loss of [*PSI*+]. Because Sis1 is essential for yeast growth, mutations in Sis1 have been shown to affect viability (20, 54, 60, 61).

Unlike the nonsense suppression assay that discerns [*PSI*+] strains, strains of the [*RNQ*+] prion are characterized by the rate at which they induce the formation of [*PSI*+], ranging from low to very high rates (62–65). Therefore, we performed two different biochemical assays to assess [*RNQ*+] propagation. First, we performed a sedimentation assay (59) which consists of separating the soluble and insoluble protein fractions using ultracentrifugation to assay the relative solubility of Rnq1. These fractions are subjected to SDS-Page and probed with an antibody specific to Rnq1. We performed this assay using two [*RNQ*+] strains, low [*RNQ*+] and v.h. [*RNQ*+]. Using this assay, we identified a defect in [*RNQ*+] propagation when Sis1-S49V is expressed in both low [*RNQ*+] and v.h. [*RNQ*+] strains (Figure 3B). In comparison, there is only a slight change in Rnq1 solubility when Sis1-E53A or Sis1-N56L are expressed, indicating that J domain variants differentially impair processing of client conformers, albeit not completely.

Furthermore, in an effort to elucidate whether Rnq1 aggregate size or SDS resistance was altered by expression of the mutants, we performed a semi-denaturing detergent agarose gel electrophoresis (SDD-AGE) (Figure 3C). This assay allows for the visualization of SDS-resistant aggregates such as aggregated prion conformers (66). Strikingly, we observed a dramatic loss of SDS-resistant Rnq1 aggregates in the low and v.h. [*RNQ*+] strains when Sis1-S49V was expressed. We also observed a slight reduction in SDS-resistant Rnq1 aggregates in low [*RNQ*+] strains expressing Sis1-E53A, while low [*RNQ*+] strains expressing Sis1-N56L appeared unaffected. In addition, v.h. [*RNQ*+] strains expressing Sis1-E53A or Sis1-N56L showed a decreased in SDS-resistant Rnq1 aggregates. Altogether, these biochemical assays present varying degrees of altered [*RNQ*+] propagation that occur in a conformer-specific manner when disease-

associated J domain mutants are expressed in these strains.

Disruption in prion propagation is due to a defect in substrate refolding. We observed different degrees of alteration in prion propagation due to the expression of the homologous LGMDD1 mutations in the J domain, which vary depending on the client and its conformation. To further understand the effect of these disease-associated mutations, we assessed how they impact protein folding of a non-prion substrate *in vivo*. Thus, we utilized the well-established firefly luciferase (FFL) refolding assay (67, 68) (Figure 4). FFL is a non-prion substrate of the Hsp40/Hsp70/Hsp104 chaperone machinery and is denatured in cells by heat shock. When FFL misfolds and aggregates it requires the Hsp40/Hsp70/Hsp104 chaperone machinery to efficiently refold. Since Hsp104 is required for efficient refolding of FFL, we used a *hsp104Δ* strain expressing FFL as a negative control. We observed a significant defect in FFL refolding when Sis1-S49V was expressed in either a [*PSI+*] [*RNQ+*] (Figure 4) or [*psi-*] [*rnq-*] strain (Supporting Figure S2A), but not when other J domain variants are expressed. We subjected other known G/F domain mutants expressed in a chimeric protein where the G/F domain of Sis1 is replaced by that of DNAJB6, referred to as SDSS (37), to this assay and saw no significant differences in luciferase refolding as compared to wild type (Supporting Figure S2B). Hence, these results highlight the importance of substrate recognition – both with the client itself, as well as conformer specificity. Most importantly, these findings emphasize the distinction of assessing native chaperone clients as these mutants displayed varying degrees of altered prion propagation in our assays. Given this specificity, care should be taken when using certain assays, or a limited number of them, when trying to understand and confirm whether a potentially pathogenic chaperone variant is disease-causing or not.

Stability of a mutated Hsp40 chaperone is rescued by expression of the [PSI+] prion. While assessing the effect of the J domain LGMDD1 mutations on [*RNQ+*] propagation we identified a growth defect that only occurred with expression of the Sis1-S49V variant (Figure 3A). Interestingly, we observed this defect when Sis1-

S49V was expressed in [*RNQ+*] [*psi-*] strains, but not when expressed in [*PSI+*] strains (Figure 2A and Figure 3A). Because *SIS1* is essential (20, 54, 60, 61), we were curious whether this growth defect was due to any change in steady-state expression levels. We assessed steady state Sis1 levels by western blot and found a 40-60% decrease in the level of Sis1-S49V in various [*RNQ+*] and [*rnq-*] strains (Figure 5A). We hypothesize that this decrease in steady-state levels is responsible for the growth defect.

To confirm this decrease is due to a lack of [*PSI+*] presence, we set out to recapitulate this phenomenon by curing [*PSI+*] strains of its prion form. Curing of [*PSI+*] was performed by growth on media containing guanidine hydrochloride (GdHCl), which inactivates the ATPase activity of Hsp104 and hinders the replication of [*PSI+*] seeds (69). After curing both a strong [*PSI+*] and a Sc37 [*PSI+*] strain (each expressing Sis1-S49V) we identified a significant decrease in Sis1 steady-state levels, and this was not observed with expression of wild type Sis1 nor the other LGMDD1 J domain variants (Figure 5B). The decrease in steady-state levels recapitulates what we observed in [*psi-*] strains expressing Sis1-S49V (Figure 5A). This striking result suggests the Sis1 client, Sup35, in its aggregated and [*PSI+*]-propagating form, interacts with this particular Hsp40 mutant in a way that stabilizes the steady-state level of Sis1-S49V. Of note, this decrease in Sis1 steady-state levels has not been observed in the context of LGMDD1-associated mutations found in the G/F domain. This interesting finding is the first known instance wherein a potentially unstable mutant chaperone is stabilized by the presence of a specific client protein in its aggregated form.

DISCUSSION

LGMDD1 is a myopathy historically characterized by mutations within the G/F domain of the chaperone DNAJB6 (6–9, 11). Here we used the yeast model system, a homologous chaperone, and native chaperone clients to demonstrate that recently reported mutations in the J domain of DNAJB6 alter canonical chaperone function. Specifically, expression of homologous disease-associated mutations in the yeast Hsp40, Sis1, identified defects in prion propagation to varying degrees depending on the disease-

associated variant, the client protein and its conformation, as well as a significant defect in refolding of a non-prion client protein. We conclude these mutations in the J domain of Sis1 lead to aberrant chaperone function, altered protein homeostasis, and potentially drive a variety of defects in chaperone machinery which may contribute to pathogenesis.

The function and interaction of the various Hsp40 domains has been a topic of interest for many years (20, 21, 70–75). Surprisingly, colleagues identified LGMDD1 patients with novel variants in the J domain of DNAJB6, a disease previously characterized by mutations in the G/F domain (12, 13). As previously mentioned, LGMDD1 disease-associated mutations in the G/F domain have been found to impair the processing of specific client conformers. These findings highlighted the G/F domain as having an important role in substrate regulation and conformer selectivity (37). By contrast, the J domain of Sis1 has a known role in regulating Hsp70 ATPase activity (76).

Here we demonstrate a novel finding by which disease-associated missense mutations in the J domain of an Hsp40 result in the chaperone having client and conformer specificity. Strikingly, this fits with previous structural studies which have highlighted the importance of the intramolecular interaction between the J and G/F domains (21). Specifically, a particular amino acid, E50, in the Sis1 J domain has been shown to interact with the EEVD(HSP70) motif, an interaction required for both chaperones to partner in protein refolding (21). Moreover, this key amino acid (E50) forms a salt bridge with R73 which is found in the Sis1 G/F domain. We and others (12, 21) believe this interaction is crucial and of functional importance for the stability and interaction between both domains and, ultimately, the interaction between Hsp40 and Hsp70.

In the context of the LGMDD1-associated mutants found in the J domain, all variants appear to affect prion propagation to varying degrees depending on client and conformation specificity. Notably, the homologous Sis1-S49V variant appears to have the most deleterious effect on the chaperone machinery and its function, probably due to its proximity to the E50:R73 salt bridge. This would also explain why only Sis1-S49V presented with defective substrate refolding in our

luciferase refolding assay. Overall, we hypothesize that the steric hindrance caused by these missense mutations may obstruct the intramolecular interaction or communication between the J and G/F domains. Moreover, these findings support the idea that it may be the interaction between these two domains that is crucial for the specificity and processing of Hsp40 client proteins.

By assessing the propagation of various Sis1 clients, we found that there are differences in prion propagation, but that propagation is not completely abolished by expression of these mutants. When compared to LGMDD1-associated mutations in the G/F domain, for example, we observe variants in the G/F domain to have a stronger effect on prion propagation in certain assays, oftentimes demonstrating complete loss of prion propagation (37). As such, although these disease-associated mutations in the J domain disrupt prion processing and protein refolding, they do so in a manner that is slightly different from that of G/F domain mutants.

Given the limited knowledge of the clients of DNAJB6 in skeletal muscle and its role in disease, we hypothesize, as we did for the G/F domain mutations, that distinct variants may be associated with different levels of disease severity due to impairment of Hsp40 function to varying degrees, or in different ways. Nonetheless, we are cautious to imply that a more drastic loss-of-propagation effect would correlate with or be causative of a more severe disease pathogenesis, as we have observed different degrees of aberrant chaperone function in the context of these disease-associated variants that are greatly dependent on client and conformation specificity. It is not uncommon for LGMDD1-associated mutations to have an effect in some but not all assays or when assaying specific client proteins (12, 37). Moreover, as we continue to understand and classify disease-associated variants as pathogenic, we should consider how variable clinical outcomes and disease severity may be influenced by factors such as variable expressivity and sex influences (11). Given these considerations, there is a need for simple models, such as the yeast system, by which we can understand if and how identified genetic variants affect chaperone function in order to better understand variable clinical outcomes within LGMDD1. Currently, we can only show that there are differences in adequate client

processing and refolding although, undoubtedly, further *in vitro* studies are required to further elucidate the mechanistic cause of these functional defects and how they translate to pathogenesis.

Lastly, we identified a novel phenomenon by which a mutated Hsp40, Sis1-S49V, appears to have a decreased steady state level of protein expression. This is, to our knowledge, the first known instance where steady-state expression of a chaperone is dependent on stabilization by a known substrate. Although serendipitous, this finding not only highlights an important function of chaperone-substrate interaction, but also hints at an additional role that specific substrates might play in chaperone function and protein homeostasis overall. Indeed, there is still much to be learned about co-chaperones, their function, interactions between themselves and with clients, and their role in disease pathogenesis.

EXPERIMENTAL PROCEDURES

Protein sequence alignment structure modeling. Protein sequences for DNAJB6b orthologs were found on UniProt and subsequently aligned to homologous disease-causing J domain mutations in Sis1. Sequences were visualized using BioEdit (77) with multiple sequences aligned through the use of ClustalW2 (77–79). Mutant DNAJB6b structures were generated through the use of using I-TASSER (80–82) and visualized using PyMOL (83). Structural comparisons and protein threading of variants to wild-type protein structures was performed using wild-type Sis1 (PBD: 4RWU) (21).

Yeast cultures and transformation. All experiments were performed in derivates of 74-D694 (*ade1-14 his3-Δ200 leu2-3,112 trp1-289 ura3-52*). Yeast strains are [*PSI+*], [*psi-*], [*RNQ+*], or [*rnq-*], kindly gifted by S. Liebman (50, 84) and J. Weissman (51). Yeast were cultured using YPD (1% yeast extract, 2% peptone, 2% dextrose), 1/4 YPD (0.25% yeast extract, 2% peptone, 2% dextrose) or synthetic defined (SD) media (0.67% yeast nitrogen base without amino acids, 2% dextrose) lacking specific amino acids using standard techniques. In order to study the function of *sis1* mutants in the absence of wild type *SIS1*, previously described plasmid shuffle strains carrying pRS316-SIS1 (37) were transformed with pRS314 (85) carrying *sis1* mutants. We selected for colonies that lost wild type *SIS1* through plasmid shuffle on plates containing 5-fluoroorotic acid (5-FOA). Plasmid transformations were performed by the PEG/LiOAC method (86). Plasmid pRS316-Sis1 was a kind gift from E. Craig (17, 18). Plasmid 316-GPD-Lux was a kind gift from J. Weissman (67). Other plasmids are described below and were constructed using standard molecular techniques.

Plasmid Construction. Oligonucleotides used for site-directed mutagenesis are listed in Table 1. Using pRS314-SIS1, the J domain LGMDD1 mutations were created by site-directed mutagenesis using the Agilent QuikChange II XL Sit'-Directed Mutagenesis Kit, as per manufacturer's instructions along with the following oligonucleotides: 1890 and 1891 (S49V), 1892 and 1893 (E53A), 1894 and 1895 (N56L). Primer sequences were generated using Agilent's online primer design program. Mutagenesis was confirmed by sequencing the entire coding region of *SIS1*.

Yeast Phenotypic Assays. For yeast spottings, cultures were grown overnight in selective media. Cultures were pelleted, washed, and suspended in water to an optical density of 0.1. The normalized yeast solutions were pipetted into a 96-well plate, and serial dilutions (1:5) were made using a multichannel pipette. Yeast were spotted onto plates using an ethanol-sterilized 48-pin replicator. To monitor cell growth or the [*PSI+*] status of yeast cells with the spread plate assay, overnight cultures were normalized by A600, serially diluted 5-fold, and spotted on the indicated media. [*PSI+*] status was assessed by colony color on large 145mm plates in rich 1⁄4 YPD media. Plates were incubated for 5 days at 30°C followed by overnight incubation at 4°C for additional color development.

Protein Analysis. For boiled gel assays, yeast strains were cultured overnight. Cells were lysed with glass beads in buffer (25mM Tris-HCl pH7.5, 100mM NaCl, 1mM EDTA, protease inhibitors) and pre-cleared at 6,000 rpm for 1 minute at 4°C. Protein concentration of cell lysates was normalized using a Bradford assay and mixed with SDS-Page sample buffer (200mM Tris-HCl pH 6.8, 4% SDS, 0.4% bromophenol blue, 40% glycerol). Samples remained un-boiled and were loaded on a 12% polyacrylamide gel and run under constant current of 110V until the dye front migrated halfway through the resolving gel. The

current was then stopped, and the gel in glass plates was sealed in plastic and boiled upright for 15mins in a 95-100°C water bath. After boiling, gels were removed from the plastic cover and were reinserted in the SDS-PAGE apparatus, where voltage was re-applied until the dye migrated to the bottom of the gel. SDS-PAGE was followed by standard western blotting with Sup35, Sis1, and Pgk1 antibodies. Sedimentation analysis for Rnq1 was performed as previously described (37, 48). Semi-denaturing agarose gel electrophoresis (SDD-AGE) for [*PSI*⁺] was performed as previously described (37, 87).

Luciferase Refolding. Luciferase refolding assays were performed as previously described (68). To monitor the ability of Sis1 to enhance refolding of luciferase *in vivo*, [*PSI*⁺] or [*rnq-*] yeast strains were transformed with the plasmid pRS316-GPD-Lux. Cultures were grown overnight in selective media and back diluted to an optical density of 0.3 in 8mL plastic culture tubes. Subsequently, cyclohexamide was added to a final concentration of 10ug/mL. Treated samples were then subjected to heat shock at 42°C for 22 minutes. Meanwhile, control samples were plated in 98-well clear bottom plates and kept at 30°C. After heat shock, 200ul of each sample was plated in triplicate on 98-well clear bottom plates. All cultures were allowed to recover at 30°C. Luminometer readings were taken at 30, 60, 90 and, 120 minutes. For data analysis, each triplicate was averaged. 2way ANOVA was performed using GraphPad Prism version 8.1.1 for Windows, GraphPad Software, La Jolla California USA, www.graphpad.com.

Acknowledgements: We thank S. Liebman, J. Weissman and E. Craig for reagents. We also thank J. Daw for assistance with site-directed mutagenesis. We are appreciative of L. Dublin and A. Bhadra for helpful discussions and comments on the manuscript.

Author contributions: MYP, CCW and HLT designed the research. MYP performed the research and analyzed the data. MYP, CCW and HLT wrote or/and edited the paper.

Conflict of interest: The authors declare that they have no conflicts of interest with the contents of this article. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

1. Jee, H. (2016) Size dependent classification of heat shock proteins: a mini-review. *J. Exerc. Rehabil.* **12**, 255–259
2. Muranova, L. K., Ryzhavskaya, A. S., Sudnitsyna, M. V., Shatov, V. M., and Gusev, N. B. (2019) Small Heat Shock Proteins and Human Neurodegenerative Diseases. *Biochem. Biokhimiia.* **84**, 1256–1267
3. Hipp, M. S., Kasturi, P., and Hartl, F. U. (2019) The proteostasis network and its decline in ageing. *Nat. Rev. Mol. Cell Biol.* **20**, 421–435
4. Straub, V., Murphy, A., Udd, B., and LGMD workshop study group (2018) 229th ENMC international workshop: Limb girdle muscular dystrophies - Nomenclature and reformed classification Naarden, the Netherlands, 17-19 March 2017. *Neuromuscul. Disord. NMD.* **28**, 702–710
5. Ruggieri, A., Saredi, S., Zanotti, S., Pasanisi, M. B., Maggi, L., and Mora, M. (2016) DNAJB6 Myopathies: Focused Review on an Emerging and Expanding Group of Myopathies. *Front. Mol. Biosci.* **3**, 63
6. Couthouis, J., Raphael, A. R., Siskind, C., Findlay, A. R., Buenrostro, J. D., Greenleaf, W. J., Vogel, H., Day, J. W., Flanigan, K. M., and Gitler, A. D. (2014) Exome sequencing identifies a DNAJB6 mutation in a family with dominantly-inherited limb-girdle muscular dystrophy. *Neuromuscul. Disord. NMD.* **24**, 431–435
7. Harms, M. B., Sommerville, R. B., Allred, P., Bell, S., Ma, D., Cooper, P., Lopate, G., Pestronk, A., Weihl, C. C., and Baloh, R. H. (2012) Exome sequencing reveals DNAJB6 mutations in dominantly-inherited myopathy. *Ann. Neurol.* **71**, 407–416
8. Sarparanta, J., Jonson, P. H., Golzio, C., Sandell, S., Luque, H., Screen, M., McDonald, K., Stajich, J. M., Mahjneh, I., Vihola, A., Raheem, O., Penttilä, S., Lehtinen, S., Huovinen, S., Palmio, J., Tasca, G., Ricci, E., Hackman, P., Hauser, M., Katsanis, N., and Udd, B. (2012) Mutations affecting the cytoplasmic functions of the co-chaperone DNAJB6 cause limb-girdle muscular dystrophy. *Nat. Genet.* **44**, 450–455, S1-2
9. Sato, T., Hayashi, Y. K., Oya, Y., Kondo, T., Sugie, K., Kaneda, D., Houzen, H., Yabe, I., Sasaki, H., Noguchi, S., Nonaka, I., Osawa, M., and Nishino, I. (2013) DNAJB6 myopathy in an Asian cohort and cytoplasmic/nuclear inclusions. *Neuromuscul. Disord. NMD.* **23**, 269–276
10. Bohlega, S. A., Alfawaz, S., Abou-Al-Shaar, H., Al-Hindi, H. N., Murad, H. N., Bohlega, M. S., Meyer, B. F., and Monies, D. (2018) LGMD1D myopathy with cytoplasmic and nuclear inclusions in a Saudi family due to DNAJB6 mutation. *Acta Myol. Myopathies Cardiomyopathies Off. J. Mediterr. Soc. Myol.* **37**, 221–226
11. Zima, J., Eaton, A., Pál, E., Till, Á., Ito, Y. A., Warman-Chardon, J., Hartley, T., Cagnone, G., Melegh, B. I., Care4Rare Canada, Boycott, K. M., Melegh, B., and Hadzsiev, K. (2019) Intrafamilial variability of limb-girdle muscular dystrophy, LGMD1D type. *Eur. J. Med. Genet.* 10.1016/j.ejmg.2019.04.012
12. Palmio, J., Jonson, P. H., Inoue, M., Sarparanta, J., Bengoechea, R., Savarese, M., Vihola, A., Jokela, M., Nakagawa, M., Noguchi, S., Olivé, M., Masingue, M., Kerty, E., Hackman, P., Weihl, C. C., Nishino, I., and Udd, B. (2019) Mutations in the J domain of DNAJB6 cause dominant distal myopathy. *Neuromuscul. Disord.* 10.1016/j.nmd.2019.11.005
13. Savarese, M., Di Fruscio, G., Mutarelli, M., Torella, A., Magri, F., Santorelli, F. M., Comi, G. P., Bruno, C., and Nigro, V. (2014) MotorPlex provides accurate variant detection across large muscle genes both in single myopathic patients and in pools of DNA samples. *Acta Neuropathol. Commun.* **2**, 100
14. Liebman, S. W., and Chernoff, Y. O. (2012) Prions in yeast. *Genetics.* **191**, 1041–1072
15. Richter, K., Haslbeck, M., and Buchner, J. (2010) The heat shock response: life on the verge of death. *Mol. Cell.* **40**, 253–266

16. Bukau, B., Weissman, J., and Horwich, A. (2006) Molecular chaperones and protein quality control. *Cell.* **125**, 443–451
17. Aron, R., Higurashi, T., Sahi, C., and Craig, E. A. (2007) J-protein co-chaperone Sis1 required for generation of [RNQ+] seeds necessary for prion propagation. *EMBO J.* **26**, 3794–3803
18. Lopez, N., Aron, R., and Craig, E. A. (2003) Specificity of class II Hsp40 Sis1 in maintenance of yeast prion [RNQ+]. *Mol. Biol. Cell.* **14**, 1172–1181
19. Sondheimer, N., Lopez, N., Craig, E. A., and Lindquist, S. (2001) The role of Sis1 in the maintenance of the [RNQ+] prion. *EMBO J.* **20**, 2435–2442
20. Yan, W., and Craig, E. A. (1999) The glycine-phenylalanine-rich region determines the specificity of the yeast Hsp40 Sis1. *Mol. Cell. Biol.* **19**, 7751–7758
21. Yu, H. Y., Ziegelhoffer, T., Osipiuk, J., Ciesielski, S. J., Baranowski, M., Zhou, M., Joachimiak, A., and Craig, E. A. (2015) Roles of intramolecular and intermolecular interactions in functional regulation of the Hsp70 J-protein co-chaperone Sis1. *J. Mol. Biol.* **427**, 1632–1643
22. Higurashi, T., Hines, J. K., Sahi, C., Aron, R., and Craig, E. A. (2008) Specificity of the J-protein Sis1 in the propagation of 3 yeast prions. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 16596–16601
23. Kryndushkin, D., and Wickner, R. B. (2007) Nucleotide exchange factors for Hsp70s are required for [URE3] prion propagation in Saccharomyces cerevisiae. *Mol. Biol. Cell.* **18**, 2149–2154
24. Winkler, J., Tyedmers, J., Bukau, B., and Mogk, A. (2012) Chaperone networks in protein disaggregation and prion propagation. *J. Struct. Biol.* **179**, 152–160
25. Tuite, M. F., and Koloteva-Levin, N. (2004) Propagating prions in fungi and mammals. *Mol. Cell.* **14**, 541–552
26. Berger, S. E., Nolte, A. M., Kamiya, E., and Hines, J. K. (2019) Three J-proteins impact Hsp104-mediated variant-specific prion elimination: a new critical role for a low-complexity domain. *Curr. Genet.* 10.1007/s00294-019-01006-5
27. Summers, D. W., Douglas, P. M., and Cyr, D. M. (2009) Prion propagation by Hsp40 molecular chaperones. *Prion.* **3**, 59–64
28. Xu, L., Gong, W., Zhang, H., Perrett, S., and Jones, G. W. (2018) The same but different: the role of Hsp70 in heat shock response and prion propagation. *Prion.* **12**, 170–174
29. Killian, A. N., Miller, S. C., and Hines, J. K. (2019) Impact of Amyloid Polymorphism on Prion-Chaperone Interactions in Yeast. *Viruses.* **11**, 349
30. Masison, D. C., Kirkland, P. A., and Sharma, D. (2009) Influence of Hsp70s and their regulators on yeast prion propagation. *Prion.* **3**, 65–73
31. Guinan, E., and Jones, G. W. (2009) Influence of Hsp70 chaperone machinery on yeast prion propagation. *Protein Pept. Lett.* **16**, 583–586
32. Sweeny, E. A., and Shorter, J. (2008) Prion proteostasis: Hsp104 meets its supporting cast. *Prion.* **2**, 135–140
33. Needham, P. G., and Masison, D. C. (2008) Prion-impairing mutations in Hsp70 chaperone Ssa1: effects on ATPase and chaperone activities. *Arch. Biochem. Biophys.* **478**, 167–174
34. Jung, G., Jones, G., Wegrzyn, R. D., and Masison, D. C. (2000) A role for cytosolic hsp70 in yeast [PSI(+)] prion propagation and [PSI(+)] as a cellular stress. *Genetics.* **156**, 559–570
35. Jones, G. W., and Masison, D. C. (2003) Saccharomyces cerevisiae Hsp70 mutations affect [PSI+] prion propagation and cell growth differently and implicate Hsp40 and tetratricopeptide repeat cochaperones in impairment of [PSI+]. *Genetics.* **163**, 495–506
36. Loovers, H. M., Guinan, E., and Jones, G. W. (2007) Importance of the Hsp70 ATPase domain in yeast prion propagation. *Genetics.* **175**, 621–630
37. Stein, K. C., Bengoechea, R., Harms, M. B., Weihl, C. C., and True, H. L. (2014) Myopathy-causing mutations in an HSP40 chaperone disrupt processing of specific client conformers. *J. Biol. Chem.* **289**, 21120–21130
38. Alberti, S., Halfmann, R., and Lindquist, S. (2010) Chapter 30 - Biochemical, Cell Biological, and Genetic Assays to Analyze Amyloid and Prion Aggregation in Yeast. in *Methods in Enzymology*,

pp. 709–734, Guide to Yeast Genetics: Functional Genomics, Proteomics, and Other Systems Analysis, Academic Press, **470**, 709–734

39. Nathan, D. F., Vos, M. H., and Lindquist, S. (1997) In vivo functions of the Saccharomyces cerevisiae Hsp90 chaperone. *Proc. Natl. Acad. Sci.* **94**, 12949–12956

40. Chernoff, Y. O. (2007) Stress and Prions: Lessons from the Yeast Model. *FEBS Lett.* **581**, 3695–3701

41. Inge-Vechtomov, S. G. (2011) Yeast prions as a model of neurodegenerative infectious amyloidoses in humans. *Russ. J. Dev. Biol.* **42**, 293

42. Dunham, M. J., and Fowler, D. M. (2013) Contemporary, yeast-based approaches to understanding human genetic variation. *Curr. Opin. Genet. Dev.* 10.1016/j.gde.2013.10.001

43. Chernova, T. A., Chernoff, Y. O., and Wilkinson, K. D. (2019) Yeast Models for Amyloids and Prions: Environmental Modulation and Drug Discovery. *Mol. Basel Switz.* 10.3390/molecules24183388

44. Lindström, M., and Liu, B. (2018) Yeast as a Model to Unravel Mechanisms Behind FUS Toxicity in Amyotrophic Lateral Sclerosis. *Front. Mol. Neurosci.* **11**, 218

45. Monahan, Z. T., Rhoads, S. N., Yee, D. S., and Shewmaker, F. P. (2018) Yeast Models of Prion-Like Proteins That Cause Amyotrophic Lateral Sclerosis Reveal Pathogenic Mechanisms. *Front. Mol. Neurosci.* **11**, 453

46. Tenreiro, S., Franssens, V., Winderickx, J., and Outeiro, T. F. (2017) Yeast models of Parkinson's disease-associated molecular pathologies. *Curr. Opin. Genet. Dev.* **44**, 74–83

47. Sporn, Z. A., and Hines, J. K. (2015) Hsp40 function in yeast prion propagation: Amyloid diversity necessitates chaperone functional complexity. *Prion.* **9**, 80–89

48. Bardill, J. P., Dulle, J. E., Fisher, J. R., and True, H. L. (2009) Requirements of Hsp104p activity and Sis1p binding for propagation of the [RNQ(+)] prion. *Prion.* **3**, 151–160

49. Tanaka, M., Chien, P., Naber, N., Cooke, R., and Weissman, J. S. (2004) Conformational variations in an infectious protein determine prion strain differences. *Nature.* **428**, 323–328

50. Derkatch, I. L., Chernoff, Y. O., Kushnirov, V. V., Inge-Vechtomov, S. G., and Liebman, S. W. (1996) Genesis and variability of [PSI] prion factors in Saccharomyces cerevisiae. *Genetics.* **144**, 1375–1386

51. Tanaka, M., Collins, S. R., Toyama, B. H., and Weissman, J. S. (2006) The physical basis of how prion conformations determine strain phenotypes. *Nature.* **442**, 585–589

52. Sharma, J., and Liebman, S. W. (2012) [PSI(+)] prion variant establishment in yeast. *Mol. Microbiol.* **86**, 866–881

53. Toyama, B. H., Kelly, M. J. S., Gross, J. D., and Weissman, J. S. (2007) The structural basis of yeast prion strain variants. *Nature.* **449**, 233–237

54. Luke, M. M., Sutton, A., and Arndt, K. T. (1991) Characterization of SIS1, a Saccharomyces cerevisiae homologue of bacterial dnaJ proteins. *J. Cell Biol.* **114**, 623–638

55. Matveenko, A. G., Barbitoff, Y. A., Jay-Garcia, L. M., Chernoff, Y. O., and Zhouravleva, G. A. (2018) Differential effects of chaperones on yeast prions: CURrent view. *Curr. Genet.* **64**, 317–325

56. Dulle, J. E., Bouttenot, R. E., Underwood, L. A., and True, H. L. (2013) Soluble oligomers are sufficient for transmission of a yeast prion but do not confer phenotype. *J. Cell Biol.* **203**, 197–204

57. Crist, C. G., Kurahashi, H., Nakayashiki, T., and Nakamura, Y. (2006) Conformation preserved in a weak-to-strong or strong-to-weak [PSI+] conversion during transmission to Sup35 prion variants. *Biochimie.* **88**, 485–496

58. Liebman, S. W., and Derkatch, I. L. (1999) The Yeast [PSI +] Prion: Making Sense of Nonsense. *J. Biol. Chem.* **274**, 1181–1184

59. Alberti, S., Halfmann, R., and Lindquist, S. (2010) Chapter 30 - Biochemical, Cell Biological, and Genetic Assays to Analyze Amyloid and Prion Aggregation in Yeast. in *Methods in Enzymology*, pp. 709–734, Guide to Yeast Genetics: Functional Genomics, Proteomics, and Other Systems Analysis, Academic Press, **470**, 709–734

60. Lee, S., Fan, C. Y., Younger, J. M., Ren, H., and Cyr, D. M. (2002) Identification of essential residues in the type II Hsp40 Sis1 that function in polypeptide binding. *J. Biol. Chem.* **277**, 21675–21682

61. Johnson, J. L., and Craig, E. A. (2001) An Essential Role for the Substrate-Binding Region of Hsp40s in Saccharomyces cerevisiae. *J. Cell Biol.* **152**, 851–856

62. Westergard, L., and True, H. L. (2014) Extracellular environment modulates the formation and propagation of particular amyloid structures. *Mol. Microbiol.* **92**, 698–715

63. Huang, V. J., Stein, K. C., and True, H. L. (2013) Spontaneous variants of the [RNQ+] prion in yeast demonstrate the extensive conformational diversity possible with prion proteins. *PloS One.* **8**, e79582

64. Derkatch, I. L., Bradley, M. E., Masse, S. V., Zadorsky, S. P., Polozkov, G. V., Inge-Vechtomov, S. G., and Liebman, S. W. (2000) Dependence and independence of [PSI(+)] and [PIN(+)]: a two-prion system in yeast? *EMBO J.* **19**, 1942–1952

65. Bradley, M. E., Edskes, H. K., Hong, J. Y., Wickner, R. B., and Liebman, S. W. (2002) Interactions among prions and prion "strains" in yeast. *Proc. Natl. Acad. Sci.* **99**, 16392–16399

66. Kryndushkin, D. S., Alexandrov, I. M., Ter-Avanesyan, M. D., and Kushnirov, V. V. (2003) Yeast [PSI+] Prion Aggregates Are Formed by Small Sup35 Polymers Fragmented by Hsp104. *J. Biol. Chem.* **278**, 49636–49643

67. Tipton, K. A., Verges, K. J., and Weissman, J. S. (2008) In vivo monitoring of the prion replication cycle reveals a critical role for Sis1 in delivering substrates to Hsp104. *Mol. Cell.* **32**, 584–591

68. Abrams, J. L., and Morano, K. A. (2013) Coupled Assays for Monitoring Protein Refolding in Saccharomyces cerevisiae. *JoVE J. Vis. Exp.* 10.3791/50432

69. Ferreira, P. C., Ness, F., Edwards, S. R., Cox, B. S., and Tuite, M. F. (2001) The elimination of the yeast [PSI+] prion by guanidine hydrochloride is the result of Hsp104 inactivation. *Mol. Microbiol.* **40**, 1357–1369

70. Hennessy, F., Nicoll, W. S., Zimmermann, R., Cheetham, M. E., and Blatch, G. L. (2005) Not all J domains are created equal: Implications for the specificity of Hsp40–Hsp70 interactions. *Protein Sci. Publ. Protein Soc.* **14**, 1697–1709

71. Ramos, C. H. I., Oliveira, C. L. P., Fan, C.-Y., Torriani, I. L., and Cyr, D. M. (2008) Conserved central domains control the quaternary structure of type I and type II Hsp40 molecular chaperones. *J. Mol. Biol.* **383**, 155–166

72. Pinheiro, G. M. S., Amorim, G. C., Iqbal, A., Almeida, F. C. L., and Ramos, C. H. I. (2019) Solution NMR investigation on the structure and function of the isolated J-domain from Sis1: Evidence of transient inter-domain interactions in the full-length protein. *Arch. Biochem. Biophys.* **669**, 71–79

73. Kaschner, L. A., Sharma, R., Shrestha, O. K., Meyer, A. E., and Craig, E. A. (2015) A conserved domain important for association of eukaryotic J-protein co-chaperones Jjj1 and Zuo1 with the ribosome. *Biochim. Biophys. Acta.* **1853**, 1035–1045

74. Ajit Tamadaddi, C., and Sahi, C. (2016) J domain independent functions of J proteins. *Cell Stress Chaperones.* **21**, 563–570

75. Schilke, B. A., Ciesielski, S. J., Ziegelhoffer, T., Kamiya, E., Tonelli, M., Lee, W., Cornilescu, G., Hines, J. K., Markley, J. L., and Craig, E. A. (2017) Broadening the functionality of a J-protein/Hsp70 molecular chaperone system. *PLOS Genet.* **13**, e1007084

76. Bukau, B., and Horwich, A. L. (1998) The Hsp70 and Hsp60 Chaperone Machines. *Cell.* **92**, 351–366

77. Tom A. Hall (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic acids symposium series.* **41**, 1979–2000

78. Hall, T., Biosciences, I., and Carlsbad, C. (2011) BioEdit: an important software for molecular biology. *GERF Bull Biosci.* **2**, 60–61

79. Chenna, R., Sugawara, H., Koike, T., Lopez, R., Gibson, T. J., Higgins, D. G., and Thompson, J. D. (2003) Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* **31**, 3497–3500

80. Roy, A., Kucukural, A., and Zhang, Y. (2010) I-TASSER: a unified platform for automated protein structure and function prediction. *Nat. Protoc.* **5**, 725–738
81. Yang, J., Yan, R., Roy, A., Xu, D., Poisson, J., and Zhang, Y. (2015) The I-TASSER Suite: protein structure and function prediction. *Nat. Methods.* **12**, 7–8
82. Zhang, Y. (2008) I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics.* **9**, 40
83. *The PyMOL Molecular Graphics System* Schrödinger, LLC
84. Bradley, M. E., and Liebman, S. W. (2003) Destabilizing interactions among [PSI(+)] and [PIN(+)] yeast prion variants. *Genetics.* **165**, 1675–1685
85. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in Saccharomyces cerevisiae. | Genetics [online] https://www.genetics.org/content/122/1/19.short (Accessed December 4, 2019)
86. Guthrie, C., and Fink, G. (1991) *Guide to Yeast Genetics and Molecular Biology.*, Academic Press, Inc, San Diego, CA
87. Dulle, J. E., Stein, K. C., and True, H. L. (2014) Regulation of the Hsp104 middle domain activity is critical for yeast prion propagation. *PloS One.* **9**, e87521

Footnotes:

This work was supported by National Institutes of Health Grant AR068797

Abbreviations and nomenclature:
LGMDD1: : Limb Girdle Muscle Dystrophy Type 1D
LGMD1D: Limb Girdle Muscle Dystrophy Type 1D
HSP: Heat shock protein
DNAJB6: DnaJ heat shock protein family (Hsp40) member B6
SDS-PAGE: sodium dodecyl sulfate–polyacrylamide gel electrophoresis
SDD-AGE: Semi-Denaturating Detergent Agarose Gel Electrophoresis
v.h.: very high
FFL: firefly luciferase
GdHCl: guanidine hydrochloride

TABLES

Table 1. Oligonucleotides used in this study

Oligonucleotide	Description	Sequence
1890	3'-Sis1-S49V	5'-aaatttcaaaggcctctactatctccttaaacttttctgtgtcacctgttgg
1891	5'-Sis1-S49V	5'-ccaacaggtgacacagaaaagtttaaggagatagtagaggcctttgaaattt
1892	3'-Sis1-E53A	5'-ttaaggagatatcagaggcctttgcaattttaaatgatcctcaaaaaagg
1893	5'-Sis1-E53A	5'-ccttttttgaggatcatttaaaattgcaaaggcctctgatatctccttaa
1894	3'-Sis1-N56L	5'-gaaaagtttaaggagatatcagaggcctttgaaattttactagatcctcaaaaaagggaaatat
1895	5'-Sis1-N56L	5'-atatttcccttttttgaggatctagtaaaatttcaaaggcctctgatatctccttaaacttttc
1964	5'-Sis1-ClaI	5'-ccccatcgatatggtcaaggagaca
1965	3'-Sis1-SpeI	5'-gcactagtttaaaaattttcatctatagcacgttt

FIGURES

Figure 1. Novel LGMDD1 disease-associated mutations were identified in the J-domain of DNAJB6. *A*, amino acid sequence comparison of DNAJB6 and orthologs, including the yeast Hsp40, Sis1, aligned using ClustlW2. *B,* protein structure identifying homologous LGMDD1-associated mutations in the Sis1 J domain.

Figure 2. Homologous LGMDD1-associated mutations in the Hsp40 Sis1 differentially impair propagation of [PSI⁺] strains. *A*, yeast 74-D694 *sis1Δ* [*PSI⁺*] strains expressing wild-type *SIS1* or mutated *SIS1* constructs were serially diluted 5-fold and spotted onto 1/4 YPD media and SD-Ade to monitor nonsense suppression of the *ade1-14* allele (n=3). *B*, [*PSI⁺*] colonies expressing the indicated constructs were isolated, grown in liquid YPD overnight, and plated on large 1/4 YPD spread-plates. An average of 626 colonies were counted and scored for strong [*PSI+*], weak [*PSI+*] or [*psi-*]/very weak phenotype. Data were collected from three separate biological replicates. *C*, yeast *sis1Δ* [*PSI+*] strains expressing wild-type *SIS1* or mutated *sis1* constructs were lysed and subjected to a boiled gel assay in order to display the amount of aggregated and monomeric Sup35. Pgk1 is shown as a loading control. Standard protein markers are inappropriate for this particular assay and therefore are not shown (n=4-7).

Figure 3. Homologous LGMDD1-associated mutations in the J domain of Sis1 differentially impair propagation of [RNQ+]. *A*, yeast *sis1Δ* cells harboring the indicated constructs and propagating low [*RNQ+*], very high [*RNQ+*] or [*rnq-*] were serially diluted 5-fold and spotted onto media in order to express the indicated *SIS1* constructs and select for either loss of wild-type Sis1 (-Sis1) or co-expression of wild-type Sis1 (+Sis1). Cells were also spotted on rich media (YPD) (n=3). *B*, sedimentation assay to separate Rnq1 into soluble and insoluble fractions with expression of the indicated constructs (n=6). Total (T), soluble (S) or insoluble (I) fractions were subjected to SDS-PAGE and western blot probing for Rnq1. *C*, SDD-AGE assay for the indicated constructs in low [*RNQ+*] and very high [*RNQ+*] strains (n=4-6).

Figure 4. A homologous LGMDD1-associated mutant in Sis1 exhibits impaired substrate refolding. *A*, the capability for refolding luciferase was measured in Sc37 [*PSI*+] *sis1Δ* yeast strains harboring the indicated construct along with a plasmid expressing firefly luciferase. Yeast were normalized, treated with cycloheximide and subjected to heat shock at 42°C for 22 minutes, followed by recovery at 30°C. Luminescence was measured at the indicated timepoints during recovery and normalized to luminescence of samples without heat shock treatment. The amount of luciferase refolding is plotted as percentage of recovery and represented as mean SEM (n=6).

Figure 5. The LGMDD1-associated J-domain mutation Sis1-S49V has reduced steady state levels in the absence of [PSI⁺]. *A*, western blot analysis showing expression of the indicated constructs in [*rnq-*], low [*RNQ⁺*], and very high [*RNQ⁺*] strains (n=3). *B*, western blot analysis of strong [*PSI⁺*] *sis1Δ* yeast expressing the indicated construct (left). Strong [*PSI⁺*] strains expressing the indicated constructs were passaged twice on 3mM GdHCl, grown on complete media and lysed for SDS-PAGE followed by western blot (right). Pgk1 is shown as a loading control, n=3.