Quantitative Proteomics and Phosphoproteomics Supports a Role for Mut9-Like Kinases
in Multiple Metabolic and Signaling Pathways in *Arabidopsis*

Margaret E. Wilson[1], Shin-Cheng Tzeng[1], Megan M. Augustin[1], Matthew Meyer[1], Xiaoyue
Jiang[2], Jae H. Choi[3], John C. Rogers[2], Bradley S. Evans[1], Toni M. Kutchan[1], Dmitri A.
Nusinow[1*]

[1]Donald Danforth Plant Science Center, St. Louis, MO, 63132, USA

[2]Thermo-Fisher Scientific, San Jose, CA, 95134, USA

[3]Thermo-Fisher Scientific, Rockford, IL, 61101 USA

*To whom correspondence should be addressed dnusinow@danforthcenter.org

Running Title: Quantitative Proteomic Analysis of MLK Family Kinases

Summary/Abstract

Protein phosphorylation is one of the most prevalent post-translational modifications found in eukaryotic systems and serves as a key molecular mechanism by which protein function is regulated in response to environmental stimuli. The Mut9-Like Kinases (MLKs) are a plant-specific family of Ser/Thr kinases that have been linked to light, circadian, and abiotic stress signaling. Here we use quantitative phosphoproteomics in conjunction with global proteomic analysis to explore the role of the MLKs in daily protein dynamics. In the absence of MLK family kinases, proteins involved in light, circadian, and hormone signaling as well as several chromatin modifying enzymes were found to have altered phosphorylation profiles. Additionally, *mlk* mutant seedlings were found to have elevated glucosinolate accumulation and increased sensitivity to DNA damage. Our analysis in combination with previously reported data supports the involvement of MLKs in a diverse set of stress responses and developmental processes, suggesting that the MLKs may serve as key regulators linking environmental inputs to developmental outputs.

Introduction

Protein phosphorylation is one of the most prevalent and well-studied post-translational modifications occurring in eukaryotic cells. This dynamic modification is key in regulating protein function and turnover, making it an integral part of complex signaling networks and regulatory processes. Rapid and reversible post-translational regulation is advantageous to plants, as they are often required to quickly adapt to changing environments. Moreover, protein phosphorylation is at the core of various biological processes including stress response, light signaling, circadian regulation, and hormone perception and transduction. In Arabidopsis, nearly 4% of protein encoding genes are kinases (Wang et al., 2014), which is a testament to the importance of phosphorylation-based protein regulation. Despite the upswing of large-scale phosphoproteomic studies in plant species (Silva-Sanchez et al., 2015), a recent study suggests that the identification

of Arabidopsis phosphoproteins and phosphosites is far from comprehensive (Vlastaridis et al.,

2017).

Recently, a four-member family of Ser/Thr protein kinases known as the MUT9-like

kinase/Photoregulatory Protein Kinases/Arabidopsis EL1-like (MLK/PPK/AEL) kinases, herein

referred to as the MLKs, has been shown to be involved in the phosphoregulation of several key

signaling proteins (Ni et al., 2017; Liu et al., 2017; Su et al., 2017; Chen et al., 2018). The MLKs

are a plant and green algae-specific family of kinases related to casein kinase I (CKI). The MLKs

show significant divergence from CKI, as similarities are restricted to their catalytic domains

(Casas-Mollano et al., 2008). MLK family kinases are capable of phosphorylating histones H3 and

H2A in *Arabidopsis* as well as the green algae *Chlamydomonas*. The *Chlamydomonas* MLK

orthologue, MUT9, is also required for transgene silencing and response to DNA damaging

agents (Casas-Mollano et al., 2008; Jeong Br et al., 2002). In addition to phosphorylating

histones, MLK family kinases phosphorylate proteins involved in multiple signaling pathways.

Early studies of a rice MLK orthologue, EARLY FLOWERING1 (EL1), have linked this kinase

family to hormone signaling and the regulation of flowering time (Dai and Xue, 2010), a role which

is at least in part conserved in *Arabidopsis* (Zheng et al., 2017; Chen et al., 2018; Huang et al.,

2016). MLKs also interact with core components of the morning (Zheng et al., 2017; Su et al.,

2017) and evening (Huang et al., 2016) loops of the *Arabidopsis* circadian clock. The association

of the MLKs with the evening complex components, EARLY FLOWERING 3 AND 4 (ELF3 and

ELF4) is dependent on the presence of the red light receptor phytochrome B (Huang et al., 2016).

Additionally, the MLKs are capable of phosphorylating the blue light receptor CRYPTOCHROME2

and the red light-regulated transcription factor PHYTOCHROME INTERACTING FACTOR 3 (Ni

et al., 2017; Liu et al., 2017). Taken together, these studies suggest that the MLKs provide a link

between light and circadian signaling, which in turn regulates subsequent growth and

development.

In this study, we used quantitative phosphoproteomic techniques to expand our understanding of

the various signaling pathways and cellular protein networks the MLK family of kinases

participate in. We combined isobaric labeling with high pH reversed-phase prefractionation and

TiO_2 based phosphopeptide enrichment to achieve an in-depth phosphoproteomic analysis of

wild-type and *mlk* mutant seedlings at two different time points, one at the end of the day (ZT12)

and the other several hours into the night (ZT14). Over 20,000 phosphosites mapping to nearly

5,000 protein groups were identified. In the MLK mutant backgrounds, enzymes involved in

glucosinolate metabolism were altered in abundance and proteins involved in a diverse set of

biological processes including RNA processing, chromatin organization, and stress responses

were differentially phosphorylated. The confluence of stress and chromatin factors suggested that

MLKs may also be involved in regulating DNA-damage responses in *A. thaliana*. This hypothesis

was addressed by assessing the sensitivity of *mlk* mutants to DNA-damaging agents.

Experimental Procedures

Plant Material

The *mlk1* (SALK_002211; AT5G18190), *mlk2* (SALK_064333; AT3G03940), and *mlk3*

(SALK_017102; AT2G25760) mutant lines were obtained from the ABRC (Ohio State University).

The *mlk4* (GABI_756G08; AT3G13670) mutant line was obtained from the Nottingham

Arabidopsis Stock Centre. All are in the Colombia (Col-*0*) background and were isolated as

previously described (Huang et al., 2016).

Tissue Collection for Mass Spectrometry

Arabidopsis wild type (Col-*0*) and mutant seedlings were grown on sterilized qualitative filter paper

(Whatman) overlaid on ½ x MS (Murashige and Skoog) plates containing 1% sucrose and 0.8%

agar at 22˚C. Seedlings were entrained under 12 h white light (100-110 μmol/m^2/s)/12 h dark

cycle. Tissue was collected on the 10[th] day of growth immediately prior to lights off [Zeitgeber 12,

(ZT12)] and after 2h of dark (ZT14).

Protein Isolation and Digestion

The seedlings were transferred into a liquid N_2 chilled 35ml ball mill and disrupted in a reciprocal

mixer mill [30 hz, 45 seconds, repeated 3 times (Retsch USA)] under liquid nitrogen. Ground

tissue was gently resuspended in 1 mL (approximately 1 packed tissue volume) of SII buffer (100

mM sodium phosphate, pH 8.0, 150 mM NaCl, 5 mM EDTA, 5 mM EGTA, 0.1% Triton X-100, 1

mM PMSF, 1x protease inhibitor cocktail [Roche], 1x Phosphatase Inhibitors II & III [Sigma], and

50 µM Mg-132 [Peptides International]) and sonicated twice at 40% power, 1 second on/off cycles

for 20 s total on ice (Fisher Scientific model FB505, with microtip probe). Extracts were clarified

by centrifugation twice at 4˚C for 10 min at ≥20,000xg. Protein concentrations were determined

by BCA protein assay (Thermo-Fisher Scientific, Rockford, IL). Protein samples were reduced

with 10 mM TCEP and alkylated with 25 mM iodoacetamide prior to trypsin digestion in 1/40

enzyme/protein ratio at 37°C overnight.

Phosphopeptide Enrichment

Phosphopeptide enrichment was performed using the High-Select™ TiO_2 Phosphopeptide

Enrichment kit (Thermo Scientific PN32993) following vendor's protocol. Briefly, dried peptides

were reconstituted in 150 µL of binding/equilibration buffer provided and applied to the TiO_2 spin

that was previously equilibrated with binding buffer/equilibration. After reapplying sample once,

the tip was sequentially washed twice with 20 µL of binding buffer and wash buffer, and once with

20 µL of LC-MS grade water. Bound peptides were eluted by two applications of 50 µL of elution

buffer (also provided). Eluates containing the enriched phosphopeptides were dried down and

subsequently resuspended with 50 µl 0.1% formic acid for peptide concentration measurement

using the Pierce Quantitative Colorimetric Assay kit (Thermo Scientific PN23275).

Tandem Mass Tag (TMT) Labeling

100 µg of each digested sample was added to 100 µL of 100 mM HEPES pH 8.5 buffer. A reference pooled sample composed of equal amounts of material from all samples was also generated to link TMT experiments. Isobaric labeling of the samples was performed using 10-plex tandem mass tag (TMT) reagents (Thermo Fisher Scientific, Rockford, IL). All individual and pooled samples were labeled according to the TMT 10-plex reagent kit instructions. Briefly, TMT regents were brought to room temperature and dissolved in anhydrous acetonitrile. Peptides were labeled by the addition of each label to its respective digested sample. Labeling reactions were incubated for 1 h at room temperature. Reactions were terminated with the addition of hydroxylamine.

High pH Reverse Phase Fractionation

High pH reverse phase fractionation was performed using the Pierce High pH Reversed-Phase Peptide Fractionation Kit (Thermo Scientific PN84868) according to manufacturer's instructions. Briefly, peptide samples were dissolved in 300 µL of 0.1% TFA solution in LC-MS grade water. and subsequently loaded onto reversed-phase fractionation spin columns also equilibrated with 0.1% TFA. Samples were then washed with 300 µL of 5% ACN/ 0.1% TEA to remove unreacted TMT reagent. Peptides were eluted into 8 peptide fractions with an ACN step gradient (i.e. 10%, 12.5%, 15%, 17.5%, 20%, 22.5%, 25%, and 50%). Samples were acidified and dried down prior to LC-MS.

LC-MS/MS Analysis

Two microliters (one microgram) of sample was injected onto a 0.075 x 500 mm EASY-Spray Pepmap C18 column equipped with a 0.100 x 5 mm EASY-Spray Pepmap C18 trap column (Thermo-Fisher Scientific, San Jose, CA) attached to an EASY-nLC 1000 (Thermo-Fisher Scientific, San Jose, CA). The peptides were separated using water (A) and acetonitrile (B)

containing 0.1% formic acid as solvents at a flow rate of 300 nL per minute with a three-hour

gradient. Data were acquired in positive ion data dependent mode on an Orbitrap Fusion Lumos

mass spectrometer (Thermo-Fisher Scientific, San Jose, CA) with a resolution of 120,000 (at m/z

200) and a scan range from m/z 380-1500. Precursor isolation was performed using the

quadrupole prior to either CID activation in the ion trap and detection in the Orbitrap at a resolution

of 30,000 or HCD activation with detection in the Orbitrap at a resolution of 60,000.

Data Analysis

All MS/MS data were analyzed using Proteome Discoverer 2.1 (Thermo-Fisher Scientific, San

Jose, CA). The search algorithm used in the study was Byonic v2.11 as part of the Proteome

Discoverer software platform. Precursor ion mass tolerance was set to 10 ppm, fragment ion

tolerance was 20 ppm; up to 2 missed cleavages were allowed. Carbamidomethylation (+57.021

Da) on cysteine and TMT tag (+229.163 Da) on peptide N-termini as well as lysine residues were

set as static modifications. Dynamic modifications included acetylation (+42.011 Da) on protein

N-termini, oxidation (+15.995 Da) on methionine and phosphorylation (+79.966 Da) on serine,

threonine, and tyrosine. Data were searched against TAIR10 database (20101214, 35,386

entries) with FDR set to 1%.

For quantitation, reporter ion intensity integration tolerance was set to 20 ppm. Reporter ion

abundances were corrected for isotopic impurities based on manufacturer's specifications. For

each peptide, a minimal average reporter S/N threshold of 2 and a co-isolation threshold of 100%

are required. The S/N values for all peptides were summed within each TMT channel, and each

channel was scaled according to the reference channel. Both unique and razor peptides were

used for quantification.

Peptides of altered abundance were identified from the Byonic output list generated from HCD

MS2 analysis using Microsoft Excel. Abundance ratios for mutant/wild type pairwise comparisons

were calculated from the average peptide abundance of mutant and wild-type biological

replicates. Only peptides identified in at least 2 biological replicates were considered for further

analysis. Statistical significance was determined by Student's *t*-test (*P*-value ≤ 0.05).

Bioinformatic Analysis

The Motif-X algorithm (Chou and Schwartz, 2011) was used to extract significantly enriched

phosphorylation motifs from *mlk1/2/3* and *mlk1/3/4* phosphopeptide data sets. Only

phosphopeptides with high confidence phosphorylation sites were used in analysis. The peptides

were aligned and extended to a width of 15 amino acids using the online utility

PEPTIDEXTENDER ver.0.2.2 alpha (schwartzlab.uconn.edu/pepextend/). The aligned peptides

were used to extract motifs. The probability threshold was set to p-value ≤ 10^{-5}; the occurrence

threshold was set to 10. The default IPI Arabidopsis Proteome data set was used as the

background data set.

Enrichment analysis of Gene ontology (GO) categories was performed with g:Profiler (Reimand

et al., 2016). AGI accession numbers for Arabidopsis were uploaded to the g:Profiler webserver

(http://biit.cs.ut.ee/gprofiler/) and GO enrichment was determine using default settings

(significance level 0.05). Enriched terms were summarized and redundancy removed using the

online tool REVIGO (Supek et al., 2011). Semantic similarity threshold (dispensability) was set to

0.5 (default) for all global proteome analysis and cellular component category of the

phosphoproteome analysis. Dispensability was increased to 0.7 for all other analysis.

Glucosinolate Extraction and Analysis by HPLC and LC-MS/MS

Arabidopsis seeds (Col-0*, mlk 1/2/3,* and *mlk 1/3/4*) were sown on ½ x MS (Murashige and Skoog)

plates containing 1% sucrose and 0.8% agar and grown under 12 h white light (100-110

μmol/m^2/s)/12 h dark cycle at 22˚C for 10 days prior to harvest at ZT12. Glucosinolates were

extracted from approximately 350 mg of whole seedlings and desulfonated (in quadruplicate) as

previously described (Crocoll *et al.* 2016) using sinigrin as an internal standard. Desulfo-GLS

extracts were analyzed by HPLC (Waters) equipped with a photodiode array detector and

separated using a Gemini C-18 column (150 X 2.00 mm, 5 µm; Phenomenex) with a flow rate of

0.5 mL per minute and the following solvents and binary gradient: solvent A-water and solvent B-

acetonitrile; where solvent B was held at 1.5% for 1 min, then 1-6 min 1.5-5% B, 6-8 min 5-7% B,

8-18 min 7-21% B, 18-23 min 21-29% B, 23-30 min 29-43% B, 30-33 min 43-100% B, 33-37 100%

B, 37-38 min 100-1.5% B, and held at 1.5% B for an additional 7 minutes. GLS peaks were

identified using previously published UV spectra in addition to reported relative retention times

and quantitated using peak areas of desulfo-GLS and internal standard along with published

response factors (Brown *et al.* 2003, Grosser and Dam 2017). GLS identities were confirmed by

LC-MS/MS (SCIEX 6500 QTRAP, Framingham, MA) using enhanced product ion (EPI; ion trap

MS/MS) scans to verify the presence of previously published fragment ions (Crocoll et al. 2016)

from each glucosinolate ion. Mass spectrometric data were collected in positive ion mode using

the same gradient/solvents/column as for HPLC-UV analysis with the following source conditions:

curtain gas, 20; ion-spray voltage, 5500 V; temperature, 500° C; gas 1, 40; gas 2, 45; declustering

potential, 80 V; entrance potential, 10 V; collision energy 20 eV.

MMS Treatment

Arabidopsis wild type (Col-*0*) and mutant seed was surface-sterilized and sown on ½ x MS

(Murashige and Skoog) plates containing 1% sucrose and 0.8% agar with or without methyl

methanesulfonate (MMS, Sigma). After stratification for 2 days at 4°C, seedlings were grown

under 12 h white light (100-110 µmol/m^2/s)/12 h dark cycle at 22˚C. For growth sensitivity assays,

seedlings were germinated on control media and after 5 days of growth were transferred to MMS

treatment media. Fresh weight was measured after 15 days of growth in the presence of MMS.

For post-germination developmental assessment, seed was germinated on ½ x MS plates

containing 1% sucrose, 0.8% agar, and 150 ppm MMS. Seedlings were imaged and scored for

arrest at 12 days after germination.

UV-C Tolerance Assay

Whole-plant sensitivity to UV-C (254 nm) was evaluated as described in Castells et al. (2010) with

the following modifications. 8-day old seedlings were irradiated with 2000 or 4000 J m^{-2} of UV-C

twice during a 48 hr time period using a Stratalinker® UV Crosslinker 1800. Following each

treatment, plants were returned to growth conditions of 12 h white light (100-110 µmol/m^2/s)/12 h

dark cycle at 22˚C. After 5 days of recovery, phenotypes were measured and seedlings were

imaged.

Results

Proteomic Analysis Reveals Changes in Stress Response Pathways in *mlk* Mutant

Seedlings

Tandem mass tag (TMT) labeling combined with high pH reversed-phase fractionation and

tandem mass spectrometry was used to quantify the regulatory effects of the MLKs on proteome

dynamics (**Fig. 1**). Wild-type and *mlk* mutant *Arabidopsis* seedlings were entrained under 12 h

light and 12 h dark conditions. We compared *mlk1/2/3* and *mlk1/3/4* mutant seedlings, as we were

unable to isolate viable *mlk1/2/3/4* mutant seed (Huang et al., 2016; Liu et al., 2017). This mutant

combination will allow us to assess potential redundancy within the MLK family and facilitate the

identification of *mlk2* or *mlk4* specific changes. As the MLKs have been previously associated

with light-signaling pathways, we collected tissue either immediately prior to lights off [Zeitgeber

12, (ZT12)] or after 2h of dark (ZT14). We identified nearly 50,000 peptides combined, mapping

to over 7,500 protein groups at both ZT12 and ZT14. Pairwise comparisons between *mlk* mutants

and WT were performed to identify peptides showing altered abundance in the MLK mutant

backgrounds at both the ZT12 and ZT14 time-points (**Supplemental Dataset S1**). Peptides were

classified as altered in abundance if both the $\log_2$ FC was at least ±1 (2 fold-change) and the p-

value < 0.05. Only 13 unique proteins met our altered abundance criteria in the *mlk1/2/3* mutant

when compared to wild type at both ZT12 and ZT14 (**Fig. 2 and Supplemental Table S1**). In the

mlk1/3/4 mutant background, more than 225 peptides mapping to over 110 unique proteins were

found to meet our altered abundance threshold at ZT12 and ZT14 when compared to wild type

(**Fig. 2 and Supplemental Table S1**). These results suggest that the *mlk1/3/4* mutant

combination has a greater impact on the global proteome than the *mlk1/2/3* mutant combination

at the light-to-dark transition.

Quantitative Phosphoproteomic Comparisons of *mlk* Mutants

The MLKs have been shown to physically interact with and phosphorylate important regulatory

proteins (Liu et al., 2017; Chen et al., 2018; Dai and Xue, 2010; Ni et al., 2017). Thus, to gain

insight into the role of the MLKs in global phosphorylation, we characterized the

phosphoproteome of wild type, *mlk1/2/3,* and *mlk1/3/4* mutant seedlings in the light (ZT12) and

after transition to dark (ZT14). We applied a TiO_2 based phosphopeptide enrichment technique to

the TMT-10plex labeled samples described above to achieve an in-depth phosphoproteomic

analysis using the Thermo Scientific Orbitrap Fusion Lumos Tribrid mass spectrometer **(Fig. 1)**.

Byonic software, run as a node within the Proteome Discover V2.1 platform was used to identify

and derive relative quantitation for phosphoproteins. Using this strategy, a combined total of

23,386 phosphosites on 15,222 unique peptides mapping to 4,854 protein groups were identified

at ZT12. Slightly fewer were identified at ZT14, with 19,947 phosphosites on 12,818 unique

peptides mapping to 4,467 protein groups being identified. At ZT12, over 80% of the identified

phosphosites were serine residues, approximately 15% were threonine, and less than 2% were

tyrosine (**Fig. 3A**), which is consistent with phosphosite distributions previously reported for

Arabidopsis (Champion et al., 2004; Sugiyama et al., 2008). The phosphosite residue distributions

were similar at ZT14 (**Figure 3A**).

We performed *mlk*-to-WT pairwise comparisons for each time point (**Supplemental Table S1 and**

Dataset S2) to identify peptides that showed altered phosphorylation in the absence of the MLKs

at either ZT12 or ZT14. Peptides were considered differentially phosphorylated if they had a

minimum $\log_2$ FC of ±0.585 (1.5 fold-change) and a p-value < 0.05. We identified 113

phosphopeptides corresponding to 93 unique protein groups that met the cutoff in the *mlk1/2/3*

ZT12 set. In the *mlk1/3/4* ZT12 set, 429 phosphopeptides corresponding to 284 unique proteins

showing differential phosphorylation were identified, which corresponds to a 300% increase (**Fig.**

3B). 170 and 274 phosphopeptides, corresponding to 149 and 215 unique protein groups, have

altered phosphorylation in the *mlk1/2/3* ZT14 and *mlk1/3/4* ZT14 analysis, respectively (**Figure.**

3B and Supplemental Table S1). The phosphosite residue distribution was similar in all datasets

analyzed (**Figure 3A**). These data show that the *mlk1/3/4* mutant combination has a larger impact

on the phosphoproteome than the *mlk1/2/3* mutant combination at each time point, particularly at

ZT12. This observation supports a role for MLK4 in regulating the phosphoproteome in a light-

dependent manner, which is in agreement with the known role of MLK4 in phosphorylating key

components of blue- and red-light signaling pathways (Liu et al., 2017; Ni et al., 2017).

Over 50% (48 out of 93) of the differentially phosphorylated proteins identified in the *mlk1/2/3*

ZT12 set were also identified in the *mlk1/3/4* ZT12 set. In contrast, over 80% of those identified

in the *mlk1/3/4* ZT12 set were specific to the *mlk1/3/4* mutant (**Fig. 3C**). When comparing the

ZT14 sets, 42 phosphoproteins were shared, accounting for 28% and 19.5% of the proteins

identified in the *mlk1/2/3* ZT14 and *mlk1/3/4* ZT14 sets, respectively (**Fig. 3D**). Many of the

proteins shared between *mlk1/2/3* and *mlk1/3/4* protein sets are involved in gene silencing and

chromatin organization (**Supplemental Dataset S2**), supporting a conserved role for the MLK

family kinases in these processes (Casas-Mollano et al., 2008; Jeong Br et al., 2002). Some of

these proteins, including SUO and SERRATE (SE) – which are involved in microRNA biogenesis

pathways, exhibit altered phosphopeptide abundance at both ZT12 and ZT14. However, proteins

involved in chromatin organization, such as Increased in Bonsai Methylation 1 (IBM1) and

SPLAYED (SYD), only showed altered phosphorylation in both *mlk1/2/3* and *mlk1/3/4* mutant

backgrounds at ZT12 (**Supplemental Dataset S2**). These results suggest that the MLKs are

involved in regulating gene expression, possibly through modulating light-dependent chromatin

organization.

MLKs Influence Diverse Kinase Signaling Networks

The proteins identified as differentially phosphorylated in the *mlk1/3/4* mutant background at both

ZT12 and ZT14 are associated with a diverse set of biological processes, suggesting a possible

disruption of multiple protein kinase networks. Motif-X (http://motif-x.med.harvard.edu/motif-

x.html; (Chou and Schwartz, 2011; Schwartz and Gygi, 2005) was used to isolate

overrepresented sequence motifs present in the phosphopeptide sets that are associated with

known kinase families. Following extension of differentially phosphorylated high-confidence,

unambiguous   peptides   using   PEPTIDEXTENDER   ver.0.2.2   alpha

(http://schwartzlab.uconn.edu/pepextend), the resulting 15-mers were submitted for motif

analysis using a significance threshold of $p < 10^{-6}$ and a minimum occurrence requirement of 20.

Peptides exhibiting increased or decreased abundance when compared to wild type were

analyzed separately. Three serine phosphorylation (Sp) motifs (S-x-x-K, R-x-x-S, and K-x-x-S)

were found to be overrepresented in peptides that were decreased in abundance at ZT12 in the

mlk1/3/4 mutant seedling background (**Fig. 4A**). The K-x-x-S motif, along with an acidic S-type

motif (S-x-x-x-x-x-E), was also found to be overrepresented at ZT14 in *mlk1/3/4* mutants (**Fig.**

4B). The CDPK-SnRK superfamily of protein kinases is known to recognize R/K-x-x-S/T basic

motifs. The R-x-x-S motif has also been associated with the AGC family kinases, PKA and PKC

(Rademacher and Offringa, 2012; Marondedze et al., 2016), which are involved in mid- to late-

day rhythmic phosphorylation (Choudhary et al., 2015). The kinase(s) responsible for

phosphorylation at the S-x-x-K site in plants is less well-characterized. However, it was recently

shown that the highly conserved eukaryotic cyclin B-dependent protein kinase Cdk1 recognizes

several non-S/T-P motifs including the S/T-x-x-R/K motif (Suzuki et al., 2015). The classical

minimal motif required for recognition by proline-directed kinases families (mitogen-activated

protein kinase (MAPK), cyclin-dependent kinase (CDK), and glycogen synthase kinase 3 (GSK-

3)), S-P, was found to be overrepresented in peptides that were increased in abundance in

mlk1/3/4 mutants at ZT12 and ZT14. While the S-x-S motif, which is associated with the receptor-

like protein kinase (RLK) family (van Wijk et al., 2014), was only found to be overrepresented at

ZT12 (**Fig. 4**). Using the same parameters, there were no overrepresented motifs identified from

the sites that decreased in abundance in neither of the *mlk1/2/3* data sets. However, if the

minimum occurrence was reduced to 10, the R-x-x-S and S-P motifs were found to be

overrepresented in peptides with decreased or increased abundance at ZT12 in *mlk1/2/3* mutants

(**Supplemental Figure S2**). The diversity of identified overrepresented kinases motifs suggests

that MLK family kinases influence numerous biological processes through systemic regulation of

multiple kinase signaling networks.

MLKs May Regulate Glucosinolate Metabolism

In silico classification using Gene Ontology (GO) analysis (https://biit.cs.ut.ee/gprofiler/) revealed

that proteins exhibiting altered abundance were associated with biotic and abiotic stresses (**Fig.**

5). To simplify the enriched GO term lists and focus on the most relevant terms, we performed

additional analysis using REVIGO (default settings, dispensability threshold = 0.7) to remove

functionally redundant terms (Supek et al., 2011). Proteins involved in glucosinolate biosynthesis

(GO:0019761) and related processes (GO:1901659, GO:0016143, and GO:0044272) were found

to have increased abundance in *mlk1/3/4* mutant seedlings at ZT12 and in both *mlk1/2/3* and

mlk1/3/4 mutant seedlings at ZT14 (**Fig. 5**). Nearly 75% of the peptides with an increased

abundance of 3-fold or greater in the *mlk1/3/4* mutant seedlings at ZT14 mapped to proteins

directly involved in glucosinolate biosynthesis (**Supplemental Dataset S1**). These proteins

include enzymes responsible for the early reactions leading to methionine-derived glucosinolates

(branched-chain aminotransferase 4 (BCAT4) and methylthioalkymalate synthase 1 (MAM1)) as

well as, desulfo-glucosinolate sulfotransferase 17 and 18 (SOT17 and SOT18), which are

involved in the final step of glucosinolate core structure biosynthesis (Sønderby et al., 2010).

Other proteins involved in glucosinolate biosynthesis that showed increased abundance in

mlk1/3/4 mutants compared to wild type include isopropymalate dehydrogenase 1 (IMD1),

iospropylmaate isomerase 2 (IPMI2), 2-isopropylmalate synthase 2 (IMS2), flavin-

monooxygenase glucosinolate S-oxygenase 1 (FMO GS-OX1) and the cytochrome P_{450} proteins

CYP83A1 and CYP79F1 (**Supplemental Dataset S1**). Proteins involved in glucosinolate

catabolism such as glucoside glucohydrolase 2 (TGG2), nitrile specifier protein 1 (NSP1), and

beta glucosidase 34 and 35 (BGLU34 and BGLU35) were decreased in abundance in *mlk1/3/4*

mutant seedlings at ZT12 compared to wild type.

To begin testing the hypothesis that MLKs are involved in the regulation of glucosinolate

metabolism, we quantified glucosinolate levels at the end of day (ZT12) when GLS levels are

expected to peak (Huseby et al., 2013). Total GLSs were extracted from whole seedlings and

analyzed by HPLC. Peaks corresponding to individual GLSs were identified by comparison with

published UV absorbance spectra and expected retention times and the identities were validated

using LbyC-MS/MS. These analyses revealed an increase in aliphatic glucosinolates (Met-

derived) in both *mlk1/2/3* and *mlk1/3/4* mutant seedlings compared to wild type, whereas the

levels of indolic glucosinolates (Trp-derived) remained unchanged in the mutant backgrounds

(**Fig. 6**). Interestingly, the first seven glucosinolates originating from the earliest steps in the Met-

derived glucosinolate biosynthetic pathway were increase 2- to 4-fold over wild type (**Fig. 6**), a

pattern which correlates with the increased abundance of glucosinolate-associated biosynthetic

enzymes (BCAT4, MAM1, SOT17/18, etc.) in the *mlk* mutants. Together these findings support a

role for the MLKs in early stages of aliphatic glucosinolate biosynthesis and overall glucosinolate

metabolism.

In addition to glucosinolate biosynthetic enzymes, several proteins involved in hormone signaling

and diverse stress responses exhibited differential abundance in *mlk1/3/4* mutant seedlings

compared to wild type at either ZT12 or ZT14, including BRI1-EMS-SUPPRESSOR 1 (BES1),

SUPER SENSITIVE TO ABA AND DROUGHT2 (SAD2), CORONATINE INDUCED 1 (CORI3),

pathogenesis-related gene 5 (PR5), lipoxygenase 2 (LOX2), thylakoidal ascorbate peroxidase

(TAPX), and cold regulated 15a and b (COR15A and COR15B). Peptides mapping to the blue

light receptor cryptochrome 2 (CRY2) also showed an increased abundance of 2-fold in the

mlk1/3/4 mutant when compared to wild type at ZT12 (**Supplemental Dataset 1**). These

observations are in agreement with the previously reported role of MLKs in hormone signaling,

stress response, and light signaling (Liu et al., 2017; Dai and Xue, 2010; Casas-Mollano et al.,

2008; Ni et al., 2017; Chen et al., 2018).

MLKs Influence the Phosphorylation Status of Nuclear Localized Proteins

Using g:Profiler (https://biit.cs.ut.ee/gprofiler/), 149 GO terms were found to be enriched in the

differentially phosphorylated proteins from the *mlk1/3/4* ZT12 set; 107 of these were classified as

biological process, 34 as cellular component, and 5 as molecular function. Forty-nine terms were

found to be enriched in the *mlk1/3/4* ZT14 set, 28 biological process and 21 cellular components.

Fewer terms were found to be enriched in the *mlk1/2/3* phosphoprotein sets, with only 8 enriched

terms at ZT12 and 18 at ZT14. A complete list of enriched GO terms can be found in

Supplemental Dataset S2. Functionally redundant terms were removed using REVIGO (default

settings, dispensability threshold = 0.7 (cellular component) or 0.5 (biological process);

Supplemental Tables S2-3; (Supek et al., 2011)). Under the cellular component category, there

was strong enrichment for terms associated with the nucleus: 'nucleus' (GO:0005634), 'nuclear

part' (GO:0044428), and 'nucleoplasm' (GO:0005654). Terms including 'chromosome'

(GO:0005694), 'chromosomal part' (GO:0044427) and 'chromatin' (GO:0000785) were also found

to be enriched in at least one of the differentially phosphorylated protein lists (**Fig. 7A**). These

results are in agreement with the known nuclear localization of the MLKs, and their role in

modifying chromatin (Wang et al., 2015a; Huang et al., 2016; Su et al., 2017).

Circadian-Associated Proteins Exhibit Altered Phosphorylation in *mlk1/3/4* Mutant
Seedlings

The *mlk1/3/4* ZT12 differentially phosphorylated protein set was enriched in proteins associated

with rhythmic processes (GO:0048511) and/or circadian rhythms (GO:0007623). These

observations agree with previous reports linking the MLKs to light signaling and circadian

regulation (Huang et al., 2016; Ni et al., 2017; Liu et al., 2017; Su et al., 2017; Zheng et al., 2017).

The core circadian clock proteins PSEUDO-RESPONSE REGULATOR 7 (PRR7), TIME FOR

COFFE (TIC), and REVEILLE 8 (RVE8) were all differentially phosphorylated in the *mlk1/3/4*

mutant background at ZT12 (**Table 1 and Supplemental Table S3**). We observed decreased

phosphorylation of TIC at S324 and PRR7 at S355 and S275, while RVE8 showed increased

phosphorylation of the C-terminal half (**Table 1**). The red-light photoreceptor phytochrome B

(PHYB) and the transcriptional master regulator ELONGATED HYPOCOTYL 5 (HY5) also

showed reduced phosphorylation at Threonine 42 and T64, respectively (**Table 1**); to the best of

our knowledge, these phosphosites are previously unreported. A complete list of circadian-

associated proteins with altered phospho-abundance in the *mlk1/3/4* mutant at ZT12 can be found

in **Table 1**.

Gene Ontology Analysis Reveals Enrichment of Proteins Involved in Chromatin
Organization

Due to the large number of enriched GO terms identified from both the *mlk1/3/4* ZT12 and

mlk1/3/4 ZT14 list, the dispensability threshold was reduced to 0.5 for REVIGO analysis. This

resulted in the identification of 34 and 15 representative and non-redundant enriched biological

process terms in *mlk1/3/4* ZT12 and *mlk1/3/4 Z*T14 altered phosphoprotein lists, respectively.

Many of the enriched GO terms and their underlying gene identifiers were shared between the

*mlk1/3/4 Z*T12 and ZT14 sets including, 'organic cyclic compound metabolism', GO:1901360,

'nitrogen compound metabolism', GO:0006807, 'chromosome organization', GO:0051276 and

'negative regulation of gene expression' GO:0006807 (**Fig. 7B-C and Supplemental Table S3**).

The chromatin modifying proteins BRAHMA (BRM), SIN3-like3 (SNL3), vernalization5/VIN3-like

(VEL1), and high mobility ground B1 (HMGB1) were shared amongst these GO terms. Additional

proteins associated with chromatin modifications were present in the *mlk1/3/4 Z*T12 list including

the histone methyltransferase EARLY FLOWERING IN SHORT DAYS (EFS)/SET DOMAIN

GROUP 8 (SDG8), the histone acetyltransferase TBP-ASSOCIATED FACTOR 1 (TAF1), as well

as IBM1, actin-related protein 4 (ARP4), alfin-like 7 (AL7), GLIOMAS 41 (GAS41/YAF9a), and

stress-induced histone H2A protein 9 (HTA9). Few biological process GO-terms were found to be

enriched in the *mlk1/2/3* data sets, but those that were ('regulation of gene expression',

'epigenetic' (ZT12) and 'chromosome organization', 'chromatin organization' and 'mitotic sister

chromatid cohesion' (ZT14)) were also found to be enriched in the *mlk1/3/4* protein list

(**Supplemental Dataset S2**). These results support a role for the MLKs in regulating chromatin

organization and gene expression at the assessed time points.

mlk1/3/4 **Mutants Show Altered Phosphorylation of Proteins Involved in Nuclear**

Organization and DNA Damage Response

Our analysis of differentially phosphorylated peptides has shown that the loss of functional

versions of *mlk1*, *mlk3*, and *mlk4* at ZT12 has the greatest impact on both the global- and

phospho- proteomes. Therefore, we chose to expand on our analysis exclusively for the *mlk1/3/4*

ZT12 data set. The phosphoproteins that were increased or decreased in abundance

independently were analyzed to further elucidate biological processes influenced by the MLKs at

the end of the day (ZT12). Of the 429 phosphopeptides found to have altered abundance in the

mlk1/3/4 mutant background at ZT12 (**Supplemental Dataset S2**), 133 were increased and 296

were decreased in abundance, mapping to 103 increased and 190 decreased unique proteins.

Interestingly, 9 gene identifiers were shared between the increased and decreased groups,

including several that are involved in various aspects of nuclear organization such as LITTLE

NUCLEI 1/ CROWED NUCLEI 1 (LINC1/CRWN1), VEL1, and HMGB1 (**Fig. 8A and**

Supplemental Dataset 2). Proteins that were increased in abundance were associated with the

representative GO-terms 'response to organic substance' (GO:0010033) and 'response to

stimulus' (GO:0050896) (**Fig. 8B**). However, we found that the majority of GO terms enriched in

the inclusive set (both increased and decreased peptides) such as 'RNA processing'

(GO:0006396), 'chromosome organization' (GO:0051276), and 'developmental processes'

(GO:0032592) are associated with decreased phosphorylation (**Fig. 8C**). The GO term 'cellular

response to DNA damage stimulus' (GO:0006974), was also found to be enriched in proteins

exhibiting decreased phosphorylation. Proteins associated with this term include a catalytic

subunit of DNA polymerase alpha INCARVATA2 (ICU2), as well as X-ray cross complementation

group4 (XRCC4) and MUTM homolog-1 (MMH-1) both of which are directly involved in DNA repair

(West et al., 2000; Ohtsubo et al., 1998; Barrero et al., 2007).

***mlk* Mutants Show Increased Sensitivity to DNA-damaging Agents**

It is well established that nuclear organization and chromatin dynamics strongly influence DNA

damage repair efficacy (Reviewed in Vergara and Gutierrez, 2017; Donà and Mittelsten Scheid,

2015). Additionally, it has been previously shown that the *Chlamydomonas* MLK orthologue,

Mut9, is required for survival when grown in the presence of genotoxic agents (Jeong Br et al.,

2002). Since many of the proteins associated with changes in phosphorylation abundance in the

mlk1/3/4 mutants at ZT12 are proteins associated with nuclear organization (LINC/CRWN family

members, SAD1/UNC-84 domain protein 2 (SUN2), BRM, and SYD) and DNA damage, we

sought to further explore what role the *Arabidopsis* MLKs might play in DNA damage response.

To do so, we evaluated the sensitivity of mutant and wild type seedlings to the genotoxic agent

methyl methane sulfonate (MMS) and UV-C. MMS is a monofunctional DNA alkylating agent

which induces replication fork stalling and subsequent double strand breaks (Ensminger et al.,

2014). In addition to the *mlk1/2/3* and *mlk1/3/4* mutants, the *mlk4* single and *mlk* quadruple

amiRNA line (amiR^{4k})(Liu et al., 2017) were included in our analysis. Five days after germination,

mutant and wild type seedlings were transferred to solid media containing increasing levels of

MMS and allowed to grow for an additional 14 days, after which point seedlings were imaged and

weighed (**Fig. 9A-B**). After 2 weeks of treatment, all genotypes showed a reduction in aerial mass

correlating with increasing levels of MMS. This reduction was greater in *mlk* mutants compared

to wild-type seedlings, which exhibited a less than 10% reduction in fresh weight (FW) when

grown in the presence of 50 ppm MMS compared to the mock treated seedlings. The *mlk4* single

mutant seedlings had a more than 20% reduction of FW compared to the mock treated seedlings.

The reductions were even greater in the *mlk1/3/4* and amiR^{4K} lines, with a FW of 61% and 56%

of mock treated seedlings, respectively (**Fig. 9B**). The *mlk1/3/4* and amiR^{4k} mutants continued to

show greater reduction of FW than wild type in all concentration tested. At 100 ppm MMS,

mlk1/2/3 seedlings showed approximately a 10% greater reduction in FW than what was observed

for wild type (i.e. 49% and 61%, respectively, **Fig. 9B**). In addition to overall growth reduction, the

appearance of chlorotic tissue was observed in the *mlk134* mutants growing on as little as 50 ppm

MMS and in the amiR^{4K} mutants at 75 PPM MMS (**Fig. 9A**). Growth in the presence of 150 ppm

MMS caused severe growth reduction and lethality in all post-germination growth development in

the presence of 150 ppm MMS genotypes assessed, thus seedlings were imaged but not

weighted (**Fig. 9A**). Post-germination growth was found to be severely impaired in the *mlk1/3/4*

mutant, with more than 65% of seedlings exhibiting complete developmental arrest compared to

approximately 10% in wild-type seedlings (**Fig. 9C-D**). The *amiR^{4k}* mutant seedlings developed

similar to wild-type seedlings when germinated in the presence of 150 ppm MMS (**Fig. 9C-D**),

which could be a result of near endogenous expression levels of MLK2 and MLK3 (Ni 2017).

Sensitivity to UV-induced DNA damage was also assessed after periodic exposure to multiple

doses of UV-C irradiation. The phenotypic impact of chronic irradiation with either 2000 or 4000

J m^{-2} on seedlings was evaluated 5 days after a recovery period. Cotyledon cell death and

reduced growth was observed in all genotypes after exposure to both doses of UV-C. However,

tissue chlorosis was only observed in *mlk1/3/4* mutant seedlings after exposure to 2000 and 4000

J m^{-2}. Minimal chlorosis was also observed in the amiR^{4k} mutants after irradiation with 4000 J m^{-}

^{2} (**Fig. 9E**). Taken together, these data suggest that *mlk* mutants have increased sensitivity to

DNA damage.

Discussion

MLK protein kinases alter developmental and stress response pathway protein
phosphorylation

The current repertoire of MLK and MLK homologue substrates is composed of a photoreceptor,

multiple transcription factors, hormone receptors, and histones (Ni et al., 2017; Liu et al., 2017;

Dai and Xue, 2010; Chen et al., 2018; Casas-Mollano et al., 2008; Su et al., 2017). In the absence

of the MLKs, effects on circadian period, hypocotyl elongation, flowering time, osmotic stress

response, seed set, chromatin organization, and hormone sensitivity have been reported (Su et

al., 2017; Casas-Mollano et al., 2008; Liu et al., 2017; Huang et al., 2016; Zheng et al., 2017;

Chen et al., 2018). These observations support a model where MLK family members function as

central regulators of numerous interconnected signaling pathways. Our quantitative analysis of

the global- and phosphoproteomes of *mlk* triple mutant seedlings supports a diverse and complex

role for the MLKs in the regulation of cellular signaling and response. Interestingly, these kinases

seem to share a balance of functional redundancy and substrate specificity which, with respect to

circadian period and hypocotyl elongation, results in opposing phenotypes (Huang et al., 2016;

Liu et al., 2017). We found *mlk4* the *mlk134* mutant displays a much more severe proteomic

phenotype relative to the *mlk123* mutant, with over 10-fold more proteins showing altered

abundance in the *mlk134* mutant (**Fig. 2**). This holds true for the phosphoproteome as well.

However, the difference was greater in tissue sampled in the light. This could be the result of

MLK4 acquiring substrate specificity or may be explained by MLK4 having a greater tendency

than other MLKs for interacting with key light signaling proteins in planta. MLK4 has been shown

to have a higher affinity for PIF3, a known substrate of MLK4, when compared to other MLKs (Ni

et al., 2017) and both phyB and HY5 were found to have altered phosphorylation only at ZT12 in

the *mlk134* mutants (**Table 1**). While some proteins with altered abundance were unique to the

mlk134 or *mlk123* mutants, the common themes of glucosinolate biosynthesis (global proteome;

Fig. 3) and chromosome organization (phosphoproteome; **Fig. 7B-C and Supplemental Dataset**

S2) are seen in both mutants, suggesting a role for the MLK family of kinases in these processes.

Further work is needed to determine if any of the proteins showing altered phosphorylation are

direct substrates of the MLKs, or whether changes in phosphorylation status is occurring indirectly

through additional kinases.

MLKs Regulate Hormone Signaling and Stress Responses

Several proteins responsible for glucosinolate metabolism showed altered abundance in *mlk*

mutant seedlings before and after dark transition (**Fig. 3** and **Supplemental Dataset S1**).

Glucosinolates are nitrogen- and sulfur-containing secondary metabolites known for their role in

plant defense (Kim et al., 2008; Kos et al., 2012; Bednarek et al., 2009; Clay et al., 2009) and for

their anticarcinogenic properties (Higdon et al., 2007). Accumulation of glucosinolates in

Arabidopsis thaliana is a rhythmic process controlled in part by circadian-regulated jasmonate

accumulation that has been linked to the activity of the basic leucine zipper (bZIP) transcription

factor, ELONGATED HYPOCOTYL 5 (HY5) activity (Goodspeed et al., 2012; Huseby et al.,

2013). HY5 is a positive regulator of light signaling and functions as a central regulator of light-

dependent growth and development by integrating various environmental signals (Gangappa and

Botto, 2016). Peak glucosinolate levels occur during the day, which may provide a protective

advantage against rhythmic herbivory. Two glucosinolate biosynthesis genes that showed

increased protein abundance in *mlk* mutants, *CYP79F1* and *SOT18*, showed reduced expression

in the hy5 mutant background (Huseby et al., 2013). The phosphorylation-status of HY5 is

associated with its activity and stability, with the non-phosphorylated form being more active

(Hardtke, 2000). Thus, it is possible that increased HY5 activity, resulting from decreased

phosphorylation in the *mlk* mutant background, could be influencing glucosinolate metabolism.

Additionally, abscisic acid (ABA) is capable of inducing glucosinolate accumulation in plants

(Wang et al., 2015b; Zhu and Assmann, 2017). MLK3 has been shown to regulate ABA signaling

through the phosphorylation of the PYR/PYL ABA receptor family of proteins (Chen et al., 2018).

In agreement with altered ABA signaling, increased phosphorylation of proteins associated with

ABA responses and the SnRK consensus motif, R/K-x-x-S/T, was found to overrepresented in

the *mlk1/3/4* mutant background. Thus, it is possible that the MLKs are involved in the regulation

of defense responses through multiple converging signaling pathways.

The Phosphorylation Status of Key Circadian and Light Signaling Components Are Altered
in the Absence of MLK Family Kinases

Several differentially phosphorylated proteins that are involved in chromatin organization also

function as core circadian clock components (RVE8) or are central regulators of clock input

pathways such as temperature and light signaling (phyB and HY5). RVE8 is a MYB-like

transcription factor that regulates the expression of the clock gene TIMING OF CAB

EXPRESSION1 (TOC1) by promoting histone 3 (H3) acetylation of its promoter (Farinas and Mas,

2011). RVE8 shares structural similarity to the core clock transcription factors CCA1 and LATE

ELONGATED HYPOCOTYL (LHY). Phosphorylation of CCA1 by the Ser/Thr protein kinase CK2

antagonistically regulates CCA1 transcriptional activity by reducing its ability to bind to the

promoters of clock gene targets, which in turn alters circadian period (Portolés and Más, 2010).

Further exploration of the impact of RVE8 phosphorylation could reveal a new avenue of post-

translation regulation of the circadian clock.

Temperature and light signaling are critical circadian inputs that allow plants to coordinate growth

and development (*e.g.* germination and photoperiodic flowering) with their environment. The phyB

photoreceptor is central to both temperature and light signaling pathways (Legris et al., 2016).

PhyB activity is regulated in part by phosphorylation of its N-terminus (Medzihradszky et al., 2013;

Nito et al., 2013). Altered phospho-status of phyB Ser86 and Y10 influences phyB rate dark-

reversion rates, hypocotyl elongation, and flowering time in Arabidopsis (Hajdu et al., 2015;

Medzihradszky et al., 2013; Nito et al., 2013). Here we report decreased phosphorylation of a

previously unidentified phyB N-terminal phosphosite, T42, in the *mlk* mutant background at ZT12

(**Table 2**). MLKs are known to associate with phyB, phosphorylate the phytochrome–interacting

factor, PIF3, and display a variety of red-light dependent growth phenotypes (Ni et al., 2017;

Huang et al., 2016). In addition to the well-established light-induced phyB-PIF signaling cascade,

there is an ample amount of evidence supporting the role of phyB in large-scale chromatin

organization (van Zanten et al., 2010; Tessadori et al., 2009). The MLK-phyB interaction may

contribute to light-dependent chromatin re-organization in addition to regulating PIF3 turnover.

We also found decreased phosphorylation in another key light signaling component, HY5 at T64.

Whether the phosphorylation of $HY5^{T64}$ and/or $phyB^{T42}$ is directly or indirectly influenced by the

MLKs and how those phosphosites fit into the existing light signaling paradigm will be an exciting

line of future research.

The Role of MLK Family Kinases in Modulating Nuclear Architecture

The role of histone modifications in the regulation of developmental processes and stress

response has been well-established, yet our understanding of the responsible modifiers,

modification crosstalk, and targeted genes is incomplete (Probst and Mittelsten Scheid, 2015;

Rosa and Shaw, 2013). Early observations have implicated the MLK family in the regulation of

environmentally-stimulated chromatin organization. MLK1, like its *Chlamydomonas* homologue

MUT9, has been shown to phosphorylate histone H3 on threonine 3 (H3T3p) and to function

redundantly with MLK2 to promote H3T3p in response to salt stress (Wang et al., 2015a; Casas-

Mollano et al., 2008). Accordingly, the *mlk1mlk2* double mutant has abnormal chromatin

organization and an increased sensitivity to osmotic stress (Wang et al., 2015a). Comparisons

have been drawn between the defects in chromosomal organization observed in the *mlk1mlk2*

mutants and those occurring in plants harboring mutations in members of the LITTLE

NUCLEI/CROWDED NUCLEI (LINC/CRWN) gene family, which are involved in controlling

nuclear size and heterochromatin organization (Wang et al., 2013; Sakamoto and Takagi, 2013).

Our analysis of the *mlk* mutant phosphoproteomes found that peptides mapping to multiple

members of the LINC/CRWN family were altered in abundance in *mlk123* and *mlk134* mutants,

suggesting that MLKs may influence nuclear organization in part through regulation of the LINC

proteins.

MLKs Are Involved in May Influence DNA Damage Repair Through Multiple Pathways

Plants are continuously subjected to DNA-damage from their external environment (e.g. ultraviolet

light, ionizing radiation, heat stress, and bacterial and fungal toxins) as well as endogenous

sources such as DNA-alkylating metabolic byproducts. Maintenance of genomic integrity requires

an efficient DNA damage repair (DDR) system that can identify, access, remove and reassemble

damaged genomic regions and requires fluidity within the context of chromatin. Mutations in

genes involved in chromatin organization and remodeling often exhibit defects in DDR and

enhanced susceptibility to DNA damaging reagents (Donà and Mittelsten Scheid, 2015).

Dysregulation of proteins involved in chromatin remodeling such as GAS41/YAFa, ARP4, BRM,

and SYD which we have observed (*vida supra*) is one route by which the increased sensitivity of

mlk mutants to DNA damaging agents (**Fig. 9**) can be explained. The *mlk134* mutant also shows

altered phosphorylation of several proteins directly involved in DDR such as MMH-1 and XRCC4

(Yuan et al., 2014; Roy et al., 2013). Additionally, there is accumulating evidence supporting a

role for small regulatory RNAs in DDR (Hawley et al., 2017); proteins associated with small RNA

metabolism are enriched in *mlk134* at ZT12 (**Fig. 7**). There is no question that full elucidation and

validation of the mechanisms linking MLKs and DDR will require further exploration. However,

taken together, our data suggests the MLKs play an important role in mitigating DNA damage

through regulation of multiple response pathways.

References

Barrero, J.M., Gonzalez-Bayon, R., del Pozo, J.C., Ponce, M.R., and Micol, J.L. (2007). INCURVATA2 Encodes the Catalytic Subunit of DNA Polymerase and Interacts with Genes Involved in Chromatin-Mediated Cellular Memory in Arabidopsis thaliana. PLANT CELL ONLINE.

Bednarek, P., Piślewska-Bednarek, M., Svatoš, A., Schneider, B., Doubský, J., Mansurova, M., Humphry, M., Consonni, C., Panstruga, R., Sanchez-Vallet, A., Molina, A., and Schulze-Lefert, P. (2009). A Glucosinolate Metabolism Pathway in Living Plant Cells Mediates Broad-Spectrum Antifungal Defense. Science (80-.). **323**: 101 LP – 106.

Casas-Mollano, J.A., Jeong, B.-R., Xu, J., Moriyama, H., and Cerutti, H. (2008). The MUT9p kinase phosphorylates histone H3 threonine 3 and is necessary for heritable epigenetic silencing in Chlamydomonas. Proc. Natl. Acad. Sci. U. S. A.

Champion, A., Kreis, M., Mockaitis, K., Picaud, A., and Henry, Y. (2004). Arabidopsis kinome: After the casting. Funct. Integr. Genomics.

Chen, H.-H., Qu, L., Xu, Z.-H., Zhu, J.-K., and Xue, H.-W. (2018). EL1-like Casein Kinases Suppress ABA Signaling and Responses by Phosphorylating and Destabilizing the ABA Receptors PYR/PYLs in Arabidopsis. Mol. Plant **11**: 706–719.

Chou, M.F. and Schwartz, D. (2011). Biological sequence motif discovery using motif-x. Curr Protoc Bioinforma.

Choudhary, M.K., Nomura, Y., Wang, L., Nakagami, H., and Somers, D.E. (2015). Quantitative Circadian Phosphoproteomic Analysis of Arabidopsis Reveals Extensive Clock Control of Key Components in Physiological, Metabolic, and Signaling Pathways. Mol. Cell.

Proteomics.

Clay, N.K., Adio, A.M., Denoux, C., Jander, G., and Ausubel, F.M. (2009). Glucosinolate

metabolites required for an Arabidopsis innate immune response. Science (80-.).

Dai, C. and Xue, H.W. (2010). Rice early flowering1, a CKI, phosphorylates della protein SLR1

to negatively regulate gibberellin signalling. EMBO J.

Donà, M. and Mittelsten Scheid, O. (2015). DNA Damage Repair in the Context of Plant

Chromatin. Plant Physiol.

Ensminger, M., Iloff, L., Ebel, C., Nikolova, T., Kaina, B., and Löbrich, M. (2014). DNA

breaks and chromosomal aberrations arise when replication meets base excision repair. J.

Cell Biol.

Farinas, B. and Mas, P. (2011). Functional implication of the MYB transcription factor

RVE8/LCL5 in the circadian control of histone acetylation. Plant J.

Gangappa, S.N. and Botto, J.F. (2016). The Multifaceted Roles of HY5 in Plant Growth and

Development. Mol. Plant.

Goodspeed, D., Chehab, E.W., Min-Venditti, A., Braam, J., and Covington, M.F. (2012).

Cozzarelli Prize Winner: Arabidopsis synchronizes jasmonate-mediated defense with

insect circadian behavior. Proc. Natl. Acad. Sci.

Hajdu, A., Ádám, É., Sheerin, D.J., Dobos, O., Bernula, P., Hiltbrunner, A., Kozma-Bognár,

L., and Nagy, F. (2015). High-level expression and phosphorylation of phytochrome B

modulates flowering time in Arabidopsis. Plant J.

Hardtke, C.S. (2000). HY5 stability and activity in Arabidopsis is regulated by phosphorylation in

its COP1 binding domain. EMBO J.

Hawley, B.R., Lu, W.-T., Wilczynska, A., and Bushell, M. (2017). The emerging role of RNAs

in DNA damage repair. Cell Death Differ.

Higdon, J. V., Delage, B., Williams, D.E., and Dashwood, R.H. (2007). Cruciferous

vegetables and human cancer risk: epidemiologic evidence and mechanistic basis.

Pharmacol. Res.

Huang, H., Alvarez, S., Bindbeutel, R., Shen, Z., Naldrett, M.J., Evans, B.S., Briggs, S.P.,

Hicks, L.M., Kay, S.A., and Nusinow, D.A. (2016). Identification of evening complex

associated proteins in arabidopsis by affinity purification and mass spectrometry. Mol. Cell.

Proteomics.

Huseby, S., Koprivova, A., Lee, B.R., Saha, S., Mithen, R., Wold, A.B., Bengtsson, G.B.,

and Kopriva, S. (2013). Diurnal and light regulation of sulphur assimilation and

glucosinolate biosynthesis in Arabidopsis. J. Exp. Bot.

Jeong Br, B., Wu-Scharf, D., Zhang, C., and Cerutti, H. (2002). Suppressors of transcriptional

transgenic silencing in Chlamydomonas are sensitive to DNA-damaging agents and

reactivate transposable elements. Proc. Natl. Acad. Sci. U. S. A.

Kim, J.H., Lee, B.W., Schroeder, F.C., and Jander, G. (2008). Identification of indole

glucosinolate breakdown products with antifeedant effects on Myzus persicae (green

peach aphid). Plant J.

Kos, M., Houshyani, B., Achhami, B.B., Wietsma, R., Gols, R., Weldegergis, B.T., Kabouw,

P., Bouwmeester, H.J., Vet, L.E.M., Dicke, M., and van Loon, J.J.A. (2012). Herbivore-

Mediated Effects of Glucosinolates on Different Natural Enemies of a Specialist Aphid. J.

Chem. Ecol.

Legris, M., Klose, C., Burgie, E.S., Rojas, C.C., Neme, M., Hiltbrunner, A., Wigge, P.A.,

Schäfer, E., Vierstra, R.D., and Casal, J.J. (2016). Phytochrome B integrates light and

temperature signals in Arabidopsis. Science (80-.).

Liu, Q. et al. (2017). Molecular basis for blue light-dependent phosphorylation of Arabidopsis

cryptochrome 2. Nat. Commun.

Marondedze, C., Groen, A.J., Thomas, L., Lilley, K.S., and Gehring, C. (2016). A

Quantitative Phosphoproteome Analysis of cGMP-Dependent Cellular Responses in

Arabidopsis thaliana. Mol. Plant.

Medzihradszky, M. et al. (2013). Phosphorylation of Phytochrome B Inhibits Light-Induced

Signaling via Accelerated Dark Reversion in Arabidopsis. Plant Cell.

Ni, W., Xu, S.-L., González-Grandío, E., Chalkley, R.J., Huhmer, A.F.R., Burlingame, A.L.,

Wang, Z.-Y., and Quail, P.H. (2017). PPKs mediate direct signal transfer from

phytochrome photoreceptors to transcription factor PIF3. Nat. Commun.

Nito, K., Wong, C.C.L., Yates, J.R., and Chory, J. (2013). Tyrosine Phosphorylation

Regulates the Activity of Phytochrome Photoreceptors. Cell Rep.

Ohtsubo, T., Matsuda, O., Iba, K., Terashima, I., Sekiguchi, M., and Nakabeppu, Y. (1998).

Molecular cloning of AtMMH, an Arabidopsis thaliana ortholog of the Escherichia coli mutM

gene, and analysis of functional domains of its product. Mol. Gen. Genet.

Portolés, S. and Más, P. (2010). The functional interplay between protein kinase CK2 and cca1

transcriptional activity is essential for clock temperature compensation in Arabidopsis.

PLoS Genet.

Probst, A. V. and Mittelsten Scheid, O. (2015). Stress-induced structural changes in plant

chromatin. Curr. Opin. Plant Biol.

Rademacher, E.H. and Offringa, R. (2012). Evolutionary Adaptations of Plant AGC Kinases:

From Light Signaling to Cell Polarity Regulation. Front. Plant Sci.

Reimand, J., Arak, T., Adler, P., Kolberg, L., Reisberg, S., Peterson, H., and Vilo, J. (2016).

g:Profiler-a web server for functional interpretation of gene lists (2016 update). Nucleic

Acids Res.

Rosa, S. and Shaw, P. (2013). Insights into Chromatin Structure and Dynamics in Plants.

Biology (Basel).

Roy, S., Choudhury, S.R., Sengupta, D.N., and Das, K.P. (2013). Involvement of AtPol in the

Repair of High Salt- and DNA Cross-Linking Agent-Induced Double Strand Breaks in

Arabidopsis. PLANT Physiol.

Sakamoto, Y. and Takagi, S. (2013). LITTLE NUCLEI 1 and 4 regulate nuclear morphology in

arabidopsis thaliana. Plant Cell Physiol.

Schwartz, D. and Gygi, S.P. (2005). An iterative statistical approach to the identification of

protein phosphorylation motifs from large-scale data sets. Nat. Biotechnol.

Silva-Sanchez, C., Li, H., and Chen, S. (2015). Recent advances and challenges in plant

phosphoproteomics. Proteomics.

Sønderby, I.E., Geu-Flores, F., and Halkier, B.A. (2010). Biosynthesis of glucosinolates--gene

discovery and beyond. Trends Plant Sci.

Su, Y., Wang, S., Zhang, F., Zheng, H., Liu, Y., Huang, T., and Ding, Y. (2017).

Phosphorylation of Histone H2A at Serine 95: A Plant-specific Mark Involved in Flowering

Time Regulation and H2A.Z Deposition. Plant Cell.

Sugiyama, N., Nakagami, H., Mochida, K., Daudi, A., Tomita, M., Shirasu, K., and

Ishihama, Y. (2008). Large-scale phosphorylation mapping reveals the extent of tyrosine

phosphorylation in Arabidopsis. Mol. Syst. Biol.

Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011). Revigo summarizes and visualizes

long lists of gene ontology terms. PLoS One.

Suzuki, K., Sako, K., Akiyama, K., Isoda, M., Senoo, C., Nakajo, N., and Sagata, N. (2015).

Identification of non-Ser/Thr-Pro consensus motifs for Cdk1 and their roles in mitotic

regulation of C2H2 zinc finger proteins and Ect2. Sci. Rep.

Tessadori, F. et al. (2009). PHYTOCHROME B and HISTONE DEACETYLASE 6 control light-

induced chromatin compaction in Arabidopsis thaliana. PLoS Genet.

Vergara, Z. and Gutierrez, C. (2017). Emerging roles of chromatin in the maintenance of

genome organization and function in plants. Genome Biol.

Vlastaridis, P., Kyriakidou, P., Chaliotis, A., Van de Peer, Y., Oliver, S.G., and Amoutzias,

G.D. (2017). Estimating the total number of phosphoproteins and phosphorylation sites in

eukaryotic proteomes. Gigascience.

Wang, H., Dittmer, T.A., and Richards, E.J. (2013). Arabidopsis CROWDED NUCLEI (CRWN)

proteins are required for nuclear size control and heterochromatin organization. BMC Plant

Biol.

Wang, Y., Liu, Z., Cheng, H., Gao, T., Pan, Z., Yang, Q., Guo, A., and Xue, Y. (2014). EKPD:

A hierarchical database of eukaryotic protein kinases and protein phosphatases. Nucleic

Acids Res.

Wang, Z., Casas-Mollano, J.A., Xu, J., Riethoven, J.-J.M., Zhang, C., and Cerutti, H.

(2015a). Osmotic stress induces phosphorylation of histone H3 at threonine 3 in

pericentromeric regions of Arabidopsis thaliana. Proc. Natl. Acad. Sci. U. S. A.

Wang, Z., Yang, R., Guo, L., Fang, M., Zhou, Y., and Gu, Z. (2015b). Effects of abscisic acid

on glucosinolate content, isothiocyanate formation and myrosinase activity in cabbage

sprouts. Int. J. Food Sci. Technol.

West, C.E., Waterworth, W.M., Jiang, Q., and Bray, C.M. (2000). Arabidopsis DNA ligase IV

is induced by γ-irradiation and interacts with an Arabidopsis homologue of the double

strand break repair protein XRCC4. Plant J.

van Wijk, K.J., Friso, G., Walther, D., and Schulze, W.X. (2014). Meta-Analysis of

Arabidopsis thaliana Phospho-Proteomics Data Reveals Compartmentalization of

Phosphorylation Motifs. Plant Cell.

Yuan, D., Lai, J., Xu, P., Zhang, S., Zhang, J., Li, C., Wang, Y., Du, J., Liu, Y., and Yang, C.

(2014). AtMMS21 regulates DNA damage response and homologous recombination repair

in Arabidopsis. DNA Repair (Amst).

van Zanten, M., Tessadori, F., McLoughlin, F., Smith, R., Millenaar, F.F., van Driel, R.,

Voesenek, L.A.C.J., Peeters, A.J.M., and Fransz, P. (2010). Photoreceptors

CRYTOCHROME2 and Phytochrome B Control Chromatin Compaction in Arabidopsis.

PLANT Physiol.

Zheng, H., Zhang, F., Wang, S., Su, Y., Jiang, P., Cheng, R., Ji, X., Hou, S., and Ding, Y.

(2017). MLK1 and MLK2 coordinate RGA and CCA1 activity to regulate hypocotyl

elongation in Arabidopsis thaliana. Plant Cell.

Zhu, M. and Assmann, S.M. (2017). Metabolic Signatures in Response to Abscisic Acid (ABA)

Treatment in Brassica napus Guard Cells Revealed by Metabolomics. Sci. Rep.

Figure Legends

Figure 1. Schematic Representation of the Quantitative Proteomics Workflow

Tissue samples were collected at ZT12 and ZT14 from wild type and mutant seedlings entrained with a 12L:12D light/dark cycle (1). Total protein was extracted and digested (2). Following TMT10plex isobaric labeling (3), samples were subjected to high pH reversed phase prefractionation (5). Phosphopeptides were enriched using a TiO2-based method (4). Both phosphopeptide-enriched and global samples were analyzed by LC-MS/MS (6).

Figure 2. Global Proteomic Analysis of *mlk* Mutant Seedlings

Volcano plots of peptides identified in mutant and wild-type seedlings at ZT12 and ZT14. The x-axis specifies the log2 fold-change (FC) of mutant/wild-type and the y-axis specifies the negative logarithm to the base 10 of the t-test p-values. Open circles represent individual peptides, with blue circles specifying those considered statistically significant. Black vertical and horizontal lines reflect the filtering criteria (log2 FC = ±1 and p-value = 0.05) for significance.

Figure 3. Analysis of Quantitative Phosphoproteomics of *mlk* Mutant Seedlings

(A) The distribution of threonine (T), serine (S), and tyrosine (Y) phosphorylation sites identified at ZT12 and ZT14. (B) Volcano plot of phosphopeptides identified in mutant and wild-type seedlings at ZT12 and ZT14. The x-axis specifies the log2 fold-change (FC) of mutant/wild-type and the y-axis specifies the negative logarithm to the base 10 of the t-test p-values. Open circles represent individual peptides, with blue circles specifying those considered statistically significant. Black vertical and horizontal lines reflect the filtering criteria (log2 FC = ±0.585 and p-value = 0.05) for significance. (C and D) Size-proportional Venn diagrams of differentially regulated phosphoproteins in *mlk1/2/3* and *mlk1/3/4* mutants at ZT12 (C) and ZT14 (D). Numbers indicate unique phosphoproteins.

Figure 4. Motif Analysis of Differentially Phosphorylated Peptides

Phosphopeptides with altered abundance in *mlk1/3/4* mutants at ZT12 (A) and ZT14 (B) where extended (http://schwartzlab.uconn.edu/pepextend) and centered. Motif-X analysis was then preformed with the probability threshold was set to p-value ≤ 10^{-6}, the occurrence threshold was set to 20, and the default IPI Arabidopsis Proteome data set was used as the background data set.

Figure 5. Gene Ontology (GO) Enrichment Analysis

GO enrichment analysis of proteins identified as having increased (A) or decreased (B) abundance in *mlk* mutant seedlings at indicated ZTs when compared to wild-type. Cluster representative GO terms identified with REVIGO (semantic similarity threshold < 0.7) in the category of Biological Process are shown.

Figure 6. *mlk* Mutant Seedlings Contain Elevated Levels of MET Derived Glucosinolates

Glucosinolate (GLS) content of 10 day old mutant and wild type seedlings at ZT12 was quantified using HPLC. GLS identity determined using UV spectra and confirmed by LC-MS/MS. The average of 4 biological replicates is presented. Error bars indicate standard deviation. *P<0.01, compared with wild type seedlings (Student's *t*-test).

Figure 7. Gene Ontology Enrichment Analysis of Differentially Phosphorylated Proteins

(A) Heat map showing the p-value significance of enriched cellular component GO categories of proteins with altered abundance in *mlk* mutant seedlings at ZT12 and ZT14. (B and C) Treemap representation of Biological Process GO categories enriched in *mlk1/3/4* mutant seedlings at ZT12 (B) and ZT14 (C). The box size correlates to the –log10 *P*-value of the GO-term enrichment. Boxes with the same color indicate related GO-terms and correspond to the

representative GO-term which is found in the upper-left of each box. REVIGO was used to

eliminate redundant GO-terms with a dispensability value ≥ 0.7 (A) or ≥ 0.5 (B and C).

Figure 8. Analysis of Differentially Phosphorylated Peptides in *mlk1/3/4* Mutants at ZT12

(A) Size-proportional Venn diagram of proteins which show increased, decreased, or both

increased and decreased abundance of identified phosphosites in *mlk1/3/4* mutant seedlings

at ZT12. Numbers indicate unique phosphoproteins. Analysis of phosphoproteins that show

increased (B) or decreased (C) in abundance in the *mlk1/3/4* mutant seedlings background at

ZT12. Treemap representations of Biological Process GO category enrichment are shown. The

box size correlates to the –log10 *P*-value of the GO-term enrichment. Boxes with the same color

indicate related GO-terms and correspond to the representative GO-term which is found in the

upper-left of each box. REVIGO was used to eliminate redundant GO-terms with a dispensability

value ≥ 0.5.

Figure 9. *mlk1/3/4* Mutants Have Increased Sensitivity to MMS Treatment

(A) Representative images of mutant and wild type seedlings 15 days after transfer to solid

media containing the indicated concentration of MMS. Chlorotic tissue is indicated with a blue

arrowhead. Scale bar = 5 mm. (B) Fresh weight of 21-day old wild type and mutant seedlings

grown in the presence of MMS relative to mock treated samples. The average of 3 biological

replicates of ≥10 seedlings each is presented. (C) Percent of seedlings exhibiting post-

germination developmental arrest after 12 days of growth in the presence of 150 PMM MMS.

The average of 3 biological replicates of ≥30 seedlings is presented. (B and C) Error bars

indicate standard deviation. *$P<0.05$, **$P<0.01$ compared with wild type seedlings (Student's *t*-

test). (D) Representative images of mutant and wild type seedlings germinated in the presence

of 150 PPM MMS. (E) Representative images of mutant and wild type seedlings irradiated with

indicated levels of UV-C. Scale bar = 2 mm.

Table 1. Phosphosites Identified in Circadian-Associated Proteins.

Protein Group Accession	Protein Names	Protein Description	Peptide Sequence	Log2 FC	P-Value
AT2G18790	PHYB	phytochrome B	GGEQAQSSGT^{42}K	-1.36	5.95E-06
AT4G30200	VEL1	vernalization5/VIN3-like protein	KPS^{506}SKNEDNNSPSVDESAAK	-0.89	3.00E-02
AT5G11260	HY5	Basic-leucine zipper (bZIP) transcription factor family protein	ESGSAT^{64}GQER	-0.81	1.46E-02
AT3G22380	TIC	time for coffee	MPS^{324}TSKQEAAGNDLTEAAK	-0.79	1.98E-04
AT5G02810	PRR7	pseudo-response regulator 7	QDNS^{355}FEK	-0.74	1.04E-05
AT2G17840	ERD7	Senescence/dehydration-associated protein-like protein	SAAS^{488}QKK	-0.69	3.95E-02
AT5G02810	PRR7	pseudo-response regulator 7	AVS^{275}LWDR	-0.62	1.93E-03
AT4G30200	VEL1	vernalization5/VIN3-like protein	LCSSALESLETIAT^{330}TPPDVAALPS^{340}PR	0.61	1.53E-02
AT4G30200	VEL1	vernalization5/VIN3-like protein	NEDNNSPS^{516}VDESAAK	0.63	1.01E-04
AT5G52310	LTI78	low-temperature-responsive protein 78 (LTI78) / desiccation-responsive protein 29A (RD29A)	NEYSPES^{387}DGGLGAPLGGNFPVR	0.64	6.53E-03
AT3G09600	RVE8	Homeodomain-like superfamily protein	GLLNVSSPSTSGMGSSSR	0.83	1.72E-03
AT3G46780	PTAC16	plastid transcriptionally active 16	ADAVGVT^{410}VDGLFNK	0.95	9.29E-03

Figure 1. Schematic Representation of the Quantitative Proteomics Workflow

Tissue samples were collected at ZT12 and ZT14 from wild type and mutant seedlings entrained with at 12L:12D light cycle (1). Total protein was extracted and digested (2). Following TMT10plex isobaric labeling (3), samples were subjected to high pH reversed phase prefractionation (5). Phosphopeptides were enriched using a TiO2-based method (4). Both phosphopeptide-enriched and global samples were analyzed by LC-MS/MS (6).

Figure 2. Global Proteomic Analysis of *mlk* Mutant Seedlings

Volcano plots of peptides identified in mutant and wild-type seedlings at ZT12 and ZT14. The x-axis specifies the log2 fold-change (FC) of mutant/wild-type and the y-axis specifies the negative logarithm to the base 10 of the t-test p-values. Open circles represent individual peptides, with blue circles specifying those considered statistically significant. Black vertical and horizontal lines reflect the filtering criteria (log2 FC = ±1 and p-value = 0.05) for significance.

Figure 3. Analysis of Quantitative Phosphoproteomics of *mlk* Mutant Seedlings

(A) The distribution of threonine (T), serine (S), and tyrosine (Y) phosphorylation sites identified at ZT12 and ZT14. (B) Volcano plot of phosphopeptides identified in mutant and wild-type seedlings at ZT12 and ZT14. The x-axis specifies the log2 fold-change (FC) of mutant/wild-type and the y-axis specifies the negative logarithm to the base 10 of the t-test p-values. Open circles represent individual peptides, with blue circles specifying those considered statistically significant. Black vertical and horizontal lines reflect the filtering criteria (log2 FC = ±0.585 and p-value = 0.05) for significance. (C and D) Size-proportional Venn diagrams of differentially regulated phosphoproteins in *mlk1/2/3* and *mlk1/3/4* mutants at ZT12 (C) and ZT14 (D). Numbers indicate unique phosphoproteins.

A.

	#	Motif	Motif Score	Foreground Matches	Foreground Size	Background Matches	Background Size	Fold Increase
Decreased	1	.......S..K....	10.36	38	188	60894	1013104	3.36
	2	....R..S.......	5.09	23	150	52253	952210	2.79
	3	....K..S.......	5.02	22	127	55173	899957	2.83
Increased	1	.......SP......	10.53	42	250	53111	1013104	3.2
	2	.......S.S.....	6.52	51	208	113600	959993	2.07

B.

	#	Motif	Motif Score	Foreground Matches	Foreground Size	Background Matches	Background Size	Fold Increase
Decreased	1	.......S.....E.	6.15	27	149	63949	1013104	2.87
	2	....K..S.......	5.12	21	122	55198	949155	2.96
Increased	1	.......SP......	11.64	25	88	53111	1013104	5.42

Figure 4. Motif Analysis of Differentially Phosphorylated Peptides

Phosphopeptides with altered abundance in *mlk1/3/4* mutants at ZT12 (A) and ZT14 (B) where extended (http://schwartzlab.uconn.edu/pepextend) and centered. Motif-X analysis was then preformed with the probability threshold was set to p-value ≤ 10^{-6}, the occurrence threshold was set to 20, and the default IPI Arabidopsis Proteome data set was used as the background data set.

Figure 5. Gene Ontology (GO) Enrichment Analysis

GO enrichment analysis of proteins identified as having increased (A) or decreased (B) abundance in *mlk* mutant seedlings at indicated ZTs when compared to wild-type. Cluster representative GO terms identified with REVIGO (semantic similarity threshold < 0.7) in the category of Biological Process are shown.

Figure 6. *mlk* **Mutant Seedlings Contain Elevated Levels of MET Derived Glucosinolates**

Glucosinolate (GLS) content of 10 day old mutant and wild type seedlings at ZT12 was quantified using HPLC. GLS identity determined using UV spectra and confirmed by LC-MS/MS. The average of 4 biological replicates is presented. Error bars indicate standard deviation. *$P<0.01$, compared with wild type seedlings (Student's *t*-test).

Figure 7. Gene Ontology Enrichment Analysis of Differentially Phosphorylated Proteins

(A) Heat map showing the p-value significance of enriched cellular component GO categories of proteins with altered abundance in *mlk* mutant seedlings at ZT12 and ZT14. (B and C) Treemap representation of Biological Process GO categories enriched in *mlk1/3/4* mutant seedlings at ZT12 (B) and ZT14 (C). The box size correlates to the –log10 *P*-value of the GO-term enrichment. Boxes with the same color indicate related GO-terms and correspond to the representative GO-term which is found in the upper-left of each box. REVIGO was used to eliminate redundant GO-terms with a dispensability value ≥ 0.7 (A) or ≥ 0.5 (B and C).

Figure 8. Analysis of Differentially Phosphorylated Peptides in *mlk1/3/4* Mutants at ZT12

(A) Size-proportional Venn diagram of proteins which show increased, decreased, or both increased and decreased abundance of identified phosphosites in *mlk1/3/4* mutant seedlings at ZT12. Numbers indicate unique phosphoproteins. Analysis of phosphoproteins that show increased (B) or decreased (C) in abundance in the *mlk1/3/4* mutant seedlings background at ZT12. Treemap representations of Biological Process GO category enrichment are shown. The box size correlates to the –log10 *P*-value of the GO-term enrichment. Boxes with the same color indicate related GO-terms and correspond to the representative GO-term which is found in the upper-left of each box. REVIGO was used to eliminate redundant GO-terms with a dispensability value ≥ 0.5.

Figure 9. *mlk1/3/4* Mutants Have Increased Sensitivity to MMS Treatment

(A) Representative images of mutant and wild type seedlings 15 days after transfer to solid media containing the indicated concentration of MMS. Chlorotic tissue is indicated with a blue arrowhead. Scale bar = 5 mm. (B) Fresh weight of 21-day old wild type and mutant seedlings grown in the presence of MMS relative to mock treated samples. The average of 3 biological replicates of ≥10 seedlings each is presented. (C) Percent of seedlings exhibiting post-germination developmental arrest after 12 days of growth in the presence of 150 PMM MMS. The average of 3 biological replicates of ≥30 seedlings is presented. (B and C) Error bars indicate standard deviation. *P<0.05, **P<0.01compared with wild type seedlings (Student's *t*-test). (D) Representative images of mutant and wild type seedlings germinated in the presence of 150 PPM MMS. (E) Representative images of mutant and wild type seedlings irradiated with indicated levels of UV-C. Scale bar = 2 mm.