Emergence and diversification of a highly invasive chestnut pathogen lineage across south-eastern Europe

Lea Stauber[1,2], Simone Prospero[1,*], Daniel Croll[2,*]

[1] Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland

[2] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel

[*] Corresponding authors: simone.prospero@wsl.ch, daniel.croll@unine.ch

Data availability: All raw sequencing data is available on the NCBI Short Read Archive (BioProject PRJNA604575)

Abstract

Invasive microbial species constitute a major threat to biodiversity, agricultural production and human

health. Invasions are often dominated by one or a small number of genotypes, yet the underlying

factors driving invasions are poorly understood. A prime example for a successful global invasion is

the recent outbreak of the chestnut blight fungus *Cryphonectria parasitica*. Native to East Asia, the

pathogen colonized North America and Europe during the first half of the 20th century. After

decimation of the American chestnut, the pathogen threatens European chestnut trees. To unravel the

mechanisms underpinning the most recent invasion of south-eastern Europe over the past decades, we

sequenced genomes of 188 predominantly European strains. Genotypes outside of the invasion zone

showed high levels of diversity with evidence for frequent and ongoing recombination. The invasive

lineage emerged from the highly diverse European genotype pool rather than a secondary introduction

from Asia. The expansion across south-eastern Europe was mostly clonal and shows distinct signs of

mutation accumulation. The lineage is also dominated by a single mating type suggesting a fitness

advantage to switch to asexual reproduction. However, we show experimentally that the lineage

retained the ability to undergo mating consistent with the low degree of recombination detected

among strains within the lineage and possibly closely related strains. Our findings show how an

intermediary, highly diverse bridgehead population gave rise to an invasive, largely clonally

expanding pathogen.

Introduction

Over the past century, a multitude of invasive species have emerged as threat to forest and agricultural ecosystems worldwide (Wingfield et al., 2010; Santini et al., 2013). Increased human activities, such as global trade, enabled invasive species to cause economic damage through reduced agricultural production, degradation of ecosystems and negative impacts on human health (Marbuah et al., 2014). A key group of invasive species in forests are fungal pathogens, which are often accidentally spread via living plants and plant products (Rossman, 2001). To successfully colonize a new environment, fungal pathogens have to overcome several invasion barriers including effective dispersal abilities, changes in available hosts, competition with other fungi, and niche availability (Gladieux et al., 2015). This may be achieved with plastic phenotypic changes followed by rapid genetic adaptation (Garbelotto et al. 2015). However, because the number of initial founders is often low, invasive populations are frequently of low genetic diversity which reduces adaptive genetic variation (Allendorf et al. 2003, Yang et al. 2012). Yet, many fungal plant pathogen invasions were successful despite low genetic diversity within founding populations (Raboin et al., 2007, Fontaine et al., 2013, Wuest et al., 2017).

A major hypothesis explaining the successful expansion of invasive populations despite low initial genetic diversity is the so-called bridgehead effect. In this model, highly adapted lineages emerge through recombination among genotypes established in an area of first introduction (Lombaert et al., 2010). Hence, the primary introduction serves as the bridgehead for a secondary and more expansive invasion. Although the bridgehead effect has been proposed as a scenario for many biological invasions (Gau et al. 2013, van Boheem et al. 2017), empirical evidence for the creation of highly adaptive genotypes within bridgehead populations is still largely missing (Bertelsmeier & Keller 2018). An alternative explanation suggests that first introductions simply serve as repeated sources of inoculum for additional secondary invasions without creating adaptive genotypes *in situ* (Bertelsmeier & Keller, 2018). This alternative scenario implies that initial populations are already composed of genotypes adapted to the new environment or have high phenotypic plasticity (Bock et

al., 2015, Gladieux et al., 2015, Vukovic et al. 2019). Dissecting whether invasive species were pre-

adapted to the new environment or gained adaptation through a bridgehead effect is crucial for

effective containment strategies, but requires a deep sampling of genotypes during the early invasion

process.

The pace of adaptive evolution leading to successful invasions is determined by several life-history

traits including the mating system (*i.e.* sexual, asexual or mixed). The low availability of mating

partners can be a major cost for individuals at the invasion front. To avoid mating costs, invasive

fungal pathogens often switch from sexual to predominantly asexual reproduction (Heitman et al.,

2013, Suehs et al., 2004). However, the switch to asexual progeny reproduction limits genetic

diversification and adaptive potential (Taylor et al. 2015, Drenth et al. 2019). Populations of some

highly successful invaders, such as the ascomycete *Ophiostoma novo-ulmi* (Paoletti et al., 2006), or

the oomycete *Phytophthora ramorum* (Grünwald et al., 2012), are dominated by a single mating type.

Low diversity in invasive lineages may be breached by secondary invasions introducing the opposite

mating type as observed in *Phytophthora infestans* in Europe. Prior to the 1980s only the A1 mating

type was present followed by the secondary introduction of the A2 mating type (Mariette et al.,

2016). Introgression from closely related species can also reintroduce a missing mating type. *O.*

novo-ulmi acquired the missing mating type from *O. ulmi* (Brasier & Webber, 2019, Paoletti et al.,

2006). Although switches in reproductive modes can be a key factor for invasion success (Philibert et

al., 2011), mechanisms underlying such switches remain poorly understood (Billiard et al., 2012).

The ascomycete *Cryphonectria parasitica* (Murr.) Barr. is the causal agent of chestnut blight, a lethal

bark disease of *Castanea* species (Rigling & Prospero 2018). The pathogen is native to eastern Asia

and was first observed in 1904 in North America on the American chestnut (*Castanea dentata*). In

the following years, the disease rapidly spread throughout the distribution range of *C. dentata*,

causing the vast decimation of this native tree species (Elliott & Swank 2008). In 1938, the fungus

was first detected on the European chestnut (*C. sativa*) near Genoa (Italy) and is now established in

all major chestnut-growing countries in Europe (Rigling & Prospero 2018). The damage to the

European chestnut may have been reduced by the presence of the *Cryphonectria hypovirus 1* (CHV-1) which acts as a biological control agent of chestnut blight (Rigling & Prospero 2018). The virus can be transmitted both vertically to asexual spores (conidia) or horizontally through hyphal anastomoses between virus-infected and virus-free strains (Heiniger and Rigling 1994). Hyphal anastomoses are controlled by a vegetative compatibility system and the virus spreads most effectively between strains of the same vegetative compatibility type (Cortesi et al. 2001).

Population genetic analyses showed that the invasion of Europe occurred through multiple introductions both from native populations in Asia and from bridgehead populations in North America (Dutech et al. 2012). Invasive European *C. parasitica* populations are characterized by lower vegetative compatibility diversity than North American and Asian populations (Milgroom & Cortesi 1999). Furthermore, European populations exhibit lower recombination rates (Dutech et al., 2010; González-Varela et al., 2011; Prospero & Rigling, 2012). Mating in *C. parasitica* is controlled by a single mating type (MAT) locus (Marra and Milgroom, 2001) and natural populations can reproduce both sexually and asexually (Marra et al., 2004). Previous analyses of south-eastern European *C. parasitica* populations based on sequence characterized amplified region (SCAR) markers suggested that the region was largely colonized by a single and likely asexual lineage also identified as S12 (Milgroom et al. 2008). The lineage belongs to the vegetative compatibility type EU-12. Within the distribution range of the lineage, sexual structures (*i.e.* perithecia) are rarely found (Sotirovski et al. 2004, Milgroom et al., 2008). Based on SCAR marker and field records, Milgroom et al. (2008) suggested that the invasive S12 lineage originated in northern Italy and a subsequently spread across south-eastern Europe (Avolio, 1978, Biraghi, 1946, Buccianti & Feliciani 1966, Karadzic et al., 2019, Myteberi et al., 2013, Robin & Heiniger, 2001). However, due to the low molecular marker resolution and challenges in relying on observational data, the origin, invasion route and genetic diversification of *C. parasitica* in south-eastern Europe remain largely unknown.

In this study, we sequenced complete genomes of a comprehensive collection of European *C. parasitica* strains with a fine-scaled sampling throughout the S12 invasion zone. Based on high-

120 confidence genome-wide single nucleotide polymorphisms (SNPs), we identified the most likely

121 origin and recapitulated the invasion process across south-eastern Europe. We show that the invasive

122 *C. parasitica* lineage arose through an intermediary, highly diverse bridgehead population. During

123 the expansion, the lineage became dominated by a single mating type but retained the ability to

124 reproduce sexually.

126
127 **Results**
128

Genome-wide polymorphism analyses and phylogenomic reconstruction

We analyzed complete genomes of 188 *C. parasitica* isolates covering the European outbreak region

of the invasive S12 lineage, as well as reference isolates from South Korea ($n = 2$) and the United

States ($n = 4$). Isolates were sequenced at a mean depth of 8-27.5X to detect high-confidence

genome-wide SNPs. A region of 179'501 -2'084'312 bp on scaffold 2 was associated to the mating

type locus based on association mapping *p*-values (Supp. Fig. 2). This region encoded known mating

type associated genes (Idnurm et al. 2015) and was characterized by a high SNP density consistent

with observations in other fungi (Taylor et al. 2015). We removed SNPs within the mating type

associated region to avoid confounding genetic structure with mating type divergence. We retained

17'873 SNPs and constructed a whole-genome maximum likelihood phylogenetic tree (Fig. 1A). The

tree revealed two major clades splitting two Asian and one Swiss isolate (vegetative compatibility

type EU-65) from the European and North American isolates. The large European/North American

clade was subdivided into three clades. The clades a and c showed mixed geographic origins

including North American isolates and European isolates of various vegetative compatibility (EU-

)types (Figure 1A, Supp. Fig. 3). In contrast, clade b consisted of genetically highly similar isolates

($n = 113$; Figure 1A) of vegetative compatibility type EU-12, which predominantly originated from

the Balkans, Italy, Turkey and Georgia. Approximately 93% ($n = 105$) of all isolates in the EU-12

clade were MAT-1 (Figure 1A) and were found in Albania, Bosnia, Bulgaria, Croatia, Georgia,

Greece, Italy, Macedonia, Serbia and Turkey (Figure 1B; Supp. Table 1).

Figure 1: **Genome-wide analyses of *Cryphonectria parasitica* isolates focussing on the invasion in south-eastern Europe**. A) Whole-genome single nucleotide polymorphism (SNP) maximum likelihood tree constructed from 188 sequenced *C. parasitica* isolates. The US/European clade is shown with a grey box. The pie charts represent the proportions of vegetative compatibility types within each of the three clades. The most frequent vegetative compatibility types are highlighted in colour (the colour scheme matches with Figure 2) . B) Map of European *C. parasitica* sampling locations. Grey circles show sampling locations of various vegetative compatibility types. Purple coloured hexagons represent sampling locations where EU-12 (mating type MAT-1) isolates were found (marked as "S12 lineage"). First observations of chestnut blight in the corresponding countries and regions are marked in a colour scheme according to decade. A tree with individual isolate labelling is shown in Supp. Fig. 3.

We performed a SplitsTree phylogenetic network analysis to account for reticulation caused by

recombination. The network showed a high diversification, with both long branching and reticulation

(Figure 2, Supp. Fig. 4). The PHI-test revealed significant evidence for recombination ($p < 0.0001$).

Despite the high level of genetic diversity, we found no evidence for geographic structure. Moreover,

we found no clustering of isolates belonging to the same vegetative compatibility type with the

exception of some EU-01, EU-02 and EU-12 (S12) genotypes from the Balkans. Nearly all *C.*

parasitica isolates representing the S12 lineage showed almost identical genotypes and tight

clustering. All most tightly clustered S12 genotypes were of mating type MAT-1 ($n = 104$).

Consistent with analyses by Milgroom et al. (2008), this group represents the invasive S12 lineage at

the origin of the expansion of *C. parasitica* across south-eastern Europe. Additionally, the

phylogenetic network revealed closely related but not identical S12 genotypes of mating type MAT-

2 ($n = 7$, Figure 2). Hence, S12 outbreak strains of MAT-2 connect the nearly uniform cluster of S12

MAT-1 strains with the remaining genetic diversity of the major European subgroup of *C. parasitica*.

The S12 cluster was furthermore connected with the remaining genotypes of the major clade by two

EU-12 isolates from Bosnia (M1808 with MAT-1) and Georgia (MAK23 with MAT-2).

Figure 2: **Phylogenetic network structure of the US/European subgroup analyzed by SplitsTree**.
The highlighted branches represent the most abundant vegetative compatibility types (colour scheme

matching Figure 1A). Isolates belonging to the S12 outbreak lineage (EU-12; mating type MAT-1, n

= 104) are marked with a purple hexagon. S12 isolates of mating type MAT-2 are highlighted in light-

purple. Additional EU-12 isolates not belonging to the S12 lineage are highlighted in blue with

information on the country of origin. Genetic donors of the S12 lineage as inferred by

fineSTRUCTURE (Figure 3, Table 1, Supp. Fig. 5) are marked with red squares.

Potential S12 founder populations

The maximum likelihood phylogenetic tree and the SplitsTree network revealed that the invasive S12

lineage has closely related genotypes occurring in Europe. Thus, to dissect the genetic origin of S12,

we performed a co-ancestry matrix analysis using fineSTRUCTURE considering all isolates of the

major US/European subgroup, including the S12 lineage (n = 185; Figure 1A). The averaged co-

ancestry matrix revealed no direct ancestors of the invasive S12 lineage among the major clade of

different vegetative compatibility types. However, we found an association with a coefficient of 24.9

– 32.3 between the recipient S12 genotypes and European donors from different locations (Figure 3,

Table 1, Supp. Fig. 5). The donors mostly originated from the North Balkans (Bosnia and Croatia), as

well as Southern Switzerland, with the exception of isolate MAK23 from Georgia.

Figure 3: **Analysis of donors to the S12 lineage genotypes**. The averaged co-ancestry matrix and
phylogenetic tree of the US/European subgroup estimated by fineSTRUCTURE. The matrix shows
recipient genotypes as rows and donors as columns. The heatmap indicates averaged coancestry
coefficients between donors and recipients (i.e. darker colours indicate stronger genetic relationship
between donors and recipients). Contributing S12 donors are marked in grey-dashed boxes. A detailed
list of contributing donors is shown in Table 1, Figure 2 and Supp. Fig. 5.

Potential S12 invasion routes and retention of mating competence

To infer potential invasion routes of the S12 outbreak lineage, we investigated intra-lineage genetic

diversity across south-eastern Europe. We focused only on S12 isolates of mating type MAT-1 to

delimit the closest genotypes contributing to the outbreak (n = 104; Figure 2). The closely related

genotypes segregated 468 high-confidence SNPs across the genome. The genetic structure assessed

by a principal component analysis showed loose clustering of genotypes across south-eastern Europe

(Figure 4A and B, Supp. Fig. 6). We assigned genotypes to five regions: Italy, Northern Balkans,

Central Balkans, Greece/Turkey and Georgia (Figure 4A). Italy, Northern and Central Balkans, as

well as Georgia harbored mainly genotypes of two dominant clusters. In contrast, the Greece/Turkey

region contained genotypes of the two dominant clusters but also a broad diversity of further

genotypes. We analyzed evidence for reticulation in the phylogenetic relationships among genotypes

but found a star-like structure. We found minor evidence for reticulation among Central Balkans

genotypes (Figure 4C). Consistent with the phylogenetic network pattern, we found significant

evidence for recombination within the S12 lineage (PHI test; $p = 0.0035$). We tested experimentally

whether S12 mating type MAT-1 isolates were still able to reproduce sexually. We confirmed

outcrossing of isolates of opposite mating type within the S12 lineage by pairing isolates from Molliq

(Kosovo) and Nebrodi (southern Italy) (Fig. 4D). Mating pairs from Molliq and Nebrodi grew

numerous perithecia, which are the fruiting bodies specific to sexual reproduction (Fig. 4E). Pairings

of Bosnian isolates showed no perithecia formation. Using molecular mating type assays, we

recovered both mating types among the ascospores produced from successful matings .

Figure 4: Fine-scale genetic diversity analyses of the S12 lineage. A) Geographic map showing the collected S12 isolates of *Cryphonectria parasitica* (n=104) according to five geographic regions. B) Principal component analysis (PCA) and C) SplitsTree of the S12 outbreak isolates. Symbols and colors are as in A). D) Scheme of successful mating pairs of S12 mating type MAT-1 isolates crossed with isolates from the opposite mating type, of the same geographic origin. Symbols are as in C). E)

Photographic images of sexual *C. parasitica* fruiting bodies (*i.e.* perithecia) emerging from crosses of
S12 mating type MAT-1 isolates with isolates of the opposite mating type after five months of
incubation under controlled conditions. Left: Perithecia embedded in a yellow-orange stromatic
tissue. Right: Cross-section of perithecia and chestnut bark. Flask-shaped structures with a long
cylindrical neck develop in yellow–orange stromatic tissue and are embedded in the bark (except for
the upper part). The ascospores are formed in sac-like structures (asci) in the basal part of the
perithecium. When mature, the ascospores are actively ejected into the air through a small opening
(ostiole) at the end of the perithecial neck.
243
244

Polymorphism and allele frequency spectra within the outbreak lineage

To gain insights into evolutionary forces shaping polymorphism in the outbreak S12 mating type

MAT-1 lineage versus non-S12 populations, we first analyzed allele frequencies across the genome

in both groups (Figure 5A). The S12 lineage segregated virtually no intermediate allele frequencies in

the range of 0.05-0.95. In contrast, the non-S12 genotypes showed overall a wide spectrum of allele

frequencies across the genome. Second, we analyzed the predicted impact on protein functions in the

S12 lineage and non-S12 populations (Figure 5B). We found 4 highly and 94 moderately deleterious

SNPs within S12 in contrast to 29 high and 773 moderately deleterious mutations in non-S12 groups

(Figure 5B). Three of the high impact SNPs in the S12 lineage were classified as stop gain mutations,

as well as one splice acceptor variant (insertion variant). Two of these high impact mutations affect

proteins of the major facilitator superfamily, as well as a protein containing a LCCL domain and an

ecdysteroid kinase. Non-S12 populations showed an over-representation of low frequency high-

impact mutations (Figure 5B). This is consistent with purifying selection reducing the frequency of

these mutations due to fitness costs. Within the S12 lineage nearly all segregating mutations were at

very low frequency. We found only modifier (*i.e.* nearly neutral) mutations rising to higher frequency

within the lineage.

Figure 5: **Polymorphism segregating in the S12 lineage**. A) Reference allele frequencies across the genome for the S12 lineage ($n = 104$) compared to all other analyzed European samples (non-S12; $n = 80$). M: mitochondria. B) Allele frequency spectra of high, moderate and modifier (i.e. near neutral) impact mutations as identified by SnpEff. The plots are shown both for the S12 outbreak lineage as well as the pool of all other analyzed European samples.

Discussion

We analyzed at the genome level how the chestnut blight pathogen *C. parasitica* successfully invaded south-eastern Europe after a first establishment on the continent. Deep sampling of the invasive genotypes showed that the invasion was caused by a single, highly homogeneous lineage consisting nearly exclusively of a single mating type. The invasive lineage likely transitioned recently from biparental mating populations to a single mating type. The spread across south-eastern Europe,

suspected to originate in northern Italy, left no clear geographic imprint. This lack of genetic

differentiation along the invasion route may be due to high levels of gene flow after the initial

colonization or a very rapid spread from the center of origin to multiple locations across south-

eastern Europe. During the invasion, sexual reproduction was likely ongoing sporadically and may

have introgressed genetic material from outside of the invasive lineage.

The establishment of a European bridgehead population

Central and south-eastern European *C. parasitica* populations likely originated from North American

lineages. Clusters of European and North American genotypes were largely overlapping indicating

significant gene flow and, possibly, multiple introductions to Europe. These genome-scale analyses

are consistent with previous findings documenting multiple North American introductions into

France, but also directly from Asia (Milgroom et al. 1996, Dutech et al. 2012, Demené et al. 2019).

Our results suggest that central and south-eastern European *C. parasitica* populations were largely

established from North American sources alone. The high diversity at the genetic and vegetative

compatibility type level suggests that Europe was colonized repeatedly over the past century.

Although genetic diversity could also have accumulated *in situ* in Europe, the establishment of a

large set of genotypes and vegetative compatibility types seems difficult to explain with population

age alone. Sexual recombination between the three most common vegetative compatibility types in

Europe (*i.e.* EU-01, EU-02 and EU-05; Robin & Heiniger 2001) could not account for the observed

vegetative compatibility type diversity. Observational records date the first introductions into Europe

to the 1930s (Robin & Heiniger, 2001). Hence, based on the observed genetic diversity, a scenario of

repeated introductions of different vegetative compatibility types since 1930s seems most plausible.

Populations from Southern Switzerland, Slovenia Croatia and Bosnia are highly diverse, have nearly

balanced mating type ratios, and a lack clustering according to vegetative compatibility types. This

strongly suggests frequent outcrossing and population admixture consistent with reports of perithecia

in the field (Ježić et al. 2012, Prospero et al. 2006, Prospero & Rigling, 2012, Trestic et al. 2001).

Low vegetative compatibility type diversity in most European *C. parasitica* populations was thought

to have contributed to low population admixture within Europe compared to Asia and North America

(Cortesi et al. 1996, Dutech et al. 2012, Prospero & Rigling, 2012). However, our genome-wide

analyses revealed frequent and ongoing *in situ* admixture in Europe. Thus, vegetative compatibility

type diversity does not necessarily underpin population admixture frequency and genetic diversity in

sexually recombining populations. Our findings show that in asexually reproducing populations, such

as in the S12 lineage, genotypes tend to cluster according to vegetative compatibility types.

Emergence of an invasive lineage from a European bridgehead

The invasive lineage S12 most likely arose from existing genotypes established in Europe. We

identified a series of closely related genotypes to the S12 lineage in Bosnia, Croatia, Georgia and

southern Switzerland. Strikingly, the closest genotypes to the dominant S12 MAT-1 were S12 MAT-

2 isolates found in Bosnia, Kosovo and southern Italy. Analyses based on a coancestry matrix

identified a group of more distantly related genotypes from Bosnia, Croatia, Switzerland and Georgia

having made the strongest genetic contributions to the S12 lineage. This shows that introductions

from outside of Europe are unlikely to explain the emergence of S12. Furthermore, the emergence of

S12 was accompanied by a striking evolutionary transition from mixed mating type populations to

single mating type outbreak populations. Human activity may have contributed to the shift towards

single mating type populations. Shipments of infected chestnut seedlings from Northern Italy and

other trading activities could have disseminated the invasive lineage further South. This would have

exposed the pathogen to the geographically more fragmented chestnut forests typically found in

south-eastern Europe where asexuality or selfing may be advantageous. Although *C. parasitica* is

able to produce asexual conidia in large quantities, these specific spores are thought to be mainly

splash dispersed by rain over short distances (Griffin, 1986). Accounting for occasional dispersal by

birds or insects (Heald & Studhalter, 1914), conidia dispersal is unlikely to contribute substantially to

the colonization of new areas.

Despite the loss of a mating type in the S12 lineage, we found genome-wide evidence for reticulation

indicating at least low levels of recombination. If mating in *C. parasitica* follows the canonical

process found in many ascomycetes, isolates of opposite mating type are required. Hence, S12

isolates of mating type MAT-1 may sporadically mate with rare S12 isolates of mating type MAT-2,

which are comparatively more diverse. The emergence of the opposite mating type at low frequency

could be the result of recombination with other genotypes and subsequent backcrossing. Combined

with experimental evidence, we show that the dominant S12 mating type MAT-1 has retained the

ability for sexual reproduction. Furthermore, in Bosnia, Croatia, Italy (Sicily) and Turkey the S12

lineage co-exists with other genotypes (*i.e.* vegetative compatibility types EU-01 and EU-02) of both

mating types, potentially enabling sexual recombination and diversification *in situ*. The invasive S12

lineage was likely pre-adapted to the south-eastern European niche as we traced the origins to a likely

Italian bridgehead population. Niche availability and benefits associated with asexual reproduction to

colonize new areas may have pre-disposed the European *C. parasitica* bridgehead population to

produce a highly invasive lineage.

Expansion and mutation accumulation within the invasive S12 lineage

The MAT-1 S12 lineage diversified largely through mutation accumulation as nearly all high-

confidence SNPs were identified as singletons. Mutation accumulation in absence of substantial

recombination resulted in star-like phylogenetic relationships. Analyses of allele frequency spectra

suggested that the broader European *C. parasitica* populations efficiently removed the most

deleterious mutations through purifying selection. In contrast, the S12 lineage shows strong skews

towards very low minor allele frequencies of all mutation categories. Interestingly, we found a

broader spread in allele frequencies for nearly neutral mutations in the S12 lineages. This suggests

that despite the largely clonal population structure, deleterious mutations can still be removed

through low levels of recombination and purifying selection. Using accumulated mutations as

markers to retrace the spatial expansion of the invasive S12 lineage, we found no indication for a

step-wise geographic expansion along potential invasion routes. A lack of genetic clustering across

south-eastern Europe may be a consequence of high levels of gene flow frequently introducing new

genotypes over large distances. However, the lack of geographic structure could also have its origins

from substantial population bottlenecks during the spread of S12 across south-eastern Europe.

Finally, the largely clonal lineage may also become exposed to processes such as Muller's Ratchet

fixing deleterious mutations over time (Felsenstein, 1978).

Conclusions

We show how a highly invasive fungal pathogen lineage can emerge from an intermediate,

genetically diverse bridgehead population. This is in line with the self-reinforcement invasion model

where initial introductions promote secondary spread (Bertelsmeier et al. 2018, Garnas et al. 2016).

However, empirical evidence for adaptation in bridgehead populations is often elusive. Additionally,

human transport (Banks et al. 2015) and host naivety of the European chestnut could have

contributed substantially to the rapid spread without the need for diversification and adaptation in the

bridgehead population. As a response to fungal pathogen invasions among bats, some host

populations evolved resistance (Langwig et al. 2017). There is no evidence for the emergence of

tolerance or resistance in the European chestnut, however deployed control measures such as the

artificial introduction of the *Cryphonectria hypovirus I* (CHV-1) can severely reduce fungal virulence

(Rigling & Prospero 2018). Interestingly, the mycoviral spread should be facilitated in asexual

populations such as the invasive S12 lineage due to the lack of vegetative compatibility barriers.

Outcrossing populations often harbor many different vegetative compatibility groups slowing

transmission (Robin & Heiniger 2001). Hence, the presence of the mycovirus may favor sexual

reproduction and the immigration of additional vegetative compatibility types. In turn, the

diversification may reduce the evolutionary advantage of the invasive lineage.

Materials and Methods

Samples of *C. parasitica*

A total of 188 virulent and mycovirus-free isolates of *C. parasitica* were sequenced. A majority of

the 182 isolates originated from Albania, Bosnia, Bulgaria, Croatia, Georgia, Greece, Italy, Kosovo,

Macedonia, Serbia, Slovenia, Switzerland, and Turkey (Fig. 1B, Supp. Table 1, Supp. Fig. 7). The

six other isolates were from South Korea (2) and North America (4) (Supp. Table 1). A total of 125

European isolates belong to the vegetative compatibility type EU-12, whereas 57 isolates represent

other vegetative compatibility types (also called EU-types) occurring in central and south-eastern

Europe. A total of 45 isolates from Bulgaria, Greece, Italy and Macedonia were already included in a

previous population-wide study on south-eastern European *C. parasitica* diversity by Milgroom et al.

(2008). All samples were collected between 1951-2018 and are stored as glycerol stocks at –80 °C in

the culture collection of the Swiss Federal Research Institute WSL.

DNA extraction and genotyping

All isolates were first inoculated onto cellophane-covered potato dextrose agar plates (PDA, 39 g/L;

BD Becton, Dickinson & Company; Franklin Lakes, USA) (Hoegger et al., 2000) and incubated for

a minimum of one week at 24 °C, at a 14 h light and 10 h darkness cycle. After a sufficient amount

of mycelium and spores had grown, the isolates were harvested by scratching the mycelial mass off

the cellophane, transferring it into 2 ml Eppendorf tubes and freeze-drying it for 24 h. For DNA

extraction, 15 – 20 mg of dried material was weighted and single tube extraction was performed

using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany). DNA quantity was measured using the

Invitrogen Qubit 3.0 Fluorometer (Thermo Fisher Scientific, Waltham (MA), USA) and DNA

quality was assessed using the Nanodrop 1000 Spectrophotometer (Thermo Fisher Scientific,

Waltham (MA), USA). Prior to sequencing, all isolates were screened for their genotype at 10

microsatellite markers (Prospero & Rigling 2012). Additionally, the isolates were screened for their

vegetative compatibility and mating type alleles in two multiplex PCRs, as described in Cornejo et

al. (2019). Allele sizes for the genotyping, as well as the vegetative compatibility and mating type

assays were scored with GeneMapper 5 (Thermo Fisher Scientific, Waltham MA, USA).

Illumina whole-genome sequencing, variant calling and filtration

Isolates were prepared for sequencing using the TrueSeq Nano DNA HT Library Preparation kit

(Illumina, San Diego, USA). The libraries were sent to the Functional Genomics Centre Zurich (ETH

Zurich and University of Zurich) and sequenced on the Illumina Hiseq 4000 platform (Illumina, San

Diego, USA). The obtained sequences were trimmed with Trimmomatic v0.36 (Bolger et al., 2014)

and aligned with Bowtie 2 v2.3.5.1 (Langmead & Salzberg, 2012) and SAMtools v1.9 (Li et al.,

2009) to the *C. parasitica* reference genome (43.9 Mb) EP155 v2.0 available at the Joint Genome

Institute (http://jgi.doe.gov/). Variant calling, selection and filtration were conducted with the

Genome Analysis Toolkit GATK v3.8 and v4.0.2.0 (McKenna et al., 2010). We retained variants

satisfying the following filtration parameters: QUAL: $\geq$ 100, MQRankSum (lower): $\geq$ -2.0, QD: $\geq$

20.0, MQRankSum (upper): $\leq$ 2.0, MQ: $\geq$ 20.0, BaseQRankSum (lower): $\geq$ -2.0, ReadPosRankSum

(lower): $\geq$ -2.0, ReadPosRankSum (upper): $\leq$ 2.0, BaseQRankSum(upper): $\leq$ 2.0 (Supp. Fig. 1).

Furthermore, we used BCFtools v1.9 (Narasimhan et al., 2016) and the R package vcfR (Knaus &

Grünwald, 2017, R Core Team, 2017) for querying and VCFtools v0.1.16 (Danecek et al., 2011) for

downstream variant filtering. Variants were additionally filtered for minor allele count (MAC) $\geq$1,

excluding all missing data. Sites were also filtered per genotype, only keeping biallelic SNPs with a

minimum depth of 3 and a genotyping quality (GQ) of 99. To exclude SNPs associated with the

mating type, we ran an association study with TASSEL 5 (Bradbury et al., 2007) and retrieved *p*-

values for each SNP across the genome. We set a *p*-value threshold to remove all SNPs with $p \leq 1 \times 10^{-10}$

with the mating type for further analysis. The SNPs showing strong association with the mating

type were located on scaffold 2 between positions 100'853 – 1'997'710 bp. (Supp. Fig. 2).

Phylogenetic reconstruction

The filtered whole-genome SNP dataset was used to build phylogenetic trees. We generated a

maximum likelihood (ML) tree with RAxML v8.2.11 (Stamatakis, 2006), applying rapid

bootstrapping generating 100 maximum likelihood (ML) trees. Phylogenetic trees were displayed

using FigTree v1.4.3. We also generated an unrooted phylogenetic network with using SplitsTree

v4.14.6. (Hudson & Bryant, 2005). SplitsTree was also used for calculating the PHI test (Bruen et

al., 2006) to test for recombination. The required file conversions for using RAxML and SplitsTree

(i.e. from VCF to FASTA format) were done with PGDSpider v2.1.1.5 (Lischer & Excoffier, 2011) .

We also performed a principal component analysis (PCA) as implemented in the R package ade4

(Bougeard & Dray, 2018).

Inference of S12 donor populations

We generated an averaged co-ancestry matrix as inferred by fineSTRUCTURE v2.1.3 (Lawson et al., 2012). The software uses a Markov-Chain-Monte-Carlo (MCMC) based algorithm to infer ancestral contributions based on patterns of haplotype similarity. We ran the fineSTRUCTURE pipeline in 'automatic mode', with 500 Expectation-Maximation (EM) and 300'000 MCMC iterations, 400'000 maximization steps to infer the best tree and with ploidy set to 1. fineSTRUCTURE input files were created with LDhat (McVean & Auton, 2007) and the R package vcfR.

Population genetic analyses and SNP impact assessment

We computed allele frequencies and estimated the allele spectrum using VCFtools. We used RStudio (RStudio Team, 2015) and ggplot2 (Wickham, 2016) for visualizations. Synonymous and non-synonymous sites were identified and annotated using SnpEff v4.3t (Cingolani et al., 2012). Variants which were classified by SnpEff as having a "high", "moderate" or "modifying" impact on fungal protein sequences, were further processed in R using the packages dplyr (Wickham et al., 2018), reshape2 (Wickham, 2007), tidyr (Wickham & Henry, 2018), as well as and ggplot2. Furthermore, we performed a genome-wide association study (GWAS) with TASSEL 5 for identifying highly associated SNPs with genetic groups such as S12.

Mating experiments

The ability of *C. parasitica* isolates belonging to S12 (of mating type MAT-1) to outcross with isolates of the opposite mating type, was assessed in an inoculation experiment. For this, we randomly selected four S12 and one non-S12 isolate with mating type MAT-1, as well as five isolates of opposite mating type MAT-2 from populations in Italy (Nebrodi), Kosovo (Molliq) and Bosnia (Konic, Vrnograč, Projsa) (Supp. Table 2). All isolates of both mating types belonged to the vegetative compatibility type EU-12 and mating pairs were only formed between isolates from the same geographic source populations. As substrate for the pairings, 40 mm long segments of dormant

chestnut (*C. sativa*) stems (15-20 mm in diameter) were split longitudinally in half. The wood pieces

were autoclaved twice, placed onto sterile petri dishes (90 mm diameter) filled with 1.5% water agar

to enclose the pieces. Mating pairs were inoculated on opposite sides of each halved stem with three

replicates per pairing. The inoculated plates were then incubated at 25 °C under a 16 h photoperiod

(2500 Lux) for 14 days. After two weeks, mating was stimulated by adding 5 ml sterile water to the

plates to suspend and distribute the conidia produced by both isolates over the stem segment. Any

excess water was subsequently removed and the plates were incubated at 18 °C under an 8 h

photoperiod (2500 Lux). After 5 months of incubation, perithecia formation was assessed under a

dissecting microscope. To confirm successful outcrossing, single perithecia were carefully extracted

from the stromata and crushed in a drop of sterile distilled water. The resulting ascospore suspensions

were plated on PDA and incubated at 25°C for 24-36 h. Afterwards, single germinating ascospores

were transferred to PDA and incubated in the dark for 3-5 days at 25°C. DNA was then extracted

from 10 mg of lyophilized mycelium using the kit and instructions by KingFisher (Thermo Fisher

Scientific, Waltham MA, USA). All single ascospore cultures were screened for mating types by

performing a multiplex PCR following the protocol described in Cornejo et al. (2019).

Acknowledgements

We are grateful to Ursula Oggenfuss for helpful comments on a previous manuscript version. We

thank Sandrine Fattore, Hélène Blauenstein, Quirin Kupper, Eva Augustiny, Dario Rüegg and Silvia

Kobel for laboratory assistance. We acknowledge the Genetic Diversity Centre (GDC), ETH Zurich,

for technical support and facility access. Ludwig Beenken and Martin Wrann helped with

documenting mating experiments. Kiril Sotirovski, Mihajlo Risteski, Marin Ježić and Seçil Akilli

kindly provided samples. We thank Pierre Gladieux and Nikhil Kumar Singh for discussions and

sharing scripts for data analyses. We are grateful for insightful discussions with Daniel Rigling.

Funding was awarded to SP by the Swiss National Science Foundation (grant 170188).

References

Allendorf, F. W. & Lundquist, L. L. (2003). Introduction: Population biology, evolution, and control of invasive species. *Conservation Biology*, *17* (1), 24–30.

Avolio, S. (1978). La castanicoltura in Calabria. Aspetti selvicolturali e linee di intervento. *L'Italia Forestale e Montana*, *2*, 278–296.

Banks, N. C., Paini, D. R., Bayliss, K. L., & Hodda, M. (2015). The role of global trade and transport network topology in the human-mediated dispersal of alien species. *Ecology Letters*, *18* (2), 188–199.

Bertelsmeier, C. & Keller, L. (2018). Bridgehead effects and role of adaptive evolution in invasive populations. *Trends in Ecology & Evolution*.

Bertelsmeier, C., Ollier, S., Liebhold, A. M., Brockerhoff, E. G., Ward, D., & Keller, L. (2018). Recurrent bridgehead effects accelerate global alien ant spread. *Proceedings of the National Academy of Sciences*, *115* (21), 5486–5491.

Billiard, S., López-Villavicencio, M., Hood, M., & Giraud, T. (2012). Sex, outcrossing and mating types: unsolved questions in fungi and beyond. *Journal of Evolutionary Biology*, *25* (6), 1020–1038.

Biraghi, A. (1946). *Cancro della corteccia del castagno*. Ramo editoriale degli agricoltori.

Bock, D. G., Caseys, C., Cousens, R. D., Hahn, M. A., Heredia, S. M., Hübner, S., Turner, K. G., Whitney, K. D., & Rieseberg, L. H. (2015). What we still don't know about invasion genetics. *Molecular Ecology*, *24* (9), 2277–2297.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30* (15), 2114–2120.

Bougeard, S. & Dray, S. (2018). Supervised multiblock analysis in R with the ade4 package. *J. Stat. Softw*, *86*, 1–17.

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., & Buckler, E. S. (2007). Tassel: software for association mapping of complex traits in diverse samples. *Bioinformatics*, *23* (19), 2633–2635.

Brasier, C. & Webber, J. (2019). Is there evidence for post-epidemic attenuation in the Dutch elm disease pathogen *Ophiostoma novo-ulmi*? *Plant Pathology*, *68* (5), 921–929.

Bruen, T. C., Philippe, H., & Bryant, D. (2006). A simple and robust statistical test for detecting the presence of recombination. *Genetics*, *172* (4), 2665–2681.

Buccianti, M. & Feliciani, A. (1966). La situazione attuale dei castagneti in italia (the present situation of chestnut stands in Italy). *Atti Convegno Internazionale sul Castagno, Cuneo*, 61–85.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., & Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, *6* (2), 80–92.

Cornejo, C., Šever, B., Kupper, Q., Prospero, S., & Rigling, D. (2019). A multiplexed genotyping assay to determine vegetative incompatibility and mating type in *Cryphonectria parasitica*. *European Journal of Plant Pathology*, 1–11.

Cortesi, P., McCulloch, C. E., Song, H., Lin, H., & Milgroom, M. G. (2001). Genetic control of horizontal virus transmission in the chestnut blight fungus, *Cryphonectria parasitica*. *Genetics*, *159* (1), 107–118.

Cortesi, P., Milgroom, M. G., & Bisiach, M. (1996). Distribution and diversity of vegetative compatibility types in subpopulations of *Cryphonectria parasitica* in Italy. *Mycological Research*, *100* (9), 1087–1093.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., et al. (2011). The variant call format and VCFtools. *Bioinformatics*, *27* (15), 2156–2158.

Demené, A., Legrand, L., Gouzy, J., Debuchy, R., Saint-Jean, G., Fabreguettes, O., & Dutech, C. (2019). Whole-genome sequencing reveals recent and frequent genetic recombination between clonal lineages of *Cryphonectria parasitica* in western Europe. *Fungal Genetics and Biology*.

Drenth, A., McTaggart, A. R., & Wingfield, B. D. (2019). Fungal clones win the battle, but recombination wins the war. *IMA fungus*, *10* (1), 18.

Dutech, C., Barrès, B., Bridier, J., Robin, C., Milgroom, M., & Ravigné, V. (2012). The chestnut blight fungus world tour: Successive introduction events from diverse origins in an invasive plant fungal pathogen. *Molecular Ecology*, *21* (16), 3931–3946.

Dutech, C., Fabreguettes, O., Capdevielle, X., & Robin, C. (2010). Multiple introductions of divergent genetic lineages in an invasive fungal pathogen, *Cryphonectria parasitica*, in France. *Heredity*, *105* (2), 220–228.

Elliott, K. J. & Swank, W. T. (2008). Long-term changes in forest composition and diversity following early logging (1919–1923) and the decline of American chestnut (*Castanea dentata*). *Plant Ecology*, *197* (2), 155–172.

Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics*, *78* (2), 737–756.

Fontaine, M. C., Gladieux, P., Hood, M. E., & Giraud, T. (2013). History of the invasion of the anther smut pathogen on *Silene latifolia* in North America. *New Phytologist*, *198* (3), 946–956.

Garbelotto, M., Rocca, G. D., Osmundson, T., di Lonardo, V., & Danti, R. (2015). An increase in transmission-related traits and in phenotypic plasticity is documented during a fungal invasion. *Ecosphere*, *6* (10), 1–16.

Garnas, J. R., Auger-Rozenberg, M.-A., Roques, A., Bertelsmeier, C., Wingfield, M. J., Saccaggi, D. L., Roy, H. E., & Slippers, B. (2016). Complex patterns of global spread in invasive insects: Eco-evolutionary and management consequences. *Biological Invasions*, *18* (4), 935–952.

Gau, R. D., Merz, U., Falloon, R. E., & Brunner, P. C. (2013). Global genetics and invasion history of the potato powdery scab pathogen, *Spongospora subterranea* f. sp. *subterranea*. *PLoS One*, *8* (6), e67944.

Gladieux, P., Feurtey, A., Hood, M. E., Snirc, A., Clavel, J., Dutech, C., Roy, M., & Giraud, T. (2015). The population biology of fungal invasions. *Molecular Ecology*, *24* (9), 1969–1986.

González-Varela, G., González, A. J., & Milgroom, M. G. (2011). Clonal population structure and introductions of the chestnut blight fungus, *Cryphonectria parasitica*, in Asturias, northern Spain. *European Journal of Plant Pathology*, *131* (1), 67–79.

Griffin, G. J. (1986). Chestnut blight and its control. *Horticultural Reviews*, *8*, 291–336.

Grünwald, N. J., Garbelotto, M., Goss, E. M., Heungens, K., & Prospero, S. (2012). Emergence of the sudden oak death pathogen *Phytophthora ramorum*. *Trends in Microbiology*, *20* (3), 131–138.

615 Heald, F. & Studhalter, R. A. (1914). Birds as carriers of the chestnut blight fungus. *Journal of*
616 *Agriculture Research.*, *2*, 405–422.
617

618 Heiniger, U. & Rigling, D. (1994). Biological control of chestnut blight in Europe. *Annual review of*
619 *Phytopathology*, *32* (1), 581–599.
620

621 Heitman, J., Sun, S., & James, T. Y. (2013). Evolution of fungal sexual reproduction. *Mycologia*,
622 *105* (1), 1–27.
623

624 Hoegger, P. J., Rigling, D., Holdenrieder, O., & Heiniger, U. (2000). Genetic structure of newly
625 established populations of *Cryphonectria parasitica*. *Mycological Research*, *104* (9), 1108–1116.
626

627 Huson, D. H. & Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies.
628 *Molecular Biology and Evolution*, *23* (2), 254–267.
629
630

Idnurm, A., Hood, M. E., Johannesson, H., & Giraud, T. (2015). Contrasted patterns in mating-type
chromosomes in fungi: Hotspots versus coldspots of recombination. *Fungal Biology reviews*, *29*
(3-4), 220–229.

Ježić, M., Krstin, L., Rigling, D., & Ćurković-Perica, M. (2012). High diversity in populations of the
introduced plant pathogen, *Cryphonectria parasitica*, due to encounters between genetically
divergent genotypes. *Molecular Ecology*, *21* (1), 87–99.
637

638 Karadžić, D., Radulović, Z., Sikora, K., Stanivuković, Z., Ćurguz, V. G., Oszako, T., & Milenković,
639 I. (2019). Characterisation and pathogenicity of *Cryphonectria parasitica* on sweet chestnut and
640 sessile oak trees in Serbia. *Plant Protection Science*, *55* (3), 191–201.
641

642 Knaus, B. J. & Grünwald, N. J. (2017). vcfr: A package to manipulate and visualize variant call
643 format data in R. *Molecular Ecology Resources*, *17* (1), 44–53.
644

645 Langmead, B. & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,
646 *9*, 357–360.
647

648 Langwig, K. E., Hoyt, J. R., Parise, K. L., Frick, W. F., Foster, J. T., & Kilpatrick, A. M. (2017). Re-
649 sistance in persisting bat populations after white-nose syndrome invasion. *Philosophical*
650 *Transactions of the Royal Society B: Biological Sciences*, *372* (1712), 20160044.
651

652 Lawson, D. J., Hellenthal, G., Myers, S., & Falush, D. (2012). Inference of population structure
653 using dense haplotype data. *PLoS genetics*, *8* (1), e1002453.
654

655 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
656 Durbin, R., & 1000 Genome Project Data Processing Subgroup (2009). The Sequence
657 Alignment/Map format and SAMtools. *Bioinformatics*, *25* (16), 2078–2079.
658

659 Lischer, H. E. & Excoffer, L. (2011). PGDSpider: an automated data conversion tool for connecting
660 population genetics and genomics programs. *Bioinformatics*, *28* (2), 298–299.
661

662 Mariette, N., Mabon, R., Corbière, R., Boulard, F., Glais, I., Marquer, B., Pasco, C., Montarry, J.,
663 & Andrivon, D. (2016). Phenotypic and genotypic changes in French populations of *Phytophthora*
664 *infestans:* Are invasive clones the most aggressive? *Plant Pathology*, *65* (4), 577–586.
665
666

667 Marra, R., Cortesi, P., Bissegger, M., & Milgroom, M. (2004). Mixed mating in natural populations
668 of the chestnut blight fungus, *Cryphonectria parasitica*. *Heredity*, *93* (2), 189–195.
669

670 Marra, R. E. & Milgroom, M. G. (2001). The mating system of the fungus *Cryphonectria parasitica*:
671 Selfing and self-incompatibility. *Heredity*, *86* (2), 134–143.
672
673

674 McGuire, I., Davis, J., Double, M., MacDonald, W., Rauscher, J., McCawley, S., & Milgroom, M.
675 (2005). Heterokaryon formation and parasexual recombination between vegetatively incompatible
676 lineages in a population of the chestnut blight fungus, *Cryphonectria parasitica*. *Molecular*
677 *Ecology*, *14* (12), 3657–3669.
678

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, 1297–1303.

McVean, G. & Auton, A. (2007). LDhat 2.1: A package for the population genetic analysis of recombination. *Department of Statistics, Oxford, OX1 3TG, UK*.

Milgroom, M. G. & Cortesi, P. (1999). Analysis of population structure of the chestnut blight fungus based on vegetative incompatibility genotypes. *Proceedings of the National Academy of Sciences*, *96* (18), 10518–10523.

Milgroom, M. G., Sotirovski, K., Risteski, M., & Brewer, M. T. (2009). Heterokaryons and parasexual recombinants of *Cryphonectria parasitica* in two clonal populations in southeastern Europe. *Fungal Genetics and Biology*, *46* (11), 849–854.

Milgroom, M. G., Sotirovski, K., Spica, D., Davis, J. E., Brewer, M. T., Milev, M., & Cortesi, P. (2008). Clonal population structure of the chestnut blight fungus in expanding ranges in southeastern Europe. *Molecular Ecology*, *17* (20), 4446–4458.

Milgroom, M. G., Wang, K., Zhou, Y., Lipari, S. E., & Kaneko, S. (1996). Intercontinental population structure of the chestnut blight fungus, *Cryphonectria parasitica*. *Mycologia*, *88* (2), 179–190.

Myteberi, I. F., Lushaj, A. B., Keča, N., Lushaj, A. B., & Lushaj, B. M. (2013). Diversity of *Cryphonectria parasitica*, hypovirulence, and possibilities for biocontrol of chestnut canker in Albania. *Int. J. Microb. Res. Rev*, *1*, 11–21.

Narasimhan, V., Danecek, P., Scally, A., Xue, Y., Tyler-Smith, C., & Durbin, R. (2016). BCFtools/RoH: A hidden Markov model approach for detecting autozygosity from next-generation sequencing data. *Bioinformatics*, *32* (11), 1749–1751.

Paoletti, M., Buck, K. W., & Brasier, C. M. (2006). Selective acquisition of novel mating type and vegetative incompatibility genes via interspecies gene transfer in the globally invading eukaryote *Ophiostoma novo-ulmi*. *Molecular Ecology*, *15* (1), 249–262.

Philibert, A., Desprez-Loustau, M.-L., Fabre, B., Frey, P., Halkett, F., Husson, C., Lung-Escarmant, B., Marçais, B., Robin, C., Vacher, C., et al. (2011). Predicting invasion success of forest pathogenic fungi from species traits. *Journal of Applied Ecology*, *48* (6), 1381–1390.

Prospero, S., Conedera, M., Heiniger, U., & Rigling, D. (2006). Saprophytic activity and sporulation of *Cryphonectria parasitica* on dead chestnut wood in forests with naturally established hypovirulence. *Phytopathology*, *96* (12), 1337–1344.

Prospero, S. & Rigling, D. (2012). Invasion genetics of the chestnut blight fungus *Cryphonectria para-sitica* in Switzerland. *Phytopathology*, *102* (1), 73–82.

R Core Team (2014). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.

Raboin, L.-M., Selvi, A., Oliveira, K. M., Paulet, F., Calatayud, C., Zapater, M.-F., Brottier, P., Luzaran, R., Garsmeur, O., Carlier, J., et al. (2007). Evidence for the dispersal of a unique lineage from Asia to America and Africa in the sugarcane fungal pathogen *Ustilago scitaminea*. *Fungal Genetics and Biology*, *44* (1), 64–76.

Rambaut, A. (2016). Figtree v1. 4.3 software. Institute of Evolutionary Biology, University of Edinburgh.

Rigling, D. & Prospero, S. (2018). *Cryphonectria parasitica*, the causal agent of chestnut blight: Invasion history, population biology and disease control. *Molecular Plant Pathology*, *19* (1), 7–20.

Robin, C. & Heiniger, U. (2001). Chestnut blight in Europe: Diversity of *Cryphonectria parasitica*, hypovirulence and biocontrol. *Forest Snow and Landscape Research*, *76* (3), 361–367.

RStudio Team (2015). *RStudio: Integrated Development Environment for R*. Boston, MA: RStudio, Inc.

Sherpa, S., Blum, M. G., Capblancq, T., Cumer, T., Rioux, D., & Després, L. (2019). Unravelling the invasion history of the Asian tiger mosquito in Europe. *Molecular Ecology*, *28* (9), 2360–2377.

Sotirovski, K., Papazova-Anakieva, I., Grünwald, N., & Milgroom, M. (2004). Low diversity of vegeta-tive compatibility types and mating type of *Cryphonectria parasitica* in the southern Balkans. *Plant Pathology*, *53* (3), 325–333.

Stamatakis, A. (2006). RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thou-sands of taxa and mixed models. *Bioinformatics*, *22* (21), 2688–2690.

Suehs, C., A□re, L., & Médail, F. (2004). Invasion dynamics of two alien *Carpobrotus* (Aizoaceae) taxa on a Mediterranean island: II. Reproductive strategies. *Heredity*, *92* (6), 550.

Taylor, J. W., Hann-Soden, C., Branco, S., Sylvain, I., & Ellison, C. E. (2015). Clonal reproduction in fungi. *Proceedings of the National Academy of Sciences*, *112* (29), 8901–8908.

Trestic, T., Uscuplic, M., Colinas, C., Rolland, G., Giraud, A., & Robin, C. (2001). Vegetative compatibility type diversity of *Cryphonectria parasitica* populations in Bosnia-Herzegovina, Spain and France. *For. Snow Landsc. Res.*, *76* (3), 391–396.

van Boheemen, L. A., Lombaert, E., Nurkowski, K. A., Gau□re, B., Rieseberg, L. H., & Hodgins, K. A. (2017). Multiple introductions, admixture and bridgehead invasion characterize the introduction history of *Ambrosia artemisiifolia* in Europe and Australia. *Molecular Ecolo*, *26* (20), 5421–5434.

Vuković, R., Liber, Z., Ježić, M., Sotirovski, K., & Ćurković-Perica, M. (2019). Link between epigenetic diversity and invasive status of south-eastern European populations of phytopathogenic fungus *Cryphonectria parasitica*. *Environmental Microbiology*.

Wickham, H. (2007). Reshaping Data with the reshape Package. *Journal of Statistical Software*, *21* (12), 1–20.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wickham, H., François, R., Henry, L., & Müller, K. (2018). *dplyr: A Grammar of Data Manipulation*. R package version 0.7.8.

Wuest, C. E., Harrington, T. C., Fraedrich, S. W., Yun, H.-Y., & Lu, S.-S. (2017). Genetic variation in native populations of the laurel wilt pathogen, *Ra□aelea lauricola*, in Taiwan and Japan and the introduced population in the United States. *Plant Disease*, *101* (4), 619–628.

Yang, X.-M., Sun, J.-T., Xue, X.-F., Li, J.-B., & Hong, X.-Y. (2012). Invasion genetics of the western flower thrips in China: Evidence for genetic bottleneck, hybridization and bridgehead e□ect. *PLoS One*, *7* (4), e34567.

789   **Table**

791   **Table 1: Donor isolates for S12 identified using FineStructure**. The geographic origin (country and
792   population), vegetative compatibility (vc) type and mating type of donors contributing to genotypes of
793   the invasive S12 lineage are given. See Figure 3 for the corresponding co-ancestry matrix.
794
795

Country	Population	vc type	Mating type	Donor isolate
Bosnia	Projsa	EU-11	MAT-1	M1412
	Projsa	EU-12	MAT-2	M1407
	Projsa	EU-12	MAT-2	M1430
	Projsa	EU-12	MAT-2	M1431
	Vrnograč	EU-12	MAT-1	M1808
	Vrnograč	EU-15	MAT-2	M1834
	Vrnograč	EU-18	MAT-2	M1797
Croatia	Kostajnica	EU-12	MAT-1	HK60B
Georgia	Mackhunceti	EU-12	MAT-2	MAK23
Switzerland	Gnosca	EU-12	MAT-1	M6697
	Biasca	EU-12	MAT-2	M4023
	Biasca	EU-12	MAT-2	M4022
	Claro	EU-12	MAT-2	M2466

A

B

PC2 (vertical axis), PC1 (horizontal axis)

Regions

- ▲ Northern Balkans
- ● Central Balkans
- + Italy
- ◆ Georgia
- ✕ Greece/Turkey

C

0.01

D

★ MUL63B	x	MUL59
MAT-1		**MAT-2**
⬟ NE202	x	NE164
MAT-1		**MAT-2**

E

1mm

500µm

A

B