The interaction of crossover formation and the dynamic architecture of the synaptonemal complex during meiosis

Simone Köhler[1,2,*], Michal Wojcik[3,4,*], Ke Xu[3,4,5,6,#] and Abby F. Dernburg[1,2,5,7,#]

1. Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley CA 94720-3220
2. Howard Hughes Medical Institute, 4000 Jones Bridge Road, Chevy Chase, MD 20815
3. Department of Chemistry, University of California, Berkeley, Berkeley CA 94720-3220
4. Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
5. California Institute for Quantitative Biosciences, Berkeley CA 94720
6. Division of Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA 94720
7. Division of Biological Systems and Engineering, Lawrence Berkeley National Laboratory, Berkeley CA 94720

* these authors contributed equally to this work.
# corresponding authors:
Ke Xu <xuk@berkeley.edu>
Abby F. Dernburg < afdernburg@berkeley.edu >

During meiosis, chromosomes pair with their homologous partners, and a structure known as the synaptonemal complex (SC) assembles between them. Evidence indicates that this material plays a role in regulating crossover recombination between homologs, although this remains controversial. Here we use three-dimensional STochastic Optical Reconstruction Microscopy (3D-STORM) to interrogate the molecular architecture of the SC in intact germline tissue from *Caenorhabditis elegans*, and analyze its ultrastructure during meiotic progression. Using a probabilistic mapping approach, we determine the position and orientation of the four known SC structural proteins. We report that a marked structural transition occurs within this material upon crossover designation. We also identify a mutation that disturbs both SC ultrastructure and crossover interference, the widespread mechanism that results in non-randomly wide spacing of crossovers between the same chromosome pair. Together with other evidence, our findings suggest that the SC is an active material, and that structural transitions may directly contribute to chromosome-wide crossover regulation.

The synaptonemal complex (SC) is a protein structure that assembles between homologous chromosomes during meiotic prophase, stabilizing their parallel alignment [1]. The SC appears as a periodic ladder-like structure in electron micrographs [2]. Recent studies have revealed that this material shows dynamic, liquid-like behavior [3-7]. The SC is required for and appears to regulate meiotic recombination [3, 8-13]. Despite its central role in meiosis, the molecular organization of the SC remains poorly characterized.

In *C. elegans,* the SC assembles between homologs even in the absence of meiotic recombination [14]. Four structural proteins essential for SC assembly have been identified: SYP-1, SYP-2, SYP-3, and SYP-4 [15-18]. Limited information about their organization within the SC has been obtained through protein-protein interaction analysis and immuno-electron microscopy [19]. Super-resolution microscopy has emerged as a powerful tool to investigate the organization of macromolecular assemblies, including the SC [20, 21]. We recently described the molecular architecture of meiotic chromosome axes, which form the substrate for SC assembly, based on 3D-STORM [22, 23] in intact *C. elegans* gonads [24]. Here, we apply this approach to analyze the organization of the SC during meiotic prophase.

Results

Using antibodies that bind either short peptides within SC proteins or epitope tags inserted by genome editing, we localized the SYP proteins along paired meiotic chromosomes using 3D-STORM and averaging methods (Extended Data Fig. S1). The orientation of the SC in our images was defined by labeling a reference protein in the chromosome axes, HIM-3 [25]. For positional mapping of SC proteins, we analyzed regions that were observed in frontal view (Fig. 1b, c). We rotated all images about the optical axis (z-axis) such that the chromosomes are extended along the y-axis, with the SC spanning the distance between the homologs on the x-axis (Extended Data Fig. S1b-c, methods). Taking advantage of the well-characterized progression of meiotic stages within the germline of adult *C. elegans*, we compared SC organization before, during and after the designation of crossover sites (Fig. 1a).

Dynamic architecture of the synaptonemal complex in *C. elegans*

The width of the SC in *x* in *C. elegans*, as measured from electron micrographs, is approximately 96 nm [3]. An antibody recognizing the C-terminal peptide of SYP-1 was resolved as two parallel strands, located at 42.3±0.8 nm and 42.0±1.2 nm off-center in early and late pachytene, respectively (Fig. 1b, m). This epitope thus lies near the outer edges of the SC and within a few nanometers of HIM-3, the most proximal known component of the chromosome axis (Table S1)[24]. Intriguingly, in cross-sectional views (*xz*-view, Fig. 1c) we observed striking differences in its distribution in *z* between early and late pachytene SCs: While the C-terminus of SYP-1 was confined near the central plane in early pachytene, it was widely distributed in *z* in late pachytene (Fig. 1d, e, m, center and Table S1).

We then asked whether the N-terminus of SYP-1 is similarly relocalized during meiotic progression. Tagging the N-terminus of SYP-1 disrupts SC assembly ([26] and data not shown), but we identified a poorly conserved region close to the N-terminus in which insertion of an HA epitope did not impair function (Table S2). Interestingly, this epitope was confined to a thin plane in the center of the SC along both the *x* and *z* axis in both early and late pachytene SCs (Fig. 1d, e, m, top). Our data support a transverse orientation for SYP-1, as previously reported [19], with its N-terminus at the center and C-terminus near the axes. Thus, SYP-1 appears to be an ortholog of "transverse filament" proteins from other organisms [27-31]. Importantly, these localization data are inconsistent with a uniform growth in thickness of the SC through e.g.

lateral stacking of SYP-1 molecules during meiotic progression, which has previously been

suggested based on observations of increased fluorescence intensity during late pachytene in

budding yeast [4] and *C. elegans* [5]. The distinct localization patterns of N- and C-terminal

epitopes in early and late SCs along the *z* axis suggest a more complex reorganization of SYP-1

during meiosis.

To further probe the orientation of SYP-1 within the SC, we mapped the N-terminal

localization events to the C-terminal distributions, assuming a rigid, rod-like conformation (Fig.

S2a-b, see methods). Our approach indicated that SYP-1 lies nearly parallel to the SC in early

pachytene, but is diagonally oriented in late pachytene (Fig. 2a-b).

We next asked how the organization of other SC components was affected by meiotic

progression. We mapped the C-termini of SYP-2, SYP-3-HA and SYP-4-HA, the N-terminus of

GFP-SYP-3[32], and an epitope inserted into the middle of SYP-4 (Table S2, Fig. 1d, e, m and

Extended Data Fig. S3). The results are summarized in Table S1. We observed consistent

differences between early and late pachytene for most epitopes, particularly for the C-terminus of

SYP-4, which moves from a central plane in early pachytene to a splayed orientation, protruding

above and below the central plane of the SC, in late pachytene (Fig. S2c and S3, Table S1).

Similarly, epitopes on SYP-2 and SYP-3 were more central in early pachytene than late

pachytene (Fig. 1d,e,m, S2c and Fig. S3). We further found that SYP-3 localized at (N) or close

to (C) the midline of the SC in *x*, suggesting a head-to-head arrangement of SYP-3 molecules

(Fig. S3 and Table S1). These findings are inconsistent with one conclusion of a previous study,

that the C-terminus of SYP-3 lies in proximity to the axes [19]. This discrepancy may reflect the

challenges of determining the orientation of the SC in samples prepared for immuno-EM.

Overall, 3D-STORM imaging and probabilistic modeling provide new insights into SC

ultrastructure and reveal a marked reorganization of SC components during meiotic progression

(Fig. S2c).

Crossover formation is accompanied by a change in synaptonemal complex organization

Programmed double-strand breaks (DSBs) are generated during early prophase. A subset

eventually gives rise to crossover (CO) recombination events. In *C. elegans* each pair of

homologous chromosomes undergoes only a single CO, while all other breaks are repaired to

yield noncrossover products. Some of the key signaling molecules that regulate CO site selection

localize to the SC [3, 11-13, 33, 34], and the integrity of this structure is important for CO

regulation [9]. Previous work from our group revealed that the SC behaves as a liquid crystal[3],

suggesting that diffusing signals and/or structural transitions could be propagated through this

material [35]. Interestingly, recent evidence suggests that the central region of the SC grows in

length and/or thickness throughout the pachytene stage of meiotic prophase, and that SC proteins

may become more stably associated with this structure upon crossover formation [4-7, 9]. Thus,

we wondered whether the dramatic reorganization of the SC during meiotic prophase might be

linked to crossover formation and might provide a structural basis for the SC's role in CO

regulation. We therefore first asked whether the timing of SC reorganization coincides with CO

designation. In *C. elegans*, designated crossovers can be visualized as bright GFP-COSA-1 foci,

which are first detected in mid-pachytene [36] (Fig. 1a). To test whether the structural transition

within the SC coincides with the appearance of bright GFP-COSA-1 foci, we imaged nuclei at

mid-pachytene within intact gonads. We inferred the crossover designation status of individual

nuclei based on the detection of bright COSA-1 foci and classified them as either mid(-)

pachytene or mid(+) pachytene. Throughout mid-pachytene, SYP-1 retains its transverse

orientation and both N- and C-termini of SYP-1 are widely distributed in *z* (Fig. 1f,g,m and

Table S1). This is consistent with a lateral stacking of SYP-1 molecules in parallel orientation, as

observed in early pachytene (Fig. 2a,b). However, the distance of the C-termini of SYP-1 from

the central axis of the SC in frontal view was drastically lower in mid(+) pachytene nuclei

compared to SCs in mid(-) pachytene nuclei.

This indicates that the designation of COs coincides with a conformational switch within

the SC. To take an unbiased approach to analyze the dynamics of SC organization during meiotic

prophase, we employed principal component analysis (PCA) of SYP-1 and SYP-2 distributions

(see methods). The first dimension in the PCA, indicative of the main changes within the SC

across different stages during meiotic prophase, corresponds to the transition from early to late

pachytene SCs (Fig. 2c). Mid(+) pachytene SCs are similar to late pachytene SCs along the first

principal component, while mid(-) SCs are in between early and late pachytene SCs, indicating

that the SC is reorganized upon CO designation.

In *C. elegans*, designation of COs triggers the asymmetrical remodeling of chromosome

axes and the SC to create two domains on either side, known as the long and the short arm.

Downstream steps in this differentiation require the recruitment of the Polo-like kinase PLK-2 to

the SC [5, 37, 38]. To assess whether the changes we observe in SC ultrastructure are dependent

on crossover designation and/or the subsequent remodeling processes, we determined the

organization of the SC in *zhp-3(jf61)* null mutants, which lack crossovers [39], and in *plk-*

2(ok1936) null mutants, which are proficient for CO designation but show pronounced delays in

downstream remodeling and arm differentiation [5, 7, 37, 40], although they do eventually

segregate most chromosomes properly [38]. We found that SCs in late pachytene in *zhp-3(jf61)*

mutants resembled those in early pachytene nuclei in wild-type hermaphrodites, while SCs at late

pachytene in *plk-2(ok1936)* mutants are very similar to wild-type (Fig. 1h,i,n, 2 and Fig. S3c,e).

Thus, structural reorganization of the SC requires CO designation but is independent of PLK-2.

We note that PLK-2 also plays an early role in homologous paring and synapsis [37, 40], and our

super-resolution images reveal severe defects in SC architecture in *plk-2(ok1936)* null mutants

despite the overall similarity to wild-type late pachytene SCs (Fig. 1i, n). Similarly, while the

global SC architecture in *zhp-3(jf61)* mutants resembled early pachytene SCs in wild type, the

SC increased in thickness along the optical axis during pachytene in *zhp-3(jf61)* oocytes, based

on the spatial distribution of SYP-1 (Fig. 1h, n), consistent with the continuous incorporation of

SC subunits even in the absence of crossover formation [5]. In summary, our data indicate that

the SC grows in thickness from early to late pachytene independent of crossovers, while the

reorganization of SC ultrastructure requires and coincides with CO designation.

Defects in SC organization are linked to defects in crossover interference

The SC plays a critical role in regulating the number and distribution of CO

recombination events [3, 8, 9, 12, 13]. We therefore wondered whether reorganization of the SC

is linked to CO regulation. Fortuitously, in our efforts to isolate a functional C-terminally tagged

syp-4 allele, we identified an unusual, partially functional allele, *syp-4(ie25)* (Fig. 3a). In *syp-*

4(ie25) homozygotes, we observed timely homologous synapsis, as determined by the length of

the transition zone, which corresponds to the leptotene and zygotene stages of meiosis. Six

bivalents were consistently observed at diakinesis, indicative of CO formation on all six

chromosome pairs (Fig. 3b, c, e). However, super-resolution microscopy indicated that the

architecture of the SC was drastically altered in *syp-4(ie25)* late pachytene after crossover

formation (Fig. 1l and 2, Extended Fig. S3). SYP-1 lies in a highly tilted orientation (Fig. 2a,b)

and the overall conformation of the SC in *syp-4(ie25)* at late pachytene is clearly distinct from

any conformation in wild-type animals in our principal component analysis (Fig. 2c). In contrast,

early pachytene SCs in *syp-4(ie25)* are more reminiscent of SC conformation of prior to CO

formation in wild-type animals (Fig. 1k and 2). These data suggest a strong defect in SC

organization in *syp-4(ie25)* that is exacerbated upon CO designation (Fig. 2c).

Notably, *syp-4(ie25)* mutant hermaphrodites exhibited high frequencies of chromosome

missegregation, resulting in embryonic lethality (38±4% survival, compared to 104% in wild-

type) and male self-progeny (6±1% compared to 0.1% in wild-type, Table S2), albeit less severe

than in a *syp-4(tm2713)* deletion allele (2.5% viability and 40% males) [18]. Importantly,

crossover formation was severely impacted in *syp-4(ie25)*: We observed a slight delay in meiotic

progression, indicated by a modest extension of CHK-2 kinase activity [41] (Fig. 3d) and an

accumulation of RAD-51 foci, which mark unrepaired recombination intermediates (Extended

Data Fig. S4). Moreover, the structure of bivalents at diakinesis was altered in *syp-4(ie25)*

oocytes (Fig. 3f), indicating that crossover formation might be affected [42, 43].

To better characterize the effects of *syp-4(ie25),* we imaged GFP-COSA-1 to quantify

designated crossover sites at late pachynema [36]. Wild-type animals display a single bright

GFP-COSA-1 focus on each of the 6 chromosome pairs, even when the number of DSBs is

markedly increased [36, 42]. In contrast to the 6±0.2 (s.d.) GFP-COSA-1 foci we observed in

wild-type and heterozygous *ie25/+* animals, we observed 10.9±1.7 GFP-COSA-1 foci at late

pachytene in homozygous *syp-4(ie25)* hermaphrodites (Fig. 3g, h). This suggested that crossover

interference, which in wild-type animals acts over distances longer than any single chromosome

[9, 42], is greatly decreased in *syp-4(ie25)* homozygotes. To test this hypothesis, we use two

different metrics to quantify the crossover interference strength, the gamma factor [44] and the

coefficient of coincidence [45]. Each of these approaches confirmed that crossover interference

is severely reduced or absent in *syp-4(ie25)* mutants (Fig. 3i,j).

To verify that GFP-COSA-1 foci indeed correspond to designated crossover sites in wild-

type and *syp-4(ie25)* animals, we also measured crossing-over genetically using whole genome

sequencing (see methods). This confirmed that crossovers in *syp-4(ie25)* homozygotes match the

number of GFP-COSA-1 foci. Thus, *syp-4(ie25)* mutants are proficient for synapsis, yet they

exhibit drastic alterations in SC architecture and cannot impose normal crossover interference.

Crossover interference acts along the physical length of the synapsed chromosomes, and

in some organisms chromosomes with longer axes undergo more crossovers than chromosomes

with shorter axes [46-49]. Therefore, we next asked whether the axis length is increased in *syp-*

4(ie25) mutants. Interestingly, axes in *syp-4(ie25)* mutants are shorter than in wild-type animals.

Thus, changes in axis length cannot account for the observed increase in COs in *syp-4(ie25)*

mutants. We hypothesize that the increase in CO number may be caused by the defects in SC

organization in *syp-4(ie25)* mutants.

Partial depletion of SC proteins in *C. elegans* by RNAi can result in a slight increase in

the number of crossovers, to about 7 GFP-COSA-1 foci per nucleus [9]. Thus, we tested whether

the defect in crossover regulation in *syp-4(ie25)* can be attributed to reduced levels of SC

proteins. The abundance of an epitope-tagged HA-SYP-1 was not altered by the *syp-4(ie25)*

mutation (Fig. S5a). However, the immunofluorescence signal of SYP-1 and SYP-2 was reduced

in *syp-4(ie25),* although staining appeared contiguous along the chromosomes (Fig. S5b, c). This

indicates the *syp-4(ie25)* mutation impairs assembly of SYP proteins between chromosomes. In

contrast, we also isolated a mutant, *syp-4(ie27)* (Fig. S6a), which reduced expression levels of

SYP-4-HA to about 30% of wild-type (Fig. S6c). Both the transition zone and CHK-2 active

zone were extended in *syp-4(ie27)* single mutants, indicating that completion of synapsis and CO

designation were delayed (Fig. 3c-d, Fig. S6b, d). We also observed a slight increase in the

number of GFP-COSA-1 foci to 7±1.1 (Fig. S6e, Fig. 3g), which is comparable to results

obtained for partial depletion of SYP-1, -2 or -3 by RNAi [9], but distinct from our results for

syp-4(ie25). This indicates that the defects in crossover interference in *syp-4(ie25)* are not a mere

consequence of reduced protein expression.

To test whether the defects in SC ultrastructure can account for the effects of *syp-4(ie25)*

on crossover regulation, we investigated the role of *syp-4(ie25)* in the regulation of COSA-1 foci

within aggregates of SC proteins known as polycomplexes. These bodies self-assemble prior to

or after synapsis in many organisms, including in early meiotic nuclei in *C. elegans*. Large

polycomplexes also assemble in meiotic nuclei in worms lacking the axis protein HTP-3, and

show periodic striations in orthogonal directions that correspond precisely to the width and

longitudinal banding of the SC [3, 50]. Intriguingly, we have found that polycomplexes in *htp-3*

null mutants recapitulate key aspects of the spatial patterning of CO proteins normally seen along

SCs, although these mutant animals lack DSBs and meiotic recombination intermediates.

Specifically, at late pachytene, most polycomplexes display a single COSA-1 focus (1.1±0.2;

Fig. 4a, b) [3]. When we combined the *syp-4(ie27)* mutation, which reduces expression of SYP-

4, with Δ*htp-3*, this resulted in smaller polycomplexes, but they showed an identical number of

GFP-COSA-1 foci at late pachytene (1.1±0.2) (Fig. 4a, b). By contrast, when *syp-4(ie25)* was

combined with Δ*htp-3,* multiple GFP-COSA-1 foci (2.1±0.8) were associated with each

polycomplex (Fig. 4a, b), mirroring the ~2-fold increase in GFP-COSA-1 foci observed along

bona fide SCs in *syp-4(ie25)* mutants.

We thus interrogated the organization of polycomplexes in *syp-4(ie25)* by EM. While

structures resembling the electron-dark lateral and brighter, striated central regions were

observed in *Δhtp-3 syp-4(ie25)*, the overall organization was dramatically altered: polycomplexes

appeared to be internally fragmented, rather than maintaining constant orientation over hundreds

of nm, as in "normal" polycomplexes [3] (Fig. 4c), and the distance between parallel electron-

dark bands was much narrower (46.3±1.2 nm, vs 97.6±1.5 nm in *Δhtp-3* polycomplexes, Fig.

4c), This likely reflects differences in organization and/or orientation of SYP proteins within

polycomplexes. Overall, the disorganized structure of polycomplexes in *Δhtp-3 syp-4(ie25)* in

EM is reminiscent of the architecture of SCs in *syp-4(ie25)* determined by 3D-STORM. These

findings indicate that structural defects within polycomplexes are sufficient to alter special

patterning of CO proteins, analogous to the structurally altered SCs that assemble in *syp-4(ie25)*

mutants.

Discussion

Together these findings establish that the correct organization of the synaptonemal

complex is required for spatial patterning of crossovers, and further corroborate previous

findings that the SC is important for crossover interference. They further suggest that CO

regulation may involve transduction of structural information through this liquid crystalline

material. This idea is reminiscent of a "beam-film" model for crossover interference, which

proposes that crossovers relieve local stress along a beam-like material [49, 51]. This mechanical

stress has been proposed to be imposed by chromatin structure and/or tethering of chromosomes;

however, our findings reinforce previous studies that implicated the SC as a crucial mediator of

crossover patterning [8, 9, 12]. In particular, we have reported that polycomplexes comprised of

SC central region proteins alone can mediate spatial patterning of GFP-COSA-1 foci even in the

absence of association with chromatin or axis proteins, and polycomplexes in *syp-4(ie25)*

mutants exhibit both defects in their internal organization and altered patterning of GFP-COSA-1

foci. Interestingly, forces applied to nematic liquid crystalline materials such as the SC [3] can

perturb the orientation of molecules within the liquid crystal [35]. Therefore, we speculate that

the reorganization of proteins within the SC that occurs concomitant with CO designation may

reflect a mechanical signaling component of CO interference.

 Figures

279
280

Figure 1: Super-resolution microscopy reveals the dynamic three-dimensional organization of the synaptonemal complex. (a) In the *C. elegans* gonad, nuclei (representative DAPI images) are arranged in a gradient of premeiotic and meiotic stages. Using highly inclined illumination

(red), we imaged SCs in the bottom section of nuclei adjacent to the coverslip (gray) in early,
mid- and late pachytene (rectangles). Mid(-) (cyan) and mid(+) (red) pachytene correspond to
nuclei without and with bright GFP-COSA-1 foci in the same field of view, respectively. (b)
Measurements based on averaged STORM images indicate that the C-terminus of SYP-1 lies
close to the chromosome axes. Colors denote localizations in z from -150 to 150 nm (color scale
bar). (c) The schematic depicts the SC, which forms a ladder-like structure approximately 100-
nm wide (purple), which links the axes (black) of paired homologous chromosomes (white rods)
in the frontal view (left). To visualize the 3D architecture of the SC, we show cross-sectional
views of the SC as depicted (right). (d-l) Averaged cross-sectional views of STORM images for
the N-terminus of HA-SYP-1 (top row), the C-terminus of SYP-1 (center row), and C-terminus
of SYP-2 (bottom row), for wild-type SCs in early (d, blue closed spheres in m), late (e, purple
closed spheres in m), mid(-) (f, cyan open squares in m), and mid(+) (g, red closed squares in m)
pachytene and for *zhp-3(jf61)* (h, green triangles, late pachytene in n), *plk-2(ok1936)* (i, pink
diamonds in n, late pachytene) and *syp-4(ie25)* (k, black closed triangles for early pachytene
orange crosses in late pachytene in n). (m, n) Histograms of localization events in frontal view
(x, left) and across the SC (z, right) are fitted with 1 or 2 Gaussians. The results of the fits are
summarized in Extended Data Table S1.

Figure 2: The synaptonemal complex undergoes a structural transition upon crossover designation. (a) The stochastic nature of STORM does not allow us to image the N- and C-terminus of a specific single molecule at the same time but will yield population averages for the localization of individual domains. We therefore used a probabilistic mapping approach to reconstruct the orientation of SYP-1 molecules for each condition and genotype. The resultant models of SYP-1 (black) are shown with C-termini depicted as cubes and N-termini as spheres. A proxy for chromatin (gray areas) is added for visualization purposes. Models are rendered using POV-ray (v3.7.0). Models show changes in the orientation of SYP-1 upon crossover formation. (b) These changes are revealed by the distributions of angles of SYP-1 with respect to the central plane of the SC in cross-sectional views. Cumulative distribution functions are shown. Gray horizontal lines mark median (50%) and 90th percentile values. (c) A principal component analysis (see methods) summarizes the SC organization at different conditions and

genotypes. The progression of SC reorganization in wild-type animals throughout pachytene is

highlighted by a gray dashed line. Dimension 1 scales with SC reorganization upon crossover

formation (colored arrow). We note that mid(-) SCs prior to designation of COs marked by

bright GFP-COSA-1 foci are closer to late pachytene than early pachytene. This suggests that

some aspects of the transition in SC reorganization may occur even prior to the appearance of

GFP-COSA-1 foci and/or nuclei designated mid(-) pachytene may include false negatives given

that our field of view in *z* is limited to 1 μm slices and does not contain whole nuclei. SC

organization in late pachytene *zhp-3(jf61)* mutants that lack CO formation is similar to wild-type

early pachytene, while SCs in *plk-2(ok1936)* null mutants resemble wild-type SCs. By contrast,

SCs in *syp-4(ie25)* mutants are distinct in both early and late pachytene.

Figure 3: Crossover interference is diminished in *syp-4(ie25)* mutants. (a) The C-terminal
amino-acid sequences of wild-type SYP-4 and SYP-4(*ie25*) are indicated. (b) Fluorescence
micrographs of DAPI-stained gonads from wild-type (*syp-4-ha(ie29)*, top) and *syp-4(ie25)*
mutants (bottom). Insets depict single oocyte nuclei, revealing 6 bivalents in both wild-type and
syp-4(ie25). (c) The *syp-4(ie25)* mutation or epitope tagging of *syp-4* alone *(ie29)* did not perturb
the length of the transition zone (solid lines in b), but this length was significantly increased in
syp-4-ha(ie27) mutants, indicative of a defect in synapsis. (d) Immunofluorescence detection of a
CHK-2-dependent phoshpoepitope provides a more consistent assay for meiotic progression. In
animals homozygous for the syp-4-HA allele *ie29,* this "CHK-2 active" zone comprises 45% of
the length of the germline containing leptotene, zygotene and pachytene stages of meiosis. This
fraction was extended to 56% and 77% in *ie25* and *ie27*, respectively. Boxplots in c and d show
mean±s.d (whiskers show minima and maxima). (e) Fluorescence micrographs reveal robust
synapsis for *ie29* (top) and *ie25* (bottom). (f) The cruciform bivalent structure, as illustrated in
the cartoon (top) is disrupted in *ie25*, likely reflecting the occurrence of multiple crossovers per

chromosome pair. Immunostaining of HTP-3 (axis; green) and SYP-4 (SC; magenta) is shown
(e-f). (g) The number of crossovers per nucleus as estimated from whole genome sequencing
(black/white boxplots, mean±s.e., whiskers are extreme values) correspond to the number of
GFP-COSA-1 foci (green dots and boxplots, mean±s.d.). (h) Fluorescence micrographs show a
marker for designated crossovers, GFP-COSA-1 (yellow), and SYP-4 (blue). (i) To measure
crossover interference strength, we fit the cumulative distribution function of inter-GFP-COSA-1
distances in *syp-4(ie25)* mutants (orange crosses) using a gamma function[44]. γ=1 denotes no
interference (gray), γ=37 corresponds to interference measured for wild-type animals[9] (purple
dashed line), while interference in *syp-4(ie25)* is severely reduced to γ=3.5 (orange line, *n*=41
chromosomes). (j) Likewise, interference strength can be described by the coefficient of
coincidence[45], which denotes the ratio of the observed number of 2 COs (GFP-COSA-1 foci)
at a given distance to the expected number assuming random CO distribution. In wild-type
animals, the coefficient of coincidence of GFP-COSA-1 foci is 0 corresponding to complete
interference (purple dashed line; *n*=88 chromosomes), while it is approximately 1 in *syp-4(ie25)*
mutants (orange, *n*=41), indicating no interference (gray line). (k) Chromosome axes lengths are
shorter in *syp-4(ie25)* mutants compared to wild-type axes in mid(-)-pachytene prior do
designation of CO sites (left) and comparable in late pachytene (right). Boxplots show mean±s.d
and whiskers are extreme values.
359
360
361

Figure 4: Irregular organization and an increase in GFP-COSA-1 foci in *syp-4(ie25)*
polycomplexes. (a) Fluorescence micrographs show that the number of GFP-COSA-1 foci
(yellow) is limited to 1 or 2 per polycomplex (SYP-2, blue) in wild-type conditions (Δ*htp-3*) and
in the presence of *syp-4(ie27)* but is elevated in *syp-4(ie25)*. DAPI is shown in gray. (b)
Quantification of GFP-COSA-1 foci. Boxplots show mean±s.d. and whiskers are extreme values.
(c) Representative electron micrographs show that the SC-like structure of polycomplexes (Δ*htp-
3*, 97 nm spacing of parallel, electron dark regions) is disturbed in Δ*htp-3 syp-4(ie25)*, which are
disorganized and 46 nm apart.

375 **Table S1: Summary of localizations of SC components determined by STORM.** Positions

376 are distances of the maxima to the center of the SC (wt = wild-type; early = early pachytene;

377 mid(-) = mid pachytene nuclei without bright GFP-COSA-1 foci; mid(+) = mid pachytene nuclei

378 with bright GFP-COSA-1 foci). HWHM denotes the half width at half maximum of the

379 distributions (Fig. 1m, gray lines)

| domain | condition | x [nm] | | z [nm] | | total length analyzed [nm] |
		position	HWHM	position	HWHM	(# images)
HA-SYP-1	wt, early	0	28.3 ± 1.8	0	33.1 ± 2.7	3438 (8)
	wt; mid(-)	0	27.5 ± 1.6	0	65.4 ± 2.8	3056 (5)
	wt; mid(+)	0	29.4 ± 2.6	0	69.7 ± 3.5	3992 (6)
	wt, late pachytene	0	27.0 ± 1.6	0	29.1 ± 3.7	4780 (9)
	zhp-3(jf61)	0	42.7 ± 2.5	0	46.9 ± 2.9	3579 (7)
	plk-2(ok1936)	0	25.5 ± 1.5	0	52.0 ± 3.9	4136 (10)
	syp-4(ie25); early	0	22.1 ± 1.4	0	28.6 ± 1.2	4351 (10)
	syp-4(ie25); late	0	19.8 ± 0.8	0	20.5 ± 1.6	3114 (6)
SYP-1(C)	wt, early	42.3 ± 0.8	22.8 ± 0.9	0	37.1 ± 2.3	4092 (7)
	wt; mid(-)	40.8 ± 1.3	28.2 ± 0.6	0	68.1 ± 4.3	3184 (7)
	wt; mid(+)	35.9 ± 0.9	31.2 ± 1.0	0	78.7 ± 2.8	3503 (7)
	wt, late pachytene	42.0 ± 1.2	28.5 ± 1.0	0	62.7 ± 3.2	5312 (9)
	zhp-3(jf61); late	41.0 ± 1.3	23.6 ± 1.1	0	48.2 ± 4.3	1851 (8)
	plk-2(ok1936); late	36.3 ± 1.4	39.6 ± 3.6	0	67.0 ± 2.5	5741 (9)
	syp-4(ie25);	38.5 ± 2.5	24.9 ± 0.8	0	43.2 ± 3.8	2768 (5)

	early					
	syp-4(ie25); late	30.5 ± 1.5	29.1 ± 1.8	0	54.0 ± 2.2	4848 (6)
SYP-2(C)	wt, early	0	27.0 ± 2.0	0	38.3 ± 3.7	5866 (10)
	wt; mid(-)	16.3 ± 7.5	27.1 ± 2.7	0	27.6 ± 1.1	8437 (9)
	wt; mid(+)	20.6 ± 6.3	26.5 ± 3.6	0	27.7 ± 1.4	8615 (12)
	wt, late pachytene	18.9 ± 2.1	21.6 ± 1.5	0	49.5 ± 5.0	4573 (9)
	wt, late (HA)	20.9 ± 0.9	20.0 ± 1.3	0	39.9 ± 2.6	2958 (8)
	zhp-3(jf61); late	0	36.1 ± 4.3	0	38.6 ± 3.8	3746 (6)
	plk-2(ok1936); late	0	37.1 ± 2.0	0	47.4 ± 2.3	8206 (9)
	syp-4(ie25); early	0	32.4 ± 2.2	0	73.6 ± 8.8	3172 (7)
	syp-4(ie25); late	0	28.9 ± 2.7	0	33.4 ± 3.2	3604 (8)
GFP-SYP-3	wt, early	0	23.3 ± 0.8	0	40.0 ± 4.5	1784 (4)
	wt, late pachytene	0	18.5 ± 1.4	0	63.2 ± 6.2	2238 (6)
SYP-3-HA	wt, early	12.3 ± 1.8	20.3 ± 1.2	0	27.1 ± 2.4	4653 (10)
	wt, late pachytene	16.6 ± 4.7	29.0 ± 2.3	0	48.1 ± 5.9	2764 (7)
	zhp-3(jf61); late	14.2 ± 2.0	24.4 ± 1.4	0	28.5 ± 2.2	2723 (6)
SYP-4-intFlag	wt, early	0	25.8 ± 0.9	0	43.3 ± 5.3	3462 (9)
	wt, late pachytene	0	30.3 ± 4.3	0	49.8 ± 4.6	2524 (4)
SYP-4-HA	wt, early	14.2 ± 2.0	23.0 ± 1.6	0	38.6 ± 2.4	5355 (10)

	wt, late pachytene	0	36.8 ± 1.1	33.1 ± 11.9	51.5 ± 8.0	5264 (9)
SYP-4(*ie25*)	syp-4(ie25); late	0	29.1 ± 1.1	0	54.9 ± 5.6	4611 (9)
HIM-3	wt, early	33.5 ± 5.2	43.4 ± 2.3	n.d.	n.d.	2548 (6)
	wt, mid(-)	41.4 ± 2.2	46.7 ± 1.8	n.d.	n.d.	3661 (7)
	wt, mid(+)	41.8 ± 1.9	45.1 ± 1.2	n.d.	n.d.	2817 (7)
	wt, late pachytene	47.3 ± 1.5	30.2 ± 1.2	n.d.	n.d.	4411 (11)
	zhp-3(jf61); late	45.9 ± 1.6	49.0 ± 2.3	n.d.	n.d.	6969 (11)
	plk-2(ok1936); late	41.5 ± 3.8	53.9 ± 4.1	n.d.	n.d.	4769 (8)
	syp-4(ie25); early	45.3 ± 3.6	48.9 ± 2.7	n.d.	n.d.	3900 (7)
	syp-4(ie25); late	35.8 ± 3.8	45.9 ± 3.1	n.d.	n.d.	2926 (6)

385 **Table S2: List of Strains used in this study and their characterization.** *p*-Values for deviation

386 from wild-type are calculated using Mann-Whitney tests.

name	genotype	egg viability [%] (*p*-value)	males [%] (*p*-value)	# of eggs
N2	*wild-type*	104±6	0.08±0.20	4773
GFP-COSA-1 [36]	*meIs8* II	107±4	0.00±0.00	1362
mEos-HIM-3 [24]	*meIs8* II; *him-3(ie33)* IV	n.d.	n.d.	
mMaple-HIM-3 [24]	*meIs8* II; *him-3(ie34)* IV	n.d.	n.d.	
HA-SYP-1	*meIs8* II; *him-3(ie33)* IV; *syp-1(ie40[syp-1(1-47)::HA::syp-1(48-489)])* V	100±4 (0.886)	0.00±0.00 (0.452)	840
SYP-2-HA	*meIs8* II; *him-3(ie33)* IV; *syp-2(ie99[syp-2::HA])* V	101±6 (0.394)	0.00±0.00 (1.00)	1405
GFP-SYP-3	*syp-3(ok758)* I; *ieSi11 [syp-3p::syp-3::EmGFP::syp-3-3'UTR + unc-119(+)]* II; *him-3(ie33)* IV	83±10 (0.031)	0.87±0.59 (0.045)	1143
SYP-3-HA	*syp-3(ie42[syp-3::HA])* I; *meIs8* II; *him-3(ie33)* IV	95±4 (0.041)	1.80±0.60 (0.004)	1457
SYP-4-HA(*ie29*)	*syp-4(ie29[syp-4::HA])* I; *meIs8* II; *him-3(ie34)* IV	100±2 (0.150)	0.72±0.41 (0.052)	1336
SYP-4-HA(*ie29*)	*syp-4(ie29[syp-4::HA])* I; *meIs8* II	105±3 (0.686)	0.00±0.00 (1.000)	951
SYP-4-intFlag	*syp-4(ie30[syp-4(1-315)::2xFlag::syp-4(319-605)])* I; *meIs8* II; *him-3(ie33)* IV	98±3 (0.032)	1.91±0.59 (0.010)	1350
SYP-4(*ie25*)	*syp-4(ie25[syp-4ΔC::3xFlag])* I; *meIs8* II	38±4 (0.008)	5.97±1.25 (0.010)	1024
SYP-4-HA(*ie27*)	*syp-4(ie29[syp-4::HA::mod3'UTR])* I; *meIs8* II	80±9 (0.002)	1.58±1.58 (0.010)	992
Δ*htp-3* [3]	*htp-3(tm3655)* I; *meIs8* II	n.d.	n.d.	
Δ*htp-3 syp-4(ie25)*	*htp-3(ie100) syp-4(ie25)* I; *meIs8* II	n.d.	n.d.	

Δhtp-3 syp-4(ie27)	*htp-3(ie101) syp-4(ie27)* I; *meIs8* II	n.d.	n.d.	
SYP-4(ie25)	*syp-4(ie25[syp-4ΔC::3xFlag])* I; *meIs8* II; *him-3(ie33)* IV	n.d.	n.d.	
SYP-4(ie25); HA-SYP-1	*syp-4(ie25)* I; *meIs8* II; *him-3(ie33)* IV; *syp-1(ie40)* V	n.d.	n.d.	
zhp-3(jf61)	*zhp-3(jf61)* I; *meIs8* II; *him-3(ie33)* IV	n.d.	n.d.	
zhp-3(jf61); HA-SYP-1	*zhp-3(jf61)* I; *meIs8* II; *him-3(ie33)* IV; *syp-1(ie40)* V	n.d.	n.d.	
zhp-3(jf61) SYP-3-HA	*syp-3(ie102[syp-3::HA]) zhp-3(jf61)* I; *meIs8* II; *him-3(ie33)* IV	n.d.	n.d.	
plk-2(ok1936); HA-SYP-1	*plk-2(ok1936)* I; *meIs8* II; *him-3(ie33)* IV; *syp-1(ie40)* V	n.d.	n.d.	

388
389

390
Extended Data Figure S1: Experimental strategy for mapping the architecture of the

synaptonemal complex. (a) The resolution in conventional fluorescence micrographs is

insufficient to visualize the structure of the axis or SC (HIM-3, magenta, left) and SC

components (SYP-2-HA, green, center) are co-localized (merge, right). (b) The structure of the

SC is resolved by STORM super-resolution microscopy. Colors denote localization in z (left and

center). Localization of HIM-3 (magenta, right) is used to determine the orientation of the SC

(SYP-2-HA(C), wild-type, late pachytene in green). White boxes in b show regions magnified in

other panels. (c) For further analysis, individual stretches in frontal view (right), which are

characterized by clearly separated HIM-3 axes, are straightened and aligned in x and z to

generate the averaged image (left). (d) The results are highly reproducible between different

preparations and different antibodies (SYP-2(C) peptide antibody, black, and SYP-2-HA epitope

antibody, green) (Table S2). Note that the widths of the distributions are narrower for the

monoclonal HA antibody compared to the polyclonal SYP-2(C) peptide antibody.

Extended Figure S2: Dynamic orientation of SYP-1 molecules. (a) To generate a model of SYP orientation within the SC, individual datapoints (magenta circle, center) in the measured distributions of the N- (blue, left) and C-terminus (gray, left) of a given component (here: SYP-1 in wild-type late pachytene) are randomly mapped to a localization event within the same percentile ±7.5% tolerance of the C-terminal distribution in *x* and *z* (purple, center). Mapped events resulting in the shortest or longest 5% of antibody to antibody distances (red, right) are disregarded. (b) The boxplots show means±s.d. of the distances between N- and C-terminal antibodies as estimated by the probabilistic mapping approach. Whiskers are extremes and the black line shows the mean across all genotypes and conditions. (c) The resultant model of SC organization generated by mapping of N- and C-terminal distributions of SYP proteins is shown for early (left) and late (right) pachytene in wild-type animals. C-termini are cubes and N-termini

406
407

418   (central domain for SYP-4) are spheres. The localization of HIM-3 in x is used as a proxy for

419   axis localization. Models are rendered using POV-ray (v3.7.0).

Extended Data Figure S3: Dynamic organization of SYP-3 and SYP-4 within the SC. (a-b) Averaged STORM images show cross-sectional views of SYP-3 (a) and SYP-4 (b) domains as indicated in wild type at early (blue) and late pachytene (purple) and *zhp-3(jf61)* (orange) and *syp-4(ie25)* (green) mutants in late pachytene. Colors denote localization in *z*. The distributions of localization events in frontal (*x*), and vertical (*z*) views are analyzed in (c). The results are summarized in Extended Data Table S2.

434
Extended Data Figure S4: Double-strand break formation is extended in *syp-4(ie25)*. (a)
Immunofluorescence images show DAPI (upper) and RAD-51 (lower), which marks double-
strand breaks. While the RAD-51 zone (gray line) is normally limited to a short region in
early/mid-pachytene in wild-type (top), it is markedly elongated in *syp-4(ie25)* (bottom). (b)
shows the average number of RAD-51 foci per nucleus (boxes are mean±s.e.) as a function of the
relative length of the gonad from the distal tip to the end of pachynema of wild-type (purple solid
line, *n*=1672 nuclei from (7) gonads), *syp-4(ie29)* (red dashed line, *n*=1407 (6)), *syp-4(ie25)*
(green dot-dashed line, *n*=1440 (6)) and *syp-4(ie27)* (blue dotted line, *n*=1162 (4)), respectively.

Extended Data Figure S5: Abundance of SC proteins in *syp-4(ie25)*. (a) The expression level

of HA-SYP-1 (magenta) is identical in wild-type and *syp-4(ie25)* strains as shown by a Western

Blot. Tubulin (green) is shown as a reference. (b) By contrast, the levels of immunofluorescence

staining for SYP-1 (left) and SYP-2 (center) are drastically reduced in *syp-4(ie25)* SCs (bottom)

compared to wild-type (*syp-4-ha(ie29)*, right). To ensure identical handling, gonads from both

genotypes were stained in a single reaction and their genotype was determined by anti-Flag

staining (right), which marks *syp-4(ie25)* but not *syp-4(ie29)*. (c) Boxplots show mean±s.d.

(whiskers are extreme values) of fluorescence intensities of late pachytene SCs (*ie29*: *n*=339,

ie25: *n*=209).

 Extended Data Figure S6: The *syp-4-ha(ie27)* **mutation phenocopies partial RNAi of other**

SC components. (a) The mutation *syp-4-ha(ie27)* (bottom) is constituted by an insertion (cyan)

in the 3' untranslated region (lower case letters) of the *syp-4* gene, but the coding sequence

(capital letters) is identical to *syp-4-ha(ie29)* (top), which is phenotypically wild-type. (b) A

DAPI stained fluorescence micrographs show that meiotic progression is delayed in *syp-4-*

ha(ie27) as indicated by an elongated transition zone (gray line) but can form 6 bivalents (inset)

indicating largely successful homologous pairing and crossover formation. (c) A western blot

shows that the level of SYP-4-HA (top) is reduced to ~ 30 % in *syp-4-ha(ie27)* compared to *syp-*

4-ha(ie29). Tubulin is shown as a reference (bottom). (d) Synapsis is delayed in *syp-4-ha(ie27)*

as shown by the incomplete overlap of immunofluorescence images of an axial protein (HTP-3,

green) and SC proteins (SYP-4-HA, magenta). (e) As has been described for partial RNAi

depletion of SYP-1, -2 and -3[9], a reduction in SYP-4 expression level results in a slight

increase in the number of GFP-COSA-1 (yellow) foci to about 7 foci/nucleus (DAPI is shown in

blue).

Methods

Worm strains and transgenes

A complete list of *C. elegans* strains used in this study can be found in Table S1. All strains were cultured at 20ºC using standard methods[52]. GFP-SYP-3[32] was inserted by MosSCI[53] and crossed into *syp-3(ok758)*. Other tags were inserted using CRISPR-Cas9 genome editing as described in[24]. Briefly, Cas9 and gRNA were delivered by microinjection, either encoded on a plasmid[54] or as *in vitro* preassembled Cas9-ribonucleoprotein complexes. Repair templates for small epitope tags were codon optimized for *C. elegans* [55] and synthesized as "Ultramers" by IDT.

Immunofluorescence

Immunofluorescence of dissected gonads from young adults 24 h post L4 was performed as described[56] with modifications described in[24]. The following primary antibodies, all of which have been previously described, were used: goat anti-SYP-1 (1:500, affinity purified)[37], mouse anti-HA (1:500, monoclonal 2-2.2.14, Thermo Fisher Scientific), rabbit anti-SYP-2 (1:500, affinity purified)[16], mouse anti-GFP (1:500, monoclonal 7.1 and 13.1, Roche), mouse anti-Flag (1:500, monoclonal M2, Sigma-Aldrich), chicken anti-HTP-3 (1:500)[57], rabbit anti-HIM-3 (1:500, SDQ4713 ModENCODE project)[58], rabbit anti-HIM-8pT64 (1:2000)[41], and rabbit anti-RAD-51 (1:500)[37]. Secondary antibodies raised in donkeys were fluorescently labelled with Alexa Fluor® 488, Cy3 or Alexa Fluor® 647 (1:500, Jackson ImmunoResearch and Invitrogen). Gonads were mounted in ProLong Gold antifade mountant (ThermoFisher Scientific) and epifluorescence images were acquired on a DeltaVision Elite microscope (Applied Precision) using a 100× N.A. 1.4 oil-immersion objective.

For quantification of proteins levels by immunofluorescence, imaging was performed on a Marianas spinning-disk confocal microscope (3i) with a 100x 1.46 NA oil immersion objective. 3D stacks of wild-type (*syp-4-ha(ie29)*) and *syp-4(ie25)* animals were taken from the same slide and quantified in Fiji. SC containing voxels are identified by thresholding the SYP-2 and SYP-4-3xFlag channels for *syp-4(ie29)* and *syp-4(ie25)*, respectively. Images shown in Fig. S6 are background subtracted maximal intensity projections with equally scaled intensity values.

STORM and PALM imaging

Super-resolution imaging of dissected, immuno-stained intact germline tissue was carried out as

described[24]. Targets for STORM were labeled with donkey or goat secondary antibodies

labelled with Alexa Fluor® 647 (1:500, Jackson ImmunoResearch and Invitrogen).

Subsequently, the fluorescently tagged internal reference protein, mEos2-HIM-3 or mMaple3-

HIM-3[24] was imaged using PALM [59] [60]. For early pachytene images, mEos2-HIM-3 was

co-stained with a rabbit anti-HIM-3 antibody and a donkey-anti-rabbit secondary antibody

(Jackson ImmunoResearch) labelled with NHS esters of CF568 to achieve a 2:1 dye-to-antibody

ratio. Aligned and averaged images (Fig. S1) were used to generate histograms of localization

events in x and z[24]. To systematically distinguish between mono- and bi-modal distributions,

we evaluated fits with one and two Gaussians using an ANOVA test in R ($p < 0.05$). Standard

deviations of fit parameters were estimated by a subsampling approach using subsets of half the

number of individual SC stretches[24]. The results are summarized in Table S2.

Statistical analysis

Sample sizes were not predetermined, and experiments were not randomized. The investigators

were not blinded to allocation during experiments and outcome assessment. For STORM

experiments, the total lengths and number of stretches analyzed are summarized in Table S2.

Analysis of of SC organization

To derive a model for SC organization from our data, we asked which orientations of SC

proteins were consistent with the observed distributions, assuming that each protein is a rigid

rod. We first corrected the raw data by removing the extreme 2.5% of the localization events

using squared Mahalanobis distances in R (version 3.4.2). We then mapped each localization

event corresponding the nth-percentile in x and the mth-percentile in z of the N- (or C-) terminal

distribution to a randomly selected localization event within the nth ± 7.5% in x and mth ± 7.5%

in z of the corresponding C- (or N-) terminal distribution. This analysis reveals the distance

between the N- and C-terminal antibodies used for imaging and the orientation of the protein

within the SC. For further analysis, we remove outliers among the mapped localization events

with extreme values corresponding to the lower and upper 2.5% of distances.

To visualize changes within SC organization, we performed principal component analysis of the

widths, and – where applicable – positions off-center, for N- and C-terminal distributions of

SYP-1 and the C-terminal distribution of SYP-2, the orientations of SYP-1 molecules as

characterized by the median and 90th percentile angles, and the average distance between N- and

C-terminal antibodies using R (version 3.4.2). We excluded SYP-3 and SYP-4 from this analysis

since experimental challenges prevented us from analyzing their localizations for all genotypes.

Recombination mapping

To map meiotic recombination events, the *syp-4(ie25)* allele (generated in Bristol N2

background) was introgressed into the divergent Hawaiian strain CB4856 by 8 sequential crosses

and selection for *ie25*. Hawaiian and Bristol wild-type and *syp-4(ie25)* strains were crossed, and

the hybrid F1 progeny were backcrossed to Hawaiian males. These cross-progeny, which have

Hawaiian paternal chromosomes and recombined Bristol/Hawaiian maternal chromosomes, were

plated individually, allowed to reproduce for one generation, and the genomic DNA of their

pooled progeny was extracted by phenol-chloroform extraction. Illumina sequencing libraries

were prepared as described in [61] and sequenced as 50 bp single reads on a HiSeq2500 Illumina

sequencer at the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley, with a

coverage of 2-4x per individual. Reads were mapped to the genomes of Bristol (Wormbase

release WS230) and Hawaiian[62] strains and genotypes called using the multiplexed shotgun

genotyping (MSG) toolbox and default parameters[63]. Aligned sequences obtained by deep

sequencing have been uploaded on the NCBI SRA database under accession number

SRP126693. Following this analysis, crossovers in oogenesis are characterized by transitions

from Hawaiian/Bristol heterozygous stretches to homozygous Hawaiian stretches, and the

calculated number of crossovers per nucleus is twice the number of mapped crossovers in the

oocytes.

Electron microscopy

High-pressure freezing, freeze-substitution, sample preparation, and microscopy was performed

as described previously[3, 64, 65]. Images were acquired on a Tecnai 12 transmission electron

microscope (120 kV, FEI, Hillsboro, OR) equipped with a Gatan Ultrascan 1000 CCD camera

(Pleasanton, CA).

 Immunoblotting

To compare expression levels of SYP proteins, 120 adult worms were lysed by boiling in 40 µL

Laemmli sample buffer (with β-mercaptoethanol) for about 5 min, until particulate matter was

not detected using a dissection stereomicroscope. Samples of whole-worm lysates were run on a

Nupage 4-12% polyacrylamide gradient gel and transferred to a PVDF membrane. Primary

antibodies were rabbit (Pierce, PA1-985, used for HA-SYP-1) or mouse anti-HA (SYP-4-HA)

and mouse anti-tubulin (EMD Millipore), each diluted 1:5,000. HRP-conjugated secondary

antibodies (Jackson Laboratory) were detected with ECL reagents (Amersham). For SYP-4-HA

blots, HA and tubulin were easily resolved and can be detected simultaneously. For HA-SYP-1,

HA was detected by HRP-anti-rabbit/ECL and tubulin is detected by Cy3-conjugated anti-mouse

secondary antibody (1:5,000). Images were recorded using a Chemidoc system (Bio-Rad) and

quantified using Fiji.

Data Availability

NGS datasets generated during the current study are available on the NCBI SRA database under

accession number SRP126693 (https://www.ncbi.nlm.nih.gov/sra/SRP126693). Any additional

data are available upon reasonable request.

Acknowledgements

We thank Lydia Smith and Rasmus Nielsen for assistance with recombination mapping, Kent

McDonald and Reena Zalpuri for help with high-pressure freezing and electron microscopy and

Ofer Rog and current members of our laboratories for helpful discussions and critical reading of

the manuscript. Some nematode strains used in this work were provided by the Caenorhabditis

Genetics Center, which is supported by the NIH - Office of Research Infrastructure Programs

(P40 OD010440). This work was funded by a postdoctoral fellowship of the Human Frontier

Science Program to SK (LT000903/2013-C), an NSF Graduate Research Fellowship (DGE

1106400) to MW, the Pew Biomedical Scholars Award and the Chan Zuckerberg Biohub to KX,

and support to AFD from the National Institutes of Health (R01 GM065591) and the Howard

Hughes Medical Institute.

Competing Interests

The authors declare no competing interests.

Author Contributions

SK and MW performed experiments and analyzed data. SK, MW, KX and AFD designed

experiments and wrote the manuscript.

References

1. Zickler D, Kleckner N (1999) Meiotic chromosomes: integrating structure and function.

Annu Rev Genet 33:603–754. doi: 10.1146/annurev.genet.33.1.603

2. Moses MJ (1968) Synaptinemal complex. Annu Rev Genet 2:363–412.

3. Rog O, Köhler S, Dernburg AF (2017) The synaptonemal complex has liquid crystalline

properties and spatially regulates meiotic recombination factors. Elife 6:e21455. doi:

10.7554/eLife.21455

4. Voelkel-Meiman K, Moustafa SS, Lefrançois P, et al (2012) Full-length synaptonemal
complex grows continuously during meiotic prophase in budding yeast. PLoS Genet
8:e1002993. doi: 10.1371/journal.pgen.1002993

5. Pattabiraman D, Roelens B, Woglar A, Villeneuve AM (2017) Meiotic recombination
modulates the structure and dynamics of the synaptonemal complex during C. elegans
meiosis. PLoS Genet 13:e1006670. doi: 10.1371/journal.pgen.1006670

6. Machovina TS, Mainpal R, Daryabeigi A, et al (2016) A Surveillance System Ensures
Crossover Formation in C. elegans. Curr Biol 26:2873–2884. doi: 10.1016/j.cub.2016.09.007

7. Nadarajan S, Lambert TJ, Altendorfer E, et al (2017) Polo-like kinase-dependent
phosphorylation of the synaptonemal complex protein SYP-4 regulates double-strand break
formation through a negative feedback loop. Elife 6:e23437. doi: 10.7554/eLife.23437

8. Hayashi M, Mlynarczyk-Evans S, Villeneuve AM (2010) The synaptonemal complex shapes
the crossover landscape through cooperative assembly, crossover promotion and crossover
inhibition during Caenorhabditis elegans meiosis. Genetics 186:45–58. doi:
10.1534/genetics.110.115501

9. Libuda DE, Uzawa S, Meyer BJ, Villeneuve AM (2013) Meiotic chromosome structures
constrain and respond to designation of crossover sites. Nature 502:703–706. doi:
10.1038/nature12577

10. Wang K, Wang C, Liu Q, et al (2015) Increasing the Genetic Recombination Frequency by
Partial Loss of Function of the Synaptonemal Complex in Rice. Mol Plant 8:1295–1298. doi:
10.1016/j.molp.2015.04.011

11. Woglar A, Villeneuve AM (2018) Dynamic Architecture of DNA Repair Complexes and the
Synaptonemal Complex at Sites of Meiotic Recombination. Cell 173:1678–1691.e16. doi:
10.1016/j.cell.2018.03.066

12. Cahoon CK, Helm JM, Libuda DE (2019) Synaptonemal complex proteins direct and
constrain the localization of crossover-promoting proteins during Caenorhabditis elegans
meiosis. bioRxiv 523605. doi: 10.1101/523605

13. Zhang L, Köhler S, Rillo-Bohn R, Dernburg AF (2018) A compartmentalized signaling
network mediates crossover control in meiosis. Elife 7:245. doi: 10.7554/eLife.30789

14. Dernburg AF, McDonald K, Moulder G, et al (1998) Meiotic recombination in C. elegans
initiates by a conserved mechanism and is dispensable for homologous chromosome
synapsis. Cell 94:387–398.

15. MacQueen AJ, Colaiácovo MP, McDonald K, Villeneuve AM (2002) Synapsis-dependent
and -independent mechanisms stabilize homolog pairing during meiotic prophase in C.
elegans. Genes Dev 16:2428–2442. doi: 10.1101/gad.1011602

16. Colaiácovo MP, MacQueen AJ, Martinez-Perez E, et al (2003) Synaptonemal complex
assembly in C. elegans is dispensable for loading strand-exchange proteins but critical for
proper completion of recombination. Dev Cell 5:463–474. doi: 10.1016/S1534-
5807(03)00232-6

17. Smolikov S, Eizinger A, Schild-Prüfert K, et al (2007) SYP-3 restricts synaptonemal
complex assembly to bridge paired chromosome axes during meiosis in Caenorhabditis
elegans. Genetics 176:2015–2025. doi: 10.1534/genetics.107.072413

18. Smolikov S, Schild-Prüfert K, Colaiácovo MP (2009) A yeast two-hybrid screen for SYP-3
interactors identifies SYP-4, a component required for synaptonemal complex assembly and
chiasma formation in Caenorhabditis elegans meiosis. PLoS Genet 5:e1000669. doi:
10.1371/journal.pgen.1000669

19. Schild-Prüfert K, Saito TT, Smolikov S, et al (2011) Organization of the synaptonemal
complex during meiosis in Caenorhabditis elegans. Genetics 189:411–421. doi:
10.1534/genetics.111.132431

20. Schücker K, Holm T, Franke C, et al (2015) Elucidation of synaptonemal complex
organization by super-resolution imaging with isotropic resolution. Proc Natl Acad Sci
112:2029–2033. doi: 10.1073/pnas.1414814112

21. Cahoon CK, Yu Z, Wang Y, et al (2017) Superresolution expansion microscopy reveals the
three-dimensional organization of theDrosophilasynaptonemal complex. Proc Natl Acad Sci
114:E6857–E6866. doi: 10.1073/pnas.1705623114

22. Rust MJ, Bates M, Zhuang X (2006) Sub-diffraction-limit imaging by stochastic optical
reconstruction microscopy (STORM). Nat Meth 3:793–795. doi: 10.1038/nmeth929

23. Huang B, Wang W, Bates M, Zhuang X (2008) Three-dimensional super-resolution imaging
by stochastic optical reconstruction microscopy. Science 319:810–813. doi:
10.1126/science.1153529

24. Köhler S, Wojcik M, Xu K, Dernburg AF (2017) Superresolution microscopy reveals the
three-dimensional organization of meiotic chromosome axes in intact Caenorhabditis elegans
tissue. Proc Natl Acad Sci 114:E4734–E4743. doi: 10.1073/pnas.1702312114
25. Zetka MC, Kawasaki I, Strome S, Müller F (1999) Synapsis and chiasma formation in
Caenorhabditis elegans require HIM-3, a meiotic chromosome core component that functions
in chromosome segregation. Genes Dev 13:2258–2270.
26. Gao J, Barroso C, Zhang P, et al (2016) N-terminal acetylation promotes synaptonemal
complex assembly in C. elegans. Genes Dev 30:2404–2416. doi: 10.1101/gad.277350.116
27. Schmekel K, Meuwissen RL, Dietrich AJ, et al (1996) Organization of SCP1 protein
molecules within synaptonemal complexes of the rat. Exp Cell Res 226:20–30. doi:
10.1006/excr.1996.0198
28. Sym M, Roeder GS (1995) Zip1-induced changes in synaptonemal complex structure and
polycomplex assembly. J Cell Biol 128:455–466.
29. Higgins JD, Sanchez-Moran E, Armstrong SJ, et al (2005) The Arabidopsis synaptonemal
complex protein ZYP1 is required for chromosome synapsis and normal fidelity of crossing
over. Genes Dev 19:2488–2500. doi: 10.1101/gad.354705
30. Meuwissen RL, Offenberg HH, Dietrich AJ, et al (1992) A coiled-coil related protein
specific for synapsed regions of meiotic prophase chromosomes. EMBO J 11:5091–5100.
31. Anderson LK, Royer SM, Page SL, et al (2005) Juxtaposition of C(2)M and the transverse
filament protein C(3)G within the central region of Drosophila synaptonemal complex. Proc
Natl Acad Sci USA 102:4482–4487. doi: 10.1073/pnas.0500172102
32. Rog O, Dernburg AF (2015) Direct Visualization Reveals Kinetics of Meiotic Chromosome
Synapsis. Cell Rep 10:1639–1645. doi: 10.1016/j.celrep.2015.02.032
33. Reynolds A, Qiao H, Yang Y, et al (2013) RNF212 is a dosage-sensitive regulator of
crossing-over during mammalian meiosis. Nat Genet 45:269–278. doi: 10.1038/ng.2541
34. Qiao H, Prasada Rao HBD, Yang Y, et al (2014) Antagonistic roles of ubiquitin ligase HEI10
and SUMO ligase RNF212 regulate meiotic recombination. Nat Genet 46:194–199. doi:
10.1038/ng.2858
35. de Gennes PG, Prost J (1993) The Physics of Liquid Crystals, 2nd ed. Clarendon Press,
Oxford

36. Yokoo R, Zawadzki KA, Nabeshima K, et al (2012) COSA-1 reveals robust homeostasis and

separable licensing and reinforcement steps governing meiotic crossovers. Cell 149:75–87.

doi: 10.1016/j.cell.2012.01.052

37. Harper NC, Rillo R, Jover-Gil S, et al (2011) Pairing centers recruit a Polo-like kinase to

orchestrate meiotic chromosome dynamics in C. elegans. Dev Cell 21:934–947. doi:

10.1016/j.devcel.2011.09.001

38. Sato-Carlton A, Nakamura-Tabuchi C, Chartrand SK, et al (2018) Phosphorylation of the

synaptonemal complex protein SYP-1 promotes meiotic chromosome segregation. The

Journal of Cell Biology 217:555–570. doi: 10.1083/jcb.201707161

39. Jantsch V, Pasierbek P, Mueller MM, et al (2004) Targeted gene knockout reveals a role in

meiotic recombination for ZHP-3, a Zip3-related protein in Caenorhabditis elegans. Mol Cell

Biol 24:7998–8006. doi: 10.1128/MCB.24.18.7998-8006.2004

40. Labella S, Woglar A, Jantsch V, Zetka M (2011) Polo kinases establish links between

meiotic chromosomes and cytoskeletal forces essential for homolog pairing. Dev Cell

21:948–958. doi: 10.1016/j.devcel.2011.07.011

41. Kim Y, Kostow N, Demburg AF (2015) The Chromosome Axis Mediates Feedback Control

of CHK-2 to Ensure Crossover Formation in C. elegans. Dev Cell 35:247–261. doi:

10.1016/j.devcel.2015.09.021

42. Hillers KJ, Villeneuve AM (2003) Chromosome-wide control of meiotic crossing over in C.

elegans. Curr Biol 13:1641–1647. doi: 10.1038/nrg1210

43. Nabeshima K, Villeneuve AM, Colaiácovo MP (2005) Crossing over is coupled to late

meiotic prophase bivalent differentiation through asymmetric disassembly of the SC. J Cell

Biol 168:683–689. doi: 10.1083/jcb.200410144

44. Broman KW, Weber JL (2000) Characterization of human crossover interference. Am J Hum

Genet 66:1911–1926. doi: 10.1086/302923

45. Muller HJ (1916) The mechanism of crossing-over IV. American Naturalist 50:421–434.

46. Drouaud J, Mercier R, Chelysheva L, et al (2007) Sex-specific crossover distributions and

variations in interference level along Arabidopsis thaliana chromosome 4. PLoS Genet

3:e106. doi: 10.1371/journal.pgen.0030106

47. Petkov PM, Broman KW, Szatkiewicz JP, Paigen K (2007) Crossover interference underlies

sex differences in recombination rates. Trends Genet 23:539–542. doi:

10.1016/j.tig.2007.08.015

48. Mets DG, Meyer BJ (2009) Condensins Regulate Meiotic DNA Break Distribution, thus

Crossover Frequency, by Controlling Chromosome Structure. Cell 139:73–86. doi:

10.1016/j.cell.2009.07.035

49. Zhang L, Liang Z, Hutchinson J, Kleckner N (2014) Crossover Patterning by the Beam-Film

Model: Analysis and Implications. PLoS Genet 10:e1004042. doi:

10.1371/journal.pgen.1004042

50. Goodyer W, Kaitna S, Couteau F, et al (2008) HTP-3 links DSB formation with homolog

pairing and crossing over during C. elegans meiosis. Dev Cell 14:263–274. doi:

10.1016/j.devcel.2007.11.016

51. Kleckner N, Zickler D, Jones GH, et al (2004) A mechanical basis for chromosome function.

Proc Natl Acad Sci USA 101:12592–12597. doi: 10.1073/pnas.0402724101

52. Brenner S (1974) The genetics of Caenorhabditis elegans. Genetics 77:71–94.

53. Frøkjær-Jensen C, Davis MW, Hopkins CE, et al (2008) Single-copy insertion of transgenes

in Caenorhabditis elegans. Nat Genet 40:1375–1383. doi: 10.1038/ng.248

54. Dickinson DJ, Ward JD, Reiner DJ, Goldstein B (2013) Engineering the Caenorhabditis

elegans genome using Cas9-triggered homologous recombination. Nat Meth 10:1028–1034.

doi: 10.1038/nmeth.2641

55. Redemann S, Schloissnig S, Ernst S, et al (2011) Codon adaptation-based control of protein

expression in C. elegans. Nat Meth 8:250–252. doi: 10.1038/nmeth.1565

56. Phillips CM, McDonald KL, Dernburg AF (2009) Cytological analysis of meiosis in

Caenorhabditis elegans. Methods Mol Biol 558:171–195. doi: 10.1007/978-1-60761-103-

5_11

57. Macqueen AJ, Phillips CM, Bhalla N, et al (2005) Chromosome sites play dual roles to

establish homologous synapsis during meiosis in C-elegans. Cell 123:1037–1050. doi:

10.1016/j.cell.2005.09.034

58. Gerstein MB, Lu ZJ, Van Nostrand EL, et al (2010) Integrative analysis of the

Caenorhabditis elegans genome by the modENCODE project. Science 330:1775–1787. doi:

10.1126/science.1196914

59. McKinney SA, Murphy CS, Hazelwood KL, et al (2009) A bright and photostable

photoconvertible fluorescent protein for fusion tags. Nat Meth 6:131–133. doi:

10.1038/nmeth.1296

60. Wang S, Moffitt JR, Dempsey GT, et al (2014) Characterization and development of

photoactivatable fluorescent proteins for single-molecule-based superresolution imaging.

Proc Natl Acad Sci 111:8452–8457. doi: 10.1073/pnas.1406593111

61. Meyer M, Kircher M (2010) Illumina sequencing library preparation for highly multiplexed

target capture and sequencing. Cold Spring Harb Protoc 2010:pdb.prot5448. doi:

10.1101/pdb.prot5448

62. Thompson OA, Snoek LB, Nijveen H, et al (2015) Remarkably Divergent Regions Punctuate

the Genome Assembly of the Caenorhabditis elegans Hawaiian Strain CB4856. Genetics

200:975–989. doi: 10.1534/genetics.115.175950

63. Andolfatto P, Davison D, Erezyilmaz D, et al (2011) Multiplexed shotgun genotyping for

rapid and efficient genetic mapping. Genome Res 21:610–617. doi: 10.1101/gr.115402.110

64. McDonald KL, Webb RI (2011) Freeze substitution in 3 hours or less. J Microsc 243:227–

233. doi: 10.1111/j.1365-2818.2011.03526.x

65. McDonald KL (2014) Rapid embedding methods into epoxy and LR White resins for

morphological and immunological analysis of cryofixed biological specimens. Microsc

Microanal 20:152–163. doi: 10.1017/S1431927613013846