Convergent network effects along the axis of gene expression during prostate cancer progression

Konstantina Charmpi[1,2,#], Tiannan Guo[3,4,#,*], Qing Zhong[5,#], Ulrich Wagner[5,#], Rui Sun[4], Nora C. Toussaint[5,6,7], Christine E. Fritz[5], Chunhui Yuan[4], Hao Chen[4], Niels J. Rupp[5], Ailsa Christiansen[5], Dorothea Rutishauser[5], Jan H. Rüschoff[5], Christian Fankhauser[5], Karim Saba[5,8], Cedric Poyet[8], Thomas Hermanns[8], Kathrin Oehl[5], Ariane L. Moore[9], Christian Beisel[9], Laurence Calzone[10], Loredana Martignetti[10], Qiushi Zhang[4], Yi Zhu[3,4], María Rodríguez Martínez[11], Matteo Manica[11], Michael C. Haffner[12], Ruedi Aebersold[3,13,*], Peter J. Wild[5,14,*], Andreas Beyer[1,2,*]

[1] CECAD, University of Cologne, Cologne, Germany
[2] Center for Molecular Medicine Cologne (CMMC), Medical Faculty, University of Cologne, Cologne, Germany
[3] Department of Biology, Institute of Molecular Systems Biology, ETH Zurich, Zurich, Switzerland
[4] School of Life Sciences, Westlake University, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China
[5] Department of Pathology and Molecular Pathology, University Hospital Zurich, University of Zurich, Zurich, Switzerland
[6] NEXUS Personalized Health Technologies, ETH Zurich, Zurich, Switzerland
[7] Swiss Institute of Bioinformatics, Zurich, Switzerland
[8] Department of Urology, University Hospital Zurich, University of Zurich, Zurich, Switzerland
[9] Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland
[10] Institut Curie, Paris, France
[11] IBM Zurich Research Laboratory, Zurich, Switzerland
[12] Fred Hutchinson Cancer Research Center, Seattle, USA
[13] Faculty of Science, University of Zurich, Zurich, Switzerland
[14] Dr. Senckenberg Institute of Pathology, University Hospital Frankfurt, Goethe-University Frankfurt, Frankfurt, Germany.

[*]Corresponding authors:

Tiannan Guo: guotiannan@westlake.edu.cn

Ruedi Aebersold: aebersold@imsb.biol.ethz.ch

Peter J. Wild: Peter.Wild@kgu.de

Andreas Beyer: abeyer2@uni-koeln.de

[#]Equal contribution

Abstract

Background

Tumor-specific genomic aberrations are routinely determined by high throughput genomic

measurements. It remains unclear though, how complex genome alterations affect molecular networks

through changing protein levels, and consequently biochemical states of tumor tissues.

Results

Here, we investigated the propagation of genomic effects along the axis of gene expression during

prostate cancer progression. For that, we quantified genomic, transcriptomic and proteomic alterations

based on 105 prostate samples, consisting of benign prostatic hyperplasia regions and malignant tumors,

from 39 prostate cancer patients. Our analysis revealed convergent effects of distinct copy number

alterations impacting on common downstream proteins, which are important for establishing the tumor

phenotype. We devised a network-based approach that integrates perturbations across different

molecular layers, which identified a sub-network consisting of nine genes whose joint activity positively

correlated with increasingly aggressive tumor phenotypes and was predictive of recurrence-free survival.

Further, our data revealed a wide spectrum of intra-patient network effects, ranging from similar to very

distinct alterations on different molecular layers.

Conclusions

This study uncovered molecular networks with remarkably convergent alterations across tumor sites and

patients, but it also exposed a diversity of network effects: we could not identify a single sub-network

that was perturbed in all high-grade tumor regions.

Keywords: molecular aberrations, network effects, prostate cancer, proteogenomic analysis, tumor
heterogeneity
60

Background

Prostate cancer (PCa) represents one of the most common neoplasm among men with almost

1,300,000 new cases and 360,000 deaths in 2018 [1] accounting for 15% of all cancers diagnosed. PCa is

the fifth leading cause of cancer death in men and represents 6.6% of total cancer mortality in men [1].

Despite earlier detection and new treatments, the lifetime risk to die of PCa has remained stable at

approximately   3%   since   1980.   (National   Cancer   Institute   SEER   data:

https://seer.cancer.gov/statfacts/html/prost.html). In many patients, PCa is indolent and slow growing.

The challenge is to identify those patients who are unlikely to experience significant progression while

offering radical therapy to those who are at risk. Current risk stratification models are based on

clinicopathological variables including histomorphologically defined grade groups, prostate-specific

antigen (PSA) levels and clinical stage. Although those variables provide important information for

clinical risk assessment and treatment planning [2, 3], they do not sufficiently predict the course of the

disease.

Extensive genomic profiling efforts have provided important insights into the common genomic

alterations in primary and metastatic PCa [4-9]. Interestingly, PCa genomes show a high frequency of

recurrent large-scale chromosomal rearrangements such as TMPRSS2-ERG [10]. In addition, extensive

copy number alterations (CNAs) are common in PCa, yet point mutations are relatively infrequent in

primary PCa compared to other cancers [6, 11]. A major complicating factor is that around 80% of PCas

are multifocal and harbor multiple spatially and often morphologically distinct tumor foci [12, 13].

Several recent studies have suggested that the majority of topographically distinct tumor foci appear to

arise independently and show few or no overlap in driver gene alterations [14-16]. Therefore, a given

prostate gland can harbor clonally independent PCas.

To allow for a more functional assessment of the biochemical state of PCa, it is necessary to go

beyond genomic alterations and comprehensively catalogue cancer specific genomic, transcriptomic and

proteomic alterations in an integrated manner [17-19]. Such an approach will provide critical

information for basic and translational research and could result into clinically relevant markers. While

hundreds of PCa genomes and transcriptomes have been profiled to date [20], little is known about the

PCa proteome. Although recent work has emphasized the need for integrated multi-omics profiling of

PCa, we still lack understanding about how genomic changes impact on mRNA and protein levels [17-19].

Especially the complex relationship between tumor grade, tumor progression and multi-layered

molecular network changes remains largely elusive.

For example, previous work has shown that copy number changes may alter transcript levels of

many genes, whereas the respective protein levels remain relatively stable [21]. Indeed, there is

compelling evidence across multiple tumor types that many genomic alterations are 'buffered' at the

protein level and are hence mostly clinically inconsequential [22]. To better understand the evolution of

PCa and to identify core networks perturbed by genomic alterations and thus central for the tumor

phenotype, it is therefore essential to investigate the transmission of CNAs to the transcriptomic and

proteomic level.

To this end, it is important to decipher which genomic alterations impact PCa proteomes, which

of those proteomic alterations are functionally relevant, and how molecular networks are perturbed at

the protein level across tumors.

To address these open questions, we performed a multi-omics profiling of radical prostatectomy

(RP) specimens at the level of the genome, transcriptome and proteome from adjacent biopsy-level

samples, using state-of-the-art technologies. Unique features of this study are (1) the utilization of PCT

(pressure cycling technology)-SWATH (Sequential Window Acquisition of all THeoretical Mass Spectra)

mass spectrometry [23, 24], allowing rapid and reproducible quantification of thousands of proteins

from biopsy-level tissue samples collected in clinical cohorts; (2) the simultaneous profiling of all omics

layers from the same tissue regions; (3) inclusion and full profiling of benign regions, which provides a

matching control for each tumor; and (4) the full multi-omics characterization of multiple tumor regions

from the same patients, thus enabling the detailed investigation of tumor heterogeneity. This design

resulted in the multi-layered analyses of 105 samples from 39 PCa patients, as well as of the exome of

corresponding peripheral blood cells yielding a comprehensive molecular profile for each patient and

identified molecular networks that are commonly altered in multiple patients. Importantly, some of the

affected genes/proteins exhibited very small individual effect sizes, suggesting that combined network

effects of multiple genes may significantly contribute to determining PCa phenotypes.

Results

Proteogenomic analysis of the sample cohort identifies known PCa biomarkers.

In this study, we analyzed 39 PCa patients (**Additional file 1: Fig. S1**) belonging to three groups

who underwent laparoscopic robotic-assisted RP. The patients were from the PCa Outcomes Cohort

(ProCOC) study [25, 26]. Tumor areas were graded using the ISUP (International Society of Urological

Pathology) grade groups [27], which range from ISUP grade group G1 (least aggressive) to G5 (most

aggressive). The more advanced grade groups G4 and G5 are considered jointly (G4/5). The cohort

tested included 12 low-grade (G1), 17 intermediate- (G2 and G3), and 10 high-grade (G4/5) patients (**Fig.**

1a, **Additional file 1: Fig. S1**, **Additional file 2: Table S1**). For low-grade PCa patients, we selected two

representative regions, one of benign prostatic hyperplasia (BPH) and one of malignant tumor (TA). Since

PCa often presents as a multifocal disease with heterogeneous grading within each prostate specimen

[24] we analyzed two different tumor regions from the 27 intermediate- and high-grade patients. In

those cases three representative regions, including BPH, the most aggressive tumor (TA1) and a

secondary, lower-grade tumor (TA2) [2] were analyzed. Thus, TA1 always represented the higher-grade

nodule compared to TA2. Note, whereas each patient was assigned a patient-specific overall grade (*i.e.*

'low', 'intermediate' or 'high'), each tumor area was additionally assigned an individual grade group

based on its histological appearance. According to current ISUP guidelines, the grading of the entire

prostate specimen depends on the size and grade of individual nodules [28]. Thus, it is possible that the

patient grading is lower than the grading of the most aggressive nodule, if another lower-grade nodule is

larger. Tumor regions contained at least 70% tumor cellularity and the distance between the analyzed

areas (TA1 versus TA2) was at least 5 mm. Altogether, we obtained 105 prostate tissue specimens

(**Additional file 2: Table S1**). Three adjacent tissue biopsies of the dimensions 0.6 x 0.6 x 3.0 mm were

punched from each representative region for exome sequencing, CNA (derived from the exome

sequencing data), RNA sequencing (RNA-seq), and quantitative proteomic analysis using the PCT-SWATH

technology [23] respectively. Proteomic analysis was performed in duplicates for each tissue sample.

Peripheral blood samples from each patient were also subjected to exome sequencing and served as the

genomic wild-type reference (**Fig. 1**). All three types of grading (*i.e.* patient-specific overall grading, TA1

grading and TA2 grading) were predictive of the recurrence-free survival (RFS) in our study.

In agreement with prior reports, we observed relatively few recurrent point mutations across

patients (**Additional file 1: Fig. S2**, **Additional file 3: Table S2**), but substantial CNAs (**Additional file 1:**

Figs. S3 and **S4**, **Additional file 4: Table S3**). In total, 1,110 genes showed copy number gains in at least

five samples or copy number losses in at least five samples (see **Additional file 1: Supplementary Text**

for details). Likewise, our data confirmed the differential expression of several transcripts/proteins that

had previously been suggested as PCa biomarkers or which are known oncogenes in other tumor types

(**Additional file 1: Fig. S5, Additional file 5: Table S4** and **Additional file 6: Table S5**) (see **Additional file**

1: Supplementary Text for details). This consistency with previously published results confirmed the

quality of our data and motivated us to go beyond previous work by performing a network-based multi-

omics multi-gene analysis.

Molecular perturbations correlate with tumor grade.

As a first step towards a cross-layer analysis, we asked if high-grade PCa would be generally

affected by stronger alterations (compared to low-grade PCa) at the genome, transcriptome, and

proteome layer [29]. Thus, we devised molecular perturbation scores that quantified the number of

affected genes/proteins and the extent to which these genes/proteins were altered in the tumor

specimens compared to their benign controls (see the '**Methods**' section for details). Higher-grade

tumors (G3 and G4/5) exhibited significantly higher molecular perturbation scores than lower-grade

tumors (G1 and G2). Those differences were statistically significant in all but one case (P value < 0.05,

one-sided Wilcoxon rank sum test, **Fig. 2**). The CNA perturbation magnitude exhibited the highest

correlation with the PCa grading, confirming prior studies documenting the tight association between

CNA, histopathological grade and risk of progression [4, 5, 30]. Previous work suggested that copy

number changes are to some extent buffered at the protein level [31]. Interestingly, we observed that

proteins known to be part of protein complexes were significantly less strongly correlated with the fold

changes (FCs) of their coding mRNAs than proteins not known to be part of protein complexes (P value <

2.6e-11, one-sided t-test, **Additional file 1: Fig. S6**). This result is consistent with the concept that protein

complex stoichiometry contributes to the buffering of mRNA changes at the level of proteins [22, 32-34].

Thus, molecular patterns in high-grade PCa are more strongly perturbed at all layers and the effects of

genomic variation are progressively but non-uniformly attenuated along the axis of gene expression.

Inter-patient heterogeneity decreases along protein biosynthesis.

Our analysis of CNA profiles (above and **Additional file 1: Supplementary Text**) already revealed

many shared CNAs across patients, suggesting that such common CNAs might represent genomic driver

changes. We therefore investigated if such a convergence towards common molecular endpoints could

also be observed at the transcript and protein level. To address this question, we first computed a

reference molecular signature that is characteristic of the molecular perturbations of tumors in a given

grade group. These 'centroid vectors' were obtained by computing the average tumor-to-benign FCs

across all samples within a grade group. Consistent with the observation above, we found that the

average effect sizes (averaged absolute centroid FCs) were increasing with the grade group for all three

layers (CNA, mRNA, and protein; **Additional file 7: Table S6**). Next, we compared each individual sample

within a group against the matching centroid of the same group. For the quantification of the similarity

between a tumor sample and the corresponding centroid we used four similarity/distance measures:

Pearson correlation, Mutual Information (MI)[35], Manhattan distance and Euclidean distance. While the

185 first two measures (*i.e.* Pearson correlation and MI) quantify the degree to which the tumor sample and

186 centroid vector co-vary, the other two measures (*i.e.* Manhattan and Euclidean distance) also take into

187 account the magnitude of the FCs in the two vectors. To illustrate this difference, imagine two patients

188 having perturbations of the same genes/proteins, but one of them exhibiting overall two-fold greater

189 FCs (*i.e.* all FCs are increased by a factor of two compared to the other patient). In such a scenario

190 Pearson correlation and MI would yield identical results for the two patients when compared to the

191 centroid, whereas Manhattan and Euclidean distance would identify them as different. Using the

192 Pearson correlation and MI, we found that high-grade PCa (G4/5) were more similar to their respective

193 centroid than low-grade PCa (G1) to their centroid (**Fig. 3**). This effect was particularly pronounced for

194 protein-level changes. This is consistent with the notion that protein levels (and not mRNA levels) are

195 subjected to stronger selection. Interestingly, when the Euclidean distance and the Manhattan distance

196 were used to characterize tumor similarity, we found that the high-grade tumors were more *dissimilar* to

197 each other than the low-grade tumors (**Additional file 1: Fig. S7**), in sharp contrast to the Pearson

198 correlation and MI. Based on the nature of the different similarity measures tested, we hypothesized

199 that there is a set of proteins commonly affected in their abundance by oncogenic alterations in high-

200 grade tumors. This would increase the similarity using the Pearson correlation or MI. However, although

201 the same proteins are affected, they are affected to a different extent in different high-grade tumors, *i.e.*

202 the FCs exhibited a high degree of variability (**Additional file 1: Fig. S7**), which would increase the

203 dissimilarity based on the Euclidean distance or Manhattan distance (see the '**Methods**' section and

204 **Additional file 1: Fig. S7** for a schematic explanation).

205   To further corroborate the notion of common endpoints, we focused on the 20 proteins with the

206 largest average absolute FCs across all tumor specimens (**Additional file 1: Fig. S7**, **Additional file 7:**

207 **Table S6**). Among them was PSA (KLK3), and several other well established PCa-associated proteins like

208 AGR2 [36], MDH2 [37], MFAP4 [38] and FABP5 [39]. We observed that for some of these top 20 proteins,

209 FCs were more extreme in the higher-grade tumors (G3 and G4/5) compared to lower-grade tumors (G1

210 and G2), such as MDH2 and SEPHS1 (up-regulation; **Fig. 3c**). RABL3 was one of the most strongly down-

211 regulated proteins (**Fig. 3c**), which is a surprising finding as RABL3 is known to be up-regulated in other

212 solid tumors [40, 41]. Interestingly, in most cases these proteins were from loci that were not subject to

213 CNAs (**Additional file 1: Fig. S7**, **Additional file 7: Table S6**), hinting that independent genomic events

214 would impact on these target proteins *via* network effects.

Effects of distinct CNAs converge on common proteins.

It has previously been suggested that mutations affecting different genes could impact common

molecular networks if the respective gene products interact at the molecular level [42]. However,

previous analyses were mostly restricted to individual molecular layers. For example, it was shown that

genes mutated in different patients often cluster together in molecular interaction networks [42]. But,

effects of these mutations on transcript and protein levels remained unexplored in this case. Here, we

aimed at a multi-layer network analysis, involving the genome, transcriptome and proteome using two

different network approaches.

First, we speculated that distinct genomic events in different patients would commonly impact

on at least some of the top 20 target proteins identified above. Among those top targets we selected

AGR2, ACPP, POSTN and LGALS3BP, because these proteins/genes had correlated protein- and mRNA

FCs; thus, protein level changes were likely caused by cognate mRNA level changes. Importantly, those

mRNA/protein level changes could not be explained by CNAs of the coding genes themselves (**Additional**

File 1: Fig. S7). Thus, these transcriptional changes were likely caused by *trans*-effects. To identify

potential regulators for each target gene, we used an independently inferred generic transcriptional

regulatory network (Leote *et al. in revision*; preprint [available on bioRxiv](#)) and selected putative

regulators at most two edges away from the target genes (see the '**Methods**' for details). Using the

ElasticNet algorithm we next fitted a linear model regressing the mRNA changes of the target genes

against the network neighbors' CNAs. Thereby we identified genes whose CNA changes were associated

with mRNA changes of the four target genes AGR2, ACPP (a.k.a ACP3), POSTN and LGALS3BP (**Additional**

file 7: Table S6). To validate our approach, we used two independent PCa cohorts (TCGA; [8] and MSKCC

[30]) and computed the association between the CNAs of each significant regulator and the

corresponding mRNA log-FC of the respective target gene. In most cases we observed an agreement in

terms of effect directions, *i.e.* the signs of association between CNA changes of the putative regulators

and the mRNA log-FCs of the respective target had the same direction as in our cohort (**Additional file 7:**

Table S6). Here we use ACPP/ACP3 as an illustrative example (**Fig. 3d)**: ACP3 (a.k.a. PAcP) is a prostate-

specific acid phosphatase with a critical role PCa etiology and has been suggested as a PCa biomarker

long before PSA [43]. ACP3 is known to inhibit cell proliferation and is therefore typically down-regulated

in PCa [44], despite elevated ACP3 protein levels in patient blood [43]. In our cohort ACP3 levels were

strongly down-regulated in all of the high-grade patients and in the vast majority of low- and

intermediate-grade patients, suggesting that ACP3 down-regulation represents an early event during

PCa evolution. Despite its established role in PCa, little is known about the oncogenic driver events

downregulating ACP3 [43]. Our network modeling identified six putative ACPP/ACP3 regulators

(ANKRD22, MS4A3, RHOV, ARL11, DEFB1, and DPYSL2), several of which are already known to be

associated with PCa [45-47]. Analyzing the CNA signatures of these six putative regulators reveals at

least two larger groups of patients (**Fig. 3d)**: the first one harboring joint deletions of ARL11 and

ANKRD22, the second one harboring joint deletions of DEFB1 and DPYSL2. The latter two genes are both

encoded on Chromosome 8 and thus, their deletion may be due to single CNA events. ARL11 and

ANKRD22 however, are encoded on different chromosomes. Importantly, these events were clonal in

most cases, *i.e.* they were mostly common to both tumor samples of a given patient. A remarkable

exception was patient M7, who had a joint deletion of ARL11 and ANKRD22 in tumor area 1 (TA1) and a

DEFB1/DPYSL2 deletion in tumor area 2 (TA2). Hence, our network analysis hints that distinct deletions

in the network vicinity of ACP3 can lead to the repression of this anti-proliferative protein. More

examples of distinct CNAs having similar effects on downstream targets were found for the other three

focus proteins (**Additional file 1: Fig. S7)**. Taken together, these findings suggest that tumor mechanisms

in different patients converged on common protein endpoints and that the expression levels of these

proteins were progressively more strongly affected during tumor evolution.

Joint network effects of CNAs drive tumor progression.

The analysis above identified molecular networks driving tumor alterations and thus indicated

altered biochemical states that were common to most tumor specimens. To identify sub-networks that

specifically distinguish high-grade from low-grade tumors, we mapped our data onto the STRING gene

interaction network [48], and employed network propagation [49, 50] separately to the CNA,

transcriptome and proteome data for each of the tumor samples. We excluded point mutations from

this analysis as their frequency was too low in our cohort. By combining published molecular

interactome data with a network propagation algorithm [42, 49], we aimed to 'enrich' network regions

with many perturbed genes/proteins. We reasoned that the convergent consequences of genomic

variants on common network regions would be indicative of specific biochemical functions that are

important for the tumor biology. We therefore identified genes/proteins in network regions that showed

a higher score (or a lower score) in high-grade (G4/5) relative to lower-grade (G1) tumor groups at all

three levels (**Fig. 4a**, **b**; '**Methods'** section). This analysis identified sub-networks consisting of over- and

under-expressed genes (relative to the benign controls). We found 57 amplified genes (**Additional file 7:**

Table S6) for which transcripts and proteins were often over-expressed in high-grade PCa (**Fig. 4a**) and

21 genes with copy number loss (**Additional file 7: Table S6**) for which transcripts and proteins were
often down-regulated compared to lower-grade tumors (**Fig. 4b**).

Among the up-regulated network nodes, we observed genes modulating the stability of
chromatin, such as chromatin-binding protein Chromobox 1 (CBX1) [51], SET Domain Bifurcated 1
(SETDB1) [52], a function linking to H3K27me3 and H3K9me3 in chromatin, and CBX3 (known as HP1-γ)
[53]. SETDB1 is an oncogene in melanoma [54] and has also been found to be over-expressed in PCa and
cell lines [55]. Further, we found genes involved in DNA damage repair, such as SMG7 [56] and ATR [57],
and PRKCZ[58], which had already been suggested as a biomarker prognostic for survival in PCa [59].
Multiple actin related proteins including ARPC1B [60], ARPC5 [61], ACTL6A [62], and CFL1 [63], which are
markers for aggressive cancers, were part of the up-regulated network nodes. Moreover, the up-
regulated genes contained proteins related to the cell cycle like BANF1 and proteins interacting with the
centrosome including LAMTOR1 and RAB7A that had already been associated with PCa [64]. Finally,
several signaling molecules with known roles in PCa were up-regulated, such as the transcription factor
Yin Yang 1 (YY1) [65], the TGF-β receptor TGFBR1 [66], and KPNA4, which promotes metastasis through
activation of NF-κB and Notch signaling [67]. Thus, up-regulated network nodes are involved in
DNA/chromatin integrity and growth control.

Likewise, several of the down-regulated genes had functions associated with PCa. For example,
the oxidative stress related gene MGST1, which is recurrently deleted in PCa [68]. ALDH1A3 is a direct
androgen-responsive gene, which encodes NAD-dependent aldehyde dehydrogenase [69]. DHCR24 is
involved in cholesterol biosynthesis and regulated by the androgen receptor [70]. Polymorphisms in
CYP1A1 are associated with PCa risk in several meta-analyses among different ethnicities [71-73].

Further, our network analysis is suggesting tumor mechanisms converging on genes that are
known contributors to PCa tumor biology. For example, the PCa-associated gene SF3B2 [74, 75] was only
weakly amplified in some of the high-grade tumors (average $\log_2$FC = 0.016) and mRNA levels showed
similarly small changes (average $\log_2$FC = 0.024). On the other hand, the SF3B2 protein levels were
consistently and more strongly up-regulated across tumors (average $\log_2$FC = 0.31), especially within the
high-grade tumors (**Additional file 1: Fig. S8**). Another example is UBE2T whose over-expression is
known to be associated with PCa [76]. Unfortunately, we could not quantify the corresponding protein
levels. However, we observed a strong and consistent mRNA over-expression across several tumors
(average $\log_2$FC = 0.73), even though at the DNA level the gene was only weakly amplified (average
$\log_2$FC = 0.023; **Additional file 1: Fig. S8**). Our findings of more heterogeneous CNAs, but more uniform

mRNA and protein alterations point on convergent evolutionary mechanisms, as we move along the axis
of gene expression.

Next, we analyzed the largest connected component with genes up-regulated in advanced
disease in more detail (see the '**Methods**' section). It consists of the nine nodes EMD, BANF1, ACTL6A,
YY1, RUVBL1, KANSL1, MRGBP, VPS72 and ZNHIT1 (**Fig. 4a**), and is referred to in the following as
Network Component 1 (**Additional file 7: Table S6**). Seven of these proteins are involved in chromosome
organization which may induce genomic alterations and influence the outcome of multiple cancers
including PCa [77]. For example, the actin-related protein ACTL6A is a member of the SWI/SNF (BAF)
chromatin remodeling complex[78], and a known oncogene and a prognostic biomarker for PCa [79].
Further, ACTL6A, RUVBL1 and MRGBP are together part of the NuA4/Tip60-HAT complex, which is
another chromatin remodeling complex involved in DNA repair [80]. Likewise, KANSL1 is involved in
histone post-translation modifications, while VPS72 is a member of histone- and chromatin remodeling
complexes [81]. Thus, Network Component 1 consists of genes involved in chromatin remodeling and
DNA repair, many of which are known to be involved in cancers.

Several samples were characterized by a small, but consistent DNA amplification of multiple
members of Network Component 1 (**Fig. 4c**). Out of the 66 tumor samples, there were 30 samples –
belonging to all grade groups – with a weak but consistent DNA amplification of Network Component 1
members, while the high-grade samples had stronger amplifications on average (*i.e.* larger effect sizes).
Importantly, gene members of Network Component 1 were dispersed across eight chromosomes
(**Additional file 7: Table S6**). The parallel DNA amplification of these genes is therefore the result of
multiple independent CNA events, while the signal on any single gene alone was too weak to be
significant in isolation. In some but not all cases, the amplifications led to a small, but consistent increase
in mRNA expression of the amplified gene loci (**Fig. 4d**). Unfortunately, only three out of the nine
proteins were detected in our proteomics experiments (**Fig. 4e**). Interestingly, patients where the DNA
amplifications led to transcript over-expression were almost always high-grade patients, whereas
patients where the amplification affected gene expression to a smaller extent were low- or
intermediate-grade patients (**Fig. 4c**, **d**). Further, we noticed that TA2 samples graded as G3 from high-
grade patients carried amplifications of Network Component 1, whereas tumor areas graded as G3 from
intermediate-grade patients did not have amplifications of this network component (**Fig. 4c**, **d**). Thus,
although the tumor areas were histologically equally classified, tumor areas from high-grade patients
carried a CNA signature and expression patterns reminiscent of the high-grade areas from the same
patients. Therefore, within the cohort tested the joint DNA amplification of this network component

along with RNA up-regulation is a signature of high-grade tumors. Curiously, the higher-grade tumor

areas of those high-grades patients (TA1) carried stronger DNA amplifications than the respective lower-

grade areas (TA2), which implies that the progressive amplification of Network Component 1 during

tumor evolution may contribute to an increasingly aggressive phenotype. In order to further corroborate

the clinical relevance of this network perturbation we analyzed published datasets of three additional

PCa cohorts (TCGA[8], MSKCC [30], and Aarhus [82]), together comprising a total of 713 patients with

known clinical outcome. We found that amplification of genes from Network Component 1 was a

significant predictor of reduced RFS in the MSKCC cohort (*P* value = 8.8e-3, log-rank test). In the TCGA

cohort, we observed the same trend although the difference in RFS was not statistically significant (*P*

value = 0.17; **Fig. 4f**). Additionally, we found that over-expression of genes from Network Component 1

was a significant predictor of reduced RFS in the TCGA cohort (*P* value = 2.1e-4, log-rank test), which was

the cohort with the largest number of patients. In the other two cohorts we observed the same trend,

although the difference in RFS was not statistically significant (*P* value = 0.30 and 0.093 for MSKCC, and

Aarhus; **Fig. 4f**). Thus, both CNA and RNA changes of Network Component 1 are predictive of the time to

relapse in independent cohorts.

In conclusion, our findings suggest that relatively weak but broad CNAs of entire network

components are associated with high-grade tumors and that the presence of some of these

perturbations in lower-grade tumors may be predictive of the future development of a more aggressive

phenotype.

Analysis of distinct tumor nodules defines intra-patient heterogeneity (TA1 versus TA2 comparison).

The CNA patterns (**Additional file 1: Fig. S4**) and the Network Component 1 analysis (**Fig. 4c**, **d**)

suggest that different tumor areas from the same patient shared several mutations. Such common

signatures are expected if different tumor nodules originate from a common clone. If this was true, we

would expect mutational signatures to be more similar between different nodules from the same patient

than between patients, even though mutated genes may be shared across patients. To compare the

intra- and inter-patient molecular heterogeneity at the levels of CNAs, transcript, and protein FCs, we

computed the Pearson correlation between tumor area 1 (TA1) and its paired tumor area 2 (TA2) for

each layer and all of the 27 patients with two characterized tumor areas (25 for the mRNA, see the

'**Methods**' section and **Additional file 1: Supplementary Text**). As a control, we also computed all

pairwise Pearson correlations between the samples within each of the grade groups (*i.e.* inter-patient

correlation). As expected, paired TA1 and TA2 from the same patient were on average more strongly
correlated to each other compared to samples from different patients within the same grade group. This
finding was consistent for all omics layers (**Fig. 5a**), and was more pronounced at the CNA and mRNA
layers compared to the protein layer.

Next, we tested whether a high correlation at the level of CNAs also implies a high correlation at
the level of mRNA and proteins. We tested this idea by 'correlating the correlations', *i.e.* we correlated
the TA1-TA2 correlation of CNA profiles with the correlation between the mRNA and protein profiles of
the same tumor areas (**Fig. 5b**). Indeed, a higher correlation of two tumor areas at the level of CNA
correlated significantly with a higher correlation at the level of mRNA (r=0.49, *P* value=0.014). In other
words, knowing how similar two tumor areas of a patient are at the CNA level supports a prediction of
their similarity at the mRNA level (and conversely). Although the correlation between protein and CNA
was not statistically significant, it followed the same trend (r=0.35, *P* value=0.076).

Comparing molecular similarity across omics layers allowed us to identify specific types of
patients. The patients H2, H4, M13 had highly correlated tumor areas at all three layers (upper right
corner in all scatterplots of **Fig. 5b**). Likely, the tumor areas of these patients have a common clonal
origin (**Additional file 1: Fig. S3**). In contrast, patients M12 and M14 had weakly correlated tumor areas
at all levels (bottom left corner in all scatterplots of **Fig. 5b**). These tumor nodules either have
independent clonal origins or they diverged at an earlier stage during tumor evolution (**Additional file 1:**
Fig. S3) [16]. For example, in the case of patient M12 large parts of the genome were not affected by
CNAs in the benign sample as well as in TA1 and TA2. However, as shown on **Additional file 1: Fig. S3**, a
large region was amplified in TA1, whereas the same region was deleted in TA2. This is consistent with a
scenario in which TA1 and TA2 show parallel evolution. A third class of patients is exemplified by the
patients M9 and M17, who showed a high correlation between their tumor areas on the CNA and mRNA
levels, but not on the protein level. Yet other patterns were apparent in patients M4, M7, and H10. They
showed similar mRNA and protein patterns in the two tumor areas, but relatively uncorrelated CNAs. M7
was the patient that we identified earlier with two different CNA signatures both reducing the levels of
the same protein (ACP3). The results here apply to global proteome patterns and therefore hint that
such convergent network effects of CNAs can be frequent. We confirmed that protein-level similarity
correlated with similar histological characteristics of the tumor areas. **Additional file 1: Fig. S9** shows
formalin-fixed paraffin-embedded (FFPE) tissue microarray images (duplicates) from the analyzed tumor
nodules (TA1 and TA2, diameter 0.6 mm), further underlining the hypothesis that ultimately protein-
level alterations are responsible for common cellular phenotypes. Although we cannot fully exclude the

possibility that some of these results were affected by technical noise in the data, our findings suggest

that transcript alterations can frequently be buffered at the level of proteins (patients M9, M17,

Additional file 1: Fig. S6) and that convergent evolutionary processes may lead to the alteration of

common proteins (patients M4, M7, H10). We also note that our findings are specific to the two tumor

areas available in this study and could be different if other nodules had been sampled for each of the

patients. However, our findings on patients with weakly correlated tumor areas at all levels like M12 and

M14 suggest that these patients might carry more than one disease [16].

Discussion

Despite twenty years of oncological research involving genome-scale (omics) technologies, we

know remarkably little about how the discovered genomic alterations affect the biochemical state of a

cell and consequently the disease phenotype. In particular, little is known about how genomic

alterations propagate along the axis of gene expression [17, 18]. Here, we have exploited recent

technological advances in data acquisition that made it possible to characterize small samples of the

same tumor specimens at the level of genomes, transcriptomes, and proteomes and advances in

computational strategies towards the network-based integration of multi-omics data.

In our study, samples were generated from small, less than 1 mm diameter punches in

immediate spatial proximity in the tumor and subsequently profiled at all three 'omics layers' (DNA, RNA,

proteome). Due to the large spatial heterogeneity of PCa [14, 24], this design - which is so far uncommon

for studies profiling multiple layers from tumor specimens - was instrumental for increasing the

comparability of the various omics layers and thus facilitated the analysis of molecular mechanisms. Our

key findings are: (1) we confirmed the importance of CNAs for PCa biology and the alteration of many

known PCa-associated genes at the transcript- and protein-level; (2) we revealed a generally elevated

molecular alteration of high-grade tumors compared to lower-grade tumors; (3) although our study

confirmed large within- and between-patient genomic heterogeneity, (4) we detected molecular

networks that were commonly altered at the mRNA and protein-level. The fact that many of those target

molecules are known drivers of PCa tumorigenesis, supports the notion that these proteins/transcripts

are subject to convergent evolutionary mechanisms.

We integrated the three omics layers using a network-based approach as opposed to directly

comparing gene perturbations (mutations) to gene products (transcripts and proteins). Using genome

data only, it had previously been hypothesized that whereas the identity of specific mutated genes may

differ between tumors, those mutations might still affect common molecular networks [42]. In other

words, tumor phenotypes are determined by the perturbation of molecular networks and not by the

perturbation of isolated genes. Our study provides experimental evidence that such network effects are

indeed propagated to subsequent molecular layers and that this effect propagation may be clinically

relevant. A very prominent example is our discovery that the long-known PCa gene ACPP (ACP3) is

downregulated through diverse CNA events, while sometimes even within the same patient different

CNA events might downregulate this critical tumor suppressor.

Our multi-omics network analysis revealed that high-grade PCa tumors distinguished themselves

from low-grade tumors in two aspects. The first is a generally higher heterogeneity and loss of controlled

gene regulation, which increased the molecular differences among high-grade tumors. It had previously

been shown that gene expression in tumors is often less coordinated than in normal samples [29]. The

increasing heterogeneity of protein concentrations suggests that this loss of coordinated expression also

affects protein levels. The second aspect is the convergence of molecular alterations towards specific

molecular sub-networks at the genomic, transcriptomic and proteomic layer along the progression from

low-grade to high-grade tumors. Thus, although we observed globally a higher degree of variability in

gene expression and proteome control among high-grade specimens, a specific subset of the observed

alterations appeared crucial for determining the aggressive tumor phenotype. Tumors are under

selective pressure acting on the biochemical function of the cells. It is generally believed that proteins

are a closer reflection of the functional state of a cell than the mRNA. Here we could show that the fold

changes of proteins like RABL3, MFAP4, and SF3B2 were more pronounced and/or uniform across high-

grade tumors than either their coding mRNAs or the underlying CNAs.

Specifically, our analysis led to the identification of Network Component 1, a sub-network

involved in chromatin remodeling and consisting of genes that were weakly amplified in intermediate-

grade (G3) tumor specimens. Signals of individual gene members of this component were virtually

indistinguishable from noise in our cohort. However, their consistent alterations across the network

region, across molecular layers and the fact that the same genes showed enhanced signals in high-grade

specimens, rendered this component highly interesting. The fact that copy number and expression

changes of Network Component 1 members were predictive for survival in independent cohorts further

supports the potential clinical relevance of this sub-network. Our network-based cross-omics analysis

identified nine other network components (**Fig. 4**) successfully capturing several known and potentially

new PCa-associated genes. However, neither Network Component 1 nor any of the other network

components was uniformly subject to CNAs across all high-grade patients. Instead, we found different

network components modified in different patients and these sub-networks were involved in cellular

processes as diverse as actin remodeling, DNA damage response, and metabolic functions, all of which

are known contributors to PCa biology. This further underlines the large inter-patient variability of PCa

and it demonstrates the diversity of molecular mechanisms leading to histologically similar phenotypes.

Future prediction models of PCa including the ISUP grade groups, PSA levels and clinical stage might be

improved by exploiting multi-omics network analyses. Detecting aggressive networks alterations in

prostate biopsies would help clinicians to advice either active surveillance or active therapy. However,

the development of such multi-dimensional biomarkers would require much larger patient cohorts.

Another distinguishing feature of this study was the simultaneous profiling of two different

tumor regions in 27 out of the 39 patients. The profiling of multiple tumor regions from the same

prostate helped to further highlight the enormous heterogeneity of PCa within patients and provided

important insights into PCa evolution. The fact that Network Component 1 was more strongly affected in

the paired higher-grade nodules of high-grade patients suggests that at least certain sub-networks are

subject to an evolutionary process, that progressively 'moves' protein levels towards a more aggressive

state. Generally, and at all molecular layers tested, the two paired tumor areas were more similar to

each other compared to two samples from the same grade group but different patients, suggesting

common evolutionary origins. Although the two tumor areas seemed to mostly originate from the same

clone, this was not always the case. In some patients, different nodules exhibited different molecular

patterns at all omics layers, suggesting early evolutionary separation. Thus, for the first time, current

diagnostic, expert-level consensus guidelines [28] are supported by detailed proteogenomic data. Our

findings support earlier claims that clonality itself might be a prognostic marker with implications for

future, more tumor-specific treatment when targeted therapies become available also for PCa [16, 83].

Our study shows that all three molecular layers (genome, transcriptome and proteome)

contributed valuable information for understanding the biology of PCa. In particular the DNA layer

informed about causal events, clonality, and genomic similarity between tumors. The transcriptome was

relevant for understanding the transmission of CNA effects to proteins and served as a surrogate in cases

where protein levels remained undetected. The proteome was crucial for revealing protein-level

buffering of CNA effects as well as for indicating convergent evolution on functional endpoints. In a

routine diagnostic context though, measuring all three layers may not be feasible for the near future due

to resource and time limitations. Thus, the identification of improved, routine-usable molecular markers

for PCa diagnostics and prognosis remains an open problem [17].

Conclusions

This study uncovered molecular networks with remarkably convergent alterations across tumor

sites and patients. In particular, we identified a sub-network consisting of nine genes whose joint activity

positively correlated with increasingly aggressive tumor phenotypes. The fact that this sub-network was

predictive for survival in independent cohorts further supports its potential clinical relevance. At the

same time though, our study also exposed a diversity of network effects: we could not identify a single

sub-network that was perturbed in all high-grade tumor regions, let alone the observed distinct intra-

patient alterations at all omics layers for some patients. Overall, our study has significantly expanded our

understanding of PCa biology and serves as a model for future work aiming to explore network effects of

mutations with an integrated multi-omics approach.

Methods

Patients and samples

A total of 39 men with localized PCa who were scheduled for RP were selected from a cohort of

1,200 patients within the ProCOC study and processed at the Department of Pathology and Molecular

Pathology, University Hospital Zurich, Switzerland [25]. Each of the selected intermediate- and high-

grade patients had two different tumor nodules with different ISUP grade groups. H&E (Hematoxylin and

Eosin)-stained fresh frozen tissue sections of 105 selected BPH and tumor regions were evaluated by two

experienced pathologists (PJW, NJR) to assign malignancy, tumor stage, and Grade Group according to

the International Union Against Cancer (UICC) and WHO/ISUP criteria. This study was approved by the

Cantonal Ethics Committee of Zurich (KEK-ZH-No. 2008-0040), the associated methods were carried out

in accordance with the approved guidelines, and each patient has signed an informed consent form.

Patients were followed up on a regular basis (every three months in the first year and at least annually

thereafter) or on an individual basis depending on the disease course in the following years. The RFS was

calculated with a biochemical recurrence (BCR) defined as a PSA ≥0.1 ng/ml. Patients were censored if

lost to follow-up or event-free at their most recent clinic visit. Patients with a postoperative PSA

persistence or without distinct follow-up data for the endpoint BCR were excluded from the analysis of

BCR.

Exome sequencing and somatic variant analysis

The exome sequencing (exome-seq) was performed using the Agilent Sure Select Exome

platform for library construction and Illumina HiSeq 2500 for sequencing read generation. We mapped

and processed the reads using a pipeline based on bowtie2 [84] (1.1.1) and the Genome Analysis Tools

Kit (GATK) [85] (3.2-2). We detected and reported nonsynonymous variants or variants causing splicing

changes using Strelka (1.0.14) and Mutect (1.1.7) combined with post-processing by the CLC Genomics

Workbench (8.0.3). In this process, all tissue samples of a patient were compared to the respective blood

sample.

Trimmomatic [86] (0.36) was used for adaptor clipping and low-quality subsequence trimming of

the FASTQ files. Subsequently, single reads were aligned to the hg19 reference genome with bowtie2

with options "--very-sensitive -k 20". We applied samtools [87] (0.1.19) and picard-tools (1.119) to sort

the resulting bam files in coordinate order, merge different lanes, filter out all non-primary alignments,

and remove PCR duplicates. Quality of the runs was checked using a combination of BEDtools [88] (2.21),

samtools, R (3.1) and FastQC (0.11.2).

Bam files containing the mapped reads were preprocessed in the following way: indel

information was used to realign individual reads using the RealignerTargetCreator and IndelRealigner

option of the GATK. Mate-pair information between mates was verified and fixed using Picard tools and

single bases were recalibrated using GATK's BaseRecalibrator. After preprocessing, variant calling was

carried out by comparing benign or tumor prostate tissue samples with matched blood samples using

the programs MuTect [89] and Strelka [90] independently. Somatic variants that were only detected by

one of the two programs were filtered out using CLC Genomics Workbench. So were those that had an

entry in the dbSNP [91] common database and those that represented synonymous variants without

predicted effects on splicing.

CNA analysis of exome-seq data

The Bam files generated during the process of somatic variant calling were processed with the

CopywriteR package (v.2.2.0) for the R software [92]. CopywriteR makes use of so-called "off-target"

reads, *i.e.* reads that cover areas outside of the exon amplicons. "Off-target" reads are produced due to

inefficient enrichment strategies. In our case on average 28.5% of the total reads were not on target.

Briefly, CopywriteR removes low quality and anomalous read pairs, then peaks are called in the

respective blood reference, and all reads in this region are discarded. After mapping the reads into bins,

those peak regions, in which reads had been removed, were compensated for. Additionally, read counts

are corrected based on mappability and GC-content. Finally, a circular binary segmentation is carried out

and for each segment the log count ratios between tissue samples and the respective blood sample are

reported as copy number gain or loss. The copy number of each gene in each sample was reported

based on the log count ratio of the respective segment in which the gene was located. The overall

performance of this CNA-calling approach was evaluated by comparing the results of the TA1 (and TA)

samples with CNA results obtained by applying the OncoScan Microarray pipeline to FFPE samples from

the same tumors (**Additional file 1: Fig. S10**).

OncoScan Microarrays

OncoScan copy number assays were carried out and analyzed as described previously [93].

Briefly, DNA was extracted from punches of FFPE cancer tissue blocks. Locus-specific molecular inversion

probes were hybridized to complementary DNA and gaps were filled in a nucleotide-specific manner.

After amplification and cleavage of the probes, the probes were hybridized to the OncoScan assay

arrays. Scanning the fluorescence intensity and subsequent data processing using the Affymetrix®

GeneChip® Command Console and BioDiscovery Nexus express resulted in log intensity ratio data

(sample versus Affymetrix reference) and virtual segmentation of the genome into areas with copy

number gain, loss or stability.

RNA Sequencing

RNA sequencing was performed at the Functional Genomics Center Zurich. RNA-seq libraries

were generated using the TruSeq RNA stranded kit with PolyA enrichment (Illumina, San Diego, CA, USA).

Libraries were sequenced with 2x126bp paired-end on an Illumina HiSeq 2500 with an average of 105.2

mio reads per sample.

Paired-end reads were mapped to the human reference genome (GRCh37) using the STAR

aligner (version 2.4.2a) [94]. Quality control of the resulting bam files using QoRTs [95] and mRIN [96]

showed strong RNA degradation[97] in a significant fraction of the samples: mRIN classified 31 samples

as highly degraded (**Additional file 1: Fig. S11**, **Additional file 5: Table S4**). In order to correct for this 3'

bias, 3 tag counting was performed as described by Sigurgeirsson et al [98] using a tag length of 1,000.

After 3' bias correction, three samples still showed a clear 3' bias: the two tumor regions (TA1 and TA2)

of the patient M5 and TA2 from patient M8 (**Additional file 1: Fig. S11**). These samples were excluded

from subsequent analyses. Additionally, the BPH region of the patient M5 was excluded due to the

exclusion of both its tumor regions.

FeatureCounts [99] was used to determine read counts for all genes annotated in ENSEMBL v75.

Genes for which no read was observed in any of the samples in the original data were excluded from the

analysis. Further, after 3 tag counting, all genes with without at least 1 read per million in N of the

samples were removed. We chose N to be 10 which corresponds to the size of the smallest grade group

(G2). In a last reduction step, all genes with more than one transcript were excluded, yielding a final set

of 14,281 genes.

Read count normalization and differential gene expression analysis was performed using the R

packages sva [100] and DESeq2 [101]. All benign tissues were considered biological replicates and

differential gene expression for the individual tumor samples was determined against all benign tissues.

Gene expression changes with an adjusted *P* value < 0.1 were considered significant.

RNA-seq - 3' bias correction

The 3 tag counting approach for 3' bias correction was used on the RNA-seq dataset [98]. This

approach requires changing of the annotation file in two steps: 1) isoform filtering and 2) transcript

length restriction. As proposed in [98] for each gene we determined the highest expressed isoform

within a set of high quality samples. As high quality samples we used all samples with an mRIN score

greater than or equal to 0.02. This set contains 7 benign and 15 tumor samples. Isoform expression was

determined using cufflinks [102]. As transcript length we chose 1,000bp.

Gene fusions

602   FusionCatcher (version 0.99.5a beta) was used to determine gene fusions for all samples.

603 Fusions classified as "probably false positive" are discarded unless they are also classified as "known

604 fusion".

PCT assisted sample preparation for SWATH-MS

We first washed each tissue sample to remove O.C.T., followed by PCT-assisted tissue lysis and

protein digestion, and SWATH-MS analysis, as described previously [23]. Briefly, a series of ethanol

solutions were used to wash the tissues each tissue, including 70% ethanol / 30% water (30 s), water (30

s), 70% ethanol / 30% water (5 min, twice), 85% ethanol / 15% water (5 min, twice), and 100% ethanol (5

min, twice). Subsequently, the tissue punches were lysed in PCT-MicroTubes with PCT-MicroPestle [103]

with 30 µl lysis buffer containing 8 M urea, 0.1 M ammonium bicarbonate, Complete protease inhibitor

cocktail (Roche) and PhosSTOP phosphatase inhibitor cocktail (Roche) using a barocycler (model

NEP2320-45k, PressureBioSciences, South Easton, MA). The lysis was performed with 60 cycles of high

pressure (45,000 p.s.i., 50 s per cycle) and ambient pressure (14.7 p.s.i., 10 s per cycle). The extracted

proteins were then reduced and alkylated prior to lys-C and trypsin-mediated proteolysis under pressure

cycling. Lys-C (Wako; enzyme-to-substrate ratio, 1:40) -mediated proteolysis was performed using 45

cycles of pressure alternation (20,000 p.s.i. for 50 s per cycle and 14.7 p.s.i. for 10 s per cycle), followed

by trypsin (Promega; enzyme-to-substrate ratio, 1:20)-mediated proteolysis using the same cycling

scheme for 90 cycles. The resultant peptides were cleaned using SEP-PAC C18 (Waters Corp., Milford,

MA) and analyzed, after spike-in 10% iRT peptides [51], using SWATH-MS following the 32-fixed-size-

window scheme as described previously [19, 21] using a 5600 TripleTOF mass spectrometer (Sciex) and a

1D+ Nano LC system (Eksigent, Dublin, CA). The LC gradient was formulated with buffer A (2%

acetonitrile and 0.1% formic acid in HPLC water) and buffer B (2% water and 0.1% formic acid in

acetonitrile) through an analytical column (75 μm × 20 cm) and a fused silica PicoTip emitter (New

Objective, Woburn, MA, USA) with 3-μm 200 Å Magic C18 AQ resin (Michrom BioResources, Auburn, CA,

USA). Peptide samples were separated with a linear gradient of 2% to 35% buffer B over 120 min at a

flow rate of 0.3 μl min^{-1}. Ion accumulation time for MS1 and MS2 was set at 100 ms, leading to a total

cycle time of 3.3 s.

SWATH assay query library for prostate tissue proteome

To build a comprehensive library for SWATH data analysis, we analyzed unfractionated prostate

tissue digests prepared by the PCT method using Data Dependent Acquisition (DDA) mode in a tripleTOF

mass spectrometer over a gradient of 2 hours as described previously [19]. We spiked iRT peptides [51] into

each sample to enable retention time calibration among different samples. We then combined these

data with the DDA files from the pan-human library project [104]. All together we analyzed 422 DDA files

using X!Tandem [52] and OMSSA [53] against three protein sequence databases downloaded on Oct 21, 2016

from UniProt, including the SwissProt database of curated protein sequences (n=20,160), the splicing

variant database (n=21,970), and the trembl database (n=135,369). Using each database, we built target-

decoy protein sequence database by reversing the target protein sequences. We allowed maximal two

missed cleavages for fully tryptic peptides, and 50 p.p.m. for peptide precursor mass error, and 0.1 Da

for peptide fragment mass error. Static modification included carbamidomethyl at cysteine, while

variable modification included oxidation at methionine. Search results from X!Tandem and OMSSA were

further analyzed through Trans-Proteomic Pipeline (TPP, version 4.6.0) [54] using PeptideProphet and

iProphet, followed by SWATH assay library building procedures as detailed previously [19, 55]. Altogether,

we identified 167,402 peptide precursors, from which we selected the proteins detected in prostate

tissue samples, and built a sample-specific library. SWATH wiff files were converted into mzXML files

using ProteoWizard [56] msconvert v.3.0.3316, and then mzML files using OpenMS [57] tool FileConverter.

OpenSWATH[105] was performed using the tool OpenSWATHWorkflow with input files including the

mzXML file, the TraML library file, and TraML file for iRT peptides.

Peptide quantification using OpenSWATH

To obtain consistent quantification of the SWATH files, we obtained the all annotated *b* and *y*

fragments from the sp, sv and tr libraries. About ten thousand redundant and low-quality assays were

removed. Then we extracted the chromatography of these fragments and MS1 signals using

OpenSWATHWorkflow, followed by curation using DIA-expert[106]. Briefly, the chromatography of all

fragments and MS1 signals were subject to scrutiny by empirically developed expert rules. A reference

sample with best q value by pyprophet was picked up to refined fragments. The peptide precursors are

further filtered based on the following criteria: i) remove peptide precursors with a q value higher than

1.7783e-06 to achieve a false discovery rate of 0.00977 at peptide level using SWATH2stats [107]; ii)

peptides with a FC higher than 2 between the reference sample and its technical replicate were

removed; iii) peptides matching to multiple SwissProt protein sequences were removed. The data matrix

was first quantile normalized, $\log_2$ transformed, followed by batch correction using the ComBat R

package [108]. Finally, for each protein and pair of technical replicates the average value was computed.

Statistical analysis

663   All plots were produced with R. Kaplan-Meier estimators were used for RFS analysis. Differences

664   between survival estimates were evaluated by the log-rank test.

Computation of molecular perturbation scores

On the genomic level (mutation and CNA), we kept the tumor samples (66 in total) that contain

FCs with respect to the blood. The mutation matrix was further discretized by setting all non-zero events

to 1. At the transcriptomics level, the FCs for the 63 tumor samples were computed as described above

(see 'RNA Sequencing'). Finally, on the proteomics level, we computed the FCs for the tumor samples (66

in total) as follows: for each protein, its mean intensity over the normal samples was subtracted from

the intensities of the tumor samples. (We chose to compute the FCs for the tumor samples with respect

to a global reference (average of all normal samples) and not with respect to their paired benign sample

in order to achieve a higher consistency with the transcriptomics level.)

We assigned to each sample two molecular perturbation scores summarizing/quantifying the

magnitude of its FCs: DE_count counts the number of mutated/differentially expressed (DE) genes, while

the DE_sum score is the sum of absolute FCs of all genes. Thus, while the first score counts the number

of events (mutations/DE genes), the second one quantifies their magnitude. A gene is regarded as

mutated/DE if its value is 1 in the mutation layer and if its absolute value is above a threshold that has

been set to 1 for the mRNA and protein layer. For the CNA layer, the corresponding threshold was set to

0.5 because the range of FCs in the CNA matrix is smaller than the mRNA and protein matrices. Both

types of scores were computed for each molecular level, except for the point mutations where only

DE_count was computed. Afterwards, the 66 DE_count scores (63 for the mRNA) and the DE_sum scores

at each layer were divided into the four grade groups G1, G2, G3 and G4/5 respectively.

Network propagation/smoothing

As a network, the STRING gene interaction network (version 10)[48] was used, after removing all

edges with combined score smaller or equal to 900 and keeping subsequently the largest connected

component. The resulting network consisted of 10,729 nodes and 118,647 (high-confidence) edges. For

the network smoothing, the weight matrix was computed as described in Vanunu et al.[49], but for an

unweighted graph and the propagation parameter was set to 0.5. The propagation was iteratively

repeated 500 times to ensure convergence of the results. For the mapping from gene symbols to STRING

identifiers (**Additional file 7: Table S6**) we used the R/Bioconductor package STRINGdb [109]. The gene

symbols with no matching STRING identifier were removed, while for those that mapped to multiple

STRING identifiers, the first mapping was kept (default choice in the package). From the multiple gene

symbols that mapped to the same STRING identifier, the first mapping was kept. The genes that were

not present in the network were removed from the datasets, while those that were present in the

network but not in the corresponding dataset were initially filled in with 0's.

Genes with very small, 'smoothed' (absolute) FCs were filtered out as follows: after the network

propagation, only network nodes that had protein measurements themselves or at least one direct

neighbor (on the filtered STRING network) with protein measurements were considered in the next

steps of this analysis. *I.e.* network nodes without measured FCs at the protein layer that had no direct

neighbor with measured protein values were removed from the subsequent analyses.

For significance testing, the one-sided Wilcoxon rank sum test comparing the smoothed FCs

between the groups G4/5 and G1 was applied to each network node (after filtering) and layer, once for

up-regulation and once for down-regulation. The resulting sub-networks (up-regulated and down-

regulated) consisted of those genes that were significant (*P* value below 0.05) at all three layers and all

of the edges connecting them on the filtered STRING network.

Network Component 1 analysis

For each tumor sample at the CNA layer, a one-sided, one-sample t-test has been applied testing

if its average FC over the genes of the Network Component 1 (and in particular those that have been

measured at the CNA) is significantly greater than 0. Due to the presence of outliers in some samples,

the non-parametric, one-sided Wilcoxon signed-rank test has been applied as well yielding very similar

results (data not shown). A result is considered to be significant if the corresponding P value is below

0.05. The analysis has been repeated for the mRNA and protein layer.

Independent cohorts validation

For the validation of Network Component 1, we used published datasets of three PCa cohorts:

TCGA, MSKCC, and Aarhus. For TCGA and MSKCC, we downloaded the CNA, mRNA with precomputed z-

scores per gene, and corresponding clinical data from cBioPortal[110] (https://www.cbioportal.org/).

There were 489 samples with $\log_2$CNA data and 493 samples with mRNA profiles in TCGA. In MSKCC,

there were 157 primary tumors with CNA data and 131 primary tumors with mRNA data. The clinical

endpoint used in TCGA was the progression-free survival time and the disease-free survival in MSKCC. All

previous samples had known survival time.

For the Aarhus study (NCBI GEO dataset GSE46602), we downloaded the mRNA matrix and

corresponding clinical information as described in Ycart et al [111]. The resulting mRNA matrix consisted

of 20,186 genes and 50 samples- 40 PCa samples with known RFS time and 10 benign samples. Once

excluding the benign samples, we computed z-scores per gene in order to have comparable values with

the other two studies. These 40 PCa samples were also considered in the subsequent survival analysis.

CNA data was not available for the Aarhus study.

We reduced all datasets to the nine genes of Network Component 1. In each of the datasets, we

computed for each sample an average copy number change (CNA) or an average z-score (mRNA) across

the nine genes of Network Component 1 (combined risk score). Subsequently, we used these combined

risk scores to split the samples of each dataset into two groups: samples with a combined risk score

larger or equal to the median combined risk score of the study were considered as 'altered' and the rest

as 'unaltered'. Kaplan-Meier curves were generated for the two groups. Due to the high level of

discretized values in MSKCC at the CNA layer, a sample is considered to be 'altered' in that dataset if its

combined risk score is above zero.

Analysis of regulators and target genes

For this analysis, we used an independently inferred (partially directed) generic transcriptional

regulatory network (Leote *et al. in revision*; preprint available on bioRxiv). For each target gene the

following procedure was applied: firstly we identified its neighborhood of order two in the

transcriptional network by considering incoming edges only, *i.e.* all nodes from which the target gene

was reachable in at most two steps (equivalently the node itself, its parents and the parents of parents).

These are potential regulators of the target under consideration. Of these nodes, genes with no copy

number measurements and genes altered (*i.e.* with $\log_2$ copy number ratio greater than 0.5 in absolute)

in fewer than two tumor samples across the 66 tumor samples were filtered out. Subsequently, we fitted

an elastic net model with alpha=0.5. We used as output variable the mRNA FC of the target gene and as

input variables the CNAs of the regulators after the filtering. The value for the regularization parameter

lambda was chosen through 10-fold cross validation (default in the R package glmnet (https://cran.r-

project.org/web/packages/glmnet/)). Predictors/regulators with a non-zero beta coefficient were

deemed significant. We have used the elastic net model with alpha=0.5 because it is a method giving

sparse solutions and can deal with correlated predictors at the same time.

For the validation of our approach, we used the two independent PCa cohorts described above

(TCGA and MSKCC) and reduced the samples to those having both CNA and mRNA profile. This resulted

in 488 samples for TCGA and 109 samples for MSKCC. Next, for each of the significant

regulators/predictors we computed the Spearman correlation between its CNAs and the corresponding

mRNA z-scores of the target gene in each of the two independent studies. Finally, for each target gene

and each study we counted how many times the sign of the Spearman correlation matched the sign of

the Spearman correlation computed for our cohort, *i.e.* there was an agreement regarding the direction

of the association (**Additional file 7: Table S6**).

Differences between similarity/distance measures

A mathematical explanation for the observed differences between the similarity/distance

measures is provided. Let $\boldsymbol{y}_i^G$ denote the vector of $\log_2$ fold changes ($\log_2$FCs) corresponding to the *i*-th

tumor sample in the grade group G ($G \in \{G1, G2, G3, G4/5\}$), *i.e.* the fold changes of protein/mRNA

concentration in the respective tumor sample versus the global benign control. Then we can write:

764
$$y_i^G = \begin{pmatrix} y_{i1}^G \\ y_{i2}^G \\ \vdots \\ y_{iN}^G \end{pmatrix} = \begin{pmatrix} \mu_1^G \\ \mu_2^G \\ \vdots \\ \mu_N^G \end{pmatrix} + \begin{pmatrix} \varepsilon_{i1}^G \\ \varepsilon_{i2}^G \\ \vdots \\ \varepsilon_{iN}^G \end{pmatrix},$$
(1)

766 where y_{ij}^G corresponds to the log$_2$FC of protein/mRNA j in the respective tumor sample (*i.e. i*-th tumor

767 sample in the grade group G), μ_j^G to its (population) mean log$_2$FC in the grade group G and ε_{ij}^G to the

768 deviation of the respective individual log$_2$FC from the population mean (j=1,…N), N being the total

769 number of proteins/genes. The population mean μ_j^G is estimated by computing the average log$_2$FC

770 across all tumor samples in that grade group for the protein/mRNA j. Equivalently, (1) can be re-written:

771
$$\begin{pmatrix} y_{i1}^G \\ y_{i2}^G \\ \vdots \\ y_{iN}^G \end{pmatrix} - \begin{pmatrix} \mu_1^G \\ \mu_2^G \\ \vdots \\ \mu_N^G \end{pmatrix} = \begin{pmatrix} \varepsilon_{i1}^G \\ \varepsilon_{i2}^G \\ \vdots \\ \varepsilon_{iN}^G \end{pmatrix}.$$

On the one hand, the Euclidean and Manhattan distance between one tumor sample and the

corresponding mean/centroid vector are functions of these error terms ε_{ij}^G's alone and thus only

dependent on the variances of the proteins/genes in the corresponding grade group – these are shown

to be higher in the high-grade tumors. For the Pearson correlation r on the other hand, one can use the

equivalent simple linear regression modeling in (1) with the beta coefficient being equal to 1. By

standardizing the vector y_i^G and the centroid vector, we have that $r =$

$1\sqrt{var(centroid\ vector)/var(y_i^G)} =$

$\sqrt{var(centroid\ vector)/(var(centroid\ vector) + var(residuals))} =$

$\sqrt{1/(1 + var(centroid\ vector)/var(residuals))}$ where *var* denotes the variance. The Pearson

correlation will thus depend both on the variance of the residuals and the variance of the predictor, *i.e.*

mean/centroid vector in the corresponding grade group. Both quantities are expected to be higher in the

high-grade samples, while our data implies that the variance of the centroids (predictors) dominates

over the variance of the residuals.

Mutual information

In order to compute the MI between each individual sample within a group against the matching

centroid of the same group, the R package infotheo (http://cran.r-

project.org/web/packages/infotheo/index.html) has been used. For the MI, the data needs to be

discrete. For that, we discretize the tumor-to-benign FCs and centroids by setting 1, if the value is above

a threshold *thr*, -1, if the value is below *-thr* and 0 else. (*thr* is equal to 1 for the mRNA and proteins and

0.5 for the CNAs.)

792   Availability of data and materials

Exome and RNA sequencing data were submitted to the Sequence Read Archive (SRA) at NCBI under

accession numbers PRJNA577801 (exome-seq) and PRJNA579899 (RNA-seq), respectively. The SWATH

proteomics data were deposited in PRIDE. Project accession code is PXD004589. The published datasets

of the two PCa cohorts (TCGA and MSKCC) analyzed during the current study can be downloaded from

cBioPortal[110] while the third (Aarhus) is available at the NCBI GEO repository under the accession

number GSE46602 .

799   Acknowledgements

800   Not applicable.

801   Funding

This work was supported by the SystemsX.ch project PhosphoNet PPM (to R.A. and P.J.W.), the Swiss

National Science Foundation (grant no. 3100A0-688 107679 to R.A.), the European Research Council

(grant no. ERC-2008-AdG 233226 and ERC-2014-AdG670821 to R.A.), the Foundation for Scientific

Research at the University of Zurich (to P.J.W.), the Westlake Startup Grant (to T.G.), Zhejiang Provincial

Natural Science Foundation of China (Grant No. LR19C050001 to T.G.), Hangzhou Agriculture and Society

Advancement Program (Grant No. 20190101A04 to T.G.). A.B. and K.C. are supported by the German

Federal Ministry of Education and Research Grants Sybacol and PhosphoNetPPM.

809   Authors' contributions

A.B., T.G., P.J.W. and R.A. designed the project. P.J.W., T.G., Q.Z., C.E.F., N.J.R., A.C, D.R, J.H.R., C.F., K.S.,

C.P., T.H., A.L.M. and C.B. procured the samples and performed the experiments. K.C., T.G., Q.Z., U.W.,

R.S., N.C.T, K.O., L.C., L.M., M.R.M, M.M and A.B. designed and performed the statistical analyses with

critical inputs from C.Y., H.C., Q.Z., Y.Z., M.H. and other authors. K.C., A.B., T.G. and R.A. interpreted the

results. K.C., T.G., P.J.W., A.B. and R.A. wrote the manuscript with inputs from all co-authors. A.B., R.A.,

P.J.W. and T.G. supported and supervised the project.

Ethics declarations

Ethics approval and consent to participate

This study was approved by the Cantonal Ethics Committee of Zurich (KEK-ZH-No. 2008-0040), the

associated methods were carried out in accordance with the approved guidelines, and each patient has

signed an informed consent form.

Consent for publication

Not applicable.

Competing interests

R.A. holds shares of Biognosys AG, which operates in the field covered by the article. The research

groups of R.A. and T.G. are supported by SCIEX, which provides access to prototype instrumentation, and

Pressure Biosciences Inc., which provides access to advanced sample preparation instrumentation.

Supplementary information

Additional file 1: Supplementary text and supplementary figures.

Additional file 2: Table S1. Clinicopathological, immunological and other molecular information of the

39 PCa patients. (a) Overall clinicopathological characteristics. (b) Detailed information for each patient.

Pat: numeric patient ID; Pat_id: patient ID grouped by the overall grade. L: low grade; M: intermediate

grade; H: high grade; Overall_Gleason_GrGp: overall ISUP grade group; pT: tumor stage; pN: nodal

status; R: surgical margin status; Age_at_OP: age at operation; PSA_at_Diag: blood PSA level at

diagnosis; Time (months): RFS time. A value of 0 corresponds to patients excluded for the reasons

explained in the '**Methods**' section (see 'Patients and samples'); Status: status indicator. 1 means

recurrence; DX name: tissue region name; ImageName: name of the scanned images; index_tumor_id:

patient ID of TA1 (or TA); TA1_GrGp: grade group for TA1; T_GrGp: grade group for TA2.

Additional file 3: Table S2. Exome analysis of the peripheral blood cells and 105 prostate tumor

punches in 39 patients. (a) Allele frequencies (AF) of somatic single nucleotide variants (SNVs) that were

called by our bioinformatics pipeline. Genes with called SNV are indicated by an $AF > 0$. A value of 0

indicates that no SNV was found in the respective genes. In our data, no gene was found with more than

one called somatic SNV. (b) Number of samples per gene with called somatic SNV. (c) Protein domain

analysis using DAVID.

Additional file 4: Table S3. Copy number analysis of 105 PCa samples. (a) Log$_2$ ratios indicating the CNA

status are shown for all genes in all samples. Values were determined by overlapping gene locations with

CNA segments as calculated by CopywriteR. In case more than one segment overlapped with a gene,

number was chosen that had the highest absolute value. (b) Genes are shown with log$_2$ ratios higher

than 0.5 or lower than -0.5 in at least one sample.

Additional file 5: Table S4. RNA-seq analysis. (a) Log$_2$FCs (relative to all benign samples) for all genes

across the tumor samples. (b) mRIN score per sample generated using mRIN (v1.2.0). (c) ETS family gene

fusions observed in tumor samples using FusionCatcher: a value of 1 means that the fusion was observed

in the respective sample but not its corresponding benign sample, otherwise the value is 0.

Additional file 6: Table S5. Proteomics data of 210 PCa samples with duplicates. (a) Sample information

includes patient ID, clinical diagnosis, sample ID and batch design. (b) Protein matrix of log$_2$ scaled

intensity of 2,371 proteins quantified in 210 PCa samples.

Additional file 7: Table S6. Integration analysis of 66 tumor samples. (a) L1-norm of the 'centroid

vectors' in the three layers (CNA, mRNA and protein) across the four grade groups. (b) Information (i.e.

reference linking them to PCa, consistency between observed and reported effect and number of tumor

samples with CNAs) for the first 10 highest-scoring proteins (those with largest average absolute FCs

across all tumor specimens). (c) For each target gene the regulators with a non-zero beta coefficient

from the elastic net model are given. Further, the proportion of the elastic net predictors with the same

sign of Spearman correlation in our cohort and the independent study are noted for the two

independent PCa cohorts. (d) Consistently up-regulated genes in the high-grade tumors: for each of

these genes, there is a significant up-regulation of its FCs after network smoothing in the group G4/5

compared to the group G1 in all three layers (CNA, mRNA and protein). (e) Consistently down-regulated

genes in the high-grade tumors: for each of these genes, there is a significant down-regulation of its FCs

after network smoothing in the group G4/5 compared to the group G1 in all three layers (CNA, mRNA

and protein). (f) Chromosome information for the gene members of Network Component 1. (g) Mapping

from gene symbols to STRING identifiers.

References

1.   Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F:
Cancer incidence and mortality worldwide: sources, methods and major patterns in
GLOBOCAN 2012. *Int J Cancer* 2015, **136:**E359-386.
2.   Epstein JI, Zelefsky MJ, Sjoberg DD, Nelson JB, Egevad L, Magi-Galluzzi C, Vickers AJ, Parwani AV,
Reuter VE, Fine SW, et al: **A Contemporary Prostate Cancer Grading System: A Validated**
Alternative to the Gleason Score. *Eur Urol* 2016, **69:**428-435.
3.   Gordetsky J, Epstein J: **Grading of prostatic adenocarcinoma: current state and prognostic**
implications. *Diagn Pathol* 2016, **11:**25.
4.   Berger MF, Lawrence MS, Demichelis F, Drier Y, Cibulskis K, Sivachenko AY, Sboner A, Esgueva R,
Pflueger D, Sougnez C, et al: **The genomic complexity of primary human prostate cancer.**
Nature 2011, **470:**214-220.
5.   Barbieri CE, Baca SC, Lawrence MS, Demichelis F, Blattner M, Theurillat JP, White TA, Stojanov P,
Van Allen E, Stransky N, et al: **Exome sequencing identifies recurrent SPOP, FOXA1 and MED12**
mutations in prostate cancer. *Nat Genet* 2012, **44:**685-689.
6.   Robinson D, Van Allen EM, Wu YM, Schultz N, Lonigro RJ, Mosquera JM, Montgomery B, Taplin
ME, Pritchard CC, Attard G, et al: **Integrative clinical genomics of advanced prostate cancer.** *Cell*
2015, **161:**1215-1228.
7.   Baca SC, Prandi D, Lawrence MS, Mosquera JM, Romanel A, Drier Y, Park K, Kitabayashi N,
MacDonald TY, Ghandi M, et al: **Punctuated evolution of prostate cancer genomes.** *Cell* 2013,
153:666-677.
8.   Cancer Genome Atlas Research N: **The Molecular Taxonomy of Primary Prostate Cancer.** *Cell*
2015, **163:**1011-1025.
9.   Wedge DC, Gundem G, Mitchell T, Woodcock DJ, Martincorena I, Ghori M, Zamora J, Butler A,
Whitaker H, Kote-Jarai Z, et al: **Sequencing of prostate cancers identifies new cancer genes,**
routes of progression and drug targets. *Nat Genet* 2018, **50:**682-692.
10. Tomlins SA, Rhodes DR, Perner S, Dhanasekaran SM, Mehra R, Sun XW, Varambally S, Cao X,
Tchinda J, Kuefer R, et al: **Recurrent fusion of TMPRSS2 and ETS transcription factor genes in**
prostate cancer. *Science* 2005, **310:**644-648.
11. Grasso CS, Wu YM, Robinson DR, Cao X, Dhanasekaran SM, Khan AP, Quist MJ, Jing X, Lonigro RJ,
Brenner JC, et al: **The mutational landscape of lethal castration-resistant prostate cancer.**
Nature 2012, **487:**239-243.
12. Boutros PC, Fraser M, Harding NJ, de Borja R, Trudel D, Lalonde E, Meng A, Hennings-Yeomans
PH, McPherson A, Sabelnykova VY, et al: **Spatial genomic heterogeneity within localized,**
multifocal prostate cancer. *Nat Genet* 2015, **47:**736-745.
13. Cooper CS, Eeles R, Wedge DC, Van Loo P, Gundem G, Alexandrov LB, Kremeyer B, Butler A,
Lynch AG, Camacho N, et al: **Analysis of the genetic phylogeny of multifocal prostate cancer**
identifies multiple independent clonal expansions in neoplastic and morphologically normal
prostate tissue. *Nat Genet* 2015, **47:**367-372.
14. Lovf M, Zhao S, Axcrona U, Johannessen B, Bakken AC, Carm KT, Hoff AM, Myklebost O, Meza-
Zepeda LA, Lie AK, et al: **Multifocal Primary Prostate Cancer Exhibits High Degree of Genomic**
Heterogeneity. *Eur Urol* 2019, **75:**498-505.
15. Lindberg J, Klevebring D, Liu W, Neiman M, Xu J, Wiklund P, Wiklund F, Mills IG, Egevad L,
Gronberg H: **Exome sequencing of prostate cancer supports the hypothesis of independent**
tumour origins. *Eur Urol* 2013, **63:**347-353.

16.   Espiritu SMG, Liu LY, Rubanova Y, Bhandari V, Holgersen EM, Szyca LM, Fox NS, Chua MLK,
918   Yamaguchi TN, Heisler LE, et al: **The Evolutionary Landscape of Localized Prostate Cancers**
919   **Drives Clinical Aggression.** *Cell* 2018, **173:**1003-1013 e1015.
920   17.   Sinha A, Huang V, Livingstone J, Wang J, Fox NS, Kurganovs N, Ignatchenko V, Fritsch K, Donmez
921   N, Heisler LE, et al: **The Proteogenomic Landscape of Curable Prostate Cancer.** *Cancer Cell* 2019,
922   **35:**414-427 e416.
923   18.   Latonen L, Afyounian E, Jylha A, Nattinen J, Aapola U, Annala M, Kivinummi KK, Tammela TTL,
924   Beuerman RW, Uusitalo H, et al: **Integrative proteomics in prostate cancer uncovers robustness**
925   **against genomic and transcriptomic aberrations during disease progression.** *Nat Commun*
926   2018, **9:**1176.
927   19.   Iglesias-Gato D, Wikstrom P, Tyanova S, Lavallee C, Thysell E, Carlsson J, Hagglof C, Cox J, Andren
928   O, Stattin P, et al: **The Proteome of Primary Prostate Cancer.** *Eur Urol* 2016, **69:**942-952.
929   20.   Attard G, Parker C, Eeles RA, Schroder F, Tomlins SA, Tannock I, Drake CG, de Bono JS: **Prostate**
930   **cancer.** *Lancet* 2016, **387:**70-82.
931   21.   Goncalves E, Fragoulis A, Garcia-Alonso L, Cramer T, Saez-Rodriguez J, Beltrao P: **Widespread**
932   **Post-transcriptional Attenuation of Genomic Copy-Number Variation in Cancer.** *Cell Syst* 2017,
933   **5:**386-398 e384.
934   22.   Liu Y, Beyer A, Aebersold R: **On the Dependency of Cellular Protein Levels on mRNA**
935   **Abundance.** *Cell* 2016, **165:**535-550.
936   23.   Guo T, Kouvonen P, Koh CC, Gillet LC, Wolski WE, Rost HL, Rosenberger G, Collins BC, Blum LC,
937   Gillessen S, et al: **Rapid mass spectrometric conversion of tissue biopsy samples into**
938   **permanent quantitative digital proteome maps.** *Nat Med* 2015, **21:**407-413.
939   24.   Guo T, Li L, Zhong Q, Rupp NJ, Charmpi K, Wong CE, Wagner U, Rueschoff JH, Jochum W,
940   Fankhauser CD, et al: **Multi-region proteome analysis quantifies spatial heterogeneity of**
941   **prostate tissue biomarkers.** *Life Sci Alliance* 2018, **1**.
942   25.   Umbehr M, Kessler TM, Sulser T, Kristiansen G, Probst N, Steurer J, Bachmann LM: **ProCOC: the**
943   **prostate cancer outcomes cohort study.** *BMC Urol* 2008, **8:**9.
944   26.   Wettstein MS, Saba K, Umbehr MH, Murtola TJ, Fankhauser CD, Adank JP, Hofmann M, Sulser T,
945   Hermanns T, Moch H, et al: **Prognostic Role of Preoperative Serum Lipid Levels in Patients**
946   **Undergoing Radical Prostatectomy for Clinically Localized Prostate Cancer.** *Prostate* 2017,
947   **77:**549-556.
948   27.   Epstein JI, Egevad L, Amin MB, Delahunt B, Srigley JR, Humphrey PA, Grading C: **The 2014**
949   **International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading**
950   **of Prostatic Carcinoma: Definition of Grading Patterns and Proposal for a New Grading System.**
951   *Am J Surg Pathol* 2016, **40:**244-252.
952   28.   Epstein JI, Amin MB, Reuter VE, Humphrey PA: **Contemporary Gleason Grading of Prostatic**
953   **Carcinoma: An Update With Discussion on Practical Issues to Implement the 2014**
954   **International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading**
955   **of Prostatic Carcinoma.** *Am J Surg Pathol* 2017, **41:**e1-e7.
956   29.   Reznik E, Sander C: **Extensive decoupling of metabolic genes in cancer.** *PLoS Comput Biol* 2015,
957   **11:**e1004176.
958   30.   Taylor BS, Schultz N, Hieronymus H, Gopalan A, Xiao Y, Carver BS, Arora VK, Kaushik P, Cerami E,
959   Reva B, et al: **Integrative genomic profiling of human prostate cancer.** *Cancer Cell* 2010, **18:**11-
960   22.
961   31.   Zhang B, Wang J, Wang X, Zhu J, Liu Q, Shi Z, Chambers MC, Zimmerman LJ, Shaddox KF, Kim S,
962   et al: **Proteogenomic characterization of human colon and rectal cancer.** *Nature* 2014, **513:**382-
963   387.

32.   Geiger T, Cox J, Mann M: **Proteomic changes resulting from gene copy number variations in**
cancer cells. *PLoS Genet* 2010, **6:**e1001090.
33.   Juschke C, Dohnal I, Pichler P, Harzer H, Swart R, Ammerer G, Mechtler K, Knoblich JA:
Transcriptome and proteome quantification of a tumor model provides novel insights into
post-transcriptional gene regulation. *Genome Biol* 2013, **14:**r133.
34.   Stingele S, Stoehr G, Peplowska K, Cox J, Mann M, Storchova Z: **Global analysis of genome,**
transcriptome and proteome reveals the response to aneuploidy in human cells. *Mol Syst Biol*
2012, **8:**608.
35.   Cover TM, Thomas JA: *Elements of information theory.* John Wiley & Sons; 2012.
36.   Zhang JS, Gong A, Cheville JC, Smith DI, Young CY: **AGR2, an androgen-inducible secretory**
protein overexpressed in prostate cancer. *Genes Chromosomes Cancer* 2005, **43:**249-259.
37.   Liu Q, Harvey CT, Geng H, Xue C, Chen V, Beer TM, Qian DZ: **Malate dehydrogenase 2 confers**
docetaxel resistance via regulations of JNK signaling and oxidative metabolism. *Prostate* 2013,
73:1028-1037.
38.   Yang J, Song H, Chen L, Cao K, Zhang Y, Li Y, Hao X: **Integrated analysis of microfibrillar-**
associated proteins reveals MFAP4 as a novel biomarker in human cancers. *Epigenomics* 2019,
11:1635-1651.
39.   Senga S, Kobayashi N, Kawaguchi K, Ando A, Fujii H: **Fatty acid-binding protein 5 (FABP5)**
promotes lipolysis of lipid droplets, de novo fatty acid (FA) synthesis and activation of nuclear
factor-kappa B (NF-kappaB) signaling in cancer cells. *Biochim Biophys Acta Mol Cell Biol Lipids*
2018, **1863:**1057-1067.
40.   Pan Y, Liu Z, Feng Z, Hui D, Huang X, Tong D, Jin Y: **The overexpression of Rabl3 is associated**
with pathogenesis and clinicopathologic variables in hepatocellular carcinoma. *Tumour Biol*
2017, **39:**1010428317696230.
41.   Zhang W, Sun J, Luo J: **High Expression of Rab-like 3 (Rabl3) is Associated with Poor Survival of**
Patients with Non-Small Cell Lung Cancer via Repression of MAPK8/9/10-Mediated
Autophagy. *Med Sci Monit* 2016, **22:**1582-1588.
42.   Hofree M, Shen JP, Carter H, Gross A, Ideker T: **Network-based stratification of tumor**
mutations. *Nat Methods* 2013, **10:**1108-1115.
43.   Muniyan S, Chaturvedi NK, Dwyer JG, Lagrange CA, Chaney WG, Lin MF: **Human prostatic acid**
phosphatase: structure, function and regulation. *Int J Mol Sci* 2013, **14:**10438-10464.
44.   Araujo CL, Quintero IB, Ovaska K, Herrala AM, Hautaniemi S, Vihko PT: **Transmembrane**
prostatic acid phosphatase (TMPAP) delays cells in G1 phase of the cell cycle. *Prostate* 2016,
76:151-162.
45.   Qiu Y, Yang S, Pan T, Yu L, Liu J, Zhu Y, Wang H: **ANKRD22 is involved in the progression of**
prostate cancer. *Oncol Lett* 2019, **18:**4106-4113.
46.   Donald CD, Sun CQ, Lim SD, Macoska J, Cohen C, Amin MB, Young AN, Ganz TA, Marshall FF,
Petros JA: **Cancer-specific loss of beta-defensin 1 in renal and prostatic carcinomas.** *Lab Invest*
2003, **83:**501-505.
47.   Siltanen S, Fischer D, Rantapero T, Laitinen V, Mpindi JP, Kallioniemi O, Wahlfors T, Schleutker J:
ARLTS1 and prostate cancer risk--analysis of expression and regulation. *PLoS One* 2013,
8:e72040.
48.   Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A,
Santos A, Tsafou KP, et al: **STRING v10: protein-protein interaction networks, integrated over**
the tree of life. *Nucleic Acids Research* 2015, **43:**D447-D452.
49.   Vanunu O, Magger O, Ruppin E, Shlomi T, Sharan R: **Associating Genes and Protein Complexes**
with Disease via Network Propagation. *Plos Computational Biology* 2010, **6**.

50. Zhou D, Bousquet O, Lal TN, Weston J, Schölkopf B: **Learning with Local and Global Consistency.**
Advances in Neural Information Processing Systems 2004, **16:**312-328.
51. Baude A, Aaes TL, Zhai B, Al-Nakouzi N, Oo HZ, Daugaard M, Rohde M, Jaattela M: **Hepatoma-**
derived growth factor-related protein 2 promotes DNA repair by homologous recombination.
Nucleic Acids Res 2016, **44:**2214-2226.
52. Guler GD, Tindell CA, Pitti R, Wilson C, Nichols K, Kaiwai CT, Kim HJ, Wongchenko M, Yan Y, Haley
B: **Repression of Stress-Induced LINE-1 Expression Protects Cancer Cell Subpopulations from**
Lethal Drug Exposure. *Cancer Cell* 2017, **32:**221.
53. Slezak J, Truong M, Huang W, Jarrard D: **HP1gamma expression is elevated in prostate cancer**
and is superior to Gleason score as a predictor of biochemical recurrence after radical
prostatectomy. *BMC Cancer* 2013, **13:**148.
54. Ceol CJ, Yariv H, Judit JV, Steve B, Orlando DA, Valentine B, Lauriane F, Lin WM, Hollmann TJ,
Fabrizio F: **The SETDB1 histone methyltransferase is recurrently amplified in and accelerates**
melanoma. *Nature* 2011, **471:**513-517.
55. Sun Y, Wei M, Ren SC, Chen R, Xu WD, Wang FB, Lu J, Shen J, Yu YW, Hou JG, et al: **Histone**
methyltransferase SETDB1 is required for prostate cancer cell proliferation, migration and
invasion. *Asian J Androl* 2014, **16:**319-324.
56. Luo H, Cowen L, Yu G, Jiang W, Tang Y: **SMG7 is a critical regulator of p53 stability and function**
in DNA damage stress response. *Cell Discovery* 2016, **2:**15042.
57. Karanika S, Karantanos T, Li L, Wang J, Park S, Yang G, Zuo X, Song JH, Maity SN, Manyam GC:
Targeting DNA Damage Response in Prostate Cancer by Inhibiting Androgen Receptor-CDC6-
ATR-Chk1 Signaling. *Cell Reports* 2017, **18:**1970.
58. Diouf B, Cheng Q, Krynetskaia NF, Yang W, Cheok M, Pei D, Fan Y, Cheng C, Krynetskiy EY, Geng
H: **Somatic deletions of genes regulating MSH2 protein stability cause DNA mismatch repair**
deficiency and drug resistance in human leukemia cells. *Nature Medicine* 2011, **17:**1298-1303.
59. Yao S, Bee A, Brewer D, Dodson A, Beesley C, Ke Y, Ambroisine L, Fisher G, Møller H, Dickinson T:
PRKC-ζ Expression Promotes the Aggressive Phenotype of Human Prostate Cancer Cells and Is
a Novel Target for Therapeutic Intervention. *Genes Cancer* 2010, **1:**444-464.
60. Zhou W, Jiang Y, Zhu M, Hang D, Chen J, Zhou J, Dai J, Ma H, Hu Z, Jin G, et al: **Low-frequency**
nonsynonymous variants in FKBPL and ARPC1B genes are associated with breast cancer risk in
Chinese women. *Mol Carcinog* 2017, **56:**774-780.
61. De Robertis M, Mazza T, Fusilli C, Loiacono L, Poeta ML, Sanchez M, Massi E, Lamorte G, Diodoro
MG, Pescarmona E, et al: **EphB2 stem-related and EphA2 progression-related miRNA-based**
networks in progressive stages of CRC evolution: clinical significance and potential miRNA
drivers. *Mol Cancer* 2018, **17:**169.
62. Zeng Z, Yang H, Xiao S: **ACTL6A expression promotes invasion, metastasis and epithelial**
mesenchymal transition of colon cancer. *BMC Cancer* 2018, **18:**1020.
63. Castro MA, Dal-Pizzol F, Zdanov S, Soares M, Muller CB, Lopes FM, Zanotto-Filho A, da Cruz
Fernandes M, Moreira JC, Shacter E, Klamt F: **CFL1 expression levels as a prognostic and drug**
resistance marker in nonsmall cell lung cancer. *Cancer* 2010, **116:**3645-3655.
64. Overbye A, Skotland T, Koehler CJ, Thiede B, Seierstad T, Berge V, Sandvig K, Llorente A:
Identification of prostate cancer biomarkers in urinary exosomes. *Oncotarget* 2015, **6:**30357-
30376.
65. Deng Z, Wan M, Cao P, Rao A, Cramer SD, Sui G: **Yin Yang 1 regulates the transcriptional activity**
of androgen receptor. *Oncogene* 2009, **28:**3746-3757.
66. Fournier PG, Juarez P, Jiang G, Clines GA, Niewolna M, Kim HS, Walton HW, Peng XH, Liu Y,
Mohammad KS, et al: **The TGF-beta Signaling Regulator PMEPA1 Suppresses Prostate Cancer**
Metastases to Bone. *Cancer Cell* 2015, **27:**809-821.

67. Yang J, Lu C, Wei J, Guo Y, Liu W, Luo L, Fisch G, Li X: **Inhibition of KPNA4 attenuates prostate**
cancer metastasis. *Oncogene* 2017, **36:**2868-2878.
68. Liu W, Xie CC, Zhu Y, Li T, Sun J, Cheng Y, Ewing CM, Dalrymple S, Turner AR, Sun J, et al:
Homozygous deletions and recurrent amplifications implicate new genes involved in prostate
cancer. *Neoplasia* 2008, **10:**897-907.
69. Trasino SE, Harrison EH, Wang TT: **Androgen regulation of aldehyde dehydrogenase 1A3**
(ALDH1A3) in the androgen-responsive human prostate cancer cell line LNCaP. *Exp Biol Med*
(Maywood) 2007, **232:**762-771.
70. Bonaccorsi L, Luciani P, Nesi G, Mannucci E, Deledda C, Dichiara F, Paglierani M, Rosati F, Masieri
L, Serni S, et al: **Androgen receptor regulation of the seladin-1/DHCR24 gene: altered**
expression in prostate cancer. *Lab Invest* 2008, **88:**1049-1056.
71. Li H, Xiao D, Hu L, He T: **Association of CYP1A1 polymorphisms with prostate cancer risk: an**
updated meta-analysis. *Mol Biol Rep* 2012, **39:**10273-10284.
72. Aktas D, Hascicek M, Sozen S, Ozen H, Tuncbilek E: **CYP1A1 and GSTM1 polymorphic genotypes**
in patients with prostate cancer in a Turkish population. *Cancer Genet Cytogenet* 2004, **154:**81-
85.
73. Acevedo C, Opazo JL, Huidobro C, Cabezas J, Iturrieta J, Quinones Sepulveda L: **Positive**
correlation between single or combined genotypes of CYP1A1 and GSTM1 in relation to
prostate cancer in Chilean people. *Prostate* 2003, **57:**111-117.
74. Cao HM, Wan Z, Wu Y, Wang HY, Guan C: **Development and internal validation of a novel**
model and markers to identify the candidates for lymph node metastasis in patients with
prostate cancer. *Medicine (Baltimore)* 2019, **98:**e16534.
75. Kawamura N, Nimura K, Saga K, Ishibashi A, Kitamura K, Nagano H, Yoshikawa Y, Ishida K,
Nonomura N, Arisawa M, et al: **SF3B2-Mediated RNA Splicing Drives Human Prostate Cancer**
Progression. *Cancer Res* 2019, **79:**5204-5217.
76. Wen M, Kwon Y, Wang Y, Mao JH, Wei G: **Elevated expression of UBE2T exhibits oncogenic**
properties in human prostate cancer. *Oncotarget* 2015, **6:**25226-25239.
77. Kleppe A, Albregtsen F, Vlatkovic L, Pradhan M, Nielsen B, Hveem TS, Askautrud HA, Kristensen
GB, Nesbakken A, Trovik J, et al: **Chromatin organisation and cancer prognosis: a pan-cancer**
study. *Lancet Oncol* 2018, **19:**356-369.
78. Lee RS, Roberts CW: **Linking the SWI/SNF complex to prostate cancer.** *Nat Genet* 2013,
45:1268-1269.
79. Chuu C-P, Lin C-Y, Huang S-H, Tsai KK-C: **MO2-10-6ACTL6A is a novel oncogene and prognostic**
biomarker for prostate cancer. *Annals of Oncology* 2019, **30**.
80. Doyon Y, Selleck W, Lane WS, Tan S, Cote J: **Structural and functional conservation of the NuA4**
histone acetyltransferase complex from yeast to humans. *Mol Cell Biol* 2004, **24:**1884-1896.
81. Cai Y, Jin J, Florens L, Swanson SK, Kusch T, Li B, Workman JL, Washburn MP, Conaway RC,
Conaway JW: **The mammalian YL1 protein is a shared subunit of the TRRAP/TIP60 histone**
acetyltransferase and SRCAP complexes. *J Biol Chem* 2005, **280:**13665-13670.
82. Mortensen MM, Hoyer S, Lynnerup AS, Orntoft TF, Sorensen KD, Borre M, Dyrskjot L: **Expression**
profiling of prostate cancer tissue delineates genes associated with recurrence after
prostatectomy. *Sci Rep* 2015, **5:**16018.
83. Hussain M, Mateo J, Fizazi K, Saad F, Shore ND, Sandhu S, Chi KN, Sartor O, Agarwal N, Olmos D,
et al: **LBA12_PRPROfound: Phase III study of olaparib versus enzalutamide or abiraterone for**
metastatic castration-resistant prostate cancer (mCRPC) with homologous recombination
repair (HRR) gene alterations. *Annals of Oncology* 2019, **30**.
84. Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods* 2012, **9:**357-
359.

85.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
1108   Gabriel S, Daly M, DePristo MA: **The Genome Analysis Toolkit: a MapReduce framework for**
1109   **analyzing next-generation DNA sequencing data.** *Genome Res* 2010, **20:**1297-1303.
1110   86.   Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data.**
1111   *Bioinformatics* 2014, **30:**2114-2120.
1112   87.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1113   Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools.**
1114   *Bioinformatics* 2009, **25:**2078-2079.
1115   88.   Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.**
1116   *Bioinformatics* 2010, **26:**841-842.
1117   89.   Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, Gabriel S, Meyerson M,
1118   Lander ES, Getz G: **Sensitive detection of somatic point mutations in impure and**
1119   **heterogeneous cancer samples.** *Nat Biotechnol* 2013, **31:**213-219.
1120   90.   Saunders CT, Wong WSW, Swamy S, Becq J, Murray LJ, Cheetham RK: **Strelka: accurate somatic**
1121   **small-variant calling from sequenced tumor-normal sample pairs.** *Bioinformatics* 2012,
1122   **28:**1811-1817.
1123   91.   Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, Smigielski EM, Sirotkin K: **dbSNP: the NCBI**
1124   **database of genetic variation.** *Nucleic Acids Res* 2001, **29:**308-311.
1125   92.   Kuilman T, Velds A, Kemper K, Ranzani M, Bombardelli L, Hoogstraat M, Nevedomskaya E, Xu G,
1126   de Ruiter J, Lolkema MP, et al: **CopywriteR: DNA copy number detection from off-target**
1127   **sequence data.** *Genome Biol* 2015, **16:**49.
1128   93.   Noske A, Brandt S, Valtcheva N, Wagner U, Zhong Q, Bellini E, Fink D, Obermann EC, Moch H,
1129   Wild PJ: **Detection of CCNE1/URI (19q12) amplification by in situ hybridisation is common in**
1130   **high grade and type II endometrial cancer.** *Oncotarget* 2017, **8:**14794-14805.
1131   94.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR:
1132   **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2013, **29:**15-21.
1133   95.   Hartley SW, Mullikin JC: **QoRTs: a comprehensive toolset for quality control and data**
1134   **processing of RNA-Seq experiments.** *BMC Bioinformatics* 2015, **16:**224.
1135   96.   Feng H, Zhang X, Zhang C: **mRIN for direct assessment of genome-wide and gene-specific mRNA**
1136   **integrity from large-scale RNA-sequencing data.** *Nat Commun* 2015, **6:**7816.
1137   97.   Shao W, Guo T, Toussaint NC, Xue P, Wagner U, Li L, Charmpi K, Zhu Y, Wu J, Buljan M, et al:
1138   **Comparative analysis of mRNA and protein degradation in prostate tissues indicates high**
1139   **stability of proteins.** *Nat Commun* 2019, **10:**2524.
1140   98.   Sigurgeirsson B, Emanuelsson O, Lundeberg J: **Sequencing degraded RNA addressed by 3' tag**
1141   **counting.** *PLoS One* 2014, **9:**e91851.
1142   99.   Liao Y, Smyth GK, Shi W: **featureCounts: an efficient general purpose program for assigning**
1143   **sequence reads to genomic features.** *Bioinformatics* 2014, **30:**923-930.
1144   100.   Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD: **The sva package for removing batch effects**
1145   **and other unwanted variation in high-throughput experiments.** *Bioinformatics* 2012, **28:**882-
1146   883.
1147   101.   Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-seq**
1148   **data with DESeq2.** *Genome Biol* 2014, **15:**550.
1149   102.   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ,
1150   Pachter L: **Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts**
1151   **and isoform switching during cell differentiation.** *Nat Biotechnol* 2010, **28:**511-515.
1152   103.   Shao S, Guo T, Gross V, Lazarev A, Koh CC, Gillessen S, Joerger M, Jochum W, Aebersold R:
1153   **Reproducible Tissue Homogenization and Protein Extraction for Quantitative Proteomics Using**
1154   **MicroPestle-Assisted Pressure-Cycling Technology.** *J Proteome Res* 2016, **15:**1821-1829.

1155   104.   Rosenberger G, Koh CC, Guo T, Rost HL, Kouvonen P, Collins BC, Heusel M, Liu Y, Caron E,
1156   Vichalkovski A, et al: **A repository of assays to quantify 10,000 human proteins by SWATH-MS.**
1157   *Sci Data* 2014, **1:**140031.
1158   105.   Röst HL, Rosenberger G, Navarro P, Gillet L, Miladinovic SM, Schubert OT, Wolski W, Collins BC,
1159   Malmstrom J, Malmstrom L, Aebersold R: **OpenSWATH enables automated, targeted analysis of**
1160   **data-independent acquisition MS data.** *Nat Biotechnol* 2014, **32:**219-223.
1161   106.   Guo T, Luna A, Rajapakse VN, Koh CC, Wu Z, Liu W, Sun Y, Gao H, Menden MP, Xu C, et al:
1162   **Quantitative Proteome Landscape of the NCI-60 Cancer Cell Lines.** *iScience* 2019, **21:**664-680.
1163   107.   Blattmann P, Heusel M, Aebersold R: **SWATH2stats: An R/Bioconductor Package to Process and**
1164   **Convert Quantitative SWATH-MS Proteomics Data for Downstream Analysis Tools.** *PLoS One*
1165   2016, **11:**e0153160.
1166   108.   Johnson WE, Li C, Rabinovic A: **Adjusting batch effects in microarray expression data using**
1167   **empirical Bayes methods.** *Biostatistics* 2007, **8:**118-127.
1168   109.   Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P,
1169   von Mering C, Jensen LJ: **STRING v9.1: protein-protein interaction networks, with increased**
1170   **coverage and integration.** *Nucleic Acids Res* 2013, **41:**D808-815.
1171   110.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson
1172   E, et al: **Integrative analysis of complex cancer genomics and clinical profiles using the**
1173   **cBioPortal.** *Sci Signal* 2013, **6:**pl1.
1174   111.   Ycart B, Charmpi K, Rousseaux S, Fournié J-J: **Large scale statistical analysis of GEO datasets.**
1175   2014, **4:**113:111-119.
1176
1177

1178   Figure legends

1179

1180   **Figure 1. Proteogenomics analysis of 105 tissue regions from 39 PCa patients. a** Representative
1181   immunohistochemistry images of prostate tissues and the selection of BPH and tumorous tissue regions
1182   for genome, transcriptome and proteome analysis. **b** Kaplan-Meier curves for our cohort when the
1183   patients are stratified by the overall grade (left), the TA1 or TA grade group (middle) and the TA2 or TA
1184   grade group (right). Point-wise 95% confidence bands are shown for the whole range of time values.

1185

Figure 2. Molecular perturbation scores for point mutations, CNAs, transcriptome and proteome data.
a Distributions of the first type of molecular perturbation scores (DE_count's) for the four grade groups
(visualized as violin plots) at the mutation layer (upper left), CNA layer (upper right), mRNA layer (lower
left) and protein layer (lower right). Points represent the actual values. The horizontal lines correspond
to the median value in each of the four grade groups. **b** Distributions of the second type of molecular
perturbation scores (DE_sum's) for the four grade groups (visualized as violin plots) at the CNA layer
(upper left), mRNA layer (upper right) and protein layer (lower left). Points represent the actual values.
The horizontal lines correspond to the median value in each of the four grade groups. *P* values (in each
of the titles) show the significance of the one-sided Wilcoxon rank sum test where the values of G3 and
G4/5 are gathered together and compared to the values of G1 and G2 (also gathered together).

1196

Figure 3. **Within-group similarity at the different layers quantified by different similarity measures. a,**
b Distributions of the similarity scores between the individual tumor samples and the centroid using the
Pearson correlation (**a**) and MI (**b**) for the four grade groups (visualized as violin plots) at the CNA (upper
left), mRNA (upper right) and protein (lower left) layers. Points represent the actual values. The
horizontal lines correspond to the median value in each of the four grade groups. *P* values from the one-
sided Wilcoxon rank sum test comparing G4/5 versus G1: 0.0014 for the CNA, 0.89 for the mRNA and
0.053 for the protein layer with the Pearson correlation, and 0.027 for the CNA, 0.0052 for the mRNA
and 0.0081 for the protein layer with the MI. **c** Density plots of the FCs in the four grade groups for three
selected proteins among the 20 highest scoring (score: mean of the absolute FCs across all tumor
samples) proteins. Vertical lines correspond to the average FC in each of the four grade groups. The
selected proteins have more extreme FCs in the high-grade tumors (G3 and G4/5). **d** Heatmap of the
CNA matrix reduced to the significant regulators of the target gene ACPP output by the fitted elastic net
model (*i.e.* those with a non-zero beta coefficient). The columns are ordered based on the grade group
while there is a hierarchical clustering of the rows. The added colorbar depicts the mRNA FCs of the
target gene ACPP.

1212

Figure 4. **Consistently dysregulated sub-networks, Network Component 1 heatmaps and validation in**
three independent cohorts. a Sub-networks consistently dysregulated in high-grade compared to low-
grade tumors. There is a significant up-regulation of the FCs of the depicted genes (colored in red) after
network smoothing in the group G4/5 compared to the group G1 in all three layers (CNA, mRNA and
protein). **b** Same as in (**a**) but here there is a significant down-regulation of the FCs of the depicted genes
(colored in blue) after network smoothing in the group G4/5 compared to G1 in all layers. Functional
annotation of the sub-networks in (**a**) and (**b**) with more than one node is given. **c** Heatmap of the CNA
matrix reduced to the Network Component 1 genes. The columns are ordered based on the grade group.

The bottom colorbar depicts the effect size of each sample, *i.e.* its average FC across the genes of the
Network Component 1. The next colorbar represents the negative logarithm in base 10 of the *P* value
from the t-test. The above colorbar shows whether the result is significant: significant results are colored
in black and the rest in white. The top colorbar depicts the grade group of each sample. A black box has
marks the 'interesting' area with amplification of Network Component 1. Gray rectangles at the bottom
show the grade group of the patients (low, intermediate, high) where the samples have (mainly) come
from. **d, e** Same as in (**c**) but for the mRNA and the proteins. The samples removed due to degradation
(mRNA) are shown in gray. **f** Kaplan-Meier curves for 'altered' and 'unaltered' samples in the three
independent studies, TCGA (left), MSKCC (middle) and Aarhus (right) using the corresponding CNA data
when available (first row) and mRNA data (second row).

1231

Figure 5. Within-patient similarity at the different layers. a Distributions of the within-group similarities
for the four grade groups (visualized as violin plots) based on the Pearson correlation at the CNA layer
(upper left), mRNA layer (upper right) and protein layer (lower left). A 'violin' with the correlations
between TA1 and paired TA2 for the different patients has been added to all three plots and colored in
purple. Points represent the actual values. The horizontal lines correspond to the median value in each
of the groups. This analysis is similar to **Fig. 3a** – but it is not identical. *P* values from the one-sided
Wilcoxon rank sum test comparing the within-patient to the within-group similarities (where all values
from the four groups are gathered together): 8.97e-09 for the CNA, 4.42e-08 for the mRNA and 6.27e-04
for the protein layer. **b** The correlations between TA1 and paired TA2 for the different patients at one
layer are plotted against the corresponding correlations at another layer for each pair of layers: mRNA
versus CNA (upper left), protein versus CNA (upper right) and protein versus mRNA (bottom left). The
points are labeled and colored based on the overall grade in all plots; r: Pearson correlation.

1244

1245

1246   **Figure 1**

a

b

1247
1248

1249   **Figure 2**

1250

1251

1252

1253   **Figure 3**

1254

1255

1256
1257

1258   **Figure 4**

1259

1260

Functional annotation

Up-regulation
1. Chromatin remodelling
2. Mitochondrial import
3. RNA polymerase 3
4. unknown function in prostate
5. Actin related complex, Arp2/3 complex
6. Protein quality control
7. mRNA splicing
8. DNA damage response

Down-regulation
1. Glutathione metabolism
2. Cholesterol biosynthesis

1261

1262

1263

1264   **Figure 4 (*continued*)**

1265

1266

1267

1268   **Figure 5**

1269

1270
1271

1272