Coral reef restoration efforts in Latin American countries and territories Elisa Bayraktarov^{1*}, Anastazia T. Banaszak², Phanor Montoya Maya³, Joanie Kleypas⁴, Jesús E. AriasGonzález⁵, Macarena Blanco⁶, Johanna Calle Triviño^{5,6,15}, Nufar Charuvi⁷, Camilo Cortés Useche^{5,15}, Victor Galván⁸, Miguel A. García Salgado⁹, Mariana Gnecco³, Sergio D. Guendulain García², Edwin A. Hernández Delgado^{10,11,12}, José A. Marín Moraga⁴, María Fernanda Maya³, Sandra Mendoza Quiroz^{2,13}, Samantha Mercado Cervantes⁸, Megan Morikawa⁶, Gabriela Nava⁹, Valeria Pizarro¹⁴, Rita

- 8 I. Sellares-Blasco¹⁵, Samuel E. Suleimán Ramos¹⁰, Tatiana Villalobos Cubero⁴, María Villalpando¹⁵,
- 9 Sarah Frías-Torres^{16,17}
- 10
- 11¹ Centre for Biodiversity and Conservation Science, The University of Queensland, Brisbane,
- 12 Queensland, Australia
- 13 ² Investigación Integral para la Conservación de Arrecifes, Universidad Nacional Autónoma de
- 14 México (UNAM), Puerto Morelos, Quintana Roo, Mexico
- 15 ³ Corales de Paz, Cali, Colombia
- 16 ⁴ Raising Coral Costa Rica, Costa Rica
- 17 ⁵ Departamento de Recursos del Mar, Centro de Investigación y de Estudios Avanzados del Instituto
- 18 Politécnico Nacional, Mérida, Yucatán, Mexico
- 19⁶ Iberostar Hotels & Resorts, *Wave of Change*, Dominican Republic, Mexico
- 20 ⁷ Fundación Calipso, Taganga, Magdalena, Colombia
- 21 ⁸ Fundación Grupo Puntacana (FGPC), Punta Cana, La Altagracia Province, Dominican Republic
- 22 ⁹ Oceanus A.C., Chetumal, Quintana Roo, Mexico
- 23 ¹⁰ Sociedad Ambiente Marino (SAM), San Juan, Puerto Rico
- 24 ¹¹ University of Puerto Rico, Department of Environmental Sciences, San Juan, Puerto Rico
- 25 ¹² University of Puerto Rico, Centre for Applied Tropical Ecology and Conservation, San Juan, Puerto
- 26 Rico
- 27 ¹³ SECORE International, Inc., Hilliard, Ohio, USA
- 28 ¹⁴ Fundación para la Investigación y Conservación Biológica Marina ECOMARES, Cali, Valle del Cauca,
- 29 Colombia
- 30 ¹⁵ Fundación Dominicana de Estudios Marinos (FUNDEMAR), Santo Domingo, Dominican Republic
- 31 ¹⁶ Vulcan, Seattle, Washington, USA
- 32 ¹⁷ Smithsonian Institution, Washington, District of Columbia, USA
- 33
- 34 * Corresponding author
- 35 Email address: e.bayraktarov@uq.edu.au

- 37
- 38

39 Abstract

40 Coral reefs worldwide are degrading due to climate change, overfishing, pollution, coastal 41 development, bleaching and diseases. In areas where natural recovery is negligible or protection 42 through management interventions insufficient, active restoration becomes critical. The Reef Futures 43 symposium in 2018 brought together over 400 reef restoration experts, businesses, and civil 44 organizations, and galvanized them to save coral reefs through restoration or identify alternative 45 solutions. The symposium highlighted that solutions and discoveries from long-term and ongoing coral 46 reef restoration projects in Spanish-speaking countries in the Caribbean and Eastern Tropical Pacific 47 were not well known internationally. Therefore, a meeting of scientists and practitioners working in 48 these locations was held to compile the data on the extent of coral reef restoration efforts, advances 49 and challenges. Here, we present unpublished data from 12 coral reef restoration case studies from 50 five Latin American countries, describe their motivations and techniques used, and provide estimates 51 on total annual project cost per unit area of reef intervened, spatial extent as well as project duration. 52 We found that most projects used direct transplantation, the coral gardening method, micro-53 fragmentation or larval propagation, and aimed to optimize or scale-up restoration approaches (51%) 54 or provide alternative, sustainable livelihood opportunities (15%) followed by promoting coral reef 55 conservation stewardship and re-establishing a self-sustaining, functioning reef ecosystem (both 56 13%). Reasons for restoring coral reefs were mainly biotic and experimental (both 42%), followed by 57 idealistic and pragmatic motivations (both 8%). The median annual total cost from all projects was 58 \$93,000 USD (range: \$10,000 USD - \$331,802 USD) (2018 dollars) and intervened a median spatial 59 area of 1 ha (range: 0.06 ha - 8.39 ha). The median project duration was 3 years; however, projects 60 have lasted up to 17 years. Project feasibility was high with a median of 0.7 (range: 0.5 - 0.8). This 61 study closes the knowledge gap between academia and practitioners and overcomes the language 62 barrier by providing the first comprehensive compilation of data from ongoing coral reef restoration 63 efforts in Latin America.

64 Introduction

Active restoration, the process of assisting the recovery of an ecosystem that has been degraded, damaged, or destroyed [1], may be increasingly necessary on coral reefs, once it has been determined that the natural recovery of corals is hindered [2]. The goal of any restoration action is to eventually establish self-sustaining, sexually reproducing populations with enough genetic variation enabling them to adapt to a changing environment [3-5].

Coral reef restoration may play a particularly important role where coral species are threatened with extinction. The Caribbean Elkhorn coral, *Acropora palmata*, and Staghorn coral, *A. cervicornis*, were once widely distributed and among the major reef-building species in the region [6]. Both species are now listed as Critically Endangered on the International Union for Conservation of Nature (IUCN) Red List [7] as a result of major losses in cover of both species throughout the Caribbean since the 1970s [8]. Management programmes have not aided in the recovery of *A. palmata* [9]. In this context, active restoration of these species is essential to recover their ecosystem functions in the Caribbean region.

77 Several techniques are used for the restoration of coral reefs. The most common techniques are based 78 on asexual methods such as direct transplantation, coral gardening, and micro-fragmentation [10]. An 79 alternative technique, larval propagation, is based on the collection of gametes and the consequent 80 culturing of embryos and larvae, after which the coral spat are either grown in ex situ aquaria to larger-81 sized colonies or are outplanted onto degraded reefs at approximately one month old [11]. While the 82 techniques used to restore coral reefs are reviewed elsewhere (e.g. [10, 12-14]), here we focus on 83 direct transplantation, coral gardening, micro-fragmentation, and larval propagation as the 84 techniques most-commonly employed by the case studies in the study area. One of the oldest 85 techniques used in coral reef restoration is direct transplantation of corals [15], which involves the 86 harvesting of coral colonies from a donor site and their immediate transplantation to a restoration 87 site or re-attaching colonies that have been dislodged by a ship grounding, storm or hurricane [16]. 88 The coral gardening approach was developed to scale-up restoration while reducing the stress on

89 donor colonies. Fragments of corals are harvested from donor colonies, grown in nurseries to a 90 threshold size [17] before being transplanted onto a degraded reef [18, 19]. Nurseries can be ocean-91 based (in situ) or land-based (ex situ). In situ nurseries are typically located at well-lit sites safe from 92 predation, storm surges, and wave energy, and are regularly maintained and cleaned by physical 93 removal of algal growth [20]. However, strategic siting of ocean nurseries can promote the 94 recruitment of fish assemblages that eat biofouling, thus may significantly reduce person-hours spent 95 in nursery cleaning [21]. In situ nurseries can have many shapes and sizes. For example, they can 96 consist of floating mid-water structures built using ropes, mesh or cages [21-24], structures placed on 97 concrete, tables or frames [25], PVC 'trees' [26], PVC grids or dead coral bommies [27]. Ex situ 98 nurseries typically use flow-through large aquaria or raceways, and require continuous access to 99 electricity, water quality monitoring, and control of temperature and light availability [28]. Micro-100 fragmentation is an approach especially useful for slow-growing massive corals. This technique 101 involves the fragmentation of parts of a massive coral donor to yield multiple ~ 1 cm² fragments. The 102 fragments are placed close to each other on either artificial substrates or on the surface of dead coral 103 colonies. The micro-fragments, as they recognize neighbouring fragments as kin, grow towards each 104 other and fuse [29]. Ideally, they are outplanted to the degraded reef at a size of ~6 cm² [29, 30]. Larval 105 propagation involves the breeding of corals from eggs and sperm. Studies describing this technique 106 typically report the use of raceways with seawater flow-through systems where coral spawn is 107 collected from the wild, fertilization is assisted, embryos are cultured to larvae, which are settled onto 108 substrates and then transported and seeded onto a degraded coral reef [31-33]. This process has also 109 been referred to as larval enhancement, sexual propagation, sexual coral cultivation or larval 110 reseeding [12]. As an emerging larval propagation technique, larval restoration concentrates coral 111 larvae over enhancement plots on the degraded reef to facilitate coral larvae settlement directly to 112 the substrate, without the need of laboratory facilities [34]. The main advantages of the larval 113 propagation techniques are that they increase the genetic diversity among restored coral populations 114 thus enabling increased rates of adaptation and improved resilience in the context of climate change

[35], and they have the potential to be used over large scales while reducing the cost [31]. Also, theydo not cause damage to the parent colonies.

117 While efforts in the USA, Australia or places where European scientists conduct their research are well 118 described in the published literature and disseminated at conferences, there is a paucity of 119 documentation on coral reef restoration projects carried out by practitioners in the Caribbean and 120 Eastern Tropical Pacific. Reasons for this lack of exchange may be the language barrier, lack of interest 121 in knowledge transfer between higher and lower income countries or cultural differences as well as 122 lack of funding. In 2018, the Reef Futures symposium was held in the Florida Keys, USA and attended 123 by over 400 delegates. The aim of this international meeting was to 'bring together experts from 124 around the world to share the latest science and techniques for coral reef restoration while kicking off 125 a global effort to dramatically scale-up the impact and reach of restoration as a major tool for coral 126 reef conservation and management'. The conference was organized by the Coral Restoration 127 Consortium, which is 'a community of practice comprised of scientists, managers, coral restoration 128 practitioners, and educators dedicated to enabling coral reef ecosystems to survive the 21st century 129 and beyond' [36]. Within the Reef Futures conference, we convened a meeting of scientists and 130 practitioners involved in active coral reef restoration in the Latin- and Centro-American Caribbean as 131 well as the Eastern Tropical Pacific to fill the knowledge gap between academia and practitioners in 132 the region and overcome the language barriers in coral reef restoration. Here, we showcase the 133 advances and share the lessons learned from 12 restoration case studies from the Caribbean and 134 Eastern Tropical Pacific. We provide a comprehensive compilation of unpublished data from coral reef 135 restoration efforts where we outline the techniques that were employed, the motivations and 136 objectives of each project, total project cost per unit area per year, spatial extent of intervention, and 137 project duration. This work provides the most complete data set on total project cost and feasibility 138 of coral reef restoration from practical cases that may guide decisions required to establish new 139 restoration projects in the future.

140 Approach

141 Data collection

- 142 The co-authors of this work contributed data and descriptions of their restoration projects which
- 143 constitute the case studies used here. The coral reef restoration projects were carried out in Latin
- 144 American countries and territories in the Caribbean and Eastern Tropical Pacific (Fig. 1). The data
- 145 obtained included estimates on total annual project cost, spatial extent of area intervened, project
- duration, and an estimate on the project reaching specific objectives within a fixed period of time. The
- 147 motivations for each restoration project were adopted from [10, 37, 38] and classified as biotic,
- 148 experimental, idealistic, legislative, and pragmatic (Table 1).
- **Table 1:** Five motivation categories for carrying out coral reef restoration projects and examples.

| Motivation category | Examples | | | | | |
|---------------------|--|--|--|--|--|--|
| Biotic | Biodiversity enhancement (e.g., native species, habitat creation, ecosystem connectivity, ecological resilience) | | | | | |
| Experimental | Improve restoration approaches, technologies, and methods. Answer ecologically-based research questions | | | | | |
| Idealistic | Cultural reasons (e.g., recreation, tourism, medicinal/ceremonial substances, spiritual importance, aesthetic value) | | | | | |
| | Social reasons (e.g., community involvement, job creation, nature education, environmental outreach) | | | | | |
| | Political reasons (e.g., raising environmental profile) | | | | | |
| Legislative | Restoration after environmental impact (e.g., ship-grounding, mining, oil spill, hurricane damage) | | | | | |
| | Biodiversity offset (e.g., threatened species, threatened ecological communities) | | | | | |
| Pragmatic | Enhance ecosystem services (e.g., fisheries production) | | | | | |
| | Enhance ecosystem services (e.g., water quality improvement, pollution prevention) | | | | | |
| | Enhance ecosystem services (e.g., coastal protection, erosion control, bank stabilisation) | | | | | |
| | Enhance ecosystem services (e.g., carbon sequestration, carbon offsets) | | | | | |

151 The objectives of coral reef restoration projects can be highly diverse and dependent on the specific 152 project as well as its location. In this study, the restoration practitioners were asked to provide the 153 objectives for their restoration projects, which were specific, measurable, achievable, repeatable and 154 time-bound (SMART: [3]). We modified the six primary objectives observed by Hein et al. [39] into the 155 following categories: 1) enhance ecosystem services for the future; 2) optimize/scale-up restoration 156 approaches; 3) promote coral reef conservation stewardship; 4) provide alternative, sustainable 157 livelihood opportunities; 5) reduce coral population declines and ecosystem degradation; and 6) re-158 establish a self-sustaining, functioning reef ecosystem.

159 The total estimated project cost includes both capital and operating costs. Capital costs are those used 160 for planning, land acquisition, construction, and financing [40]. These may also include costs for 161 laboratory/infrastructure, boats and dive equipment. Operating costs are those used for maintenance, 162 monitoring, equipment repair and replacement [40] and may include salaries, housing for 163 scientific/implementation teams, air for SCUBA tanks, gasoline for boat engines, and replacement of 164 computers. Coral reef practitioners were asked to estimate the total cost for restoration interventions 165 based on the guidelines for standardised reporting of costs for management interventions for 166 biodiversity conservation [41] and are provided as United States Dollars (USD) per hectare of coral 167 reef intervened per year in 2018 USD.

The project spatial extent is the coral reef area intervened by the restoration project and is reported in hectares. Spatial extent is not provided for each project since not all restoration case studies have an objective to increase the area of restored habitat. For instance, some projects are aimed at developing new restoration techniques, using coral nurseries as a tool to stimulate public awareness and engagement, for educational purposes, or as a tourist attraction.

173 The project duration is the time during which the restoration project has existed until the present, or 174 the time during which the restoration cost was budgeted for and is provided in years. All projects 175 described here are ongoing and active throughout 2019.

176 The feasibility is the likelihood that each specific project objective can be reached successfully with 177 the interventions at hand and within the outlined project duration. It is ideally measured as the 178 likelihood of success in returning the ecosystem function and resilience of an ecosystem through 179 restoration [42]. This overall restoration project feasibility is rarely reported in the published literature 180 because a standardised method to measure restoration success is largely missing [40]. Here, 181 restoration practitioners estimated the feasibility of the restoration interventions they employed to 182 achieve their specific project objectives. Feasibility is given as a ratio between 0 and 1 and can be 183 interpreted as the likelihood of success to reach a specific objective within the duration of the 184 restoration project. Practitioners provided a minimum, maximum and the best guess for the project 185 feasibility.

186 Results

187 Data from a total of 12 coral reef restoration projects carried out by practitioners in the Spanish-188 speaking Caribbean and Eastern Tropical Pacific were compiled and are summarised in Table 2. The 189 supplementary material contains more detailed information about each restoration case study. 190 Information was gathered from Colombia (Alianza Coralina Taganga, Corales de Paz, and ECOMARES), 191 Costa Rica (Raising Coral Costa Rica), the Dominican Republic (FUNDEMAR, the Iberostar Group, and 192 Fundación Grupo Puntacana), Mexico (Oceanus A.C., CORALIUM at Universidad Nacional Autónoma 193 de México, and the Iberostar & CINVESTAV Group), and Puerto Rico (Sociedad Ambiente Marino) 194 (Figure 1). Note that the Fundación Grupo Puntacana has two restoration programs of which one is 195 focused on coral gardening (Program 1) and one is directed towards micro-fragmentation (Program 196 2). These were treated as independent projects for analytical purposes. The restoration projects use 197 techniques that include direct transplantation, coral gardening, micro-fragmentation, and larval 198 propagation (Figure 2; Supplementary information Table S1).

199 **Table 2**: Summary of the 12 restoration projects in the Caribbean and Eastern Tropical Pacific. Cost values are given in 2018 USD. More detailed information

200 can be found in the supplementary material. Abbreviations: Fundación Dominicana de Estudios Marinos, Inc. (FUNDEMAR), Fundación Grupo Puntacana

201 (FGPC), and Sociedad Ambiente Marino (SAM).

| Country, Location, Organization | Technique employed (type of nursery) | Targeted coral species | Motivations | Specific project objectives | Strategy for outplanting | Spatial extent of project | Estimated project budget and funding bodies/partners | Estimated project feasibility |
|--|--|--|--|--|--|---|--|--|
| | | | | Implemented and in progress as | of 2019 | | | |
| Colombia, Taganga, Caribbean Sea, Alianza Coralina Taganga | A floating mid-water nursery built of PVC tubes is located 5 m below the water surface and anchored to the sandy bottom at 13 m depth. Coral fragments are produced by micro- fragmentation of donor colonies, which are then attached to cement cookies and outplanted once they reach a diameter of 7 cm. Each cement cookie is connected via a plastic screw to a mesh frame in the coral nursery with a carrying capacity of 50 – 80 cookies per frame. Corals of opportunity are presently used. | Montastraea cavernosa; Porites porites; Millepora sp. | The primary motivation is idealistic following social reasons such as community education and engagement. The secondary motivation is experimental i.e., to improve management and develop standardized restoration protocols. | 1) to develop a training centre for the sustainable use of marine resources and ecological restoration; 2) to establish a community-based coral reef monitoring system for Taganga Bay and coral nurseries therein with the possible expansion of monitoring to other areas; 3) to develop a management plan for Taganga Bay as a marine reserve, which is governed by the local community; and 4) to create a financed organization, which aims to facilitate long-term ecological reef restoration and research in Taganga Bay | Corals grown in the mid- water nursery will be outplanted by drilling holes in the natural substrate with a pneumatic drill and inserting the plastic nails of the cement cookies carrying the coral fragments into the holes (supported by epoxy glue where necessary). All outplanted corals at the restoration site will be monitored at least once per month while they reattach to the natural substrate. | The spatial extent of the project is currently a matter of negotiation that depends on the capacity to recruit coral gardeners from the local community and to obtain a permit to carry out the ecological restoration work in Taganga Bay. | An estimated budget for the project is \$500,000 USD over the next two years. Forty percent of this budget is self-funded by local stakeholders to accelerate the capacity of coral growth and maintenance of coral outplanting through local capacity building. A large proportion of the estimated project budget will be directed to activities such as education, community engagement and training while a minor part will be focused on growing and outplanting corals to the restoration site. Members from the local community are trained as coral gardeners to identify corals of opportunity, carry out coral micro-fragmentation and maintain the nursery. | best guess = 0.5; minimum = 0.2; maximum = 0.9 |
| Colombia, San Andres and Providencia Islands, Caribbean Sea, Corales de Paz | The project employs the coral gardening technique. Rope nurseries are floating at 4 to 6 m below the water surface (Fig. 2a). Micro- fragmentation is also | Acropora palmata; A. cervicornis; Porites porites; Madracis decactis; Pseudodiploria | The primary project motivation is biotic i.e., to enhance coral reef biodiversity, while the secondary motivation is pragmatic i.e., to enhance the ecosystem | 1) to generate an annual stock of > 5,000 coral fragments from four reef-building species per island; 2) to transplant 5,000 coral colonies per ha ⁻¹ yr ⁻¹ per island from year two of the project, for a total of 30,000 coral transplants over six | Nursery-grown corals as well as micro-fragments will be outplanted to the reef using a unique mix of marine cement and a colloid adjuvant to improve fluidity and | The total spatial extent is six hectares (area of outplant) by year four distributed over three hectares at each of the two islands. | The total estimated budget is \$900,000 USD resulting in an annual expenditure of \$37,500 USD ha ⁻¹ yr ⁻¹ . Partners include the Secretary of Agriculture and Fisheries from the Government of the Archipelago of San Andrés, | best guess = 0.6; minimum = 0.5; maximum = 0.9 |

| | being employed during the outplanting phase. | clivosa; P. astreoides; Millepora complanata | services of fisheries, tourism and coastal protection by the local coral reefs. | hectares in three years; 3) to achieve a 25% increase in selected coral reef health indicators (i.e., live coral cover, coral settlement, fish biomass, and rugosity) at the intervened sites; 4) to design and implement an effective system of protection and restoration of intervened reef areas that encourages conservation and contributes to the sustainability of benefits derived from these reefs together with relevant social actors; and 5) to quantify the ecosystem services of intervened reef areas in current and future scenarios of intervention, variability and climate change | reduce runoff. Outplanting density will be 5,000 individuals per ha of reef. | | Providencia and Santa Catalina, the provincial environmental authority CORALINA, Conservation International Colombia, and the NGO Corales de Paz. Both islands are within the Seaflower Biosphere Reserve in the Colombian Caribbean. The first phase of the project was financed by all participating organizations with support from MasBosques and BanCO2. | |
|---|--|---|---|--|--|--|---|--|
| Costa Rica, Golfo Dulce, Eastern Tropical Pacific, Raising Coral Costa Rica | The project employs the techniques coral gardening and micro- fragmentation. Coral fragments are often collected as fragments of opportunity and are grown in tree and rope nurseries, after which they are outplanted to the restoration site (Fig. 2f). | Pocillopora sp.; Porites evermanni; P. lobate; Pavona gigantea; Pavona frondifera; Psammocora sp. | The primary motivation of the project is experimental with the rationale to improve coral propagation techniques for growing corals in the Eastern Tropical Pacific, with an emphasis on answering questions of ecological concern. The secondary motivation is biotic i.e., to enhance biodiversity, ecosystem connectivity, and ecological resilience. | to define the best coral propagation and restoration techniques; 2) to establish a coral restoration program in Costa Rica; to facilitate coral reef research to improve restoration work; and to integrate local communities into coral restoration projects | Branching corals grown in the nursery are outplanted onto the substrate with cable ties attached to large nails. Future endeavours will attach corals grown on ropes in rope-nurseries directly to the substrate without separating them from the ropes. Massive and encrusting corals, which are grown on ceramic plugs, are outplanted by drilling holes into the substrate and inserting the stem of the plug with a small amount of marine epoxy or cement. | The project aims to restore 10 reef patches of 200-500 m ² each within the next three years equalling a maximum intervened area of 0.5 ha. | The total project cost over the last 2.5 years was \$120,000 USD. If in kind support (such as accommodation, university technical support, volunteer time, etc.) is included, these costs would be 100% higher, i.e., a total of \$240,000 USD. The annual project budget was \$35,000 USD for 2018, which was mostly composed of salaries (\$15,000 USD) and logistics such as travel and boat rental (\$15,000 USD). The remaining \$5,000 were needed for material and consumables. The project is mainly financed by private donations and Raising Coral Costa Rica is currently initiating a fundraising campaign call to restore several thousand corals for Costa Rica and to scale- up coral propagation and restoration efforts. | best guess = 0.8; minimum = 0.6; maximum = 0.9 |
| Dominican Republic, | The techniques coral gardening (rope and steel | Acropora cervicornis; | The primary motivation of the project is biotic | 1) to propagate coral tissue of the endangered <i>A. cervicornis</i> using | Corals grown in eight underwater nurseries | FUNDEMAR's restoration project | FUNDEMAR is a largely self- sustainable organization that has | best guess = 0.7; minimum = 0.4; |

| Bayahibe, Caribbean Sea, FUNDEMAR | rod nurseries (Fig. 2b) and larval propagation (seeding coral recruits after cultivation using <i>in</i> <i>situ</i> SECORE Intdesigned floating pools (Fig. 2c)) and <i>ex situ</i> in a wet lab) are used to restore local coral populations. | A. palmata; Colpophyllia natans; Diploria labyrinthiformis; Orbicella annularis; O. faveolata; | with the rationale of biodiversity enhancement. The secondary motivation is legislative focused on restoration after environmental impact and as a biodiversity offset. However, the project has also idealistic motivations for cultural, social and political reasons. | the genetically diverse coral nurseries; 2) to enhance coral populations' genetic diversity and resilience to environmental changes by outplanting substrates (either SECORE's cement or ceramic substrates or FUNDEMAR's cement "cookies") with settled coral recruits. | (holding more than 3 km of tissue) are outplanted with nails, cable ties and epoxy glue techniques. FUNDEMAR has already carried out coral outplanting at 12 restoration sites (Table S3 , supplementary material). The project is monitoring two spawning sites used to deliver the spawning stocks for rearing coral larvae in an <i>ex situ</i> facility (Table S2 , supplementary material). Corals reared by larval propagation settle on hand-made cement or ceramic substrates which are nailed or seeded on the reef. | aims to intervene on at least 0.5 hectares of degraded coral reef per year on a restoration schedule of one coral colony per square meter, transplanting around 2,000 <i>A. cervicornis</i> coral fragments of around 20-30 cm in diameter and seeding 2,000 recruit substrates (from 3-5 different species). | developed strategic alliances with private and public national and international institutions and with financial support for implementation of new projects. The total project cost per year is around \$51,800 USD which includes costs for maintenance, staff salaries, boats, and keeping up the facility but excludes in-kind support from volunteers. Part of this funding (\$18,400 USD per year) comes from the local, national and international alliances that FUNDEMAR has established. The coral reef restoration project has been financed by two grants and alliances with other organizations carrying out coral reef restoration. | maximum = 0.9 |
|---|---|---|---|--|--|---|--|--|
| Dominican Republic, Bayahibe, Caribbean Sea, The Iberostar Group | The Iberostar Group uses the coral gardening technique to restore coral reefs and is currently in charge of two coral nurseries at two locations in the Dominican Republic. One of the nurseries is an <i>in situ</i> nursery and the other one is an <i>in situ</i> nursery connected to a land-based facility for research and to evaluate the genetic diversity of corals in the nursery. | Acropora cervicornis; A. palmata; Diploria labyrinthiformis; Porites porites; P. astreoides; Orbicella annularis; O. faveolata; Agaricia agaricites | Reef restoration in one of the main pillars of the <i>Wave of Change</i> movement initiated in 2018 by the international hotel chain Iberostar. <i>Wave of</i> <i>Change</i> aims at contributing to the conservation of the oceans by engaging with the tourism sector. The main motivations of this project are experimental, biotic and idealistic. | 1) to determine current intraspecific diversity; 2) to enhance intra- and inter-specific diversity; 3) to maintain <i>in situ</i> and <i>ex situ</i> genetic bank; 4) to engage hotel clients and staff; 5) to characterise individual physiological traits of corals; 6) to enhance resilience in restored reefs | Despite the nursery still being developed, fragments of <i>A</i> . <i>cervicornis, P. porites</i> and <i>A. agaricites</i> have been maintained with 100% survival rate for three months to date. | No information on the spatial extent of area intervened is available yet, because the transplantation strategy is currently being developed. As part of this strategy, the group aims to identify coral genotypes potentially less susceptible to environmental stress. | The estimated budget spent from the beginning of the project in May 2018 to March 2019 is \$100,000 USD, including materials and construction of the land-based facility and salaries. In 2018 alone, \$40,000 USD were spent on construction (excluding salaries). The project is privately financed with annual funds destined to the <i>Wave of Change</i> initiative. From 2016 onwards, the group has taken over the responsibility of an <i>in situ</i> coral nursery that was set up and maintained in collaboration with FUNDEMAR. Research will be carried out by the scientific team of <i>Wave of Change</i> as well as by collaborating with international | best guess = 0.5; minimum = 0.2; maximum = 0.8 |

| | | | | | | | scientists who can use the facility to conduct their studies. The interspecific diversity of nursery- grown corals is being addressed through genetic analyses in collaboration with the University of California at Santa Barbara. Finally, the facility will be used as an outreach centre to teach and raise awareness about topics such as coral biology, the importance of reefs, threats to marine ecosystems, etc. to hotel clients and staff. | |
|---|--|--|---|---|--|--|--|--|
| Dominican Republic, Punta Cana, Caribbean Sea, Fundación Grupo Puntacana (Program 1) | Nursery fragments are grown on A-Frames, tables and ropes at water depths between 3.5 and 5 m. | Acropora. cervicornis; A. palmata; Orbicella spp.; Porites spp.; Pseudodiploria spp. | The primary motivation of the project program is biotic and is focused on the enhancement of biodiversity and resilience. The secondary motivation is idealistic and concentrates on social reasons (e.g. development of alternative income opportunities for local communities and improved user experience for tourism, fisheries, ecosystem services etc.). | 1) to prevent a potential local or regional disappearance of coral species through enhancement of successful sexual reproduction using fast growing, genetically diverse, nursery-reared fragments; 2) to reduce local environmental problems such as marine pollution, unsustainable wastewater treatment, uncontrolled fisheries and tourist carrying capacity; 3) to train local community members such as fishermen or dive centre staff in the installation and maintenance of coral nurseries and outplanting of nursery-grown corals; 4) to replicate the lessons learned in other parts of the Dominican Republic and other Caribbean island nations to improve coral reef restoration in Punta Cana; 5) to generate alternative income opportunities for members of the local community, especially for local fishermen | Nursery fragments are outplanted on the local, patchy, fringing reef using cable ties and galvanized nails at similar depths to fragments growing in the nursery. | Since 2014, a total of 8,810 A. cervicornis fragments (representing 5,394 linear meters of coral tissue) have been transplanted over almost 0.44 ha of degraded reef. Sexual reproduction has been consistently observed at both the nursery and surrounding outplanted sites. | The total estimated budget for 2018 was around \$93,000 USD resulting in \$211,363 USD ha ⁻¹ yr ⁻¹ when extrapolated from the actual area intervened (0.44 ha). This budget includes salaries, material, equipment, consumables, fixed- assets, infrastructure upkeep, and project-related expenses. For the next 3 years (2019 – 2021), if grant proposals submitted are approved, there is a plan to scale-up coral reef restoration efforts. These include an increase in the number of <i>A. cervicornis</i> fragments outplanted to approximately 5,000 fragments per year. FGPC estimates that over the next 3 years about 15,000 fragments can be transplanted over one ha of natural coral reef. The total estimated budget for the time interval 2019 – 2021 will be approximately \$950,000 USD, thus equalling the total cost of \$313,500 USD ha ⁻¹ yr ⁻¹ . The coral reef restoration programs are supported by the general budget of FGPC. Additional support is provided by private donations, | best guess = 0.8; minimum = 0.5; maximum = 0.9 |

| | | | | | | | national and international grants and institutions such as Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ), The Nature Conservancy (TNC), Counterpart International (CPI), Caribbean Hotel and Tourism Association, Global Giving, and InterAmerican Development Bank (IDB) among others. | |
|---|---|--|--|--|--|---|--|--|
| Dominican Republic, Punta Cana, Caribbean Sea, Fundación Grupo Puntacana (Program 2) | The donor colonies (fragments of opportunity) are cut by a diamond band saw into approximately one cm ² pieces, which are then attached to cement discs made in-house and deposited into flow- through raceways. This program consists of three phases. The first phase identifies the best conditions for high survival and fast growth of the micro-fragments in the <i>ex situ</i> nursery and develops the protocols for the approach. The second phase identifies adequate restoration sites and develops outplanting protocols. During this phase, methods, tools and equipment are tested. The third phase will scale-up outplanting efforts with micro-fragments. | Pseudodiploria strigosa; P. clivosa; Porites astreoides; P. furcata; Orbicella annularis; Montastraea cavernosa | The primary motivation of this project program is biotic (biodiversity enhancement), while the secondary is experimental (improve restoration approach, technology and methods). A tertiary motivation is idealistic (environmental education and outreach for the local community and tourists). | As above | Micro-fragments will be outplanted using established protocols. | By the end of the third phase, an estimate of 5,000 micro-fragments will be outplanted annually covering up to 200 m ² per year. | The total budget for 2018 was around \$30,000 USD. The project duration is three years and the total estimated budget is \$850,000 USD (pending grant approvals). Funding bodies/partners are as above. | best guess = 0.6; minimum = 0.4; maximum = 0.9 |
| Mexico, Chetumal, Caribbean Sea, Oceanus A.C. | This project uses the coral gardening approach where corals are grown in <i>in situ</i> nurseries. Fragments of opportunity | Acropora palmata, A. cervicornis; A. prolifera; Porites spp.; Agaricia | The primary motivation of this program is pragmatic i.e., to recover reef ecosystem health and promote | 1) to promote the rehabilitation of coral reefs through transplantation of 10,000 colonies every year at different sites in the Gulf of Mexico and the Mexican | The restoration sites are selected according to a set of established criteria. Every new site requires between three | The spatial extent of total area intervened for all restoration activities since 2014 is | The total project budget was estimated to average \$150,000 USD per year since 2014 to outplant 10,000 colonies every year with an outplanting schedule | best guess = 0.8; minimum = 0.5; maximum = 0.9 |

are rescued from donor areas and grown in the nurseries. Small concrete bases are attached to the reef and then corals from the nurseries are fixed to these structures (Fig. 2d). To increase the diversity at the restoration sites and promote natural resilience to climate change and local stressors, the program identifies the genetic material (genotypes) from healthy donor populations using the microsatellite technique. At least five genotypes are combined at each restoration site.

spp.; Orbicella recovery of spp.; Diploria spp. environmen

environmental services of the reef as well as associated species populations and biomass with special emphasis on recovering protected and no-take areas. The secondary motivation is legislative, i.e., to restore coral reefs after environmental impacts such as ship-grounding or hurricanes depending on the location and site.

Caribbean; 2) to strengthen the resilience and adaptation potential of coral reefs by increasing diversity on restoration sites through the identification of genetic material from healthy donor populations that could be naturally resilient to climate change and local stressors; 3) to secure community and reef managers' engagement to build local restoration groups that work based on a self-sustainable strategy to multiply efforts, increasing benefits to local communities in the short and midterm as well as helping the activities of the program to be maintained for a longer term

until colonies of the first and second generation have grown to reproduce sexually. Every year, monitoring is carried out before and after transplantation at each of the sites to evaluate the survival and growth of restored corals. The overall average of transplant survival has been about 80%. At the oldest restoration sites initiated from 2013 onwards and maintained by the program, the outplanted coral fragments, which initially had average sizes of between 7 and 10 cm, have now (in 2019) grown to an average size of 30 cm in diameter. Some outplants have reached a diameter of up to 110 cm (Fig. 2e). About 30% of the transplants evaluated in 2019 at all sites had a size of 20 cm in diameter on average indicating that they have reached a reproductive size.

and five years of work

estimated as 6.3 ha to date.

of one coral colony per square meter. Therefore, the annual budget was estimated at \$150,000 USD ha⁻¹ yr⁻¹. The restoration work of Oceanus A.C. is mainly artisanal and requires intensive maintenance to achieve results. Therefore, restoration efforts can only be sustained if the local community is involved to guarantee restoration success. The restoration program has initiated the formation of local restoration groups mainly consisting of members of the local fishing communities and other local organizations as well as the private sector (e.g., hotels) to support the restoration efforts. The program also seeks to engage local communities, service providers such as diving shops, hoteliers and managers to build local restoration groups and form a restoration network that helps increase restoration efforts along the Mesoamerican Reef. Establishing this network and applying different restoration strategies depending on the local stakeholder involved is envisioned to allow the program to become self-sustainable in the long term. The main partners of Oceanus A.C. for the development and scalingup of the program have been the Comisión Nacional de Áreas Naturales Protegidas (CONANP), Summit Foundation, the Mesoamerican Reef Fund, Fairmont Mayakobá and OHL Group, with local partners such as Acuario de Veracruz, Fundación de Parques y Museos de Cozumel, hotels from Playa del Carmen

Mexico, Mexican Caribbean, Caribbean Sea, CORALIUM, Universidad Nacional Autónoma de México

reproduction with the production of sexual recruits for restoration efforts beginning in 2011. Subsequently, it has focused on the development of low-cost techniques to scale-up the production of coral sexual recruits. This involves gamete collection in the wild, assisted fertilization and embryo husbandry in ex situ aquaria followed by outplanting of the sexual recruits to degraded reef sites. Restoration trials involve outplanting sexual recruits produced annually in the laboratory since 2011. From 2011 to 2014, the recruits were grown to juvenile size (up to 10 cm maximum diameter) in ex situ aquaria located in the Xcaret Ecopark. These colonies are now sexually mature as evidenced by the production of gametes in 2019. Since 2014, the research, in collaboration with SECORE International, has focused on scaling-up production and reducing costs by

Since 2007. CORALIUM

has been studying the

basic biology of coral

Acropora palmata; Orbicella faveolata; O. annularis; Diploria labyrinthiformis; Pseudodiploria strigosa This project undertakes science-based research to promote and scaleup best practices for coral restoration using sexual recruits (experimental motivation) and to increase genetic

motivation) and to increase genetic diversity in restoration efforts in the face of global climate change (biotic motivation). 1) to reduce costs of techniques using larval propagation of corals 100-fold; 2) to conduct research to improve survivorship of sexual recruits 20-fold; 3) to scale-up coral restoration techniques to ecologically significant scales over a 10-year period

To reduce costs. coral larvae are settled onto the artificial substrates and outplanted two weeks post-settlement (one-polyp stage). The substrates are placed manually into natural gaps formed by the reef framework without using cement or resin. New substrate designs are in the process of being tested to increase recruit survival from 0.1% at one-year postsettlement currently to a target of 10% and to improve substrate retention in the reef framework.

CORALIUM. in collaboration with SECORE International and Experiencias XCARET Aquarium have outplanted coral sexual recruits with sizes ranging from one polyp to colonies with an estimated volume of 500 cm³ on eight degraded reefs along the Mexican Caribbean from Cancun to Xcalak (Table S7, supplementary material). In total, the area of outplants corresponds to 0.15

hectares.

(Mayakobá chain) and from Mahahual and Xcalak, the Xcalak community, and tourist services providers from Cozumel, Puerto Morelos and Veracruz.

The costs for the production and outplanting of sexual recruits between 2014 and 2018 is estimated at \$15,000 USD per year and equals \$100,000 USD ha-1 yr-1. CORALIUM's research and restoration efforts have been funded by Universidad Nacional Autónoma de México, Comisión Nacional de Áreas Naturales Protegidas, Consejo Nacional de Ciencia y Tecnología, Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, Alianza World Wildlife Fund - Fundación Carlos Slim, SECORE International, The Nature Conservancy and Experiencias XCARET. Through communication and outreach, CORALIUM has promoted the message that coral reef conservation efforts and adaptive protected areas management are key to any coral restoration efforts. In collaboration with SECORE International and The Nature Conservancy, CORALIUM has implemented an annual coral reproduction course focused at students, coral restoration practitioners and other stakeholders.

best guess = 0.7; minimum = 0.6; maximum = 0.9

outplanting sexual recruits at the one polyp stage settled onto tetrapodshaped substrates, designed by SECORE International.

Coral nurseries were

established in year 2003

palmata; A.

Puerto Rico, Culebra Island, Caribbean Sea, Sociedad Ambiente Marino

prolifera; and operated continuously for 15 years Dendrogyra until the devastating cylindrus; O. impacts by hurricanes annularis; O. Irma and María in 2017. faveolata; Following the hurricanes, Madracis tree coral nurseries were aurentenra; established at a depth of 9 Eusmilia m. Additional tree nurseries were fastigiata established at a depth of 6-8 m, and at a depth of 7-12 m, to prevent further damage from coral bleaching and storm swells. Microfragmentation methods and direct coral cuttings have also been employed since the recent expansion. Direct transplantation has been conducted for emergency outplanting of fragments and/or detached colonies generated by vessel groundings, winter swells or hurricanes.

The primary motivation A. cervicornis; A. of the project is biotic (i.e., to enhance biodiversity, coral reef connectivity and ecosystem resilience). The secondary motivation is experimental (i.e., Porites divaricata; testing alternative methods and designs, with aims to answer ecological research questions). The tertiary motivation is pragmatic (i.e., enhance the ecosystem services by improving shallowwater essential fish habitat, restoring depleted fisheries, restoring hurricane mechanical impacts, recovering shallow reef accretion following the 2005 post-bleaching mass coral mortality event; enhancing carbon sequestration, tourism. and coastal protection of local coral reefs). Also, an important local motivation is to restore coral reef ecological functions within areas formerly impacted by

military training

1) to expand the annual stock in the nurseries of A. cervicornis to 8,000 colonies, of A. palmata to 2,500 colonies, D. cylindrus to 500 colonies, and O. annularis to 500 colonies; 2) to restore approximately 3 ha of degraded reef per year till 2022; 3) to outplant a minimum of 20,000 colonies of four species grown in the nurseries by year 2022, including 13,300 colonies of A. cervicornis, 5,000 colonies of A. palmata, 1,200 colonies of D. cylindrus, and 500 colonies of O. annularis; 4) to achieve a 25% increase in selected coral reef health indicators (i.e., live coral cover, fish biomass, and rugosity) at intervened sites for A. cervicornis and A. palmata; 5) to design and implement an effective community-based plan for the rehabilitation of intervened reef areas, which encourages conservation and rehabilitation of ecosystem functions, and to contribute to the sustainability of the benefits of coral reefs; 6) to quantify the ecosystem services of intervened reef areas in current and future scenarios of intervention, variability and climate change

Overall, in the time span of 2003-2017 approximately 60,000 coral colonies (mostly A. cervicornis) were harvested and outplanted to coral reefs in Culebra Island. Nursery-grown corals, fragments of opportunity of multiple species, as well as micro-fragments and cuttings are directly outplanted to the reef using Portland marine cement mixed with lime to neutralize pH. Cable ties and masonry nails are also used in the case of A. cervicornis. An outplanting schedule with a density of one individual per square meter of reef for A. cervicornis and of one colony per four square meters for other species is often followed.

The project has intervened an area of ca. 6 ha, but many of these corals were lost during the 2017 hurricanes. The projected spatial extent of reef rehabilitation by year 2023 in total will be 8.4 ha, with a potential to increase the area intervened depending on funding and on communitybased volunteer support.

The funds projected towards restoration for the period of 2019 to 2023 are \$1,327,206 (2018 USD), resulting in \$158,189 USD per restored ha per year or an estimated investment of \$50.26 USD per coral colony. These figures are based on the direct funds spent without accounting for in-kind contributions from the community. The real total estimated budget (including community-based in-kind support) for the period of 2019 to 2022 is \$2,311,280 (2018 USD) resulting in a total annual expenditure of \$275,480 (2018 USD) ha-1 or a total estimated expense of \$87.53 per coral colony. Coral nurseries have been historically managed by SAM. and since 2011. in direct collaboration with the Centre for Applied Tropical Ecology and Conservation (CATEC) of the University of Puerto Rico – Río Piedras Campus, under a memorandum of agreement with the Puerto Rico Department of Natural and Environmental Resources (PRDNER), and NOAA Restoration Centers (NOAA-RC). The first 4-year sub-project of the post-hurricane long-term phase of the project will be financed through multiple sources, as described above. It will also involve extensive volunteer work. through a combination of

best guess = 0.7; minimum = 0.5; maximum = 0.9

activities. Finally, the project is motivated by an idealistic rationale due to cultural reasons (i.e., community-based aim to restore formerly bombarded grounds by the U.S. Navy which used local coral reefs in Culebra Island to support naval training activities between 1901 and 1975, rescue and stewardship of local coral reefs) and due to social reasons (i.e., fostering increased community involvement, job creation, nature education, environmental outreach, hands-on training in coral farming and reef rehabilitation methods). More recently, the project is being motivated by legislative reasons (i.e., restoration of A. palmata and A. cervicornis as part of mitigatory compensation project).

strategies involving students, fishermen, NGOs, and an internship program. SAM also plans to involve the hospitality sector. There will also be a large focus on a combination of outreach, educational and hands on strategies to prepare the next generation of coral farmers and coral reef restoration researchers in Puerto Rico.

Planned work


| Colombia, Gorgona National Natural Park, Eastern Tropical Pacific, ECOMARES | The project is presently gathering scientific information for future needs on coral restoration in the area. | Pocillopora damicornis; Pavona clavus | The project is motivated by experimental reasons to improve restoration approaches for their use at Gorgona National Natural Park once restoration efforts are necessary. | 1) to determine the feasibility of coral nurseries in the area; 2) to determine the minimum coral fragment size of <i>P. damicornis</i> for successful survival and growth in coral nurseries; 3) to find the optimal fragment size for outplanting in terms of survival | The group's expertise in outplanting has been focused in the most abundant coral species in the area, <i>P.</i> <i>damicornis</i> . For this coral species, Portland cement mixed with sand | N/A | The cost for running the projects have been lower than expected because they are mostly experimental and have not carried out formal coral reef restoration activities yet. In 2018 the budget was \$10,000 USD. Partners: Universidad del Valle, Universidad | best guess = 0.7; minimum = 0.5; maximum = 0.9 |
|--|--|---|--|---|--|-----|--|--|
|--|--|---|--|---|--|-----|--|--|

| | | | | and coral growth; 4) to determine the effect of fish predation on <i>P.</i> <i>damicornis</i> during the outplanting; 5) to evaluate the use of enriched substrates for the massive coral species <i>P. clavus</i> | and freshwater was used. So far, no information is available to determine the spatial extent (area) of restored habitat that will be obtained. | | Javeriana de Cali, and Gorgona National Natural Park. | |
|---|--|---|---|--|--|-----|---|-----|
| Mexico, Cozumel National Natural Park, Caribbean Sea, The Iberostar & CINVESTAV Group: | The project aims to start with the development of four genotyped coral nurseries, two for <i>Acropora palmata</i> (3 and 5 m water depth) and two for <i>A. cervicornis</i> (10 and 13 m water depth). Each nursery will have 5 structures with a carrying capacity of approximately 40 fragments each enabling growth of 800 corals at a time. | Acropora palmata; A. cervicornis; Pseudodiploria spp.; Siderastrea spp.; Diploria labyrinthiformis; Orbicella spp. | Coral reef restoration envisioned by both groups is mainly motivated by experimental, biotic (i.e., enhance biodiversity, ecosystem connectivity, and ecological resilience), idealistic and pragmatic reasons (i.e., enhanced water quality and ecosystem services, shallow-water essential fish habitat, restore depleted fisheries, enhanced tourism, and coastal protection of local coral reefs. | to develop genotyped coral nurseries, which represent the coral diversity at Cozumel Island; to establish sufficient material in the coral nurseries to develop activities for education, research, technological innovation, recreation and tourism; 3) to yield sufficient material for the establishment of transplant zones; to collect gametes during the spawning season for larval rearing and use the larval propagation technique to grow sexual recruits at the transplantation site | N/A | N/A | This is a collaboration between the Iberostar Group and CINVESTAV Group with the Cozumel National Natural Park and the Mexican Secretariat of Environment and Natural Resources (SEMARNAT). The program will engage with local communities, universities, government entities and tourism service providers to gather sustained funding into the future. The group is open to new partners interested in participating in the project. | N/A |

The primary motivations to carry out the coral reef restoration projects are biotic and experimental to equal parts (41.7%), followed by idealistic and pragmatic reasons (both 8.3%). Biotic (36.3%) and experimental (27.3%) reasons were important secondary motivations, followed by legislative reasons (18.2%), and pragmatic/idealistic motivations (both 9.1%). All except for one of the projects reported secondary motivations. The tertiary motivations reported by 5 of the 12 projects were mainly pragmatic (80.0%) and idealistic (20.0%).

Most projects have specific objectives to optimize/scale-up restoration approaches (51.1%), followed by providing alternative, sustainable livelihood opportunities (14.9%), and then in equal parts to promote coral reef conservation stewardship and re-establish a self-sustaining, functioning reef ecosystem (12.8%). The objectives to enhance ecosystem services for the future and the reduction of population decline and ecosystem degradation accounted for only 4.2% each of the specific project objectives. The median total cost from all projects per year is \$93,000 USD (± \$32,731 SE) ranging between

\$10,000 USD and \$331,802 USD. The median spatial extent of coral reef restoration intervention is 1.0
ha (± 1.3 ha SE) ranging between 0.06 ha and 8.39 ha. Project duration was as short as 1 year and
could be as long as 17 years with the median project duration of 3 years (± 1.5 years SE) to reach the
project targets. Projects reported a median feasibility of 0.7 (± 0.03 SE) ranging from 0.5 to 0.8 (Table
3).



- 222 Figure 1: Map of coral reef restoration projects in Spanish-speaking Latin American countries and
- 223 territories.

221



224

Figure 2: Types of nurseries described in the text. a) Floating rope nurseries used in San Andrés and Providencia islands for large-scale coral gardening (Photo: Corales de Paz); b) rope nurseries by FUNDEMAR in Dominican Republic (Photo: Greg Asner); c) FUNDEMAR's floating *in situ* coral larvae rearing tank (Photo: Paul Selvaggio); d) Oceanus A.C. diver outplants nursery grown corals in Veracruz,

Mexico (Photo: Oceanus A.C.); e) outplanted *Acropora palmata* coral in Puerto Morelos, Mexico
(Photo: Oceanus A.C.); Raising Coral Costa Rica's tree nurseries in Costa Rica (Photo: David Garcia).

Table 3: Summary of total annual project costs, spatial extent of coral reef area intervened, project
 duration and feasibility from 12 case studies in the Spanish-speaking Caribbean and Eastern Tropical
 Pacific (Fundación Grupo Puntacana's restoration programs were treated as two independent
 projects). Error is given as standard error (± SE). Abbreviation: number of observations (N).

| | Total cost per year (2018 USD) | Spatial extent (ha) | Project duration (yrs) | Feasibility (best guess) |
|--------|-----------------------------------|------------------------|---------------------------|-----------------------------|
| Median | 93,000 (± 32,731) | 1.00 (± 1.30) | 3.0 (± 1.5) | 0.7 (± 0.03) |
| Min | 10,000 | 0.06 | 1 | 0.5 |
| Max | 331,802 | 8.39 | 17 | 0.8 |
| Ν | 11 | 7 | 12 | 11 |

236 Discussion

Here we present the first comprehensive assessment of coral reef restoration projects in Spanishspeaking countries and territories of the Caribbean and Eastern Tropical Pacific (ETP), which are already being implemented or are in the initiation phase. These projects were identified through an open call for participation at the Reef Futures conference in December 2018, which aimed to bring together a large international community to develop and implement solutions to the global coral reef crisis.

We describe 12 coral reef restoration case studies in the Caribbean and Eastern Tropical Pacific that employ coral reef restoration techniques including direct transplantation, coral gardening, microfragmentation and larval propagation (Supplementary information, **Table S1**). With a median total project cost per year of \$93,000 USD, spatial extent of 1 ha, duration of 3 years and overall project feasibility of 0.7, we show that coral reef restoration projects in these countries are more costeffective, have overcome the barriers of scaling-up restoration interventions, are persistent through time, and have a higher likelihood of success than reported from previous literature [10, 12, 40]. For instance, the most recent published literature review on coral reef restoration presented a median value of \$400,000 (2010 USD) to restore 1 ha (10,000 m²) of coral reef, project duration of 1 year, an area intervened of 0.01 ha (108 m²), and survival of restored corals as an item-based success indicator of 0.61 [10].

254 The objectives for coral reef restoration are often undocumented in the published literature, thus 255 extracting data on the objectives from published papers may lead to skewed results. For example, 256 Hein et al. [39] reviewed 83 published coral reef restoration studies and observed that 60% of the 257 studies reported on evaluating the biological response of the coral reef ecosystem to transplantation 258 (outplanting) as a main project objective. The remaining 40% of studies included the following 259 objectives: 1) to accelerate reef recovery post-disturbance (18%), 2) to re-establish a self-sustaining, 260 functioning reef ecosystem (48%), 3) to mitigate coral loss prior to a known disturbance (18%), and 4) 261 to reduce population declines and ecosystem degradation (15%). In comparison, we observed that 262 when data are elicited directly from restoration practitioners, most coral reef restoration projects in 263 the Caribbean and Eastern Tropical Pacific had the following objectives: 1) to optimize or scale-up 264 restoration approaches (51.1%), followed by 2) to provide alternative, sustainable livelihood 265 opportunities (14.9%). Similarly, the projects presented here were mostly motivated by biotic reasons 266 such as to enhance biodiversity and experimental reasons (both 41.7%), followed by 267 idealistic/pragmatic reasons (both 8.3%). In contrast, most motivations to restore coral reefs extracted 268 from the published literature were dominated by experimental reasons, such as to improve the 269 restoration approach and answer ecological research questions (65.3%) [10]. Many restoration 270 projects presented here focused on harnessing social or economic benefits from coral reef restoration 271 such as involving the community through inclusion in activities or educational programs to raise 272 awareness or to provide alternative, sustainable livelihood opportunities for local communities. An 273 assessment of social, economic, and cultural benefits derived from the restoration of coral reefs has 274 been largely ignored by the published literature, which has mostly concentrated on outcomes related to the ecology or described endeavours to improve restoration technology [10]. The present work is an attempt to bridge the gap between academics and practitioners. Academics tend to be more focused on small-experimental coral reef restoration attempts to answer questions of ecological concern, whereas practitioners are more focused on optimising and scaling-up restoration. Bridging the gap between academics and practitioners has been identified as critical for many fields of conservation [43, 44].

281 Coral reef restoration in the Caribbean and Eastern Tropical Pacific face challenges similar to those of 282 restoration efforts elsewhere in the world. For instance, the Intergovernmental Panel on Climate 283 Change (IPCC) concluded that, if no action is taken to reduce CO_2 emissions, coral reefs would decline 284 by 70-90 % with global warming of 1.5°C above pre-industrial levels, whereas virtually all coral reefs 285 (> 99 percent) would be lost with 2°C warming within the next 50 years [45]. Thus, while actions to 286 reduce CO_2 emissions are drastically needed, restoration with heat resilient species is regarded as a 287 key strategy to rehabilitate the ecological function and ecosystem services provided by coral reefs 288 [35]. In addition to climate change, coral reef restoration in the Caribbean and ETP face other 289 challenges such as overfishing, sedimentation, pollution, and non-sustainable coastal development 290 [46-51]. The recent outbreak of Scleractinian Coral Tissue Loss Disease (SCTLD) has decimated coral 291 populations and is of major concern to those attempting to restore corals in the Caribbean. Since its 292 onset in 2017, SCTLD has caused widespread mortality of corals, especially in the Florida Reef Tract 293 and the Gulf of Mexico [52, 53]. The vectors causing this disease or how it can be prevented are 294 currently unknown but are most likely bacterial [52]. A further challenge to the restoration of coral 295 reefs in the Caribbean and ETP is the apparent lack of funding and funding strategies. None of the 296 countries have cohesive national plans for the restoration of coral reefs similar to the Reef Restoration 297 and Adaptation Plan in Australia which has invested AUD \$100 million in 2018 to develop, trial, and 298 deploy coral reef restoration interventions for the Great Barrier Reef (GBR) [54].

299 Despite the impediment of limited financial resources, considerable advances in coral reef restoration, 300 both scaling-up of interventions and optimisation of techniques, have been achieved in Colombia, 301 Costa Rica, Dominican Republic, Mexico and Puerto Rico. For instance, one of the largest and longest 302 running projects (18 years) has plans to restore up to 8.4 ha, requiring outplanting 10.000 corals or up 303 to 8,000 coral settlement bases with coral larvae per year. These interventions were led by pioneering 304 environmental NGOs and foundations, who often procured un-paid volunteers to carry out much of 305 the work. The interventions were also enabled by strong partnerships initiated by the champion 306 organization with universities (e.g. Universidad Nacional Autónoma de México, University of Puerto 307 Rico, Universidad del Valle, Universidad Javeriana de Cali, Universidad de Costa Rica), conservation 308 management bodies and regulators (e.g. Natural Parks administrations, Departments of Natural and 309 Environmental Resources and the United States National Oceanic and Atmospheric Administration), 310 associations (e.g. Fishers Association, Caribbean Hotel and Tourism Association), national and 311 international business partners (e.g. SECORE International), international environmental NGOs (e.g. 312 Conservation International, The Nature Conservancy), tourist service providers (e.g. the Iberostar 313 Group), private donations (e.g. Global Giving), international grant schemes (e.g. from Deutsche 314 Gesellschaft für Internationale Zusammenarbeit, Counterpart International, InterAmerican 315 Development Bank (IDB)) and in large part with local community groups. Coral reef restoration still 316 remains an underfunded area in the Spanish-speaking countries and territories of the Caribbean and 317 ETP despite the ecosystem services restored coral reefs could provide for the regions such as food, 318 tourism income, protection against storms and wave surges [55, 56], and reduction in insurance 319 premiums by offering coastal protection [57].

There are a few caveats that need to be considered when assessing the data within the present work. First, this review does not contain an exhaustive list of interventions in the Spanish-speaking countries and territories of the Caribbean and ETP. Additional projects exist or are planned, but were not aware of, or chose to not participate in our open call. Second, the projects presented here varied in their specific objectives, best practice protocols, and monitoring, which hindered their comparison. For 325 example, some projects were designed to improve and optimise the restoration approach 326 (experimental projects), while others were more operational, i.e., aimed to scale-up the restoration 327 of coral reefs by using already established restoration techniques. Furthermore, the projects used 328 different best practice protocols or key indicators of restoration success, such as size of transplant and 329 density of transplants which made a direct comparison between the projects difficult. Some projects 330 lacked monitoring milestones to evaluate the survival, cover and health conditions of outplanted 331 corals beyond year one. Yet, post-restoration monitoring is an imperative method needed to confirm 332 that outplanted corals are self-sustaining which, from an evolutionary perspective, is the ultimate goal 333 of any restoration effort [3-5]. Third, evaluation of the overall project feasibility or the likelihood of 334 success to reach specific project objectives is naturally linked to local conditions and circumstances, 335 thus may be a subjective measure directly related to the experience of the practitioner. More 336 quantitative measures of overall project feasibility (e.g., based on measurements) would be a 337 considerable improvement over the qualitative (derived from expert elicitation) approach.

338 Prior to any conservation action, a prioritisation of interventions based on decision-support 339 frameworks is recommended to help practitioners increase their planning rigor, project accountability, 340 stakeholder participation, transparency in decisions, and learning [58]. Cost-effectiveness analysis is 341 such a tool that allows for the evaluation and prioritisation of conservation interventions [59]. This 342 analysis relates the costs of a project to its key outcomes or benefits i.e., the specific measures of 343 project effectiveness [59, 60]. Although this work includes all data required for a cost-effectiveness 344 analysis (see Supplementary material), we considered that comparing the different projects against 345 each other will be inappropriate given the variety of their project objectives (e.g. experimental vs. 346 operational) and the lack of standardisation in reporting on cost, feasibility and key outcomes.

Future collaborations between academics, local communities and practitioners will be crucial if we want to achieve restoration at meaningful ecological, spatial and social scales [61]. Unfortunately, the language barrier often inhibits such collaborations. For instance, Amano et al. [62] argues that

languages are still a major barrier to global science by showing that more than 35% of the knowledge in conservation is missed by those who only look at peer-reviewed literature in English. Many practitioners who carry out large-scale coral restoration projects only convey their knowledge in the form of unpublished reports and grey literature [10], which adds another level of complexity to the loss of information on restoration efforts. Here we close this gap by accessing this knowledge and overcoming the language barrier.

356

357 Conclusions

358 Although not previously highlighted by the published literature, there are many coral reef restoration 359 projects currently in progress in the Spanish-speaking countries and territories of the Caribbean and 360 Eastern Tropical Pacific. Most of these projects are being carried out by pioneering civil organizations 361 often in strong partnerships with universities, conservation management bodies and regulators, 362 tourism operators, the private sector, associations, and local community groups. While coral reef 363 restoration has been portrayed as too expensive and challenging with regards to spatial scale, 364 duration, and success, the projects presented here have shown that many of these barriers have 365 already been overcome. These pioneering endeavours were often possible by in-kind commitments 366 of staff and volunteers as well as involvement of the local community and tourism operators, thus 367 socio-economic aspects play a substantial role in coral reef restoration in the Caribbean and Eastern 368 Tropical Pacific. Strong national plans for restoration in conjunction with national and international 369 funding are needed to multiply the already existing activities made by Latin-American organisations 370 to improve the health and status of coral reefs in the Caribbean and Eastern Tropical Pacific. From this 371 compilation of data and knowledge, it is apparent that it would be beneficial for coral reef restoration 372 practitioners in this area to coordinate their efforts with each other and make sure they are sharing 373 and implementing their best practices protocols to standardise efforts and track restoration progress 374 by specific, measurable, achievable and repeatable metrics of success through time.

375 Acknowledgements

- 376 We would like to thank Nufar Charuvi, the pioneer driving the Alianza Coralina Taganga project, who,
- although no longer with us, continues to inspire by his example, dedication, and passion he served
- 378 over the last decade for Colombian coral reefs and the local community. This manuscript has been
- 379 developed upon in-kind time of the authors and has not received any financial support. We
- 380 acknowledge the Iberostar Group for covering the open access publication fees of this manuscript.

382 References

- 383 1. SER. The SER primer on ecological restoration. Tucson, Arizona, USA: Group SPW; 2004.
- 2. Perrow M, Davy A. Handbook of ecological restoration. Cambridge, UK: Cambridge University

385 Press; 2002.

- 386 3. McDonald T, Gann G, Jonson J, Dixon K. International standards for the practice of ecological
- 387 restoration including principles and key concepts.2016.
- 388 4. Edwards A. Reef rehabilitation manual. St Lucia, Australia: Coral Reef Targeted Research &
- 389 Capacity Building for Management Program; 2010.
- 390 5. Baums IB, Baker AC, Davies SW, Grottoli AG, Kenkel CD, Kitchen SA, et al. Considerations for
- 391 maximizing the adaptive potential of restored coral populations in the western Atlantic. Ecol Appl.
- 392 2019;0(0):e01978. doi: 10.1002/eap.1978.
- Jeremy BCJ. Pleistocene Perspectives on Coral Reef Community Structure. American Zoologist.
 1992;32(6):719-31.
- 395 7. IUCN. The IUCN Red List of Threatened Species 2019 [Downloaded on 10 January 2019].
 396 Available from: http://www.iucnredlist.org.
- Aronson RB, Precht WF. White-band disease and the changing face of Caribbean coral reefs.
 Hydrobiologia. 2001;460:25-38. doi: 10.1023/a:1013103928980. PubMed PMID:
 WOS:000173249100003.
- 400 9. Rodriguez-Martinez RE, Banaszak AT, McField MD, Beltran-Torres AU, Alvarez-Filip L.
 401 Assessment of Acropora palmata in the Mesoamerican Reef System. PLoS One. 2014;9(4):7. doi:
 402 10.1371/journal.pone.0096140. PubMed PMID: WOS:000335505000064.
- 403 10. Bayraktarov E, Stewart-Sinclair PJ, Brisbane S, Bostrom-Einarsson L, Saunders MI, Lovelock CE,
- 404 et al. Motivations, success, and cost of coral reef restoration. Restor Ecol. 2019;27(5):981-91. doi:
- 405 10.1111/rec.12977. PubMed PMID: WOS:000484997000012.

406 11. Barton JA, Willis BL, Hutson KS. Coral propagation: a review of techniques for ornamental
407 trade and reef restoration. Rev Aquac. 2017;9(3):238-56. doi: 10.1111/raq.12135. PubMed PMID:
408 WOS:000411513000003.

Boström-Einarsson L, Ceccarelli D, Babcock RC, Bayraktarov E, Cook N, Harrison P, et al. Coral
restoration in a changing world - A global synthesis of methods and techniques - A report for the Reef
Restoration and Adaptation Program, Subproject 1a - Review of existing technologies/pilots and new
initiatives. Townsville: James Cook University, 2018.

Hernández-Delgado E, Mercado-Molina A, Suleimán-Ramos S. Multi-disciplinary lessons
learned from low-tech coral farming and reef rehabilitation practices. I. Best management practices.
In: Duque-Beltrán C, Tello-Camacho E, editors. Corals in a Changing World: InTech Publ; 2018. p. 213-

416 43.

417 14. Rinkevich B. The Active Reef Restoration Toolbox is a Vehicle for Coral Resilience and
418 Adaptation in a Changing World. Journal of Marine Science and Engineering. 2019;7(7):18. doi:
419 10.3390/jmse7070201. PubMed PMID: WOS:000478581900009.

420 15. Edwards AJ, Clark S. Coral transplantation: A useful management tool or misguided meddling?
421 Mar Pollut Bull. 1998;37(8-12):474-87. PubMed PMID: WOS:000082454300013.

422 16. Precht W. Coral Reef Restoration: An Overview. Coral Reef Restoration Handbook: CRC Press;
423 2006. p. 55-76.

424 17. Rinkevich B. Conservation of coral reefs through active restoration measures: Recent
425 approaches and last decade progress. Environ Sci Technol. 2005;39(12):4333-42. doi:
426 10.1021/es0482583. PubMed PMID: WOS:000229724100007.

18. Rinkevich B. Restoration strategies for coral reefs damaged by recreational activities: The use
of sexual and asexual recruits. Restor Ecol. 1995;3(4):241-51. doi: 10.1111/j.1526100X.1995.tb00091.x. PubMed PMID: WOS:A1995TL38900003.

430 19. Rinkevich B. Steps towards the evaluation of coral reef restoration by using small branch
431 fragments. Mar Biol. 2000;136(5):807-12. doi: 10.1007/s002270000293. PubMed PMID:
432 WOS:000088186100006.

433 20. Johnson M, Lustic C, Bartels E, Baums I, Gilliam D, Larson L, et al. Caribbean Acropora
434 restoration guide: best practices for propagation and population enhancement. Arlington, VA: The
435 Nature Conservancy; 2011.

Frias-Torres S, Goehlich H, Reveret C, Montoya-Maya PH. Reef fishes recruited at midwater
coral nurseries consume biofouling and reduce cleaning time in Seychelles, Indian Ocean. Afr J Mar
Sci. 2015;37(3):421-6. doi: 10.2989/1814232x.2015.1078259. PubMed PMID:
WOS:000363342200014.

440 22. Bowden-Kerby A. Low-tech coral reef restoration methods modeled after natural
441 fragmentation processes. Bull Mar Sci. 2001;69(2):915-31. PubMed PMID: WOS:000172609000051.

Shaish L, Levy G, Gomez E, Rinkevich B. Fixed and suspended coral nurseries in the Philippines:
Establishing the first step in the "gardening concept" of reef restoration. J Exp Mar Biol Ecol.

444 2008;358(1):86-97. doi: 10.1016/j.jembe.2008.01.024. PubMed PMID: WOS:000255448300010.

445 24. Ferse SCA, Kunzmann A. Effects of concrete-bamboo cages on coral fragments: Evaluation of
446 a low-tech method used in artisanal ocean-based coral farming. Journal of Applied Aquaculture.
447 2009;21(1):31-49. doi: 10.1080/10454430802694538.

Mbije NE, Spanier E, Rinkevich B. A first endeavour in restoring denuded, post-bleached reefs
in Tanzania. Estuar Coast Shelf Sci. 2013;128:41-51. doi: DOI 10.1016/j.ecss.2013.04.021. PubMed
PMID: WOS:000321414100005.

451 26. Nedimyer K, Gaines K, Roach S. Coral tree nursery©: an innovative approach to growing corals
452 in an ocean-based field nursery. AACL Bioflux. 2011;4(4):442-6.

453 27. dela Cruz DW, Rinkevich B, Gomez ED, Yap HT. Assessing an abridged nursery phase for slow 454 growing corals used in coral restoration. Ecol Eng. 2015;84:408-15. doi: 455 10.1016/j.ecoleng.2015.09.042. PubMed PMID: WOS:000363948000051.

456 28. Ng CSL, Lim SC, Ong JY, Teo LMS, Chou LM, Chua KE, et al. Enhancing the biodiversity of coastal
457 defence structures: transplantation of nursery-reared reef biota onto intertidal seawalls. Ecol Eng.
458 2015;82:480-6. doi: http://dx.doi.org/10.1016/j.ecoleng.2015.05.016.

459 29. Forsman ZH, Page CA, Toonen RJ, Vaughan D. Growing coral larger and faster: micro-colony460 fusion as a strategy for accelerating coral cover. Peerj. 2015;3:16. doi: 10.7717/peerj.1313. PubMed
461 PMID: WOS:000364330400004.

462 30. Page CA, Muller EM, Vaughan DE. Microfragmenting for the successful restoration of slow
463 growing massive corals. Ecol Eng. 2018;123:86-94. doi: 10.1016/j.ecoleng.2018.08.017. PubMed
464 PMID: WOS:000446253500009.

465 31. Chamberland VF, Petersen D, Guest JR, Petersen U, Brittsan M, Vermeij MJA. New Seeding
466 Approach Reduces Costs and Time to Outplant Sexually Propagated Corals for Reef Restoration. Sci
467 Rep. 2017;7:12. doi: 10.1038/s41598-017-17555-z. PubMed PMID: WOS:000418644100035.

Banaszak A, Schutter M, Guendulain García S, Mendoza Quiroz S, Gómez Campo K. Guía
práctica para la restauración con base en la producción de reclutas sexuales de corales con énfasis en
Acropora palmata. México: 2018.

33. Calle-Trivino J, Cortes-Useche C, Sellares-Blasco RI, Arias-Gonzalez JE. Assisted fertilization of
threatened Staghorn Coral to complement the restoration of nurseries in Southeastern Dominican
Republic. Reg Stud Mar Sci. 2018;18:129-34. doi: 10.1016/j.rsma.2018.02.002. PubMed PMID:
WOS:000427936600014.

475 34. dela Cruz DW, Harrison PL. Enhanced larval supply and recruitment can replenish reef corals
476 on degraded reefs. Sci Rep. 2017;7:13. doi: 10.1038/s41598-017-14546-y. PubMed PMID:
477 WOS:000413700200024.

478 35. Van Oppen MJH, Gates RD, Blackall LL, Cantin N, Chakravarti LJ, Chan WY, et al. Shifting
479 paradigms in restoration of the world's coral reefs. Glob Change Biol. 2017;23(9):3437-48. doi:
480 10.1111/gcb.13647. PubMed PMID: WOS:000406812100003.

481 36. CRC. Coral Restoration Consortium 2019 [cited 2019 10 January 2019]. Available from:

482 <u>http://reefresilience.org/restoration/restoration-introduction/coral-restoration-consortium/.</u>

483 37. Clewell AF, Aronson J. Motivations for the restoration of ecosystems. Conserv Biol.

484 2006;20(2):420-8. doi: 10.1111/j.1523-1739.2006.00340.x. PubMed PMID: WOS:000236064200020.

485 38. Hagger V, Dwyer J, Wilson K. What motivates ecological restoration? Restor Ecol.
486 2017;25(5):832-43. doi: 10.1111/rec.12503. PubMed PMID: WOS:000411686900019.

487 39. Hein MY, Willis BL, Beeden R, Birtles A. The need for broader ecological and socioeconomic

488 tools to evaluate the effectiveness of coral restoration programs. Restor Ecol. 2017;25(6):873-83. doi:

489 10.1111/rec.12580. PubMed PMID: WOS:000414808200004.

490 40. Bayraktarov E, Saunders MI, Abdullah S, Mills M, Beher J, Possingham HP, et al. The cost and
491 feasibility of marine coastal restoration. Ecol Appl. 2016;26(4):1055-74. doi: 10.1890/15-1077.
492 PubMed PMID: WOS:000378092900008.

493 41. Iacona GD, Sutherland WJ, Mappin B, Adams VM, Armsworth PR, Coleshaw T, et al.
494 Standardized reporting of the costs of management interventions for biodiversity conservation.
495 Conserv Biol. 2018;32(5):979-88. doi: doi:10.1111/cobi.13195.

496 42. Ruiz-Jaen MC, Mitchell Aide T. Restoration Success: How Is It Being Measured? Restor Ecol.
497 2005;13(3):569-77. doi: 10.1111/j.1526-100X.2005.00072.x.

498 43. Sunderland T, Sunderland-Groves J, Shanley P, Campbell B. Bridging the Gap: How Can
499 Information Access and Exchange Between Conservation Biologists and Field Practitioners be
500 Improved for Better Conservation Outcomes? Biotropica. 2009;41(5):549-54. doi: 10.1111/j.1744501 7429.2009.00557.x. PubMed PMID: WOS:000269700500005.

502 Milner-Gulland EJ, Fisher M, Browne S, Redford KH, Spencer M, Sutherland WJ. Do we need 44. 503 to develop а more relevant conservation literature? Oryx. 2010;44(1):1-2. doi: 504 10.1017/s0030605309991001. PubMed PMID: WOS:000274770200001.

45. IPCC. Global Warming of 1.5 °C. An IPCC special report on the impacts of global warming of
506 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context

507 of strengthening the global response to the threat of climate change, sustainable development, and 508 efforts to eradicate poverty. Geneva, Switzerland: 2018.

509 46. Cortés J. Eastern Tropical Pacific Coral Reefs. In: Hopley D, editor. Encyclopedia of Modern

510 Coral Reefs: Structure, Form and Process. Dordrecht: Springer Netherlands; 2011. p. 351-8.

511 47. Hubbard D. Eastern Caribbean Coral Reefs. In: Hopley D, editor. Encyclopedia of Modern Coral

512 Reefs: Structure, Form and Process. Dordrecht: Springer Netherlands; 2011. p. 338-48.

513 48. Rogers CS. RESPONSES OF CORAL REEFS AND REEF ORGANISMS TO SEDIMENTATION. Mar

514 Ecol-Prog Ser. 1990;62(1-2):185-202. doi: 10.3354/meps062185. PubMed PMID:
515 WOS:A1990CZ63900020.

516 49. Cloern JE. Our evolving conceptual model of the coastal eutrophication problem. Mar Ecol-

517 Prog Ser. 2001;210:223-53. doi: 10.3354/meps210223. PubMed PMID: WOS:000167175000019.

50. Hernandez-Delgado EA. The emerging threats of climate change on tropical coastal ecosystem 519 services, public health, local economies and livelihood sustainability of small islands: Cumulative 520 impacts and synergies. Mar Pollut Bull. 2015;101(1):5-28. doi: 10.1016/j.marpolbul.2015.09.018.

521 PubMed PMID: WOS:000367630700015.

51. Jackson J, Donovan M, Cramer K, Lam VV. Status and Trends of Caribbean Coral Reefs: 19702012. Gland, Switzerland: 2014.

524 52. Meyer JL, Castellanos-Gell J, Aeby GS, Häse CC, Ushijima B, Paul VJ. Microbial Community 525 Shifts Associated With the Ongoing Stony Coral Tissue Loss Disease Outbreak on the Florida Reef Tract.

526 Frontiers in Microbiology. 2019;10(2244). doi: 10.3389/fmicb.2019.02244.

527 53. AGRRA. Coral disease outbreak: Atlantic and Gulf Rapid Reef Assessment; 2019 [cited 2019
528 13/10/2019]. Available from: https://www.agrra.org/coral-disease-outbreak/.

529 54. RRAP. Reef Restoration and Adaptation Program 2019 [cited 2019 13/10/2019]. Available 530 from: https://www.gbrrestoration.org/the-program.

531 55. Woodhead AJ, Hicks CC, Norström AV, Williams GJ, Graham NAJ. Coral reef ecosystem services

532 in the Anthropocene. Functional Ecology. 2019;33(6):1023-34. doi: 10.1111/1365-2435.13331.

533 56. Ferrario F, Beck MW, Storlazzi CD, Micheli F, Shepard CC, Airoldi L. The effectiveness of coral 534 reefs for coastal hazard risk reduction and adaptation. Nat Commun. 2014;5(1):3794. doi: 535 10.1038/ncomms4794.

536 57. Beck MW, Losada IJ, Menéndez P, Reguero BG, Díaz-Simal P, Fernández F. The global flood 537 protection savings provided by coral reefs. Nat Commun. 2018;9(1):2186. doi: 10.1038/s41467-018-

538 04568-z.

539 58. Schwartz MW, Cook CN, Pressey RL, Pullin AS, Runge MC, Salafsky N, et al. Decision Support
540 Frameworks and Tools for Conservation. Conserv Lett. 2018;11(2):e12385. doi: 10.1111/conl.12385.

541 59. Riegg Cellini S, Kee E. Cost-Effectiveness and Cost-Benefit Analysis. In: Newcomer K, Hatry H,

542 Wholey J, editors. Handbook of Practical Program Evaluation2015. p. 636-72.

60. Beher J, Possingham HP, Hoobin S, Dougall C, Klein C. Prioritising catchment management
projects to improve marine water quality. Environmental Science & Policy. 2016;59:35-43. doi:
10.1016/j.envsci.2016.02.005. PubMed PMID: WOS:000373548500005.

61. Gillies CL, Fitzsimons JA, Branigan S, Hale L, Hancock B, Creighton C, et al. Scaling-up marine
restoration efforts in Australia. Ecological Management & Restoration. 2015;16(2):84-5. doi:
10.1111/emr.12159. PubMed PMID: WOS:000354644800002.

62. Amano T, González-Varo JP, Sutherland WJ. Languages Are Still a Major Barrier to Global
Science. PLoS Biol. 2016;14(12):e2000933. doi: 10.1371/journal.pbio.2000933.