**Temporal and spatial dynamics of peat microbiomes in drained and rewetted**

**soils of three temperate peatlands**

**Haitao Wang[1]\*, Micha Weil[1], Dominik Zak[2, 3], Diana Mu☐nch[1], Anke Günther[4], Gerald Jurasinski[4],**

**Tim Urich[1]\***

[1] Institute of Microbiology, University of Greifswald, Germany

[2] Department of Bioscience, Aarhus University, Denmark

[3] Department of Biogeochemistry and Chemical Analytics, Leibniz-Institute of Freshwater Ecology and

Inland Fisheries Berlin, Germany

[4] Chair of Landscape Ecology and Site Evaluation, University of Rostock, Germany

**Contacting details:**

H. Wang: haitao.wang@uni-greifswald.de

M. Weil: micha.weil@uni-greifswald.de

D. Zak: doz@bios.au.dk

D. Mu☐nch: diana.muench1@gmx.de

A. Günther: anke.guenther@uni-rostock.de

G Jurasinski: gerald.jurasinski@uni-rostock.de

Tim Urich: tim.urich@uni-greifswald.de

\* Corresponding to: haitao.wang@uni-greifswald.de or tim.urich@uni-greifswald.de

**Abstract**

**Background:** Drainage of high-organic peatlands for agricultural purposes has led to increased

greenhouse gas emissions and loss of biodiversity. In the last decades, rewetting of peatlands is on the rise

worldwide, to mitigate these negative impacts. However, it remains still questionable how rewetting

would influence peat microbiota as important drivers of nutrient cycles and ecosystem restoration. Here,

we investigate the spatial and temporal dynamics of the diversity, community composition and network

interactions of prokaryotes and eukaryotes, and the influence of rewetting on these microbial features in

formerly long-term drained and agriculturally used fens. Peat-soils were sampled seasonally from three

drained and three rewetted sites representing the dominating fen peatland types of glacial landscapes in

Northern Germany, namely alder forest, costal fen and percolation fen.

**Results:** Costal fens as salt-water impacted systems showed a lower microbial diversity and their

microbial community composition showed the strongest distinction from the other two peatland types.

Prokaryotic and eukaryotic community compositions showed a congruent pattern which was mostly

driven by peatland type and rewetting. Rewetting decreased the abundances of fungi and prokaryotic

decomposers, while the abundance of potential methanogens was significantly higher in the rewetted

sites. Rewetting also influenced the abundance of ecological clusters in the microbial communities

identified from the co-occurrence network. The microbial communities changed only slightly with depth

and over time. According to structural equation models rewetted conditions affected the microbial

communities through different mechanisms across the three studied peatland types.

**Conclusions:** Our results suggest that rewetting strongly impacts the structure of microbial communities

and, thus, important biogeochemical processes, which may explain the high variation in greenhouse gas

emissions upon rewetting of peatlands. The improved understanding of functional mechanisms of

rewetting in different peatland types lays the foundation for securing best practices to fulfil multiple

restoration goals including those targeting on climate, water, and species protection.

**Keywords:** Prokaryotes, Eukaryotes, Rewetting, Peatland soils, Alder forest, Coastal fen, Percolation fen

**Background**

Peatlands store over 30% of the earth's soil carbon although they only cover nearly 3% of the global land

area [1]. Waterlogging contributes to the high level of soil organic carbon since peat, consisting of

incompletely decomposed organic matters derived mostly from plant residues, accumulates when the

oxygen is deficient [2]. However, drainage of wetlands including peatlands for agricultural use

contributed to half of the loss of the global wetlands [1, 3]. Drainage of peatlands results in decreased

emissions of methane ($CH_4$) but significantly increased emissions of carbon dioxide ($CO_2$) and potentially

increased emissions of nitrous oxide ($N_2O$), ultimately contributing to increased warming potential effect

[2, 4-6]. The conversion of natural fens to agricultural lands also leads to biodiversity loss due to

intensive dehydration, tillage and fertilization [1]. Rewetting of drained peatlands aims at mitigating these

negative impacts by restoring peatlands. By increasing the water level, the existing peat carbon pool is

conserved as a result of the quickly reestablished anoxia. Under anoxic conditions phenol oxidase

becomes inactivated promoting the accumulation of recalcitrant phenolic compounds released by plants

which in turn limits the activity of hydrolase enzymes and eventually leads to low decomposition rates

[7-9]. This benefit, however, might be offset since deficiency of oxygen can also promote carbon loss

through elevated production and emission of $CH_4$ in particular if sites become inundated and colonized by

dense stands of hydrophytes and/or helophytes [10]. Which effect prevails in a given rewetted peatland is

strongly driven by the prevailing microbial decomposition processes. Therefore, a better understanding of

the microbial communities and their processing of organic matter is paramount to an informed and

optimized management of rewetted peatlands.

Fungi are considered to be the major agents of plant litter decomposition, while bacteria as well as some

invertebrates can, to lesser extent, also be significant contributors [11]. The decomposition of litter and

deadwood eventually leads to the release of $CO_2$ to the atmosphere. This process is called soil respiration,

and is a crucial element of all carbon cycling on Earth [12]. The fate of organic carbon and nitrogen is

also influenced by microbial communities associated with methane cycling and nitrogen cycling in which

$CH_4$ and $N_2O$ are produced [13, 14]. While decent efforts were invested to investigate the influence of

rewetting on the emissions of greenhouse gases (GHGs) [15-18], extracellular enzyme activities [19],

dissolve organic carbon [20, 21], and water chemistry [15, 19], little is known about the changes of

microbial communities driving these processes.

Since microbes are both the most diverse organism group on the planet and the major agents of

biogeochemical cycling [22], their diversity and community structures are of great importance to

ecosystem functioning. It is especially yet unclear how rewetting impacts the peat communities regarding

all domains of life (archaea, bacteria and eukaryotes). Furthermore, unraveling the dynamics of

community composition together with relevant environmental factors helps to carve out the predictable

spatial and temporal patterns [23], thus better perceiving the ecological consequences resulting from these

patterns. It has been shown that decomposition rates and GHG emissions follow seasonal patterns

[24-27], but the temporal dynamics of prokaryotic and eukaryotic communities in peat-soils are still

poorly understood. A better understanding of microbial community dynamics in rewetted peatlands could

lay the foundation for maintaining peat-soil quality and health, thus guiding future management of

peatland restoration.

About two decades ago a peatland restoration program was initiated in the state of

Mecklenburg-Vorpommern (M-V) in Northern Germany, and in this course over 20,000 ha of peat-soils

in this state were rewetted [28]. The relatively high number of rewetted sites in different peatland types

provides an excellent opportunity to study the influence of rewetting on peat microbial community

dynamics in the three most relevant peatland types of the region, namely alder forest, coastal fen and

percolation fen. The temporal dynamics of prokaryotic (archaeal and bacterial) and eukaryotic

communities were assessed through high-throughput sequencing with 16S and 18S rRNA gene

amplicons, respectively. Additional edaphic variables, including water content, pH, organic carbon and

nutrients, were monitored. The diversities and community compositions of prokaryotes and eukaryotes

were compared between the rewetted and drained sites within each peatland type. Our objectives are 1) to

investigate the temporal changes of microbial diversities and community compositions in different

seasons, 2) to reveal the influence of rewetting on the diversities, community compositions, network

interactions and the potential functions of relevant biogeochemical processes, and 3) to reveal the

functional mechanisms of rewetting on shaping the microbial communities and functions in the three

different peatland types. We hypothesize that rewetting drained fens will influence diversity and

community composition of both prokaryotes and eukaryotes, and changes in these communities will

follow a seasonally temporal pattern. Moreover, the functional mechanisms of rewetting will be different

in these three peatland types *inter alia* due to differences in salinity, vegetation and soil organic matter

composition. This study finally intends to provide a theoretical guide for future planning and land

management of drained peatlands.

**Results**

**Study sites and soil edaphic properties**

The six study sites are located in M-V in Northeastern Germany (Additional file 1: Fig. S1) and cover the

three major peatland types (alder forest, coastal fen and percolation fen) in M-V. For each peatland type,

two sites were selected. Both of them were drained in the past but one of the two has been rewetted. The

alder forest presumably formed under water-logged conditions. Both sites were drained in the past and

used as managed forest for many decades. The drained alder forest site (AD) is still drained, and the

rewetted site (AW) was getting rewetted when the nearby Bauernmoor was rewetted in 2003. The coastal

fen sites are located near Greifswald at the Karrendorf peninsula. The whole area was drained already in

1850 and has been used for pasture since. From the 1960s drainage was enforced and the areas were used

as intensive grasslands. The drained site (CD) is still behind a dike but is now in an area that is used for

extensive cattle grazing, the rewetted site (CW) is located in a part of the area that has been rewetted in

1993 by removing the old dikes. These parts are now flooded regularly, mostly during autumn and winter.

The two percolation fens are located in one percolation fen complex at the rivers Recknitz and Trebel,

respectively. Both sites also have a long drainage history with early drainage in the 18th century and

much more intense drainage in the 1960s. Both sites were used as intensive grassland. While the drained

site (PD) is still under medium intensive grassland use, the wet site (PW) has been rewetted in 1997 and

was not actively managed since then.

The soil properties varied significantly among the six sites (Additional file 2: Table S1). Rewetting

generally increased the soil moisture in all three peatland types (Additional file 1: Fig. S2 and Additional

file 1: Fig. S3), with a significant increase in alder forest (Additional file 2: Table S1). Since September

2017, the water level was always higher in the rewetted sites compared with their drained counterparts

with the exception that CW and CD showed similar water level fluctuations since May 2018 (Additional

file 1: Fig. S4). The soil temperature followed a clear seasonal pattern from August 2017 to July 2018

(Additional file 1: Fig. S5). Moreover, dissolved organic matter (DOM) was higher in the wet sites,

especially in the alder forest and coastal fen and the concentrations of the three considered DOM

compounds were in most of the rewetted sites (Additional file 2: Table S1). The concentrations of DOM

and single compounds also increased sharply from April to November in 2017 and then decreased to

lower levels in February 2018 (Additional file 1: Fig. S2). All rewetted sites showed lower nitrate

concentrations compared to their still-drained counterparts, while this was only the case in AW and CW

for nitrite and ammonium (Additional file 2: Table S1). Nitrate showed a similar temporal pattern as

DOM in AD but a contrasting trend in the other sites, while ammonium concentrations were increasing

since April in all sites except for CD (Additional file 1: Fig. S2). Water-extractable P content was higher

in AW than in AD (Additional file 2: Table S1), and it significantly decreased since April in AW, CD and

PD (Additional file 1: Fig. S2). pH and salinity were significantly higher in the rewetted sites compared

with their drained counterparts (Additional file 2: Table S1). pH and salinity were not compared among

seasons since they were measured with different methods in April and the other seasons. However, there

were less variations of these soil properties with depths, with only a few changes observed in certain sites

(Additional file 1: Fig. S3).

**Microbial diversities**

The diversities of both prokaryotes and eukaryotes were lower in the coastal sites compared with the other

sites (Fig. 1a, Additional file 1: Fig. S6, and Additional file 2: Table S1). Prokaryotic diversities changed

significantly across seasons in AD, CD, PD and PW, while eukaryotic diversities, including fungi, protists

and metazoa, showed less significant changes across seasons (Fig. 1a). Under wet conditions prokaryotic

diversity was higher in coastal fen and protist diversity was higher in alder forest and coastal fen

(Additional file 2: Table S1). Metazoan diversity, however, was lower in all wet sites while diversities of

fungi and protists were lower in the wet percolation fen (Additional file 2: Table S1). Prokaryotic

diversity declined with depth in AW and CW (Additional file 1: Fig. S6) as well as the diversities of fungi

and metazoa (Additional file 1: Fig. S6). Both prokaryotic and eukaryotic diversities showed positive

correlation with nitrate and negative correlation with salinity and prokaryotic diversity was also positively

correlated with other soil properties (Additional file 1: Fig. S7).

**Microbial community compositions**

The prokaryotic communities were dominated by Acidobacteria (19.4%), Betaproteobacteria (11.7%),

Alphaproteobacteria (11.1%), Actinobacteria (7.69%) and Chloroflexi (6.41%) (Additional file 1: Fig.

S8). The taxa distributions showed similar patterns during these four seasons, but obvious differences

among sites were observed. For instance, the Acidobacteria were more abundant in the drained sites

compared with the rewetted sites, while Betaproteobacteria showed a contrasting trend (Additional file 1:

Fig. S8). Actinobacteria were more abundant in the coastal fens while Cloroflexi were more abundant in

the other two peatland types (Additional file 1: Fig. S8). These variations were reflected in the nonmetric

multidimensional scaling (NMDS) plot where the study sites grouped very well (Fig. 1b). However, the

samples from percolation fen and alder forest showed considerable overlap resulting from the numerous

shared amplicon sequence variants (ASVs) between these two peatland types (Fig. 1b). In addition, these

two peatland types shared much less ASVs with the coastal fen than they shared with each other (Fig. 1b).

The changes in prokaryotic community composition were mostly driven by peatland types

(PERMANOVA, $R^2$=0.255, $P$=0.001). The hydrological state (drained/rewetted) also showed a

significant but less strong influence (PERMANOVA, $R^2$=0.080, $P$=0.001) on the prokaryotic community.

However, season showed no significant impact on prokaryotic community composition (PERMANOVA,

$R^2$=0.017, $P$=0.168, Additional file 1: Fig. S9a), while depth showed a significant but weak impact

(PERMANOVA, $R^2$=0.026, $P$=0.001, Additional file 1: Fig. S9b). NMDS was also conducted separately

for each site to exclude the site effect, and the results showed that the interaction of depth and season

exhibited strong impact on community composition in all sites except AD (PERMANOVA, $P$<0.05,

Additional file 1: Fig. S10).

The eukaryotic communities were dominated by fungi (Ascomycota, 25.6%; Basidiomycota, 12.5%;

Glomeromycota, 3.09%), Nematoda (10.5%), Cercozoa (9.53%), Apicomplexa (5.01%) and Arthropoda

(4.47%) (Additional file 1: Fig. S11). Specifically, Nematoda and Mortirellales were more abundant in

the drained sites compared with the rewetted sites and Apicomplexa were more abundant in the

percolation fens compared with the other two peatland types (Additional file 1: Fig. S11). The eukaryotic

community composition exhibited a congruent pattern as prokaryotes but with less overlapping between

percolation fen and alder forest (Fig. 1c). There were significant differences between peatland types

(PERMANOVA, $R^2$=0.189, $P$=0.001), and between drained and rewetted sites (PERMANOVA,

$R^2$=0.071, $P$=0.001), while there was no significant impact of season (PERMANOVA, $R^2$=0.018,

$P$=0.069, Additional file 1: Fig. S9c) and only a weak impact of depth (PERMANOVA, $R^2$=0.018,

$P$=0.002, Additional file 1: Fig. S9d). Similar to prokaryotes, we also found that the interaction between

depth and season exhibited a strong impact on community composition in all sites except AD

(PERMANOVA, $P$<0.05, Additional file 1: Fig. S12).

The prokaryotic and eukaryotic community compositions were driven by soil properties in a similar way.

Salinity, which was higher in the coastal fen sites, differentiated the coastal from the other fens, while

nitrate was more abundant and related with alder forest (Fig. 1b and c). Moisture, pH, DOM and its

components, P and ammonium content, were generally highest in PW, and they also differentiated the
rewetted from the drained sites (Fig. 1b and c). The community composition of eukaryotes was
significantly correlated with that of prokaryotes (Mantel test, r=0.852, *P*=0.001). By analyzing plant
ASVs separately, we also found distinct clusters of the samples grouped by sites (Additional file 1: Fig.
S9e). The Mantel tests showed that the plant community composition was strongly and significantly
correlated with both prokaryotic (r=0.551, *P*=0.001) and eukaryotic (r=0.634, *P*=0.001) community
compositions.

**Differentially abundant taxa and predicted functions**

The prokaryotic ASVs were collapsed into functional groups based on their taxonomies using
FAPROTAX. 8 dominant and representative functions regarding carbon and nitrogen cycles, and
decomposition were analyzed (Fig. 2). Rewetting significantly promoted functions including
methanogenesis, denitrification, sulfate respiration and fermentation in certain peatland types, while the
others including cellulolysis, nitrification and nitrogen fixation were over-represented in the drained sites
(Fig. 2). Methanotrophy was over-represented in the rewetted sites of alder forest and percolation fen but
in the drained site of coastal fen (Fig. 2). While the predicted functions showed few temporal patterns
(Additional file 1: Fig. S13), there were more differences between depth sections (Additional file 1: Fig.
S14). Some differentially abundant ASVs were dominant and could be assigned to certain functions
according to FAPROTAX. Within Actinobacteria, several dominant ASVs of *Acidothermus* genus
involved in cellulolysis were more abundant in AD and PD than in their rewetted counterparts (Fig. 3)
supporting the finding that microbes conducting cellulolysis were more abundant in the drained sites (Fig.
2). Two *Flavobacterium* ASVs with high fold changes were more abundant in CD than in CW, and they
are aerobic chemoheterotrophs (Fig. 3). One ASV of *Desulfobulbus* driving sulfate reduction was more
abundant in CW than in CD (Fig. 3), which is in line with the result that sulfate respiration was stronger
in the rewetted sites (Fig. 2). Some ASVs of *Sulfurimonas* were also more abundant in the rewetted site of
coastal fen (Fig. 3). The ASVs belonging to phylum Nitrospirae were mostly more abundant in the
rewetted sites, while several ASVs of *Ca.* Nitrosotalea were more important in the drained sites (Fig. 3).

While most of the fungal ASVs could not be matched to the database, 15 functional guilds were identified
with FUNGuild, and only 8 of them showed significant differences between the drained and the rewetted
sites (Additional file 1: Fig. S15). Rewetting promoted the fungi that might be plant pathogens, algal or

fungal parasites, and soil or plant saprotroph in the alder forest and the percolation fen (Additional file 1:
Fig. S15). Soil saprotroph and arbuscular mycorrhizal fungi (AMF) were abundant and were
over-represented in CW, while AMF or fungi that might be some certain saprotrophs were more
important in AD and PD (Additional file 1: Fig. S15). The analysis of differentially abundant taxa
supported these results. The ASVs of *Archaeorhizomyces* were identified as soil saprotrophs, and one of
these abundant ASVs was more important in PW than in PD (Fig. 4). The ASVs belonging to Tremellales
(Fungal Parasite-Undefined Saprotroph) were more abundant in the drained site (Fig. 4). The
Glomeromycota phylum was identified as AMF. One Archaeosporales ASV was dominant in AD, and the
two differentially abundant AMF ASVs were found in PD, while there were more abundant AMF ASVs
in CW than in CD (Fig. 4). For protists, the Apicomplexa including ASVs of Eugregarinorida, Gregarina
and *Monocystis agills* was more important in the drained sites compared to the rewetted sites (Fig. 4). For
metazoa, the abundant ASVs of Acari belonging to Arthropoda were found in AD and in both CW and
CD, while several nematoda ASVs belonging to Rhabditida, Tylenchida and Enplea were mostly found to
be more abundant in the drained sites (Fig. 4).

Since microbes carrying out cellulolysis showed high abundance (Additional file 1: Fig. S13 and S14), we
are considering them as the major prokaryotic decomposers. Both the prokaryotic (cellulolysis) and
eukaryotic decomposers (fungi) were significantly and negatively correlated with the water content
(Additional file 1: Fig. S16).

**Co-occurrence network**

In total, 6,290 significant correlations were captured among 1,769 prokaryotic and 708 eukaryotic ASVs,
and all of them were positive (Fig. 5a). These correlations could indicate the potential interactions among
different microbial taxa. We identified 8 major ecological clusters (modules) from this network which
might potentially drive some specific functions in the ecosystem (Fig. 5a). These clusters were differently
distributed across sites, and the abundances of prokaryotes and eukaryotes within each module showed
consistent patterns (Fig. 5b). Module I and III were dominant in some rewetted sites, while module IV, V
and VI showed higher abundances in the drained sites. Module II was more abundant in the alder forest
while module VII was more abundant in the coastal fen. Elements of Module VIII were mainly
distributed in AW and PD. However, there was no distinct change of these modules with depths or
seasons (Fig. 5b). Since all prokaryotes and eukaryotes with a certain module responded similarly, we

averaged their abundances for each module. The correlation analysis exhibited that all these modules

were significantly correlated with soil moisture, nitrate concentration and salinity (Additional file 1: Fig.

S7). Some of them were also correlated with DOM and pH (Additional file 1: Fig. S7).

**Structural equation modelling (SEM)**

We designed a *priori* theoretical model based on the causal relationships among the variables (Additional

file 1: Fig. S17). By conducting the group analysis and comparing the constrained and unconstrained

models, we observed significant differences of the models for the three peatland types. In the alder forest,

rewetting showed no direct effects on prokaryotic and eukaryotic diversities but had direct effects on the

other response variables. Rewetting could also indirectly influence these microbial parameters through

changing soil moisture and then DOM which showed significant impacts on these responses (Fig. 6).

DOM also covaried with Nitrate, pH and plant community composition, indicating that DOM might also

be controlled by these properties (Fig. 6). Moreover, the different plant community compositions between

drained and rewetted sites also influenced the eukaryotic community composition, ecological clusters and

predicted functions (Fig. 6). In the coastal fen, rewetting directly influenced all the responses. The

indirect effects of rewetting were mainly mediated by nitrate, DOM and pH, but not by soil moisture (Fig.

6). As salinity strongly covaried with nitrate and differed significantly between CD and CW (Additional

file 1: Fig. S7and Additional file 2: Table S1), salinity in addition to pH might be the key driver of

rewetting effect in coastal fen. The plant community compositions only mediated the influence of

rewetting on eukaryotic functions and ecological clusters (Fig. 6). In the percolation fen, rewetting

showed direct effects on all the response variables excluding prokaryotic diversity. It also initially

influenced the soil moisture which then influenced nitrate, DOM and pH. The pH was an important

mediator which was influenced directly by rewetting and moisture, and influenced all the responses

excluding eukaryotic communities (Fig. 6). However, DOM and plant community composition showed

less impact on the response variables (Fig. 6). The $R^2$ values of eukaryotic diversity were low in all study

sites, and the prokaryotic diversity was also weakly explained in alder forest and percolation fen (Fig. 6).

**Discussion**

Our results show that rewetting drained fens influences the diversity and community composition of both

prokaryotes and eukaryotes, but different patterns were found for the three peatland types. In the

following temporal and spatial aspects are considered in detail besides certain specific factors driving the

change of microbial communities. Finally, the overall impact of rewetting on microbial communities is

discussed in the broader context of ecological restoration and GHG emissions.

**1. Temporal and vertical patterns of the prokaryotic and eukaryotic communities**

In our study, most of the detected soil parameters showed significant temporal changes in addition to soil

temperature (Additional file 1: Fig. S2 and S5), the most important indicator of seasonality. These

parameters probably drove the temporal patterns observed in the prokaryotic and some of the eukaryotic

diversities (Fig. 1a), as well as some of the temporal changes of predicted functions (Additional file 1:

Fig. S13). However, both prokaryotic and eukaryotic community compositions at sites showed no

obvious temporal changes, indicating that peat microbial communities do not react strongly to weather

driving temporal changes of environmental conditions. This resilience to external forces such as

temperature and nutrient supply might result from the internal feedback mechanisms including

competition, viral infection and predator-prey interlinkages, that maintain a relatively stable state of

community in seasonally changing environments [23]. Moreover, the water level we monitored since

September 2017 (Additional file 1: Fig. S4) and the water content (Additional file 1: Fig. S2) showed no

obvious seasonal patterns. The stable water status among seasons might also have contributed to the slow

changes in the microbial communities since soil moisture was one of the most important environmental

factors shaping the community compositions (Fig. 1b and c). Also, if water-saturation or high-water

levels occur, respectively it may create homogeneity and weak niche differentiation that leads to stronger

interactions between microbes [29], resulting in high resilience to environmental stresses. These results

support that peatlands can stand and be resilient to gradual, long-term changes in climate and water

conditions [2].

The decrease of fungal and metazoan diversities with increasing depth (Additional file 1: Fig. S6)

suggests the general pattern that these two eukaryotic groups prefer to live in upper soil layers where

oxygen is sufficiently available. It has been demonstrated that environmental gradients drive changes in

microbial community composition along soil depth [30]. Our study found that depth showed a weak yet

significant effect on the microbial community compositions. While many soil parameters we recorded

changed only slightly with depth, distinct moisture depth profiles were observed in sites with higher water

content (Additional file 1: Fig. S3). This indicated that oxygen related to water content might contribute

to depth profiles of microbial communities, which was also supported by significant changes in

oxygen-sensitive microbial processes with depths, including methanogenesis, nitrification, cellulolysis

and fermentation (Additional file 1: Fig. S14). Despite of the weak impact of depth and season, their

interaction showed strong and significant effect on both prokaryotic and eukaryotic community

compositions in all sites except AD (Additional file 1: Fig. S10 and S12), suggesting that the influence of

these two factors was covered by the influence of site. It also indicated that temporal shifts of

compositions only existed at certain depths in a single site. However, one year of observation is a limited

basis for understanding seasonal patterns as these patterns include an annual repeating pattern, not just

changes over a short period [23]. In addition, we experienced quite unusual weather patterns over the

course of our study period with very wet end of 2017 followed by an exceptionally dry year 2018.

Therefore, a longer-term monitoring of microbial dynamics in peatlands at a smaller scale is needed in

future studies.

**2. Major factors driving changes in prokaryotic and eukaryotic communities**

By investigating the dynamics of prokaryotic and eukaryotic communities in peat soils, we observed a

congruent pattern between these two communities as well as their responses to the environmental factors,

which is supported by a previous study showing a similar pattern [31]. In the coastal sites we found a

significant lower prokaryotic and eukaryotic diversity than in the two freshwater peatland types, which

was mainly due to the varying salinity that was negatively correlated to both diversities (Additional file 1:

Fig. S7). Salinity also differentiated the coastal sites from the other two types in terms of community

composition. Since salinity is a key factor in influencing biogeochemical cycles, increased salinity can

shape specific microbial guilds [32-34]. Given that CW is frequently flooded by the brackish seawater

from the Baltic Sea, the high saline condition in CW strongly increased the abundance of microbes

carrying out sulfate respiration while decreasing the abundance of those carrying out methanogenesis

(Additional file 1: Fig. S13). Since sulfate is the main electron acceptor in saline habitats, sulfate reducers

with a higher affinity with substrates and a higher energy gain outcompete methanogens under these

conditions [35]. Moisture was another important factor that shaped the communities, and it was

significantly and positively correlated with diversities of prokaryotes and protists, as well as with other

properties including pH and concentrations of nutrients including organic compound, nitrogen and P

(Additional file 1: Fig. S7). This is in line with a previous study showing that increased water level by

permanent inundation released large amounts of nutrients into pore water, especially phosphorus and

ammonium [15]. Therefore, higher moisture might initially increase the availability of nutrients, which

then increases the microbial diversities and shapes the community structures in years of rewetting.

Interestingly, nitrate concentrations were much higher in AD and thus differentiated AD from the other

sites (Fig. 1b and 2c). This might be due to either the high nitrification activity generating nitrate and/or

due to higher external nitrate import via groundwater or precipitation. Moreover, vegetation was also an

important factor that cannot be neglected. According to our data, the plant community composition was

strongly correlated with both prokaryotic and eukaryotic community compositions, suggesting that plants

might be the driving factor of the congruent patterns observed between prokaryotes and eukaryotes.

**3. Influence of rewetting on microbial communities regarding ecological consequences**

Rewetting potentially promoted methanogenesis in AW and PW, but not in CW (Fig. 2), due to the high

amount of sulfate reducers in CW which is regularly flooded with sea water. The higher abundance of

methanogens in rewetted sites seems to be the main reason for the increase of $CH_4$ production and thus

potentially emission after rewetting [36]. It has been shown that $CH_4$ dominated the balances of annual

greenhouse gases in decades after rewetting a bog in Lower Saxony [16]. The dominance of $CH_4$ was also

observed in riparian wetlands after 12-year rewetting [37]. Since the fens in our study all have been

rewetted for over 12 years before our measurements commenced, the rewetting could have contributed to

high $CH_4$ emissions, which should to be tested with measurements of GHG exchange in the future.

However, it is also noteworthy that potential methanotrophs consuming $CH_4$ were also more abundant in

AW and PW (Fig. 2). ANME groups were only found in AW in our study, with a relative abundance of

0.15%, resulting in the highest methanotrophy abundance in AW, and this indicated that rewetting in this

site potentially accumulated the anaerobic methane oxidation. Hence, the alder forest and the coastal fen

seem to be the better choice to be rewetted regarding consumption and production of $CH_4$, respectively,

but this is of course a rather narrow perspective since the whole GHG budgets have to be considered.

One purpose of rewetting drained peatlands is to restore the carbon storage by cutting off oxygen supply

with inundation, thus decreasing the emitted $CO_2$ from soil respiration. One major process is the

decomposing of litter and deadwood by decomposers comprising some major bacteria and fungi [11, 12].

Microbes that carry out cellulolysis seemed to be the major prokaryotic decomposers in our study due to

their high abundance (Additional file 1: Fig. S13 and S14). We found that abundances of both cellulolysis

and fungi were strongly and negatively correlated with water content (Additional file 1: Fig. S16),

suggesting an underlying mechanism of the long-known fact that water content is a key factor controlling

the microbial decomposition activity in peat-soils. Considering that rewetting generally elevated the water

level, our results also illustrate the known fact that rewetting of drained peat-soils could inhibit the

microbial decomposition, thus conserving the carbon. This is further supported by some studies showing

that rewetting can reduce the carbon loss from peatlands by investigating net ecosystem changes of $CO_2$

and $CH_4$ [15, 38, 39].

Nitrification and denitrification are sources of $N_2O$ and the function prediction supported the changes in

these processes under wet conditions (Fig. 2). Since the relative abundances of potential denitrifiers were

much lower compared with those of nitrifiers (Additional file 1: Fig. S13 and S14), the production of $N_2O$

in these fens seems to stem primarily from nitrification. Potential nitrifiers were over-represented in

drained sites of all peatland types (Fig. 2), supporting the finding that rewetting generally reduces the

emission of $N_2O$ [17, 39].

Rewetting also showed influences on the eukaryotic communities. The lower diversities of fungi and

metazoa in rewetted sites might have been driven by the deficiency of oxygen under wet conditions since

both groups prefer aerobic living conditions. Some protists including Eugregarinorida, Gregarina and

*Monocystis* are known parasites of invertebrates, and they were over-represented in drained sites (Fig. 4).

Some metazoa known as parasites of plants and animals, including Tylenchida, Rhabditida and Enpleas,

were also over-represented in drained sites (Fig. 4). These results illustrate that rewetting might help

survivals of certain animal or plant species living in that area.

**Conclusions**

Our study showed the significant influence of rewetting on both prokaryotic and eukaryotic communities

in peat soils. To our knowledge, this is the first study that comprehensively described the changes of peat

communities regarding all domains of life across seasons after long-term rewetting, which bridges the gap

between rewetting practice and the induced ecological consequences. Congruent responses of prokaryotic

and eukaryotic communities to rewetting were observed and were driven by important abiotic and biotic

factors. Significant changes in potential microbial functions, fungi and animals indicate important

environmental consequences of rewetting relevant to carbon balance, GHG productions and parasitism.

Rewetting also induced different mechanistic relationships between microbial parameters and

environmental factors between different types of peatlands, suggesting that habitat-specific responses of

soil and water properties to rewetting should be aware, given that these are important factors driving

biogeochemical processes. Nevertheless, lack of temporal dynamics of the peat communities needs to be

verified with future studies in a longer term of investigation.

**Methods**

**Soil sampling**

The peat soils were sampled from these six sites in April 2017 (spring), August 2017 (summer),

November 2017 (autumn) and February 2018 (winter). At each site, samples were taken at three spots as

replicates at depths of 5-10 cm, 15-20 cm and 25-30 cm from the top soils. Samples were taken using a

soil core and then stored in sterilized Nasco Whirl-PAK baggies. The collected soils were immediately

blended and transported to the laboratory with an ice box. Soils were kept at 4 $^\circ$C before being processed

on the next day of sampling. In total, 216 samples (3 depths $\times$ 6 sites $\times$ 4 seasons $\times$ 3 replicates) were

collected.

**Soil edaphic properties**

Soils moisture was gravimetrically measured by drying the soil over night at 90 $^\circ$C until mass constancy.

Dissolved organic carbon (DOC) was extracted from 3 g fresh soil in 30-mL 0.1 M NaCl solution on a

shaker for 30 min at 200 rpm. Then the extracts were filtered using 0.45 µm sodium acetate filters. To

minimize possible artifacts acetate filters were carefully pre-rinsed with ca. 100 ml deionized water and

filtered samples were stored not longer than 1 week at 4°C before conducting DOM analysis [40]. A

size-exclusion chromatography (SEC) with organic carbon and organic nitrogen detection

(LC-OCD-OND analyzer, DOC-Labor Huber, Karlsruhe, Germany) was used to detect the concentrations

and composition of DOC based on size categories [41]. The detected organic compounds were

categorized into three groups: (i) biopolymers (BP-S), (ii) humic or humic-like substances including

building blocks (HL-S), and (iii) low molecular-weight substances (LM-S). The total amount of these

three groups was calculated as DOM. The ammonium molybdate spectrometric method (DIN EN 1189

D11) was used to determine soluble reactive phosphorus with a Cary 1E Spectrophotometer (Varian). The

photometry CFA method (Skalar SAN, Skalar Analytical B.V., The Netherlands) was used to

calorimetrically determine N-NH$_4^+$ and N-NO$_x^-$ concentrations according to the guidelines in EN ISO

11732 (DEV-E 23) and EN ISO 13395 (DEV, D 28), respectively. The pH and salinity of water samples

taken from wells installed at each study site were measured manually by digital meters in April 2017. The

data of the other three seasons were manually measured at groundwater wells with the water sensors

(Aquaread AP-2000 / AP-2000-D). Water levels and soil temperatures were also continuously monitored

at each site since then using Campbell Scientific CR300 Dataloggers (Logan, USA) and HOBO

Dataloggers (Bourne, USA), respectively.

**DNA extraction and sequencing**

DNA was extracted from 0.25 g soil using the DNeasy PowerSoil Kit (QIAGEN, Hilden, Germany)

according to the manufacturer's instructions with some modification. Vortex in the bead beating step was

replaced with a FastPrep®-24 5G instrument (MP Biomedicals, Santa Ana, USA), with an intensity of 5.0

m/s for 45 s. The extracted DNA samples were sent to LGC Genomics GmbH (Berlin, Germany) for 16S

rRNA and 18S rRNA gene amplicon sequencing. Primers pairs of 515YF

(5'-GTGYCAGCMGCCGCGGTAA-3')/B806R (5'-GGACTACNVGGGTWTCTAAT-3') [42] and

1183F (5'-AATTTGACTCAACRCGGG-3')/1443R (5'-AATTTGACTCAACRCGGG-3') [43] were

used to amplify 16S rRNA (prokaryotes) and 18S rRNA (eukaryotes) genes, respectively. Amplicons

were sequenced with Illumina Miseq 300 bp paired-end platform.

**Sequencing data analysis**

16S rRNA and 18S rRNA gene amplicon sequences were processed separately. For each processing, the

raw sequence reads were demultiplexed with barcodes, adapters and primers removed using the Illumina

*bcl2fastq* software. The data were then processed with *dada2* (v1.8.0) pipeline [44] in R v3.5. The

qualities of the sequences were checked, and sequences failing to meet the filter scores (maxEE=2,

truncQ=2, maxN=0) were discarded. The filtered sequences were de-replicated and clustered into ASVs,

and paired-end sequences were merged. The chimeric sequences were then de-novo checked and

removed. The final representative sequence of each ASV was assigned to taxonomy against a modified

version of the SILVA SSUref_NR_128 database [45]. ASVs with only one sequence were removed.

ASVs of 16S rRNA amplicons that were assigned to Chloroplast or mitochondria were also removed.

Several samples were discarded due to their low sequence numbers (<1,000), resulting in a total of 209

samples for further analysis. Finally, 16S rRNA and 18S rRNA sequences were clustered into 25,864 and

15,937 ASVs, respectively.

Plant sequences accounted for ~33% of the 18S rRNA amplicons, since peat-soil is mainly formed with

dead plants. We split plant sequences from eukaryotic ASVs to increase the resolution of the other

eukaryotes, but the plant sequences were also analyzed accordingly and separately. The diversity

(Shannon index) of both prokaryotes and eukaryotes (fungi, protists and metazoa) were calculated. The

tables of ASV counts were normalized using metagenomeSeq's CSS [46], and the community

compositions were analyzed using nonmetric NMDS based on Bray-Curtis dissimilarity distances. The

16S rRNA taxonomic profiles were converted to putative functional profiles using FAPROTAX which

maps prokaryotic clades to established metabolic or other ecologically relevant functions [47]. However,

some modifications were made. The ASVs involved in nitrification were further verified by blasting

against NCBI database according to a previous study [48]. The anaerobic methanotrophic groups ANME

were excluded from methanogenesis. Similarly, the fungal taxonomic profiles were also converted to

ecological guilds using FUNGuild [49].

A co-occurrence network was constructed to explore the potential interactions between species, including

prokaryotes, plants and other eukaryotes. ASVs with relative abundances lower than 0.01% were filtered.

The pairwise Spearman's rank correlations were performed with *Hmisc* package [50] and all *P*-values

were adjusted by the Benjamini and Hochberg false discovery rate (FDR) method. The cutoffs of

correlation coefficient and adjusted *P*-value were 0.8 and 0.01, respectively. The significant correlations

were visualized using *igraph* package [51]. Network consists of modules which are intensively connected

themselves but sparsely connected to other modules. These ecological clusters (modules) were identified

from this network using *igraph*. The relative abundances of prokaryotes and eukaryotes in each module

were calculated as sum of the relative abundances of the prokaryotic and eukaryotic ASVs belonging to

this module, respectively.

**Statistical analysis**

The statistical analyses were done with R v3.5. The significance of the difference among different seasons

or depths was identified with non-parametric Kruskal-Wallis test using *vegan* package [52].

Kruskal-Wallis posthoc tests were conducted to compare the means of alpha diversity, soil properties,

predicted functions and ecological clusters between each two groups of a factor using *PMCMR* package

[53]. The ASVs (relative abundance > 0.01%) that were differently abundant in drained and rewetted sites

were identified with differential expression analysis based on the Binomial distribution using *DESeq2*

package [54]. The same differential expression analysis was also performed on predicated prokaryotic and

fungal functions. Permutational Multivariate Analysis of Variance (PERMANOVA) was performed to

test the significance of impacts of rewetting, peatland type, depth, season and their interactions on the

prokaryotic and eukaryotic community compositions using *vegan* package. Mantel test was used to test

the correlations between prokaryotic, eukaryotic and plant community compositions with *vegan* package.

The pairwise Spearman's rank correlations were conducted to examine the relationships between

diversities, soil properties, community compositions, predicted functions and ecological clusters using

*Hmisc* package, and were plotted with *corrplot* package [55]. The soil properties were fitted into NMDS

ordination, and the significant ones were kept using the envfit function in the *vegan* package. All the

*P*-values for multiple comparisons were adjusted by FDR method and the null hypothesis was rejected

while *P*-values were less than 0.05.

SEM was performed to further find out the mechanism of rewetting effect on the microbial features. We

built a *priori* theoretical model based on the causal relationships among the variables (Additional file 1:

Fig. S17). We assume that rewetting could influence the microbial communities either directly or

indirectly through changing the water content and then the other soil parameters and the plant community

composition. To reduce the complexity of the model, 4 soil parameters were chosen according to the

correlation analysis (Additional file 1: Fig. S7). pH possibly drove DOM as well as other properties,

while nitrate showed less relevance with other properties but was driven by salinity (Additional file 1:

Fig. S7). We also reduced the dimensions of predicted functions and ecological clusters using NMDS

based on Bray-Curtis dissimilarities. The axes of NMDS were utilized in this model. SEM was

constructed using covariance-based method with *lavaan* package [56]. Since we found that peatland type

mostly drove the microbial communities (Fig. 1b and c), it is interesting to find out the different

functional mechanisms of rewetting across these 3 peatlands. We therefore implemented the group

analysis using the *sem* function with peatland type as the group in *lavaan*. Before that, all variables were

checked for normality, and the non-normally distributed ones were transformed using a two-step method

[57]. Since the multivariate normality of the final dataset still showed significant multivariate skew and

kurtosis, we used the Satorra-Bentler (S-B) procedure to correct the fitting statistics. The initial model

showed a weak yet adequate fit to the data (S-B $\chi^2$=4.752, *P*=0.191). To improve this model, we then

constrained the coefficients of two paths that were not significant in all 3 peatland types to zero

(Additional file 1: Fig. S17). The constrained model and unconstrained model showed no significant

difference with scaled chi-square difference test ($P$=0.786). Using the same method, we also constrained

that regressions among different fens were equal, and the significant difference ($P$<0.001) between

constrained and unconstrained models indicated that our models were different between these 3 peatlands.

**Acknowledgements**

We thank John Couwenberg from the University of Greifswald for collecting ideas and integrating the

data. We also thank Florian Beyer from the University of Rostock for providing the site maps.

**Funding**

This study was supported by the European Social Fund (ESF) and the Ministry of Education, Science and

Culture of Mecklenburg-Western Pomerania (Germany) within the scope of the project WETSCAPES

(ESF/14-BM-A55-0034/16 and ESF/14-BM-A55-0030/16).

**Availability of data and material**

All the sequencing files were deposited to the European Nucleotide Archive of EMBL (European

Molecular Biology Laboratory). The study accession number is PRJEB36764.

**Authors' contributions**

HW, MW and TU conceived and designed the research; HW, MW and DM collected the samples and

performed the lab work; DZ measured the soil edaphic properties, and AG provided the environmental data;

GJ helped to generate the structure of this manuscript; HW analyzed the data and wrote the first draft of the

manuscript, and all authors contributed to revisions. All authors read and approved the final manuscript.

**Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Author details**

[1]Institute of Microbiology, University of Greifswald, Germany. [2]Department of Bioscience, Aarhus

University, Denmark. [3]Department of Biogeochemistry and Chemical Analytics, Leibniz-Institute of

Freshwater Ecology and Inland Fisheries Berlin, Germany. [4]Chair of Landscape Ecology and Site

Evaluation, University of Rostock, Germany.

**References**

1.   Lamers LP, Vile MA, Grootjans AP, Acreman MC, van Diggelen R, Evans MG, Richardson CJ,

Rochefort L, Kooijman AM, Roelofs JG *et al*. Ecological restoration of rich fens in Europe and

North America: from trial and error to an evidence-based approach. Biol Rev Camb Philos Soc.

2015;90(1):182-203.

2.   Page SE, Baird AJ. Peatlands and Global Change: Response and Resilience. Annu Rev Env

Resour. 2016;41(1):35-57.

3.   Zedler JB, Kercher S. WETLAND RESOURCES: Status, Trends, Ecosystem Services, and

Restorability. Annu Rev Env Resour. 2005;30(1):39-74.

4.   Haddaway NR, Burden A, Evans CD, Healey JR, Jones DL, Dalrymple SE, Pullin AS.

Evaluating effects of land management on greenhouse gas fluxes and carbon balances in

boreo-temperate lowland peatland systems. Environ Evidence. 2014;3(1):5.

5.   Martikainen PJ, Nykänen H, Crill P, Silvola J. Effect of a lowered water table on nitrous oxide

fluxes from northern peatlands. Nature. 1993;366(6450):51.

6.   Kasimir□Klemedtsson Å, Klemedtsson L, Berglund K, Martikainen P, Silvola J, Oenema O.

Greenhouse gas emissions from farmed organic soils: a review. Soil Use Manage.

1997;13:245-250.

7.   Freeman C, Ostle N, Kang H. An enzymic'latch'on a global carbon store. Nature.

2001;409(6817):149.

8.   Wetzel RG. Gradient-dominated ecosystems: sources and regulatory functions of dissolved

organic matter in freshwater ecosystems. In: Dissolved organic matter in lacustrine ecosystems.

Springer; 1992. p. 181-198.

9.   Zak D, Roth C, Unger V, Goldhammer T, Fenner N, Freeman C, Jurasinski G. Unravelling the

importance of polyphenols for microbial carbon mineralization in rewetted riparian peatlands.

Front Env Sci Eng. 2019;7:147.

10.    Zak D, Reuter H, Augustin J, Shatwell T, Barth M, Gelbrecht J, McInnes RJB. Changes of the
$CO_2$ and $CH_4$ production potential of rewetted fens in the perspective of temporal vegetation
shifts. Biogeosciences. 2015;12(8):2455-2468.

11.    Bani A, Pioli S, Ventura M, Panzacchi P, Borruso L, Tognetti R, Tonon G, Brusetti L. The role of
microbial community in the decomposition of leaf litter and deadwood. Appl Soil Ecol.
2018;126:75-84.

12.    Van Geffen KG, Poorter L, Sass-Klaassen U, Van Logtestijn RS, Cornelissen JH. The trait
contribution to wood decomposition rates of 15 Neotropical tree species. Ecology.
2010;91(12):3686-3697.

13.    Nazaries L, Murrell JC, Millard P, Baggs L, Singh BK. Methane, microbes and models:
fundamental understanding of the soil methane cycle for future predictions. Environ Microbiol.
2013;15(9):2395-2417.

14.    Thamdrup B. New pathways and processes in the global nitrogen cycle. Annu Rev Ecol
Evolution S. 2012;43:407-428.

15.    van de Riet BP, Hefting MM, Verhoeven JTA. Rewetting Drained Peat Meadows: Risks and
Benefits in Terms of Nutrient Release and Greenhouse Gas Exchange. Water Air Soil Poll.
2013;224(4).

16.    Vanselow-Algan M, Schmidt SR, Greven M, Fiencke C, Kutzbach L, Pfeiffer EM. High methane
emissions dominated annual greenhouse gas balances 30 years after bog rewetting.
Biogeosciences. 2015;12(14):4361-4371.

17.    Wilson D, Blain D, Couwenberg J, Evans C, Murdiyarso D, Page S, Renou-Wilson F, Rieley J,
Sirin A, Strack M. Greenhouse gas emission factors associated with rewetting of organic soils.
Mires Peat. 2016;17.

18.    Vroom RJE, Xie F, Geurts JJM, Chojnowska A, Smolders AJP, Lamers LPM, Fritz C. Typha
latifolia paludiculture effectively improves water quality and reduces greenhouse gas emissions in
rewetted peatlands. Ecol Eng. 2018;124:88-98.

19.    Peacock M, Jones TG, Airey B, Johncock A, Evans CD, Lebron I, Fenner N, Freeman C. The
effect of peatland drainage and rewetting (ditch blocking) on extracellular enzyme activities and
water chemistry. Soil Use Manage. 2015;31(1):67-76.

20.    Zak D, Gelbrecht JJB. The mobilisation of phosphorus, organic carbon and ammonium in the
599           initial stage of fen rewetting (a case study from NE Germany). Biogeochemistry.
600           2007;85(2):141-151.

601    21.    Cabezas A, Gelbrecht J, Zak D. The effect of rewetting drained fens with nitrate-polluted water
602           on dissolved organic carbon and phosphorus release. Ecol Eng. 2013;53:79-88.

603    22.    Fuhrman JA. Microbial community structure and its functional implications. Nature.
604           2009;459(7244):193-199.

605    23.    Fuhrman JA, Cram JA, Needham DM. Marine microbial community dynamics and their
606           ecological interpretation. Nat Rev Microbiol. 2015;13(3):133-146.

607    24.    Gunadi B, Verhoef HA, Bedaux JJ. Seasonal dynamics of decomposition of coniferous leaf litter
608           in a forest plantation (*Pinus merkusii*) in Central Java, Indonesia. Soil Biol Biochem.
609           1998;30(7):845-852.

610    25.    Epron D, Le Dantec V, Dufrene E, Granier A. Seasonal dynamics of soil carbon dioxide efflux
611           and simulated rhizosphere respiration in a beech forest. Tree Physiology. 2001;21(2-3):145-152.

612    26.    Wang H, Liao G, D'Souza M, Yu X, Yang J, Yang X, Zheng T. Temporal and spatial variations
613           of greenhouse gas fluxes from a tidal mangrove wetland in Southeast China. Environ Sci Pollut
614           Res Int. 2016;23(2):1873-1885.

615    27.    Yu K, Faulkner SP, Baldwin MJ. Effect of hydrological conditions on nitrous oxide, methane,
616           and carbon dioxide dynamics in a bottomland hardwood forest and its implication for soil carbon
617           sequestration. Global Change Biol. 2008;14(4):798-812.

618    28.    Zerbe S, Steffenhagen P, Parakenings K, Timmermann T, Frick A, Gelbrecht J, Zak D.
619           Ecosystem service restoration after 10 years of rewetting peatlands in NE Germany. Environ
620           Manage. 2013;51(6):1194-1209.

621    29.    Faust K, Raes J. Microbial interactions: from networks to models. Nat Rev Microbiol.
622           2012;10(8):538-550.

623    30.    Fierer N, Schimel JP, Holden PA. Variations in microbial community composition through two
624           soil depth profiles. Soil Biol Biochem. 2003;35(1):167-176.

625    31.    Weil M, Wang H, Bengtsson M, Günther A, Jurasinski G, Couwenberg J, Negassa W, Zak D,
626           Urich T. Rewetting of three drained peatlands drives congruent compositional changes in pro-

and eukaryotic microbiomes through environmental filtering. bioRxiv. 2019:848192. doi:
http://dx.doi.org/10.1101/848192.

32.   Herlemann DP, Lundin D, Andersson AF, Labrenz M, Jurgens K. Phylogenetic Signals of
Salinity and Season in Bacterial Community Composition Across the Salinity Gradient of the
Baltic Sea. Front Microbiol. 2016;7:1883.

33.   Fan LF, Tang SL, Chen CP, Hsieh HL. Diversity and composition of sulfate- and sulfite-reducing
prokaryotes as affected by marine-freshwater gradient and sulfate availability. Microb Ecol.
2012;63(1):224-237.

34.   Wang H, Gilbert JA, Zhu Y, Yang X. Salinity is a key factor driving the nitrogen cycling in the
mangrove sediment. Sci Total Environ. 2018;631-632:1342-1349.

35.   Purdy K, Nedwell D, Embley T. Analysis of the sulfate-reducing bacterial and methanogenic
archaeal populations in contrasting Antarctic sediments. Appl Environ Microbiol.
2003;69(6):3181-3191.

36.   Wen X, Unger V, Jurasinski G, Koebsch F, Horn F, Rehder G, Sachs T, Zak D, Lischeid G,
Knorr K-H *et al*. Predominance of methanogens over methanotrophs in rewetted fens
characterized by high methane emissions. Biogeosciences. 2018;15(21):6519-6536.

37.   Kandel TP, Lærke PE, Hoffmann CC, Elsgaard L. Complete annual $CO_2$, $CH_4$, and $N_2O$ balance
of a temperate riparian wetland 12 years after rewetting. Ecol Eng. 2019;127:527-535.

38.   Karki S, Elsgaard L, Kandel TP, Laerke PE. Carbon balance of rewetted and drained peat soils
used for biomass production: a mesocosm study. GCB Bioenergy. 2016;8(5):969-980.

39.   Karki S, Elsgaard L, Audet J, Lærke PE. Mitigation of greenhouse gas emissions from reed
canary grass in paludiculture: effect of groundwater level. Plant Soil. 2014;383(1-2):217-230.

40.   Heinz M, Zak D. Storage effects on quantity and composition of dissolved organic carbon and
nitrogen of lake water, leaf leachate and peat soil water. Water Res. 2018;130:98-104.

41.   Huber SA, Balz A, Abert M, Pronk W. Characterisation of aquatic humic and non-humic matter
with size-exclusion chromatography-organic carbon detection–organic nitrogen detection
(LC-OCD-OND). Water Res. 2011;45(2):879-885.

42.   Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA, Jansson
JK, Caporaso JG, Fuhrman JA. Improved bacterial 16S rRNA gene (V4 and V4-5) and fungal

internal transcribed spacer marker gene primers for microbial community surveys. mSystems.
2016;1(1):e00009-00015.

43. Hadziavdic K, Lekang K, Lanzen A, Jonassen I, Thompson EM, Troedsson C. Characterization
of the 18S rRNA gene for designing universal eukaryote specific primers. PLoS One.
2014;9(2):e87624.

44. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
high-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581.

45. Lanzén A, Jørgensen SL, Huson DH, Gorfer M, Grindhaug SH, Jonassen I, Øvreås L, Urich T.
CREST–classification resources for environmental sequence tags. PloS One. 2012;7(11):e49334.

46. Paulson JN, Stine OC, Bravo HC, Pop M. Differential abundance analysis for microbial
marker-gene surveys. Nat Methods. 2013;10(12):1200.

47. Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean
microbiome. Science. 2016;353(6305):1272-1277.

48. Arce MI, von Schiller D, Bengtsson MM, Hinze C, Jung H, Alves RJE, Urich T, Singer G.
Drying and Rainfall Shape the Structure and Functioning of Nitrifying Microbial Communities in
Riverbed Sediments. Front Microbiol. 2018;9:2794.

49. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG.
FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild.
Fungal Ecol. 2016;20:241-248.

50. Frank E Harrell Jr, with contributions from Charles Dupont and many others. Hmisc: Harrell
Miscellaneous. R package version 4.1-1. 2018. https://CRAN.R-project.org/package=Hmisc

51. Csardi G, Nepusz T: The igraph software package for complex network research, InterJournal,
Complex Systems 1695. 2006. http://igraph.org

52. Jari Oksanen, F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan
McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H.
Stevens, Eduard Szoecs and Helene Wagner. vegan: Community Ecology Package. R package
version 2.5-1. 2018. https://CRAN.R-project.org/package=vegan

53. Pohlert T. The Pairwise Multiple Comparison of Mean Ranks Package (PMCMR). R package.
2014. https://CRAN.R-project.org/package=PMCMR

54.    Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biol;2014;15(12):550.

55.    Taiyun Wei, Viliam Simko. R package "corrplot": Visualization of a Correlation Matrix (Version
0.84). 2017. https://github.com/taiyun/corrplot

56.    Rosseel Y. Lavaan: An R package for structural equation modeling and more. Version 0.5–12
(BETA). J Stat Softw. 2012;48(2):1-36.

57.    Templeton GF. A two-step approach for transforming continuous variables to normal:
implications and recommendations for IS research. CAIS. 2011;28(1):41-58.

**Figure legends**

**Fig. 1** (**a**) Changes in Shannon index of prokaryotes and eukaryotes including fungi, protists and metazoa
in different seasons. Data are shown as mean values and the error bars represent standard errors. Asterisks
indicate the significant changes in different seasons in a corresponding site (Kruskal-Wallis test, *$P$<0.05,
**$P$<0.01). (**b**) Prokaryotic and (**c**) eukaryotic community compositions based on the Bray-Cutis
dissimilarities shown as NMDS plots. Venn diagrams show the number of shared prokaryotic and
eukaryotic ASVs associated with different peatland types. The main ordinations in NMDS plots show
similarity between samples and the arrows show correlations between environmental variables and
ordination axes. DOM, dissolved organic matter; BP-S, biopolymer substances; HL-S, humic-like
substances; LM-S, low-molecular substances; P, phosphorus.

**Fig. 2** The differentially abundant predicted functions of prokaryotes between rewetted and drained sites
in three different mires. Positive log2-fold change values indicate significantly higher abundances in
rewetted site (adjusted $P$<0.05), while negative values indicate significantly higher abundances in drained
site (adjusted $P$<0.05). The size of the points indicates the relative abundance of the functional groups.

**Fig. 3** The differentially abundant ASVs of prokaryotes between rewetted and drained sites in three
different mires. Positive log2-fold change values indicate significantly higher abundances in rewetted site

(adjusted *P*<0.01), while negative values indicate significantly higher abundances in drained site

(adjusted *P*<0.01). The size of the points indicates the relative abundance of the ASVs.

**Fig. 4** The differentially abundant ASVs of eukaryotes between rewetted and drained sites in three

different mires. Positive log2-fold change values indicate significantly higher abundances in rewetted site

(adjusted *P*<0.01), while negative values indicate significantly higher abundances in drained site

(adjusted *P*<0.01). The size of the points indicates the relative abundance of the ASVs.

**Fig. 5** (**a**) Co-occurrence network and identified modules. Prokaryotic nodes are shown as cycles while

eukaryotic nodes are shown as triangles. (**b**) The distribution of relative abundances of prokaryotic and

eukaryotic nodes within each module in different sites, depths or seasons. M, module; pro, prokaryote;

euk, eukaryote.

**Fig. 6** Structural equation model showing the direct and indirect effect of rewetting on the diversities,

community compositions, network ecological clusters and the predicted microbial functions. Paths that

showed significant relationships (*P*<0.05) are shown. The numbers along with arrows are standardized

path coefficients, and the numbers in the brackets indicate the number of axis (nMDS-axis 1 or axis 2).

Direct effects of rewetting and moisture are shown as red and blue arrows, respectively. The gray

double-sided arrows indicate the covariant relationships between the variables. Amount of variance

explained by the model ($R^2$) is listed for all response variables. DOM, dissolved organic matter; Pro,

prokaryote; Euk, eukaryote; Eco-clusters, ecological clusters derived from network analysis;

Micro-functions, microbial functions derived from FAPROTAX.

**Additional files**

**Additional file 1: Fig. S1.** Map showing locations of sampling sites. **Fig. S2.** Changes in soil edaphic

properties in different seasons. **Fig. S3.** Changes in soil edaphic properties in different depths. **Fig. S4.**

Groundwater level monitored every 15 min, shown as mean value of each day from 2017-9-22 to

2018-7-31. **Fig. S5.** Soil temperature monitored every 15 min at two depths (5 cm and 15 cm), shown as

mean value of two depths and of each day from 2017-7-24 to 2018-7-31. **Fig. S6.** Changes in Shannon

index of prokaryotes and eukaryotes including fungi, protists and metazoa in different depths. **Fig. S7.**

Pairwise Spearman's rank correlations between diversities, soil properties, community compositions,

predicted functions and ecological clusters. **Fig. S8.** The distributions of prokaryotic phyla or classes in

different sites, depths and seasons. **Fig. S9.** NMDS plots based on the Bray-Cutis dissimilarities showing

prokaryotic community composition among seasons (**a**) and depths (**b**), eukaryotic community

composition among seasons (**c**) and depths (**d**), and plant community composition (**e**) and fungal

community composition (**f**) associated with peatland types and water status. **Fig. S10.** NMDS plots based

on the Bray-Cutis dissimilarities showing prokaryotic community composition among seasons and depths

in AD (**a**), AW (**b**), CD (**c**), CW (**d**), PD (**e**) and PW (**f**). **Fig. S11.** The distributions of eukaryotic phyla in

different sites, depths and seasons. **Fig. S12.** NMDS plots based on the Bray-Cutis dissimilarities showing

eukaryotic community composition among seasons and depths in AD (**a**), AW (**b**), CD (**c**), CW (**d**), PD

(**e**) and PW (**f**). **Fig. S13.** Changes in predicted prokaryotic functions in different seasons. **Fig. S14.**

Changes in predicted prokaryotic functions in different depths. **Fig. S15.** The differentially abundant

predicted functions of fungi between rewetted and drained sites in three different peatlands. **Fig. S16.**

Linear regression between gravimetric water content and relative abundance of prokaryotes carrying

cellulolysis (**a**) or fungi (**b**). **Fig. S17.** A *priori* structural equation model. (PDF 16.6M)

**Additional file 2: Table S1.** Soil edaphic properties and Shannon indices of prokaryotes, fungi, protists

and metazoa in different sites, in different depths, and in different seasons. (XLXS 12KB)

**Fig. 1** (**a**) Changes in Shannon index of prokaryotes and eukaryotes including fungi, protists and metazoa in different seasons. Data are shown as mean values and the error bars represent standard errors. Asterisks indicate the significant changes in different seasons in a corresponding site (Kruskal-Wallis test, \*$P$<0.05, \*\*$P$<0.01). (**b**) Prokaryotic and (**c**) eukaryotic community compositions based on the Bray-Cutis dissimilarities shown as NMDS plots. Venn diagrams show the number of shared prokaryotic and eukaryotic ASVs associated with different peatland types. The main ordinations in NMDS plots show similarity between samples and the arrows show correlations between environmental variables and ordination axes. DOM, dissolved organic matter; BP-S, biopolymer substances; HL-S, humic-like substances; LM-S, low-molecular substances; P, phosphorus.

**Fig. 2** The differentially abundant predicted functions of prokaryotes between rewetted and drained sites in three different mires. Positive log2-fold change values indicate significantly higher abundances in rewetted site (adjusted $P<0.05$), while negative values indicate significantly higher abundances in drained site (adjusted $P<0.05$). The size of the points indicates the relative abundance of the functional groups.

**Fig. 3** The differentially abundant ASVs of prokaryotes between rewetted and drained sites in three different mires. Positive log2-fold change values indicate significantly higher abundances in rewetted site (adjusted *P*<0.01), while negative values indicate significantly higher abundances in drained site (adjusted *P*<0.01). The size of the points indicates the relative abundance of the ASVs.

**Fig. 4** The differentially abundant ASVs of eukaryotes between rewetted and drained sites in three different mires. Positive log2-fold change values indicate significantly higher abundances in rewetted site (adjusted *P*<0.01), while negative values indicate significantly higher abundances in drained site (adjusted *P*<0.01). The size of the points indicates the relative abundance of the ASVs.

**Fig. 5** (**a**) Co-occurrence network and identified modules. Prokaryotic nodes are shown as cycles while eukaryotic nodes are shown as triangles. (**b**) The distribution of relative abundances of prokaryotic and eukaryotic nodes within each module in different sites, depths or seasons. M, module; pro, prokaryote; euk, eukaryote.

**Fig. 6** Structural equation model showing the direct and indirect effect of rewetting on the diversities, community compositions, network ecological clusters and the predicted microbial functions. Paths that showed significant relationships (*P*<0.05) are shown. The numbers along with arrows are standardized path coefficients, and the numbers in the brackets indicate the number of axis (nMDS-axis 1 or axis 2). Direct effects of rewetting and moisture are shown as red and blue arrows, respectively. The gray double-sided arrows indicate the covariant relationships between the variables. Amount of variance explained by the model ($R^2$) is listed for all response variables. DOM, dissolved organic matter; Pro, prokaryote; Euk, eukaryote; Eco-clusters, ecological clusters derived from network analysis; Micro-functions, microbial functions derived from FAPROTAX.