Original Article

Immunoinformatic design of multi epitopes peptide-based universal cancer vaccine using matrix metalloproteinase-9 protein as a target.

Abdelrahman H. Abdelmoneim[*1], Mujahed I. Mustafa[2], Miysaa I. Abdelmageed[3], Naseem S. Murshed[4], Enas A. Dawoud[5], Enas M. Ahmed[6], Rahma M. Kamal eldin[6], Nafisa M. Elfadol[7], Anfal Osama M. Sati[5] and Abdelrafie M. Makhawi[2]

[1]Faculty of Medicine, Alneelain University, Khartoum, Sudan.
[2]Department of Biotechnology, University of Bahri, Khartoum, Sudan.
[3]Department of Pharmacy, University of Khartoum, Khartoum, Sudan
[4]Department of Microbiology, International University of Africa, Khartoum, Sudan.
[5]Faculty of Medical laboratory sciences, Razi University, Khartoum, Sudan.

[6]Faculty of Medicine, Karary University, Khartoum, Sudan.
[7]Department of Molecular biology, National University Biomedical, Khartoum, Sudan.

*Corresponding author: abduhamza009@gmail.com

Abstract:

Background: Cancer remains a major public health hazard despite the extensive research over the years on cancer diagnostic and treatment , this is mainly due to the complex pathophysiology and genetic makeup of cancer. A new approach toward cancer treatment is the use of cancer vaccine, yet the different molecular bases of cancers reduce the effectiveness of this approach. In this work we aim to use matrix metalloproteinase-9 protein (MMP9) which is essential molecule in the survival and metastasis of all type of cancer as a target for universal cancer vaccine design. **Method:** reference sequence of matrix metalloproteinase-9 protein was obtained from NCBI databases along with the related sequence, which is then checked for conservation using BioEdit, furthermore the B cell and T cell related peptide were analyzed using IEDB website. The best candidate peptide were then visualized using chimera software. **Result:** Three Peptides found to be good candidate for interactions with B cells (SLPE, RLYT, and PALPR), while ten peptides found as a good target for interactions with MHC1 (YRYGYTRVA, YGYTRVAEM, YLYRYGYTR, WRFDVKAQM, ALWSAVTPL, LLLQKQLSL, LIADKWPAL, KLFGFCPTR , MYPMYRFTE, FLIADKWPA) with world combined coverage of 94.77% . In addition, ten peptides were also found as a good candidates for interactions with MHC2 (KMLLFSGRRLWRFDV, GRGKMLLFSGRRLWR, RGKMLLFSGRRLWRF, GKMLLFSGRRLWRFD, TFTRVYSRDADIVIQ, AVIDDAFARAFALWS, FARAFALWSAVTPLT, MLLFSGRRLWRFDVK, GNQLYLFKDGKYWRF, NQLYLFKDGKYWRFS), with world combined coverage of 90.67%. **CONCLUSION:** 23

peptide-based vaccine was designed for use as a universal cancer vaccine which has a high world population coverage for MHC1(94.77%) and MHC2 (90.67%) related alleles.

1. Introduction:

Cancer is a chronic disease with varying degree of manifestations and high mortality and morbidity rates, in addition it is considered the second leading cause of death worldwide.(1, 2) All types of cancers cause common clinical symptoms including weight-loss, fatigue, pain, nausea, vomiting and constipation, etc.(3)Additional symptoms are specific to the site of malignancy.(4) Breast cancer is the most common site of cancer in women worldwide.(5) Lung cancer is the second most common cancer in both men and women while colorectal is the third most common type cancer in both sexes. (6)

Although many trials of therapeutic and preventive drugs to cancer has been carried out throughout the years. All of them have failed to obtain a positive response from all types of cancers and the FDA approved none.(7, 8) The reason for this failure is the fact that no effective intervention is available that covers all the pathological pathways of all types of cancer caused by the massive tumor heterogeneity.(9)

There are approximately 28 subtypes of matrix metalloproteinases (MMPs). They are responsible of maintaining the components of the extracellular matrix (ECM), and controlling metastasis of cancerous cells. *Matrix metalloproteinases-9 (MMP-9)* gene (MIM 120361) is located at chromosome 20q12-13. It encodes Matrix metalloproteinase- 9 protein, a major enzyme for the degradation and absorption of ECM, signaling of wound healing, bone resorption, angiogenesis and plays a crucial role in various types of cancer progression. Such as breast cancer, colorectal cancer, papillary thyroid cancer, urinary cancers, lymphomas and gastric cancer.(10-13) In addition, MMP-9 is considered as a biomarker as its level increases in inflammatory diseases, i.e., Rheumatoid arthritis, asthmas, and abdominal aortic aneurysm.(14, 15) This study aims to predict a safe multi-epitope based anti-cancer vaccine using human MMP-9 protein as a target.

MMP-9 was tested in mouse before for several purpose,(16, 17) but to the best of our knowledge, this is the first immunoinformatic study to use human MMP9 protein as an immunogenic target for a peptide-based vaccine design for all cancer types. Peptide based vaccines are safe, stable with minimal exposure to hazardous or allergenic substances, cheaper and less laboring.(18, 19) In Addition, reverse vaccinology techniques have proven its efficiency in general and in cancer field in particular, as many of the designed vaccines has made it into clinical trials.(20-22)

2. Materials and Methods:

2.1 Sequences retrieval:

The whole protein sequence of the MM9 in FASTA format, accessed on November 3,2019 (NP-004985.2) was retrieved from NCBI database (https://www.ncbi.nlm.nih.gov/protein).(23) On the other hand, the candidate epitopes were analyzed by IEDB analysis tools (http://www.iedb.org/).(24)

2.2 Identification of conserved regions:

Sequences were aligned to identify the conserved regions using BioEdit tool package version 7.2.5. Furthermore, epitope conservancy investigation for each epitopes was predicted by IEDB analysis resource. (http://tools.immuneepitope.org/tools/conservancy).(25)

2.3 B-cell epitope prediction:

B cell epitope is the part of a vaccine which interacts with B cell receptor. The epitopes of interest were investigated using numerous B-cell prediction algorisms to identify the antigenicity, flexibility, hydrophilicity and surface accessibility; the selection of highest immunogenic protein sequence is a prerequisite for epitope -based peptide vaccine design.(26)

2.3.1 Prediction of linear B-cell epitopes:

BepiPred from immune epitope database (http://toolsiedb.ofg/bcell/) [28] was used as linear B-cell epitopes prediction from the conserved region with a default threshold value of 0.35.all epitopes with values less than this threshold were chosen.

2.3.2 Prediction of surface accessibility:

By using Emini surface accessibility prediction tool of the immune epitope database (IEDB) (http://tools.immuneepitope.org/tools/bcell/iedb). [29] The surface accessible epitopes were predicted from the conserved region holding the default threshold value 1.000. we selected all peptides with values less than this threshold for further analysis.

2.3.3 Prediction of epitopes antigenicity :

The kolaskar and tongaonker antigenicity method was used to determine the antigenic sites with a default threshold value of 1.028 It is available at (http://tools.immuneepitope.org/bcell/),[30] all antigenic peptide with values less than the threshold were selected for further analysis.

2.4 T cell epitope prediction:

2.4.1 MHC class I binding predictions:

Peptide binding to MHC class I molecules was analyzed by the IEDB MHC I prediction tool (http://tools.iedb.org/mhc1.) ; based on Artificial Neural Network (ANN) .this tool has different algorithms to show the ability of the submitted sequence to bind to a specific MHC class 1 molecule.(24, 27-29) All epitope lengths were set as 9 amino acids, all conserved epitopes that

bind to alleles at score equal or less than 100 half-maximal inhibitory concentration (IC50) is selected for further analysis.(30)

2.4.2 MHC class II binding predictions:

Peptide binding to MHC class II molecules was analyzed by the IEDB MHC II prediction tool (http://tools.iedb.org/mhcII) (31, 32) the three major types of HLA alleles (DR, DQ and DP) were chosen for analysis based on NN-align, and all peptide with IC50 less than 500 were chosen for further workout.

2.5 Population coverage:

Diverse HLA alleles are very distributable world-wide population. Hence, the HLA-alleles distribution among the world population is critical for effective multi-epitope vaccine design. Predicted population coverage analysis of above-mentioned MHC class I and II epitopes were done by using IEDB tool against world-wide population. (http://tools.iedb.org/tools/population/iedb_input).(33)

2.6 3D Modeling:

The full 3D structure of MM9 was generated by RaptorX (http://raptorx.uchicago.edu/).(34) The accession number of the reference sequence is NP-004985.2 then UCSF Chimera was used to visualize the position of suggested peptides.(35)

3. Results:

Prediction of B cell epitope:

The metalloproteinase-9 protein (MMP-9) reference sequence was examined by using Bepipred linear epitope prediction, Kolaskar and Tongaonkar antigenicity and Emini surface accessibility methods in IEDB, to confine the probability of specific regions in the protein to bind to B cell receptor, being in the surface and immunogenic, respectively.

In bepipred linear epitope prediction method which was the first method used to determine the B cell epitopes, the average binder's score of MMP-9 protein to B cell was 0.321, with a minimum score of -0.006 and a maximum score of 3.227, Basically all values that equal or greater than the default threshold 0.350 were potentially linear epitopes. as shown in (table 1 and figure 1).

Table1: describe the predicted antigenic B cell epitopes, 23 antigenic sites were identified from MMP-9 protein

No.	Start	End	Peptide	Length
1	21	24	PRQR	4
2	59	67	EMRGESKSL	9
3	79	88	SLPETGELDS	10

4	96	102	TPRCGVP	7
5	129	135	SEDLPRA	7
6	173	184	AEHGDGYPFDGK	12
7	192	205	FPPGPGIQGDAHFD	14
8	221	228	RFGNADGA	8
9	241	269	YSACTTDGRSDGLPWCSTTANYDTDDRFG	29
10	275	288	RLYTQDGNADGKPC	14
11	299	312	YSACTTDGRSDGYR	14
12	318	321	ANYD	4
13	331	343	TRADSTVMGGNSA	13
14	358	369	YSTCTSEGRGDG	12
15	377	390	SNFDSDKKWGFCPD	14
16	425	437	FTEGPPLHKDDVN	13
17	443	515	YGPRPEPEPRPPTTTTPQPTAPPTVCPTGPPTVHPSERPTAGPTGPPSAGPTGPPTAGPSTATTVPLSPVDDA	73
18	544	553	EGRGSRPQGP	10
19	561	565	PALPR	5
20	616	619	ALRS	4
21	641	650	MVDPRSASEV	10
22	657	660	VPLD	4
23	685	690	RSELNQ	6

In Emini surface accessibility prediction, for an efficient B-cell epitope the average surface accessibility areas of the protein was scored as 1.000, with a maximum of 6.034 and a minimum of 0.031, so all the values equal or greater than the default threshold 1.000 were potentially in the surface. As well as, Kolaskar and Tongaonkar antigenicity prediction's the average of antigenicity was 1.028, with a maximum of 1.307 and minimum of 0.869; all values equal to or greater than the default threshold 1.028 were potential antigenic determinants. The results of conserved predicted B cell epitopes that passed both Emini & kolaskar tests are shown in (table 2 and figure 2&3).

Table 2: List of conserved B cell epitopes with their surface accessibility and antigenicity scores

Start	End	Peptide	Length	Conservation	EMINi score	Prediction	KOLSKAR score	Prediction
79	82	SLPE	4	C	1.079	P	1.044	P
275	278	RLYT	4	C	1.331	P	1.048	P
561	565	PALPR	5	C	1.104	P	1.063	P

Figure 1: Bepipred linear epitope prediction, the yellow peaks above the red line(Threshold)are proposed to be a part of B cell epitopes and the green peaks are not a part.

Figure 2: Emini Surface Accessibility Prediction, Yellow peaks above the red line(Threshold) are proposed to be a part of B cell epitopes while green peaks are not a part.

Figure 3: Kolaskar and Tongaonkar Antigenicity Prediction, Yellow peaks above the red line(Threshold) are proposed to be a part of B cell epitope while green peaks are not a part.

Prediction of T cell epitope:

MHC-I binding profile prediction for T cytotoxic cell conserved epitopes:

227 epitopes were anticipated to interact with diꢀerent MHC-1 alleles. The core epitopes (YRYGYTRVA/ YGYTRVAEM) were noticed to be the dominant binders with 7 alleles for each (HLA-C*06:02,HLA-C*07:01,HLA-C*12:03,HLA-C*14:02,HLA-B*27:05,HLA-C*07:02,HLA-B*39:01/HLA-C*03:03,HLA-C*14:02,HLA-C*12:03,HLA-C*06:02,HLA-B*08:01,HLA-C*15:02,HLA-C*07:01), followed by (YLYRYGYTR, WRFDVKAQM, ALWSAVTPL, LLLQKQLSL, LIADKWPAL, KLFGFCPTR, MYPMYRFTE, FLIADKWPA) which binds with (8, 5, 5, 3, 3, 5, 1, 3) alleles. (Table 3).

Table 3: List of most promising epitopes that had a good binding affinity with MHC-I alleles in terms of IC50.

Peptide	start	end	No. of MHC-I allele hits
YRYGYTRVA	50	58	7
YGYTRVAEM	52	60	7
YLYRYGYTR	48	56	8
WRFDVKAQM	633	641	5
ALWSAVTPL	146	154	5
LLLQKQLSL	72	80	3
LIADKWPAL	555	563	3
KLFGFCPTR	324	332	5
MYPMYRFTE	419	427	1
FLIADKWPA	554	562	3

MHC-II binding profile prediction for T helper cell conserved epitopes:

479 conserved predicted epitopes were found to interact with MHC-II alleles. The core epitopes (KMLLFSGRRLWRFDV) is thought to be the top binder as it interacts with 18 alleles; (HLA-

DRB1*01:01,HLA-DRB1*03:01,HLA-DRB1*04:02,HLA-DRB1*07:01,HLA-
DRB1*04:04,HLA-DRB1*08:01,HLA-DRB1*09:01,HLA-DRB1*13:01,HLA-
DRB1*11:01,HLA-DRB1*13:02,HLA-DRB1*12:01,HLA-DRB1*15:01,HLA-
DRB1*16:02,HLA-DRB3*01:01,HLA-DRB4*01:03,HLA-DRB5*01:01,HLA-
DRB3*03:01,HLA-DRB4*01:01).   Followed   by   (GRGKMLLFSGRRLWR,
RGKMLLFSGRRLWRF,   GKMLLFSGRRLWRFD,   TFTRVYSRDADIVIQ,
AVIDDAFARAFALWS,   FARAFALWSAVTPLT,   MLLFSGRRLWRFDVK,
GNQLYLFKDGKYWRF, NQLYLFKDGKYWRFS) which binds to (17, 17, 18, 14, 14, 17, 15,
14, 14) alleles. (Table 4).

Table 4: List of the most promising core sequence epitopes that had a strong binding affinity
with MHC-II in terms of IC50.

Peptide Sequence	Start	End	No. of MHC-II allele hits
KMLLFSGRRLWRFDV	623	637	18
GRGKMLLFSGRRLWR	620	634	17
RGKMLLFSGRRLWRF	621	635	17
GKMLLFSGRRLWRFD	622	636	18
TFTRVYSRDADIVIQ	155	169	14
AVIDDAFARAFALWS	135	149	14
FARAFALWSAVTPLT	141	155	17
MLLFSGRRLWRFDVK	624	638	15
GNQLYLFKDGKYWRF	528	542	14
NQLYLFKDGKYWRFS	529	543	14

Population coverage analysis

Population coverage test was performed to detect the world Coverage of all epitopes binds to
MHC1 alleles and MHC11 alleles, Exhibit an exceptional coverage with percentages of 94.77%,
90.67%. Respectively, as well as selected most promising epitopes for each test.
Population coverage for MHC1: ten epitopes are given to interact with the most frequent MHC
class I alleles (YRYGYTRVA, YGYTRVAEM, YLYRYGYTR, WRFDVKAQM,
ALWSAVTPL, LLLQKQLSL, LIADKWPAL, KLFGFCPTR, MYPMYRFTE and
FLIADKWPA), representing a considerable coverage against the whole world population. The
maximum population coverage percentage over these epitopes worldwide was found to be
62.74% for YRYGYTRVA (table 5, figure 4).

Table 5: Global Population coverage of promising epitopes binding to MHC class I alleles

Peptide	MHC I world coverage
YRYGYTRVA	62.74%

YGYTRVAEM	58.08%
YLYRYGYTR	54.43%
WRFDVKAQM	54.29%
ALWSAVTPL	50.41%
LLLQKQLSL	46.87%
LIADKWPAL	45.42%
KLFGFCPTR	43.03%
MYPMYRFTE	43.03%
FLIADKWPA	42.53%

Figure 4. Illustrates the global coverage for the top ten MHC-I peptides (YRYGYTRVA, YGYTRVAEM, YLYRYGYTR, WRFDVKAQM, ALWSAVTPL, LLLQKQLSL, LIADKWPAL, KLFGFCPTR, MYPMYRFTE and FLIADKWPA). Note: In the graph, the line (-o-) represents the cumulative percentage of population coverage of the epitopes; the bars represent the population coverage for each epitope.

Population coverage for MHC11: In the case of MHC class II, ten epitopes were assumed to interact with the most frequent MHC class II alleles (KMLLFSGRRLWRFDV, GRGKMLLFSGRRLWR, RGKMLLFSGRRLWRF, GKMLLFSGRRLWRFD, TFTRVYSRDADIVIQ, AVIDDAFARAFALWS, FARAFALWSAVTPLT, MLLFSGRRLWRFDVK, GNQLYLFKDGKYWRF, NQLYLFKDGKYWRFS), inferring a massive global coverage. The highest population coverage percentage of these epitopes worldwide was that of KMLLFSGRRLWRFDV with percentage of 83.74% % (table 6, figure 5).

Table 6: Global Population coverage of promising epitopes in isolated MHC class II

Peptide Sequence	MHC II world coverage
KMLLFSGRRLWRFDV	83.74%
GRGKMLLFSGRRLWR	83.46%
RGKMLLFSGRRLWRF	83.46%
GKMLLFSGRRLWRFD	83.14%
TFTRVYSRDADIVIQ	81.42%
AVIDDAFARAFALWS	80.82%
FARAFALWSAVTPLT	79.23%
MLLFSGRRLWRFDVK	78.71%
GNQLYLFKDGKYWRF	78.50%
NQLYLFKDGKYWRFS	78.50%

Figure 5. Illustrates the global proportion for the top ten MHC-II epitopes (KMLLFSGRRLWRFDV, GRGKMLLFSGRRLWR, RGKMLLFSGRRLWRF, GKMLLFSGRRLWRFD, TFTRVYSRDADIVIQ, AVIDDAFARAFALWS, FARAFALWSAVTPLT, MLLFSGRRLWRFDVK, GNQLYLFKDGKYWRF, NQLYLFKDGKYWRFS). Notes: In the graph, the line (-o-) represents the cumulative percentage of population coverage of the epitopes; the bars represent the population coverage for each epitope.

Figure 6. Present 3D structure of metalloproteinase-9 protein visualizing top ten T-cell peptides binding to MHC class I using chimera (version 1.14).yellow color indicate the chosen peptides while the purple color indicate the rest of the molecules.

Figure 7. Present 3D structure of metalloproteinase-9 protein visualizing top ten T-cell peptides binding to MHC class II, using chimera (version 1.14).yellow color indicate the chosen peptides while the purple color indicate the rest of the molecules.

Figure 8. Present 3D structure of metalloproteinase-9 protein visualizing most promisisng B-cell peptides that have high suface accessibility and antigenicity scores, using chimera (version 1.14).yellow color indicate the chosen peptides while the purple color indicate the rest of the molecules.

4. **Discussion:**

Multi Epitope peptides will be the next generation of cancer immunotherapy. However, an efficient vaccine progression and production are costly and time consuming, but the development of immunoinformatics tools has made a broader way in developing vaccines through understanding the immune response within a short amount of time.(36) Many research aimed to design universal cancer vaccines using different approaches, yet until now, there is no FDA approved universal cancer vaccine.(37) The current work used immunogenic protein (matrix metalloproteinase-9) that is essential for the survival of all type of cancer as a target for vaccine design, where three Peptides found to be good candidate for interactions with B cells, in addition ten peptides found as a good target for interactions both MHC1 and MHC2.

Most cancer vaccines depend on T cell interactions but recent research indicates the significant role B cell plays in the immunology of cancer vaccine, B cell produces antibodies and help in the regulation of other immune reaction.(38)furthermore, several B cell based cancer vaccine studies show Promising results in chronic lymphocytic leukemia and other types of tumors.(39) In the current work, Linear B cell epitopes of matrix metalloproteinase-9 analyzed by Bepipred linear epitope prediction tool. After checking for conservation using BioEdit software **(figure 1),** the epitopes were checked for surface accessibility and antigenicity by using the Emini surface accessibility prediction method and Kolaskar and Tongaonkar antigenicity method respectively. The following peptides have the highest score and were approved as the most promising B cell epitopes (SLPE, RLYT, and PALPR).

T cell receptor can only identify epitopes if it is presented on major histocompatibility complex, Epitopes binds in linear form to MHCs, they linked together into the binding groove of MHC class I and class II through interactions with the amino acids located in the pockets on the floor of MHC.(40, 41)

Cytotoxic T cell(CD8) can only recognize antigen when it is presented on MHC1, and so we analyzed the peptides that has the best binding to MHC1 and found 10 candidates(YRYGYTRVA, YGYTRVAEM,YLYRYGYTR,WRFDVKAQM, ALWSAVTPL, LLLQKQLSL, LIADKWPAL, KLFGFCPTR , MYPMYRFTE, FLIADKWPA)**(Table 1).** It is interesting to note that among the associated alleles related to these peptides, we found the most frequent reported allele in Caucasians which is HLA-A*0201.(42) This further proves the significant coverage of these peptides.

In the case of T helper cells, it can only recognize antigen when it is presented in MHC2. Therefore, we predicted the peptide of MMP9 that can firmly interact with MHC2 and end up with the following 10 peptides sequence (KMLLFSGRRLWRFDV, GRGKMLLFSGRRLWR, RGKMLLFSGRRLWRF, GKMLLFSGRRLWRFD, TFTRVYSRDADIVIQ, AVIDDAFARAFALWS, FARAFALWSAVTPLT, MLLFSGRRLWRFDVK, GNQLYLFKDGKYWRF, and NQLYLFKDGKYWRFS). (Table 2)

The MHC is divided into 3 functionally distinct regions: class I, II and III regions, the population coverage of the epitopes and their respective HLA alleles are crucial, as HLA alleles are highly polymorphic in nature.(43)

We aim for this vaccine to benefit patient from different parts of the world, therefore, we analyzed the whole word population coverage for our predicted peptide, where we found MHC1 and MHC2 to have population coverage of 94.77% and 90.67% Respectively **(Table 1 and 2).** Therefor our vaccine is assumed to elicit a strong comprehensive immune response against our targeted protein matrix Metallopeptidase 9 (MMP9) in the majority of vaccinated population.

All peptides were visualized using chimera software **(Figures 2 and 3)** which shows the predicted structures and positions of these peptides in MMP9 protein.

In a study done by Roomi et al, where they used MMP9 as anticancer vaccine in mice, they found 80% reduction in tumor volume after using this vaccine.(44)Which further proves the predicted effectiveness of our proposed vaccine. We took further step in our work to analyze this target using human database.which confirm our hypothesis about the role of MMP9 as agood candidate for vaccine.

Previous research by Dosset et al. designed a universal peptide based cancer vaccine using telomerase reverse transcriptase (TERT) which showed improve response against CD4 and CD8, but the role of B cell was not investigated in this research.(45)

This study is limited due to the few numbers of reference sequence we were able to acquire from NCBI, along with the 6 alleles(HL-DRB3*03:01, HL-DRB3*02:01, HL-DRB4*01:03, HL-DRB3*01:01, HL-DRB4*01:01, HL-DRB5*01:01) that was not applicable for prediction in IEDB.in addition to the limited therapeutic activity of peptide vaccine in general.(46)

In the end, 23 peptides were chosen as a target for universal cancer vaccine design, yet Wet lab studies are required to confirm these results, in order to move another step towered the goal of eradicating cancer from the face of the earth.

5. Conclusion:

The current work designs multi epitopes peptide based universal cancer vaccine using MMP9 protein as a target where three Peptides found to be good candidate for interactions with B cells, and ten peptides found as a good target for interactions with each MHC1 and MHC2.

6. Conflict of interest:

The authors decline the existence of any conflict of interest regarding this paper.

7. Acknowledgement:
The authors acknowledge the Deanship of Scientific Research at University of Bahri for the supportive cooperation.

1.   Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. CA: A Cancer Journal for Clinicians. 2019;69(1):7-34.
2.   Nagai H, Kim YH. Cancer prevention from the perspective of global cancer burden patterns. J Thorac Dis. 2017;9(3):448-51.

3.	Miaskowski C, Lee KA. Pain, Fatigue, and Sleep Disturbances in Oncology Outpatients Receiving Radiation Therapy for Bone Metastasis: A Pilot Study. Journal of Pain and Symptom Management. 1999;17(5):320-32.

4.	Scheel BI, Holtedahl K. Symptoms, signs, and tests: The general practitioner's comprehensive approach towards a cancer diagnosis. Scand J Prim Health Care. 2015;33(3):170-7.

5.	Momenimovahed Z, Salehiniya H. Epidemiological characteristics of and risk factors for breast cancer in the world. Breast Cancer (Dove Med Press). 2019;11:151-64.

6.	Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. CA: A Cancer Journal for Clinicians. 2020;70(1):7-30.

7.	Maeda H, Khatami M. Analyses of repeated failures in cancer therapy for solid tumors: poor tumor-selective drug delivery, low therapeutic efficacy and unsustainable costs. Clin Transl Med. 2018;7(1):11-.

8.	Dalgleish AG, Stern PL. The failure of radical treatments to cure cancer: can less deliver more? Therapeutic Advances in Vaccines and Immunotherapy. 2018;6(5-6):69-76.

9.	Aurisicchio L, Pallocca M, Ciliberto G, Palombo F. The perfect personalized cancer therapy: cancer vaccines against neoantigens. Journal of Experimental & Clinical Cancer Research. 2018;37(1):86.

10.	Felizi RT, Veiga MG, Carelli Filho I, Souto RPd, Fernandes CE, Oliveira E. Association between Matrix Metallopeptidase 9 Polymorphism and Breast Cancer Risk. Rev Bras Ginecol Obstet. 2018;40(10):620-4.

11.	Wang R-X, Chen S, Huang L, Shao Z-M. Predictive and prognostic value of Matrix metalloproteinase (MMP) - 9 in neoadjuvant chemotherapy for triple-negative breast cancer patients. BMC Cancer. 2018;18(1):909.

12.	Schlom J, Arlen PM, Gulley JL. Cancer vaccines: moving beyond current paradigms. Clinical Cancer Research. 2007;13(13):3776-82.

13.	Zarkesh M, Zadeh-Vakili A, Akbarzadeh M, Fanaei SA, Hedayati M, Azizi F. The role of matrix metalloproteinase-9 as a prognostic biomarker in papillary thyroid cancer. BMC cancer. 2018;18(1):1199.

14.	Maguire EM, Pearce SWA, Xiao R, Oo AY, Xiao Q. Matrix Metalloproteinase in Abdominal Aortic Aneurysm and Aortic Dissection. Pharmaceuticals (Basel). 2019;12(3):118.

15.	Tokito A, Jougasaki M. Matrix Metalloproteinases in Non-Neoplastic Disorders. International Journal of Molecular Sciences. 2016;17(7).

16.	Fields GB. Mechanisms of Action of Novel Drugs Targeting Angiogenesis-Promoting Matrix Metalloproteinases. Frontiers in immunology. 2019;10:1278-.

17.	Huang H. Matrix Metalloproteinase-9 (MMP-9) as a Cancer Biomarker and MMP-9 Biosensors: Recent Advances. Sensors (Basel). 2018;18(10):3249.

18.	Li W, Joshi M, Singhania S, Ramsey K, Murthy A. Peptide vaccine: progress and challenges. Vaccines. 2014;2(3):515-36.

19.	Rahat MA. Targeting Angiogenesis With Peptide Vaccines. Frontiers in Immunology. 2019;10:1924.

20.	Okuno K, Sugiura F, Hida J-i, Tokoro T, Ishimaru E, Sukegawa Y, et al. Phase I clinical trial of a novel peptide vaccine in combination with UFT/LV for metastatic colorectal cancer. ExpErimEntal and thErapEutic mEdicinE. 2011;2(1):73-9.

21.	Obara W, Kanehira M, Katagiri T, Kato R, Kato Y, Takata R. Present status and future perspective of peptide-based vaccine therapy for urological cancer. Cancer science. 2018;109(3):550-9.

22.	Sawada Y, Yoshikawa T, Nobuoka D, Shirakawa H, Kuronuma T, Motomura Y, et al. Phase I trial of a glypican-3–derived peptide vaccine for advanced hepatocellular carcinoma: immunologic evidence and potential for improving overall survival. Clinical Cancer Research. 2012;18(13):3686-96.

23.	NCBI protein sequence database. 2019.

24.   Kim Y, Ponomarenko J, Zhu Z, Tamang D, Wang P, Greenbaum J, et al. Immune epitope database analysis resource. Nucleic acids research. 2012;40(Web Server issue):W525-30.

25.   Bui HH, Sidney J, Li W, Fusseder N, Sette A. Development of an epitope conservancy analysis tool to facilitate the design of epitope-based diagnostics and vaccines. BMC bioinformatics. 2007;8:361.

26.   Larsen JE, Lund O, Nielsen M. Improved method for predicting linear B-cell epitopes. Immunome research. 2006;2:2.

27.   Kolaskar AS, Tongaonkar PC. A semi-empirical method for prediction of antigenic determinants on protein antigens. FEBS letters. 1990;276(1-2):172-4.

28.   Nielsen M, Lundegaard C, Worning P, Lauemoller SL, Lamberth K, Buus S, et al. Reliable prediction of T-cell epitopes using neural networks with novel sequence representations. Protein science : a publication of the Protein Society. 2003;12(5):1007-17.

29.   Peters B, Sette A. Generating quantitative models describing the sequence specificity of biological processes with the stabilized matrix method. BMC bioinformatics. 2005;6:132.

30.   Sidney J, Assarsson E, Moore C, Ngo S, Pinilla C, Sette A, et al. Quantitative peptide binding motifs for 19 human and mouse MHC class I molecules derived using positional scanning combinatorial peptide libraries. Immunome research. 2008;4:2.

31.   Wang P, Sidney J, Dow C, Mothe B, Sette A, Peters B. A systematic assessment of MHC class II peptide binding predictions and evaluation of a consensus approach. PLoS computational biology. 2008;4(4):e1000048.

32.   Wang P, Sidney J, Kim Y, Sette A, Lund O, Nielsen M, et al. Peptide binding predictions for HLA DR, DP and DQ molecules. BMC bioinformatics. 2010;11:568.

33.   Bui HH, Sidney J, Dinh K, Southwood S, Newman MJ, Sette A. Predicting population coverage of T-cell epitope-based diagnostics and vaccines. BMC bioinformatics. 2006;7:153.

34.   Wang S, Li W, Liu S, Xu J. RaptorX-Property: a web server for protein structure property prediction. Nucleic acids research. 2016;44(W1):W430-5.

35.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera--a visualization system for exploratory research and analysis. Journal of computational chemistry. 2004;25(13):1605-12.

36.   Khan M, Khan S, Ali A, Akbar H, Sayaf AM, Khan A, et al. Immunoinformatics approaches to explore Helicobacter Pylori proteome (Virulence Factors) to design B and T cell multi-epitope subunit vaccine. Sci Rep. 2019;9(1):13321.

37.   Lokhov PG, Balashova EE. Design of universal cancer vaccines using natural tumor vessel-specific antigens (SANTAVAC). Hum Vaccin Immunother. 2015;11(3):689-98.

38.   Wennhold K, Shimabukuro-Vornhagen A, von Bergwelt-Baildon M. B Cell-Based Cancer Immunotherapy. Transfus Med Hemother. 2019;46(1):36-46.

39.   Biagi E, Rousseau R, Yvon E, Schwartz M, Dotti G, Foster A, et al. Responses to human CD40 ligand/human interleukin-2 autologous cell vaccine in patients with B-cell chronic lymphocytic leukemia. Clin Cancer Res. 2005;11(19 Pt 1):6916-23.

40.   Zvyagin IV, Tsvetkov VO, Chudakov DM, Shugay M. An overview of immunoinformatics approaches and databases linking T cell receptor repertoires to their antigen specificity. Immunogenetics. 2019.

41.   Kennedy PR, Barthen C, Williamson DJ, Davis DM. HLA-B and HLA-C Differ in Their Nanoscale Organization at Cell Surfaces. Front Immunol. 2019;10:61.

42.   Pelte C, Cherepnev G, Wang Y, Schoenemann C, Volk HD, Kern F. Random screening of proteins for HLA-A*0201-binding nine-amino acid peptides is not sufficient for identifying CD8 T cell epitopes recognized in the context of HLA-A*0201. J Immunol. 2004;172(11):6783-9.

43.   Glanville KP, Coleman JRI, Hanscombe KB, Euesden J, Choi SW, Purves KL, et al. Classical Human Leukocyte Antigen Alleles and C4 Haplotypes Are Not Significantly Associated With Depression. Biol Psychiatry. 2019.

44.   Niedzwiecki A, & Rath, M. . Peptide vaccines directed against human Metalloproteinases (MMPs) with anti-tumor efficacy in vitro and in vivo. (2018).

45.   Dosset M, Godet Y, Vauchy C, Beziaud L, Lone YC, Sedlik C, et al. Universal cancer peptide-based therapeutic vaccine breaks tolerance against telomerase and eradicates established tumor. Clin Cancer Res. 2012;18(22):6284-95.

46.   Bezu L, Kepp O, Cerrato G, Pol J, Fucikova J, Spisek R, et al. Trial watch: Peptide-based vaccines in anticancer therapy. Oncoimmunology. 2018;7(12):e1511506.