Pollutants in Hong Kong Soil: As, Cd, Cr, Cu, Hg, Pb and Zn

M.K. Chung, K.C. Cheung, and M.H. Wong[*]

Croucher Institute for Environmental Sciences and Department of Biology, Hong Kong Baptist University, Kowloon Tong, Kowloon, Hong Kong SAR, PR China

[*] Corresponding author. E-mail: mhwong@hkbu.edu.hk (M.H. Wong); tel.: +852-3411-7746; fax: +852-3411-7743

Pollutant in Hong Kong soils series:

1.

Chung, M. K., Hu, R., Cheung, K. C. & Wong, M. H. Pollutants in Hong Kong soils:

Polycyclic aromatic hydrocarbons. Chemosphere 67, 464–473 (2007).

2.

Chung, M. K., Cheung, K. C. & Wong, M. H. Pollutants in Hong Kong Soil: As, Cd,

Cr, Cu, Hg, Pb and Zn

https://doi.org/10.1101/2020.02.16.951558

3.

Chung, M. K., Hu, R., Cheung, K. C. & Wong, M. H. Pollutants in Hong Kong Soils:

Organochlorine Pesticides and Polychlorinated Biphenyls

https://doi.org/10.1101/2020.02.16.951541

Current first author contact:

Ming-Kei CHUNG; mingk@alumni.cuhk.net

Abstract

Six heavy metals (Hg, Cu, Cd, Cr, Pb, Zn) and 1 metalloid (As) in surface soils of Hong

Kong were investigated in 10 land use categories (urban park, greening area, country

park, rural area, restored landfill, agricultural farmland, orchard farm, crematorium,

industrial and near highway area). Edaphic Hg concentration in Hong Kong was firstly

reported here.   Clustering of land uses was observed based on total pollutants

concentrations (sum of 7 metals). The most polluted cluster consisted of industrial and

highway areas (median: 617 to 833 mg kg$_{-1}$) and the runner-up cluster included urban

park, greening area and restored landfill (median: 400 to 500 mg kg$_{-1}$). However, this

general finding was not observed for Hg, where higher concentration was found in

agricultural farmland (median 109 µg kg$_{-1}$). The use of low quality fertilizers, together

with the contribution from exhausts and wearable parts from automobiles were believed

to be the major sources of Cr, Cu and Zn in Hong Kong, while the application of Hg-

containing agrochemicals maybe the main mechanism of Hg contamination in

agricultural soil. Based on the daily intake assumption of 0.2 g d$_{-1}$ of soil particles by

USEPA, direct ingestion of Hg-containing soils is not a major exposure pathway for

population in Hong Kong. When comparing the edaphic heavy metal concentrations

with Dutch soil quality guidelines demonstrated that Hg, Cd and Pb were not in level

of health concerns, while Cu, Cr and Zn in less than 6% of total samples were found to

exceed the Dutch intervention values sporadically. In contrast, suburban soils from

northern and northeastern Hong Kong were mostly contaminated with As (10% of total

samples) at concentration that could be potentially causing adverse health impacts to

the nearby population.

72   Keywords: Urban soils, South China, Heavy Metals, Contamination, Pollution

1. Introduction

Environmental contamination of heavy metals has been reported world-wide, and excessive exposure to toxic metals has been commonly known to be hazardous to human health (Agency for Toxic Substances and Disease Registry, 2006). Among these toxic metals, Hg is of particular in concern as it is characterized by high vapour pressure. This unique feature makes it ubiquitous in the environment and become a global pollutant. In addition, its toxicity to human, especially childbearing women, also highlighted the concern from government agency to work on Hg reduction (Srivastava *et al.*, 2006). In countries under the European Union, Hg is classified as a dangerous chemical because of its mobility, volatility and its bioaccumulative properties within organisms and along the food chains (Mukherjee *et al.*, 2004).

A recent review on Hg contamination in China (Zhang and Wong, 2006) revealed that soil Hg contents in most cities (70 to 700 μg kg^{-1}, 12 out of 14 cities reviewed) exceeded the background edaphic Hg value in China (65 μg kg^{-1}) (State Environmental Protection Administration of China, 1990). Enrichment of Hg and other metals was observed in agricultural crop soils (Wong *et al.*, 2002) and sediments (Cheung *et al.*, 2003) around the Pearl River Delta. This led to higher concentrations of all the toxic metals found in bivalves and freshwater fish collected (including from the field and available in markets) within the region (Fang *et al.*, 2001, 2003; Kong *et al.*, 2005; Zhou and Wong, 2000), including seafood in Hong Kong (Tam and Mok, 1991).

In Hong Kong, it has been observed that Hg was bioaccumulated in cetaceans and Indo-Pacific hump-backed dolphins (*Sousa chinensis*) (Parsons, 1998, 1999). It was noted that the mean value of Hg in adult human hair was 3.3 μg g^{-1} which was higher the

mean value for US counterpart (1.5 µg g-1) (Dickman and Leung, 1998), also the

recommended limit for Hg in hair set by USEPA is 1 µg g-1 (Gallagher, 2006). It was

indicated that the elevated Hg levels were linked to subfertility in Hong Kong males

(Dickman *et al.*, 1998). In addition, higher Hg levels in blood and hair of children in

Hong Kong were also observed to be correlated with the frequency of fish consumption

(Ip *et al.*, 2004).

Soils can act as both sinks, via dry atmospheric deposition; as well sources of tosic

metals, via re-emission of semi-volatile pollutants and wind-blown of contaminated

soil materials. By direct contact and inhalation of soil particles, toxic metals would

pose risks to human health. In addition, leaching is also one of the important pathways

to transfer toxic metals to water bodies and therefore accumulate in various aquatic

organisms. Ultimately the toxic metals can enter human via consumption of food such

as crops and fish, and thus understanding the levels and potential sources of toxic metals

are essential to secure public health. Modeling of exposure mechanisms such as dermal

contact and inhalation of dust of soil pollutants for risk assessment requires intensive

data (U.S. Environmental Protection Agency, 1996c), and there is a lack of data on

toxic metals especially Hg in Hong Kong soils. A more comprehensive survey of toxic

metals (except Hg) in Hong Kong soils was conducted almost 10 years ago (Chen *et*

al., 1997), and therefore there seems to be a need to provide update information for all

the toxic metals contained in soils.

The present study was aimed to address the concerns mentioned above by providing

current status of heavy metal and metalloid concentrations (Hg, As, Cu, Cd, Cr, Pb and

Zn) in Hong Kong soils, with a special focus on Hg. To our knowledge, this is the first

report on edaphic Hg levels in Hong Kong and the nearby Pearl River Delta. Results

are valuable as they partially filled the lacking edaphic metals (especially Hg)

information in the Delta and can act as reference for other studies in the region where

contamination of Hg is in raising concern by the public. Potential sources of these toxic

metals are also discussed.

2. Materials and Methods

1. Sampling and Analysis

The sampling was based on 10 different land uses in Hong Kong: urban park, country

park, rural area, restored landfill, agricultural farmland, orchard farm, crematorium,

industrial area and nearby highway. All together there were 138 composite soil samples

that taken from the depth of 0 to 5 cm from surface by a stainless steel soil core.

Samples were stored in plastic bags and subsequently air-dried for 2 weeks and sieved

through a 2-mm mesh.

Chemical analyses of metal contents in soils were based on standard method. 0.25 g of

soil sample was mixed with 9 mL nitric acid, 3 mL hydrofluoric acid and 1 mL

hydrochloric acid and subjected to microwave-assisted acid digestion (USEPA 3052)

(U.S. Environmental Protection Agency, 1996a). The solutions were then filtered

through Advantec 5C filter paper, diluted and made up with deionized water in a 50-ml

plastic volumetric flask. Concentrations of As, Cu, Cd, Cr, Pb, and Zn were determined

by inductively coupled plasma - optical emission spectrometry (ICP-OES) (Perkin-

Elmer Optima 3000 DV), while Hg was quantified by Flow Injection Mercury System

(FIMS) (Perkin-Elmer FIMS-400) based on the cold-vapor atomic absorption

spectrometry (CVAA) (U.S. Environmental Protection Agency, 1996b). Limit of

detection (LOD) for Hg was 0.5 µg kg-1, while 50 µg kg-1 for Cd, Cr, Cu and Zn, and

100 µg kg-1 for As and Pb.

2. Quality assurance and Data analysis

Standard Reference Material (SRM) 2711 was obtained from National Institute of

Standards and Technology (NIST, USA). An analytical blank and the SRM were

included in every batch of microwave acid digestion to assess the recoveries and

performance of extraction.

Mean individual recoveries were: $83 \pm 2\%$ (Hg), $102 \pm 5\%$ (As), $88 \pm 1\%$ (Cd), $97 \pm$

1% (Cu), $107 \pm 4\%$ (Pb) and $89 \pm 2\%$ (Zn). On average, the recoveries of all the

investigated elements in SRM were all $> 94\%$. Statistical analyses including

descriptive statistics, correlation analysis, and PCA analysis were conducted with

Statistica (version 6.0 from StatSoft). Not detected values were substituted with half

of lowest limit of detection (LOD) only for descriptive statistics.

3. Results and Discussion

1. Concentration of pollutants in Hong Kong

Kriged maps were constructed to show the spatial distribution of investigated pollutants

(Figure 1), and it is observed that most of the hotspots for pollutants were found in the

northern part of Hong Kong. In addition, clustering of soil pollutant concentrations in

land uses was also observed (Table 1). Total heavy metal concentrations were highest

in industrial area and area nearby highway (median 617 and 833 mg kg-1); while similar

for urban park, greening area and restored landfill (median 400 to 500 mg kg-1), and the

rest of the land uses are least contaminated (median 200 to 350 mg kg-1). The general

findings that soils in industrial area and adjacent to highways were most contaminated

can also be observed when considering the pollutant individually, but excluding Hg.

Variations in pollutant concentration were usually greatest in urban park, which

spanned up to 3 orders of magnitude, and large variations were found in most of the

land uses, which reflects the heterogeneity of pollutants concentrations is under the

strong influence of local activities or pollution sources.

Hg concentration was ranged from non detectable (N.D.) to 3790 µg kg^{-1}, and the 10

most contaminated soil samples were found in urban parks, greening areas and farms

(29 and 3790 µg kg^{-1}). The 5 locations with highest Hg levels were: urban parks in

Kwun Tong (633 µg kg^{-1}), Central (985 µg kg^{-1}) and Tuen Mun (3785 µg kg^{-1}) and

agricultural farm in Sha Tin (762 µg kg^{-1}) and Tai Po (2196 µg kg^{-1}). The usual contents

of Hg in soils are in the range of 0.01 to 0.03 µg kg^{-1} (Senesi *et al.*, 1999). For

contaminated areas such as Hg mine, Hg concentrations in soils were a thousand folds

more (Loredo *et al.*, 1999). Median Hg levels in urban soil in Korea and Norway were

45 and 130 µg kg^{-1} respectively (Kim and Kim, 1999; Reimann and Caritat, 1998). The

mean and median Hg concentrations in Hong Kong were 135 and 70.5 µg kg^{-1}

respectively, which were broadly in line with the Hg concentrations observed in major

cities in China (Beijing: 509 µg kg^{-1}, Chongqing: 319 µg kg^{-1}, Wuhan: 314 µg kg^{-1})

(Liu *et al.*, 1998; Wang, 2001; Wang *et al.*, 2005). In addition, the concentration ranges

of Cd (N.D. to 4.11 mg kg^{-1}), Cr (N.D. to 2500 mg kg^{-1}), and Pb (11 to 490 mg kg^{-1}) in

the present study (Table 1) were similar to those reported in Shenyang, Beijing, Nanjing

and Xi'an, China (Fang *et al.*, 2004; Wang *et al.*, 2001). However, the range of As

(N.D. to 336 mg kg^{-1}) was generally higher for an order of magnitude when compared

with those reported in major cities in China (Wang *et al.*, 2001). This implied that there

are significant sources of As in Hong Kong that are absent from the aforementioned

cities. Mean concentrations of Cu (37.2 mg kg-1) and Zn (276 mg kg-1) in Hong Kong

were closed to those found in Nanjing (Cu: 40.4 mg kg-1, Zn: 280 mg kg-1) (Wu *et al.*,

2003), but higher than those reported in Guangzhou (Cu: 9.62 mg kg-1, Zn: 115.4 mg

kg-1) (Guan *et al.*, 2001).

2. Statistical analyses among pollutants and their potential sources

The correlations among Cu, Cr and Zn were also identified by principal factor 1 (PC 1)

in the PCA plots shown in Figure 2a. Cadmium was excluded for PCA because of a

large set of not-detected value. PC 1 was able to explain 56% of the variance while PC

2 explained 16%. Together they extracted 72% of the total variance from the present

study. However, PC2 represented an antagonistic relationship between Hg and As.

Figure 2b shows the projection of sampling points to the factor plane. Samples from 2

different urban parks in New Territories contained very high level of metals (Cu, Cr

and Zn) and Hg. Certain agricultural farms and soils from rural and adjacent to highway

in the New Territories were best explained by PC2, implied that they are either high in

Hg or As concentration. Arsenic level was reported to be higher in industrial and heavy

traffic sites (Deb *et al.*, 2002), and the present study also indicated higher level of

edaphic As in the vicinity of highways.

Arsenic has both natural and anthropogenic sources, and their anthropogenic origins

included agrochemicals such as herbicides and pesticides (in form of monosodium

methanearsonate), and wood preservative (arsenic trioxide) (USGS, 2006). Since high

levels of As (>100 mg kg-1) were found chiefly in soil samples collected nearby

highways, it is expected that automobiles could be the major contributor of As through

burning of fossil fuel and wearing of the As-containing babbitt bearings. Many of the

soil samples with As ranged from 30 to 100 mg kg^{-1} were collected in the northern rural

part of Hong Kong, in which historical use of agrochemicals containing As would be

the main contributor of As in soils.

Higher Hg levels were observed in soils of parks and farms with plantation (Table 1).

Similar to As, the origin of Hg can be both anthropogenic and natural, such as ore

mining and forest fire. Certain fertilizers, pesticides and fungicides are known to

contain Hg (Matthews *et al.*, 1995; Nakagawa and Hiromoto, 1997). Therefore, the Hg

concentrations found in farmlands, orchard farms and urban parks maybe due to the use

of agrochemicals. The world-wide average of Hg content in coal is 0.1 ± 0.01 mg kg^{-1},

whereas coals from southern China were enriched in Hg by 1 to 2 orders of magnitude

(Yudovich and Ketris, 2005). Due to the fact that Hong Kong is located at the southern

tip of PRD, which is known for its electrical and electronic manufacturing industry, the

high power demand and the associated emission of Hg is likely to create a regional Hg

problem (Wang *et al.*, 2006), and hence contributes to Hg level in Hong Kong soils.

Higher levels of Hg in human hair leading to subfertility in males have been suspected

to link with higher rates of fish consumption in Hong Kong (Dickman *et al.*, 1998;

Dickman and Leung, 1998). The average consumption rate of seafood is about 60 kg

yr^{-1} person^{-1}, which is equivalent to 167 g d^{-1} person^{-1}, and the mean Hg levels in marine

and freshwater fish available in markets were 120 and 80 µg kg^{-1} respectively (Dickman

and Leung, 1998). The present study showed that the average Hg level of 135 µg kg^{-1}

in Hong Kong soils is slightly higher than those reported in fish. However, the assumed

ingestion amount of soil particles for children (15 kg in weight) is 0.2 g d^{-1} when

calculating the soil screening level for residential exposure of soil pollutants (U.S.

Environmental Protection Agency, 1996c), which is 833 folds less than the average

intake rate of fish (167 g d-1). It is therefore believed that the direct ingestion of Hg-

contaminated soils is not a major health concern in Hong Kong.

Table 2 shows the most prominent correlations were Cu-Cr, Cu-Pb and Cu-Zn, which

were found in 5 out of 10 different land uses. Soil samples collected adjacent to

highway, country park, agricultural farmland, crematorium and industrial area did not

show more than 2 significant correlations. It is a common practice to use compost as

soil conditioner in urbanized areas. In Hong Kong, the sources for composting are

largely derived from livestock wastes from pig and poultry farms under the free

livestock waste collection service provided by the government. Pig manure contains

high levels of Cu and Zn (Bowland, 1990) as common additives in pig feed to increase

the feed conversion efficiency and economic returns (Jin *et al.*, 1995). Cases of

excessive addition of Cu and Zn in feeds were noted (Kessler *et al.*, 1994), and

considerable amounts of these metals were also reported in local composts (Wong,

1990). Apart from compost, fertilizers are also known to contain As, Cd, Cr, Pb and

Zn (Guan *et al.*, 2001; Renner, 2004). Land application of sewage sludge was also

reported to be the principle sources of heavy metals, especially Cd and As (Chu and

Wong, 1984; Elinder, 1985), but this possibility can be ruled out since sludge is

commonly dumped in domestic landfills in Hong Kong. In England and Wales soil,

greatest inputs of Zn and Cu were from animal manure and greatest inputs of Cr were

from industrial wastes (McGrath, 2000). The significant correlations of Cu-Cr, Cu-Pb

and Cu-Zn implied that they are derived from the same sources, and the most likely

source of these metals in Hong Kong soil is from low quality fertilizers, because of its

ease of application and more stable quality than compost. In addition, Zn and Cu are

pollutants associated with automobiles (Viklander, 1997). Approximately 3% of ZnO

is commonly added to the tyres of vehicles as a vulcanization agent and the wear of

tyres can be a significant source of Zn in urban areas (Friedlander, 1973). Other heavy

metal compounds including Cu, Cd and Pb (~0.002%, <0.001% and <0.005%

respectively) are identified in tyres (UNEP, 2000). Other wearable parts of vehicles

such as brake and brake lining also contained high contents of Cu, Pb and Zn (80 to 24

000 mg kg$_{-1}$) (Westerlund, 2001) and therefore contribute a significant portion of heavy

metals in soils.

Lead pollution in cities was commonly recognized as one of the major pollutants caused

by vehicle emissions (Yang *et al.*, 2000). Hong Kong government introduced unleaded

petrol (ULP) in 1991 and banned the supply, sale and dispensing of leaded petrol as

well as any fuel additives containing Pb in 1999 (Hong Kong Environmental Protection

Department, 1999), resulting in a decline of Pb concentration in street dust of Hong

Kong from 1300 ± 1400 mg kg$_{-1}$ (Yim and Nau, 1987) to 180 ± 93 mg kg$_{-1}$ (Li *et al.*,

2001).

Atmospheric deposition from nearby regions also represents another important input of

heavy metals such as Cr, Cu, Pb and Zn to surface soils. According to a quality

monitoring program in China (General Administration of Quality Supervision

Inspection and Quarantine of the People's Republic of China, 2004), only about 70%

of the unleaded petrol samples in China was found to comply with the national standard.

In some cases, Pb level was exceeded more than 200 times to the standards. Study on

atmospheric deposition in the PRD revealed that the deposition of Cr, Cu, Pb and Zn

(6.43 ± 3.19, 18.6 ± 7.88, 12.7 ± 6.72 and 104 ± 36.4 mg m$_{-1}$ yr$_{-1}$) was significantly

higher when compared with Europe and North America (Wong *et al.*, 2003). Long-

range transport of air-borne pollutants or wind-blown contaminated soil particles from

Mainland China by the northeast monsoon was reported (Lee and Hills, 2003).

Moreover, atmospheric input was reported to be the major contributor of Pb, Cd, As

and Hg in agricultural soils in England and Wales (Nicholson *et al.*, 2006), and thus

atmospheric deposition, either locally or regionally, may play a significant role for the

presence of particle-bound pollutants in soils.

3. Comparison of soils cleanup criteria

The soil quality guideline values on the various pollutants investigated imposed by

Netherlands (Dutch Guidelines) (Ministry of Housing Spatial Planning and

Environment, 2000), Sweden (Soil Remediation Goals) (Swedish Environmental

Protection Agency, 2002), England (Kelly Indices) (Contaminated Land Assessment &

Remediation Research Centre, 2004) and China (Environmental Quality Standard)

(State Environmental Protection Administration of China, 1995) are summarized in

Table 3. As Dutch guideline is the most comprehensive and commonly used,

comparisons the present findings are made chiefly with the Dutch values. Mercury, Cd

and Pb in 138 samples were below the Dutch intervention values, suggesting that the

concentrations of these metals in soils were not hazardous to human. In terms of As,

Cu, Cr and Zn, most of their levels did not exceed the Dutch intervention values, but

there were sporadic soil samples containing levels of As, Cu, Cr and Zn exceeded the

intervention values (14, 3, 2, and 8 out of 138 samples correspondingly). Nine out of

10 suburban samples from the northern and northeastern New Territories contained As

concentrations greater than the Dutch intervention value. Thus, it is suspected that a

large part of northern and northeastern in Hong Kong is contaminated with As at

concentration that can impose adverse effects on human health. Soil samples with Zn

concentration greater than the intervention value were mainly noted in industrial areas,

and this is also true for Cu and Cr. For the scarcity and remoteness of the hotspots (Cr,

Cu, and Zn), their potential adverse impacts to the general public were kept to

minimum. In England, soil remediation is often required before further development

on brownfield soils as they are typically contaminated with high levels of heavy metals

due to past industrial activities (French *et al.*, 2006). Although leaching of toxic metals

to underground water is not a major concern as Hong Kong relies mainly on river water

transported from the mainland as well as rainwater collected locally, the fact that more

than 10% of the soil samples were highly contaminated with As warrants further

investigation. This is especially true if any of these sites (north and north-east of the

New Territories) are used for residential development in the future.

4. Conclusions

In terms of total concentrations of all the metals (metalloids), industrial and highway

areas were the most contaminated. This finding is also true for individual elements

(As, Cd, Cr, Cu, Pb and Zn) other than Hg. It was found that Hg concentration was the

highest in soil collected from agricultural farmland, which could be attributed to the

application of Hg-containing agrochemicals. The use of low quality fertilizers is also

believed to be the main source of As, Cu and Zn, while substantial contributions of

pollutants by exhausts and wearable parts from automobiles are also suspected.

Atmospheric deposition from local and nearby regions is also believed to be a major

source of edaphic metals in Hong Kong. A reduction of Pb in soils during the past 10

to 15 year was chiefly due to the use of Pb-free petrol. It is expected all the metals

(metalloids) would not cause any potential health impacts to the general public, except

As, due to its high concentrations in northern and northeastern suburbs in Hong Kong.

Acknowledgments

The authors are grateful to Mr. Y.Y. Chin from the Leisure and Cultural Services

Department (LCSD, HKSAR) for providing technical assistance. Financial support

from the Strategic Research Fund, Science Faculty, HKBU and the Area of Excellence

(AoE) Scheme (CITYU/AoE/03-04/02) under the University Grants Committee of

Hong Kong is gratefully acknowledged.

 Figure Captions:

Figure 1. Kriged maps of pollutants (As, Cd, Cr, Cu, Hg, Pb, and Zn) concentrations

(mg kg $_{-1}$) in surface soils of Hong Kong.

Figure 2. Plots with PC 1 and PC 2 from principal component analysis as X and Y

axis on various pollutants and sampling points. PC 1 was able to explain 49% while

PC 2 accounted for another 19% of the total variance. Note that Cd were excluded

from PCA because of a large set of not-detected value. A: Biplot showing the loading

of 6 pollutants on PC 1 and PC 2. B: Scatter plot of sampling points projecting on the

PC 1 and PC 2 plane.

400   Table 1. Mean, median, and range of concentration of studied pollutants (mg kg-1) in Hong Kong soils. Note that concentration unit of Hg µg kg-1.

Classified soil categories	Sample no.	Hg			As			Cu			Cd		
		Mean	Median	Range	Mean	Median	Range	Mean	Median	Range	Mean	Median	Range
		Concentration in soils (µg/kg)						Concentration in soils (mg/kg)					
Urban park	39	232	75.0	[N.D. - 3785]	19.4	16.5	[N.D. - 122]	81.41	13.0	[N.D. - 2129]	0.56	0.01	[N.D. - 2.93]
Greening area	14	104	66.0	[N.D. - 311]	26.6	26.6	[N.D. - 97]	21.46	15.1	[N.D. - 59]	0.50	0.01	[N.D. - 2.25]
Country park	9	52	25.0	[N.D. - 128]	36.1	20.3	[3.6 - 95]	11.0	0.01	[N.D. - 93]	0.36	0.01	[N.D. - 2.42]
Rural area	19	75	81.0	[N.D. - 202]	27.6	20.5	[3.2 - 143]	7.32	6.08	[N.D. - 28]	0.37	0.01	[N.D. - 2.91]
Restored landfill	11	50	25.0	[N.D. - 119]	22.0	22.8	[N.D. - 47]	19.3	11.0	[N.D. - 62]	0.85	0.01	[N.D. - 2.86]
Agricultural farmland	9	402	109	[N.D. - 2196]	27.2	15.2	[N.D. - 109]	12.0	6.61	[N.D. - 66]	0.14	0.01	[N.D. - 0.77]
Orchard farm	5	154	72.0	[N.D. - 555]	34.3	30.4	[7.3 - 93]	8.86	5.60	[N.D. - 27]	0.56	0.01	[N.D. - 1.6]
Crematorium	10	75	78.0	[N.D. - 146]	13.8	11.5	[N.D. - 50]	4.29	0.61	[N.D. - 16]	0.13	0.01	[N.D. - 1.21]
Industrial area	18	74	73.0	[N.D. - 237]	25.5	25.4	[N.D. - 61]	52.34	33.3	[N.D. - 396]	1.19	0.67	[N.D. - 4.11]
Nearby highway	4	134	117	[N.D. - 277]	174.6	141	[80.5 - 336]	18.75	15.4	[3.62 - 41]	0.01	0.01	[N.D. - N.D.]

Classified soil categories	Sample no.	Cr			Pb			Zn			Total pollutants		
		Mean	Median	Range	Mean	Median	Range	Mean	Median	Range	Mean	Median	Range
		Concentration in soils (mg/kg)											
Urban park	39	50.5	21.9	[1.85 - 601]	141	130	[11 - 305]	293	197	[13 - 3508]	586	401	[29 - 6553]
Greening area	14	20.8	18.2	[1.23 - 44]	142	144	[104 - 188]	367	300	[114 - 1182]	578	540	[300 - 1351]
Country park	9	7.18	2.00	[N.D. - 23]	75	68	[56 - 115]	75	85	[38 - 110]	204	204	[101 - 357]
Rural area	19	17.0	12.1	[0.81 - 57]	136	124	[53 - 244]	183	135	[43 - 555]	371	377	[155 - 763]
Restored landfill	11	25.1	24.9	[11.8 - 37]	189	180	[47 - 393]	170	146	[79 - 347]	426	425	[221 - 680]
Agricultural farmland	9	22.7	17.9	[7.36 - 54]	121	120	[79 - 161]	197	152	[63 - 564]	380	303	[234 - 845]
Orchard farm	5	16.3	12.4	[5.73 - 34]	104	91	[64 - 186]	116	95	[45 - 243]	280	236	[145 - 479]
Crematorium	10	27.3	21.1	[11.5 - 88]	118	96	[65 - 277]	140	126	[89 - 221]	304	263	[180 - 515]
Industrial area	18	167	29.1	[11.3 - 2486]	221	183	[92 - 493]	529	298	[75 - 1631]	996	617	[214 - 3033]
Nearby highway	4	35.5	35.4	[15.7 - 55]	147	134	[115 - 205]	556	349	[158 - 1367]	932	833	[409 - 1653]

Table 2. Correlation matrix of investigated metals in different land uses in Hong Kong.

Urban park

Greening area	Hg	As	Cu	Cd	Cr	Pb	Zn
Hg		-0.06	0.04	0.2	0.01	0.12	0.03
As	0.46		-0.04	0.12	-0.07	-0.16	-0.76
Cu	0.13	0.54*		-0.09	0.87*	0.47*	0.98*
Cd	0.14	0.47	0.53*		-0.17	0.15	-0.08
Cr	0.14	0.46	0.80*	0.26		0.49*	0.82*
Pb	-0.4	-0.16	0.53	0.19	0.35		0.54*
Zn	-0.08	-0.2	0.1	-0.23	0.33	0.13	

Agricultural farmland

Orchard farm	Hg	As	Cu	Cd	Cr	Pb	Zn
Hg		-0.2	0.16	-0.23	0.13	0.25	0.08
As	-0.61		-0.15	0.36	-0.45	-0.55	-0.43
Cu	-0.19	0.83		-0.21	0.62	0.53	0.86*
Cd	0.4	0.84	0.73		-0.2	-0.65	-0.18
Cr	-0.14	0.15	0.66	0.15		0.48	0.90*
Pb	-0.1	-0.35	0.23	-0.27	0.87		0.6
Zn	-0.17	0.82	0.99*	0.74	0.62	0.2	

Country park

Rural area	Hg	As	Cu	Cd	Cr	Pb	Zn
Hg		-0.32	-0.21	-0.51	0.67*	0.52	0.18
As	0.31		0.3	0.44	0.03	0.09	0.18
Cu	0.41	0.49*		0.01	0.43	0.51	0.1
Cd	0.02	0.70*	0.43		-0.23	-0.31	0.47
Cr	0.37	0.4	0.27	-0.03		0.79*	0.6
Pb	-0.16	-0.22	-0.08	-0.28	-0.1		0.29
Zn	0.48*	-0.03	0.12	-0.2	0.27	0.14	

Crematorium

Industrial area	Hg	As	Cu	Cd	Cr	Pb	Zn
Hg		-0.08	0.25	0.18	0.33	-0.16	0.07
As	0.05		0.09	0.85*	0.1	0.14	-0.04
Cu	0.78*	-0.08		-0.18	0.82*	0.13	0.72*
Cd	0.17	0.08	-0.19		-0.05	-0.16	-0.35
Cr	0.07	-0.42	0.02	0.22		-0.05	0.53
Pb	-0.26	-0.14	0.14	-0.11	-0.15		0.61
Zn	0.2	-0.31	0.43	-0.21	-0.06	0.4	

Restored landfill

Nearby highway	Hg	As	Cu	Cd	Cr	Pb	Zn
Hg		0.07	-0.36	-0.62*	-0.21	-0.23	0.39
As	-0.76		-0.43	-0.09	-0.79*	-0.54	-0.4
Cu	-0.68	0.77		0.27	0.80*	0.61*	0.51
Cd	Nil	Nil	Nil		0.35	0.39	-0.55
Cr	-0.84	0.75	0.96*	Nil		0.80*	0.47
Pb	-0.67	0.84	0.99*	Nil	0.92		0.19
Zn	0.07	-0.42	0.25	Nil	0.27	0.11	

Pearson correlation coefficients were shown. Values with * indicated that significant correlations were found at p=0.05.

405
406
407
408 Table 3. Soil quality guidelines from Netherlands, Sweden, United Kingdom and China, and their recommended values for Hg, As, Cu, Cd, Cr, Pb and
409 Zn in soils.

Country of implementation	Quality guidelines	Hg	As	Cu	Cd	Cr	Pb	Zn
		Concentration in soils (mg/kg)						
Netherlands	Dutch target value	0.3	29	36	0.8	100	85	140
	Dutch intervention value	10	55	190	12	380	530	720
Sweden	Soil remediation goals for sensitive land use	N.A.	15	100	0.4	120	80	350
	Soil remediation goals for less sensitive land use	N.A.	40	200	12	250	300	700
United Kingdom	Kelly	1	30	N.A.	1	100	500	N.A.
China	Environmental quality standard	0.5	30	200	0.6	300	300	250

Figure 1. Kriged maps of pollutants (As, Cd, Cr, Cu, Hg, Pb, and Zn) concentrations
(mg kg $^{-1}$) in surface soils of Hong Kong.

Figure 2. Plots with PC 1 and PC 2 from principal component analysis as X and Y axis on various pollutants and sampling points. PC 1 was able to explain 49% while PC 2 accounted for another 19% of the total variance. Note that Cd were excluded from PCA because of a large set of not-detected value. A: Biplot showing the loading of 6 pollutants on PC 1 and PC 2. B: Scatter plot of sampling points projecting on the PC 1 and PC 2 plane.

References

Agency for Toxic Substances and Disease Registry. (2006). Toxicological Profile
Information Sheet Retrieved August 21, 2006, from Agency for Toxic Substances
and Disease Registry Web site: http://www.atsdr.cdc.gov/toxpro2.html

Bowland, J.P. (1990). Copper as a performance promoter for pigs. Pig News Inf., 11(2),
163-167.

Chen, T.B., Wong, J.W., Zhou, H.Y., Wong, M.H. (1997). Assessment of trace metal
distribution and contamination in surface soils of Hong Kong. Environ. Pollut.,
96(1), 61-68.

Cheung, K.C., Poon, B.H., Lan, C.Y., Wong, M.H. (2003). Assessment of metal and
nutrient concentrations in river water and sediment collected from the cities in the
Pearl River Delta, South China. Chemosphere, 52(9), 1431-1440.

Chu, L.M., Wong, M.H. (1984). Application of refuse compost: Yield and metal uptake
of three different food crops. Resour. Conserv. Recycl., 7(2-4), 221-234.

Contaminated Land Assessment & Remediation Research Centre. (2004). Kelly
indices: Guidelines for classification of contaminated soils Retrieved December 2,
2005, from Contaminated Land Assessment & Remediation Research Centre Web
site:
http://www.clarrc.ed.ac.uk/index.php?option=com_docman&task=docclick&Item
id=57&bid=19&limitstart=0&limit=20

Deb, M.K., Thakur, M., Mishra, R.K., Bodhankar, N. (2002). Assessment of
atmospheric arsenic level in airborne dust particulates of an urban city of central
India. Water Air Soil Pollut., 140(1-4), 57-71.

Dickman, M.D., Leung, C.K., Leong, M.K. (1998). Hong Kong male subfertility links

to mercury in human hair and fish. Sci. Total Environ., 214, 165-174.

Dickman, M.D., Leung, K.M.C. (1998). Mercury and organochlorine exposure from

fish consumption in Hong Kong. Chemosphere, 37(5), 991-1015.

Elinder, C. (1985). Cadmium: Uses, occurrence and intake. In Friberg, L., Elinder,

C.G., Kjellstrom, T. (Eds.), Cadmium and health: A toxicological and

epidemiological appraisal. Volume I. Exposure, dose, and metabolism. Effects and

response (pp. 23-64). Boca Raton, FL: CRC Press.

Fang, F., Wang, Q., Li, J. (2004). Urban environmental mercury in Changchun, a

metropolitan city in Northeastern China: Source, cycle, and fate. Sci. Total

Environ., 330(1-3), 159-170.

Fang, Z.Q., Cheung, R.Y., Wong, M.H. (2001). Heavy metal concentrations in edible

bivalves and gastropods available in major markets of the Pearl River Delta. J.

Environ. Sci. (China), 13(2), 210-217.

Fang, Z.Q., Cheung, R.Y., Wong, M.H. (2003). Heavy metals in oysters, mussels and

clams collected from coastal sites along the Pearl River Delta, South China. J.

Environ. Sci. (China), 15(1), 9-24.

French, C.J., Dickinson, N.M., Putwain, P.D. (2006). Woody biomass

phytoremediation of contaminated brownfield land. Environ. Pollut., 141(3), 387-

395.

Friedlander, S.K. (1973). Chemical element balances and identification of air pollution

sources. Environ. Sci. Technol., 7(3), 235-240.

Gallagher, B. (2006). EQI highlights mercury threat Retrieved March 21, 2006, from

University   of   North   Carolina   Web   site:

http://www.unca.edu/banner/060223/features.html

General Administration of Quality Supervision Inspection and Quarantine of the

People's Republic of China. (2004). Retrieved April 4, 2004, from General

Administration of Quality Supervision, Inspection and Quarantine of the People's

Republic of China Web site: http://www.aqsiq.gov.cn

Guan, D., Chen, Y., Yuan, G. (2001). Study on heavy metal concentrations and the

impact of human activity on them in urban and suburb soils of Guangzhou. Acta

Sci. Nat. Univ. Sunyatseni., 40(4), 93-101.

Hong Kong Environmental Protection Department. (1999). Ban on leaded petrol

gazetted Retrieved February 13, 2004, from Press release Web site:

http://www.epd.gov.hk/epd/english/news_events/press/press_990205.html

Ip, P., Wong, V., Ho, M., Lee, J., Wong, W. (2004). Environmental mercury exposure

in children: South China's experience. Pediatr. Int., 46(6), 715-721.

Jin, R.B., Cui, H.M., Wang, S.X., Mao, J.D., Zhang, M., Chu, X.S. (1995). Experiment

on feeding amino acid chelated copper to growing-finishing pigs. J. Jiangsu Agric.

Coll., 16(3), 47-50.

Kessler, J., Zogg, M., Bachler, E. (1994). Phosphorus, copper and zinc in the pig trough.

Agrarforschung, 1(11-12), 480-483.

Kim, K.H., Kim, S.H. (1999). Heavy metal pollution of agricultural soils in central

regions of Korea. Water Air Soil Pollut., 111(1-4), 109-122.

Kong, K.Y., Cheung, K.C., Wong, C.K., Wong, M.H. (2005). Residues of DDTs, PAHs

and some heavy metals in fish (tilapia) collected from Hong Kong and mainland

China. J. Environ. Sci. Health Part A Toxic-Hazard. Subst. Environ. Eng., 40(11),

2105-2115.

Lee, Y.C., Hills, P.R. (2003). Cool season pollution episodes in Hong Kong, 1996-

2002. Atmos. Environ., 37(21), 2927-2939.

Li, X., Poon, C.S., Pui, S.L. (2001). Heavy metal contamination of urban soils and street

dusts in Hong Kong. Appl. Geochem., 16(11-12), 1361-1368.

Liu, J.H., Wang, W.H., Peng, A. (1998). Pollution and sources of mercury in top soil

in two district of Beijing. Acta. Sci. Circum., 18, 331-3336 (In Chinese).

Loredo, J., Ordonez, A., Gallego, J.R., Baldo, C., Garcia-Iglesias, J. (1999).

Geochemical characterisation of mercury mining spoil heaps in the area of Mieres

(Asturias, northern Spain). J. Geochem. Explor., 67(1-3), 377-390.

Matthews, S., McCracken, I., Lonergan, G. (1995). Mercury contamination of golf

courses due to pesticide use. Bull. Environ. Contam. Toxicol., 55(3), 390-397.

McGrath, S.P. (2000). Risk assessment of metals. In Luo, W., McGrath, S.P. (Eds.),

Proceedings of SoilRem 2000, International Conference of Soil Remediation.

Hangzhou, China.

Ministry of Housing Spatial Planning and Environment, 2000. Circular on target values

and intervention values for soil remediation. Ministry of Housing, Spatial Planning

and Environment, Netherlands.

Mukherjee, A.B., Zevenhoven, R., Brodersen, J., Hylander, L.D., Bhattacharya, P.

(2004). Mercury in waste in the European Union: sources, disposal methods and

risks. Resour. Conserv. Recycl., 42(2), 155-182.

Nakagawa, R., Hiromoto, M. (1997). Geographical distribution and background levels

of total mercury in air in Japan and neighbouring countries. Chemosphere, 34(4),

801-806.

Nicholson, F.A., Smith, S.R., Alloway, B.J., Carlton-Smith, C., Chambers, B.J. (2006).

Quantifying heavy metal inputs to agricultural soils in England and Wales. Wat.

Environ. J., 20(2), 87-95.

Parsons, E.C.M. (1998). Trace metal pollution in Hong Kong: Implications for the

health of Hong Kong's Indo-Pacific hump-backed dolphins (Sousa chinensis). Sci.

Total Environ., 214(1), 175-184.

Parsons, E.C.M. (1999). Trace element concentrations in the tissues of cetaceans from

Hong Kong's territorial waters. Environ. Conserv., 26, 30-40.

Reimann, C., Caritat, P.d. (1998). Chemical elements in the environment: Factsheets

for the geochemist and environmental scientist. New York: Springer.

Renner, R. (2004). Arsenic and lead leach out of popular fertilizer. Environ. Sci.

Technol., 38(20), 382A.

Senesi, G.S., Baldassarre, G., Radina, B., Senesi, N. (1999). Trace element inputs into

soils by anthropogenic activities and implications for human health. Chemosphere,

39(2), 343-377.

Srivastava, R.K., Hutson, N., Martin, B., Princiotta, F., Staudt, J. (2006). Control of

mercury emissions from coal-fired electric utility boilers. Environ. Sci. Technol.,

40(5), 1385-1393.

State environmental Protection Administration of China. (1990). The Background

Levels of Element in Soil in China. Beijing: Chinese Environmental Science Press

(In Chinese).

State Environmental Protection Administration of China, 1995. Environmental quality

standard for soils. State Environmental Protection Administration of China, China.

Swedish Environmental Protection Agency. (2002). Assessment of contamination level

Retrieved December 2, 2005, from Swedish Environmental Protection Agency Web

site: http://www.internat.naturvardsverket.se/index.php3?main=/doc

uments/legal/assess/assedoc/contdoc/pollevl.htm

Tam, S.Y., Mok, C.S. (1991). Metallic contamination in oyster and other seafood in

Hong Kong. Food Addit. Contam., 8(3), 333-342.

U.S. Environmental Protection Agency, 1996a. Method 3052: Microwave assisted acid

digestion of siliceous and organically based matrices SW-846. Test methods for

evaluating solid wastes. Physical/chemical methods. U.S. Environmental

Protection Agency, Washington, DC.

U.S. Environmental Protection Agency, 1996b. Method 7471B: Mercury in solid or

semisolid waste (manual cold-vapor technique). SW-846. Test methods for

evaluating solid wastes. Physical/chemical methods. U.S. Environmental

Protection Agency, Washington, DC.

U.S. Environmental Protection Agency, 1996c. Soil screening guidance: User's guide.

U.S. Environmental Protection Agency, Washington, DC.

UNEP, 2000. Technical guidelines on the identification and management of used tyres.

Basel Convention series/SBC No. 02/10. Secretariat of the Basel Convention,

Chatelaine.

USGS. (2006). Arsenic   Retrieved July 12, 2006, from USGS Web site:

http://minerals.usgs.gov/minerals/pubs/commodity/arsenic/

Viklander, M. (1997). Snow quality in urban areas. Unpublished doctoral dissertation,

Luleå University of Technology, Luleå, Sweden.

Wang, D.Y. (2001). Distribution and behaviour of mercury in terrestrial ecosystem in

acid deposition area. Chongqing, P.R. China: Xinan Agriculture University (In

Chinese).

Wang, J.Y., Tong, Z.D., Yan, J.B. (2005). Study on the relationship between contents

of poison in fishes and the levels of ocean pollutants in Zhoushan fishery. Chin. J.

Epidemiol., 26, 18-21 (In Chinese).

Wang, Q., Dong, Y., Cui, Y., Liu, X. (2001). Instances of soil and crop heavy metal

contamination in China. Soil Sediment Contam., 10(5), 497-511.

Wang, Z., Zhang, X., Chen, Z., Zhang, Y. (2006). Mercury concentrations in size-

fractionated airborne particles at urban and suburban sites in Beijing, China. Atmos.

Environ., 40(12), 2194-2201.

Westerlund, K.G., 2001. Metal emission from Stockholm traffic - wear of brake linings.

The Stockholm Environment and Health Protection Administration, Stockholm,

Sweden.

Wong, M.H. (1990). Anaerobic digestion of pig waste mixed with sewage sludge. Biol.

Waste., 31, 223-230.

Wong, S.C., Li, X.D., Zhang, G., Qi, S.H., Min, Y.S. (2002). Heavy metals in

agricultural soils of the Pearl River Delta, South China. Environ. Pollut., 119(1),

33-44.

Wong, S.C., Li, X.D., Zhang, G., Qi, S.H., Peng, X.Z. (2003). Atmospheric deposition

of heavy metals in the Pearl River Delta, China. Atmos. Environ., 37(6), 767-776.

Wu, X., Li, L., Pan, G., Ju, Y., Jiang, H. (2003). Soil pollution of Cu, Zn, Pb and Cd in

different city zones of Nanjing. Environ. Sci., 24(3), 105-111.

Yang, S.J., Dong, J.Q., Cheng, B.R. (2000). Characteristics of air particulate matter and

their sources in urban and rural area of Beijing, China. J. Environ. Sci., 12(4), 402-

409.

Yim, W.W.S., Nau, P.S. (1987). Distribution of lead, zinc, copper and cadmium in dust

from selected urban areas of Hong Kong. Hong Kong Eng., 15(1), 7-14.

Yudovich, Y.E., Ketris, M.P. (2005). Mercury in coal: A review: Part 1. Geochemistry.

Int. J. Coal Geol., 62(3), 107-134.

Zhang, L., Wong, M.H. (2006). Environmental mercury contamination in China:

Sources and impacts. Environ. Int., In Press, Corrected Proof.

Zhou, H.Y., Wong, M.H. (2000). Mercury accumulation in freshwater fish with

emphasis on the dietary influence. Water Res., 34(17), 4234-4242.

596
597