Adaptation of pancreatic cancer cells to nutrient deprivation is reversible and requires glutamine

synthetase stabilization by mTORC1

Pei-Yun Tsai[1,2], Min-Sik Lee[1,2,3], Unmesh Jadhav[4,5,6,7], Insia Naqvi[1], Shariq Madha[4,5], Ashley Adler[1,3],

Meeta Mistry[8], Caroline A. Lewis[9], Daniel S. Hitchcock[3], Frederick R. Roberts[10], Peter DelNero[1,11],

Thomas Hank[12], Kim C. Honselmann[12], Vicente Morales Oyarvide[4], Mari Mino-Kenudson[13], Clary B.

Clish[3], Ramesh A. Shivdasani[4,5,6,7,14], and Nada Y. Kalaany[1,2,3*]

[1]Division of Endocrinology and Research Center for Metabolic Health, Boston Children's Hospital,

Boston, MA, USA.

[2]Department of Pediatrics, Harvard Medical School, Boston, MA, USA.

[3]Broad Institute of MIT and Harvard, Cambridge, MA, USA.

[4]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA.

[5]Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, MA, USA.

[6]Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA.

[7]Department of Medicine, Harvard Medical School, Boston, MA, USA.

[8]Bioinformatics Core, Harvard T.H. Chan School of Public Health, Boston, MA, USA.

[9]Whitehead Institute for Biomedical Research, Cambridge, MA, USA.

[10]FUJIFILM VisualSonics Inc., Toronto, ON, Canada.

[11]Cancer Prevention Fellowship Program, Division of Cancer Prevention, National Cancer Institute,

Bethesda, MD, USA.

[12]Department of Surgery, Massachusetts General Hospital and Harvard Medical School, Boston, MA,

USA.

[13]Department of Pathology, Massachusetts General Hospital and Harvard Medical School, Boston, MA,

USA.

[14]Harvard Stem Cell Institute, Cambridge, MA, USA.

*email: nada.kalaany@childrens.harvard.edu

Abstract

Pancreatic ductal adenocarcinoma (PDA) is a lethal, therapy-resistant cancer that thrives in a highly

desmoplastic, nutrient-deprived microenvironment. Several studies investigated the effects of depriving

PDA of either glucose or glutamine alone. However, the consequences on PDA growth and metabolism

of limiting both preferred nutrients have remained largely unknown. Here, we report the selection for

clonal human PDA cells that survive and adapt to limiting levels of both glucose and glutamine. We find

that adapted clones exhibit increased growth *in vitro* and enhanced tumor-forming capacity *in vivo*.

Mechanistically, adapted clones share common transcriptional and metabolic programs, including amino

acid use for *de novo* glutamine and nucleotide synthesis. They also display enhanced mTORC1 activity

that prevents the proteasomal degradation of glutamine synthetase (GS), the rate-limiting enzyme for

glutamine synthesis. This phenotype is notably reversible, with PDA cells acquiring alterations in open

chromatin upon adaptation. Silencing of GS suppresses the enhanced growth of adapted cells and

mitigates tumor growth. These findings identify non-genetic adaptations to nutrient deprivation in PDA

and highlight GS as a dependency that could be targeted therapeutically in pancreatic cancer patients.

Introduction

With a steady rise in incidence and a poor 5-year survival rate of ~ 9%, PDA is now the third leading

cause of cancer death in the US[1, 2]. Oncogenic *KRAS* mutations are highly prevalent in PDA (> 90%),

driving many of its distinctive metabolic features[3, 4]. These include the macropinocytic uptake of

extracellular macromolecules, the recycling of intracellular components[5-7], and the re-direction of

glycolytic and glutaminolytic intermediates into anabolic and redox-controlling pathways required for

PDA growth[8, 9]. However, whether these alterations are sufficient to account for the ability of PDA to

thrive in a seemingly hostile *in vivo* microenvironment, where abnormal vasculature and exuberant

stroma restrict nutrient access, remains unknown.

Because nutrient availability largely dictates metabolic behavior[10], the relevance of studying

tumor metabolism within the native environment has been recently underscored[11-15]. Although culture

media composition can be modulated to mimic circulating metabolite levels[16, 17], this modulation may

not accurately reflect the tumor metabolic microenvironment. In particular, levels of glucose and

glutamine, two of the most abundant and tumor-preferred nutrients in the circulation, can be limiting in

PDA tumors, compared to benign adjacent tissue[7], and significantly lower within tumor cores, compared

to the periphery[18, 19]. Paradoxically, both glucose and glutamine are routinely supplemented in culture

media at levels that are significantly higher than the circulation: 11mM glucose and 2mM glutamine in

RPMI compared to ~ 5.5mM glucose and 0.6mM glutamine in serum[20]. Although prior studies focused

on depriving tumor cells of either glucose or glutamine alone[9, 21-24], how PDA cells survive and

proliferate under limiting levels of both major nutrients is not well understood.

To investigate this, we selected for clonal PDA cells that survive and adapt to limiting levels of

both glucose and glutamine. We find that the adapted clones have enhanced proliferation *in vitro* and

tumor-forming capacity *in vivo*. They also share common signaling, transcriptional and metabolic

alterations that are acquired upon adaptation. These include a post-translational role for mTORC1

signaling in the stabilization of glutamine synthetase (GS), and the use of amino acids for the synthesis

of glutamine and nucleotides. Of note, this phenotype is reversible, implicating epigenetic alterations in

the adaptation process, and highlighting GS as a candidate for therapeutic targeting in pancreatic cancer.

Results

PDA cells adapted to low glucose / low glutamine have enhanced growth *in vitro* and *in vivo*

To test whether pancreatic cancer cells can survive and adapt to harsh nutrient-deprived conditions, we

cultured seven human PDA cell lines (AsPC-1, BxPC-3, HPAC, MIA PaCa-2, PANC-1, SUIT-2 and

PA-TU-8988T, herein termed 8988T) in customized low glucose (0.5mM) and low glutamine (0.1mM),

medium (L-L). These modifications represent ~20-fold reduction from standard RPMI medium, herein

referred to as high glucose-high glutamine (H-H) and 5- to 10-fold decrease from human serum levels[20]

(Fig. 1a); both L-L and H-H media were supplemented with 10% dialyzed fetal bovine serum (FBS).

Consistent with the requirement of glucose and glutamine for PDA growth[8, 9], most PDA cells died

within two weeks of culture in L-L medium, despite media replenishment every 2-3 days. However, rare

surviving cells were observed in 2 of the 7 PDA cell lines, SUIT-2 and 8988T. These cells gradually

gave rise to adapted clones, 3 of which (A-C4, C-5 and C-6) were selected per cell line. Adapted clones

were then studied in comparison to non-adapted clones (NA-C1, C-2 and C-3), which were isolated

from the corresponding parental PDA lines grown under H-H conditions (Fig.1a). All clones were

subsequently passaged every 3-4 days for use in experiments.

To assess their *in vitro* growth potential, adapted and non-adapted cells were first subjected to a

7-day proliferation assay performed without media replenishment. Whereas no significant differences

were detected in H-H medium conditions (Supplementary Fig. 1a,b), adapted cells from both lines

showed enhanced proliferative capacity (Fig. 1b) and decreased cell death (Fig. 1c), compared to non-

adapted clones, in L-L medium. In a colony formation assay, adapted 8988T clones generated a larger

number of colonies independent of the medium, whereas adapted SUIT-2 clones tended to form more

colonies in L-L but not in H-H medium. (Fig. 1d and Supplementary Fig. 1c). When plated under 3-

dimensional (3-D) culture conditions, only adapted 8988T cells formed spheres independent of the

medium (Fig. 1e and Supplementary Fig. 1d-f). SUIT-2 clones formed a similar number of spheres,

independent of the medium or adaptation (Supplementary Fig. 1d,e). However, SUIT-2 spheres tended

to be larger in adapted, compared to non-adapted clones, when cultured under L-L, but not H-H

conditions (Fig. 1e and Supplementary Fig. 1f).

Because autophagy and macropinocytosis are well-recognized dependencies in PDA[3, 5, 6], we

assessed their potential contribution to the enhanced proliferation of the adapted clones under nutrient-

limiting conditions. No differential increase in autophagic flux was observed in the adapted cells,

compared to non-adapted controls in L-L medium (Supplementary Fig. 2a,b). Moreover, uptake of

tetramethylrhodamine (TMR)-labeled dextran, an indicator of macropinocytosis, was not further

increased in the adapted, compared to non-adapted cells, under L-L conditions (Supplementary Fig. 2c).

These data indicate that surviving PDA cells that adapt to chronic glucose and glutamine deprivation

acquire a proliferative potential that depends on processes other than autophagy or macropinocytosis.

To test whether the adaptation-mediated *in vitro* phneotype could be translated *in vivo*, adapted

and non-adapted 8988T PDA clonal cells were orthotopically injected into the pancreas of B6.Rag1[-/-]

mice. Within the span of 33 days, only 1 out of 14 non-adapted cell injections derived from 2 distinct

clones formed a tumor (100 mm^3 in volume). In contrast, 14 of 14 adapted clonal cell injections

produced tumors that were larger in size, with average volumes of 600 mm^3 and 250 mm^3 for clones 4

and 5, respectively (Fig. 1f and Supplementary Fig. 1g). These results demonstrate that metabolic

adaptation to a nutrient-deprived environment can endow cancer cells with higher growth fitness,

allowing tumors to survive and thrive during periods of nutrient starvation.

Adapted PDA clones display altered amino acid and nucleotide metabolism coupled with

enhanced mTORC1 signaling

To identify transcriptional alterations that convey enhanced proliferative potential to PDA upon

adaptation to nutrient deprivation, we used RNA-Seq to profile transcripts in SUIT-2 and 8988T clones

under L-L conditions. Principal component analysis (PCA) of the 500 most variable genes indicated

significant segregation (83% variance) of non-adapted clones, based on cell type (Supplementary Fig.

3a). This was however minimized in adapted clones, which clustered together (8% variance), implying

induction of a common transcriptional adaptation program, independent of the genetic background.

Gene set enrichment analysis (GSEA) revealed significant enrichment in the adapted clones for

pathways involved in amino acid synthesis and degradation, as well as purine and pyrimidine synthesis

(Supplementary Fig. 3b).

Metabolic profiling of SUIT-2 and 8988T clones treated for 24 hours with H-H or L-L medium

yielded an adaptation signature that largely overlaps with the altered transcriptomes (Supplementary Fig.

4a,b). Compared to H-H, all clones under L-L conditions displayed markedly higher levels of

intracellular amino acids other than glutamine, glutamate and aspartate (Supplementary Fig. 4a),

consistent with glutamine deprivation and indicative of enhanced autophagy. This increase in amino

acids was significantly suppressed in the adapted clones, compared to non-adapted controls, suggesting

enhanced utilization of intracellular amino acids (Supplementary Fig. 4a). Furthermore, non-adapted

cells showed significant accumulation of nucleotides and their biosynthetic precursors, reflecting

decreased DNA synthesis. In contrast, the adapted clones displayed a nucleotide biosynthesis profile that

closely mirrors that of PDA clones under H-H conditions, consistent with enhanced proliferation. This

profile included lower levels of mono- and di-nucleotide phosphates, used to generate tri-nucleotide

phosphates, which differentially accumulate in the adapted clones (Fig. 2a and Supplementary Fig. 4).

Indeed, although *de novo* pyrimidine synthesis was higher under H-H compared to L-L conditions, a

differential increase upon adaptation was observed only under L-L conditions (Fig. 2b). This was

evident in doubling of ^{14}C incorporation into DNA, 40 hours after labeling of cells with ^{14}C-aspartate

(Fig. 2b).

Signaling downstream of the mammalian target of rapamycin complex 1 (mTORC1) can induce

de novo pyrimidine synthesis through activation of the rate-limiting enzyme carbamoyl-phosphate

synthetase 2, aspartate transcarbamoylase, dihydroorotase (CAD)[25]. We therefore investigated a role for

mTORC1 in the adaptation of the PDA clones to nutrient deprivation. Compared to non-adapted clones,

adapted SUIT-2 and 8988T cells showed increased phosphorylation levels of CAD on Serine 1859, a

site of ribosomal S6-Kinase (S6K1) activity (Fig. 2c). This increase was particularly striking under L-L

conditions, where non-adapted clones showed low (SUIT-2) to undetectable (8988T) CAD

phosphorylation. Importantly, this was mirrored by enhanced mTORC1 activity, as evidenced by

sustained phosphorylation of its downstream effectors S6K1 and 4E-binding protein (4E-BP1), despite

nutrient deprivation (Fig. 2c). On the other hand, AKT phosphorylation levels were decreased in adapted

cells consistent with mTORC1-AKT negative feedback loops[26-28] (Supplementary Fig. 5a).

Adapted PDA clones have enhanced glutamine synthesis

Glutamine is a pleiotropic amino acid required for protein synthesis and numerous anabolic reactions in

proliferating cells[21]. We therefore asked whether the adapted clones have induced *de novo* glutamine

synthesis under L-L conditions. mRNA levels of glutamine synthetase (GS, also termed GLUL), which

catalyzes the condensation of glutamate and ammonium to generate glutamine (Fig. 3a), were only

moderately increased in adapted clones derived from SUIT-2, but not 8988T cells (Supplementary Fig.

6a). However, GS protein levels were markedly induced in all adapted clones from both PDA cell lines,

compared to non-adapted controls (Fig. 3b), where GS levels either remained undetectable (8988T) or

were minimally detected (SUIT-2). This occurred only under L-L conditions, as no clones harbored

detectable levels of GS in H-H medium, independent of adaptation (Fig. 3b). Although these findings

are consistent with a glutamine-dependent post-translational regulation of GS[29, 30], it is noteworthy that

non-adapted clones only displayed a mild to no increase in GS levels 24 hours following exposure to L-

L medium (Fig. 3b), implying differential metabolic rewiring in the PDA clones upon adaptation to

prolonged glutamine deprivation. Importantly, GS protein levels were mirrored by its enzymatic activity

levels, as demonstrated by metabolic tracing of ^{15}N-ammonium chloride (NH$_4$Cl) in SUIT-2 cells (Fig.

3c). Because glutamate dehydrogenase (GDH, also termed GLUD) catalyzes the reductive amination of

alpha-ketoglutarate (α-KG) to generate glutamate (Fig. 3a), a moderate increase was observed in the

^{15}N-glutamate fraction between H-H and L-L conditions in SUIT-2 cells (~18% vs 25%), independent of

adaptation (Fig. 3c). This suggested the use by the adapted cells of newly generated glutamate to

synthesize glutamine. Indeed, adapted clones cultured in L-L medium had a marked increase in ^{15}N-

glutamine (M+1) fraction, generated from either condensation of ^{15}N-ammonium and unlabeled

glutamate, or unlabeled ammonium and GDH-derived ^{15}N-glutamate, compared to non-adapted controls

(30% vs 8%, respectively). This increase was also observed in the (M+2) labeled fraction, which is

generated when both substrates are labeled (8% in adapted vs 2% in non-adapted cells). On the other

hand, ^{15}N-glutamine levels were either undetectable or minimally detected in the PDA clones under H-H

conditions, independent of adaptation (Fig. 3c). These data indicate that only PDA clones that survive

and adapt to long-term glutamine deprivation are able to induce GS protein levels under L-L conditions,

so as to increase GS-mediated glutamine synthesis.

Adapted PDA cells use amino acids as a nitrogen source for glutamine and nucleotide synthesis

In addition to reductive amination of α-KG through GDH (Fig. 3a), the transamination reaction

catalyzed by branched chain amino acid (BCAA) transaminase (BCAT2) could contribute to the

generation of mitochondrial glutamate that is then used for glutamine synthesis. Moreover, aspartate

transaminase (GOT2, also termed AST2) could transfer the nitrogen from glutamate to aspartate, a key

substrate in nucleotide biosynthesis (Fig. 3d). Global metabolite profiling of the PDA clones revealed a

significant decrease in amino acid levels, including the branched chain amino acids (BCAA) leucine,

valine and isoleucine in all adapted clones, suggesting enhanced amino acid utilization by the more

proliferative cells (Fig. 3e and Supplementary Fig. 4a). Although the PDA clones did not seem to share a

common transcriptional regulation of transaminases in both cell lines under L-L conditions

(Supplementary Fig. 6b), protein levels of BCAT2 and GOT2 were significantly increased in at least 2

of the 3 SUIT-2 and 8988T adapted clones (Supplementary Fig. 6c). ^{15}N-leucine tracing demonstrated

that the labeled nitrogen incorporates at significantly higher levels into glutamate, aspartate and

intermediates in nucleotide synthesis in adapted, compared to non-adapted clones under L-L conditions

(Fig. 3f and Supplementary Fig. 7a). Moreover, the ^{15}N-labeled fraction of glutamine was minimal (1%)

under nutrient-replete H-H conditions, whereas a marked 5-fold increase in ^{15}N-glutamine was detected

under L-L conditions, specifically in the adapted clones (Fig. 3f and Supplementary Fig. 7a). These data

imply that BCAAs are a significant source for *de novo* synthesis of glutamine and nucleotides in PDA

cells adapted to L-L conditions.

mTORC1 activation and enhanced proliferative potential are reversible in adapted PDA clones

Because the adapted clones sustain active mTORC1 signaling independent of nutrient deprivation, we

asked whether the adaptation process had selected for cells with inherent mutations that constitutively

activate mTORC1. We cultured the adapted clones in nutrient-replete H-H medium for 4 or more

consecutive passages (a process we termed "adaptation reversal") followed by treatment for 24 hours

with L-L medium (Fig. 4a). Reverse-adapted (RA) clones displayed gradual suppression of mTORC1

activity (Fig. 4b and Supplementary Fig. 8a,b), indicating that the enhanced anabolic signaling acquired

upon adaptation is unlikely to result from genetic mutations selected for in the adapted clones.

Importantly, *in vitro* survival and growth of adapted clones were partially reversed, with all RA clones

exhibiting decreased proliferation compared to adapted clones and two 8988T clones displaying

enhanced cell death (Fig. 4c,d and Supplementary Fig. 8c). Furthermore, among a pool of 1,842 genes

that were significantly up-regulated ($\geq$3-fold, q <0.05) upon adaptation in both SUIT-2 and 8988T cells,

reversal was associated with partial suppression of this transcriptional response (Fig. 4e), consistent with

decreased mTORC1 signaling and proliferative capacity (Fig. 4b-d and Supplementary Fig. 8a-c).

These findings led us to investigate a potential epigenetic basis for PDA cells' adaptation to

nutrient deprivation. Assays for transposase-accessible chromatin with high throughput sequencing

(ATAC-seq) identified a significantly altered landscape of open chromatin in adapted clones, with gains

and losses of chromatin accessibility at thousands of genomic sites (Fig. 4f), including 3,807 enhancers

common to adapted 8988T and SUIT-2 clones (Supplementary Fig. 8d). Adaptation-induced changes in

chromatin accessibility correlated with gene expression changes in both lines (Fig. 4g and

Supplementary Fig. 8e). Among the 1,842 transcripts altered upon adaptation and linked to enhancers

modulated in both cell lines, gains of chromatin access and transcription outnumbered the losses (Fig.

4g). Notably, alterations in open chromatin were largely preserved in RA cells (Fig. 4f and

Supplementary Fig. 8f, left) and reversion was infrequent (Fig. 4f and Supplementary Fig. 8f, right),

indicating possible acquisition of epigenetic memory when PDA cells adapt robustly to nutrient

deprivation.

mTORC1 prevents proteasomal degradation of GS in adapted PDA clones

Because the induction of GS protein in adapted cells consistently mirrored mTORC1 signaling (Fig. 2c

and 3b), including reversibility upon de-adaptation (Fig. 4b and Supplementary Fig. 8a-b), we asked

whether mTORC1 regulates GS levels. Upon treatment with the mTOR kinase inhibitor Torin1, *GS*

mRNA levels were increased in adapted clones under L-L conditions (Supplementary Fig. 9a). In

contrast, consistent with post-transcriptional GS regulation, Torin1 markedly decreased GS protein

levels (Fig. 5a). This decrease was not observed upon treatment with cycloheximide, but only when

Torin1 was added to the cells, indicating that mTORC1 regulation of GS is post-translational (Fig. 5b

and Supplementary Fig. 9b). Indeed, whereas treatment with the proteasomal inhibitor MG-132 did not

significantly affect GS levels in adapted clones over 2-12 h, it induced GS in all non-adapted clones

under L-L conditions (Fig. 5c). Importantly, the Torin1-induced decrease in GS levels in adapted cells

was rescued upon treatment with MG-132 (Fig. 5d). Furthermore, mTORC1 inhibition promoted GS

poly-ubiquitination, as evidenced in immunoprecipitated GS from adapted cells co-treated with Torin1

and MG-132 (Fig. 5e). Altogether, these data demonstrate that GS is ubiquitinated and targeted for

proteasomal degradation in non-adapted clones, even under L-L conditions, and that mTORC1 prevents

this degradation, leading to GS stabilization in adapted clones.

PDA cells adapted to nutrient-deprived conditions are sensitive to GS inhibition

To assess the contribution of glutamine synthesis to the enhanced growth of the adapted cells, PDA

clones were treated with the GS inhibitor L-methionine sulfoximine (MSO). Adapted cell proliferation

was mitigated under L-L, but not H-H conditions, underscoring the relevance of glutamine synthesis in

the nutrient-deprived state (Supplementary Fig. 10a). Genetic silencing of GS similarly suppressed

adapted cell proliferation and colony formation (Fig. 6a-c, Supplementary Fig. 10b,c). To ask whether

the contribution of GS to enhanced *in vitro* PDA growth extends *in vivo*, we induced knockdown of GS

in orthotopic PDA transplants (Fig. 6d,e). Five days following injection of adapted 8988T expressing

control or *GS* hairpins into the pancreas, gene silencing was induced by treating mice with doxycycline

(Dox) in the drinking water. Two distinct *GS* hairpins attenuated PDA growth (Fig. 6d and

Supplementary Fig. 10e,f), resulting in 6.1- to 9.2-fold lower tumor weights upon Dox treatment (Fig.

6d). These findings indicate that GS-mediated glutamine synthesis is required for PDA cells to survive

and adapt to the characteristic nutrient-deprived PDA environment and highlight GS as a metabolic

target in treating pancreatic cancer (Fig. 6f).

To extend our findings to human pathology, tissue microarrays (TMAs) of resected tumors from

127 pancreatic cancer patients were immunostained and scored for GS protein levels. As a reference for

intensity, stained control cores of normal pancreas showed high GS levels in islets and modest levels in

acinar cells. Within tumor cores, PDA cells displayed heterogeneous cytoplasmic staining that varied

widely from strong to moderate to weak, with an overall score lower than that of acinar cells. As a

result, no significant association was found between PDA GS levels and tumor size, lymph node

involvement (pT or pN staging, the American Joint Commission on Cancer, AJCC 8[th] edition[31]) or

patient survival (Supplementary Fig. 11a-c) Although these results seem discordant with a recent report

that GS expression is a feature of *KRAS*-driven PDA[32], the pathological heterogeneity and variability in

GS protein levels is consistent with an inverse correlation with nutrient deprivation (Figs. 3, 4 and

Supplementary Fig. 8). Indeed, nutrient distribution is recognized to be heterogeneous in the tumor

microenvironment, with glutamine levels being most depleted within tumor cores, compared to the

periphery[18, 19]. Therefore, gradients of GS protein levels may exist among distinct PDA cell clusters

within the tumor mass, depending on their precise location and proximity to blood vessels. Nevertheless,

based on our results and those of Bott *et al.*[32], targeting GS in the nutrient-deprived PDA tumors holds

significant therapeutic value in pancreatic cancer.

Discussion

The *in vitro* dependence of proliferating cells on glutamine[21] was recently questioned in the *in vivo*

setting, where tumors preferably utilize glucose and other metabolites, rather than circulating glutamine,

as sources of tricarboxylic acid (TCA) cycle carbon[13, 33-35]. This observation is supported by *in vivo*

resistance of tumors to inhibitors of glutaminolysis[36]. Differences in nutrient levels between cell culture

media and human serum, in addition to the *in vivo* tumor environment, might partially explain this

discrepancy[11, 37]. However, although glutamate is a major source of TCA cycle carbon and nitrogen for

transamination reactions and glutathione synthesis, glutamine *per se* is required for protein synthesis and

is an indispensable source of amide nitrogen for nucleotide and hexosamine synthesis. Therefore,

whereas proliferating tumor cells may resort to alternative sources of glutamate, such as transamination

reactions instead of glutaminolysis, they can only survive in glutamine-deprived conditions if they can

synthesize glutamine *de novo*.

Moreover, although glutamine may not seem to be depleted in extracts of whole tumors or tumor

interstitial fluid[15], tumor cores hold lower levels than the periphery[18, 19]. Thus, the precise location of a

cancer cell within the hypoxic, stromally dense tumor may dictate its dependence on glutamine at

various times. Distinct clusters of tumor cells likely exist within PDA tumors with differential nutrient

access depending on the proximity to blood vessels and density of the surrounding stroma. The

deprivation of PDA tumor cells of nutrients may also change over time, as tumor growth reshapes 3D

cellular organization and the proximity of cancer cells to stroma and endothelial cells. It is therefore not

surprising that, unlike Bott *et al.*[32], we found no significant association between average GS levels and

tumor grade or patient survival (Supplementary Fig. 11). We find instead that rare clonal cells are able

to survive and adapt to low levels of the tumor-preferred nutrients glucose and glutamine; these cells

induce glutamine synthesis, which is required for their growth *in vitro* and *in vivo*.

Despite chronic nutrient deprivation in the adapted cells, we find that activated mTORC1

induces GS. Although GS is known to be regulated post-translationally[29, 30], the signaling basis for this

regulation was unknown. We show that mTORC1 plays a key post-translational role in stabilizing GS by

preventing its ubiquitination and proteasomal degradation. mTORC1 inhibition was previously shown to

enhance proteasomal degradation of long-lived proteins[38], but GS was not identified as a specific target,

perhaps because the experiments were performed under glutamine-replete conditions. What induces

mTORC1 activity in the absence of glucose and glutamine in adapted PDA cells is a question worthy of

future investigation. One plausible explanation is that epigenetic alterations that occur upon adaptation

to nutrient deprivation (Fig. 4 and Supplementary Fig. 8) may influence genes that directly or indirectly

activate mTORC1.

We propose that PDA cells that survive and adapt to nutrient deprivation acquire an epigenetic

state that endows them with the flexibility to switch between metabolic states that are optimal for growth

under nutrient-replete or nutrient-deprived conditions. A key contributing factor to the adaptation to

nutrient-depleted conditions is activation of mTORC1, which in turn stabilizes GS, thus enabling

essential glutamine synthesis for pancreatic cancer growth (Fig. 6f). Consequently, targeting of GS,

which expression is induced in nutrient-deprived cell clusters within PDA tumors, holds therapeutic

benefit to pancreatic cancer patients.

Methods

Reagents. Antibodies for western blotting were from Abcam: BCAT1 (ab107191), GOT2 /

FABP-1 (ab171739); BD Biosciences: GS / GLUL (BD-61057); Cell Signaling Technology

(CST): AKT (pan, C67E7, # 4691), pS473-AKT (193H12, # 4058), pT308-AKT (244F9, #

4056), BCAT2 (# 9432), CAD (# 11933), pS1859-CAD (# 12662), 4EBP1 (53H11, # 9644),

pS65-4EBP1 (# 9451), LC3B (# 2775), pThr389-S6K (# 9234); Novus Biologicals: GOT1

(NBP1-54778); Proteintech: GLUD1 / GDH1 (14299-1-AP); Santa Cruz Biotechnology: S6Kα

(H-9, sc-8418), GAPDH (FL-335, sc-25778), β-Actin (sc-47778). All western blot antibodies

were used at 1:1,000 dilution, except for β-Actin (1: 20,000). Antibodies for

immunoprecipitation, ubiquitination or immunohistochemistry are described in the respective

Methods sections below. Chloroquine (Sigma-Aldrich, C6628); Cycloheximide (Sigma-Aldrich,

C4859); DMSO (Sigma-Aldrich D2650); MG-132 (Selleck Chemicals, S2619); Doxycycline

(Sigma-Aldrich; D9891); L-Methionine Sulfoximine, MSO (Sigma-Aldrich; M5379); Torin 1

(Selleck Chemicals, S2827).

Cell culture. Human PDA cell lines were from the American Type Culture Collection or ATCC

(AsPC-1, BxPC-3, HPAC, MIA PaCa-2, PANC-1), Japanese Collection of Research

Bioresources (SUIT-2), or the German Collection of Microorganisms and Cell Cultures (PA-TU-

8988T) and were authenticated by STR profiling at ATCC. All cell lines tested negative for

mycoplasma using LookOut Mycoplasma PCR Kit (Sigma, MP0035). All cells were maintained

at 37 °C in a humidified incubator with 5% CO_2 and were grown in media supplemented with

10% dialyzed fetal bovine serum (Sigma-Aldrich F0392). For adaptation or reverse adaptation to

"low glucose, low glutamine" (L-L) medium, PDA cells were cultured in RPMI 1640 medium

lacking both glucose and L-glutamine (US Biological, R9011-01) that was supplemented with

0.5 mM glucose (Thermo Fisher Scientific, A2494001) and 0.1 mM glutamine (Thermo Fisher

Scientific, 25030-81). Non-adapted cells were cultured in the same RPMI 1640 medium that was

however supplemented with 11mM glucose and 2 mM glutamine, and was termed "high glucose,

high glutamine" (H-H) medium.

Gene knockdown. For small interfering RNA knockdown, ON-TARGETplus Human GS

(GLUL) siRNA with target sequence: GCACACCUGUAAACGGAUA (J-008228-09) or control

non-targeting siRNA: UGGUUUACAUGUUGUGUGA (D-001810-02-05) were purchased from

Dharmacon, Horizon Discovery (Cambridge, UK) and transfected according to the

manufacturer's protocol. For generation of PDA cells with stable doxycycline-inducible GS or

control (Scramble) knockdown, the corresponding hairpins were cloned into Tet-pLKO-puro

plasmid (21915; Addgene). Lentiviral supernatants were generated by transfecting the plasmids

into 293T cells and used to infect 8988T cells that were then selected for at least 7 days with 2

µg ml^{-1} puromycin (Sigma-Aldrich, P8833). The RNAi consortium clone IDs for the hairpins

used were: shGS #1 (G05) TRCN0000343992 (CACACCTGTAAACGGATAATG); shGS #2

(G06) TRCN0000344059 (ATAACCACTGCTTCCATTTAA); Control Scramble shRNA

sequence (CCTAAGGTTAAGTCGCCCTCGCTCGAGCGAGGGCGACTTAACCTTAGG)

was the same as that from Addgene Plasmid # 1864.

Cell proliferation and cell death assays. Cells were seeded on day 0 in their respective

maintenance culture medium in 96-well plates, at a density of 3,000 cells per well, and incubated

overnight. On day 1, the cells were washed once with phosphate buffered saline (PBS) and

incubated in their experimental medium supplemented with propidium iodide (2 µg ml^{-1}) for up

to 7 days. Live and dead cells were counted on the indicated days using the Celigo Image

Cytometer (Nexcelom Bioscience), and either normalized to day 1 (live cell proliferation

curves), or presented as percentage of propidium iodide-positive cells per total cell number (cell

death).

Clonogenic assay. 600 cells were seeded per well in 6-well plates in their appropriate

maintenance medium and incubated overnight. The next day, cells were washed once with PBS

and incubated in their experimental medium for 7-10 days. Colonies were then either directly

counted using the Celigo Image Cytometer (Nexcelom Bioscience), or first stained and then

counted using the following protocol: media were removed and the cells fixed in a solution of

0.2% Crystal Violet (Sigma-Aldrich, C0775) in 80% methanol for 15-30 min. The wells were

then washed in distilled water and allowed to dry overnight, before colonies were counted using

the Celigo Image.

Sphere formation assay. 50 cells were seeded per well in 96-well plates and incubated

overnight in their appropriate maintenance medium supplemented with 4% matrigel (Corning,

356231). The next day, cells were washed once gently with PBS and treated with the

experimental medium. Spheres were counted after 7-10 days using the Celigo Image Cytometer

(Nexelcom Bioscience).

Macropinocytosis assay. Cells were seeded onto glass coverslips (Bellco Glass, 1943-10012A)

in 24-well plates and incubated overnight in their maintenance culture medium. The next day, the

medium was replaced with fresh L-L medium containing 10% dialyzed FBS, and the cells were

incubated for 24 h. Macropinosomes were assayed as previously described[39] and outlined below.

70-kDa TMR-dextran (Thermo Fisher Scientific, D1818) was added to the cells at a final

concentration of 1 mg ml^{-1} and the cells incubated for 30 min at 37°C, in the absence or presence

of 25 μM EIPA (Sigma-Aldrich; A3085). Cells were then rinsed 3 times on ice with ice-cold

PBS and immediately fixed in 4% paraformaldehyde (EMS, 15710-S) for 30 min. After staining

the cells with 1 μg ml^{-1} DAPI (Thermo Scientific, D3571) in PBS for 15min, the coverslips were

mounted onto glass slides (VWR, 48311-703) using Fluoroshield mounting medium (Sigma,

F6182). Image acquisition was performed using a Nikon Eclipse 90i Advanced Automated

Research Microscope equipped with a standard optical filter set including DAPI and

Rhodamine/Texas Red. Images were captured with a 40x objective and the NIS-Elements

Advanced Research Microscope Imaging Software (Nikon). Exposure time for DAPI acquisition

was 100 ms and that for TMR acquisition was 400 ms. Representative images were captured in

at least 3 different fields per well in duplicate wells, per condition and analyzed using ImageJ

software (NIH). At least 200 cells were counted per condition. Macropinosomes were visualized

after setting the threshold of the "Auto Contrast" feature in Image J to 134. The number and area

of labeled macropinosomes were then determined using the "Analyze Particles" feature, and

macropinocytic uptake was computed by the total particle number and area per cell.

U-^{14}C-asparate incorporation into DNA

Cells were seeded overnight (57,000 per well) in 6-well-plates with triplicate wells per condition.

The cells were then washed twice with PBS and incubated with either H-H or L-L medium

supplemented with 1μCi per well of L-[U-^{14}C]-Aspartic acid (PerkinElmer,

NEC268E050UC) for 40 h. The cells were washed twice with PBS before DNA was extracted

using PureLink® Genomic DNA Mini Kit (Thermo Fisher-K182001) and quantified using a

spectrophotometer. Equal volumes of DNA were added to scintillation vials and radioactivity

was measured by liquid scintillation counting and normalized to total DNA concentration.

Immunoblotting. For western blotting, cells were rinsed once in ice-cold PBS and collected in

lysis buffer containing 50 mM HEPES KOH, pH 7.4, 40mM NaCl, 2mM EDTA, 1.5mM

orthovanadate, 50mM NaF, 10mM pyrophosphate, 10mM glycerophosphate, EDTA-free

protease inhibitors (Thermo Fisher Scientific, 88266) and 1% Triton X-100. Proteins from total

lysates were resolved by 8–12% SDS–PAGE, transferred to polyvinylidene difluoride (PVDF)

and the blot was exposed to film.

Immunoprecipitation and ubiquitination assay. To assess GS protein ubiquitination, cells

were treated with either control vehicle DMSO or MG132 (10 μM) for 8 h. Cells were then lysed

under denaturing conditions using the lysis buffer (40 mM HEPES KOH, pH 7.4, 1% SDS)

containing 5 mM N-ethylmaleimide (NEM, Sigma-Aldrich, E1275) to prevent deubiquitination

and boiled for 5 min. The lysates were diluted 10-fold in IP buffer (40 mM HEPES KOH pH 7.4,

50 mM NaF, 10 mM Sodium pyrophosphate, 10 mM Sodium beta-glycerophosphate, 1.5 mM

Sodium Orthovanadate, 2.5 mM MgCl2, EDTA-free protease inhibitors, and 0.5% Triton X-

100). IP was performed by incubating the samples with 2 μg of normal rabbit IgG control (Santa

Cruz, sc-2027) or 2μg GS antibody (Abcam, ab73593) overnight with rotation at 4°C. Protein G

agarose bead slurry (Invitrogen, 15920-010) was added to the pre-cleared lysates (60 μl, 50:50)

and incubated with rotation for 1.5-2 hr at 4°C. The beads were then washed once with IP buffer,

followed by 3 additional washes with IP buffer supplemented with 500 mM NaCl. Loading

buffer was added to the immunoprecipitated proteins, which were then boiled, resolved by SDS-

PAGE and transferred to a PVDF membrane. Ubiquitinated GS was identified by HRP-

conjugated anti-Ubiquitin antibody (1:5,000; Enzo Life Sciences, FK2, BML-PW0150-0100).

Quantitative real-time PCR. Total mRNA was isolated with RNA STAT-60 (Tel-Test, Cs-111)

according to the manufacturer's instructions, treated with DNase I (RNase-free, Roche

Molecular Biochemicals**,** 04716728001) and reverse-transcribed into cDNA with random

hexamers using the SuperScript II First-Strand Synthesis System (Invitrogen, 18064071).

Primers listed below were validated and PCR reactions performed as previously described[40].

Quantitative PCR reactions were performed in triplicates using an Applied Biosystems ViiA 7

Real-Time PCR system. PCR reactions contained cDNA resulting from reverse transcription of

25ng total RNA, 150nM of each primer and 5 μl 2X-Jump Start SYBR Green PCR Mix

(Invitrogen) in10 μl total volume. Relative mRNA levels were calculated using the comparative

CT method and normalized to cyclophilin. The QPCR primers used were:

BCAT1: 5'-GTGGAGTGGTCCTCAGAGTTT-3' and 5'-AGCCAGGGTGCAATGACAG-3'

BCAT2: 5'-GCTCAACATGGACCGGATG-3' and 5'-CCGCACATAGAGGCTGGTG-3'

CyclophilinA (CYPA): 5'-GGAGATGGCACAGGAGGAA-3' and 5'-

GCCCGTAGTGCTTCAGCTT-3'

GOT1: 5'-ATTTCTTAGCGCGTTGGTACA-3' and 5'-ACACAGCATTGTGATTCTCCC-3'

GOT2: 5'-AGCCTTACGTTCTGCCTAGC-3' and 5'- AAACCGGCCACTCTTCAAGAC-3'

GS: 5'-AAGAGTTGCCTGAGTGGAATTTC-3' and 5'-AGCTTGTTAGGGTCCTTACGG-3'**.**

RNA-Seq sample preparation: Cells were seeded overnight at 700,000 per 10 cm plate. The

cells were then washed once with PBS and incubated in L-L medium for 24 h. RNA was

isolated and purified using miRNeasy Mini Kit (Qiagen-217004). Total RNA (> 150 ng) was

sent to the Broad Institute Genomics Platform for library preparation. RNA quality and insert-

size was assessed by Caliper LabchipGXII producing a RQS value (equivalent to RIN). RNA

quantity was determined by PicoGreen. mRNA libraries were prepared using the Illumina

TruSeq Stranded mRNA kit modified for improved performance and multiplexing. Libraries

were pooled and sequenced with the Illumina HiSeq 4000 using 101-cycle, pair-end settings to

obtain 50 million read sequencing depth per sample.

RNA read processing and Quantification. All samples were processed using an RNA-Seq

pipeline implemented in the bcbio-nextgen project (https://bcbio-

nextgen.readthedocs.io/en/latest/). Raw reads were examined for quality issues using FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to ensure library generation and

sequencing data were suitable for further analysis. Reads were aligned to the hg19 build of

the human genome using STAR[41]. Quality of alignments was assessed by checking for evenness

of coverage, rRNA content, genomic context of alignments, complexity and other quality checks.

Expression quantification was performed with Salmon[42] to identify transcript-level abundance

estimates and then collapsed down to the gene-level using the R Bioconductor package

tximport[43]. Principal components analysis (PCA) and hierarchical clustering methods validated

clustering of samples by cell line and group. Differential expression was performed at the gene

level using the R Bioconductor package DESeq2[44]. Differentially expressed genes were

identified using the Wald test and significant genes were obtained using an FDR threshold of

0.05. GSEA[45] was carried out on the full expression data matrix using the gene sets collection

from Kyoto Encyclopedia of Genes and Genomes (KEGG) to assess the enrichment of the

"adaptation to nutrient deprivation" signature. A pairwise GSEA was performed by creating

ranked lists of genes using the difference of class means to calculate fold change for log scale

data between the Adapted and Non-Adapted PDA clones (Supplementary Fig. 3) and P values

were obtained from permuting the gene set (1000 permutations).

ATAC-Seq. Cells were seeded overnight at 700,000 per 10 cm plate, washed once with PBS,

and incubated in L-L medium for 24 h. The assay[46] was performed on 10^6 cells per condition.

Cells were washed twice in ice-cold PBS and resuspended in 50 μl ice-cold ATAC lysis buffer

(10 mM Tris.Cl, pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% (v/v) Igepal CA-630), and

centrifuged at 500 g at 4°C to isolate nuclear pellets that were treated in 50 μL reactions with

Nextera Tn5 Transposase (Illumina, FC-121-1030) for 30 min at 37°C. Column-purified DNA

(QIAGEN) was stored at 20°C or amplified immediately in 50 μl reactions with high-fidelity 2X

PCR Master Mix (New England Biolabs) using a common forward primer and different reverse

primers with unique barcodes for each sample. From the reaction mix, 45 μl was kept on ice after

5 cycles of PCR, while 5 μl was amplified by qPCR for 20 additional cycles; the remaining 45 μl

was then amplified for the 5-7 cycles required to achieve 1/3 of the maximum qPCR

fluorescence intensity. Amplified DNA was purified over columns and primer dimers (<100 bp)

were removed using AMPure beads (Beckman Coulter). Size distribution of the amplified DNA

was analyzed using High-sensitivity Qubit dsDNA Assay Kit (ThermoFisher). High throughout

sequencing was performed by Novogene Corporation, Inc. (Sacramento, CA). Computational

analysis was performed as described previously[47].

Metabolite tracing, extraction and quantification. For stable isotope tracing experiments, cells

were incubated for 24 h in L-L or H-H medium supplemented with either 0.8mM [15]N-

ammonium chloride (99% enrichment, NLM-467-1) or 0.4mM [15]N-leucine (98% enrichment,

NLM-142-0.5) purchased from Cambridge Isotope Laboratories. Cells were then rinsed in ice-

cold PBS and metabolites extracted with ice-cold 80% methanol, vortexed for 10min at 4°C, and

centrifuged at top speed (10min, 10,000 x g, 4°C). Supernatants were transferred to fresh

512 Eppendorf tubes and dried using a SpeedVac. Dried extracts were suspended in 100 μl water and

513 centrifuged at top speed for 10 min, and the supernatant was analyzed using LC-MS. For Figures

514 2a, 3c and e, and Supplementary Figure 4a, data analysis and natural abundance correction of the

515 stable isotope labeling data were performed as previously described[48]. For Figure 3f and

516 Supplementary Figure 7, data were acquired using a hydrophilic interaction liquid

517 chromatography (HILIC)-positive ion mode mass spectrometry (MS) method operated on

518 Nexera X2 UHPLC (Shimadzu Scientific Instruments, Marlborough, MA) coupled to a Q

519 Exactive orbitrap mass spectrometer (Thermo Fisher Scientific, Waltham, MA) as described

520 previously[49], with full scan high resolution MS data collected over 70–800 m/z. For global

521 steady-state metabolite profiling, data were analyzed using MetaboAnalyst 4.0[50], median-

522 normalized, log-transformed, mean-centered and divided by the S.D. of each variable.

524 **Animal work.** All mouse studies and procedures were approved by the Animal Care and Use

525 Committee at Boston Children's Hospital. For orthotopic xenografts, 750,000 cells suspended in

526 25 μl 33% Matrigel (Corning, 356231) in Hank's Balanced Salt Solution (HBSS) were injected

527 into the pancreata of male B6.129S7-Rag1[tm1Mom]/J mice termed Rag1$^{-/-}$ mice (Jackson

528 Laboratory #002216). *In vivo* ultrasound imaging (Vevo 2100, MS550D Scanhead) was

529 performed on mice to detect and quantify tumor growth at the Small Animal Imaging Lab

530 (SAIL) at Boston Children's Hospital. Following experimental endpoint euthanasia, tumors

531 were collected, their dimensions (**a**, **b** and **c**) were measured with a caliper and tumor volume

532 was estimated according to the ellipsoid formula[51]: 4/3 x π x (a/2 x b/2 x c/2). Tumor tissue was

533 then either immediately frozen in liquid nitrogen, or fixed in formalin for later processing.

Immunohistochemistry and analysis of human samples. Formalin-fixed paraffin-embedded

PDA tissue microarrays (TMAs) with 2–3 cores per case, were sectioned and immunostained

with GS antibody (abcam, ab-73593), according to the manufacturer's protocol. A pathologist

scored, in a blinded fashion, the intensity of staining on a scale of 0–3 that represented none (0),

weak (1), moderate (2), or strong (3) based on the PDA cell staining. Scores for each tumor core

were estimated using the following formula: H-score = 0 x (% area of tumor with no staining) +

1 x (% tumor area with weak staining) + 2 x (% tumor area with moderate staining) + 3 x (%

tumor area with strong staining), where 300 would be the highest score. TMAs were derived

from 127 patients with surgically resected PDA from Massachusetts General Hospital. Informed

patient consent was waived by the Institutional Review Board or IRB (Protocol Number

2009P001838), as the TMAs were generated from discarded material following clinical

diagnosis. For analyses of human data, the following variables were from electronic medical

records: age, sex, pN and pT stage based on the American Joint Committee on Cancer 8th

edition, tumor size, location and grade of differentiation, and presence of lymphovascular and

perineural invasion. Survival analyses were performed using overall survival (OS) as the

outcome of interest, defined as the time between surgical resection and death from any cause or

date of last clinical contact. Survival outcomes were compared based on tertiles of GS staining

intensity as assessed by H-scores using the Kaplan-Meier method. Tertile thresholds were 115

and 150 H-score units. As secondary analyses, GS intensity was evaluated for its association

with baseline clinicopathological characteristics at the time of surgery using the Kruskal-Wallis

test. Statistical significance was set at P <0.05. Data analysis was performed using SAS 9.4

(version 9.4, SAS Institute, Cary, NC)."

Statistical Analysis

Data were presented as mean ± S.D. or ± S.E.M., unless otherwise indicated. When comparing

two groups, a two-tailed non-paired Student's t-test was conducted. For three or more groups,

one-way ANOVA was conducted except for proliferation curves, where two-way ANOVA was

conducted. ANOVA was followed by post hoc Tukey's multiple-comparison test. $p < 0.05$ was

considered statistically significant.

Acknowledgements

We thank members of the Kalaany lab and the Division of Endocrinology for helpful

comments and discussions. This work was supported by the American Cancer Society

Research Scholar Grant (RSG-17-070-01-TBG) to N.Y.K. and funds from Boston Children's

Hospital. N.Y.K. is also supported by NIH/NCI Grant R01 CA211944. P-Y.T. was partly

funded by Top University Strategic Alliance (TUSA) fellowship from Taiwan. Work by M.M.

was supported by Harvard Catalyst, The Harvard Clinical and Translational Science Center

(NIH award #UL1 RR 025758) and computations were run on the Orchestra cluster supported

by HMS Research Computing Group.

Author Contributions

N.Y.K. conceived, designed and supervised the study. P-Y.T. performed the experiments with

help from lab members and collaborators. M-S.L. performed biochemical assays for GS

stabilization and growth assays for stable GS knockdown with help from I.N. who also assisted

in macropinocytosis quantification and western blotting. A.A. assisted P-Y.T. with tumor cell

injection into mice. M.M. analyzed RNA-Seq data for PCA and helped P-Y.T. in pathway

analyses. R.A.M. supervised the ATAC-Seq experiment and analyses, contributing to the

interpretation of the results. U.J. performed the ATAC-Seq assay and analyzed the ATAC-Seq

and RNA-Seq data with help from S.M. C.A.L. performed metabolite quantification and tracing

analyses (Figs 2a and 3c,d and Supplementary Fig. 4). D.S.H., with supervision from C.B.C.,

performed metabolite quantification and tracing analyses (Fig. 3f and Supplementary Fig. 7).

F.R.R. assisted with mouse ultrasound imaging. M.M.-K. provided TMAs and pathological

assessment and scoring for immunostained human pancreatic tissue. T.H. and K.C.H. helped

retrieve patient data. P.D. performed the statistical analyses on human data with consultation

from V.M.-O. who contributed to the discussion of the results. N.Y.K. and P.-Y.T. wrote the

manuscript with feedback from all authors.

Supplementary information

Supplementary Figures 1-11 accompany this manuscript.

Data availability

The RNA-Seq and ATAC-Seq data have been deposited in NCBI's Gene Expression Omnibus

and will be accessible through GEO Series accession number GSE144833.

Competing interests

The authors declare no competing financial interests.

References

1.   Siegel, R.L., Miller, K.D. & Jemal, A. Cancer statistics, 2019. *CA Cancer J Clin* **69**, 7-34
(2019).
2.   Rahib, L. *et al.* Projecting cancer incidence and deaths to 2030: the unexpected burden of
thyroid, liver, and pancreas cancers in the United States. *Cancer Res* **74**, 2913-2921
(2014).
3.   Perera, R.M. & Bardeesy, N. Pancreatic Cancer Metabolism: Breaking It Down to Build
It Back Up. *Cancer Discov* **5**, 1247-1261 (2015).
4.   Ying, H. *et al.* Genetics and biology of pancreatic ductal adenocarcinoma. *Genes Dev* **30**,
355-385 (2016).
5.   Yang, S. *et al.* Pancreatic cancers require autophagy for tumor growth. *Genes Dev* **25**,
717-729 (2011).
6.   Commisso, C. *et al.* Macropinocytosis of protein is an amino acid supply route in Ras-
transformed cells. *Nature* **497**, 633-637 (2013).
7.   Kamphorst, J.J. *et al.* Human pancreatic cancer tumors are nutrient poor and tumor cells
actively scavenge extracellular protein. *Cancer Res* **75**, 544-553 (2015).
8.   Ying, H. *et al.* Oncogenic Kras maintains pancreatic tumors through regulation of
anabolic glucose metabolism. *Cell* **149**, 656-670 (2012).
9.   Son, J. *et al.* Glutamine supports pancreatic cancer growth through a KRAS-regulated
metabolic pathway. *Nature* **496**, 101-105 (2013).
10.   Muir, A. & Vander Heiden, M.G. The nutrient environment affects therapy. *Science* **360**,
962-963 (2018).
11.   Davidson, S.M. *et al.* Environment Impacts the Metabolic Dependencies of Ras-Driven
Non-Small Cell Lung Cancer. *Cell Metab* **23**, 517-528 (2016).
12.   Hensley, C.T. *et al.* Metabolic Heterogeneity in Human Lung Tumors. *Cell* **164**, 681-694
(2016).
13.   Maher, E.A. *et al.* Metabolism of [U-13 C]glucose in human brain tumors in vivo. *NMR*
Biomed **25**, 1234-1244 (2012).
14.   Marin-Valencia, I. *et al.* Analysis of tumor metabolism reveals mitochondrial glucose
oxidation in genetically diverse human glioblastomas in the mouse brain in vivo. *Cell*
Metab **15**, 827-837 (2012).
15.   Sullivan, M.R. *et al.* Quantification of microenvironmental metabolites in murine cancers
reveals determinants of tumor nutrient availability. *Elife* **8** (2019).
16.   Cantor, J.R. *et al.* Physiologic Medium Rewires Cellular Metabolism and Reveals Uric
Acid as an Endogenous Inhibitor of UMP Synthase. *Cell* **169**, 258-272 e217 (2017).
17.   Vande Voorde, J. *et al.* Improving the metabolic fidelity of cancer models with a
physiological cell culture medium. *Sci Adv* **5**, eaau7314 (2019).
18.   Pan, M. *et al.* Regional glutamine deficiency in tumours promotes dedifferentiation
through inhibition of histone demethylation. *Nat Cell Biol* **18**, 1090-1101 (2016).
19.   Lee, S.W. *et al.* EGFR-Pak Signaling Selectively Regulates Glutamine Deprivation-
Induced Macropinocytosis. *Dev Cell* (2019).
20.   Psychogios, N. *et al.* The human serum metabolome. *PLoS One* **6**, e16957 (2011).
21.   Altman, B.J., Stine, Z.E. & Dang, C.V. From Krebs to clinic: glutamine metabolism to
cancer therapy. *Nat Rev Cancer* **16**, 749 (2016).

22.   Gebregiworgis, T. *et al.* Glucose Limitation Alters Glutamine Metabolism in MUC1-
Overexpressing Pancreatic Cancer Cells. *J Proteome Res* **16**, 3536-3546 (2017).
23.   Tan, H.W.S., Sim, A.Y.L. & Long, Y.C. Glutamine metabolism regulates autophagy-
dependent mTORC1 reactivation during amino acid starvation. *Nat Commun* **8**, 338
(2017).
24.   Yan, L., Raj, P., Yao, W. & Ying, H. Glucose Metabolism in Pancreatic Cancer. *Cancers*
(Basel) **11** (2019).
25.   Ben-Sahra, I., Howell, J.J., Asara, J.M. & Manning, B.D. Stimulation of de novo
pyrimidine synthesis by growth signaling through mTOR and S6K1. *Science* **339**, 1323-
1328 (2013).
26.   Harrington, L.S. *et al.* The TSC1-2 tumor suppressor controls insulin-PI3K signaling via
regulation of IRS proteins. *J Cell Biol* **166**, 213-223 (2004).
27.   Hsu, P.P. *et al.* The mTOR-regulated phosphoproteome reveals a mechanism of
mTORC1-mediated inhibition of growth factor signaling. *Science* **332**, 1317-1322
(2011).
28.   Shah, O.J., Wang, Z. & Hunter, T. Inappropriate activation of the TSC/Rheb/mTOR/S6K
cassette induces IRS1/2 depletion, insulin resistance, and cell survival deficiencies. *Curr*
Biol **14**, 1650-1656 (2004).
29.   Nguyen, T.V. *et al.* Glutamine Triggers Acetylation-Dependent Degradation of
Glutamine Synthetase via the Thalidomide Receptor Cereblon. *Mol Cell* **61**, 809-820
(2016).
30.   Nguyen, T.V. *et al.* p97/VCP promotes degradation of CRBN substrate glutamine
synthetase and neosubstrates. *Proc Natl Acad Sci U S A* **114**, 3565-3571 (2017).
31.   Allen, P.J. *et al.* Multi-institutional Validation Study of the American Joint Commission
on Cancer (8th Edition) Changes for T and N Staging in Patients With Pancreatic
Adenocarcinoma. *Ann Surg* **265**, 185-191 (2017).
32.   Bott, A.J. *et al.* Glutamine Anabolism Plays a Critical Role in Pancreatic Cancer by
Coupling Carbon and Nitrogen Metabolism. *Cell Rep* **29**, 1287-1298 e1286 (2019).
33.   Yuneva, M.O. *et al.* The metabolic profile of tumors depends on both the responsible
genetic lesion and tissue type. *Cell Metab* **15**, 157-170 (2012).
34.   Tardito, S. *et al.* Glutamine synthetase activity fuels nucleotide biosynthesis and supports
growth of glutamine-restricted glioblastoma. *Nat Cell Biol* **17**, 1556-1568 (2015).
35.   Issaq, S.H., Mendoza, A., Fox, S.D. & Helman, L.J. Glutamine synthetase is necessary
for sarcoma adaptation to glutamine deprivation and tumor growth. *Oncogenesis* **8**, 20
(2019).
36.   Biancur, D.E. *et al.* Compensatory metabolic networks in pancreatic cancers upon
perturbation of glutamine metabolism. *Nat Commun* **8**, 15965 (2017).
37.   Muir, A. *et al.* Environmental cystine drives glutamine anaplerosis and sensitizes cancer
cells to glutaminase inhibition. *Elife* **6** (2017).
38.   Zhao, J., Zhai, B., Gygi, S.P. & Goldberg, A.L. mTOR inhibition activates overall protein
degradation by the ubiquitin proteasome system as well as by autophagy. *Proc Natl Acad*
Sci U S A **112**, 15790-15797 (2015).
39.   Commisso, C., Flinn, R.J. & Bar-Sagi, D. Determining the macropinocytic index of cells
through a quantitative image-based assay. *Nat Protoc* **9**, 182-192 (2014).
40.   Bookout, A.L. & Mangelsdorf, D.J. Quantitative real-time PCR protocol for analysis of
nuclear receptor signaling pathways. *Nucl Recept Signal* **1**, e012 (2003).

41.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
42.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast
and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).
43.   Soneson, C., Love, M.I. & Robinson, M.D. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2015).
44.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
45.   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-
15550 (2005).
46.   Buenrostro, J.D., Wu, B., Chang, H.Y. & Greenleaf, W.J. ATAC-seq: A Method for
Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21 29 21-
29 (2015).
47.   Jadhav, U. *et al.* Dynamic Reorganization of Chromatin Accessibility Signatures during
Dedifferentiation of Secretory Precursors into Lgr5+ Intestinal Stem Cells. *Cell Stem Cell*
21, 65-77 e65 (2017).
48.   Kanarek, N. *et al.* Histidine catabolism is a major determinant of methotrexate sensitivity.
Nature **559**, 632-636 (2018).
49.   Mascanfroni, I.D. *et al.* Metabolic control of type 1 regulatory T cell differentiation by
AHR and HIF1-alpha. *Nature medicine* **21**, 638-646 (2015).
50.   Chong, J., Wishart, D.S. & Xia, J. Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. *Curr Protoc Bioinformatics* **68**, e86 (2019).
51.   Wapnir, I.L., Wartenberg, D.E. & Greco, R.S. Three dimensional staging of breast
cancer. *Breast Cancer Res Treat* **41**, 15-19 (1996).

 Figure Legends

Figure 1. PDA cells adapted to nutrient deprivation exhibit enhanced growth *in vitro* and *in*

***vivo*. a**, Schematic depicting selection of adapted (A) clones (C4-C6) derived from 8988T or

SUIT-2 human PDA cells that survived and grew in low glucose-low glutamine (L-L) medium,

and control non-adapted (NA) clones (C1-C3) derived from parental cells plated at low dilution

in high glucose-high glutamine (H-H) medium. **b**, Proliferation curves of clonal cells described

in **a** that were grown in L-L medium for 7 days (n= 5). **c**, Percent cell death in clones described

in **a** (n= 5) that were grown in L-L medium for 4 days (SUIT-2) or 5 days (8988T). **d**, Colony

formation assay showing number of colonies formed by 8988T or SUIT-2 cells described in **a**

that were grown in L-L medium for 10 days or 7 days, respectively (n= 6). **e**, Three-dimensional

(3D) growth assay showing area of spheres formed by 8988T or SUIT-2 cells described in **a** that

were grown for 10 days or 7 days, respectively, in L-L medium supplemented with 4% matrigel

(n= 6). **f**, Volumes of orthotopic xenograft PDA tumors derived from 8988T clones described in

a that were injected (75×10^4 cells) into the pancreas of 4-6 week old Rag1$^{-/-}$ mice and grown for

33 days (n=7 for NA-C1; n=6 for NA-C2 and n=7 for A-C4, 5 clones). Data represent the mean

± S.E.M. in **b**, or mean ± S.D. in **c-f**. $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$, two-

way ANOVA for **b** and one-way ANOVA for **c-f**, followed by Tukey test.

Figure 2. Adapted PDA clones display enhanced *de novo* DNA synthesis and mTORC1

signaling under nutrient-deprived conditions. a, Heatmap listing in descending order of

statistical significance ($p < 0.05$ by *t* test) metabolites in the purine and pyrimidine synthesis

pathways for non-adapted or adapted PDA clones, that were treated for 24 h with L-L medium

(n= 4 replicates per clone, except for 8988T C5: n= 3). Red indicates higher level and blue lower

level, relative to the median for each metabolite across all groups. **b**, Relative incorporation of

radiolabeled U-^{14}C-aspartate into DNA synthesis in non-adapted (NA-C) and adapted (A-C)

SUIT-2 clones treated with either L-L or H-H medium for 40 h. Data are presented as relative

fold change normalized to NA-C1 under L-L conditions and represent the mean ± S.D. (n= 3

replicates per clone). *$p < 0.05$; **$p < 0.01$, one-way ANOVA followed by Tukey test. **c**,

Immunoblots of pT389-S6K, pS65-4EBP1, pS1859-CAD and total S6K, 4EBP1, CAD in 8988T

or SUIT-2 clones treated with H-H or L-L medium for 24 h. β-actin was used as a loading

control.

Figure 3. Adapted PDA cells induce GS and have enhanced incorporation of nitrogen from

amino acids into glutamine and nucleotide synthesis. a, Schematic illustrating glutamate

synthesis from α-ketoglutarate (α-KG) and ammonia by glutamate dehydrogenase (GDH) and

glutamine synthesis from glutamate and ammonia by glutamine synthetase (GS). In red, are

ammonia nitrogens incorporating into glutamate and then glutamine. **b**, Immunoblots of GS (low

or high exposures) and GDH in 8988T or SUIT-2 clones treated with H-H or L-L medium for 24

h. β-actin was used as a loading control. **c**, Fractional labeling of ^{15}N-glutamate and glutamine in

SUIT-2 cells treated for 24 h with H-H or L-L medium supplemented with 0.8mM ^{15}N-

ammonium chloride. Data are corrected for natural abundance and represent the average of 4

replicates per clone per condition ± S.D. *$p < 0.05$; ****$p < 0.0001$, one-way ANOVA followed

by Tukey test. **d**, Schematic illustrating the incorporation of leucine nitrogen (red) into glutamate

and then aspartate via transamination reactions catalyzed by branched chain amino acid

transaminase BCAT and aspartate transaminase GOT. Glutamate is used to synthesize glutamine

via GS and aspartate nitrogen incorporates into newly synthesized nucleotides. **e**, Heatmap

listing in descending order of statistical significance ($p < 0.05$ by t test) amino acids in non-

adapted or adapted PDA clones that were treated for 24 h with L-L medium (n= 4 per clone per

condition, except for 8988T C5: n= 3). Red indicates higher level and blue lower level, relative

to the median for each metabolite across all groups. **f**, Fractional labeling of metabolites in

glutamine and nucleotide synthesis pathways in cells described in **b** grown in media

supplemented with 0.4mM ^{15}N-leucine for 24 h. Data represent the average of 4 replicates per

clone per condition ± S.D. *p < 0.05; ****p < 0.0001, one-way ANOVA followed by Tukey

test.

Figure 4. PDA adaptation to nutrient deprivation is reversible and associated with changes

in open chromatin. a, Schematic depicting the adaptation reversal process where adapted PDA

clones are grown in nutrient replete (H-H) conditions for at least 4 passages, prior to treatment

with H-H or L-L medium for 24 h. **b**, Immunoblots of pT389-S6K (low and high exposures),

pS65-4EBP1, pS1859-CAD and total S6K, 4EBP1, CAD and GS in SUIT (C5) or 8988T (C4)

clones that were reverse-adapted over 4 passages and then treated with H-H or L-L medium for

24 h. β-actin was used as a loading control. **c**, Proliferation curves of SUIT-2 or 8988T cells that

are either non-adapted (NA), adapted (A) or reverse adapted (RA) for 18 passages (SUIT-2) or

30 passages (8988T) and grown under L-L conditions for 7 days (n= 6). **d**, Percent cell death in

clones described in **c** that were grown under L-L medium for 5 days (n= 6). Data represent the

mean ± S.E.M. in **c**, and mean ± S.D. in **d**. **p < 0.01; ***p < 0.001; ****p < 0.0001, two-way

ANOVA for **c** and one-way ANOVA for **d**, followed by Tukey test. In **c**, stars indicate statistical

significance for A vs NA; A vs RA; RA vs NA; ns indicates non-significance for A vs RA (day

3) and RA vs NA (day 7) in 8988T cells. **e.** Relative mRNA levels of 1,842 genes upregulated

(>3-fold, *q* <0.05) in both SUIT-2 and 8988T adapted clones, compared to non-adapted clones;

mRNA levels of the corresponding genes in reverse-adapted (RA) clones are shown alongside.

Differentially expressed genes were identified by unsupervised k-means clustering. Red, high

expression; blue, reduced expression relative to mean expression levels for each gene across all

groups (n= 3 clones per group). **f**, ATAC-seq data showing significant (>2-fold, *q* <0.05)

changes in chromatin access at enhancers (located >500 bp from transcription start sites) in each

of 2 independent non-adapted (NA), adapted (A), and reverse-adapted (RA) 8988T or SUIT-2

clones. **g**, Correlation of changes in gene expression and nearby (<25 kb) chromatin accessibility

for the 1,842 genes induced upon adaptation to nutrient deprivation in both SUIT-2 and 8988T

cells. Each dot represents a gene, and those linked to 3,807 enhancers with altered chromatin

access in both cell lines are marked in purple. More gene-linked enhancers (<25 kb) show gains

than show losses in accessibility.

Figure 5. mTORC1 stabilizes GS protein levels under glutamine deprivation.

a, Immunoblots of GS (low and high exposures), pT389-S6K and total S6K in SUIT-2 or 8988T

adapted clones (C4-C6) that were treated with vehicle control (DMSO) or Torin 1 (40 nM) in L-

L medium for 24 h. **b**, Immunoblots of GS, pS65-4EBP1, pT389-S6K, total 4EBP1 or S6K in

cells from **a** that were treated under L-L medium with control DMSO or Torin 1 (200 nM) for a

total of 8 h, in the absence or presence of cycloheximide (CHX, 20 μg ml^{-1}) for the indicated

times. **c**, GS protein levels in non-adapted and adapted SUIT-2 or 8988T clones treated in L-L

medium with MG-132 (10μM) for the indicated times. **d**, GS protein levels in SUIT-2 or 8988T

adapted clones treated in L-L medium with either Torin 1 (200 nM) or MG-132 (10μM) alone, or

in combination for 12 h (C4) or 8 h (C5, C6). **e**, Co-immunoprecipitation (IP) blots showing the

ubiquitination status of GS in SUIT-2 or 8988T adapted clones treated in L-L medium, with

either Torin 1 (200 nM) or MG-132 (10μM) alone, or in combination for 8 h. Input refers to

immunoblots of total GS, pT389-S6K and total S6K levels in whole cell lysates. In **a-e**, β-actin

or GAPDH were used as loading controls.

Figure 6. GS is required for the adaptation-induced growth fitness of PDA cells under

nutrient deprivation. a, Proliferation curves of adapted 8988T cells transfected with

Doxycycline (Dox)-inducible shScrambled control or shGS hairpins 1 and 2, grown in L-L

medium for 7 days in the absence or presence of 1 μg ml^{-1} Dox (n= 8). **b**, Percent cell death in

cells described in **a** (n= 8) that were grown in L-L medium for 7 days. **c**, Colony formation assay

for cells in **a** that were grown for 10 days, under L-L conditions (n= 6). **d**, Weights of orthotopic

xenograft PDA tumors derived from 8988T adapted cells (C4) stably expressing Dox-inducible

hairpins described in **a** that were injected (75 x 10^4 cells) into the pancreas of 4-6 week old Rag1$^-$

$^{/-}$ mice. Dox was supplemented in the drinking water (2 mg ml^{-1}) 5 days post-injection of cells,

and tumors were harvested 33 days later (n=5 shScr -Dox; n= 7 shScr + Dox; n= 8 shGS#1; n= 7

shGS#2). Data represent the mean ± S.E.M. in **a**, and mean ± S.D. in **b-d**. **$**p < 0.01$; ***$p <$

0.001; ****$p < 0.0001$, two-way ANOVA for **a**, one-way ANOVA for **b-d**, followed by Tukey

test. **e**, Immunoblots of GS in cells used for xenografts in **d** showing decreased protein levels, 48

h following treatment with Dox (1 μg ml^{-1}) for inducible knockdown . β-actin was used as a

loading control. **f**, Model illustrating a role for GS stabilization in PDA cell adaptation to nutrient

deprivation, which leads to enhanced growth fitness. The sketch was created using

Biorender.com.

Supplementary Figure Legends

Supplementary Figure 1. Adaptation to nutrient deprivation enhances PDA cell growth potential.

A, Proliferation curves of adapted 8988T or SUIT-2 PDA clones (A-C4, -C5, -C6) and control non-adapted (NA-C1, -C2, -C3) clones that were grown in H-H medium for 7 days (n= 6). **B**, Percent cell death in clones described in **a** that were grown in H-H medium for 5 days (n= 6). **C**, Colony formation assay showing number of colonies formed by cells in **a** that were grown for 10 days (8988T) or 7 days (SUIT-2) under H-H conditions (n= 6). **D-f**, 3D growth assay showing number (**d**, **e**) or area (**f**) of spheres formed by cells in **a** that were grown for 10 days or 7 days, respectively, in L-L (**d**) or H-H (**e, f**) medium supplemented with 4% matrigel (n= 6). **G,** Representative images of mice bearing orthotopic transplants of adapted or non-adapted PDA clones. Boxed areas are enlarged on the right, demonstrating the presence of a large pancreatic tumor derived from the adapted cells and absence of tumors on the pancreas that was previously injected with non-adapted cells. Data represent the mean ± S.E.M. in **a**, and mean ± S.D. in **b-f**. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$, two-way ANOVA for **a** and one-way ANOVA for **b-f**, followed by Tukey test.

Supplementary Figure 2. Autophagy and macropinocytosis are not further induced in PDA cells upon adaptation to nutrient deprivation. A and b, Immunoblots of the autophagy marker LC3B-II in SUIT-2 (**a**) or 8988T (**b**) non-adapted (NA) or adapted (A) clones treated under L-L medium, with the autophagy inhibitor chloroquine (CQ, 10 μM) for 2, 6 or 24 h. Blots demonstrate similar or decreased autophagic flux in adapted compared to non-adapted cells. **C,** Representative images for macropinocytic uptake of TMR-dextran by non-adapted and adapted

8988T and SUIT-2 cells in the presence or absence of the inhibitor EIPA (top) with

quantification of macropinocytic area per cell (bottom). Scale bar, 25 μm. Data represent the

mean ± S.D. for n= 2 independent experiments with approximately 165 cells (8988T) or 285

cells (SUIT-2) scored per experiment. ****$p < 0.0001$, one-way ANOVA followed by Tukey

test.

Supplementary Figure 3. Adaptation to nutrient deprivation minimizes transcriptional

variance between PDA cell lines. A, Principal Component Analysis (PCA) of the 500 most

variable genes in adapted and non-adapted PDA clones; n = 3 clones per group. Samples

segregate based on cell line or PC1, which explains 83% of the variance. Segregation is

minimized upon adaptation to nutrient deprivation (PC2, 8% variance). **B**, Heatmaps (left) of

genes enriched in metabolic pathways in adapted compared to non-adapted SUIT-2 or 8988T

clones (n = 3 per group), with corresponding GSEA plots (right). Red indicates higher

expression, and blue lower expression, relative to the mean expression level for each metabolite

across all groups. NES normalized enrichment scores, FDR false discovery rate, Nom. Nominal.

Supplementary Figure 4. Nutrient-deprived adapted PDA cells display metabolite levels

closer to those under nutrient-replete conditions. a, Heatmap listing in descending order of

statistical significance ($p < 0.05$ by t test), amino acids, metabolites in the nucleotide synthesis

pathway, and metabolites in the TCA cycle, glycolysis/gluconeogenesis pathways in non-adapted

or adapted PDA clones from SUIT-2 or 8988T cells that were treated for 24 h with H-H or L-L

medium. Data represent 3 clones per condition with n= 4 replicates per clone). Red indicates

higher level and blue lower level, relative to the median for each metabolite across all groups. **B**,

List of top 27 metabolic pathways in descending order of statistical significance (-log $p \geq 2.18$;

FDR $\leq$ 10E-10) that are enriched in adapted, compared to non-adapted PDA clones from both

cell lines, under L-L medium. In bold are pathways involving metabolites in **a**. FDR, false

discovery rate.

Supplementary Figure 5. AKT phosphorylation is decreased in adapted PDA clones under

nutrient deprivation. A, Immunoblots of pS473-AKT, pT389-AKT and total AKT in non-

adapted (NA) and adapted (A) SUIT-2 clones (C1-C6) under H-H and L-L medium. GAPDH

was used as loading control.

Supplementary Figure 6. Transaminase expression is induced in adapted PDA cells under

nutrient deprivation. A and **b,** Relative mRNA levels of GS (**a**), and transaminases BCAT1/2

and GOT1/2 (**b**) in adapted (A) or non-adapted (NA) SUIT-2 and 8988T cells treated for 24 h

with either H-H or L-L medium. Data represent the mean of 3 clones per condition $\pm$ S.D., n = 3

replicates per clone. **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$, one-way ANOVA followed by

Tukey test. **C**, Levels of BCAT1/2, GOT1/2 proteins in cells described in **a** and **b** under L-L

medium. β-actin was used as loading control.

Supplementary Figure 7. Branched chain amino acids serve as a source of nitrogen for

glutamine synthesis in adapted PDA clones. A, Fractional labeling of metabolites in non-

adapted (NA) or adapted (A) 8988T clones (C1-C6) treated with L-L medium supplemented with

0.4mM ^{15}N-leucine for 24 h. Data represent the average of 4 replicates per clone per condition $\pm$

S.D. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$, one-way ANOVA followed by

Tukey test.

Supplementary Figure 8. Reverse adaptation of PDA cells to nutrient deprivation results in

reversal of enhanced growth and maintenance of the adaptation-induced chromatin

accessibility changes. a, **b** Immunoblots of pT389-S6K (low and high exposures), pS65-4EBP1,

pS1859-CAD and total S6K, 4EBP1 and GS in adapted SUIT-2 (**a**) or 8988T (**b**) cells that were

reverse-adapted over at least 4 passages and then treated with H-H or L-L medium for 24 h.

GAPDH or β-actin was used as a loading control. **c**, Proliferation curves of SUIT-2 and 8988T

cells that are either non-adapted (NA), adapted (A) or reverse-adapted (RA) for 18 passages

(SUIT-2) or 30 passages (8988T) and grown under L-L conditions for 7 days (n= 6). Data

represent the mean ± S.E.M. *$p < 0.05$; **$p < 0.01$; ****$p < 0.0001$, two-way ANOVA followed

by Tukey test, and indicate significant differences for A vs NA; A vs RA; RA vs NA; ns, non-

significant for RA vs NA (day 3, SUIT-2); RA vs NA (day 5 and day 7, 8988T).). **d**, Overlap of

enhancers with altered chromatin accessibility in adapted 8988T and SUIT-2 cells identifies

3,807 sites changed in both cell lines. **e**, Gains and losses in chromatin accessibility during

adaptation (NA, non-adapted; A, adapted) of 8988T and SUIT-2 cells, revealing significant

correlation (BETA analysis) with increases and decreases, respectively, in gene expression. **f**,

Representative loci showing concordant changes in ATAC-seq and RNA-seq signals at *POLE2*

and *RRM2*. Dotted boxes demarcate enhancer elements linked to genes that show gain or loss of

chromatin accessibility and concordant changes in gene expression in cells adapted to nutrient

deprivation. Reversal of chromatin accessibility changes upon reverse adaptation was infrequent

(e.g., *RRM2*); enhancers that gained chromatin accessibility upon adaptation generally remained

accessible, despite partial reversal in mRNA expression (e.g. *POLE2* in SUIT-2 cells).

Supplementary Figure 9. Protein levels of glutamine synthetase are decreased upon Torin 1

treatment at a post-translational level. a, Relative mRNA levels of GS in SUIT-2 and 8988T

adapted clones (C4, C5, C6) treated with L-L medium in the absence or presence of Torin 1 (200

nM) for 24 h. Data represent the mean ± S.D. (n= 3 replicates per clone per condition). **$p <$

0.01; ***$p < 0.001$; ****$p < 0.0001$, one-way ANOVA followed by Tukey test.

b, Immunoblots of GS, pT389-S6K and total S6K in cells from **a** that were treated under L-L

medium with control DMSO or Torin 1 (200 nM) for a total of 8 h, in the absence or presence of

cycloheximide (CHX, 20 µg ml^{-1}) for the indicated times prior to cell lysis. β-actin was used as

loading control.

Supplementary Figure 10. Inhibition of glutamine synthetase suppresses the growth of

PDA cells adapted to nutrient deprivation. a, Fold change in cell number of 8988T or SUIT-2

adapted clones, 72 h following treatment with the GS inhibitor (MSO) compared to vehicle

(water) control, showing decrease in cell number upon GS inhibition under L-L but not H-H

conditions (n= 5 per condition). **b**, Proliferation curves of adapted 8988T cells transfected with

control non-targeting siRNA or siGS, that were grown in L-L medium for 4 days (n= 6). **c**,

Colony formation assay for cells described in b that were grown for 8 days, under L-L conditions

(n= 6). **d**, Immunoblots showing GS levels in cells from **b**, 24h post-transfection of siRNA. **e**,

Volumes quantified by ultrasound on day 29, of orthotopic PDA xenograft tumors derived from

8988T clonal adapted cells (C4) stably expressing Doxycycline (Dox)-inducible hairpins for GS

or control Scrambled hairpin. Cells (750 x 10^3) were injected on day 0 into the pancreas of 4-6

week old Rag1$^{-/-}$ mice and Dox was supplemented in the drinking water (2 mg ml^{-1}) on day 5

(n=5 shScr -Dox; n= 7 shScr + Dox; n= 8 shGS#1; n= 7 shGS#2). **f**, Representative ultrasound

images on day 29 of adapted 8988T orthotopic xenograft tumors described in **e**. Data represent

the mean ± S.D. in **a, c-e**, and mean ± S.E.M. in **b**. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p <$

0.0001, one-way ANOVA for **a** and **e,** two-way ANOVA for **b**, followed by Tukey test, and

Student's t-test for **c**.

Supplementary Figure 11. Glutamine synthetase average protein levels do not associate

with human PDA tumor grade or patient survival. a, Kruskal-Wallis test for associations

between GS level (H-score) and tumor size (pT), lymph node involvement (pN), or grade (tumor

differentiation level) at time of resection. Low grade indicates well-differentiated tumors;

Intermediate grade, moderately differentiated tumors; High grade, poorly differentiated tumors.

b, Representative images of human normal pancreatic parenchyma with GS staining in acinar

cells (red arrowhead), duct cells (black arrowhead), islets (black arrows) and acinar ductal

metaplasia (ADM, circled). Staining in human PDA shows no correlation of GS with tumor

grade, as many lower-grade tumors display high GS protein levels and higher-grade tumors

display low GS levels. **c**, Kaplan Meier survival curves stratified by GS level, categorized as

tertiles. Follow-up period is indicated in months. Median survival is slightly shorter for patients

with low GS compared to patients with medium or high GS levels. However, the differences

between the survival curves are not statistically significant, based on the Log-Rank test ($P =$

0.18). These results are consistent with a multivariate Cox proportional hazard regression model

that adjusted for age, sex, and tumor grade.

Figure 1

Figure 2

a

b

Figure 3

Figure 4

Figure 5

Figure 6

a

b

Cell Proliferation

shScr

shGS #1

shGS #2

Cell Death

c

Colony formation

d

Tumor Weight

e

f

Adapted

mTORC1

Non-Adapted

Glu

Gln

Decreased Growth Fitness

Enhanced Growth fitness

Supplementary Figure 1

Supplementary Figure 2

a

b

c

Supplementary Figure 3

a

b

Supplementary Figure 4

a

b

Pathways	-log (p)	FDR
Porphyrin and chlorophyll metabolism	5.49E+01	8.85E-23
Phenylalanine metabolism	5.34E+01	2.02E-22
Ubiquinone and other terpenoid-quinone biosynthesis	5.14E+01	9.56E-22
Valine, leucine and isoleucine degradation	**4.95E+01**	**4.57E-21**
Valine, leucine and isoleucine biosynthesis	**4.74E+01**	**3.14E-20**
Aminoacyl-tRNA biosynthesis	4.60E+01	1.07E-19
Phenylalanine, tyrosine and tryptophan biosynthesis	4.44E+01	4.67E-19
Tryptophan metabolism	4.16E+01	6.48E-18
Cyanoamino acid metabolism	4.06E+01	1.52E-17
Sulfur metabolism	3.81E+01	1.64E-16
Arginine and proline metabolism	3.78E+01	2.19E-16
Tyrosine metabolism	3.65E+01	6.84E-16
Propanoate metabolism	3.64E+01	6.84E-16
Nitrogen metabolism	3.62E+01	8.46E-16
Glycine, serine and threonine metabolism	3.58E+01	1.11E-15
Cysteine and methionine metabolism	3.54E+01	1.59E-15
Thiamine metabolism	3.50E+01	2.26E-15
Nicotinate and nicotinamide metabolism	3.40E+01	5.59E-15
Pantothenate and CoA biosynthesis	3.35E+01	8.99E-15
Sphingolipid metabolism	3.23E+01	2.84E-14
Methane metabolism	3.02E+01	2.10E-13
Pyrimidine metabolism	**2.93E+01**	**5.26E-13**
Purine metabolism	**2.64E+01**	**8.51E-12**
Citrate cycle (TCA cycle)	**2.39E+01**	**1.01E-10**
Histidine metabolism	2.34E+01	1.61E-10
D-Arginine and D-ornithine metaoblism	2.26E+01	3.56E-10
Glycolysis and Gluconeogenesis	**2.18E+01**	**7.84E-10**

Supplementary Figure 5

a

Supplementary Figure 6

a

b

c

Supplementary Figure 7

a

Supplementary Figure 8

Supplementary Figure 9

a

GS

SUIT-2

8988T

b

Supplementary Figure 10

Supplementary Figure 11

a

TMA by pT, pN and grade

Variable	Level	N	Median H-Score	P-value
pT	1	14	150	0.61
	2	89	133	
	3	24	150	
pN	0	32	150	0.34
	1	62	125	
	2	33	150	
Grade	Low	6	195	0.32
	Intermediate	67	138	
	High	52	142	

b

Grade

Normal Parenchyma / Low / Intermediate / High

c

Kaplan Meier Survival Curves by GS Level

Log-Rank *P* = 0.18

GS level H-score	N (%)	Median Survival	95% CI
Low (1)	44 (35)	20	(15, 34)
Medium (2)	43 (34)	36	(18, 86)
High (3)	40 (32)	35	(23, 57)

Survival Probability

Follow-up in Months

H3 — 1 — 2 — 3

+ Censored