Title: The tRNA pseudouridine synthase TruB1 regulates the maturation and function of

let-7 miRNA

Ryota Kurimoto[1],Tomoki Chiba[1], Yoshiaki Ito[1,2], Takahide Matsushima[1], Yuki Yano[1],

Kohei Miyata[3], Tsutomu Suzuki[4], Kozo Tomita[5], Hiroshi Asahara[1,6,7, *].

[1]Department of Systems BioMedicine, Graduate School of Medical and Dental Sciences,

Tokyo Medical and Dental University (TMDU), Bunkyo-ku, Tokyo 113-8520, Japan;

[2]Research Core, Research Facility Cluster, Institute of Research, Tokyo Medical and

Dental University (TMDU), Bunkyo-ku, Tokyo 113-8520, Japan; [3]Department Obstetrics

and Gynecology, Faculty of Medicine, Fukuoka University, Fukuoka 814-0180, Japan.;

[4]Department of Chemistry and Biotechnology, Graduate School of Engineering,

University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan; [5]Department of Computational

Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of

Tokyo, Kashiwa, Chiba 277-8562, Japan; [6]Department of Molecular and Experimental

Medicine, The Scripps Research Institute, San Diego, CA, USA. [7]Lead Contact.

Corresponding author: Hiroshi Asahara

Address: 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, JAPAN

Telephone number: (81)-03-5803-5015

Fax number: (81)-03-5803-5810

E-mail address: asahara@scripps.edu or asahara.syst@tmd.ac.jp

Running title: TruB1 enhances let-7 processing

Abstract:

Let-7 is an evolutionary conserved microRNA that mediates post-transcriptional gene silencing to regulate a wide range of biological processes, including development, differentiation, and tumor suppression. Let-7 biogenesis is tightly regulated by several RNA-binding proteins, including Lin28A/B, which represses let-7 maturation. To identify new regulators of let-7, we devised a cell-based functional screen of RNA-binding proteins using a let-7 sensor luciferase reporter, and identified the tRNA pseudouridine synthase, TruB1. TruB1 enhanced maturation specifically of let-7 family members. Rather than inducing pseudouridylation of the miRNAs, HITS-CLIP (High throughput sequencing crosslinking immunoprecipitation) and biochemical analyses revealed direct

binding between endogenous TruB1 and the stem-loop structure of pri-let-7, which also

binds Lin28A/B. TruB1 selectively enhanced the interaction between pri-let-7 and the

microprocessor DGCR-8, which mediates miRNA maturation. Finally, TruB1 suppressed

cell proliferation, which was mediated in part by let-7. Altogether, we reveal an

unexpected function for TruB1 in promoting let-7 maturation and function.

Key words

microRNA/ let-7/ RNA binding protein/ TruB1/ pseudouridine.

Introduction

MicroRNAs (miRNA) are short non-coding RNAs of about 22 bases that

regulate gene expression by inhibiting either the stability of target mRNAs or protein

synthesis in association with Argonaute (AGO) family proteins (Lee et al., 1993;

Wightman et al., 1993). Lin-4 and let-7 were the first reported miRNAs in *C. elegans*

(Lee et al., 1993). Since then, more than 1,500 conserved miRNAs have been identified

across many species including mammals (Ambros., 2004). The expression and

maturation of miRNAs are tightly regulated by multi-step mechanisms involving

protein-RNA associations with enzymatic activities, and molecular shuttling, specifically:

i) Pol-II mediated primary miRNA (pri-miRNA) transcription (Lee et al., 2004; Cai et al.,

2004), ii) Drosha/ DiGeorge Syndrome Critical Region 8 (DGCR8)-mediated cleavage

processing producing a precursor-miRNA (pre-miRNA) in the nucleus (Denli et al.,

2004; Gregory et al., 2004; Han et al., 2004; Landthaler et al., 2004; Lee et al., 2003;

Zeng et al., 2005), iii) nuclear export of pre-miRNA by Exportin-5 (Yi et al., 2003;

Bohnsack et al., 2004; Lund et al., 2004), iv) additional cleavage by the RNase III

enzyme Dicer in the cytoplasm (Bernstein et al., 2001; Grishok et al., 2001; Hutvágner

et al., 2001; Ketting et al., 2001; Knight and Bass, 2001), and v) incorporation into

the RNA-induced silencing complex (RISC) with AGO family proteins to generate

the final mature miRNA (Hammond et al., 2001; Mourelatos et al., 2002; Tabara et al.,

1999). Although the first transcription step for pri-miRNA expression can be driven by

transcription factors and RNA Pol-II as well as mRNA, the following maturation steps,

ii-v), are also critical and unique to miRNA biogenesis and dynamics.

The comparison of gene expression in tumor tissues showed discrepancies in

the levels of mature miRNAs and their pri-miRNAs (Thomson et al., 2006). In particular,

mutually exclusive and reciprocal regulation between the let-7 miRNA and the RNA

binding protein (RBP) Lin-28 at these maturation process have been well demonstrated

in ES cells, *C.elegans*, and cancer pathology (Newman et al., 2008; Piskounova et al.,

2008; Heo et al., 2008; Viswanathan et al., 2008; Chang et al., 2009). Lin28A/B

specifically binds to the preE loop sequence of pri- / pre-let-7 and suppresses the

processing of let-7 by causing oligo-uridylation of the let-7 precursor through TUT4/7

activity (Heo et al., 2008; Piskounova et al., 2011). Based on these fundamental findings,

the factors mediating let-7 multi-step regulation, which may be critical for various

biological and pathological events, have been extensively explored by several

approaches (Newman et al., 2008; Trabucchi et al., 2009; Treiber et al., 2017);

however, to date, only a few factors have been shown to regulate let-7 family-specific

maturation (Newman et al., 2008; Trabucchi et al., 2009; Michlewski et al., 2010;

Choudhury et al., 2014; Pandolfini et al., 2019).

Taking advantage of the completion of genome-wide comprehensive full-length cDNA

libraries (Iourgenko et al., 2003; Chanda et al., 2003; Conkright et al., 2003; Huang et

al., 2004; Liu et al., 2005), we and others successfully developed cell-based screening

systems to quantify miRNA targets and functions, including destabilization of target

mRNA and suppression of protein synthesis (Wolter et al., 2014: Wolter et al., 2015; Ito

et al., 2017). Here we applied the above strategy and introduced a new cell-based

screening to identify genes that regulate let-7 miRNA by using a luciferase reporter

assay with a set of expression plasmid libraries mainly encoding proteins with RNA

binding properties. We identified a new regulatory mechanism: TruB1, an

RNA-modifying enzyme, selectively regulates let-7 activity in an enzymatic

activity-independent manner.

Results

Functional screening to detect RNA binding proteins (RBPs) that promote let-7 expression

We developed a cell-based, functional screen using an expression plasmid library and a let-7 sensor reporter to identify proteins that regulate let-7 biogenesis. This type of screen enables the identification of proteins that regulate endogenous let-7 function, as opposed to other aspects of let-7 biology, such as binding, which is assayed using affinity-based (pull-down) screens. The gain-of-function format also enables the identification of regulatory factors that may cause lethality under loss of function conditions, such as with Crispr-Cas9 or RNAi-KD screens. The let-7 sensor reporter was designed to monitor let-7 expression levels by inserting a let-7a target site into the luciferase 3'UTR region driven by the SV-40 promoter in pLuc2 plasmids (Fig 1 A). Thus, luciferase activity is repressed in the presence of mature let7 miRNA (Fig EV1). In this screen, we focused on identifying molecules directly regulating the miRNA maturation process, thus we prepared a sib-selection library covering molecules annotated with RNA binding protein (RBP) features from the Center for Cancer systems Biology

(CCSB)-Broad Lentiviral over-expression library (Yang et al., 2011). As many

zinc-finger-proteins have not been directly tested for their RNA or DNA binding

preference, the genes with zinc-finger domains were also included in the sib-selection

library. Ultimately, 1469 genes were selected and prepared for the screening. This

library was transfected into HEK293FT cells, along with the let-7 sensor reporter. If let-7

maturation is inhibited by coexpression of one of the cDNAs in the RBP library,

expression of the luciferase reporter would be induced. As a positive control for the

assay, we over-expressed Lin28A from the same backbone plasmid transfection along

with the let-7 sensor reporter with or without overexpression of pri-let-7a in HEK293FT

cells. This significantly promoted luciferase activity reflecting reduction of endogenous

mature let-7 expression (Fig. 1B). Next, we screened the 1,469 genes in our library

utilizing the luciferase assay with the let-7 sensor reporter in HEK293FT cells seeded

into 384 well plates. pRL-SV40 Renilla luciferase activity was used as a transfection

efficiency control. We only included genes that did not change the luciferase activity of

the let-7 sequence minus the reporter (0.5 <, > 2.0) (results are shown in Appendix

Table S1). Of these, we found that over-expression of 259 genes significantly reduced

relative luciferase activity (< 0.50） compared with the GFP expression plasmid control

(Fig. 1D), indicating that they potentially promote let7 miRNA maturation. From the top

five hits, we performed a larger scale (96 well plate) luciferase reporter assay for

validation, whereby four of the top five genes reproducibly repressed the luciferase

activity of the let-7 sensor (Fig. 1E). To confirm the effect on endogenous let-7a

expression, we examined the expression levels of a panel of ten miRNAs including

let-7a by qPCR. Consistent with the screening results, overexpression of all of the top

five candidate genes significantly promoted endogenous mature let-7a expression. Only

TruB1 selectively induced let-7 expression but did not increase expression of the other

miRNAs, whereas the other four hits, SF3A3, LARP7, GLTSCR2, and EF1E1, also

increased the other miRNAs (Fig 1F). This suggests the specific promotion of let-7

biogenesis by TruB1. TruB1 is among the RNA-modifying enzymes that mediate

pseudouridylation of tRNA. TruB1 has also recently been shown to regulate mRNA via

pseudouridylation (Schwartz et al., 2014; Safra et al., 2017), but, to date, there has

been no reported function of TruB1 in miRNA biogenesis.

TruB1 selectively promotes the expression of let-7 families

Although a few proteins, such as Lin28A/B, KSRP, hnRNPA1, have been

shown to be involved in the biogenesis of a specific set of miRNAs, most other miRNA

regulators act without specificity, or their miRNA selectivity remains unclear. To test

whether TruB1-dependent let-7 promotion is specific for the let-7 family, as our

preliminary results suggested, we comprehensively analyzed the effect of TruB1

knockdown by siRNA on miRNA biogenesis in HEK293FT cells by TaqMan array. TruB1

knockdown down regulated the expression of five mature let-7 family members (let-7a,

b, c, f and g) (Fig 2A, Fig EV2A, Appendix Table S2). These data were further confirmed

by large scale TaqMan PCR for miRNA and northern botting (NB), revealing that all nine

endogenous mature let-7 family genes were significantly downregulated upon TruB1

knockdown(Fig 2B-D). In contrast, TruB1 knockdown increased the expression levels of

endogenous, immature, primary-let7 (Fig 2E). We confirmed these findings in both A549

and HeLa cells (Fig EV2B,C). These results indicate that TruB1 promotes let-7 family

miRNA biogenesis specifically at their maturation step, rather than at the transcriptional

level.

TruB1 promotes miRNA processing independently of its enzymatic activity

Therefore, we analyzed whether let-7 regulation by TruB1 is dependent on the

pseudouridine enzyme activity of TruB1. Critical residues involved in enzymatic activity

(D48, D90) and RNA binding ability (K64) have been identified in *E. coli* TruB (Wright et

al., 2011; Friedt et al., 2014; Keffer-Wilkes et al., 2016). Given the TruB amino acid

sequences for these enzyme activity and RNA binding ability are highly conserved

(Zucchini et al., 2003), we were able to insert mutations to modify the function of TruB1

by amino acid substitution. We generated two TruB1 mutants: Mt1, with inactivated

enzyme activity, and mt2, with suppressed RNA binding ability (Fig 3A). An *in vitro*

enzymatic activity assay with a tRNA^{phe} substrate using the wt, mt1 and mt2

recombinant proteins showed that the pseudouridylation enzyme activities of both mt1

and mt2 were completely attenuated (Fig 3B). An EMSA study using recombinant

proteins revealed a physical interaction between tRNA^{phe} and wt TruB1 and mt1, but not

with mt2 (Fig EV3A). Next, we tested the function of these mutants in cells. Western

blotting (WB) revealed that all proteins were overexpressed largely to the same levels

(Fig EV3B). Overexpression of Wt and mt1 TruB1 significantly increased let-7a

maturation, whereas mt2 did not (qPCR, NB). These results suggest that promotion of

let-7 maturation by TruB1 is independent of its enzyme activity, but dependent on RNA

binding (Fig 3C-E).

Next, we performed an in vitro enzyme assay to verify that let-7 could not be

pseudouridylated by recombinant TruB1. Pseudouridine of tRNA (Phe) was clearly

increased in the assay, whereas pseudouridylation of pri-let-7 and another primary

miRNA was not observed (Fig 3F). We also examined the presence of pseudouridine

directly by treatment with CMC (N-cyclohexyl-N0 -(2-morpholi- noethyl)carbodiimide

metho-p-toluenesulfonate) followed by a primer extension assay. Although we could

detect known pseudouridine sites in the tRNA, no pseudouridine site was found in let-7

(Fig 3G, Fig EV3C). We also synthesized RI-labeled pri-let-7a1 in which pseudouridine

was randomly introduced using UTP : pseudouridine at a ratio of 1:1. An in vitro

processing assay using this labelled pri-let-7a1 showed that the presence of

pseudouridine in pri-let-7a1 did not affect its processing (Fig EV3D,E). These results

indicate that the regulation of let-7 by TruB1 does not depend on its enzyme activity.

TruB1 binds directly to primary-let-7

To determine how TruB1 regulates let-7 processing, we first tested whether

TruB1 binds to pri-let-7. RNA immunoprecipitation (RIP) assays revealed that

overexpression of wt and mt1 TruB1 immunoprecipitated pri-let-7a1, whereas mt2 did

not (Fig 4A). An EMSA study using recombinant proteins revealed a physical interaction

between pri-let7a1 and wt TruB1, but not with mt2 (Fig 4B). Furthermore, EMSA was

performed using mutant RNA in which the loop structure of pri-let-7a1 was modified

(loop mt). As a result, no binding to TruB1 was observed in the loop mt (Fig 4B, Fig

EV4A). To comprehensively survey the RNA-protein binding property under

physiological conditions, we performed high-throughput cross-linking

immunoprecipitation (HITS-CLIP) using a cell line in which the 3xFLAG tag was

knocked into the TruB1 N-terminal end by using the CRISPR/Cas9 system (Fig 4C, Fig

EV4B-F). This revealed the expected, but not previously shown, direct interaction

between TruB1 and its known substrate tRNA (positive control). We also observed a

physical interaction between TruB1 and pri-let-7a1 (Fig 4D, Appendix Table S3). Direct

binding sites were also found in other miRNA sequences, including miR-29b, miR-139

and miR-107 whose expression levels of their mature forms were also specifically

decreased upon TruB1 KD in the TaqMan array (Fig 4E, Appendix Table S3). Looking at

the entire mapped reads, mRNA, lncRNA, and miRNA were detected in addition to

tRNA (Fig 4F). It has been reported that mRNA is modified by TruB1, which is consistent

with this result (Schwartz et al., 2014; Safra et al., 2017). When a read mapped to tRNA

was extracted and motif analysis was performed, the sequence was similar to the

pseudouridylation site reported for *Saccharomyces cerevisiae* PUS4 (Fig 4G), which is

homologous with TruB1 (Fig 4H) (Becker et al., 1997). Although reads mapped to

let-7a1 bound mainly near the terminal loop, no sequence similar to the tRNA motif was

found in the let7 loop sequence. tRNA and let-7 are similar in that they form a loop

structure (Fig 4I). Thus, we considered that TruB1 binds to let-7 in a

structure-dependent manner. Interestingly, the TruB1 binding site on pri-let-7a1 also

contained the sequence (GGAG), which is a binding motif of the CCHC-zinc domain of

Lin28A / B. This suggests that TruB1 and Lin28 both bind to the same stem loop

structure on *let-7*, and may compete for binding.

TruB1 enhances the affinity between microprocesser and primary let-7 and promotes microprocessing

The above results indicated that TruB1 directly binds to pri-let-7a1 under physiological conditions. To elucidate the function of this molecular interaction, we analyzed whether TruB1 was involved in the microprocessing step of miRNA biogenesis using an in vitro processing assay. RI-labeled pri-let-7a1 was incubated with a cell extract obtained from HEK293FT cells overexpressing different forms of TruB1. We found that the extract from cells overexpressing TruB1 promoted maturation of pri-let-7a1 into pre- / mature-let-7a, whereas the extract from cells overexpressing mutant TruB1(mt2) did not. We also found that knockdown of endogenous TruB1 by siRNA inhibited maturation of pri-let-7a (Fig. 5A, B). These results suggest that let-7a biogenesis is promoted at the level of conversion from the pri-miRNA to the pre-miRNA. DGCR-8 is the microprocessor involved in the cleavage of pri-miRNA in the nucleus to form pre-miRNA. Thus, we tested the effect of TruB1 on the interaction between pri-let-7 and DGCR-8 by RIP affinity assay using an antibody to endogenous DGCR-8.

Pri-let-7a1 was efficiently precipitated with DGCR-8, as expected. However, TruB1 KD

by siRNA treatment reduced this interaction between pri-let-7a1 and DGCR-8 (Fig 5C),

without affecting the expression level of DGCR-8. These data suggest that TruB1 is

involved in enhancing complex formation between DGCR-8 and pri-let-7a1. Next, we

tested the potential interaction between TruB1/let-7 and Lin28B, which also binds to the

pri-let-7a1 stem loop region recognized by TruB1. In Lin28B knockdown cells, the ratio

of TruB1-bound pri-let-7a1 over total pri-let-7a1 was increased, without altering TruB1

protein levels, as assessed by RIP (Fig 5D, Fig EV5). Thus, Lin28B inhibits the binding

between TruB1 and pri-let-7a1.

TruB1 suppresses cell growth through the promoting let-7

Finally, we examined the cellular function of this new TruB1/let-7 axis by

analyzing cell growth. Let-7 coordinates cell proliferation by targeting a set of genes,

including KRAS, which is a proto-oncogene (Johnson et al., 2005; Wang et al., 2013).

To test whether TruB1 can affect KRAS expression via let-7 regulation, we generated a

Luc-reporter plasmid comprising the let-7 target region of KRAS in its 3'UTR, and

performed a luciferase reporter assay (Fig 6A). As a result, overexpression of wt TruB1

suppressed the luciferase activity of the KRAS reporter, whereas mt2 did not (Fig 6B).

Next, the effect of TruB1 overexpression on HEK-293FT cell growth was evaluated by

real-time glo assay. Cell proliferation was significantly suppressed by TruB1

overexpression. To test whether this was mediated via let-7, we introduced RNAi to

target the common seed sequence in the let-7 family. KD of the let-7 family partially

rescued the suppressive effect of TruB1 overexpression on cell proliferation (Fig 6C, D,

Fig EV6A). Taken together, these results indicate that cell proliferation can be regulated

by the TruB1 and let-7 molecular cascade.

Discussion

In this study, we performed a systematic cell-based gain of function screen by combining an arrayed overexpression plasmid library and reporter system detecting mature let-7 expression and discovered a new function for TruB1, which is a pseudouridine RNA-modifying enzyme.

To date, various genes regulating miRNA maturation have been identified as co-immunoprecipitated factors from biotin-labeled miRNA hairpin, pri-miRNAs or microprocessors (Chendrimada et al., 2005; Lee et al., 2006; Auyeung et al., 2013; Treiber et al., 2017); however, only a few of them have shown selectivity for miRNA biogenesis. In this study, we could identify a set of molecules regulating let-7 biogenesis. Our strategy to use a cell-based and functional assay with an arrayed plasmid library has the following advantages: 1. The identified candidates are screened by their ability to regulate endogenous let-7 expression. Thus, the molecular function of the candidate proteins identified is already ensured, compared to a molecular affinity-based screening strategy, which only identifies binding. 2. The use of a prepared set of genes from the library enables us to perform a quantitative and more efficient screen. All the raw data

obtained in this approach is informative. 3. The gain of function approach can identify

factors that are necessary for cell survival, whereas loss of function screens, such as

those using shRNA or CRISPR/Cas9, may be unable to detect them. This is particularly

relevant when performing screens with molecules performing fundamental functions,

such as let-7.

In our approach, we identified a set of proteins that significantly enhance endogenous

let-7 expression. Among them, TruB1 is shown to play a critical role in let-7 specific

miRNA maturation, whereas the other candidates did not show such selectivity.

TruB1 is a pseudouridine synthase (Zucchini et al., 2003), which is responsible for

pseudouridylation, the first RNA modification found in tRNA and ribosomes (Cohn and

Volkin., 1951; Charette and Gray., 2000). Pseudouridylation is thought to be required for

modifying RNA structure by enhancing base-to-base stacking (Charette and Gray,

2000）. TruB1 introduces pseudouridine at position 55 of tRNAs during the early stage of

tRNA maturation (Becker et al., 1997; Zucchini et al., 2003). TruB1 mediated-

pseudouridine is also found in mRNA (Schwartz et al., 2014; Carlile et al., 2014; Li et al.,

2015; Safra et al., 2017). Notably, our CLIP analysis for endogenous TruB1 revealed a

physical association of TruB1 with a series of mRNAs, although the functional

significance of these interactions and the potential pseudouridylation were not

determined. *E. coli* TruB has been reported to function as a chaperone for tRNA that is

not mediated by its enzymatic activity (Keffer-Wilkes et al., 2016). This reflects our

finding that let-7 maturation promoted by TruB1 is also independent of its intrinsic

pseudouridylation enzyme activity. It would thus be interesting to test whether the

interactions we observed between TruB1 and mRNA are also independent of its

pseudouridylation enzyme activity.

Examination of the crystal structures of bacterial TruB and tRNA reveals that the base

of U55 in the TUC stem-loop structure near position 55 of the tRNA is flipped to bind to

the catalytic site of TruB (Hoang and Ferré- D'Amaré., 2001). In our HITS-CLIP, TruB1

was also bound to the stem-loop structure of let-7. Although the stem loop also exists in

other miRNAs, only the let-7 stem loop is selectively regulated by binding of regulatory

factors such as Lin28 and KSRP (Newman et al., 2008; Trabucchi et al., 2009).

Consistent with this unique property of the let-7 the stem loop, TruB1 also recognizes

this region, which may cause physical competition among these specific regulators.

More detailed analyses are needed to determine the structural effects of TruB1 binding

to let-7 and how it affects the microprocessing step of let-7 maturation.

As a cell biological function of the TruB1 and let-7 interaction, we found that TruB1

controlled cell proliferation by promoting let-7 maturation. The expression level of TruB1

correlates with cancer progression and/or oncogenesis in prostate cancer and

pancreatic cancer, suggesting that TruB1 could be involved in the pathogenesis of

cancer (Fig EV6B-D) (Edgar et al., 2002; Data ref: Edgar et al., 2002; Varambally et al.,

2005; Data ref: Varambally et al., 2005; Pei et al., 2009; Data ref: Pei et al., 2009; Wang

et al., 2015; Data ref: Wang et al., 2015). On the other hand, TruB2, which has the same

enzyme activity site at TruB1, preferentially mediates pseudouridinylation of

mitochondrial RNA, and is essential for cell survival by controlling glycolysis (Arroyo et

al., 2016; Antonicka et al., 2017). TruB KO *E. coli* were weaker than wild-type cells

when they were co-cultured; however this was not caused by decreased survival

(Gutgsell et al., 2000). In this way, despite having highly conserved enzyme activity

sites between species and homologs, their functions appear to be quite diverse.

PUS7, PUS10, DKC1, TruB1, TruB2, amongst others, are also pseudouridine

synthases, and among them, PUS10 was recently reported to promote miRNA

maturation also independently of its enzyme activity (Song et al., 2019). PUS10

introduces pseudouridine at position 55 of tRNA (Roovers et al., 2006), similar to TruB1,

and position 54 of tRNA in archaea (Deogharia M, et al. RNA. 2019). PUS10 increases

the affinity of microprocessors for various miRNAs and promotes their maturation (Song

et al., 2019). However, these two pseudouridine synthases showed very different

substrate selectivity. While TruB1 selectively and specifically interacts with pri-let-7 and

promotes let-7 expression, PUS10 appears to non-specifically increase maturation of

various miRNAs (Song et al., 2019). Consistent with this, our endogenous HITS-CLIP

for TruB1, and the published PUS10 PAR-CLIP analysis, reveal that TruB1 binds to the

terminal loop of let-7, while PUS10 binds to a site relatively far from the terminal loop of

miRNAs (Song et al., 2019). Furthermore, TruB1 acts suppressively on cell proliferation

due to the selective promotion of let-7 maturation, whereas PUS10 tended to promote

cell proliferation (Song et al., 2019). Thus, TruB1 functions differently to PUS10.

   TruB is known to be an RNA modifying enzyme that possesses two distinct molecular

functions; RNA folding and subsequent RNA modification. There are other such

examples of RBPs with dual functions in addition to their intrinsic enzymatic activity (i.e.,

ADRA1 / 2, METTL3, METTL16) (Ota et al., 2013; Lin et al., 2016; Pendleton et al.,

2017). Our strategy and data reveal an unexpected function of TruB1 in miRNAs

biogenesis with let-7 specificity, and present a new aspect of miRNA regulation by RNA

binding proteins.

Materials and Methods

Plasmid construction

The pLuc2-KAP-MCS vector was generated by a previously reported method (Ito et al., 2017). To create the let-7 sensor vector, the chemically synthesized let-7 complementary sequence was annealed and inserted between the EcoRI and XhoI sites. We used the empty pLus2-KAP-MCS vector as the control sensor (-) vector. The KRAS mRNA vector was generated by inserting the let-7a target site of Kras mRNA into pLuc2-KAP-MCS at the EcoR1 and Kpn1 sites. The sequence encoding human TruB1 ORF was cloned by PCR from 293FT cell cDNA. pcDNA-TruB1 was generated by inserting the ORF of human TruB1 with a Flag sequence into pcDNA.3.1(+) (Thermo Fisher Scientific) at the Nhe1 and EcoR1 sites. The pCLT vector was designed and created by modifying pCS 2 (RIKEN) using the tet-on system. The sequence is shown in Appendix Table S5. pCLT-TruB1 and pCLT-GFP were generated by inserting the ORF of human TruB1 and GFP with a Flag sequence into the pCLT vector at the Nhe1 and EcoR1 sites. pCLT-mt1 and mt2 were generated by PrimeSTAR Mutagenesis Basal Kit (Takara). pET22b-TruB1, mt1 and mt2 were generated by inserting the ORF of human TruB1 and its mutants into the pET22b vector (Merck Millipore) at the NdeI and XhoI

sites. pcDNA-pri-let-7a1 was generated by inserting the human pri-let-7a1 sequence

into pcDNA.3.1(+) at the BamH1 and Not1 sites. Loop mt was generated by inverse

PCR from pcDNA-pri-let-7a1. The nucleotide sequences of the primers used for the

PCR are shown in Appendix Table S4.

Cell culture.

HEK293FT, A549 and HeLa cells were maintained in DMEM medium (Corning)

supplemented with 10% FBS (Gibco) and 1% penicillin–streptomycin (Wako) at 37 °C

with 5% CO_2. siRNAs [(TruB1 (NM_139169.3), TruB2 (NM_015679.1), or negative

controls (SN-1002); Bioneer), (LIN28B (s52477) or negative control (AM4611); Ambion,

Life Technologies) and 2'-O-methylated antisense inhibitor (let-7 family or negative

control: detailed sequences shown in Appendix Table S4, Nippon-shinyaku) were

transfected into cells using Lipofectamine RNAiMAX (Thermo Fisher Scientific) using

the Manufacturer's recommended protocol. For transient overexpression, pcDNA

vectors (pcDNA-TruB1-Flag) were transfected with FuGENE HD (Promega) into cells

according to the Manufacturer's instructions. For steady state expression, lentiviral

vectors [pCLT vectors (pCLT-GFP, TruB1 and its mutants)] were used with doxycycline

(Clontech) treatment (1.0 µg/ml) for 5 days. For virus production, HEK293FT cells were

transfected with the lentiviral vector pCLTs together with the packaging plasmids

pCMV-VSVG-RSV-Rev (addgene) and pCAG-HIVgp (addgene) at a 1:0.5:0.5 ratio

using PEI Max reagent (Polysciences) according to the Manufacturer's instructions.

Supernatants were harvested 48 h after transfection. Cells (5×10^5) were mixed in 2 ml

viral supernatant supplemented with 8 µg/ml polybrene (Merck Millipore) and further

incubated at 37°C. The infected cells were selected by 1 µg/ml puromycin (Thermo

Fisher Scientific) and kept in selection medium for 7 days.

Genome editing

TruB1-Flag tagged HEK293FT cells were established by using CRISPR–Cas9 system.

ALT-R XT CRISPR RNA (crRNA) (Integrated DNA Technologies) and ALT-R

trans-activating crRNA (tracrRNA) (Integrated DNA Technologies) were resuspended in

Nuclease-free-duplex buffer (Integrated DNA Technologies) to a final concentration of

100 µM. crRNA and tracRNA with equal volumes were mixed and heated for 5 minutes

at 95 °C, followed by gradually cooling down to room temperature. Mixed RNAs were

incubated at room temperature for 20 minutes with ALT-R Cas9 Nuclease V3 (61 µM)

(Integrated DNA Technologies) and single-stranded oligo DNA (ssODN) including

sequences of HiBIT, 3 x Flag and complementary sequence of N-terminal in TruB1

genome. Next, this RNA, protein and ssODN were co-transfected into HEK-293FT cells

using a NEPA21 Super Electroporator (NEPAGENE). We used the HiBIT system to

detect the knocked-in cells (Schwinn et al., 2018) with dilution methods. After

transfection, knocked-in cells were detected by Nano Glo HiBiT Lytic Detection System

(Promega) using the Manufacturer's recommended protocol. Knocked-in cells were

verified by PCR and Western blotting with anti-Flag antibody (MBL).

Cell-based screening

We created 384-well reporter library plates for cell-based screening. In detail, library

plasmids were diluted to 4 ng/µL in elution buffer (10 mM Tris·HCl; pH 8.5) in 96-well

plates and were then dispensed into 384-well culture plates at 5 µL per well. Cell-based

screening for let-7 regulator genes was performed by modifying a previously described

method (Ito et al., 2017). Five microliters of OPTI-MEM (Gibco) containing 0.15 µL of

PEI-Max (Polyscience), 20 ng of sensor vector or control sensor (-) vector, and 5 ng of

pRL-SV40 (Promega) Renilla luciferase construct was added to the 384-well reporter

library plates and incubated for 20 min. Then 293FT cells (Thermo Fisher Scientific) in

40 µL of DMEM containing 10% of FBS were added into each well using an automated

multidispenser ECO DROPPER III (AS ONE). The cells were cultured in a 5% CO_2

incubator at 37 °C for 48 h. Luciferase activity was measured by means of an ARVO X3

(PerkinElmer) and the Dual-Glo Luciferase Assay System (Promega).

Luciferase reporter assay

Luciferase assays were performed by incubating 20 µL of OPTI-MEM containing 0.4 µL

of FuGENE HD, 25 ng of reporter plasmids and 5 ng of pRL- SV40 for 10 min in 96-well

culture plates. Then 293FT cells in 100 µL of DMEM containing 10% of FBS were added

to each well and cultured for 48 h. Luciferase activity was measured using an ARVO X3

(PerkinElmer) and the Dual-Glo Luciferase Assay System (Promega).

Reverse transcription and RT-qPCR

Total RNA samples were extracted using ISOGEN (NIPPON GENE). To evaluate mRNA

expression, reverse transcription was carried out using oligo dT primers (Thermo Fisher

Scientific) and SuperScript III (Thermo Fisher Scientificic). Quantitative RT-PCR

(qRT-PCR) was run using THUNDERBIRD SYBR qPCR Mix (TOYOBO). The results

are expressed as mRNA levels normalized to GAPDH mRNA expression in each

sample. To evaluate microRNA expression, reverse transcription was performed using

the TaqMan MicroRNA Reverse Transcription Kit (Thermo Fisher Scientific) and

TaqMan MicroRNA Assays (Thermo Fisher Scientific). Quantitative RT-PCR (qRT-PCR)

was run using THUNDERBIRD Probe qPCR Mix (TOYOBO). The results are expressed

as RNA levels normalized to RNU6B expression in each sample. The primer sequences

and assays used for the real-time PCR are listed in Table S4. To determine RNA levels,

at least three independent RNA samples were analyzed.

Northern blotting

LNA/DNA probes (PAGE grade, fasmac, Table SX) were labelled with 32P-γ-ATP by

means T4PNK (Takara), followed to filter by MicroSpin G-25 Columns (GE healthcare).

4 ug of small RNA purified by the mirVana miRNA Isolation kit (Thermo Fisher Scientific) was heated at 90 °C for 20 sec with gel loading buffer II (Thermo Fisher Scientific) and placed on ice, followed by running on a 15% TBE-Urea PAGE. RNA was transferred to a Hybond-N+ membrane (GE Healthcare) at 4mA/cm^2 for 45 min and fixed by UV-crosslinking. The membrane was dried and pre-hybridized at 37 °C for 30 min in ExpressHyb (Clontech). Hybridization was performed in ExpressHyb containing RI-labeled probe at 37 °C overnight with rotation. Subsequently, the membrane was washed twice in low stringency buffer (2X SSC, 0.05% SDS) at room temperature for 30 min and twice in high stringency buffer (0.5X SSC, 0.1% SDS) at room temperature for 40 min. Membranes were exposed to an autoradiography film.

TaqMan Array

Total RNA was isolated using the mirVana miRNA isolation kit (Thermo Fisher Scientific).

3μL of RNA (1000 ng) were reverse transcribed by Megaplex RT Primers, Pool A (human) (Thermo Fisher Scientific) using the TaqMan® microRNA Reverse

Transcription Kit (Thermo Fisher Scientific). For qRT-PCR, RT product was mixed with

TaqMan® Universal PCR Master Mix, No AmpErase UNG (Thermo Fisher Scientific)

and the appropriate amount of water, and subsequently loaded into the ports of the

TaqMan array cards. RealTime PCR was performed on a 7900HT Fast Real-Time PCR

System with TaqMan® Array Block (Thermo Fisher Scientific), using universal cycling

conditions (95 °C/10 min, then [95 °C/15 sec, 60 °C/60 sec] for 40 cycles). Raw Ct

values were normalized to RNU6B.

Western blotting

Total protein samples were extracted by radioimmunoprecipitation assay (RIPA) buffer

[50 mM Tris·HCl (pH 8.0), 150 mM NaCl, 0.5% deoxycholate (DOC), 0.1% SDS, 1%

Nonidet P-40]. Proteins in the cell lysates were separated by SDS/PAGE followed by

semidry transfer to a PVDF membrane. Membranes were blocked for 60 min with

Blocking One (Nacalai Tesque), incubated with an anti-Flag (M185-3L, MBL),

anti-DGCR-8 (ab90579, abcam), anti-Lin28B (11965S, Cell signaling technologies) or

anti–β-actin (A5316; Sigma Aldrich) at 4 °C overnight, rinsed, and then incubated for 1 h

with ECL mouse IgG HRP-conjugated whole antibody (GE Healthcare) or rabbit IgG

HRP-conjugated whole antibody (GE Healthcare). The blot was developed with the ECL

Select Western Blotting Detection Reagent (GE Healthcare). The protein levels are

normalized to β-actin levels in each sample.

HITS-CLIP

Flag-tagged cells were collected and crosslinked with 254 nm UV-crosslinking at 400

mJ/cm2 and 200mJ/cm2 on ice in a Ultraviolet Crosslinker (UVP). The cells were

collected, centrifuged and resuspended in Lysis buffer (1xPBS, 0.1% SDS, 0.5% DOC,

0.5% NP-40). After incubation with rotation for 10 min at 4°C, cells were sonicated

30sec/30sec (on/off) for 5cycles by Bioruptor (Cosmo BIO). Then, whole cell lysate was

collected by centrifugation at 13,000 × g for 15 min at 4 °C and treated with 5 U/ml

RNase T1 (Thermo Fisher Scientific) and Turbo DNase (Thermo Fisher Scientific) for 5

min at 37°C. Three different doses of RNaseT1 and one non-crosslinked cells were

used as controls. For immunoprecipitation, we used mouse anti-FLAG antibody (MBL)

and normal mouse IgG (Sigma Aldrich) as an additional negative control. Antibodies

were coupled to Dynabeads Protein G (Thermo Fisher Scientificc) following the

Manufacturer's protocol. Coupled antibodies were washed with PBST. RNaseT1 treated

whole cell lysate and coupled antibodies were mixed and rotated at 4°C for 2h.

Antibody-bound proteins and RNA-protein complexes were isolated by magnetic stand

and washed with wash buffer (1xPBS, 0.1% SDS, 0.5% DOC, 0.5% NP-40) and

high-salt-wash buffer (5xPBS, 0.1% SDS, 0.5% DOC, 0.5% NP-40). Purified RNA

fragments were treated with CIP (Takara) and ligated with 3' linker adaptor by using T4

RNA ligase (Thermo Fisher Scientific) at 16°C overnight. Treated RNA fragments were

labelled with 32p-γ-ATP, followed by electrophoresis with SDS-PAGE (NuPAGE,

Thermo Fisher Scientific) and transferred to nitrocellulose membranes. RI-labelled

RNAs were detected by autoradiography. Appropriate bands or smear-bands were

picked up from the membrane and treated with Proteinase K (Roche). RNA fragments

were purified by Acid phenol :CHCL3 (Thermo Fisher Scientific). Purified TruB1-bound

RNA fragments were ligated with 5′ linker RNA, followed by the reverse-transcriptase

reaction using RT primers and SuperScriptIII (Thermo Fisher Scientific). cDNA libraries

were amplified by means of Phusion High-Fidelity DNA Polymerase (New England

Biolabs). Quantification of libraries was calculated using Quibit3.0 (Thermo Fisher

Scientific), TapeStation (Agilent) with High Sensitivity DNA Kit, and qPCR analysis by

NEB Next Library Quant Kit for Illumina (New England Biolabs). All sequencing was

performed using an Illumina MiSeq (Illumina) with MiSeq Reagent Kit v3 (Illumina).

HITS-CLIP analysis

Raw reads (fastq file) obtained from the Illumina pipeline were checked, trimmed and

clipped using Bioconductor, FastQC, FASTX-toolkit and cutadapt (v.1.2.1) and were

subsequently mapped to the human genome (build hg38) using bowtie2. To eliminate

reads in which the adaptor was ligated more than once, adaptor removal was performed

3 times. Duplicate reads were filtered by IGV tools. All data analysis was done using a

combination of custom and publicly available tools (featureCounts, SAMtools, RSeQC,

bedtools, tRNAscan-SE 2.0 and R) including resources available from the UCSC

Genome Browser and Galaxy Bioinformatics (http://main.g2.bx.psu.edu/). All aligned

reads shorter than 10nt that mapped to more than a single position in the genome were

excluded from further analysis. In trimmed data, the 500-bp flanking sequences of

known pre-miRNAs were extracted as pri- miRNAs. Transcript annotation was acquired

from UCSC. The read count is plotted for miRNAs with TaqMan array data in this report.

Data was normalized by expression levels of each microRNA in control samples of the

TaqMan array.

Motif Analysis

We identified motifs for sequences mapped to tRNAs. Reads were randomly selected

within 5000 reads by seqkit (https://bioinf.shenwei.me/seqkit/) to perform the following

analysis. Multiple Em for Motif Elicitation (MEME) was performed using v.4.9.1 of the

MEME browser application.

RNA immunoprecipitation (RIP)

Cells were collected, centrifuged and resuspended in Lysis buffer (1xPBS, 0.1% SDS,

0.5% DOC, 0.5% NP-40). After incubation with rotation for 10 min at 4°C, cells were

sonicated 30sec/30sec (on/off) for 5 cycles. Then, whole cell lysate was collected by

centrifugation at 13,000 × g for 15 min at 4 °C. For immunoprecipitation, we used

mouse anti-FLAG (MBL), anti-DGCR-8 (abcam) and normal mouse IgG (Sigma Aldrich)

as an additional negative control. Antibodies were coupled to Protein G Dynabeads

(Thermo Fisher Scientific) following the Manufacturer's protocol. Coupled antibodies

were washed with PBST. Whole cell lysate and coupled antibodies were mixed and

rotated at 4°C for 2h. Antibody-bound proteins and RNA-protein complex were isolated

by magnetic stand and washed with wash buffer (1xPBS, 0.1% SDS, 0.5% DOC, 0.5%

NP-40) and high-salt-wash buffer (5xPBS, 0.1% SDS, 0.5% DOC, 0.5% NP-40). RNA

was eluted by Acid phenol :CHCL3 (Thermo Fisher Scientific). Pri-miRNA levels were

analyzed by qRT-PCR. Relative ratio of protein bound pri-miRNA was calculated as

compared to its input sample.

In vitro processing assay

pri-miRNAs were in vitro transcribed using MEGA script T7 Transcription Kit (Thermo

Fisher Scientific) with a 32p-α-UTP (RI) label, followed by PAGE purification. We also

synthesized RI-labeled pri-let-7a1 in which pseudouridine was randomly introduced

using UTP: pseudouridine (Carbosynth) at a ratio of 1:1. Cells were collected,

centrifuged and resuspended in 200 μL of Lysis buffer (WCL) (20mM Tris-HCL(pH 8.0),

100mM KCL, 0.5mM EDTA, 5% glycerol, 0.5mM PMSF and 5mM DTT). After incubation

with rotation for 10 min at 4 °C, cells were sonicated 15sec/90sec (on/off) for 3 cycles.

Whole cell lysate was collected by centrifugation at 18,000 × g for 15 min at 4 °C.

Labelled RNA was treated with whole cell lysate and 6.4mM $MgCL_2$ at 37 °C for 90 min.

RNA was eluted by Acid phenol :CHCL3 (Thermo Fisher Scientific), and run on 15%

TBE-Urea PAGE. RNA was visualized by autoradiography after gel drying.

Electrophoretic mobility shift assay (EMSA)

pri-miRNAs were in vitro transcribed using MEGA script T7 Transcription Kit (Thermo

Fisher Scientific) with 32p-α-UTP to label by RI, followed by PAGE purification. Labelled

RNA and recombinant proteins were mixed and incubated at 37 °C for 1 h, followed by

naïve PAGE. RNA and RNA-protein complexes were visualized by autoradiography.

Expression and purification of recombinant proteins

E. coli BL21 (DE3) (Novagen) was transformed by the plasmids, and the transformants

were grown at 37 °C until the A600 reached 1.0. The expression of the TruB1 proteins

was induced by adding isopropyl-β-D thiogalactopyranoside at a final concentration of

0.1 mM and incubating the cultures for 18 h at 18 °C. The cells were collected and lysed

by sonication in buffer, containing 20 mM Tris- HCl, pH 7.0, 500 mM NaCl, 10 mM

β-mercaptoethanol, 20 mM imidazole, 0.1 mM phenylmethylsulfonyl fluoride, and 5%

(v/v) glycerol. The proteins were first purified by chromatography on a Ni-NTA agarose

column (QIAGEN), and then further purified on a HiTrap Heparin column (GE

Healthcare). Finally, the proteins were purified by chromatography on a HiLoad 16/60

Superdex 200 column (GE Healthcare), in buffer containing 20 mM Tris- HCl, pH 7.0,

200 mM NaCl, and 10 mM β-mercaptoethanol. The purified proteins were concentrated

and stored at - 80 °C until use. The hTruB1 mutants were purified in a similar manner.

In vitro enzymatic activity assay

pri-miRNAs were in vitro transcribed using MEGA script T7 Transcription Kit (Thermo

Fisher Scientific) with 32p-α-UTP, followed by PAGE purification. Labelled RNA was

treated with recombinant TruB1 or its mutants in reaction buffer (20mM TrisHCL (pH

8.0), 0.1mM EDTA, 100mM NaCl, 5mM DTT and 100mM NH4Cl) at 37 °C for 60 min.

RNA was eluted by Acid phenol :CHCL3 (Thermo Fisher Scientific), and treated with

Nuclease P1 (New England BioLab) at 37 °C for 3 h. RNA was run on 1D-thin layer

chromatography on cellulose plates (TLC, 20 cm, Merck Millipore) with solvent

(isopropanol/HCl/water (70/15/15); v/v/v) for 6 h at room temperature. RNA was

visualized by autoradiography.

CMC Primer extension assay

Primers (PAGE grade, fasmac, Appendix Table S4) were labelled with 32P-γ-ATP by

T4PNK (Takara), followed by filtering through MicroSpin G-25 Columns (GE healthcare).

4 ug of total RNA was treated with 0.17M N-cyclohexyl-N0 -(2-morpholi-

noethyl)carbodiimide metho-p-toluenesulfonate (CMC, Tokyokasei) in reaction buffer

(7M urea, 50mN Bicine and 4mM EDTA) at 37 °C for 20 min. Reaction was stopped with

100 μL of stop buffer [0.3 M NaOAc and 0.1 mM EDTA, pH 5.6] and 700 μL of EtOH.

After incubation at -80 °C for 1 h. The treated RNA was centrifuged, and washed twice

with 70% EtOH. RNA pellet was dissolved in 50mN Na2CO3 (pH 10.4) and incubated at

37 °C for 4 h. Reaction was stopped in the same manner. RNA pellet was dissolved in

water and purified using the RNeasy MinElute Cleanup kit (QIAGEN). Treated RNA was

reverse transcribed with RI-labelled primer by SuperScript III (Thermo Fisher Scientific)

with or without ddATP solution (Thermo Fisher Scientific). cDNA was eluted by

phenol/chloroform/isoamyl alcohol (Thermo Fisher Scientific), and run on 10%

TBE-Urea PAGE. cDNA was visualized by autoradiography after gel drying.

Evaluation of cell viability

Cell viability was determined using RealTime-Glo MT Cell Viability Assay (Promega). A

total of 1000 cells/well were plated in 96-well culture plates. After 24 h of incubation, the

medium was changed to a growth medium with MT Cell Viability Substrate (Promega)

and Nano Luc Enzyme (Promega). Then, the cells were incubated for an additional 0,

12, 24, 48, 72 h of culture. The luminescence in the resulting solution was measured

using an ARVO X3 (PerkinElmer) at each point. Cell viability was determined by dividing

the luminescence activity by that of the control cells. These experiments were

conducted on at least three independent samples.

Data analysis from public databases

We browsed several series (GEO accession GSE3325, GDS4102, GSE64333_at;

Varambally et al., 2005; Pei et al., 2009; Wang et al., 2015) of the GEO Profiles

database (Edgar et al., 2002).

QUANTIFICATION AND STATISTICAL ANALYSIS

Blot signals were quantified using ImageJ (NIH) (Schneider et al., 2012). Data are

presented as the means ± standard errors, and differences between groups were

evaluated using the Student's t-test. A P-value of <0.05 was considered statistically

significant. Asterisk in figures indicate differences with statistical significance as follows:

*p < 0.05. All the experiments were conducted in close adherence to the institutional

regulations.

Data Availability

All the data required to reproduce this study are included in this published article and

supplementary information. The raw data of TaqMan array and HITS-CLIP were

deposited in the GEO under accession number GSE143508 and GSE143510,

respectively. Further information and requests for resources and reagents should be

directed to and will be fulfilled by Hiroshi Asahara (asahara@scripps.edu or

asahara.syst@tmd.ac.jp).

Acknowledgements

We thank Shuji Takada, Satoshi Hara and Asuteka Nagao for technical assistance and support, and Yutaro Uchida, Hiroki Tsutsumi, Maiko Inotsume, Haruka Hosogai, Hiroto Yamamoto and all other members of the Department of Systems BioMedicine in Tokyo Medical and Dental University for their support, and Helen Pickersgill at Life Science Editors for scientific critical reading and editorial assistance. This research was supported by AMED-CREST from AMED (Japan Agency for Medical Research and Development) (JP19gm0810008 to H.A.), JSPS KAKENHI (Grant numbers: 17K15018, 19K16794 to R.K. 18H03980 to K.T. 26113008, and 15H02560, 18K19603, 19KK0227 to H.A.), and grants from the National Institutes of Health (Grant numbers: AR050631, AR065379) to H.A.

Author contributions

R.K., T.C., Y.I., T.M., K.M., T.S., K.T., and H.A. designed the study; R.K. performed experiments, analyzed data; K.T. created and purified recombinant proteins; R.K., T.C., and Y.Y. generated sequencing data. T.S. and K.T. supervised the project; R.K., T.S.,

K.T., and H.A. drafted the manuscript; H.A. critically reviewed the manuscript; and all

authors approved the final manuscript.

Conflict of Interests

The authors declare no competing financial interests.

Reference

Ambros V (2004) The functions of animal microRNAs. *Nature* 431: 350-5

Antonicka H, Choquet K, Lin ZY, Gingras AC, Kleinman CL, Shoubridge EA (2017) A pseudouridine synthase module is essential for mitochondrial protein synthesis and cell viability. *EMBO Rep* 18: 28-38

Arroyo JD, Jourdain AA, Calvo SE, Ballarano CA, Doench JG, Root DE, Mootha VK (2016) A Genome-wide CRISPR Death Screen Identifies Genes Essential for Oxidative Phosphorylation. *Cell Metab* 24: 875-885

Auyeung VC, Ulitsky I, McGeary SE, Bartel DP (2013) Beyond secondary structure: primary-sequence determinants license pri-miRNA hairpins for processing. *Cell* 152: 844-58

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS (2009) MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* 37: W202-8

Becker HF, Motorin Y, Planta RJ, Grosjean H (1997) The yeast gene YNL292w encodes a pseudouridine synthase (Pus4) catalyzing the formation of psi55 in both mitochondrial and cytoplasmic tRNAs. *Nucleic Acids Res* 25: 4493-9

Bernstein E, Caudy AA, Hammond SM, Hannon GJ (2001) Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409: 363-6

Bohnsack MT, Czaplinski K, Gorlich D (2004) Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* 10: 185-91

Cai X, Hagedorn CH, Cullen BR (2004) Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* 10: 1957-66

Carlile TM, Rojas-Duran MF, Zinshteyn B, Shin H, Bartoli KM, Gilbert WV (2014) Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. *Nature* 515: 143-6

Chanda SK, White S, Orth AP, Reisdorph R, Miraglia L, Thomas RS, DeJesus P, Mason DE, Huang Q, Vega R, Yu DH, Nelson CG, Smith BM, Terry R, Linford AS, Yu Y, Chirn GW, Song C, Labow MA, Cohen D *et al.* (2003) Genome-scale functional profiling of the mammalian AP-1 signaling pathway. *Proc Natl Acad Sci U S A* 100: 12153-8

Chang TC, Zeitels LR, Hwang HW, Chivukula RR, Wentzel EA, Dews M, Jung J, Gao P, Dang CV, Beer MA, Thomas-Tikhonenko A, Mendell JT (2009) Lin-28B transactivation

is necessary for Myc-mediated let-7 repression and proliferation. *Proc Natl Acad Sci U S A* 106: 3384-9

Charette M, Gray MW (2000) Pseudouridine in RNA: what, where, how, and why. *IUBMB Life* 49: 341-51

Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K, Shiekhattar R (2005) TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature* 436: 740-4

Choudhury NR, Nowak JS, Zuo J, Rappsilber J, Spoel SH, Michlewski G (2014) Trim25 Is an RNA-Specific Activator of Lin28a/TuT4-Mediated Uridylation. *Cell Rep* 9: 1265-72

Cohn WE, Volkin E (1951) Nucleoside-5′-Phosphates from Ribonucleic Acid. *Nature* 167: 483-484

Conkright MD, Canettieri G, Screaton R, Guzman E, Miraglia L, Hogenesch JB, Montminy M (2003) TORCs: transducers of regulated CREB activity. *Mol Cell* 12: 413-23

Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ (2004) Processing of primary microRNAs by the Microprocessor complex. *Nature* 432: 231-5

Deogharia M, Mukhopadhyay S, Joardar A, Gupta R (2019) The human ortholog of

archaeal Pus10 produces pseudouridine 54 in select tRNAs where its recognition

sequence contains a modified residue. *RNA* 25: 336-351

Edgar R, Domrachev M, Lash AE (2002) Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. *Nucleic Acids Res* 30: 207-10

Friedt J, Leavens FM, Mercier E, Wieden HJ, Kothe U (2014) An arginine-aspartate

network in the active site of bacterial TruB is critical for catalyzing pseudouridine

formation. *Nucleic Acids Res* 42: 3857-70

Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R

(2004) The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432:

235-40

Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G,

Mello CC (2001) Genes and mechanisms related to RNA interference regulate

expression of the small temporal RNAs that control C. elegans developmental timing.

Cell 106: 23-34

Gutgsell N, Englund N, Niu L, Kaya Y, Lane BG, Ofengand J (2000) Deletion of the

Escherichia coli pseudouridine synthase gene truB blocks formation of pseudouridine 55 in tRNA in vivo, does not affect exponential growth, but confers a strong selective disadvantage in competition with wild-type cells. *RNA* 6: 1870-81

Hammond SM, Boettcher S, Caudy AA, Kobayashi R, Hannon GJ (2001) Argonaute2, a link between genetic and biochemical analyses of RNAi. *Science* 293: 1146-50

Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* 18: 3016-27

Heo I, Joo C, Cho J, Ha M, Han J, Kim VN (2008) Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol Cell* 32: 276-84

Hoang C, Ferre-D'Amare AR (2001) Cocrystal structure of a tRNA Psi55 pseudouridine synthase: nucleotide flipping by an RNA-modifying enzyme. *Cell* 107: 929-39

Huang da W, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57

Huang da W, Sherman BT, Lempicki RA (2009) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37: 1-13

Huang Q, Raya A, DeJesus P, Chao SH, Quon KC, Caldwell JS, Chanda SK,

Izpisua-Belmonte JC, Schultz PG (2004) Identification of p53 regulators by

genome-wide functional analysis. *Proc Natl Acad Sci U S A* 101: 3456-61

Hutvagner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD (2001) A

cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7

small temporal RNA. *Science* 293: 834-8

Iourgenko V, Zhang W, Mickanin C, Daly I, Jiang C, Hexham JM, Orth AP, Miraglia L,

Meltzer J, Garza D, Chirn GW, McWhinnie E, Cohen D, Skelton J, Terry R, Yu Y,

Bodian D, Buxton FP, Zhu J, Song C *et al*. (2003) Identification of a family of cAMP

response element-binding protein coactivators by genome-scale functional analysis in

mammalian cells. *Proc Natl Acad Sci U S A* 100: 12147-52

Ito Y, Inoue A, Seers T, Hato Y, Igarashi A, Toyama T, Taganov KD, Boldin MP,

Asahara H (2017) Identification of targets of tumor suppressor microRNA-34a using a

reporter library system. *Proc Natl Acad Sci U S A* 114: 3927-3932

Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E,

Reinert KL, Brown D, Slack FJ (2005) RAS is regulated by the let-7 microRNA family.

Cell 120: 635-47

Keffer-Wilkes LC, Veerareddygari GR, Kothe U (2016) RNA modification enzyme TruB

is a tRNA chaperone. *Proc Natl Acad Sci U S A* 113: 14306-14311

Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, Plasterk RH (2001) Dicer

functions in RNA interference and in synthesis of small RNA involved in developmental

timing in C. elegans. *Genes Dev* 15: 2654-9

Knight SW, Bass BL (2001) A role for the RNase III enzyme DCR-1 in RNA interference

and germ line development in Caenorhabditis elegans. *Science* 293: 2269-71

Krol J, Loedige I, Filipowicz W (2010) The widespread regulation of microRNA

biogenesis, function and decay. *Nat Rev Genet* 11: 597-610

Landthaler M, Yalcin A, Tuschl T (2004) The human DiGeorge syndrome critical region

gene 8 and Its D. melanogaster homolog are required for miRNA biogenesis. *Curr Biol*

14: 2162-7

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat*

Methods 9: 357-9

Lee RC, Feinbaum RL, Ambros V (1993) The C. elegans heterochronic gene lin-4

encodes small RNAs with antisense complementarity to lin-14. *Cell* 75: 843-54

Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Radmark O, Kim S, Kim VN (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425: 415-9

Lee Y, Hur I, Park SY, Kim YK, Suh MR, Kim VN (2006) The role of PACT in the RNA silencing pathway. *EMBO J* 25: 522-32

Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, Kim VN (2004) MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* 23: 4051-60

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078-9

Li X, Zhu P, Ma S, Song J, Bai J, Sun F, Yi C (2015) Chemical pulldown reveals dynamic pseudouridylation of the mammalian transcriptome. *Nat Chem Biol* 11: 592-7

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30: 923-30

Lin S, Choe J, Du P, Triboulet R, Gregory RI (2016) The m(6)A Methyltransferase

METTL3 Promotes Translation in Human Cancer Cells. *Mol Cell* 62: 335-345

Liu J, Bang AG, Kintner C, Orth AP, Chanda SK, Ding S, Schultz PG (2005) Identification of the Wnt signaling activator leucine-rich repeat in Flightless interaction protein 2 by a genome-wide functional analysis. *Proc Natl Acad Sci U S A* 102: 1927-32

Lowe TM, Chan PP (2016) tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res* 44: W54-7

Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U (2004) Nuclear export of microRNA precursors. *Science* 303: 95-8

Martin M (2011) Cutadapt removes adapter sequences from high- throughput sequencing reads. *EMBnetjournal* 17

Michlewski G, Caceres JF (2010) Antagonistic role of hnRNP A1 and KSRP in the regulation of let-7a biogenesis. *Nat Struct Mol Bio*l 17: 1011-8

Mourelatos Z, Dostie J, Paushkin S, Sharma A, Charroux B, Abel L, Rappsilber J, Mann M, Dreyfuss G (2002) miRNPs: a novel class of ribonucleoproteins containing numerous microRNAs. *Genes Dev* 16: 720-8

Newman MA, Thomson JM, Hammond SM (2008) Lin-28 interaction with the Let-7

precursor loop mediates regulated microRNA processing. *RNA* 14: 1539-49

Ota H, Sakurai M, Gupta R, Valente L, Wulff BE, Ariyoshi K, Iizasa H, Davuluri RV, Nishikura K (2013) ADAR1 forms a complex with Dicer to promote microRNA processing and RNA-induced gene silencing. *Cell* 153: 575-89

Pandolfini L, Barbieri I, Bannister AJ, Hendrick A, Andrews B, Webster N, Murat P, Mach P, Brandi R, Robson SC, Migliori V, Alendar A, d'Onofrio M, Balasubramanian S, Kouzarides T (2019) METTL1 Promotes let-7 MicroRNA Processing via m7G Methylation. *Mol Cell* 74: 1278-1290 e9

Pei H, Li L, Fridley BL, Jenkins GD, Kalari KR, Lingle W, Petersen G, Lou Z, Wang L (2009) Gene Expression Omnibus GDS4102 (https://www.ncbi.nlm.nih.gov/sites/GDSbrowser?acc=GDS4102). [DATASET]

Pei H, Li L, Fridley BL, Jenkins GD, Kalari KR, Lingle W, Petersen G, Lou Z, Wang L (2009) FKBP51 affects cancer cell response to chemotherapy by negatively regulating Akt. *Cancer Cell* 16: 259-66

Pendleton KE, Chen B, Liu K, Hunter OV, Xie Y, Tu BP, Conrad NK (2017) The U6 snRNA m(6)A Methyltransferase METTL16 Regulates SAM Synthetase Intron

Retention. *Cell* 169: 824-835 e14

Piskounova E, Polytarchou C, Thornton JE, LaPierre RJ, Pothoulakis C, Hagan JP, Iliopoulos D, Gregory RI (2011) Lin28A and Lin28B inhibit let-7 microRNA biogenesis by distinct mechanisms. *Cell* 147: 1066-79

Piskounova E, Viswanathan SR, Janas M, LaPierre RJ, Daley GQ, Sliz P, Gregory RI (2008) Determinants of microRNA processing inhibition by the developmentally regulated RNA-binding protein Lin28. *J Biol Chem* 283: 21310-4

Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26: 841-2

Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP (2011) Integrative genomics viewer. *Nat Biotechnol* 29: 24-6

Roovers M, Hale C, Tricot C, Terns MP, Terns RM, Grosjean H, Droogmans L (2006) Formation of the conserved pseudouridine at position 55 in archaeal tRNA. *Nucleic Acids Res* 34: 4293-301

Safra M, Nir R, Farouq D, Vainberg Slutskin I, Schwartz S (2017) TRUB1 is the predominant pseudouridine synthase acting on mammalian mRNA via a predictable

and conserved code. *Genome Res* 27: 393-406

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of

image analysis. *Nat Methods* 9: 671-5

Schwartz S, Bernstein DA, Mumbach MR, Jovanovic M, Herbst RH, Leon-Ricardo BX,

Engreitz JM, Guttman M, Satija R, Lander ES, Fink G, Regev A (2014)

Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation

of ncRNA and mRNA. *Cell* 159: 148-162

Schwinn MK, Machleidt T, Zimmerman K, Eggers CT, Dixon AS, Hurst R, Hall MP,

Encell LP, Binkowski BF, Wood KV (2018) CRISPR-Mediated Tagging of Endogenous

Proteins with a Luminescent Peptide. *ACS Chem Biol* 13: 467-474

Shen W, Le S, Li Y, Hu F (2016) SeqKit: A Cross-Platform and Ultrafast Toolkit for

FASTA/Q File Manipulation. *PLoS One* 11: e0163962

Song J, Zhuang Y, Zhu C, Meng H, Lu B, Xie B, Peng J, Li M, Yi C (2019) Differential

roles of human PUS10 in miRNA processing and tRNA pseudouridylation. *Nat Chem

Biol*, 16: 160-169

Tabara H, Sarkissian M, Kelly WG, Fleenor J, Grishok A, Timmons L, Fire A, Mello CC

(1999) The rde-1 gene, RNA interference, and transposon silencing in C. elegans. *Cell* 99: 123-32

Thomson JM, Newman M, Parker JS, Morin-Kensicki EM, Wright T, Hammond SM (2006) Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes Dev* 20: 2202-7

Trabucchi M, Briata P, Garcia-Mayoral M, Haase AD, Filipowicz W, Ramos A, Gherzi R, Rosenfeld MG (2009) The RNA-binding protein KSRP promotes the biogenesis of a subset of microRNAs. *Nature* 459: 1010-4

Treiber T, Treiber N, Plessmann U, Harlander S, Daiss JL, Eichner N, Lehmann G, Schall K, Urlaub H, Meister G (2017) A Compendium of RNA-Binding Proteins that Regulate MicroRNA Biogenesis. *Mol Cell* 66: 270-284 e13

Varambally S, Yu J, Laxman B, Rhodes DR, Mehra R, Tomlins SA, Shah RB, Chandran U, Monzon FA, Becich MJ, Wei JT, Pienta KJ, Ghosh D, Rubin MA, Chinnaiyan AM (2005) Gene Expression Omnibus GSE3325 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE3325). [DATASET]

Varambally S, Yu J, Laxman B, Rhodes DR, Mehra R, Tomlins SA, Shah RB,

Chandran U, Monzon FA, Becich MJ, Wei JT, Pienta KJ, Ghosh D, Rubin MA, Chinnaiyan AM (2005) Integrative genomic and proteomic analysis of prostate cancer reveals signatures of metastatic progression. *Cancer Cell* 8: 393-406

Viswanathan SR, Daley GQ, Gregory RI (2008) Selective blockade of microRNA processing by Lin28. *Science* 320: 97-100

Wang BD, Ceniccola K, Yang Q, Andrawis R, Patel V, Ji Y, Rhim J, Olender J, Popratiloff A, Latham P, Lai Y, Patierno SR, Lee NH (2015) Gene Expression Omnibus GSE64333 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE64333). [DATASET]

Wang BD, Ceniccola K, Yang Q, Andrawis R, Patel V, Ji Y, Rhim J, Olender J, Popratiloff A, Latham P, Lai Y, Patierno SR, Lee NH (2015) Identification and Functional Validation of Reciprocal microRNA-mRNA Pairings in African American Prostate Cancer Disparities. *Clin Cancer Res* 21: 4970-84

Wang L, Wang S, Li W (2012) RSeQC: quality control of RNA-seq experiments. *Bioinformatics* 28: 2184-5

Wang XR, Luo H, Li HL, Cao L, Wang XF, Yan W, Wang YY, Zhang JX, Jiang T, Kang

CS, Liu N, You YP, Chinese Glioma Cooperative G (2013) Overexpressed let-7a

inhibits glioma cell malignancy by directly targeting K-ras, independently of PTEN.

Neuro Oncol 15: 1491-501

Wightman B, Ha I, Ruvkun G (1993) Posttranscriptional regulation of the heterochronic

gene lin-14 by lin-4 mediates temporal pattern formation in C. elegans. *Cell* 75: 855-62

Wolter JM, Kotagama K, Babb CS, Mangone M (2015) Detection of miRNA Targets in

High-throughput Using the 3'LIFE Assay. *J Vis Exp*: e52647

Wolter JM, Kotagama K, Pierre-Bez AC, Firago M, Mangone M (2014) 3'LIFE: a

functional assay to detect miRNA targets in high-throughput. *Nucleic Acids Res* 42:

e132

Wright JR, Keffer-Wilkes LC, Dobing SR, Kothe U (2011) Pre-steady-state kinetic

analysis of the three Escherichia coli pseudouridine synthases TruB, TruA, and RluA

reveals uniformly slow catalysis. *RNA* 17: 2074-84

Yang X, Boehm JS, Yang X, Salehi-Ashtiani K, Hao T, Shen Y, Lubonja R, Thomas SR,

Alkan O, Bhimdi T, Green TM, Johannessen CM, Silver SJ, Nguyen C, Murray RR,

Hieronymus H, Balcha D, Fan C, Lin C, Ghamsari L *et al*. (2011) A public

genome-scale lentiviral expression library of human ORFs. *Nat Methods* 8: 659-61

Yi R, Qin Y, Macara IG, Cullen BR (2003) Exportin-5 mediates the nuclear export of

pre-microRNAs and short hairpin RNAs. *Genes Dev* 17: 3011-6

Zeng Y, Yi R, Cullen BR (2005) Recognition and cleavage of primary microRNA

precursors by the nuclear processing enzyme Drosha. *EMBO J* 24: 138-48

Zucchini C, Strippoli P, Biolchi A, Solmi R, Lenzi L, D'Addabbo P, Carinci P,

Valvassori L (2003) The human TruB family of pseudouridine synthase genes,

including the Dyskeratosis Congenita 1 gene and the novel member TRUB1.

Int J Mol Med 11: 697-704

Main Figure Legends

Figure 1. Functional screen using a luciferase reporter assay to identify RBPs

inducing let-7-expression

(A) Design of the let-7 sensor reporter and sequence of let-7a. Sensor (+) vector has

the antisense sequence of let-7a inserted into the 3'UTR of the luciferase gene. (B)

Relative luciferase activity of the let-7 sensor reporter (sensor (+)) or the negative

control sensor (-) reporter contransfected with pLX-Lin28A or the GFP vector (Ctrl) in

HEK293FT cells with or without pcDNA-pri-let-7a transfection. Error bars show SD; n=3.

(C). The screening model. A screening in 384-well plates with a library of 1469 genes

was performed. (D). Screening results: relative luciferase activity of each gene in the

library. Only genes that did not affect the luciferase activity of the sensor (-) reporter (0.5

<, > 2.0) were analyzed. The full results of the screen are also in Appendix TableS1. (E).

Relative luciferase activity of the sensor (+) reporter or sensor (-) reporter induced by

expression vectors of the top 5 genes identified in the screen or the GFP vector in

HEK293-FT cells. Error bars show SD; n=3. (F). Heat map of relative miRNA expression

of various miRNAs in HEK-293FT cells transfected with expression vectors of the top 5

genes from the screen, or the ctrl (GFP) vector. Red color represents suppressed

expression compared with ctrl (GFP). n=3.

Figure 2. TruB1 selectively promotes the expression of let-7 family miRNAs

(A). Volcano plot of TaqMan array showing the mean average expression and p-value of

miRNA expression profiles between TruB1 KD and ctrl (scramble). N=3. MiRNAs

significantly suppressed are enclosed by the red square. (B). Northern blotting for let-7a

and RNU6B in HEK293FT cells with TruB1 KD or ctrl (siRNA). (C). Hybridization

intensities of (B) were quantified and normalized to ctrl. (D). Relative expression of let-7

family miRNAs and other miRNAs in HEK293FT cells with TruB1 KD or ctrl (siRNA)

determined by qRT-PCR. (E). Relative expression of primary-let-7 family miRNAs as in

(D) determined by qRT-PCR. All experiments were performed in triplicate. Error bars

show SD; n=3.

Figure 3. TruB1 promotes let-7 processing independently of its enzymatic activity.

(A). Amino acid sequences encoding for the enzyme activity and RNA binding ability in

E.coli TruB and human TruB1 (Top). Design of mutant 1 (mt1) and mutant 2 (mt2)

(bottom). (B). In vitro enzyme activity assay. ^{32}p-UTP-labelled tRNA^{phe} were treated with

recombinant TruB1, mt1, or mt2. The strong upper bands represent UTP, and the lower

weaker bands represent pseudouridine (Ψ) on autoradiographs of the TLC plate. (C).

Relative miRNA expression of let-7a in HEK-293 cells infected with

tetracycline-inducible expressing lentiviruses for TruB1, mt1, mt2 or GFP 5 days after

doxycycline treatment, as determined by qRT-PCR. (D). Northern blotting for let-7a and

RNU6B in HEK-293 cells infected with lentiviruses encoding tetracycline-inducible

expression of TruB1, mt1, mt2, or GFP, 5 days after doxycycline treatment. (E).

Hybridization intensities of (D) were quantified and normalized to ctrl (GFP). (F).

Pseudouridylation activity of TruB1 for tRNA and pri-miRNAs. ^{32}p-UTP-labelled tRNA^{phe},

pri-let-7a1 or pri-miR10a were treated with recombinant TruB1. Upper bands represent

UTP, lower bands represent pseudouridine (Ψ) in autoradiographs of the TLC plate. (G).

Location of pseudouridine sites detected by the CMC primer extension method. Total

RNA purified from HEK-293FT cells were treated with CMC. CMC treated RNA were

reverse-transcribed with RI-labelled specific primers for tRNA^{phe} or pri-let-7b. ddATP

was used for sequence control. Pseudouridines are indicated by black arrows. All

experiments were performed in triplicate. Error bars show SD; n=3.

Figure 4. TruB1 binds to tRNA and primary-let-7

(A). RIP analysis of pri-let-7a1 and Flag-TruB1 from HEK-293FT cells. RNA was

extracted from IP material and analyzed by qRT-PCR. HEK-293FT cells were infected

with lentiviruses encoding tetracycline-inducible expression of TruB1, mt1, mt2, or GFP

and treated with doxycycline. Error bars show SD; n=3. (B). EMSA of ^{32}p-ATP-labelled

pri-let-7a1 or pri-let-7a1 loop mt mixed with recombinant TruB1 or mt2 at several doses.

RNP: Ribonucleoprotein complexes. (C). Design of Flag-labelled TruB1 knock-in (KI)

cells. HiBIT and 3 x Flag sequences were inserted into the N-terminus of the TruB1

gene in HEK-293FT cells. (D) Sequencing clusters obtained from HITS-CLIP

experiment. Left are for let-7, and right for tRNAs. (E). Tag densities represented as

dots. Density of miRNA clusters in HITS-CLIP were normalized by miRNA expression as

determined by TaqMan array. (F). The proportions of different types of mapped

transcripts from the HITS-CLIP experiment. (G). Sequence motifs from reads mapped

to tRNAs from the HITS-CLIP experiment. (H). TruB-modified site in tRNAs. Location of

pseudouridine is colored in red. (I). Sequence of the terminal loop in pri-let-7a1. Lin28B

binding site is colored in red.

Figure 5. TruB1 promotes the microprocessing of primary let-7

and enhances binding of the microprocesser to primary let-7

(A). In vitro processing assay for RI-labelled pri-let-7a1. Autoradiographs of gels

showing pri-let-7a1 treated with whole cell lysate (WCL) from HEK-293FT cells

transfected with GFP, TruB1, mt1, or mt2 (overexpression, left), and TruB1 KD or ctrl

(siRNA, right). (A). RI intensities of (B) were quantified and normalized to ctrl. Relative

processing rate of pri-let-7a1 into pre- and mature- are shown. (C). RIP assay of

pri-let-7a1 and DGCR-8 from HEK-293FT cells. Western blotting for input or

immunoprecipitate (IP) using anti-DGCR-8 antibody and anti-actin antibody are shown

on top. RNA was extracted from IP material and analyzed by qRT-PCR (bottom). (D).

RIP assay of pri-let-7a1 and TruB1 from HEK-293FT cells with Lin28B KD or ctrl

(siRNA). Western blotting for input or IP material using anti-Flag antibody, anti-Lin28B

antibody, and anti-actin antibody are shown on top. RNA was extracted from IP material

and analyzed by qRT-PCR (bottom). All experiments were performed in triplicate. Error

bars show SD; n=3.

Figure 6. TruB1 suppresses cell growth by regulating let-7

(A). Design of luciferase reporter vector comprising the let-7 target region of KRAS

mRNA (KRAS reporter). (B). Relative luciferase activity of KRAS reporter or Ctrl (empty)

reporter in HEK293FT cells infected with lentiviruses expressing tetracycline-inducible

TruB1, mt1, mt2, or GFP 5 days after doxycycline treatment. (C). Relative expression of

let-7 families determined by qRT-PCR in HEK-293FT cells with KD of let-7 family

members or ctrl (using 2'-O-methylated antisense inhibitor). (D). Real-time glo assay for

HEK293FT cells infected with lentiviruses expressing tetracycline-inducible TruB1 or

GFP, 5 days after doxycycline treatment, with or without KD of let-7 family members. (E).

Schematic model for TruB1-dependent induction of let-7. All experiments were

performed in triplicate. Error bars show SD; n=3.

Expanded View Figure Legends

Figure EV1. Schematic model of cell-based functional screen

Repressors of maturation such as Lin28 lead to high luciferase activity, and activators

lead to low luciferase activity. This cell-based screen is assaying for a function directly

on endogenous let-7 maturation, rather than affinity/binding. The gain-of-function

screen also enables to identify regulatory factors that may cause lethality under loss of

function conditions, such as with Crispr-Cas9 or RNAi-KD screens.

Figure EV2. Knockdown of TruB1 repressed expression of relative miRNAs in

various human cells.

(A). qRT-PCR analysis for TruB1 mRNA expression normalized to GAPDH. TruB1 KD

by siRNA significantly repressed the expression of TruB1 mRNA in HEK-293FT cells.

Error bars show SD; n=3. (B, C). qRT-PCR analysis for miRNAs expression in A549

cells (B) and Hela cells (C) with TruB1 KD. TruB1 KD by siRNA significantly repressed

the expression of the let-7 family. Error bars show SD; n=3.

Figure EV3. **Impact of pseudouridylation enzyme activity on microprocessing of let-7.**

(A). EMSA of 32p-ATP-labelled tRNA [Phe] mixed with recombinant TruB1, mt1 or mt2 at several doses. RNP: Ribonucleoprotein complexes. (B). Protein expression of Flag-tagged TruB1 and its mutants. Protein expression was evaluated by Western blotting. Protein was isolated from HEK-293FT cells infected with tetracycline-inducible lentiviruses expressing TruB1, mt1, or mt2, 5 days after doxycycline treatment. (C). Location of pseudouridine sites were detected by CMC primer extension. Total RNA purified from HEK-293FT cells was treated with CMC. CMC-treated RNA were reverse-transcribed with RI-labelled specific primers for tRNAphe or pri-let-7a1. ddATP was used as a sequence control. (D,E). In vitro processing analysis for pri-let-7 with pseudouridine. 32p-ATP-labelled pri-let-7a1 was synthesized using UTP : pseudouridine at a ratio of 1:1 or 1:0. This labelled RNA was treated with whole cell lysate from TruB1 expressing HEK293FT cells transfected with pcDNA3.1-TruB1. Autoradiographed image (D) and the relative processing rate (E) are shown.

Figure EV4. loop mutant of pri-let7a1 and HITS-CLIP for TruB1.

(A). Loop sequence of mutant RNA in which the loop structure of pri-let-7a1 was

modified (loop mt).

(B). Protein expression of 3 x Flag-tagged TruB1 in genome edited cells. Flag-tagged

TruB1 was most highly expressed in ED-9 cells.

(C). Western blotting for input and IP samples in HITS-CLIP.

(D). Autoradiography of RI-labelled RNA in a complex with TruB1   (black frame).

(E). Agilent data of cDNA library show smeared band at 150 – 250 bp.

(F). Mapped reads of sequenced data to human genome.

Figure EV5. Relative pri-let-7a1 expression in HEK-293 cells with knockdown of

Lin28B .

qRT-PCR analysis for pri-let-7a1 expression normalized to GAPDH. Lin28B KD by

siRNA tended to increase the expression of pri-let-7a1, but this was not statistically

significant. Error bar shows SD; n=3.

Figure EV6. TruB1 suppressed tumor progression.

(A). TruB1 and mt1 suppressed cell proliferation. Real-time glo assay for HEK-293 cells infected with tetracycline-inducible lentiviruses expressing TruB1, mt1, or GFP, 5 days after doxycycline treatment. TruB1 and mt1 suppressed cell proliferation. Error bars show SD; n=3.

(B-D). Relationship between TruB1, let-7, and tumor progression in various cancers. (B). Published microarray and microRNA-sequencing in prostate biopsy samples from patients with prostate cancer (data accessible at NCBI GEO database (Edgar et al., 2002), accession GSE64333, Wang et al., 2015; Data ref: Wang et al., 2015). In these data, there were correlations between the expression levels of TruB1 and both let-7b and let-7i. Regarding let-7a, a statistically significant correlation was not found, but the same tendency was observed. (C, D). TruB1 inversely correlates with tumor progression or oncogenesis in prostate cancer (C) and in pancreatic cancer (D). LD: local disease. Data from (Varambally et al., 2005; Data ref: Varambally et al., 2005; Pei et al., 2009; Data ref: Pei et al., 2009)

Appendix table title and legends

Appendix tableS1. Raw data of cell-based screening

Appendix tableS2. Raw data of TaqMan array

Appendix tableS3. Mapped miRNA and tRNA in HITS-CLIP

(A) Tag densities were representative as table. Density of miRNA clusters in HITS-CLIP

were normalized by miRNA expression as determined by TaqMan array. (B). Mapped

tRNA in HITS-CLIP.

Appendix tableS4. Sequences of oligonucleotide, DNA and RNA used in this

paper.

Appendix tableS5. Sequences of pCLT vector.

Figure 1

A

TaqMan array (n=3)

let-7a
let-7c
let-7g
let-7f
let-7b

p-value

Relative expression TruB1 KD / ctrl

B

ctrl   TruB1 KD

pre-let-7a

mature-let-7a

RNU6B

C

Relative mature-let-7 expression

D

■ ctrl ■ TruB1 KD

mature-miR /RNU6B

a b c d e f g i 98
Number of let-7 families

mature-miR /RNU6B

10b 15a 26b 99b 146a 200c
Number of other microRNAs

E

■ ctrl ■ TruB1 KD

pri-miR /GAPDH

a1 a2 a3 b c d e f1 f2 g i miR98
Number of let-7 families

Figure 2

Figure 3

Figure 4

A

WCL(-)

WCL(+): GFP, WT, mt1, mt2

WCL(+): ctrl, TtruB1 KD

pri-let-7a
pre-let-7a

mature-let-7a

B

Relative processing rate of pri-let-7a1

■ pre/total   ■ mature/total

GFP   WT   mt1   mt2

*

*

*

*

1.5

1.0

0.5

ctrl   truB1 KD

*

C

input   IgG   DGCR8
TruB1   Ctrl KD Ctrl KD Ctrl KD

IB : DGCR8

IB : actin

Relative pri-let-7a1 ratio (RIP / input)

1.5

0.5

IgG DGCR8   IgG DGCR8
Ctrl   TruB1 KD

*
*
*

D

input   IgG   Flag
Lin28B   Ctrl KD Ctrl KD Ctrl KD

IB : Flag

IB : actin

IB : Lin28B

Relative pri-let-7a1 ratio (RIP/ input)

0.9

0.6

0.3

IgG   Flag   IgG   Flag
Ctrl   Lin28B KD

*
*

Figure 5

A
Luciferase reporter assay (KRAS)

SV40   Luc2   miR target

KRAS mRNA position : 3466 to 3644

5'…AACUCUUCGAUCAAGCUACUUUA…3'
     | | | | | | | | | | | | | | | | |
let7a 3'…UUGAUAUGUUGGAU - GAUGGAGU…5'

B

C   Relative let-7 expression
        (/RNU6B)

Number of let-7 families

D

E
TruB1   Microprocessor
         DGCR8
         Drosha

primary-let-7   mature-let-7

Figure 6