Identification of characteristic genomic markers in human hepatoma Huh7 and Huh7.5.1-8

cell lines

Masaki Kawamoto[1], Toshiyuki Yamaji[2], Kyoko Saito[2], Kazuhiro Satomura[3], Toshinori Endo[3],

Masayoshi Fukasawa[2], Kentaro Hanada[2,*], and Naoki Osada[3, 4,*]

*Correspondence to hanak@nih.go.jp and nosada@ist.hokudai.ac.jp

1. Graduate School of Information Science and Technology, Hokkaido University, Sapporo

060-0814, Japan

2. Department of Biochemistry & Cell Biology, National Institute of Infectious Diseases,

Tokyo 162-8640, Japan

3. Faculty of Information Science and Technology, Hokkaido University, Sapporo 060-0814,

Japan

4. Global Station for Big Data and Cybersecurity, GI-CoRE, Hokkaido University, Sapporo

060-0814, Japan

Supplementary Info: Supplementary Tables 1, 2

Supplementary Dataset 1: List of missense and LoF mutations in Huh7 and Huh7.5.1-8

Abstract

The human hepatoma-derived Huh7 cell line and its derivatives (Huh7.5 and Huh7.5.1) have been widely used as a convenient experimental substitute for primary hepatocytes. In particular, these cell lines represent host cells suitable for propagating the hepatitis C virus (HCV) *in vitro*. The Huh7.5.1-8 cell line, a subline of Huh7.5.1, can propagate HCV more efficiently than its parental cells. To provide genomic information for cells' quality control, we performed whole-genome sequencing of Huh7 and Huh7.5.1-8 and identified their characteristic genomic deletions, some of which are applicable to an in-house test for cell authentication. Among the genes related to HCV infection and replication, 53 genes were found to carry missense or loss-of-function mutations likely specific to the Huh7 and/or Huh7.5.1-8. Eight genes, including *DDX58* (*RIG-I*), *BAX*, *EP300*, and *SPP1* (osteopontin), contained mutations observed only in Huh7.5.1-8 or mutations with higher frequency in Huh7.5.1-8. These mutations might be relevant to phenotypic differences between the two cell lines and may also serve as genetic markers to distinguish Huh7.5.1-8 cells from the ancestral Huh7 cells.

Introduction

Quality control of cells is crucial for biopharmaceutical manufacturing as well as research. In research, many cell lines have been misidentified (http://iclac.org/wp-content/uploads/ Cross-Contaminations-v8_0.pdf), and various academic journals have begun requesting proper authentication of cells used in studies. Cell lines can be identified by genotyping multiple common genetic variants, such as microsatellite markers and single nucleotide variants (SNVs)[1]. However, such common markers would not be useful for discriminating between cell lines that originated from the same ancestral cell line. Thus, rational, easy in-house methods of identifying cell lines are widely desired. In particular, the mutation rate of deletion is much lower than that of point mutations[2]. Cell line-specific deletions are applicable to cell line authentication of African green monkey kidney-derived Vero cells[3,4], which are widely used in virologic research and human vaccine production.

HuH-7 (hereafter Huh7) is a permanent cell line established from male hepatoma tissue, which was surgically removed from a 57-year-old Japanese male in 1982[5]. Huh7 and its derivatives have been used as a convenient experimental substitute for primary hepatocytes. Approximately 80% of hepatocellular carcinoma incidents in humans are caused by hepatitis viruses; in all hepatocellular carcinoma incidents, ~25% were caused by hepatitis C virus (HCV) and ~53% were caused by hepatitis B virus (HBV)[6]. HCV is a positive-stranded RNA virus that infects humans, causing acute and chronic liver diseases. Chronic infection eventually leads to severe symptoms, such as hepatitis and hepatocellular carcinoma. Thus, permanent cell lines suitable for investigating human hepatitis viruses have always been invaluable. After the establishment of Huh7, several studies have attempted to derive cell lines that are more permissive to HCV than Huh7[7,8]. Huh7.5, a subline of Huh7, was established as a highly permissive cell line to replicate subgenomic and genomic HCV

RNA in 2002[7]. Interestingly, Huh7.5, but not the ancestral Huh7, has a missense mutation in *DDX58*

(or *RIG-I*) gene, which participates in intracellular antiviral defense[9]. Although cell culture systems

that recapitulate the HCV life cycle have not been developed for a long time, the JFH-1 HCV strain

has been found to produce infective progenitor virions after introduction to Huh7.5 cells[10]. Huh7.5.1,

a subline of Huh7.5 cells, was subsequently established as a cell line with intent to generate a cell

line that cultures JFH-1 more efficiently than Huh7.5[11]. Although Huh7.5.1 displayed faster viral

replication kinetics than Huh7, Huh7 and Huh7.5.1 eventually produced a similar level of viral

titers[11]. We previously found that the expression level of CD81, which is a plasma membrane protein

essential for HCV infection[12], was not uniform in the Huh7.5.1 cells, and we thus obtained

Huh7.5.1-8, a subclone of Huh7.5.1, in which CD81 is stably expressed on cell surfaces[13].

Huh7.5.1-8 cells exhibit ~10-fold greater permissiveness to HCV than Huh7.5.1 cells[13]. HCV culture

systems have provided robust assay systems to screen anti-HCV compounds, resulting in the

development of various anti-HCV medicines that are currently marketed, although anti-HCV

vaccines have not been developed.

We determined the whole-genome sequences of Huh7 and Huh7.5.1-8 cell lines and found

characteristic genomic deletions in Huh7 and Huh7.5.1-8, some of which are applicable to an in-house

test for cell authentication. In addition, among the genes related to HCV infection and replication, 53

genes were found to carry missense or loss-of-function (LoF) mutations that were not registered in the

public germline mutation databases, but were found in the Huh7 and/or Huh7.5.1-8. Among them, 8

genes, including *DDX58* (*RIG-I*), *BAX*, *EP300*, and *SPP1* (osteopontin), contained mutations

observed only in Huh7.5.1-8 or mutations with higher frequency in Huh7.5.1-8.

Results and Discussion

Identification of mutations in Huh7 lineages

We obtained 41.0- and 41.1-fold coverages of whole-genome sequence reads from Huh7

(JCRB0403) and Huh7.5.1-8 cells, respectively. No sequences related to HBV and HCV were

detected in the genomes using the VirusFinder2 pipeline, as reported previously for HBV using *in*

situ hybridization[14]. To identify novel mutations, we filtered out mutations matched to the previously

known germline SNVs and indels from further analyses (see Materials and Methods). These

mutations were not found in currently available human mutation database, suggesting that most (even

if not all) of these newly-identified SNVs specifically occurred during the establishment of the Huh7

cell line and/or the development of the liver cancer in the patient, from which the cell line was

established. For the following analysis, we further classified mutations into three categories: 1)

Huh7-predominant mutations, of which   frequency is higher in Huh7 than in Huh7.5.1-8, 2)

Huh7.5.1-8-predominant mutations, of which mutation frequency is higher in Huh7.5.1-8 than in

Huh7, and 3) shared mutations, of which frequency is almost equal between the two cell lines. The

SNVs of higher frequency in Huh7 than in Huh7.5.1-8, with statistical significance (false discovery

rate of 0.05), were classified as Huh7-predominant SNVs and vice versa. The SNVs that did not pass

the criteria were further classified into shared SNVs when the variant frequencies in both samples

exceeded 0.25. Table 1 summarizes the number of mutations, including SNVs, insertions, and

deletions.

As expected, many new mutations had higher frequencies in Huh7.5.1-8 than in Huh7

(Huh7.5.1-8-predominant class), because Huh7.5.1-8 should have experienced a larger number of

cell passages. However, Huh7 also had a substantial number of mutations that were not found in

Huh7.5.1-8. The observation was, presumably, accounted for by the fact that the seed stock of Huh7

cells we sequenced was not a direct ancestral cell seed that was used to establish Huh7.5. In total, we

identified 394,568 new mutations, including 227,447 SNVs and 29,050 insertions, and 136,930

deletions.

Most of the mutations were shared between the two cell lines. These shared mutations were most

likely to have arisen before the establishment of the Huh7 cell line. Heterogeneity of mutations is often

seen not only among different cancer types but also among cancer cells of different patients diagnosed

with the same cancer type. Indeed, the amount and pattern of new mutations in liver cancer tissues

were shown to be considerably diverse[15]. In our study, Huh7 and Huh7.5.1-8 contained a relatively

large number of new mutations, although it is impossible to obtain germline cells of the patient from

whom Huh7 derived to sequence their genome, and some of the mutations might have occurred in the

germline cells. Notably, the number of insertions and deletions was considerably high, and the ratio

of indels to SNVs was much higher than the ratio observed in germline cells[2]. Our validation, using

polymerase chain reaction (PCR), confirmed that all six amplified regions contained specific

deletions in the cell line DNA, indicating that the false-positive rate would not be substantially high

in our variant calling pipeline.

Interestingly, we identified one nonsynonymous mutation (K70R) in *POLD3* gene, which plays an

important role in high-fidelity DNA replication[16] in both Huh7 and Huh7.5.1-8. In all 12 nonhuman

vertebrates we analyzed, POLD3 had lysine residues at site 70, indicating that this amino acid site

has a very important function in the protein. *POLD3* mutations have been identified repeatedly in

cancer cells[17], and they contribute to chromosomal instability[18]. We suspected that K70R mutations

in POLD3 partly explain the large number of SNVs and indels and the frequent chromosome copy

number changes in Huh7[19].

    Deletion markers for Huh7 cell lines

To establish efficient deletion markers characteristic of Huh7 cells, we designed PCR primer sets

that test the presence of deletion in genomic DNA using the dataset of homozygous deletions

identified in whole-genome sequencing. Four long deletions, ranging from 7,400 to 262,700 bp in

length, and four short deletions, ranging from 645 to 1018 bp in length, were targeted for PCR

amplification. We designated regions harboring long and short deletions as DL and DS regions,

respectively. In our genome sequencing analysis, Five indels were expected to be present only in

Huh7.5.1-8, whereas the others were expected to be shared between Huh7 and Huh7.5.1-8. We used

a DNA sample from HeLa cells as a control. Primer sequences and detailed information for each

region are presented in Supplementary Table 2.

We successfully amplified six regions by genomic PCR and confirmed that they all harbored

deletions of expected sizes in Huh7 and/or Huh7.5.1-8, but did not showed the signature deletion in

the HeLa cells (Figure 1). In total, four deletions (DL1–3 and DS2) were present only in Huh7 and

Huh7.5.1-8, and two deletions (DS1 and DS3) were observed only in Huh7.5.1-8.

Classification of newly identified SNVs

The newly identified SNVs were further classified into missense mutations, nonsense mutations,

mutations at splicing signal sites, and other mutations. In addition, frameshift and large indels

harboring protein-coding sequences (including gene fusions) were classified as LoF mutations.

Nonsense mutations and mutations at splicing signal sites were also classified as LoF mutations.

Among these mutations, we selected genes that have missense and/or LoF mutations. We

confirmed that missense mutations are present in the RNA-seq read data, which means that mutated

alleles are actually expressed in the cells. In total, 255 and 431 genes were found to have missense

and LoF mutations, respectively. The complete list of genes is shown in Supplementary Dataset 1.

We narrowed down the gene list and chose several genes that might be relevant to the HCV

replication process. We reviewed previous research and selected keywords that are related to HCV

infection and replication[20,21]: autophagy, apoptosis, antiviral defense, hepatitis C, innate immune

response, and serine protease. In addition, we surveyed the genes involved in protein-protein

association networks with nine core genes that showed a strong influence on HCV infection and

replication (*CD36*, *CD81*, *CLDN1*, *EGFR*, *EPHA2*, *LDLR*, *OCLN*, *PPIA*, and *SCARB1*).

We identified 53 candidate genes, including 12 autophagy-related genes, 22 apoptosis-related

genes, 3 antiviral defense genes, 14 HCV-related genes, 8 innate immune-related genes, 4 serine

proteases, and 4 genes involved in the HCV core-gene network. Eleven genes were categorized into

two different classes. The list of genes and mutations is summarized in Table 2. Two HLA genes,

HLA-DRB1 and *HLA-DRB5*, related to an acquired immune system, contained several LoF mutations

shared between Huh7 and Huh7.5.1-8, albeit those genes were not included in the 53 genes in Table 2

since cultured cells lack the acquired immune system. Interestingly, among the 53 candidate genes, 8

genes contained Huh7.5.8-1+ mutations, but none carried Huh7-predominant mutations. Among the

8 genes, 4 genes (*BAX*, *COL6A3*, *DEFB104B*, and *SIRPB1*) had LoF mutations, whereas the other 4

genes (*DDX58*, *EP300*, *SPP1*, and *ZNF654*) had missense mutations (Table 2). Future studies will be

required to elucidate whether these mutations are relevant to the phenotypes of the Huh7 cell lineage.

One of the genes, *DDX58*, which is also known as *RIG-I*, encodes an RNA helicase and has an

important function for innate antiviral response[22]. The mutation is, therefore, a strong candidate for

the high permissiveness of HCV in the Huh7.5 lineage. It is noteworthy that our whole-genome

sequence approach also identified this gene. However, we found a difference in the mutation

frequency between the results of this and previous studies. This incongruence is discussed in the next

subsection.

Because we do not have sufficient space to discuss all 53 candidate genes for the higher

permissiveness of HCV in detail, we here selected three key genes that have important viral

replication functions.

BAX, which has an LoF mutation observed only in Huh7.5.1-8, forms a dimer with BCL2 and

activates apoptosis[23]. In the Huh7 cell lineage, *TP53* gene, which encodes P53 protein and controls

the apoptotic process through BAX[24], has a frameshift deletion (Table 2). Therefore, BAX may

positively regulate apoptosis in Huh7. In contrast, in Huh7.5.1-8, 22% of *BAX* alleles carried

frameshift insertions, which would decrease *BAX*'s overall activity. The apoptotic activity, therefore,

might have decreased in Huh7.5.1-8 compared with Huh7, which might result in HCV's higher

replication efficiency.

Another candidate gene, *EP300*, which encodes adenovirus early region 1A-associated protein

p300[25,26], harbored two missense mutations (D985G and V1764I) in Huh7.5.1-8. *EP300* has the

function of suppressing the autophagy process[27]. Previous studies have shown that HCV and other

flaviviruses replicate in host cells using autophagosomes[28,29]. We found that the valine residue at site

1764 was highly conserved among vertebrates. The mutation may have reduced *P300*'s function for

autophagy suppression and contributed to efficient HCV replication.

The last gene we present here is *SPP1* (osteopontin), which is involved in the remodeling process

of bones [30]. Previous studies found that (1) the gene expression of *SPP1* in liver tissue was increased

according to the progress of hepatitis C[31], (2) *SPP1* enhanced autophagy in human hepatocellular

carcinoma cells[32], and (3) *SPP1* interacted with HCV proteins and helped replicate and assemble

HCV[33]. Indeed, it was reported that *SPP1* expression is up-regulated in Huh7.5 compared with

Huh7[34]. Our study found that 46% of *SPP1* alleles in Huh7.5.1-8 carried missense mutations of

K200T, whereas the frequency of mutation in Huh7 was 0%. The frequency of mutations in both

Huh7.5.1 and Huh7.5.1-8 transcripts was close to 50%, indicating the K200T mutation had already

been acquired in Huh7.5.1, and the expression of transcript with threonine allele would not be

up-regulated by HCV infection in an allele-specific manner. The missense mutation, as well as the

overall elevation of gene expression, may have increased the replication efficiency of the HCV in

Huh7 derivatives.

Missense mutation in RIG-I (T55I) was heterologous in the Huh7.5.1-8 genome

As described in the above subsection, we found a T55I mutation in RIG-I that could contribute to

the high permissiveness to HCV in Huh7.5.1-8. However, both genome sequencing and RNA-seq

data showed the mutation is heterologous in Huh7.5.1-8, contradicting previous findings that the

mutation is homozygous in Huh7.5[9]. Therefore, we resequenced the genomic DNA from Huh7 and

Huh7.5.1-8 using a Sanger sequencer and verified that the mutation is absent in Huh7 but

heterologous in Huh7.5.1-8 (Figure 2). Our recent study showed that a 55T allele transcript harbors

an allele-specific large deletion and, presumably, nonfunctional in Huh7.5.1-8 (Saito et al.,

unpublished results). The results showed that, although T55I mutation is heterozygous in the

Huh7.5.1-8 genome, all full-length RIG-I proteins have the T55I mutation.

In summary, whole-genome sequencing of Huh7 and Huh7.5.1-8 provided various genetic

characteristics of these cell lines, and some of them are, presumably, applicable to an in-house test for

authentication of the Huh7 cell lineage. In addition, 53 genes were found to carry missense or LoF

mutations specifically found in the Huh7 cell lineage. Among them, 8 genes contained mutations

observed only in Huh7.5.1-8 or mutations with higher frequency in Huh7.5.1-8. These mutations

might be relevant to the phenotypic differences between Huh7 and Huh7.5.1-8 and might also serve as

genetic markers to distinguish Huh7.5.1-8 cells from the ancestral Huh7 cells.

Materials and Methods

DNA sample preparation and sequencing

DNA samples of Huh7 hepatocarcinoma cell line (JCRB0403) and Huh7.5.1-8 cell line were

obtained from the Japanese Collection of Research Bioresources (JCRB) Cell Bank. We confirmed

that the HCV replication is slower in JCRB0403 than Huh7.5.1-8 (data not shown). For both samples,

fragment libraries (average fragment sizes of ~560 bp) were constructed using TruSeq DNA

PCR-Free LT Library Prep Kit (Illumina, San Diego, CA, USA). Paired-end sequences of 150 bp

long were determined using HiSeq X (Illumina). Approximately 1.8 billion sequence reads were

obtained from each sample. We examined whether viral sequences of HBV and HCV are integrated

into the two genomes using VirusFinder 2 software[35].

RNA-seq data

We retrieved previously obtained RNA-seq data from Huh7.5.1 and Huh7.5.1-8 (DRR018792 and

DRR018793, respectively). Because the initial quality check for the RNA-seq data showed some of

the read bases had relatively low quality, we applied Trimmomatic software (version 0.36)[36] to filter

out low-quality reads. After trimming low-quality bases of average quality score <20 (window size:

4), reads shorter than 75 bp were filtered out. RNA-seq reads were mapped to the reference human

genome using HISAT2[37].

SNV calling

Genomic reads were mapped to the reference human genome sequence (GRCh38 primary

assembly downloaded from the Ensembl database) using the BWA MEM algorithm (version

0.7.15-r1140)[38] with a default parameter setting. The read mapping rates of Huh7 and Huh7.5.1-8

were 99.88% and 99.87%, respectively. To generate alignment files for variant calling, GATK Best

Practices Pipeline 3.0, which includes duplicated read filtering, realignment around indels, and

recalibration of base quality score, was applied to the initial alignments[39].

The sites with read depths of ≥14 and ≤100 were used for the following variant calling. The

number of sites with ≥14 read depth covered >95% of the reference human genome in our dataset.

Initial SNV calling was performed using VarScan (version 2.4.3)[40] with a base quality score cutoff of

15 and a variant allele frequency cutoff of 0.1%. Multiallelic SNVs were removed from the

analytical pipeline. Known germline SNVs registered in the public database (dbSNP_149) were

filtered out using the SelectVariants program in the GATK package[39]. However, when a variant

nucleotide in Huh7 and/or Huh7.5.1-8 was different from that in the dbSNP, those variants were kept

for the further analyses as a novel variant.

We defined three categories of newly identified SNVs according to their frequency in Huh7 and

Huh7.5.1-8. We tested whether the SNV frequency is higher in Huh7 or Huh7.5.1-8 with statistical

significance. Statistical significance was evaluated using the χ^2 test and a false discovery rate of

0.05[41]. The SNVs with higher frequency in Huh7 and Huh7.5.1-8 were categorized into

"Huh7-predominant" and "Huh7.5.1-8-predominant" categories, respectively. The SNVs that did not

pass the criteria were further classified into "shared" SNVs when the variant frequencies in both

samples exceeded 0.25.

Structural variant calling

Indels shorter than 50 bp were identified, using VarScan, with the same pipeline for SNV calling.

Indels that exactly matched to known germline indels (Mills and 1000G gold standard indels[42]) were

filtered out from further analyses. Accordingly, short indels were classified into three categories:

Huh7-predominant, Huh7.5.1-8-predominant, and shared short indels.

Long indels (≥50 bp) were identified using Manta (version 1.1.0)[43]. As with the SNV calling, we

only considered the sites with read depths of ≥14 and ≤100. The estimated structural variant

frequency in Manta was difficult when we did not obtain a sufficient number of reads spanning

breakpoints. Therefore, we classified the structural variants into the above three categories without

considering variant frequencies. Large insertions that match to the insertions in Mills and 1000G

gold standard dataset and large deletions that overlapped (≥50% length of the estimated deletion size

in the cell lines) with the deletions of Mills and 1000G gold standard and/or 1000 Genome Phase 3

strucutral variants (downloaded from ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/phase3/) were filtered

out.

Functional annotation

Functional annotation of mutations was performed using the snpEff software[44] with an annotation

data snpEff_v4_3_GRCh38.86. Using the annotation information, SNVs with strong phenotypic

effects (missense mutations, nonsense mutations, and mutations at splicing acceptor-donor sites)

were extracted. Functional annotation of genes was performed using the DAVID bioinformatics

resource[45]. We used the following five fields in DAVID output: disease, functional categories, gene

ontology, pathways, and protein domains. The protein-protein association networks were retrieved

from the STRING database[46].

Sequence conservation

We investigated the evolutionary conservation at mutated amino acid sites using the amino acid

sequences of 12 nonhuman vertebrate species: *Pan troglodytes*, *Pan paniscus*, *Gorilla gorilla, Pongo*

abelii, *Macaca mulatta*, *Cricetulus griseus*, *Felis catus*, *Desmodus rotundus*, *Phascolarctos cinereus*,

and *Gallus gallus.* Sequence data were retrieved and aligned using MEGAX[47], and sequence

alignment was performed with the MUSCLE algorithm[48].

Experimental validation of deletions

The HeLa cervical carcinoma cell line (JCRB9004) was obtained from the JCRB Cell Bank and

used as a control. Genome DNA was prepared from the cell lines using the Blood Genomic DNA

Extraction Mini Kit (Favorgen, Ping-Tung, Taiwan, ROC). Deletions in the genomes of Huh7 and

Huh7.5.1-8 cells were verified by PCR experiments using a previously reported procedure (Sakuma

et al., 2017). Briefly, PrimeSTAR GXL DNA Polymerase (Takara Bio, Otsu, Japan) was used for

amplification. The reaction mixture, containing 60 ng genome DNA, was denatured at 98°C for 1

min and then subjected to 40 cycles, consisting of 98°C for 10 s, 61°C for 15 s, and 68°C, for 1 min.

The amplified products were electrophoresed on an agarose gel and visualized with the Gel Doc EZ

imager (Bio-Rad, Hercules, CA, USA). Then, 1 kb Plus DNA Ladder was used as a molecular

marker (Thermo Fisher Scientific, Waltham, MA, USA).

Sanger sequencing of DDX58 gene

Genomic DNA was prepared from 1×10^6 cells using the Blood Genomic DNA Extraction Mini Kit

(Favorgen) and used as a template to amplify a 446 bp fragment corresponding to bases 30,196 to

30,641 of the human DDX58 genic region (National Center for Biotechnology Information accession

number NG_046918.1; 78,023 bp) using PCR. PCR was performed with 35 cycles of amplification

(95°C for 10 s, 57°C for 5 s, and 72°C for 5 s) on the PCR Thermal Cycler Dice (Takara Bio) using

PrimeSTAR   MAX   DNA   polymerase   (Takara   Bio),   forward   primer

5′-GTGGCTTGGTGAAGAATGGGCACAG-3′ (bases 30,196–30,220), and reverse primer

5′-CTCAGACTAAGAGGCATGAACTATAAGTGG-3′ (complementary to bases 30,612–30,641).

The resulting fragments were separated in an agarose gel, purified using with a gel/PCR extraction kit

(FastGene; Nippon Genetics, Tokyo, Japan), and directly sequenced by the Sanger method using

forward primer 5′-CCCTATTTGGGAAGGTCTGGTGATC-3′ (bases 30,291–30,315) and reverse

primer 5′-CACTTTTACAGTATTGTCAAGCAGC-3′ (complementary to bases 30,564–30,588) by

Eurofins Genomics K.K. (Tokyo, Japan).

Acknowledgments

We thank the JCRB Cell Bank for providing DNA samples of Huh7 and Huh7.5.1-8. This work

was supported by the Japan Society for the Promotion of Science, KAKENHI (grant 17H04003 to K.H.

and grant 18H05511 to N.O.) and AMED-CREST (grant JP18gm0910005j0004 to K.H.).

Author Contributions

Study design: T.E., M.F., K.H., and N.O., performing experiments: T.Y. and K.Sai: data analysis,

M.K., K. Sat, and N.O., writing manuscript: M.K., T.Y., K.Sai., K.H., and N.O.

Additional Information

The authors declare no competing interests.

Data availability

The sequences were deposited in the DDBJ DRA database under project ID PRJDB7928. The

Python scripts used in this study were deposited in GitHub

(https://github.com/J1011Ibb14029/kwmtools).

References

1   Gilbert, D. A. *et al.* Application of DNA fingerprints for cell-line individualization.
Am. J. Hum. Genet. **47**, 499-514 (1990).

2   Wang, J. *et al.* The diploid genome sequence of an Asian individual. *Nature* **456**,
60-65, doi:10.1038/nature07484 (2008).

3   Osada, N. *et al.* The Genome Landscape of the African Green Monkey
Kidney-Derived Vero Cell Line. *DNA Res.* **21**, 673-683, doi:10.1093/dnares/dsu029 (2014).

4   Sakuma, C. *et al.* Novel endogenous simian retroviral integrations in Vero cells:
implications for quality control of a human vaccine cell substrate. *Scientific Reports* **8**, 644,
doi:10.1038/s41598-017-18934-2 (2018).

5   Nakabayashi, H., Taketa, K., Miyano, K., Yamane, T. & Sato, J. Growth of Human
Hepatoma Cell Lines with Differentiated Functions in Chemically Defined Medium. *Cancer Res.*
42, 3858-3863 (1982).

6   Perz, J. F., Armstrong, G. L., Farrington, L. A., Hutin, Y. J. F. & Bell, B. P. The
contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver
cancer worldwide. *J. Hepatol.* **45**, 529-538, doi:https://doi.org/10.1016/j.jhep.2006.05.013
(2006).

7   Blight, K. J., McKeating, J. A. & Rice, C. M. Highly Permissive Cell Lines for
Subgenomic and Genomic Hepatitis C Virus RNA Replication. *J. Virol.* **76**, 13001-13014,
doi:10.1128/jvi.76.24.13001-13014.2002 (2002).

8   Feigelstock, D. A., Mihalik, K. B., Kaplan, G. & Feinstone, S. M. Increased
susceptibility of Huh7 cells to HCV replication does not require mutations in RIG-I. *Virol. J.* **7**, 44,
doi:10.1186/1743-422x-7-44 (2010).

9   Sumpter, R. *et al.* Regulating Intracellular Antiviral Defense and Permissiveness to
Hepatitis C Virus RNA Replication through a Cellular RNA Helicase, RIG-I. *J. Virol.* **79**,
2689-2699, doi:10.1128/jvi.79.5.2689-2699.2005 (2005).

10   Wakita, T. *et al.* Production of infectious hepatitis C virus in tissue culture from a
cloned viral genome. *Nat. Med.* **11**, 791, doi:10.1038/nm1268

https://www.nature.com/articles/nm1268#supplementary-information (2005).

11   Zhong, J. *et al.* Robust hepatitis C virus infection in vitro. *Proc. Natl.*
Acad. Sci. U. S. A. **102**, 9294-9299, doi:10.1073/pnas.0503596102 (2005).

12   Petracca, R. *et al.* Structure-function analysis of hepatitis C virus envelope-CD81
binding. *J. Virol.* **74**, 4824-4830 (2000).

13   Shirasago, Y. *et al.* Isolation and Characterization of an Huh.7.5.1-Derived Cell Clone
Highly   Permissive   to   Hepatitis   C   Virus.   *Jpn.   J.   Infect.   Dis.*   **68**,   81-88,
doi:10.7883/yoken.JJID.2014.231 (2015).

14   Tay, N., Chan, S.-H. & Ren, E.-C. Detection of integrated hepatitis B virus DNA in
hepatocellular carcinoma cell lines by nonradioactive in situ hybridization. *J. Med. Virol.* **30**,
266-271, doi:doi:10.1002/jmv.1890300407 (1990).

15   Fujimoto, A. *et al.* Whole-genome mutational landscape of liver cancers displaying
biliary phenotype reveals hepatitis impact and molecular diversity. *Nat. Commun.* **6**, 6120,
doi:10.1038/ncomms7120

https://www.nature.com/articles/ncomms7120#supplementary-information (2015).

16   Johansson, E. & MacNeill, S. A. The eukaryotic replicative DNA polymerases take
shape. *Trends Biochem. Sci.* **35**, 339-347, doi:https://doi.org/10.1016/j.tibs.2010.01.004 (2010).

17   Wang, K. *et al.* Whole-genome sequencing and comprehensive molecular profiling
identify new driver mutations in gastric cancer. *Nat. Genet.* **46**, 573, doi:10.1038/ng.2983

https://www.nature.com/articles/ng.2983#supplementary-information (2014).

18   Minocherhomji, S. *et al.* Replication stress activates DNA repair synthesis in mitosis.
Nature **528**, 286, doi:10.1038/nature16139

https://www.nature.com/articles/nature16139#supplementary-information (2015).

19   Kasai, F., Hirayama, N., Ozawa, M., Satoh, M. & Kohara, A. HuH-7 reference
genome profile: complex karyotype composed of massive loss of heterozygosity. *Hum. Cell*,
doi:10.1007/s13577-018-0212-3 (2018).

20   Scheel, T. K. H. & Rice, C. M. Understanding the hepatitis C virus life cycle paves the
way for highly effective therapies. *Nat. Med.* **19**, 837, doi:10.1038/nm.3248 (2013).

21   Bukh, J. The history of hepatitis C virus (HCV): Basic research reveals unique
features in phylogeny, evolution and the viral life cycle with new perspectives for epidemic

control. *J. Hepatol.* **65**, S2-S21, doi:https://doi.org/10.1016/j.jhep.2016.07.035 (2016).

22   Yoneyama, M. *et al.* The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat. Immunol.* **5**, 730, doi:10.1038/ni1087

https://www.nature.com/articles/ni1087#supplementary-information (2004).

23   Oltval, Z. N., Milliman, C. L. & Korsmeyer, S. J. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programed cell death. *Cell* **74**, 609-619, doi:https://doi.org/10.1016/0092-8674(93)90509-O (1993).

24   Toshiyuki, M. & Reed, J. C. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell* **80**, 293-299, doi:https://doi.org/10.1016/0092-8674(95)90412-3 (1995).

25   Eckner, R. *et al.* Molecular cloning and functional analysis of the adenovirus E1A-associated 300-kD protein (p300) reveals a protein with properties of a transcriptional adaptor. *Genes Dev.* **8**, 869-884, doi:10.1101/gad.8.8.869 (1994).

26   Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. & Nakatani, Y. The Transcriptional Coactivators p300 and CBP Are Histone Acetyltransferases. *Cell* **87**, 953-959, doi:https://doi.org/10.1016/S0092-8674(00)82001-2 (1996).

27   Pietrocola, F. *et al.* Spermidine induces autophagy by inhibiting the acetyltransferase EP300. *Cell Death Differ.* **22**, 509, doi:10.1038/cdd.2014.215

https://www.nature.com/articles/cdd2014215#supplementary-information (2014).

28   Fahmy, A. M. & Labonté, P. The autophagy elongation complex (ATG5-12/16L1) positively regulates HCV replication and is required for wild-type membranous web formation. *Scientific Reports* **7**, 40351, doi:10.1038/srep40351

https://www.nature.com/articles/srep40351#supplementary-information (2017).

29   Levine, B., Mizushima, N. & Virgin, H. W. Autophagy in immunity and inflammation. *Nature* **469**, 323, doi:10.1038/nature09782 (2011).

30   Kahles, F., Findeisen, H. M. & Bruemmer, D. Osteopontin: A novel regulator at the cross roads of inflammation, obesity and diabetes. *Molecular Metabolism* **3**, 384-393, doi:https://doi.org/10.1016/j.molmet.2014.03.004 (2014).

31 Asselah, T. *et al.* Liver Gene Expression Signature of Mild Fibrosis in Patients With
Chronic Hepatitis C. *Gastroenterology* **129**, 2064-2075,
doi:https://doi.org/10.1053/j.gastro.2005.09.010 (2005).

32 Liu, G. *et al.* Osteopontin induces autophagy to promote chemo-resistance in human
hepatocellular carcinoma cells. *Cancer Lett.* **383**, 171-182,
doi:https://doi.org/10.1016/j.canlet.2016.09.033 (2016).

33 Iqbal, J. *et al.* Osteopontin Regulates Hepatitis C Virus (HCV) Replication and
Assembly by Interacting with HCV Proteins and Lipid Droplets and by Binding to Receptors
αVβ3 and CD44. *J. Virol.* **92**, e02116-02117, doi:10.1128/jvi.02116-17 (2018).

34 Choi, Steve S. *et al.* Osteopontin is up-regulated in chronic hepatitis C and is
associated with cellular permissiveness for hepatitis C virus replication. *Clin. Sci.* **126**, 845-855,
doi:10.1042/cs20130473 (2014).

35 Wang, Q., Jia, P. & Zhao, Z. VERSE: a novel approach to detect virus integration in
host genomes through reference genome customization. *Genome Med.* **7**, 2,
doi:10.1186/s13073-015-0126-6 (2015).

36 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).

37 Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low
memory requirements. *Nat. Methods* **12**, 357, doi:10.1038/nmeth.3317

https://www.nature.com/articles/nmeth.3317#supplementary-information (2015).

38 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

39 McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303,
doi:10.1101/gr.107524.110 (2010).

40 Koboldt, D. C. *et al.* VarScan 2: Somatic mutation and copy number alteration
discovery in cancer by exome sequencing. *Genome Res.* **22**, 568-576, doi:10.1101/gr.129684.111
(2012).

41 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*

(Methodological) **57**, 289-300 (1995).

42   Mills, R. E. *et al.* An initial map of insertion and deletion (INDEL) variation in the
human genome. *Genome Res.* **16**, 1182-1190, doi:10.1101/gr.4565806 (2006).

43   Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline
and cancer sequencing applications. *Bioinformatics* **32**, 1220-1222,
doi:10.1093/bioinformatics/btv710 (2016).

44   Cingolani, P. *et al.* A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff. *Fly (Austin)* **6**, 80-92, doi:10.4161/fly.19695 (2012).

45   Jiao, X. *et al.* DAVID-WS: a stateful web service to facilitate gene/protein list analysis.
Bioinformatics **28**, 1805-1806, doi:10.1093/bioinformatics/bts251 (2012).

46   Szklarczyk, D. *et al.* STRING v10: protein–protein interaction networks, integrated
over the tree of life. *Nucleic Acids Res.* **43**, D447-D452, doi:10.1093/nar/gku1003 (2015).

47   Knyaz, C., Stecher, G., Li, M., Kumar, S. & Tamura, K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **35**, 1547-1549,
doi:10.1093/molbev/msy096 (2018).

48   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).

Figure Legends

Figure 1

Evaluation of deletions identified in Huh7 and Huh7.5.1-8 using PCR. Samples in lanes a to c are

DNA from Huh7, Huh7.5.1-8, and HeLa cells, respectively. Size markers are indicated by horizontal

lines. The designs for PCR amplification and expected results are shown on the bottom right. The

region name is given for each gel image. Prefixes DL and DS represent whether the expected

deletion length is long (>7.4 kbp) or short (<0.9 kbp). For DL regions, primer pairs used for the

experiments are shown below each image. For DS regions, primer pairs s and as1 were used for all

regions. Each gel image was cropped from the full-sized gels, but different lanes were not merged

into a single image.

Figure 2

Sequence chromatograms of *DDX58* in Huh7 and Huh7.5.1-8. The genomic regions harboring the

exon 2 of *DDX58* were amplified from Huh7 and Huh7.5.1-8 DNA samples using PCR and sequenced

by the Sanger method. The arrows indicate a base 30,443 on NG_046918.1, and C-to-T transition at

this position causes a T55I substitution of the DDX58 amino acid sequence.

Tables

Table 1. Summary of newly identified variants in Huh7 and Huh7.5.1-8

Mutation type	Huh7-predominant	Huh7.5.1-8-predominant	Shared
SNV	4,094	39,497	183,856
Insertion	911	933	27,206
Deletion	1,898	8,948	126,084

     Table 2 Mutations potentially contributing HCV permissiveness to Huh7 and Huh7.5.1-8

Gene	Status	Type*	Ensembl transcript (AA type)	Function †
BAX	Huh7.5.1-8-predominant	CID	N/A	2
COL6A3	Huh7.5.1-8-predominant	SSM	N/A	7
DEFB104B	Huh7.5.1-8-predominant	CID	N/A	6
SIRPB1	Huh7.5.1-8-predominant	GFM	N/A	6
ACIN1	Shared	CID	N/A	2
AR	Shared	CID	N/A	5
ATG3	Shared	CID	N/A	1
ATM	Shared	SDA	N/A	1, 2
ATP6AP2	Shared	CID	N/A	2
CD34	Shared	CID	N/A	5
CHMP4A	Shared	CID	N/A	1, 2
CTSA	Shared	CID	N/A	1
EGF	Shared	CID	N/A	5, 4
EIF2AK3	Shared	CID	N/A	5
EIF4G3	Shared	CID	N/A	1
GPX1	Shared	CID	N/A	2

H3CR1	Shared	CID	N/A	5
HTT	Shared	CID	N/A	1, 2
IFNA10	Shared	SSM	N/A	3, 5, 6
IFT20	Shared	SDA	N/A	1
KIR2DL4	Shared	CID, SDA	N/A	5
M6PR	Shared	CID	N/A	4
MAP3K1	Shared	CID	N/A	2
MMP17	Shared	CID	N/A	5
NOTCH4	Shared	CID	N/A	5
OVOS2	Shared	CID	N/A	7
PHLDA1	Shared	CID	N/A	2
PHLPP1	Shared	CID	N/A	2
PRUNE2	Shared	CID	N/A	2
PTEN	Shared	SDA	N/A	2
SNCA	Shared	SDA	N/A	1
TMBIM4	Shared	GFM	N/A	2
TNRC18	Shared	CID, SSM	N/A	7
TP53	Shared	CID	N/A	5, 2, 4
UBE2D3	Shared	CID	N/A	2
DDX58	Huh7.5.1-8-predominant	Missense	ENST00000379883.2 (T55I)	3, 5, 6
EP300	Huh7.5.1-8-predominant	Missense	ENST00000263253.8 (D985G)	1
EP300	Huh7.5.1-8-predominant	Missense	ENST00000263253.8 (V1764I)	1

	nt			
SPP1	Huh7.5.1-8-predomina nt	Missense	ENST00000508233.5 (K200T)	5
ZNF654	Huh7.5.1-8-predomina nt	Missense	ENST00000636215.1 (D1072N)	4
ASAH2	Shared	Missense	ENST00000489640.5 (C74Y)	2
ATP6V0A2	Shared	Missense	ENST00000613625.4 (G82E)	1, 7
C4A	Shared	Missense	ENST00000498271.1 (S347Y)	6
CASP8AP2	Shared	Missense	ENST00000551025.3 (K1245N)	2
CYLD	Shared	Missense	ENST00000427738.7 (T822N)	6
DAB2IP	Shared	Missense	ENST00000408936.7 (R1164C)	6, 2
DIDO1	Shared	Missense	ENST00000266070.8 (S1019C)	2
DPF2	Shared	Missense	ENST00000531989.1 (G41V)	2
IL4R	Shared	Missense	ENST00000563002.5 (A82T)	5
PAK4	Shared	Missense	ENST00000360442.7 (R175W)	2
PIP4K2C	Shared	Missense	ENST00000540759.6 (A300G)	1
TBC1D25	Shared	Missense	ENST00000376771.8 (N277S)	1
TNFRSF10 B	Shared	Missense	ENST00000276431.8 (V191A)	5, 2
TP53	Shared	Missense	ENST00000509690.5 (Y88C)	5, 2, 4
TRIM56	Shared	Missense	ENST00000306085.10 (R395W)	3, 6

478 *Classes of LoF mutations: SDA, splice donor/acceptor mutation; GFM, gene fusion mutation; CID,

479 coding sequence InDel; Missense, missense mutation; SSM, start/stop codon mutation.

480 † 1, autophagy-related gene; 2, apoptosis-related gene; 3, antivirus-related gene; 4, genes involved in

481 protein-protein association networks, with nine core genes that showed strong influence on HCV

482 infection and replication; 5, known hepatitis C-related gene; 6, innate immune-related gene; 7, serine

483 protease-related gene.

	Forward primer	Reverse primer	Base 30,443
Huh7			C
Huh7.5.1-8			C/T