The Gene Expression Profile of Uropathogenic *Escherichia coli* in Women with

Uncomplicated Urinary Tract Infections Is Recapitulated in the Mouse Model

Arwen E. Frick-Cheng[1*], Anna Sintsova[1,2*], Sara N. Smith[1], Michael Krauthammer[2], Kathryn A.

Eaton[1] and Harry L. T. Mobley[1]

[1] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, USA

[2] Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

[*] Authors contributed equally. Author order was determined alphabetically.

Correspondence should be addressed to Harry L. T. Mobley (hmobley@med.umich.edu)

Keywords: UPEC, transcriptome, human infection, mouse model of UTIs

Abstract

Uropathogenic *Escherichia coli* (UPEC) is the primary causative agent of uncomplicated

urinary tract infections (UTIs). UPEC fitness and virulence determinants have been evaluated in a

variety of laboratory settings that include a well-established mouse model of UTI. However, the

extent to which bacterial physiology differs between experimental models and human infections

remains largely understudied. To address this important question, we compared the transcriptomes

of three different UPEC isolates in human infection and a variety of laboratory conditions

including LB culture, filter-sterilized urine culture, and the UTI mouse model. We observed high

correlation in gene expression between the mouse model and human infection in all three strains

examined (Pearson correlation coefficient of 0.86-0.87). Only 175 of 3,266 (5.4%) genes shared

by all three strains had significantly different expression levels, with the majority of them (145

genes) down-regulated in patients. Importantly, gene expression of both canonical virulence

factors and metabolic machinery were highly similar between the mouse model and human

infection, while the *in vitro* conditions displayed more substantial differences. Interestingly,

comparison of gene expression between the mouse model and human infection hint at differences

in bladder oxygenation as well as nutrient composition. In summary, our work strongly validates

the continued use of this mouse model for the study of the pathogenesis of human UTI.

Importance

Different experimental models have been used to study UPEC pathogenesis including *in vitro*

cultures in different media, tissue culture, as well as mouse models of infection. The latter is

especially important since it allows evaluation of mechanisms of pathogenesis and potential

therapeutic strategies against UPEC. Bacterial physiology is greatly shaped by environment and it

is therefore critical to understand how closely bacterial physiology in any experimental model

relates to human infection. In this study, we found a very strong correlation in bacterial gene

expression between the mouse model and human UTI using identical strains, suggesting that the

mouse model accurately mimics human infection, definitively supporting its continued use in UTI

research.

Introduction

Urinary tract infections (UTIs) are one of the most common bacterial infections in otherwise healthy individuals. Over 50% of women will experience at least one UTI in their lifetime, and half of these women will experience a recurrent infection within a year (1, 2). These infections affect 150 million people per year and result in annual medical costs of $3.5 billion in the US alone (3). Uropathogenic *Escherichia coli* (UPEC) is responsible for 80% of uncomplicated UTIs (1) and deploy diverse strategies to survive and replicate in the human host. These comprise an array of virulence factors including, but not limited to, iron acquisition systems (siderophores and heme receptors), fimbriae and other adhesins, flagella, and toxins (4-7). The importance of these systems to bacterial fitness has been studied in detail using multiple models including cultures in laboratory media, human urine cultures, tissue culture, and a mouse model first established over 30 years ago (8). However, animal models can fail to recapitulate important aspects of the human response to disease (9). Whether the mouse model accurately reflects the native environment found during human infection has not been adequately addressed. Therefore, it is vitally important to determine if the mouse model of ascending UTI recapitulates human UTI since defining mechanisms of pathogenesis and the development of UTI therapies relies on this assumption (10).

Previous studies compared the mouse model to human UTIs using microarrays to assess differences in bacterial gene expression (11, 12). Initially, urine from mice infected with UPEC type strain CFT073 was collected over a period of ten days, pooled and analyzed using a microarray based on the CFT073 genome (11). In a follow-up study, urine was collected from eight women with complicated UTIs and bacterial gene expression in the human host was analyzed, again using microarrays based on the CFT073 genome (12). Relative expression levels

of 46 fitness genes were compared between the mouse model and human UTI. This comparison

demonstrated a Pearson's correlation coefficient of 0.59 and was strongest for iron acquisition

systems and weakest for adhesin and motility systems (12). While encouraging, this study did not

provide conclusive evidence that the mouse model closely replicated human UTI. A key weakness

of the previous comparison was that genetic differences between currently circulating isolates and

strain CFT073 used for the mouse infections would obscure strain-specific responses, either due

to differences in mouse *versus* human UTI or because the CFT073-specific microarrays would not

detect expression of genes that are not encoded by that strain.

We have recently used RNA sequencing (RNA-seq) to quantify the UPEC transcriptome

during acute infection in 14 female patients (13). Importantly, RNA-seq is a more comprehensive

platform to analyze the transcriptome of clinical UPEC strains since, unlike microarrays, it is not

limited by strain-specific probes. In this study, we report the transcriptome during murine UTI for

three of the 14 clinical strains using RNA-seq and directly compare the gene expression patterns

for these identical strains between human UTI and the mouse model. We observed a high

correlation between human infection and mouse infection (Pearson correlation coefficient ranging

from 0.86-0.87) with only 175 of 3,266 shared genes being differentially expressed. Gene

expression of classical virulence factors as well as metabolic genes in the mouse model closely

resembled those observed during human UTI. Our study is the first of its kind to directly compare

the bacterial transcriptomes between human and mouse UTI using identical strains. We conclude

that the mouse model accurately reflects bacterial gene expression observed during human

infection.

Results

Study Design. We previously sequenced the transcriptomes of 14 UPEC strains isolated directly from the urine of patients with uncomplicated UTIs (hUTI) and immediately stabilized with RNAprotect (13)**.** Three out of 14 strains (HM43, HM56, and HM86) were chosen to conduct transcriptomic studies in the prevailing mouse model of UTI (mUTI) (8). We selected strains whose hUTI transcriptomes had the highest proportion of bacterial reads to eukaryotic reads (**Supplemental Table 1**) and that possessed a prototypical UPEC virulence factor profile. All three strains belong to the B2 phylogroup (13), where the majority of UPEC strains reside, and encode a range of siderophores, heme receptors, as well as multiple fimbrial types (**Fig. 1**).

To compare UPEC gene expression during mUTI against hUTI, 40 mice were transurethrally inoculated with each UPEC strain and mouse urine was collected directly into RNAprotect, 48 hours post inoculation, for RNA isolation and sequencing. Animals were then sacrificed and the bacterial burden of their urine, bladder and kidneys was quantified. All three strains successfully colonized the animals with bacterial burdens ranging between $5.0 \times 10^3 - 4.4 \times 10^4$ CFU/g in the bladder and $1 \times 10^4 - 1.2 \times 10^6$ CFU/g in the kidneys (**Fig. 2A**), levels of colonization that are consistent with an active UTI. We also assessed levels of inflammation (on a scale from 0 to 3) in the bladders and kidneys of these infected mice, comparing them to mice that were mock-infected with PBS (**Fig. 2B, Supplemental Fig.1**). After 48 hours, infection with all of the three UPEC strains resulted in mild levels of inflammation in the bladder (median inflammation scores of 1.0, 0.25, and 0.5 for HM43, HM56 and HM86, respectively) and slightly higher levels in the kidneys (median inflammation scores of 1.25, 1.5, and 1.0 for HM43, HM56 and HM86, respectively). These similar scores indicated that the general host responses were comparable across these three different strains.

In addition to isolating RNA from mouse urine during mUTI, we also isolated and

sequenced RNA from HM43, HM56 and HM86 cultured to mid-logarithmic phase in both filter-

sterilized human urine and lysogeny broth (LB). All samples processed in this study underwent

identical treatments to deplete eukaryotic mRNA, prepare libraries, and conduct sequencing (see

Methods).

The bacterial transcriptome is highly correlated between human and mouse infections. First,

we assessed how UPEC gene expression during hUTI compared to that during *in vitro* conditions

and mUTI. For each strain, we compared $\log_2$ transcripts per million (TPMs) of every gene

between LB and hUTI, human urine culture and hUTI, and finally mUTI and hUTI. Gene

expression during hUTI and mUTI was most highly correlated with the Pearson correlation

coefficient (*r*) ranging from 0.86 to 0.87 (**Fig. 3**). In contrast, the *in vitro* human urine culture when

compared to hUTI exhibited lower correlation values of 0.73-0.80 (**Fig. 3**), consistent with our

previous report (13). Interestingly, gene expression correlation between LB and hUTI was higher

than the correlation between urine and hUTI (*r* between 0.80-0.88) (**Fig 3**). Our data demonstrate

that murine infection is the most reliable and consistent model to recapitulate the conditions that

are observed during human infection.

Contribution of growth rate to gene expression patterns. We have recently shown that diverse

UPEC strains show a conserved gene expression pattern in human patients with uncomplicated

UTIs (13). Since we saw such strong correlation between gene expression in patients and in mice

for each of the UPEC strains (**Fig 3**), we hypothesized that we would also observe a conserved

pattern of gene expression between different UPEC strains during mUTI. To address this question,

we performed principal component analysis (PCA) on gene expression of the 3,266 genes present

in all three UPEC strains (**Fig. 4A**). We observed four distinct clusters that corresponded to the

two *in vitro* growth conditions (LB and filter-sterilized human urine cultures) and the two infection

sites (human patients and mice) all displaying condition-specific gene expression programs.

Samples from patients and mice clustered closer to each other than to *in vitro* samples, suggesting

that there is an infection-specific gene expression pattern conserved between the two hosts.

We further wanted to estimate the contribution of growth rate to similarity of gene

expression profiles between mUTI and hUTI since a major hallmark of the conserved

transcriptional program of UPEC in humans is rapid growth (13-15). When cultured *in vitro,* there

is no appreciable difference in the growth rate of the UPEC strains cultured in human urine

(**Supplemental Figure 2A, B**). There appears to be an extremely subtle difference in growth rate

in LB between HM43 and HM86 (**Supplemental Figure 2C, D**), which is interesting given that

these two strains had the largest difference in correlation between hUTI and LB (*r* values of 0.81

and 0.88 respectively). However, while HM86 had the highest correlation, it grew more slowly

than HM43, indicating that it is potentially more than just growth rate that is driving this similarity.

We also performed PCA using only ribosomal protein expression data (**Fig. 4B**), since

expression of ribosomal proteins is directly correlated with bacterial growth rate (16, 17) as another

method to dissect the contribution of growth rate. While this analysis still showed a clear difference

between urine samples and other conditions, it failed to clearly separate LB, mUTI and hUTI

samples (**Fig. 4B**). This suggests that while growth rate undoubtedly shapes the gene expression

pattern in mUTI and hUTI, growth rate alone is insufficient to explain gene expression patterns

observed in **Fig. 4A** and **Fig. 3**. Additionally, we looked at how much each of the ribosomal genes

contributed to the either Principal Component 1 (PC1) or Principal Component 1 (PC2) (**Fig. 4C**).

We found that all of the ribosomal genes contributed more to PC1 than PC2 values. Thus, we

hypothesize that PC1 might in fact separate samples based on growth rate, while other unknown

factors account for sample separation along PC2.

The conclusions of PCA analysis (clear separation of samples based on condition rather

than on strain) were independently confirmed using Ward's hierarchical clustering of $\log_2$ TPM

values (**Fig 4D**). We also demonstrate that growth in rich medium (LB) more closely mimics hUTI

than growth in nutrient-poor human urine, in agreement with the previously demonstrated rapid

growth of UPEC in the host (both mouse and human) compared to slow growth in human urine

(13-15, 18).

Differentially regulated genes between human and mouse infection suggest nutritional

disparities. Despite the high concordance of hUTI and mUTI gene expression data, we wanted to

determine whether any genes are differentially regulated between human and mouse infection. To

answer this question, we used the R package DEseq2 (19) to find significant differences in gene

expression between the two different infections. Strikingly, only 175 genes, representing 5.4% of

the 3,266 genes analyzed, were differentially regulated (30 upregulated, 145 downregulated) in

human infection compared to the mouse model (**Fig. 5A, Table 1** and **2, Supplemental Table 2**).

The upregulated gene with the highest $\log_2$ fold-change difference (4.3) between human

and mouse was *cspA*, which encodes an RNA chaperone initially identified as a cold shock protein

(20, 21). However, this protein may have other functions as it is highly expressed during early

exponential phase (22) and during the introduction of fresh nutrient sources (23). In addition, one

of the genes responsible for cell division, *ftsB,* and a major regulator of ribosomal RNA

transcription, *fis,* were upregulated in hUTI as compared to mUTI.

The majority of differentially regulated genes were downregulated in patients compared to

mice, and several of these genes span operons encompassing specific systems (**Table 2** and

Supplemental Table 2). For example, during mUTI, we observed: increased expression of the

citrate lyase operon *citCDEFGTX*, which is responsible for the conversion of citrate oxaloacetate

and acetate and feeds into the production of acetyl-COA under anaerobic conditions (24); the

pathway for allantoin breakdown (*allABDC*); and ethanolamine utilization

(*eutABCDEGHJLMNPQST*) (**Table 2** and **Supplemental Table 2**). In addition, genes encoding

transporters for the uptake of L-arabinose (*araADFGH)*, L-ascorbate (*ulaABCDEDF*), and

allantoin (*ybbW*) were transcribed at higher levels during mUTI (**Table 2** and **Supplemental Table**

2). Furthermore, several genes related to anaerobic metabolism or fermentation (*hycBDF* and

frdAB) were more highly expressed in mice (**Table 2** and **Supplemental Table 2**). All of these

results indicate subtle nutrient differences between the mouse and human urinary tract.

Infection-specific gene expression. Additionally, we were interested in identifying genes that

behaved similarly during both human and mouse infection, *i.e*., genes that were up- or down-

regulated in both mouse and human UTI when compared to either of the *in vitro* conditions (LB

or filter-sterilized human urine). There were 54 downregulated genes in both mouse and human

UTI when compared to LB (**Fig. 5B, Supplemental Table 3**) and there were 67 upregulated genes

during both mUTI and hUTI when compared to LB (**Fig. 5B, Supplemental Table 4**).

Interestingly, both chemotaxis (*cheABWYZ*) and flagellar machinery *(flgCFGLM* and *fliS*) were

downregulated during infection, which may be attributed to the fact that the UPEC strains we are

analyzing were isolated from the urine of infected individuals; motility genes tend to be

upregulated when UPEC enters the ureters to ascend to the kidneys (25). In contrast, *nrdEFHI*

genes are upregulated in both mice and humans compared to LB. These genes are ribonucleotide

reductases required for DNA synthesis, and therefore often associated with fast growth, fitting the

previously established paradigm of UPEC's rapid *in vivo* growth rate during human and murine

infection (13-15, 18).

There were 82 genes that were downregulated during either mUTI or hUTI when compared

to urine (**Fig. 5C**, **Supplemental Table 5**). These included branched-chain amino acid

biosynthesis (*ilvCDEMN*) and leucine biosynthesis (*leuABCD*) operons, consistent with previous

literature indicating that UPEC scavenges amino acids and peptides during infection(13, 26). In

contrast, there were 72 genes that were upregulated in humans and mice when compared to urine

(**Fig. 5C, Supplemental Table 6**). As previously reported (13), we observed 16 genes associated

with ribosomal subunit production as well as the master regulator *fis*, which activates rRNA

transcription, together reinforcing our observation that bacteria are growing rapidly in the host (13,

14).

Expression of fitness factors during murine infection is predictive of gene expression during

human infection. Finally, we wanted to determine whether previously identified UPEC virulence

factors that have been studied using *in vivo* mouse models would show comparable levels of

expression during both mouse and human infections. We focused on three major functional groups

of fitness factors: iron acquisition systems, adhesins, and metabolism (**Supplemental Table 7**).

We plotted $\log_2$ TPM of the genes in each functional group, comparing expression between hUTI

and LB, hUTI and urine, as well as hUTI and mUTI for each of the UPEC strains (**Fig. 6**,

Supplemental Fig. 3, Supplemental Fig 4.)

As expected, iron acquisition gene expression is much higher during human infection than

during growth in rich LB medium (**Fig. 6A**). The expression levels are more similar between urine

culture (an iron-poor medium) and human infection, but murine infection provides the most

analogous profile (**Fig. 6A**). The only adherence gene cluster that was highly expressed in any of

the assayed conditions was the *fim* operon, which encodes type 1 fimbriae. Expression of *fim* genes

was higher in patients compared to either of the *in vitro* conditions, but almost perfectly matched

the expression levels observed during mouse infection (**Fig. 6B**).

229   Anaerobic metabolic genes showed a major difference between human infection and *in*

vitro growth. The converse is also true; aerobic respiration genes, in particular, were expressed at

lower levels in humans than in either *in vitro* condition. Importantly, we also observed that the

expression levels of aerobic respiration genes align concordantly between human and mice (**Fig.**

6C), while anaerobic respiration gene expression was elevated in mice compared to humans. This

observation corroborates results from **Fig. 5A**, **Table 2** and **Supplemental Table 2**, where several

genes involved in anaerobic metabolism were expressed at higher levels in mice compared to

hUTI. Overall, we conclude, with only limited exceptions, that the mouse model of UTI not only

shows a strong global correlation of gene expression with hUTI, but also closely reflects the

expression of virulence and fitness genes that are known to contribute to UPEC fitness during

human infection.

Discussion

UPEC virulence factors as well as potential therapeutic strategies have been studied in detail using a well-established mouse model of ascending UTI that involves transurethral inoculation of UPEC into the bladder. This mouse model has been extensively used in the field, and the original papers defining this model (8, 27) have been cited nearly 500 times. While this model has been the gold-standard in the field to relate scientific discovery to human health, there are some differences between the experimental model and what occurs in human infection. In the murine model $>10^7$ CFU are inoculated directly into the mouse bladder, while in human infection the number of infecting bacteria is likely to be far lower, and bacteria are not directly introduced into the bladder. Rather, in humans, it is thought that the periurethral region is transiently colonized by UPEC, and bacteria ascend to the bladder and in some cases to the kidneys (28). Furthermore, there are some differences in the immune responses between mice and humans. For example, mice express Toll-like receptor 11 (TLR11), which induces host inflammation in response to UPEC and provides the kidneys with a modest level of protection from bacterial colonization (29). However, humans do not express TLR11 (29), indicating that not all murine responses will precisely mimic the human response. Therefore, it is of the utmost importance to validate the murine model of UTI and understand the extent to which it recapitulates human disease. Until now, there has been no direct comparison of global bacterial gene expression between human and mouse studies using the identical strain, and it is essential to understand how the mouse model relates to human disease. This study is the first to define the bacterial transcriptome from infected patients and infected mice using the same UPEC strain, thus presenting a direct comparison between the murine model and human infection. Our study demonstrates that the UTI mouse model accurately recapitulates the human disease with respect to the bacterial transcriptional response.

We compared three UPEC strains (HM43, HM56, and HM86) that were isolated in 2012

from women with symptoms of cystitis and documented significant bacteriuria (30). We isolated

bacterial RNA, stabilized immediately, from their urine to conduct RNA-seq and define the core

bacterial transcriptome during acute human infection (13). The same strains were then used for

mouse infection, followed by urine collection, bacterial RNA isolation and sequencing. We

consistently observed an extraordinarily high correlation between the bacterial transcriptome

during mouse and human infection, with the Pearson correlation coefficient ranging from 0.86 to

0.87. This correlation is not strain-specific, as infections with all three stains showed similar

results. Expression of virulence and metabolic genes was also found to be very similar between

human and mouse infection. While all three strains belonged to B2 phylogroup, we do not believe

this to be a phylogroup-specific effect as we have previously shown few differences in gene

expression between different UPEC phylogroups (13). This provides strong evidence that the

mouse model is an accurate representation of the infection that occurs in humans, and by

understanding which genes do not correlate between human and mouse, we can further understand

the limitations of the mouse model.

Mounting evidence suggests that UPEC in the human host are rapidly dividing (13, 14),

and we have recently shown that this rapid growth rate is recapitulated in the mouse model,

although to a lesser degree (13). This difference in growth rate between human and mouse UTIs

potentially can be understood by examining the genes that are differentially expressed between

human and mouse infections. Most of the differentially expressed genes were expressed at a lower

level during hUTI compared to mUTI (145 of 175 genes). Many of these 145 genes are involved

in anaerobic metabolism. Several of them were clustered in operons encoding oxidoreductases

involved in fumarate or nitrite reduction. Additionally, genes involved in nutrient usage under

anaerobic conditions were also expressed at a lower level during hUTI, such as the *all* operon,

which encodes the catabolic pathway for allantoin degradation (a step in purine catabolism (31)),

or the *ula* operon, which encodes both an L-ascorbate transporter and the corresponding enzymes

for L-ascorbate utilization, a compound that can be present in urine due to its water soluble nature

(32). Therefore, we hypothesize that the human bladder is better oxygenated than the mouse

bladder. Indeed, a higher oxygen level in human bladders might also account for the higher levels

of replication observed in hUTI (13), since an aerobic lifestyle can support more rapid growth.

There were also differences in transport systems involved in nutrient acquisition between hUTI

and mUTI. Arabinose transport (*araADFGH*) was expressed at a lower level in humans; this sugar

has been shown to be present in human bladders in μM amounts when normalized to creatine (32).

It is also present in murine bladders (33), but has never been precisely quantified. It would be

interesting if arabinose is present in lower amounts in humans compared to mice, accounting for

this difference in regulation. One of the few genes that was upregulated in humans compared to

mice was *dsdX*, which encodes a D-serine transporter. Interestingly, D-serine is present in

micromolar amounts in human urine (34), D-serine utilization is associated with uropathogenic

strains (35, 36) and accumulation of D-serine leads to a "hypervirulent" phenotype (36, 37) in the

urinary tract. Since there was an upregulation in D-serine transport, and not in the deaminase

required for its breakdown (*dsdA*), perhaps this presents a mechanism to increase the intracellular

levels of D-serine specific to hUTI.

We also compared the gene expression profiles of UPEC during infection with the two

most common *in vitro* models, LB cultures and pooled filter-sterilized human urine cultures.

Surprisingly, even though urine might seem to be the more physiologically relevant medium for

in vitro experimentation, LB overall provided a better model for infection compared to urine

cultures. Indeed, the correlations of strains grown in LB approached the correlations when

comparing human infection to the mouse model (**Fig. 3**). However, pathway analysis using the

online tool DAVID (38) revealed several statistically significantly enriched (enrichment score

≥1.5) pathways that were enriched when comparing either hUTI with LB (**Supplemental Table**

8) or mUTI (**Supplemental Table 9**). Several pathways were shared in both comparisons,

specifically iron homeostasis or acquisition, as well as motility and various metabolism pathways

(TCA, arginine and proline, and sulfur) were differentially regulated between the LB and *in vivo*

conditions. (**Supplemental Table 8** and **9**). Not only do these results corroborate the results from

Fig. 6 they also point to shared differences when comparing either infection model to *in vitro*

further underscoring the similarities between mUTI and hUTI.

A major difference between *in vitro* growth of UPEC in LB or urine is growth rate, and

one of the major hallmarks of the transcriptional program of UPEC during infection is rapid

growth. Subsequently, when comparing the pathways enriched in the differentially regulated genes

between hUTI or mUTI to urine, ribosomes are enriched (**Supplemental Table 10** and **11**).

However, that is just one of seven other statistically significantly pathways that are enriched; there

is a variety of factors influencing this outcome. Furthermore, several of these pathways

(biosynthesis of amino acids, flavin adenine dinucleotide binding, and nitrogen

metabolism/nitrogen utilization) are shared between mUTI vs urine and hUTI vs urine, implying

that there are similarities between the two infections that are not observed in *in vitro* growth in

urine (**Supplemental Table 10** and **11**).

There are several factors that vary between infection conditions and *in vitro* growth that

could account for the lower correlation between hUTI and urine culture gene expression. For

example, a nutrient limitation specific to *in vitro* growth. The bladder is akin to a chemostat, with

fresh urine constantly being introduced into the organ, a condition that is not recapitulated in the

in vitro conditions. Furthermore, the collected urine is typically filter-sterilized and this method

excludes exfoliated bladder epithelial cells, likely another major source of nutrients for the

pathogen during human infection. In future studies, we could add lysed bladder cells from cell

culture to supplement the filter-sterilized urine and determine if this represents a better model of

the nutrient milieu. However, when studying specific systems, such as iron acquisition, urine is a

better model than LB, since it more accurately recapitulates the iron-limited environment of the

host.

In summary, while both *in vitro* models have advantages and disadvantages, the mouse

model offers a holistic representation of infection and accurately models UPEC pathway

expression. Detailed examination of gene expression profiles between mice and humans allows us

to understand which parts of the model are not recapitulated in patients, so we may tailor our

studies accordingly. This way the mouse model can serve as a platform to answer questions that

are more difficult or impossible to assess when working with human patients.

Methods

Bacterial culture conditions. Clinical UPEC strains HM43, 56 and 86 (30) were cultured

overnight in LB medium at 37°C with aeration. The next morning, cultures were centrifuged and

the pellets washed twice with PBS, then diluted 1:100 into either fresh LB medium or human urine.

The human urine was collected and pooled from at least four healthy female volunteers and passed

through a 0.22 μm filter for sterilization. Bacteria were cultured at 37°C with aeration to mid-

exponential phase (~3 hours), then stabilized in RNAprotect (Qiagen). Bacterial pellets were stored

at -80°C until RNA isolation.

Mouse infection. Forty female CBA/J mice were transurethrally inoculated, using the previously

established ascending model of UTI (8), with 10^8 CFU of either HM43, 56, or 86 that had been

cultured in LB overnight with shaking. The infection was allowed to progress for 48 hours. Urine

from five mice was collected to enumerate bacterial burden, while the rest was collected for RNA

(see below for method). Mice were sacrificed, and their bladders and kidneys aseptically removed,

homogenized, and plated to determine bacterial burden. Mouse urine was collected as previously

described (13). Briefly, 48 HPI, urine was directly collected into RNAprotect, pooled, and pelleted.

This was repeated every 45 minutes five more times resulting in a total of six pellets. These pellets

were stored at -80°C until RNA isolation.

RNA isolation and library preparation. RNA was isolated as previously described (13). Briefly,

all bacterial pellets were treated with both lysozyme and proteinase K, and then total RNA was

extracted using a RNeasy kit (Qiagen). Genomic DNA was removed using the Turbo DNA-free

kit (ThermoFisher). Eukaryotic mRNA was depleted using Dynabeads covalently linked with

oligo dT (ThermoFisher). The *in vitro* samples underwent the same treatment with Dynabeads to

reduce any potential biases this procedure might introduce to the downstream sequencing. The

supernatant was collected from this treatment, and RNA was concentrated and re-purified using

the RNA Clean and Concentrator kit (Zymo).

To compare the results of the new RNA-sequencing experiment to the published

expression data obtained from the human samples (13), library preparation method needs to be

identical to avoid batch effects. The original sequencing data (hUTI) were obtained using the

ScriptSeq Complete kit (Bacteria) to prepare the cDNA library. However, at the time of this

study, Illumina had discontinued this kit. Therefore, we needed to use any leftover kits that had

the same base preparation method (Scriptseq). To accomplish this, we used ScriptSeq Complete

Gold Kit (Epidemiology), which also contains rRNA removal for bacteria and eukaryotes for the

HM86 mouse sample, and all three HM43 samples (mouse, LB, and urine). ScriptSeq Complete

(Bacteria) was used on the HM56 mouse sample, where the kit contained rRNA removal for

bacteria; mammalian rRNA was removed with ThermoFisher's mammalian rRNA removal kit

(cat #457012). Then the *in vitro* samples from both HM56 and HM86 were prepared using the

ScriptSeq Complete (bacteria) kit, which removes bacterial rRNA.

RNA-sequencing. *E. coli* HM43 was sequenced using an Illumina HiSeq2500 (single end, 50 bp

read length) and *E. coli* HM56 and HM86 were sequenced using the Nextseq-500 with identical

conditions (single end, 50 bp read length).

Histology and tissue processing. Bladders were removed and halved by cutting on the transverse

plane, while kidneys were cut on the sagittal plane. One half of each organ was used to enumerate

CFU, while the other halves were placed into tissue cassettes and immersion-fixed in 10% formalin

for at least 24 hours. They were then embedded in paraffin, cut into thin sections and stained with

hematoxylin and eosin (H&E) by the *In Vivo* Animal Core at the University of Michigan. Tissue

sections were scored as described in **Table 3**. Briefly, this is a scale of 0-3, where 0 was no

inflammation, and 3 was severe inflammation. Each organ section was scored by two different

people in a blinded manner and the scores averaged together.

Table 3: Scoring criteria for histopathological analysis.				
	Score			
Lesion	0	1	2	3
Cystitis	No scorable lesions	very rare PMNs in stroma or lumen or occasional perivascular lymphoid cuffs	many PMNs and moderate edema	Many PMNs; widespread, marked edema, transmural inflammation
Pyelonephritis	No scorable lesions	very occasional PMNs in lumen or peripelvic tissue	Rafts of PMNs in the pelvis and/or scattered focal aggregates of PMNs in peripelvic tissue	many PMNs in all sections, or a single large focus of PMNs in one section

402 **RNAseq Data Processing.** A custom bioinformatics pipeline was used for the analysis

403 (github.com/ASintsova/rnaseq_analysis). Raw fastq files were processed with Trimmomatic (21)

404 to remove adapter sequences and analyzed with FastQC to assess sequencing quality. Mapping

405 was done with bowtie2 aligner (39) using default parameters. Alignment details can be found in

406 **Supplemental Table 1**. Read counts were calculated using HTseq htseq-count (40).

Pearson correlation coefficient calculation, PCA analysis and hierarchical clustering analysis. For PCA and correlation analysis, transcript per million (TPM) was calculated for each gene; TPM distribution was then normalized using $\log_2$ transformation and these normalized data were used for both PCA and correlation, and hierarchical clustering analysis. Pearson correlation and PCA were performed using Python sklearn library. Hierarchical clustering was performed using Ward's method and Euclidean distance. All of the analysis we also performed using centered log-ratio transformation (instead of $\log_2$ transformation), which has recently proposed for compositional data and saw similar results (these results can be found in Jupyter notebook associated with this manuscript). Jupyter notebooks used to generate the figures are available at https://github.com/ASintsova/upec_mouse_model.

Differential expression analysis. Differential expression analysis was performed using DESeq2 R package (19). Genes with $\log_2$ fold change of greater than 1 or less than -1 and adjusted *p* values (Benjamini-Hochberg adjustment) of less than 0.05 were considered to be differentially expressed. Pathway analysis was performed using R package topGO (41).

Data access. Jupyter notebooks as well as all the data used to generate the figures in this paper are available on github: https://github.com/ASintsova/upec_mouse_model.

426
References

1. Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. 2015. Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. Nature Reviews
Microbiology 13:269-284.

2.   Foxman B. 1990. Recurring urinary tract infection: incidence and risk factors. Am J
Public Health 80:331-3.
3.   Foxman B. 2010. The epidemiology of urinary tract infection. Nature reviews Urology
7:653-660.
4.   Terlizzi ME, Gribaudo G, Maffei ME. 2017. Uropathogenic *Escherichia coli* (UPEC)
Infections: Virulence Factors, Bladder Responses, Antibiotic, and Non-antibiotic
Antimicrobial Strategies. Frontiers in Microbiology 8:1566.
5.   Subashchandrabose S, Mobley HLT. 2015. Virulence and Fitness Determinants of
Uropathogenic *Escherichia coli*. Microbiology Spectrum 3.
6.   Sivick KE, Mobley HLT. 2010. Waging war against uropathogenic *Escherichia coli*:
winning back the urinary tract. Infection and Immunity 78:568-585.
7.   Alteri CJ, Mobley HLT. 2015. Metabolism and Fitness of Urinary Tract Pathogens.
Microbiology Spectrum 3.
8.   Hagberg L, Engberg I, Freter R, Lam J, Olling S, Svanborg Edén C. 1983. Ascending,
unobstructed urinary tract infection in mice caused by pyelonephritogenic *Escherichia*
coli of human origin. Infection and Immunity 40:273-283.
9.   Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker HV, Xu W, Richards DR,
McDonald-Smith GP, Gao H, Hennessy L, Finnerty CC, López CM, Honari S, Moore
EE, Minei JP, Cuschieri J, Bankey PE, Johnson JL, Sperry J, Nathens AB, Billiar TR,
West MA, Jeschke MG, Klein MB, Gamelli RL, Gibran NS, Brownstein BH, Miller-
Graziano C, Calvano SE, Mason PH, Cobb JP, Rahme LG, Lowry SF, Maier RV,
Moldawer LL, Herndon DN, Davis RW, Xiao W, Tompkins RG, Inflammation, Host
Response to Injury LSCRP. 2013. Genomic responses in mouse models poorly mimic
human inflammatory diseases. Proceedings of the National Academy of Sciences of the
United States of America 110:3507-3512.
10.   O'Brien VP, Hannan TJ, Nielsen HV, Hultgren SJ. 2016. Drug and Vaccine Development
for the Treatment and Prevention of Urinary Tract Infections. Microbiology Spectrum 4.
11.   Snyder JA, Haugen BJ, Buckles EL, Lockatell CV, Johnson DE, Donnenberg MS, Welch
RA, Mobley HLT. 2004. Transcriptome of uropathogenic *Escherichia coli* during urinary
tract infection. Infection and Immunity 72:6373-6381.
12.   Hagan EC, Lloyd AL, Rasko DA, Faerber GJ, Mobley HLT. 2010. *Escherichia coli*
global gene expression in urine from women with urinary tract infection. PLoS pathogens
6:e1001187.
13.   Sintsova A, Frick-Cheng AE, Smith S, Pirani A, Subashchandrabose S, Snitkin ES,
Mobley H. 2019. Genetically diverse uropathogenic *Escherichia coli* adopt a common
transcriptional program in patients with UTIs. eLife 8:e49748.
14.   Forsyth VS, Armbruster CE, Smith SN, Pirani A, Springman AC, Walters MS,
Nielubowicz GR, Himpsl SD, Snitkin ES, Mobley HLT. 2018. Rapid Growth of
Uropathogenic *Escherichia coli* during Human Urinary Tract Infection. mBio 9.
15.   Bielecki P, Muthukumarasamy U, Eckweiler D, Bielecka A, Pohl S, Schanz A, Niemeyer
U, Oumeraci T, von Neuhoff N, Ghigo J-M, Häussler S. 2014. *In vivo* mRNA profiling of
uropathogenic *Escherichia coli* from diverse phylogroups reveals common and group-
specific gene expression profiles. mBio 5:e01075-01014.
16.   Dennis PP, Ehrenberg M, Bremer H. 2004. Control of rRNA Synthesis in Escherichia
coli: a Systems Biology Approach. Microbiology and Molecular Biology Reviews
68:639-668.

17.   Klumpp S, Scott M, Pedersen S, Hwa T. 2013. Molecular crowding limits translation and
478   cell growth. Proceedings of the National Academy of Sciences 110:16754-16759.
479   18.   Burnham P, Dadhania D, Heyang M, Chen F, Westblade LF, Suthanthiran M, Lee JR, De
480   Vlaminck I. 2018. Urinary cell-free DNA is a versatile analyte for monitoring infections
481   of the urinary tract. Nature Communications 9:2412.
482   19.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
483   for RNA-seq data with DESeq2. Genome Biology 15:550.
484   20.   Jones PG, VanBogelen RA, Neidhardt FC. 1987. Induction of proteins in response to low
485   temperature in *Escherichia coli*. Journal of bacteriology 169:2092-2095.
486   21.   Goldstein J, Pollitt NS, Inouye M. 1990. Major cold shock protein of *Escherichia coli*.
487   Proceedings of the National Academy of Sciences 87:283-287.
488   22.   Brandi A, Pon CL. 2012. Expression of *Escherichia coli cspA* during early exponential
489   growth at 37 °C. Gene 492:382-388.
490   23.   Yamanaka K, Inouye M. 2001. Induction of CspA, an *E. coli* major cold-shock protein,
491   upon nutritional upshift at 37 degrees C. Genes to cells : devoted to molecular & cellular
492   mechanisms 6:279-290.
493   24.   Pos KM, Dimroth P, Bott M. 1998. The *Escherichia coli* Citrate Carrier CitT: a Member
494   of a Novel Eubacterial Transporter Family Related to the 2-Oxoglutarate/Malate
495   Translocator from Spinach Chloroplasts. Journal of Bacteriology 180:4160-4165.
496   25.   Lane MC, Alteri CJ, Smith SN, Mobley HLT. 2007. Expression of flagella is coincident
497   with uropathogenic *Escherichia coli* ascension to the upper urinary tract. Proceedings of
498   the National Academy of Sciences 104:16669-16674.
499   26.   Alteri CJ, Smith SN, Mobley HLT. 2009. Fitness of Escherichia coli during Urinary
500   Tract Infection Requires Gluconeogenesis and the TCA Cycle. PLOS Pathogens
501   5:e1000448.
502   27.   Hagberg L, Hull R, Hull S, Falkow S, Freter R, Svanborg Edén C. 1983. Contribution of
503   adhesion to bacterial persistence in the mouse urinary tract. Infection and immunity
504   40:265-272.
505   28.   Kunin CM, Polyak F, Postel E. 1980. Periurethral bacterial flora in women. Prolonged
506   intermittent colonization with Escherichia coli. Jama 243:134-9.
507   29.   Zhang D, Zhang G, Hayden MS, Greenblatt MB, Bussey C, Flavell RA, Ghosh S. 2004.
508   A Toll-like Receptor That Prevents Infection by Uropathogenic Bacteria. Science
509   303:1522-1526.
510   30.   Subashchandrabose S, Hazen TH, Brumbaugh AR, Himpsl SD, Smith SN, Ernst RD,
511   Rasko DA, Mobley HLT. 2014. Host-specific induction of *Escherichia coli* fitness genes
512   during human urinary tract infection. Proceedings of the National Academy of Sciences
513   of the United States of America 111:18327-18332.
514   31.   Xi H, Schneider BL, Reitzer L. 2000. Purine catabolism in *Escherichia coli* and function
515   of xanthine dehydrogenase in purine salvage. Journal of Bacteriology 182:5332-5341.
516   32.   Bouatra S, Aziat F, Mandal R, Guo AC, Wilson MR, Knox C, Bjorndahl TC,
517   Krishnamurthy R, Saleem F, Liu P, Dame ZT, Poelzer J, Huynh J, Yallou FS, Psychogios
518   N, Dong E, Bogumil R, Roehring C, Wishart DS. 2013. The human urine metabolome.
519   PloS One 8:e73076.
520   33.   Justice SS, Lauer SR, Hultgren SJ, Hunstad DA. 2006. Maturation of intracellular
521   *Escherichia coli* communities requires SurA. Infection and Immunity 74:4793-4800.

522   34.   Brückner H, Haasmann S, Friedrich A. 1994. Quantification of D-amino acids in human
523   urine using GC-MS and HPLC. Amino Acids 6:205-211.
524   35.   Korte-Berwanger M, Sakinc T, Kline K, Nielsen HV, Hultgren S, Gatermann SG. 2013.
525   Significance of the D-serine-deaminase and D-serine metabolism of *Staphylococcus*
526   *saprophyticus* for virulence. Infection and immunity 81:4525-4533.
527   36.   Schaeffer AJ. 2004. Uropathogenic *Escherichia coli* use d-Serine Deaminase to Modulate
528   Infection of the Murine Urinary Tract. Journal of Urology 172:1571-1571.
529   37.   Anfora AT, Haugen BJ, Roesch P, Redford P, Welch RA. 2007. Roles of serine
530   accumulation and catabolism in the colonization of the murine urinary tract by
531   *Escherichia coli* CFT073. Infection and immunity 75:5298-5304.
532   38.   Anonymous. 2008. Systematic and integrative analysis of large gene lists using DAVID
533   bioinformatics resources. doi:10.1038/nprot.2008.211.
534   39.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nature
535   Methods 9:357.
536   40.   Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-
537   throughput sequencing data. Bioinformatics (Oxford, England) 31:166-169.
538   41.   Alexa A, Rahnenfuhrer J. 2018. topGO: Enrichment Analysis for Gene Ontology. R
539   package version 2.34.0.
540

Table 1: Genes differentially upregulated in human patients compared to mice[a]

Gene	Annotation	log$_2$ FC	Locus Tag
ahpC	alkyl hydroperoxide reductase, AhpC component	2.6	b0605
cspA	cold shock protein CspA	4.3	b3556
dsdX	D-serine transporter	2.7	b2365
fis	DNA-binding transcriptional dual regulator Fis	2.4	b3261
ftsB	cell division protein FtsB	2.3	b2748
gntK	D-gluconate kinase, thermostable	2.7	b3437
gpt	xanthine-guanine phsophoribosyltransferase	2.4	b0238
gspH	hypothetical type II secretion protein GspH	3.0	UTI89_C3381
gspL	hypothetical type II secretion protein GspL	3.3	UTI89_C3377
hpt	hypoxanthine phosphoribosyltransferase	2.1	b0125
ibaG	acid stress protein IbaG	2.2	b3190
lysP	lysine:H(+) symporter	2.1	b2156
opgC	protein required for succinyl modification of osmoregulated periplasmic glucans	2.6	b1047
ribE	6,7-dimethyl-8-ribityllumazine synthase	2.1	b0415
rpmE	50S ribosomal subunit protein L31	2.7	b3936
suhB	inositol-phosphate phosphatase	2.6	b2533
yajG	putative lipoprotein YajG	2.6	b0434
yceA	UPF0176 protein YceA	3.5	b1055
yciB	inner membrane protein	2.3	b1254
ydiE	PF10636 family protein YdiE	2.1	b1705
yecJ	DUF2766 domain-containing protein YecJ	2.5	b4537
yejL	DUF1414 domain-containing protein YejL	2.2	b2187
yfaZ	putative porin YfaZ	2.9	b2250
yfhL	putative 4Fe-4S cluster-containing protein YfhL	2.6	b2562
yghD	putative type II secretion system M-type protein	3.6	b2968
yghG	lipoprotein YghG	3.3	b2971
yifK	putative transporter YifK	3.6	b3795
yqcC	DUF446 domain-containing protein YqcC	2.5	b2792
yqgF	ribonuclease H-like domain containing nuclease	2.3	b2949
yqiA	esterase YqiA	2.3	b3031

[a] FC: Fold Change

Table 2: Top 30 genes differentially downregulated in human patients compared to mice[a]			
Gene	Annotation	log$_2$ FC	Locus Tag
allB	allantoinase	-7.0	b0512
allD	ureidoglycolate dehydrogenase	-6.0	b0517
araA	L-arabinose isomerase	-5.6	b0062
citC	citrate lyase synthetase	-6.2	b0618
citD	citrate lyase acyl carrier protein	-5.6	b0617
citG	triphosphoribosyl-dephospho-CoA synthase	-5.5	b0613
citX	apo-citrate lyase phosphoribosyl-dephospho-CoA transferase	-5.9	b0614
eutG	putative alcohol dehydrogenase in ethanolamine utilization	-5.7	b2453
eutM	putative structural protein, ethanolamine utilization microcompartment	-6.1	b2457
eutN	putative carboxysome structural protein	-6.6	b2456
fdrA	putative acyl-CoA synthetase FdrA	-6.2	b0518
frdA	fumarate reductase flavoprotein subunit	-5.6	b4154
frdB	fumarate reductase iron-sulfur protein	-5.3	b4153
frdC	fumarate reductase membrane protein FrdC	-6.1	b4152
glxR	tartronate semialdehyde reductase 2	-7.7	b0509
hycA	regulator of the transcriptional regulator FhlA	-5.3	b2725
hycB	formate hydrogenlyase subunit HycB	-6.0	b2724
ompW	outer membrane protein W	-6.4	b1256
ssnA	putative aminohydrolase	-5.3	b2879
tdcA	DNA-binding transcriptional activator TdcA	-6.2	b3118
tdcB	catabolic threonine dehydratase	-5.5	b3117
ulaA	L-ascorbate specific PTS enzyme IIC component	-6.1	b4193
ulaB	L-ascorbate specific PTS enzyme IIB component	-5.8	b4194
ulaC	L-ascorbate specific PTS enzyme IIA component	-5.4	b4195
ybbW	putative allantoin transporter	-6.7	b0511
ygeW	putative carbamoyltransferase YgeW	-6.9	b2870
ygeY	putative peptidase YgeY	-6.1	b2872
ygfK	putative oxidoreductase, Fe-S subunit	-5.8	b2878
yhjX	putative pyruvate transporter	-5.4	b3547
ylbE	DUF1116 domain-containing protein YlbE	-5.6	b4572

[a] FC: Fold Change, N/A: not available

Figure Legends

Fig. 1 Virulence factors present in select clinical UPEC strains. Presence or absence of these
42 genes was determined via BLAST ($\geq$80% coverage and ($\geq$90% identity). White indicates
absence of gene, while black indicates presence. Color coding on gene names identifies the
function of each virulence factor. Goldenrod is iron acquisition, green is motility, blue is adhesins,
while red is toxins. Sequence types were determined computationally and reported in Sintsova et
al (13).

Fig. 2 Murine colonization and inflammatory response of selected clinical UPEC strains.
CBA/J mice were transurethrally inoculated with 10^8 CFU of the indicated strain (HM43, HM56
of HM86). (A) Bacterial burden was enumerated from urine, bladder and kidneys 48 hours post
infection. Symbols are individual animals and bars represent the median. Dotted line indicates
limit of detection. A two-tailed Mann-Whitney test was performed to test significance, ** P <0.01,
***P <0.005 ****P <0.0001. (B) Inflammation was assessed using histopathological analysis of
stained thin sections of each specified organ. Inflammation was scored on a 0-3 scale, with zero
being no inflammation, and 3 being severe. Mice were mock-infected with PBS to serve as a
negative control. Symbols are individual animals and bars represent the median.

Fig. 3. UPEC gene expression during mouse and human infections is highly correlated.
Normalized gene expression (log$_2$ TPM) for three UPEC strains: HM43 (A), HM56 (B), and HM86
(C) was compared between LB culture and human infection (hUTI | LB); urine culture and human
infection (hUTI | urine); and mouse infection and human infection (hUTI | mUTI). Pearson
correlation coefficient (r) is shown in top left corner of each plot.

Fig. 4. Host-associated gene expression is distinct from that of *in vitro* culture. (A) Principal
component analysis of normalized gene expression of 3 clinical UPEC strains during human
infection (hUTI), during mouse infection (mUTI), during *in vitro* LB culture (LB), and *in vitro*
urine cultures (Urine). (B) Principal component analysis of normalized r-protein expression of 3
clinical UPEC strains during human infection (hUTI), during mouse infection (mUTI), during *in
vitro* LB culture (LB), and *in vitro* urine cultures (Urine). (C) Percent contribution from r-proteins
to PC1 and PC2 from PCA analysis shown in (A). (D). Hierarchical clustering of *in vitro*, murine
and patient samples based on normalized gene expression of genes present in all 3 strains
(n=3266).

**Figure 5. Differential expression analysis reveals infection-specific gene expression
responses.** (A) The DESeq2 R package was used to compare UPEC gene expression during m
UTI to that in patients. Each UPEC strain was considered an independent replicate (n = 3). Genes
were considered up-regulated (down-regulated) if log$_2$ fold change in expression was higher
(lower) than 1 (vertical lines), and P value < 0.05 (horizontal line). Using these cutoffs, we
identified 30 upregulated genes and 145 downregulated genes in patients. GO/pathway analysis
showed a number of transporters and catabolic enzymes among differentially expressed genes. (B
and C) Identification of genes differentially expressed during infection (hUTI or mUTI) compared
to LB (B) or urine (C). Genes were considered to be up/downregulated in both mouse and human
if log$_2$ fold change was higher/lower than 1, and P value < 0.05 in both cases. Genes that were
upregulated during infection when compared to LB (B) or urine (C) are shown in red, genes that
were downregulated during infection compared to LB are shown in blue.

**Figure 6. Gene expression of virulence factors as well as metabolic machinery is highly
consistent between mouse model of UTI and human UTI.** Normalized gene expression of iron
acquisition operons (A), adherence genes (B), and metabolic pathways (C) for HM43 was
compared between LB and human infection (LB | hUTI), urine and human infection (urine |
hUTI), and mouse UTI and human UTI (mUTI | hUTI).

Fig.1

Gene	Sequence Type Novel HM43	538 HM56	127 HM86
entB	Present	Present	Present
cirA	Present	Present	Present
fepA	Present	Present	Present
fiu	Present	Present	Present
iha	Absent	Absent	Absent
ireA	Absent	Absent	Absent
iroB	Absent	Present	Absent
iroN	Absent	Present	Absent
irp1	Present	Present	Present
fyuA	Present	Present	Present
iucC	Absent	Absent	Present
fitA	Present	Present	Present
iutA	Absent	Present	Present
fhuA	Present	Present	Present
chuA	Present	Present	Present
hma	Absent	Present	Present
sitA	Absent	Present	Present
tonB	Present	Present	Present
cheW	Present	Present	Present
cheY	Present	Present	Present
flgM	Present	Present	Present
motA	Present	Present	Present
motB	Present	Present	Present
csgA	Present	Present	Present
aufA	Present	Absent	Present
focA	Absent	Present	Present
c1936	Present	Present	Present
c2395	Absent	Present	Present
fimH	Present	Present	Present
papG	Absent	Present	Present
pixC	Absent	Present	Absent
ppdD	Present	Present	Absent
yadN	Present	Present	Present
yehA	Present	Present	Present
ygiL	Present	Present	Present
yfcV	Present	Present	Present
cnf1	Absent	Absent	Present
hlyA	Absent	Absent	Present
picU	Absent	Absent	Absent
sat	Absent	Absent	Absent
tosA	Absent	Absent	Absent
vat	Absent	Present	Present

Legend:
- ■ Present
- □ Absent
- Iron Acquisition
- Motility
- Adhesin
- Toxin

Fig.2

Fig. 3

Fig. 4

Fig. 5

Fig. 6