The transcriptional landscape of Venezuelan equine encephalitis virus (TC-83) infection

Short title: Single-cell transcriptional dynamics of alphavirus infection

Zhiyuan Yao[1, 2, ¶], Fabio Zanini[3, 4, ¶, *], Sathish Kumar[1, 2], Nuttada Panpradist[1, 2, 3, 5], Avery Muniz[1, 2],

Stephen R. Quake[3,6,7], and Shirit Einav[1, 2, *]

[1]Department of Microbiology and Immunology, Stanford University School of Medicine,

Stanford, CA

[2]Division of Infectious Diseases and Geographic Medicine, Department of Medicine, Stanford

University School of Medicine, Stanford, CA, USA

[3]Department of Bioengineering, Stanford University, Stanford, CA, USA

[4]Lowy Cancer Research Centre, University of New South Wales, Sydney, Australia

[5]Department of Bioengineering, University of Washington, Seattle, WA, USA

[6]Chan Zuckerberg Biohub, San Francisco, CA, USA

[7]Department of Applied Physics, Stanford University, Stanford, CA, USA

[*]Corresponding authors: Shirit Einav (email: seinav@stanford.edu) and Fabio Zanini

(fabio.zanini@unsw.edu.au)

[¶]: these authors contributed equally

Abstract

Venezuelan Equine Encephalitis Virus (VEEV) is a major biothreat agent that naturally causes

outbreaks in humans and horses particularly in tropical areas of the western hemisphere, for

which no antiviral therapy is currently available. The host response to VEEV and the cellular

factors this alphavirus hijacks to support its effective replication or evade cellular immune

responses are largely uncharacterized. We have previously demonstrated tremendous cell-to-cell

heterogeneity in viral RNA (vRNA) and cellular transcript levels during flaviviral infection using

a novel virus-inclusive single-cell RNA-Seq approach. Here, we used this unbiased, genome-

wide approach to simultaneously profile the host transcriptome and vRNA in thousands of single

cells during infection of human astrocytes with the live-attenuated vaccine strain of VEEV. Host

transcription was profoundly suppressed, yet "superproducer cells" with extremely high vRNA

abundance emerged during the first viral life cycle and demonstrated an altered transcriptome

relative to both mock-infected cells and cells with high vRNA abundance harvested at later time

points. Additionally, cells with increased structural-to-nonstructural transcript ratio exhibited

upregulation of intracellular membrane trafficking genes at later time points. Loss- and gain-of-

function experiments confirmed pro- and antiviral host factors among the products of transcripts

that positively or negatively correlated with vRNA abundance, respectively. Lastly, comparison

with single cell transcriptomic data from other viruses highlighted common and unique pathways

perturbed by infection across evolutionary scales. This study provides a high-resolution

characterization of the VEEV-host interplay, identifies candidate antiviral targets, and establishes

a comparative single-cell approach to study the evolution of virus-host interactions.

9
10

Author Summary

Little is known about the host response to Venezuelan Equine Encephalitis Virus (VEEV) and the cellular factors this alphavirus hijacks to support effective replication or evade cellular immune responses. Monitoring dynamics of host and viral RNA (vRNA) during viral infection at a single-cell level can provide insight into the virus-host interplay at a high resolution. Here, a single-cell RNA sequencing technology that detects host and viral RNA was used to investigate VEEV-host interactions during the course of infection of human astrocytes. Virus abundance and host transcriptome were heterogeneous across cells from the same culture. Subsets of differentially expressed genes, positively or negatively correlating with vRNA abundance, were identified and subsequently validated as proviral and antiviral factors, respectively. In the first replication cycle, "superproducer" cells exhibited rapid increase in vRNA abundance and unique gene expression patterns. At later time points, cells with increased structural-to-nonstructural transcript ratio demonstrated upregulation of intracellular membrane trafficking genes. Lastly, comparing the VEEV dataset with published datasets on other RNA viruses revealed unique and overlapping responses across viral clades. Overall, this study improves the understanding of VEEV-host interactions, reveals candidate targets for antiviral approaches, and establishes a comparative single-cell approach to study the evolution of virus-host interactions.

71
72

13
14

Introduction

For more than a century, Venezuelan Equine Encephalitis Virus (VEEV), a member of the

Alphavirus genus, has been the causative agent of outbreaks of febrile neurological disease in

both animals and humans in Central and South America (1,2). The incidence of VEEV infection

is underestimated since early symptoms are non-specific (2). While typically transmitted via a

mosquito bite, VEEV is also infectious as an aerosol, hence it is considered a major bioterrorism

threat (3). To date, no US FDA approved drugs or vaccines against VEEV are available. A

deeper understanding of VEEV biology in human cells is required to advance the development of

effective countermeasures against VEEV.

Because VEEV is a biosafety level 3 pathogen, TC-83, a live-attenuated vaccine strain, is

commonly used for research purposes (4). Although attenuated, VEEV TC-83 replicates rapidly:

viral protein production is observed as early as 6 hours postinfection (hpi) of human astrocytoma

cells (U-87 MG) at multiplicity of infection (MOI) of 2, and over 10^{10} copies of intracellular viral

RNA (vRNA) can be detected by 24 hpi (5). It remains unknown, however, whether a large

number of cells, each producing a small number of virions, or a few "superproducer" cells drive

this effective virus production. Productive replication is associated with profound shutdown of

host gene transcription (6). Nevertheless, since the virus relies on cellular machineries, it is

important to identify which host factors are "spared" from this shutdown, as they may represent

essential factors for effective viral replication.

The genome of VEEV is an ~11.5 kb single-stranded positive-sense RNA. The genomic RNA

contains two domains. The 5' two-thirds of the genome constitutes the first open reading frame

(ORF), which encodes the nonstructural (ns) proteins required for viral RNA synthesis (nsP1-4).

The 3' one-third of the genome is the structural protein domain. The structural proteins (capsid,

envelope glycoproteins E1-3, 6k, and transframe (TF) protein) are translated from a second ORF

that is expressed through the production of a subgenomic mRNA from an internal promoter in

the negative-strand RNA replication intermediate and function in the assembly of new virions

and their attachment and entry into cells (7). While the stoichiometry of the genomic and

subgenomic transcripts in the setting of VEEV infection has not been characterized, the

transcription of the subgenomic RNA of a related alphavirus, Sindbis virus (SINV), was shown

to be ~3-fold higher than the genomic RNA during late stages of the viral lifecycle (8,9),

supporting a switch towards increased synthesis of structural proteins required for virion

formation over nonstructural proteins required primarily for viral RNA replication (10,11).

The understanding of the alphavirus life cycle is largely based on studies conducted with the

non-pathogenic SINV and Semliki forest virus (SFV). Alphaviruses enter their target cells via

clathrin-mediated endocytosis and release their nucleocapsid into the cytoplasm via fusion with

endosomal membranes, followed by translation and processing of the nonstructural polyprotein

(12). Viral RNA replication occurs within membrane invaginations called spherules that are

thought to be derived from the plasma membrane, endoplasmic reticulum and late endosomes

and are subsequently incorporated into type 1 cytopathic vacuoles (CPV)-I composed of

modified endosomes and lysosomes (13–16). Production of genomic RNA and subsequently

subgenomic RNA are followed by polyprotein translation and processing. The current model of

infectious alphavirus production suggests that the genomic RNA is packaged by the capsid in the

cytoplasm, and that the viral glycoproteins traffic via membrane structures, presumed to be

*trans*Golgi-derived (CPV-II), to budding sites on the plasma membrane, followed by membrane

curving and scission, facilitating envelopment of the nucleocapsid (16–18).

Although VEEV is predicted to extensively interact with cellular factors to effectively replicate

and evade cellular immune responses, like other small RNA viruses, little is known about these

interactions. A recent small interfering RNA (siRNA) screen revealed a requirement for actin-

remodeling pathway proteins including ARF1, RAC1, PIP5K1-α, and ARP3 in VEEV infection

and specifically in promoting viral glycoprotein transport to the plasma membrane (19). Various

other cellular factors, such as DDX-1 and -3 (20), have been reported to have proviral functions,

whereas IFITM3 (21) and members of the PARP protein family (22), were shown to be antiviral

factors. Nevertheless, to the best of our knowledge, the interplay between VEEV and the human

host has not been studied to date via an unbiased, genome-wide approach.

Single cell RNA sequencing (scRNA-Seq) has demonstrated utility for understanding the

heterogeneity of both viral and cellular transcriptome dynamics at a high resolution. We have

recently developed virus-inclusive single-cell RNA-Seq (viscRNA-Seq), an approach to

simultaneously profile host and viral gene expression in thousands of single cells (23). The

studies we and others have conducted in cell lines infected with dengue (DENV), Zika (ZIKV),

influenza A (IAV) (24,25) and West Nile (WNV) viruses (26) and our results in samples from

DENV-infected patients (27) revealed a tremendous cell-to-cell heterogeneity in both vRNA

abundance and levels of host factors that support or restrict infection. Moreover, we have

demonstrated the utility of this approach in identifying novel cellular factors that support or

restrict viral infection (23). We have therefore hypothesized that studying VEEV (TC-83)

transcriptome dynamics at a single cell resolution may overcome challenges related to the high

viral replication rate, thereby highlighting specific transcriptomic signatures above the

suppressed transcriptional landscape and identifying novel cellular factors that support or restrict

VEEV replication.

We conducted a longitudinal study of virus-host cell interactions across 24 hours of VEEV

infection in U-87 MG cells via viscRNA-Seq. We detected extreme heterogeneity in vRNA

abundance and host transcriptome across cells from the same culture. To overcome the challenge

presented by this uneven and rapid viral replication, we stratified cell populations based on

vRNA abundance rather than time postinfection and correlated cellular gene expression with

both (i) total vRNA and (ii) the ratio of total (genomic + subgenomic) to genomic vRNA. These

approaches enabled identification of genes whose expression is altered during VEEV infection,

many of which were then confirmed via loss-of-function and gain-of-function experiments to

have pro- and antiviral roles, respectively. Moreover, we revealed a small population of

"superproducer cells" that drives the rapid increase in vRNA in the first replication cycle and a

cell population that harbors excess of the structural over nonstructural viral ORFs at late stages

of viral infection, both associated with distinct host gene expression patterns. Lastly, comparison

of the VEEV dataset with published data on other RNA viruses revealed unique and overlapping

host gene responses across viral clades, highlighting the utility of comparative single-cell

transcriptomics.

Materials and methods

Cells

U-87 MG and BHK-21 cell lines were obtained from ATCC (Manassas, VA). Cells were grown

in Dulbecco's Modified Eagle's medium (DMEM, Mediatech, Manassas, VA), supplemented

with 1% Penicillin-Streptomycin solution, 1% L-glutamine 200 mM (Thermo Fisher Scientific,

Waltham, MA) and 10% Fetal Bovine Serum (FBS, Omega Scientific, INC, Tarzana, CA). Cells

were maintained in a humidified incubator with 5% CO_2 at 37 ˚C. Cells were tested negative for

mycoplasma by the MycoAlert mycoplasma detection kit (Lonza, Morristown, NJ).

29
30

Plasmids and virus constructs

The plasmids encoding infectious VEEV TC-83 with a GFP reporter (VEEV TC-83-Cap-eGFP-

Tav, hereafter VEEV-TC-83-GFP) or a nanoluciferase reporter (VEEV TC-83-Cap-NLuc-Tav,

hereafter VEEV-TC-83-nLuc), were a gift from Dr. William B. Klimstra (Department of

Immunology, University of Pittsburgh) (28). Open reading frames (ORFs) encoding 11 hits were

selected from the Human ORFeome library of cDNA clones (Open Biosystems) (29) and

recombined into a FLAG (for FLAG tagging) vector using Gateway technology (Invitrogen).

Virus production

Viral RNA (vRNA) (VEEV-TC-83-GFP or nLuc) was transcribed *in vitro* from cDNA plasmid

templates linearized with MluI via MegaScript Sp6 kit (Invitrogen #AM1330) and electroporated

into BHK-21 cells. VEEV was harvested from the supernatant 24 hours postelectroporation,

clarified from cell debris by centrifugation, and stored at -80 °C. Virus stock titers were

determined by standard BHK-21 cell plaque assay, and titers were expressed as PFU/ml.

Infection assays

U-87 MG cells were infected with VEEV-TC-83-GFP at various MOIs (0, 0.1, and 1) and

harvested at various time points postinfection. For the functional screens, U-87 MG cells were

infected with VEEV-TC83-nLuc in 8 replicates at MOI of 0.01. Overall infection was measured

at 18 hpi via a nanoluciferase assay using a luciferin solution obtained from the hydrolysis of its

O-acetylated precursor, hikarazine-103 (prepared by Dr. Yves Janin, Pasteur Institute, France)

as a substrate (30,31).

Loss-of-function assays

siRNAs (1 pmol) were transfected into cells using lipofectamine RNAiMAX transfection reagent

(Invitrogen) 96 hours prior to infection with VEEV-TC-83-nLuc at MOI of 0.01. Custom

Cherry-Pick ON-TARGETplus siRNA library against 11 genes was purchased from Dharmacon

(see Supplementary Table 1 for gene and siRNA sequence details).

Gain-of-function assays

Individual plasmids encoding 11 human genes or empty control vector were transfected

individually into U-87 MG cells with lipofectamine-3000 (Invitrogen) 48 hours prior to infection

with VEEV-TC-83-nLuc at MOI of 0.01.

Viability assays

Viability was measured using alamarBlue reagent (Invitrogen) according to the manufacturer's

protocol. Fluorescence was detected at 560 nm on an Infinite M1000 plate reader (Tecan).

Detection of infected cells using VEEV-specific capture oligo

To optimize the viscRNA-Seq protocol for a wide dynamic range of vRNA amount per VEEV-

infected cells, we designed and screened eight oligo capture (**S1 Table**).

To screen these capture oligo, we first generated cDNA from VEEV-infected cells in the

presence of each or combinations of VEEV-specific capture oligo. Specifically, 30 pg of both

vRNA and cellular RNA purified from VEEV-infected cells was reverse-transcribed to cDNA in

a reaction containing SuperScript™ IV reverse transcriptase, 1X First Strand buffer (Invitrogen),

5 mM DTT, 1 M betaine, 6 mM $MgCl_2$, 1 µM oligo dT and each or combinations of 100 nM

reverse VEEV oligo capture. Subsequently, cDNA underwent 21-cycle PCR amplification using

ISPCR primers. cDNA was then purified using Ampure XP beads (Beckman Coulter) at the ratio

of 0.8 and eluted in 15 µL EB buffer. Fragments of purified, concentrated cDNA were visualized

and quantified using bioanalyzer (DNA High Sensitivity kit, Agilent Technologies). To quantify

the amount of vRNA captured by each or combinations of oligo capture, these purified cDNA

were also subjected to qPCR (Hot-start OneTaq (New England Biolabs), 1x Standard Taq buffer,

1x Evagreen (Biotium), forward primer: ATTCTAAGCACAAGTATCATTGTAT and reverse

primer: TTAGTTGCATACTTATACAATCTGT located upstream of all the capture oligos.

VEEV_1 and VEEV_2 yielded the highest copies of viral cDNA and did not generate significant

primer dimers. Therefore, this combination of the capture oligo was selected for downstream

experiments.

Single cell sorting

At each time point, cells were trypsinized for 10 min, spun and resuspended in 1 mL fresh media.

Within 15 min, cells were pelleted again and resuspended in 2 ml 1X phosphate-buffered saline

(PBS) buffer at a concentration of 10^6 cells per ml. Cells were filtered through a 40 μm filter into

a 5 ml FACS tube and sorted on a Sony SH800 sorter using SYTOX™ Blue dead cell stain

(ThermoFisher) to distinguish living cells from dead cells and debris. VEEV harboring cells

were sorted based on GFP signal. Cells were sorted into 384-well PCR plates containing 0.5 μl

of lysis buffer using 'Single cell' purity mode. A total of 12 384-well plates of single cells were

sorted for the VEEV time course.

Lysis buffer, reverse transcription, and PCR

To capture and amplify both mRNA and vRNA from the same cell, the Smart-seq2 protocol was

adapted (Picelli et al., 2014). All volumes were reduced by a factor of 12 compared to the

original protocol to enable high-throughput processing of 384-well plates. ERCC spike-in RNA

was added at a concentration of 1:10 of the normal amount. The lysis buffer contained 100nM of

oligo-dT primer, 100 mM of virus specific capture oligo mix (i.e. VEEV_1 and VEEV_2) to

capture the positive-stranded virus RNA.

Other virus-specific primers and higher primer concentrations were tested but resulted in a large

fraction of primer dimers. In order to reduce interference between the virus-specific primer and

the Template Switching Oligo (TSO) used to extend the RT products, a 5'-blocked biotinylated

TSO was used at the standard concentration. Reverse transcription (RT) and PCR of the cDNA

were performed in a total volume of 1 µl and 2.5 µl for each well respectively. The resulting

cDNAs were amplified for 21 cycles. Lambda exonuclease was added to the PCR buffer at a

final concentration of 0.0225 U/µl and the RT products were incubated at 37 ˚C for 30 min

before melting the RNA-DNA hybrid (as it was observed that this reduced the amount of low-

molecular weight bands from the PCR products). The cDNA was then diluted 1 to 7 in EB buffer

for a final volume of 17.5 µl. All pipetting steps were performed using a Mosquito HTS robotic

platform (TTP Labtech).

cDNA quantification

To quantify the amount of cDNA in each well after PCR, a commercial fluorometric assay was

used (ThermoFisher Quant-It™ Picogreen). Briefly, 1 µl of cDNA and 50 µl of 1:200 dye-buffer

mix were pipetted together into a flat-bottom 384-well plate (Corning 3711). For each plate, six

wells were used as standard wells. 1 µl dd H2O was added into one standard well as blank. The

standard solutions were diluted into 5 concentrations (0.1, 0.2, 0.4, 0.8, 1.6 ng/µl) and added 1µl

into the remaining 5 standard wells. The plate was vortexed for 2 min, centrifuged, incubated in

the dark for 5 min, and measured on a plate reader at wavelength 550 nm. cDNA concentrations

were calculated via an affine fit to the standard wells.

Library preparation and sequencing

For each time point, one plate was sent for library preparation and sequencing. In total, 6 plates

(2304 cells) were prepared. Sequencing libraries were prepared using the illumina Nextera XT

kit according to the manufacturer's instructions, with the following exceptions: (1) we used a

smaller reaction volume (around 1 µl per cell); (2) we chose a slightly higher cDNA

concentration (0.4 ng/µl) as input, to compensate for the lack of bead purification upstream; (3)

we used the commercial 24 i7 barcodes and the 64 new i5 barcode sequences. We noticed a low

level of cross-talk between these barcodes, indicated by up to five virus reads found in a few

uninfected cells. However, considering that a sizeable fraction of cells in the same sequencing

run (late infected and high MOI) had thousands of virus reads, the amount of cross-talk between

barcodes appears to be of the order of 1 in 10,000 or less. We used Illumina Novaseq sequencer

for sequencing.

Bioinformatics pipeline

Sequencing reads were mapped against the human GRCh38 genome with supplementary ERCC

sequences and TC-83-VEEV-GFP genome using STAR Aligner (32) . Genes were counted using

htseq-count (33). The Stanford high performance computing cluster Sherlock 2.0 was used for

the computations. Once the gene/virus counts were available, the downstream analysis was

performed on laptops using the packages Seurat (34) and singlet (https://github.com/iosonofabio/

singlet), as well as custom R and Python scripts. Ggplot2 (35), matplotlib (36) and seaborn (37)

were used for plotting.

For the mutational analysis, all reads mapping to VEEV were extracted from all cells with a

unique identifier of the cell of origin, and all four possible alleles at each nucleotide were

counted by custom scripts based on pysam (https://github.com/pysam-developers/pysam) and

wrapped in an xarray Dataset (38). The analysis was restricted to infected cells with an average

of 100 or more reads per viral genomic site to reduce shot noise.

Comparison with flaviviruses was performed as follows. First, host genes with similar expression

(within a factor of 10) in counts per millions (cpm) were identified. Within that class,

correlations with vRNA for VEEV, DENV, ZIKV were computed separately. Host factors with

the highest discrepancies between pairs of viruses were identified. For Figs 5A-C, a gene was

chosen from the most discrepant genes exemplifying the different behaviors observed and the

cells were scattered using vRNA abundance and gene expression axes, and colored by virus. For

Fig 5D, the host counts for each gene from all three experiments (in cpm) were added and

fractions belonging to each experiment were computed. Because the sum is constrained to be

100%, ternary plots could be used for plotting the three different fractions in two dimensions.

For figs 5E-F, for each gene shown we computed its percentile in correlation with DENV and

ZIKV vRNA, i.e. the percentage of other host genes with a correlation less than this focal gene.

This transformation emphasizes the top correlates/anticorrelates against batch effects and

different multiplicities of infection in the DENV and ZIKV experiments. For figs 5G-I,

published tables of counts and metadata were downloaded from links present in each publication,

normalized to counts per millions, and filtered for low-quality cells. We computed the

correlation of host gene expression and vRNA in each experiment, then features were selected

that had a high rank in at least one virus and the selected correlation coefficients were centered

and normalized between -1 and 1 for each virus to enable meaningful cross-experiment

comparison. Principal Component Analysis (PCA), UMAP, similarity graphs, and Leiden

clustering (Traag et al. 2019) were computed and plotted.

Cell selection and normalization

The criteria to select cells were as follows: total reads > 300,000, gene counts > 500 and a ratio

of ERCC spike-in RNA to total reads ratio < 0.05. Based on these criteria, 2004 out of 2301 cells

were selected for downstream analysis. Due to the high viral copies of VEEV in cells infected

for 12 and 24 hrs (more than 10%), traditional normalization (dividing by total reads) caused a

bias which underestimated the expression of host genes. To avoid this, we normalized gene

counts to ERCC total reads, since these are not affected by the virus. Each gene count column

(including virus reads) was thus divided by ERCC total reads and then log transformed.

Data and code availability

The single cell RNA-Seq data for this study is available on GEO at submission number:

GSE145815 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE145815). The code used

in the computational analyses can be found at https://github.com/saberyzy/VEEV-single_cell.

Processed count and metadata tables are also available on FigShare at

https://figshare.com/articles/Untitled_Item/11874198.

Results

viscRNA-Seq reveals cell-to-cell heterogeneity in VEEV and host gene expression.

To characterize the relation between viral and host cell transcriptional dynamics over the course

of VEEV infection, human astrocytoma cells (U-87 MG) (39) were infected with VEEV (TC-83,

attenuated vaccine strain) conjugated to GFP (28) at MOIs of 0.1 and 1 or mock infected, and

harvested at six time points: 0.5, 1.5, 4, 6, 12, and 24 hpi **(Fig 1A).** Single cells were then

isolated and processed by viscRNA-Seq, as described previously (23). Since the VEEV RNA is

polyadenylated, it can be captured by the standard poly-T oligonucleotide that hybridizes with

host transcripts. Nevertheless, to improve vRNA capture and ensure coverage at the 5' end of the

viral genome, two specific viral capture oligonucleotides, at positions 352 and 1,742 of the

VEEV genome, were added to the reaction (see Methods). In total, 4608 cells were processed, of

which 2301 cells were sequenced with approximately 1 million reads/cell **(S1A Fig).** 2004 cells

passed quality controls and were analyzed (see Methods).

To identify a proper cutoff for defining infected cells, we analyzed both GFP signal and vRNA

reads. During cell sorting (the first step of viscRNA-seq) the GFP signal was recorded using the

fluorescein (FITC) gate, enabling measurement of cellular GFP expression levels. The GFP

signal was comparable in cells harboring 1 to 1000 viral reads, yet it sharply increased in cells

harboring over 1000 viral reads **(Fig 1B)**. The lower sensitivity of GFP signal relative to viral

reads is likely due to the lag of protein expression after RNA synthesis, and indicates that virus

reads can be used as an effective indicator for VEEV infection. Next, we sought to define a

cutoff to distinguish infected from bystander cells (uninfected but derived from the sample that

was exposed to the virus). We set multiple cutoffs between 1 and 100 viral reads, selected only

cells with viral read number greater than these cutoffs, and calculated the correlation coefficient

between GFP expression and viral reads **(S1B Fig)**. The correlation between GFP expression and

viral reads first increased with the cutoffs and then stabilized once the cutoff reached 10 viral

reads, with correlation coefficients greater than 0.8 via both Spearman's and Pearson

correlations. We therefore defined the presence of 10 or more viral reads as the cutoff to

distinguish VEEV-infected from bystander cells. Similar findings were observed upon plotting

the relationship between GFP expression and virus/total reads ratio (vs. raw viral reads) **(S1C**

Fig), indicating that the selected threshold of 10 viral reads (or 0.00001 virus/total reads) is not

affected by differences in sequencing depth between cells.

The fraction of VEEV-infected cells increased with both time and MOI and saturated at 12 and

24 hpi with MOI 1 and 0.1, respectively **(Fig 1C).** A rapid increase in the ratio of both viral/total

reads and GFP expression was observed within single cells over time **(Fig 1D)**. Notably, the

distributions of virus/total reads and GFP expression were particularly wide at 12 hpi when

analyzing either the entire infected cell population or infected cells separated by the two MOIs

(S1D and S1E Figs). At 24 hpi, the observed increase in vRNA reads was associated with a

decline in cellular transcripts. The normalized cellular mRNA reads (calculated by dividing the

absolute number of reads by the sum of External RNA Controls Consortium (ERCC) spike-in

reads) declined in the infected cell group at 24 hpi relative to the corresponding mock infected

cell group and the same infected cell group at 12 hpi (**Fig 1E**). To avoid an artificial decline in

host gene reads in cells with high vRNA abundance, rather than normalizing cellular gene reads

by the total reads, we normalized by ERCC reads for most downstream analyses. This

transformation is akin to an estimate of the actual number of mRNA molecules for each gene (up

to a constant factor).

Altered expression of cellular factors and pathways during VEEV infection.

The wide distributions of virus/total reads observed at 12 hpi suggested that to more precisely

characterize the phenotype of cells from VEEV-infected samples, cells should be divided based

on the virus/total read content rather than time postinfection or MOI. To identify host genes

whose expression is altered during VEEV infection, we integrated differential gene expression

and correlation analyses. First, we combined cells harvested at different time points. Since the

GFP signal started to increase significantly with a virus/total read ratio greater than 0.001 (**S1C**

Fig), we divided cells into the following three groups based on this cutoffs: infected cells with

high vRNA (>0.001 virus/total reads), infected cells with low vRNA (<0.001 virus/total reads),

and mock-infected controls (**S2A Fig**). Since GFP expression and viral reads correlated well in

the high vRNA group, we focused on differences between the high vRNA cell group and the

mock-infected group. Computing differential expression at the distribution level (Mann-Whitney

U test) revealed 1734 host genes, whose expression level significantly differed between the high

vRNA group and the mock-infected group. To test the robustness of the population division, we

applied a set of cutoffs (ranging from 0.0001 virus/total reads to 0.01 virus/total reads) and

computed differential expression between the high vRNA group and the mock-infected controls

based on each of these cutoffs **(S2B Fig)**. The number of differentially expressed genes (DEGs)

identified increased up to a cutoff of 0.001 virus/total reads and then plateaued. Moreover, DEGs

identified by a cutoff of 0.001 largely overlapped (over 90%) with those detected with higher

cutoffs, confirming that the cutoff of 0.001 is robust in distinguishing between infected cells with

high and low vRNA abundance. We predicted that differential expression of some genes might

be related to time effect resulting from differences in incubation duration rather than from viral

infection. To control for such confounders, we calculated Spearman's correlation coefficients

between gene expression and time postinfection. Genes whose expression was similarly altered

over time between infected and mock-infected cells were thought to represent time effect. 1707

of the 1734 DEGs between the high vRNA and mock-infected groups passed this additional filter

(Fig 2A).

In parallel, we computed Spearman's rank correlation coefficients between gene expression and

vRNA abundance across all cells, as done previously for flaviviruses (23). Our data indicate that

the majority of host genes are negatively correlated with vRNA abundance **(S2C Fig)**.

Stratifying host genes by expression level in mock-infected cells indicated a stronger negative

correlation for highly expressed genes with vRNA abundance **(S2D Fig)**, suggesting that cellular

functions relying on highly expressed genes are more vulnerable to VEEV infection. To identify

genes that are both differentially expressed between infected and mock-infected cells and

correlated with vRNA, we computed the intersection between the 1707 DEGs with the top 600

genes that either positively (n=300) or negatively (n=300) correlated with vRNA. 263

overlapping genes emerged from this analysis (**Fig 2A**).

Gene Ontology (GO) enrichment analysis of these 263 genes via metascape (40) highlighted

metabolism of RNA as the most enriched molecular function term (**Fig 2A**). Shown in **Figs 2B,**

2C and S2E are representative genes that were overexpressed in high vRNA cells vs. mock-

infected and low vRNA and positively correlated with vRNA (TNFAIP3), underexpressed and

negatively correlated with vRNA (TAF7), or not differentially expressed and were uncorrelated

with vRNA (COPZ2). The expression level of these genes did not change over time in mock-

infected cells, supporting that their altered levels represent actual differences between the groups

rather than a time effect (**S2F Fig**).

Early infected "superproducer'' cells show distinct patterns of host gene expression. During

cell processing, we noticed that 2% of the cells infected with an MOI of 1 at 6 hpi, the duration

of a single cycle of VEEV replication (7,41), showed stronger GFP signals (FITC gate readout >

1000) than the remaining cells in the same condition. To probe the relevance of this unexpected

finding, we specifically sorted these cells. In correlation with their GFP expression, the majority

of these cells harbored ~100-fold higher virus/total reads ratio than the remaining cells in the

same condition, suggesting that once initiated, viral replication proceeded extremely fast in these

"superproducer" cells **(Fig 3A).** 11 cells were defined as "superproducer" cells based on the

following criteria: harboring > 0.001 vRNA/total reads and GFP readout > 1000 at 6 hpi (MOI =

1) (**Fig 3A**). To elucidate whether these "superproducer" cells exhibit a distinct gene expression

pattern, we conducted differential gene expression analysis (Mann-Whitney U test) between

these 11 cells and mock-infected cells as well as low vRNA harboring cells, both harvested at the

same time point (6 hpi). A total of 16 DEGs were identified showing a distinct expression pattern

only in these "superproducers", with representative overexpressed and underexpressed genes

shown in **Fig 3B and Fig 3C**. Notably, these genes were also differentially expressed between

the "superproducer" cells and high vRNA cells harvested at 24 hpi, suggesting that they do not

represent a general response to high vRNA abundance, but rather a unique feature of this cell

population. Among the overexpressed genes were SYTL3, a protein that directly binds to

Rab27A to regulate Rab27-dependent membrane trafficking; KDM3B, a lysine demethylase;

SNX29, a member of the sorting nexin family; and COG5, a component of Golgi-localized

complex that is essential for Golgi function. Among the underexpressed genes were ZMAT5, an

RNA-binding protein belonging to the CCCH zinc finger family of proteins implicated in

antiviral immune regulation (42); VPS37A, a component of the ESCRT-I protein complex; and

AC087343.1, a ribosomal protein L21 pseudogene. These findings provide evidence that a small

subset of "superproducer" cells largely drives VEEV replication during the first viral life cycle

and demonstrates a distinct gene expression pattern. These results also point to SYTL3, KDM3B,

SNX29 and COG5 as candidate proviral factors, and to ZMAT5, VPS37A and AC087343.1 as

potential antiviral factors.

The expression of genes involved in intracellular membrane trafficking correlates with the

ratio of 3' to 5' vRNA reads.

By including both a poly-T and a 5'-end specific capture oligonucleotides in the viscRNA-Seq,

good read coverage at both ends of the VEEV genome was obtained (**Fig 4A**). We defined 5'

RNA reads as those corresponding to the first 1,700 bases (encoding nonstructural proteins), and

thus derived from the genomic vRNA only, and 3' RNA reads as those corresponding to the last

third of the genome (encoding structural proteins), derived from both the genomic and

subgenomic vRNAs (**Fig 4B**). The stoichiometry of the 3' and 5' RNAs was highly

heterogeneous between cells. While at early stages of infection, the 3' to 5' (structural to

nonstructural) vRNA read ratio (3'/5' read ratio), as defined by these genomic regions, was

below or around 1, at late stages, it reached up to 4 and was correlated with total vRNA

abundance (**Fig 4C**). In contrast, the read ratio between two segments we selected as internal

controls at the 5' end of the vRNA (5'a/5'b read ratio) and between two segments at the 3' end

(3'a/3'b read ratio) did not correlate with the cellular vRNA abundance (**Figs 4D-E**). To test the

hypothesis that differences in vRNA stoichiometry are associated with distinct host responses,

we measured the Spearman's correlation coefficients of all host genes with the 3'/5' read ratio in

the same cell. The resulting histogram distribution curve revealed a tail of host genes whose

expression increased with the 3'/5' read ratio (**Fig 4F**), in contrast to the distribution of host

genes in correlation with the total vRNA reads (**S2C Fig**). Positively correlated genes were

mostly involved in various aspects of intracellular trafficking and included factors previously

reported to be required for VEEV infection via an siRNA screen including ARP3 (19), RAC2, a

paralog of RAC1 [19], and DDX5, a member of the DEAD box family of RNA helicases (20).

Novel factors among the positively correlated genes included factors involved in late endosomal

trafficking (RAB7A (43), the accessory ESCRT factor (BROX) (44), and the SNARE protein

VAMP7 (45)), ER to Golgi trafficking (SEC22B) (46), regulation of secretion (PIP4K2A) (47),

lysosome function and autophagy (LAMP2) (48), actin polymerization (PFN2) (49), and

acidification of intracellular organelles for protein sorting (ATP6V1B2) (50) (**Fig 4G**).

Accordingly, pathway analysis on the top 300 correlated genes identified macroautophagy,

regulated exocytosis, membrane trafficking and vesicle organization as the highly enriched

functions (**Fig 4H**). Notably, these genes were only positively correlated with the 3'/5' read

vRNA ratio and not with the total vRNA reads. These findings indicate that the late stages of

VEEV infection are characterized by heterogeneous stoichiometry of structural (3') and

nonstructural (5') vRNAs and upregulation of intracellular trafficking pathways previously

implicated in assembly and egress of various RNA viruses in cells with an excess of structural

vRNA. Moreover, these results highlight the unique opportunity to discover candidate proviral

factors for VEEV infection by correlating gene expression with specific viral genome

stoichiometry via viscRNA-Seq.

In addition to enabling quantification of the 5' and 3' vRNA reads, the high coverage of the viral

genome provided by viscRNA-Seq revealed rare structural viral read variants. The most common

among these variants was a 36-base gap within the coding region of the 6K protein, whose

presence was predicted to form a stable hairpin structure **(S1 Text and S3 Fig).** While the

biological relevance of this finding remains to be elucidated, and we cannot currently exclude

that this gap could be a result of polymerase errors during library preparation, stable RNA

structures play essential roles in viral replication and tropism across multiple viruses.

Validation of candidate proviral and antiviral factors. Next, we probed the functional

relevance of 11 genes that either strongly or moderately correlated with vRNA abundance for

viral infection. We first conducted loss-of-function screens by measuring the effect of siRNA-

mediated depletion of these 11 individual genes on VEEV infection and cellular viability in U-87

MG cells (**Figs 5A and S4**). Depletion of CXCL3, ATF3, TNFAIP3, and CXCL2, four out of

five genes tested that positively correlated with vRNA abundance via viscRNA-Seq (orange

bars), reduced VEEV infection by more than 40%, respectively, as measured by luciferase assays

18 hpi with a nano-luciferase reporter TC-83 virus and normalized to cellular viability in two

independent screens, suggesting that they are proviral factors. In contrast, depletion of 3 of 6

genes tested that negatively correlated with vRNA (grey bars) enhanced VEEV infection,

suggesting that these proteins may function as antiviral factors. Suppression of PPP2CA

demonstrated no effect on VEEV infection, suggesting that it is either non-essential or not

restricting (possibly due to redundancy in host factors requirement) or that the level of

knockdown was insufficient to trigger a phenotype.

Next, we conducted gain-of-function screens by ectopically expressing the same 11 individual

gene products in U-87 MG cells followed by VEEV infection (**Fig 5B**). Using a cutoff of greater

than 40% change in viral infection normalized to cell viability in two independent screens,

overexpression of most genes resulted in an inverse effect to that observed with the siRNA, i.e. if

knockdown inhibited viral infection, overexpression enhanced it and vice versa. Overexpression

of CXCL3, ATF3, TNFAIP3 and CXCL2 increased VEEV infection, indicating rate limitation

associated with these candidate proviral factors. In contrast, overexpression of the majority of the

anticorrelated gene products reduced VEEV infection, consistent with an antiviral phenotype.

While the transcriptional level of TRMT10C and EIF4A3 anticorrelated with vRNA abundance,

their gene products demonstrated a proviral phenotype. This may either result from regulation of

these genes at the translational level or from downstream effects of these multifunctional genes.

ARRDC3, a member of the arrestin family (51), was positively correlated with vRNA

abundance, yet its depletion increased infection and its overexpression decreased infection, in

contrast with the other four positively correlated genes tested. To probe this discrepancy, we

measured the correlation of ARRDC3 expression with the 5' and 3' vRNA reads separately.

Notably, ARRDC3 reads positively correlated with the 3' vRNA reads but negatively correlated

with the 5' vRNA reads. In contrast, the other four proviral candidates positively correlated with

both the 5' and 3' vRNA reads (**Fig 5C**). This finding suggests that ARRDC3 might have a dual

function during VEEV infection. Together, these findings highlight the utility of viscRNA-Seq in

identifying candidate proviral and antiviral factors.

Comparative viscRNA-Seq analysis across five RNA viruses reveals distinct and common

cellular pathways affected by viral infection.

To define which elements of the host response are unique to VEEV and which are common

across multiple viruses, we first compared the VEEV dataset with our previously published

viscRNA-Seq data on human hepatoma (Huh7) cells infected with DENV and ZIKV (23). Since

the baseline gene expression levels in astrocytes (VEEV) are different from those in hepatocytes

(DENV, ZIKV), we limited the analysis to genes that were similarly expressed (within a 10-fold

change) in uninfected Huh7 and U-87 MG cells. We selected cells with greater than 2 vRNA

reads per million joint (viral + host) reads and monitored how the expression of host genes

changes with increasing vRNA abundance across the three infections. In all three viral

infections, the majority of host genes were not correlated with vRNA abundance. Nevertheless, a

number of host genes exhibited correlations with one or more viruses. Three robust patterns were

identified (**Figs 6A-C**): genes, such as HSPA5, that were upregulated in DENV infection and

downregulated in ZIKV and VEEV infections (**Fig 6A**); genes like NRBF2 that were

upregulated only during ZIKV infection (**Fig 6B**); and genes, such as SERP1, that were

downregulated only in VEEV infection (**Fig 6C**). No genes that were upregulated only in VEEV

infection could be identified. Beyond these general categories, the resulting patterns of viral and

host expression were, however, quite complex.

To circumvent the masking effect of VEEV transcriptional shutdown, we then compared the

genes that positively correlated with the 3'/5' VEEV RNA ratio with those positively or

negatively correlating with DENV or ZIKV vRNA (**Fig 6D**). This analysis revealed genes, such

as BROX, GEM, and RNF114 that are positively correlated with the respective vRNA in all

three viral infections, genes, such as CTSB and SPTLC1 that are positively correlated with 3'/5'

VEEV RNA and ZIKV but not DENV vRNA, and genes that are positively correlated with 3'/5'

VEEV RNA but negatively correlated with DENV and ZIKV vRNA, such as PFN2 and

DPYSL2. In contrast, no large correlations were observed when a comparable number of random

genes were similarly analyzed (**Fig 6E**). Pathway analysis on genes that are positively correlated

with both the 3'/5' VEEV RNA ratio and the two flaviviral RNAs identified ER processing,

glycosylation, SELK (part of Endoplasmic-Reticulum-Associated Degradation), tRNA synthesis,

protein folding, virion assembly, and intracellular transport as the highly enriched functions

(**S5A Fig**). In contrast, cell cycle and apoptosis regulation were the most highly enriched

functions in pathway analysis on genes that were positively correlated with 3'/5' VEEV RNA

ratio but negatively correlated with the two flaviviral RNA (**S5B Fig**). These results provide

evidence that complex temporal dynamics exist across different RNA viral infections, and

highlight both common and unique cellular pathways that are altered by VEEV and flaviviruses.

Next, we expanded our comparative analysis by including published datasets derived from

single-cell transcriptomic studies on different cell lines infected with IAV (24) and WNV (26)

generated via 10x Genomics and Smart-seq2, respectively. Because different cell lines were used

for different viruses, we calculated the ranks of the correlation coefficients between the

expression of each host gene and vRNA for each virus, restricted the selection to the top and

bottom 200 genes, and normalized the results between -1 and 1 for each virus. We then

calculated the network of similarities between genes (52). Uniform Manifold Approximation and

Projection for Dimension Reduction (53) and Leiden clustering (54) of the genes highlighted 8

gene clusters with different expression patterns during various viral infections (**Fig 6F**). To

understand the meaning of these clusters, we performed double hierarchical clustering and

observed that clusters 2, 4, 0, and 3 were upregulated, while clusters 7, 5, 1, and 6 were mostly

downregulated during viral infection (**Fig 6G**). DENV and ZIKV shared clusters for both

upregulation and downregulation, as expected from their evolutionary proximity. The

dendrogram of the five viruses was qualitatively consistent with the known phylogeny as derived

from viral genomic sequences, which could indicate ancestral phenotypic signatures.

Overall, our analysis indicates that although comparing single cell viral infection data across

species, cell lines, and technologies still presents challenges, this approach is informative in

highlighting host genes and pathways that are commonly affected across very different viral

families.

Discussion

We and others have recently characterized the cellular response in virally infected cell lines

(23,24), primary cells (26,55) and patient samples (27) via single-cell RNA-seq approaches.

Moreover, we reported unique and overlapping determinants in the host response to two related

flaviviruses at a single cell resolution (23). Nevertheless, the host transcriptomic response to

infection by alphaviruses, which induces a profound transcriptional shutdown of host genes, has

not been previously characterized at a single cell level, and the single-cell transcriptomic

responses of unrelated viruses have not been compared. By applying viscRNA-Seq to study the

temporal infection dynamics of VEEV (TC-83) in human astrocytes, we revealed large cell-to-

cell heterogeneity in VEEV and host gene expression, transcriptomic signatures in distinct cell

subpopulations, and candidate proviral and antiviral factors, some of which we then validated.

Additionally, we established a role for viscRNA-Seq in comparative evolutionary virology by

demonstrating structural variants within the VEEV genome as well as unique and overlapping

host gene responses across multiple RNA viral clades. These findings provide insights into the

virus-host determinants that regulate VEEV infection and highlight the utility of virus-inclusive

RNA-seq approaches and comparative single-cell transcriptomics.

A prominent feature of VEEV infection is a profound suppression of cellular transcription (6).

Nevertheless, it remained unknown whether this transcriptional shutdown globally affects all

host mRNAs. Computing the distributions of vRNA expression in correlation with 5 groups of

genes, distinguished by the level of gene expression in uninfected cells, demonstrated that highly

expressed genes are more likely to be negatively correlated with vRNA abundance than genes

that are expressed at a lower level. The cellular energy and machinery required to maintain a

high level of gene expression likely play a role in increasing the vulnerability of highly

expressed cellular genes to VEEV-induced transcriptional shutdown.

We have previously reported the utility of viscRNA-Seq in discovering functional transcriptomic

signatures and candidate pro- and antiviral factors of DENV and ZIKV infections (23,27).

Nevertheless, the high replication rate of VEEV and the transcriptional shutdown it induces

challenged our ability to detect alterations in gene expression and identify pro- and antiviral

factors. To overcome these challenges, we used several strategies. First, since the viscRNA-Seq

analysis revealed large differences in vRNA abundance between cells infected with the same

MOI and harvested at the same time point, we stratified cell populations based on vRNA

abundance rather than time postinfection. Integrating differential gene expression and correlation

analyses of vRNA abundance with gene expression across the entire human transcriptome

facilitated the discovery of 263 genes that were both differentially expressed between the high

and mock infected controls and correlated with total vRNA. siRNA-mediated depletion and

overexpression of a subset of these genes revealed that overall, genes involved in cytokine

production, plus ATF3, a transcription factor commonly expressed in response to cellular stress,

and TNFAIP3, an inhibitor of NFκB signaling, demonstrated a phenotype consistent with a rate-

limiting proviral function, whereas a variety of regulatory genes, such as TAF7, were rate-

limiting antiviral factors. ARRDC3, one of 5 genes that were both differentially expressed and

positively correlated with total vRNA, demonstrated a phenotype consistent with antiviral rather

than a proviral effect. Interestingly, when studied in correlation with the individual vRNA

transcripts, ARRDC3, a signaling arrestin family protein and a cargo-specific endosomal

adaptor, was positively correlated with the 3' vRNA but negatively correlated with the 5' vRNA,

suggesting that it may have a proviral effect during later stages and an antiviral effect in earlier

stages of replication. By capturing such complex dynamics and not relying on averaging signals

at distinct time points postinfection for stratification, the viscRNA-Seq approach has an

advantage over bulk sample knockdown or knockout approaches in identifying factors required

for or restrictive of VEEV infection.

The high resolution provided by viscRNA-Seq enabled us to further focus on distinct cell

populations, which facilitated identification of additional transcriptomic signatures. We

discovered a subpopulation of cells demonstrating unusually high viral replication upon

completion of a single cycle of viral replication. Importantly, this cell subpopulation is

associated with host cell gene expression that is distinct from cells harboring lower vRNA at the

same time. It is intriguing to speculate that overexpression of the identified hits involved in

intracellular membrane trafficking (such as SYTL3, SNX29 and COG5) concurrently with

underexpression of factors implicated in antiviral immune responses (such as ZMAT5) in this

cell population drive the rapid increase in viral replication during the first viral lifecycle.

To further increase the resolution of our analysis, we took advantage of the ability of viscRNA-

Seq to detect the two VEEV transcripts. A prior study on IAV has detected different levels of

various segments of the viral genome across cells and investigated how this finding relates to

successful virion production (24). Similarly, analysis of the stoichiometry of the 5' and 3' RNA

reads of VEEV, a non-segmented virus, revealed a large cell-to-cell heterogeneity. Moreover, the

3'/5' vRNA ratio substantially increased at late stages of infection, consistent with a previous

report in another alphavirus, SINV (10). Remarkably, the histogram distribution curve of the

Spearman's correlation coefficients of all host genes with the 3'/5' read ratio in the same cell

revealed a long tail of host genes whose expression increased with the 3'/5' read ratio. Our

findings indicate that these changes in stoichiometry of the vRNA transcripts during late stages

of VEEV infection are associated with upregulation of distinct genes, particularly those involved

in intracellular trafficking pathways. Notably, detection of these factors was only possible by

correlating their expression specifically with the 3'/5' vRNA ratio and not the total vRNA reads.

The involvement of these factors specifically in cells harboring high 3'/5' vRNA read ratio thus

makes it experimentally challenging to further study them via bulk sample approaches.

Nevertheless, it is tempting to speculate that some of the discovered late endosomal trafficking

and lysosomal proteins (RAB7A (43), BROX (44), VAMP7 (45) and LAMP2 (48)) may be

involved in forming the CPV-I composed of modified endosomes and lysosomes in which

VEEV RNA replication occurs (13–15,56–59), and that ATP6V1B2 (50) may mediate the

acidification of this acidic intracellular compartment (41). Moreover, the positive correlation of

proteins involved in ER to Golgi trafficking (SEC22B) (46), regulation of secretion (PIP4K2A)

(47), autophagy (LAMP2) (48), actin polymerization (PFN2) (49), and ESCRT machinery

(BROX, a Bro1 domain-containing protein like ALIX) (44,60), TSG101 and STAM2) with the

3'/5' vRNA read ratio proposes roles for these factors in late stages of the VEEV lifecycle, such

as trafficking of the CPV-IIs to the plasma membrane, virion assembly, and/or budding (16–18).

These results propose a model wherein specific genes are upregulated within the profound

transcriptional downregulation in a stoichiometry-dependent manner, and further illuminate the

utility of viscRNA-Seq in identifying candidate proviral and antiviral factors, including

druggable candidates for host-targeted antiviral approaches.

113
114

Comparative evolutionary virology is an ideal application for single cell technologies because of

the degree of genomic and functional diversity of infections. As a proof of concept, we compared

the effect of unrelated human RNA viruses on the host cell in permissive cell lines. To address

the confounding effect of different cell line backgrounds, we restricted the analyses in **Figs 6A-F**

to genes with a similar baseline expression level across cell lines. We compared genes that

positively correlated with the 3'/5' VEEV RNA ratio with those correlating with DENV or ZIKV

vRNA and found concordant signal for genes involved in protein processing and transport,

whereas some cell cycle and apoptosis genes appeared to be specific to VEEV. When comparing

data on five different viruses derived using different cell lines and technologies, we observed that

while the closely related flaviviruses DENV and ZIKV affect a highly overlapping set of genes

in both up and downregulation, more distant evolutionary relationships between the viruses lead

to essentially distinct lists of dysregulated host genes. Moreover, the "correct" viral phylogeny

grouping all three flaviviruses as a monophyletic group could be recovered purely from the host

transcriptome perturbations, i.e. without using viral genomic information, which is intriguing.

More viruses across the viral phylogeny should be assessed to evaluate whether this signal is the

result of conserved ancestral function or, alternatively, of convergent functional evolution.

Overall, our study uncovered global and gene-specific host transcriptional dynamics during

VEEV (TC-83) infection at single cell resolution and presented a novel approach to elucidate the

evolution of virus-host interactions.

117
118

Acknowledgements

This work was supported by HDTRA11810039 from the Defense Threat Reduction Agency

(DTRA)/Fundamental Research to Counter Weapons of Mass Destruction, by the Chan

Zuckerberg Biohub, and by a Stanford Bio-X Interdisciplinary Initiative Program Award. ZY

was supported by the Maternal and Child Health Research Institute, Lucile Packard Foundation

for Children's Health). NP was supported by the University of Washington School of Medicine

Guy Tribble and Susan Barnes Graduate Discovery Fellowship. We thank investigators who

have provided plasmids (see Methods).The opinions, interpretations, conclusions, and

recommendations are those of the authors and are not necessarily endorsed by the U.S. Army or

the other funders.

Competing interests: The authors declare that no competing interests exist.

References:

1. Sharma A, Knollmann-Ritschel B. Current Understanding of the Molecular Basis of
Venezuelan Equine Encephalitis Virus Pathogenesis and Vaccine Development [Internet].
Vol. 11, Viruses. 2019. p. 164. Available from: http://dx.doi.org/10.3390/v11020164

2. Aguilar PV, Estrada-Franco JG, Navarro-Lopez R, Ferro C, Haddow AD, Weaver SC.
Endemic Venezuelan equine encephalitis in the Americas: hidden under the dengue
umbrella. Future Virol. 2011;6(6):721–40.

3. Hawley RJ, Eitzen EM Jr. Biological Weapons—a Primer for Microbiologists [Internet].
Vol. 55, Annual Review of Microbiology. 2001. p. 235–53. Available from:
http://dx.doi.org/10.1146/annurev.micro.55.1.235

4. Berge TO, Banks IS, Tigertt WD, Others. Attenua-lion of Venezuelan Equine
Encephalomyelitis Virus by ire vitro Cultivation in Guinea-Pig Heart Cells. Am J Hyg.
1961;73(2):209–18.

5. Keck F, Kortchak S, Bakovic A, Roberts B, Agrawal N, Narayanan A. Direct and indirect
pro-inflammatory cytokine response resulting from TC-83 infection of glial cells [Internet].
Vol. 9, Virulence. 2018. p. 1403–21. Available from:
http://dx.doi.org/10.1080/21505594.2018.1509668

6. Garmashova N, Atasheva S, Kang W, Weaver SC, Frolova E, Frolov I. Analysis of
Venezuelan equine encephalitis virus capsid protein function in the inhibition of cellular
transcription. J Virol. 2007 Dec;81(24):13552–65.

7. Strauss JH, Strauss EG. The alphaviruses: gene expression, replication, and evolution.
Microbiol Rev. 1994 Sep;58(3):491–562.

8. Shirako Y, Strauss JH. Cleavage between nsP1 and nsP2 initiates the processing pathway of
Sindbis virus nonstructural polyprotein P123. Virology. 1990 Jul;177(1):54–64.

9. Lemm JA, Rümenapf T, Strauss EG, Strauss JH, Rice CM. Polypeptide requirements for
assembly of functional Sindbis virus replication complexes: a model for the temporal
regulation of minus- and plus-strand RNA synthesis [Internet]. Vol. 13, The EMBO Journal.
1994. p. 2925–34. Available from: http://dx.doi.org/10.1002/j.1460-2075.1994.tb06587.x

10. Raju R, Huang HV. Analysis of Sindbis virus promoter recognition in vivo, using novel
vectors with two subgenomic mRNA promoters. J Virol. 1991 May;65(5):2501–10.

11. Levis R, Schlesinger S, Huang HV. Promoter for Sindbis virus RNA-dependent subgenomic
RNA transcription. J Virol. 1990 Apr;64(4):1726–33.

12. Kielian M, Chanel-Vos C, Liao M. Alphavirus Entry and Membrane Fusion. Viruses. 2010
Mar 26;2(4):796–825.

13. Spuul P, Balistreri G, Kaariainen L, Ahola T. Phosphatidylinositol 3-Kinase-, Actin-, and
Microtubule-Dependent Transport of Semliki Forest Virus Replication Complexes from the
Plasma Membrane to Modified Lysosomes [Internet]. Vol. 84, Journal of Virology. 2010. p.
7543–57. Available from: http://dx.doi.org/10.1128/jvi.00477-10

14. Grimley PM, Berezesky IK, Friedman RM. Cytoplasmic structures associated with an
arbovirus infection: loci of viral ribonucleic acid synthesis. J Virol. 1968 Nov;2(11):1326–
38.

15. Kujala P, Ikäheimonen A, Ehsani N, Vihinen H, Auvinen P, Kääriäinen L. Biogenesis of the
Semliki Forest virus RNA replication complex. J Virol. 2001 Apr;75(8):3873–84.

16. Garoff H, Wilschut J, Liljeström P, Wahlberg JM, Bron R, Suomalainen M, et al. Assembly
and entry mechanisms of Semliki Forest virus. Arch Virol Suppl. 1994;9:329–38.

17. Griffiths G, Quinn P, Warren G. Dissection of the Golgi complex. I. Monensin inhibits the
transport of viral membrane proteins from medial to trans Golgi cisternae in baby hamster
kidney cells infected with Semliki Forest virus. J Cell Biol. 1983 Mar;96(3):835–50.

18. Soonsawad P, Xing L, Milla E, Espinoza JM, Kawano M, Marko M, et al. Structural
evidence of glycoprotein assembly in cellular membrane compartments prior to Alphavirus
budding. J Virol. 2010 Nov;84(21):11145–51.

19. Radoshitzky SR, Pegoraro G, Chī XO, D Ng L, Chiang C-Y, Jozwick L, et al. siRNA
Screen Identifies Trafficking Host Factors that Modulate Alphavirus Infection. PLoS
Pathog. 2016 Mar;12(3):e1005466.

20. Amaya M, Brooks-Faulconer T, Lark T, Keck F, Bailey C, Raman V, et al. Venezuelan
equine encephalitis virus non-structural protein 3 (nsP3) interacts with RNA helicases
DDX1 and DDX3 in infected cells. Antiviral Res. 2016 Jul;131:49–60.

21. Gupta P, Sharma A, Han J, Yang A, Bhomia M, Knollmann-Ritschel B, et al. Differential
host gene responses from infection with neurovirulent and partially-neurovirulent strains of
Venezuelan equine encephalitis virus. BMC Infect Dis. 2017 Apr 26;17(1):309.

22. Atasheva S, Akhrymuk M, Frolova EI, Frolov I. New PARP gene with an anti-alphavirus
function. J Virol. 2012 Aug;86(15):8147–60.

23. Zanini F, Pu S-Y, Bekerman E, Einav S, Quake SR. Single-cell transcriptional dynamics of
flavivirus infection. Elife [Internet]. 2018 Feb 16;7. Available from:
http://dx.doi.org/10.7554/eLife.32942

24. Russell AB, Trapnell C, Bloom JD. Extreme heterogeneity of influenza virus infection in
single cells. Elife [Internet]. 2018 Feb 16;7. Available from:
http://dx.doi.org/10.7554/eLife.32303

25. Russell AB, Elshina E, Kowalsky JR, Te Velthuis AJW, Bloom JD. Single-Cell Virus
Sequencing of Influenza Infections That Trigger Innate Immunity. J Virol [Internet]. 2019
Jul 15;93(14). Available from: http://dx.doi.org/10.1128/JVI.00500-19

26. O'Neal JT, Upadhyay AA, Wolabaugh A, Patel NB, Bosinger SE, Suthar MS. West Nile
Virus-Inclusive Single-Cell RNA Sequencing Reveals Heterogeneity in the Type I
Interferon Response within Single Cells. J Virol [Internet]. 2019 Mar 15;93(6). Available
from: http://dx.doi.org/10.1128/JVI.01778-18

27. Zanini F, Robinson ML, Croote D, Sahoo MK, Sanz AM, Ortiz-Lasso E, et al. Virus-

inclusive single-cell RNA sequencing reveals the molecular signature of progression to
severe dengue. Proc Natl Acad Sci U S A. 2018 Dec 26;115(52):E12363–9.

28. Sun C, Gardner CL, Watson AM, Ryman KD, Klimstra WB. Stable, high-level expression
of reporter proteins from improved alphavirus expression vectors to track replication and
dissemination during encephalitic and arthritogenic disease. J Virol. 2014;88(4):2035–46.

29. Rual J-F, Hirozane-Kishikawa T, Hao T, Bertin N, Li S, Dricot A, et al. Human ORFeome
version 1.1: a platform for reverse proteomics. Genome Res. 2004 Oct;14(10B):2128–35.

30. Coutant EP, Goyard S, Hervin V, Gagnot G, Baatallah R, Jacob Y, et al. Gram-scale
synthesis of luciferins derived from coelenterazine and original insights into their
bioluminescence properties. Org Biomol Chem. 2019 Apr 10;17(15):3709–13.

31. Coutant EP, Gagnot G, Hervin V, Baatallah R, Goyard S, Jacob Y, et al. Bioluminescence
Profiling of NanoKAZ/NanoLuc Luciferase Using a Chemical Library of Coelenterazine
Analogues [Internet]. Chemistry – A European Journal. 2019. Available from:
http://dx.doi.org/10.1002/chem.201904844

32. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

33. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput
sequencing data. Bioinformatics [Internet]. 2015; Available from: https://academic.oup.com/
bioinformatics/article-abstract/31/2/166/2366196

34. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. Nat Biotechnol. 2018
Jun;36(5):411–20.

35. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer; 2016. 260 p.

36. Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007 May 1;9(3):90–
5.

37. Waskom ML, Kumaran D, Gordon AM, Rissman J, Wagner AD. Frontoparietal
representations of task context support the flexible control of goal-directed cognition. J
Neurosci. 2014 Aug 6;34(32):10743–55.

38. Hoyer S, Hamman J. xarray: ND labeled arrays and datasets in Python. Journal of Open
Research Software [Internet]. 2017;5(1). Available from:
https://openresearchsoftware.metajnl.com/articles/148/

39. PontÉN JAN, Macintyre EH. Long term culture of normal and neoplastic human glia. Acta
Pathol Microbiol Scand. 1968;74(4):465–86.

40. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape
provides a biologist-oriented resource for the analysis of systems-level datasets. Nat
Commun. 2019 Apr 3;10(1):1523.

41. Jose J, Taylor AB, Kuhn RJ. Spatial and Temporal Analysis of Alphavirus Replication and

Assembly in Mammalian and Mosquito Cells. MBio [Internet]. 2017 Feb 14;8(1). Available
from: http://dx.doi.org/10.1128/mBio.02294-16

42. Fu M, Blackshear PJ. RNA-binding proteins in immune regulation: a focus on CCCH zinc
finger proteins. Nat Rev Immunol. 2017 Feb;17(2):130–43.

43. Verhoeven K, De Jonghe P, Coen K, Verpoorten N, Auer-Grumbach M, Kwon JM, et al.
Mutations in the small GTP-ase late endosomal protein RAB7 cause Charcot-Marie-Tooth
type 2B neuropathy. Am J Hum Genet. 2003 Mar;72(3):722–7.

44. Mu R, Dussupt V, Jiang J, Sette P, Rudd V, Chuenchor W, et al. Two distinct binding
modes define the interaction of Brox with the C-terminal tails of CHMP5 and CHMP4B.
Structure. 2012 May 9;20(5):887–98.

45. Pryor PR, Jackson L, Gray SR, Edeling MA, Thompson A, Sanderson CM, et al. Molecular
basis for the sorting of the SNARE VAMP7 into endocytic clathrin-coated vesicles by the
ArfGAP Hrb. Cell. 2008 Sep 5;134(5):817–27.

46. Zhang T, Wong SH, Tang BL, Xu Y, Hong W. Morphological and functional association of
Sec22b/ERS-24 with the pre-Golgi intermediate compartment. Mol Biol Cell. 1999
Feb;10(2):435–53.

47. Rozenvayn N, Flaumenhaft R. Protein kinase C Mediates Translocation of Type II
Phosphatidylinositol 5-Phosphate 4-Kinase Required for Platelet α-Granule Secretion. J Biol
Chem. 2003 Mar 7;278(10):8126–34.

48. Hubert V, Peschel A, Langer B, Gröger M, Rees A, Kain R. LAMP-2 is required for
incorporating syntaxin-17 into autophagosomes and for their fusion with lysosomes. Biol
Open. 2016 Oct 15;5(10):1516–29.

49. Honoré B, Madsen P, Andersen AH, Leffers H. Cloning and expression of a novel human
profilin variant, profilin II. FEBS Lett. 1993 Sep 13;330(2):151–5.

50. Bernasconi P, Rausch T, Struve I, Morgan L, Taiz L. An mRNA from human brain encodes
an isoform of the B subunit of the vacuolar H (+)-ATPase. J Biol Chem.
1990;265(29):17428–31.

51. Qi S, O'Hayre M, Gutkind JS, Hurley JH. Structural and biochemical basis for ubiquitin
ligase recruitment by arrestin-related domain-containing protein-3 (ARRDC3). J Biol Chem.
2014 Feb 21;289(8):4743–52.

52. Maaten L van der, Hinton G. Visualizing Data using t-SNE. J Mach Learn Res.
2008;9(Nov):2579–605.

53. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection
for Dimension Reduction [Internet]. arXiv [stat.ML]. 2018. Available from: http://arxiv.org/
abs/1802.03426

54. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected
communities [Internet]. Vol. 9, Scientific Reports. 2019. Available from:
http://dx.doi.org/10.1038/s41598-019-41695-z

55. Gorman MJ, Caine EA, Zaitsev K, Begley MC, Weger-Lucarelli J, Uccellini MB, et al. An
Immunocompetent Mouse Model of Zika Virus Infection. Cell Host Microbe. 2018 May
9;23(5):672–85.e6.

56. Salonen A, Vasiljeva L, Merits A, Magden J, Jokitalo E, Kääriäinen L. Properly folded
nonstructural polyprotein directs the semliki forest virus replication complex to the
endosomal compartment. J Virol. 2003 Feb;77(3):1691–702.

57. Friedman RM, Levin JG, Grimley PM, Berezesky IK. Membrane-associated replication
complex in arbovirus infection. J Virol. 1972 Sep;10(3):504–15.

58. Pietilä MK, van Hemert MJ, Ahola T. Purification of Highly Active Alphavirus Replication
Complexes Demonstrates Altered Fractionation of Multiple Cellular Membranes. J Virol
[Internet]. 2018 Apr 15;92(8). Available from: http://dx.doi.org/10.1128/JVI.01852-17

59. Pietilä MK, Hellström K, Ahola T. Alphavirus polymerase and RNA replication [Internet].
Vol. 234, Virus Research. 2017. p. 44–57. Available from:
http://dx.doi.org/10.1016/j.virusres.2017.01.007

60. Zhai Q, Landesman MB, Robinson H, Sundquist WI, Hill CP. Structure of the Bro1 domain
protein BROX and functional analyses of the ALIX Bro1 domain in HIV-1 budding. PLoS
One. 2011 Dec 5;6(12):e27466.

946 **Figures and legends**

947
948

949
Fig 1. Cell-to-cell heterogeneity during VEEV infection. (A) Schematic of the experimental
setup. (B) A scatter plot showing VEEV cDNA sequencing reads and GFP expression measured
via FACS (FITC gate) in cells harboring 1 or more viral reads. The dotted line represents the
cutoff of infected cells. Cells harboring more than 10 viral reads are considered infected. (C) The
fraction of VEEV-TC-83-infected U-87 MG cells over time for two MOIs. (D) Box plots
depicting the ratio of virus to total cDNA reads (left) and GFP expression level (right) over time.
The horizontal dotted line represents the threshold dividing cells into "low vRNA" and "high
vRNA" harboring cells (see text). (E) Box plots showing host cDNA to ERCC read ratio in
infected and mock-infected cells derived from different time points postinfection. *p < 0.05 by
Mann-Whitney U test. HPI, hours postinfection; MOI, multiplicity of infection; ERCC, External
RNA Controls Consortium.

961
962

Fig 2. Host genes and pathways are altered during VEEV infection. (A) A Venn diagram
showing the number of unique and overlapping genes that emerged from the differentially gene
expression analysis and host RNV/vRNA correlation analysis. Molecular function terms and P
values derived from Gene Ontology (GO) enrichment analysis of 263 genes that are both
differentially expressed between high vRNA and mock-infected cells and correlated with vRNA.
(B) Ridge plots of representative host genes that are differentially expressed between high vRNA
and mock-infected cells and a gene (COPZ2) whose level is unaltered. 50 cells from each group
were selected for plotting. Dash lines indicate median expression level of the corresponding
genes. Gene expression was normalized using the following the formula: ln ((gene counts /
ERCC counts) + 1). (C) Representative scatter plots of host gene expression versus vRNA
abundance and corresponding Rho Spearman's correlation coefficients. Each dot is a single cell
colored by the time postinfection, and the shaded contours indicate cell density (greyscale,
darker is higher). HPI, hours postinfection; MOI, multiplicity of infection; ERCC, External RNA
Controls Consortium; TNFAIP3, Tumor Necrosis Factor Alpha-Induced Protein 3; TAF7,
TATA-Box Binding Protein Associated Factor 7; COPZ2, COPI Coat Complex Subunit Zeta 2.

149
150

A

B

C

151
152

153
154

Fig 3. "Superproducers" cells exhibit altered gene expression patterns. (A) Scatter plot depicting GFP expression level and virus/total reads in cells at 6 hours postinfection with VEEV (TC-83) at an MOI of 1. The horizontal and vertical dash lines indicate the cutoffs of GFP signal and virus/total read ratio, respectively (see text). Each dot represents a cell. Orange, cells with a GFP signal readout that is greater than 1000 and virus/total read ratio greater than 0.001 defined as "superproducers" (n = 11); blue: cells not meeting these criteria. (B and C) Representative violin plots showing genes that are upregulated (B) or downregulated (C) specifically in "superproducers" cells relative to either mock-infected cells, low vRNA cells harvested at 6 hpi or high vRNA cells harvested at 24 hpi. HPI, hours postinfection; MOI, multiplicity of infection. SYTL3, Synaptotagmin Like 3; KDM3B, Lysine Demethylase 3B; SNX29, Sorting Nexin 29; COG5, Component Of Oligomeric Golgi Complex 5; ZMAT5, Zinc Finger Matrin-Type 5; VPS37A, Vacuolar Protein Sorting-Associated Protein 37A; AC087343.1, Ribosomal Protein L21 (RPl21) Pseudogene.

1002

Fig 4. The expression of genes involved in intracellular membrane trafficking correlates with the ratio of 3' to 5' vRNA reads. (A) Coverage of viral reads over the entire VEEV genome. Each line is a cell, and the red line is a scaled average across all cells. (B) Genome architecture of VEEV highlighting the nonstructural (yellow) and structural (green) protein domains. (C) Scatter plot showing positive correlation of VEEV 3'/5' read ratio with cellular vRNA abundance. Each dot is an infected cell. (D-E) Scatter plots showing no correlation between the 3'a/3'b read ratio (D) and 5'a/5'b read ratio (E) and cellular vRNA abundance. (F) Histogram of Spearman's correlation coefficients between all host genes and the 3'/5' read ratio. (G) Representative scatter plots of host gene expression versus vRNA 3'/5' read ratio and corresponding Rho Spearman's correlation coefficients. Each dot is a cell and contour plots indicate cell density (low to high, green to red). (H) Gene enrichment analysis of top 300 genes positively correlated with the 3'/5' read ratio. ORF, opening reading frame; PFN2, Profilin 2; BROX, BRO1 Domain- And CAAX Motif-Containing Protein; ATP6V1B2, ATPase H+ Transporting V1 Subunit B2; BNIP3, BCL2 Interacting Protein 3; LAMP2, Lysosomal Associated Membrane Protein 2; PIP4K2A, Phosphatidylinositol-5-Phosphate 4-Kinase Type 2 Alpha; VAMP7, Vesicle Associated Membrane Protein 7; RAB7A, Ras-Related Protein Rab-7a; SEC22B, SEC22 Homolog B.

A

B

C

169
170

Fig 5. Validation of candidate VEEV proviral and antiviral genes. VEEV infection relative
to non-targeting (NT) siRNA (A) or empty plasmid (B) controls following siRNA-mediated
knockdown (A) or overexpression (B) of the indicated host factors measured by luminescence
assays at 18 hpi (MOI = 0.01) of U-87 MG cells with VEEV-TC-83-NLuc and normalized to cell
viability. Columns are color-coded based on the correlation of the respective gene with vRNA
abundance via viscRNA-Seq: yellow for genes that are positively correlated with vRNA and
grey for genes that are negatively correlated with vRNA. Both data sets are pooled from two
independent experiments with six replicates each. Shown are means ± SD; *p < 0.05, **p < 0.01,
***p < 0.001 relative to the respective control by 1-way ANOVA followed by Dunnett's post
hoc test. The dotted lines represent the cutoffs for positivity. Cellular viability measurements are
shown in S3 Fig. (C) Correlation coefficients between proviral candidates with the 3' (grey) and
5' (orange) vRNA reads.

1039

1040

Fig 6. Comparative viscRNA-Seq analysis across five RNA viruses. (A-C) Scatter plots of
representative host gene expression versus vRNA in single cells during DENV (orange), ZIKV
(blue), and VEEV (green) infection. Dots indicate single cells, lines are running averages at
increasing vRNA abundances. (D, E) Correlation between host gene expression and vRNA
abundance during DENV versus ZIKV infection of the top genes that positively correlate with
the VEEV 3'/5' read ratio (D) or a similar number of random genes (E). Each dot is a gene and
the axis coordinate is the percentage of genes with a correlation with vRNA smaller than the the
gene of interest. For (D), size of the dot increases with the correlation with VEEV 3'/5' read
ratio (top correlated gene is largest). (F) UMAP (53)embedding of host genes correlation with
vRNA during infection by 5 individual RNA viruses. Blue and red indicate downregulation and
upregulation during each infection, respectively. Several clusters of genes are observed (0-7).
(G) Hierarchical clustering of host gene clusters highlights that gene upregulation is mostly
virus-specific and is consistent with the known phylogeny. cpm, count per million; WNV, West
Nile virus; IAV, influenza A virus.

181
182

Supporting Information

S1 Text. Rare structural viral read variants correlate with expression of specific host genes.

S1 Fig. Quality control and definition of infected cells.

S2 Fig. Subgrouping cells based on viral load, representative differentially expressed genes

(DGEs) and correlation analysis.

S3 Fig. VEEV gap reads identified via viscRNA-Seq.

S4 Fig. Validation of proviral and antiviral factors.

S5 Fig. Pathway analysis for genes that positively correlated with VEEV 3'/5' read ratio and

positively (A) or negatively (B) correlated with DENV and ZIKV.

S1 Table. DENV capture oligonucleotides.

1103