µ-Lat: A Mouse Model to Evaluate Human Immunodeficiency Virus Eradication Strategies

Hannah S. Sperber[1,2,3*], Padma Priya Togarrati[1*], Kyle A. Raymond[1,3], Mohamed S. Bouzidi[1,3], Renata Gilfanova[1], Alan G. Gutierrez[1], Marcus O. Muench[1,3], and Satish K. Pillai[1,3+]

[1]Vitalant Research Institute, San Francisco, California, United States of America

[2]Free University of Berlin, Institute of Biochemistry, Berlin, Germany

[3]University of California, San Francisco, California, United States of America

*These authors contributed equally

+Corresponding author, e-mail: satish.pillai@ucsf.edu

14   <u>**Nonstandard abbreviations**</u>

ABBREVIATION	DEFINITION
ART	antiretroviral therapy
BM	bone marrow
GFP	green fluorescence reporter
GVHD	graft versus host disease
HIV	human Immunodeficiency virus
HSC	hematopoietic stem cell
IEL	intraepithelial lymphocytes
LN	lymph node
LRA	latency reversal agent
LTR	long terminal repeats
MFI	mean fluorescence intensity
NHP	nonhuman primates
NSG	NOD.Cg-$Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/SzJ
NSG-3GS	NOD.Cg-$Prkdc^{scid}$ $Il2rg^{tm1Wj}$l Tg(CMV-IL3,CSF2,KITLG)1Eav/MloySzJ
PBMC	peripheral blood mononuclear cell
PMA	phorbol-myristate-acetate
r.-o.	retro-orbital
SCID	Severe combined immunodeficiency
SIV	simian immunodeficiency virus
TNF	tumor necrosis factor

Abstract

A critical barrier to the development of a human immunodeficiency virus (HIV) cure is the lack of a scalable animal model that enables robust evaluation of eradication approaches prior to testing in humans. We established a humanized mouse model of latent HIV infection by transplanting "J-Lat" cells, Jurkat cells harboring a latent HIV provirus encoding an enhanced green fluorescent protein (GFP) reporter, into irradiated adult NOD.Cg-*Prkdc*^{scid} *Il2rg*^{tm1Wjl}/SzJ (NSG) mice. J-Lat cells exhibited successful engraftment in several tissues including spleen, bone barrow, peripheral blood, and lung, in line with the diverse natural tissue tropism of HIV. Administration of tumor necrosis factor (TNF)-α, an established HIV latency reversal agent, significantly induced GFP expression in engrafted cells across tissues, reflecting viral reactivation. These data suggest that our **mu**rine **lat**ency ("μ-Lat") model enables efficient determination of how effectively viral eradication agents, including latency reversal agents, penetrate and function in diverse anatomical sites harboring HIV *in vivo.*

Keywords: Humanized mouse, HIV, HIV latency, latency reversal, shock and kill, antiviral gene therapy

Introduction

The advent of antiretroviral therapy (ART) has dramatically reduced morbidity and mortality for human immunodeficiency virus (HIV)-infected individuals with access to healthcare in resource-rich countries. However, despite years of potent therapy, eradication of infection is not achieved due to the persistence of HIV latently-infected cells during treatment(1). Accumulating evidence suggest that "non-AIDS" cardiovascular, renal and hepatic diseases are amplified by HIV infection, and the immune system may exhibit premature senescence even among patients with complete viral suppression(2). Moreover, although enormous progress has been made to provide ART in resource-limited settings, there are huge economic, political and operational challenges to reach the goal of universal access to lifelong treatment. These realities have created a pronounced interest in developing HIV cure strategies.

A number of approaches to achieve a sterilizing or functional cure for HIV infection are currently under investigation, including therapeutic vaccination, immunomodulatory approaches, therapeutic HIV latency reversal (the "shock and kill" strategy), as well as a number of gene therapy approaches(3,4). In all scenarios, it will be critical to have proper diagnostic tools and models in place to comprehensively evaluate performance and safety prior to deployment in a clinical setting. A critical barrier to the development of an HIV cure is the lack of an accessible and scalable preclinical animal model that enables robust evaluation of candidate eradication approaches prior to testing in humans(5). As a result, many promising curative approaches never graduate past the petri dish stage. Infection of nonhuman primates (NHP) with simian immunodeficiency virus (SIV) is an option and has been utilized extensively to study HIV/AIDS pathogenesis(6,7). Recent advancements have been made in optimizing ART

regimens to achieve durable virus suppression and thus enable evaluation of HIV cure

strategies in the SIV-NHP model(8–10). However, biological limitations remain since this model

uses SIV and might not recapitulate human host-HIV interactions and HIV latency

mechanisms(11–15). In addition, NHP experiments involve complex ethical considerations, and

the high costs and labor requirements only allow small numbers of animals to be utilized in any

given trial, limiting statistical power and generalizability(11).

Mouse models represent another alternative with lower costs, more convenient husbandry

requirements, as well as greater scalability. In the context of HIV studies, a wide range of small

animal models have been developed comprising knockout mouse (16–18), transgenic

mouse(19–23), and humanized mouse models(6,24–29). Humanized mice are established by

xenotransplantation of human cells or tissues in immunodeficient mouse strains. Most strains

used in HIV research are derivatives of severe combined immunodeficient (SCID) mice, which

harbor mutations in the gene coding for a DNA-dependent protein kinase catalytic subunit

(Prkdc). These mice are "humanized" using two approaches: 1) human cells are injected with or

without prior irradiation of mice or 2) portions of tissue are surgically implanted. Different cells

have been used for injection, including human peripheral blood mononuclear cells (PBMCs), as

in the *hu-PBL-SCID* mouse model(30), obtained from healthy(31) or HIV-infected ART

suppressed individuals(32), or human hematopoietic stem cells (HSCs), as in the *hu-HSC* mouse

model(33), or the more recently developed *T-cell-only*(34) and *myeloid-only*(35) mouse models

(*ToM* and *MoM*). Implantation of fetal thymus and liver tissue fragments are used for the *SCID-*

hu thy/liv(36,37) and *bone marrow/liver/thymus* (BLT)(38,39) mouse models. Although all these

model systems have contributed to our understanding of HIV pathogenesis and persistence, key

limitations remain that need to be addressed in order to fully exploit the potential of these

small animal models in HIV cure research. *Hu-PBL-SCID* mice struggle with the development of

graft versus host disease (GVHD) which renders this model inapplicable for long-term studies

involving HIV persistence. The generation of *SCID-hu thy/liv* and BLT mice is limited due to the

need for surgical implantation and a limited supply of tissue. Moreover, these models as well as

hu-HSC, *ToM* and *MoM* mice rely on the engraftment of cells or tissues typically derived from

human fetal specimens, which in light of recent changes in U.S. federal policies face significant

challenges, as ethical, legal, and political considerations surrounding the use of fetal tissue in

scientific research have made it increasingly difficult to obtain such material(40). A major

limitation shared by all these models remains the low frequency of HIV latently-infected cells,

which impacts the applicability of these models as robust *in vivo* test bases for HIV cure

strategies.

In the present study, we therefore pursued a new approach and transplanted J-Lat 11.1 cells (J-

Lat cells) into irradiated adult NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wjl}*/SzJ (NSG) mice. J-Lat cell clones have

been derived from Jurkat cells, an immortalized human T lymphocyte cell line. Each cell harbors

a single latent HIV provirus encoding an enhanced green fluorescent protein (GFP) reporter in

place of the *nef* gene(41). The proviruses within the J-Lat clones are typically latent due to

epigenetic repression(42) and were selected to be responsive to TNF-α stimulation, resulting in

viral LTR-driven GFP expression. J-Lat 11.1 cells were chosen for this model as they displayed

the greatest viral reactivation upon TNF-α treatment in cell culture settings with respect to

other clones(43). Our data show robust engraftment of J-Lat cells in several tissues three weeks

after injection as well as significant reactivation of these latently-infected cells *in vivo* upon

intravenous administration of the established latency reversal agent (LRA) tumor necrosis

factor (TNF)-α. By applying an established and widely-utilized HIV latency reporter cell

line(41,43,44), we circumvent the need for fetal or any other donor-derived tissue, and achieve

high frequencies of HIV latently-infected cells *in vivo* on a short experimental time scale. In

addition, the presence of the GFP reporter cassette in the integrated viral genome provides for

quick and convenient assessment of viral reactivation using flow cytometry or microscopy.

Moreover, by taking advantage of an HIV reporter cell line, our **mu**rine **lat**ency (μ-Lat) model

represents a highly reproducible platform that allows for a simple transition from *in vitro*

screening approaches to an *in vivo* system enabling the evaluation of bioavailability and

pharmacokinetics of candidate regimens. Although not intended to serve as a pathophysiology

model, we present this approach as a scalable, accessible, and cost-effective preclinical testbed

to evaluate the safety, tolerability, and performance of HIV cure strategies in distinct

anatomical niches.

Materials and Methods

Cell culture and treatment: J-Lat 11.1 cells (kindly provided by Dr. Eric Verdin) contain an integrated latent full-length HIV genome harboring a mutation in the *env* gene and GFP in place of the *nef* gene as a reporter for transcriptional activity of the provirus[45]. J-Lat 11.1 cells were grown in media composed of RPMI 1640 (Gibco) supplemented with 10% fetal bovine serum (FBS) (Corning) and 1% Penicillin-Streptomycin (Gibco). Cells were cultured at 37°C in a humidified incubator containing 5% CO_2. To test the potency of latency reversal agents (LRAs), 1 x 10^6 cells in 1 ml RPMI were left untreated (negative control), incubated in 0.5% DMSO (vehicle control), 20 nM PMA/1 µM Ionomycin (positive control) or 20 ng/µl TNF-α for 24h.

Mice: The work was approved by the Institutional Animal Care and Use Committee guidelines at Covance Laboratories, Inc. (San Carlos, CA) under Animal Welfare Assurance A3367-01 and protocol number IAC 2185 / ANS 2469. Animal husbandry was carried out according to the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Mice were sacrificed in accordance with the guidelines from the American Veterinary Medical Association.

Adult female and male mice (≥8 weeks old) were included in this study and were maintained at Vitalant Research Institute (VRI). Breeding pairs of NSG mice were obtained from Jackson Laboratories (Bar Harbor, ME), and were bred and maintained at VRI in a vivarium free from >40 murine pathogens as determined through biannual nucleic acid testing (Mouse Surveillance Plus PRIA; Charles River) of sentinel mice exposed to mixed bedding. Mice were maintained in sterile, disposable microisolator cages (Innovive, Inc.), which were changed every 14 days.

Environmental enrichment was provided by autoclaved cotton Nestlets (Ancare Corp.) and GLP-

certified Bio-Huts (Bio-Serv). Feed consisted of sterile, irradiated diet of Teklad Global 19%

protein diet (Envigo) with free access to sterile-filtered, acidified water (Innovive, Inc.). Several

days prior to radiation and 3 weeks following radiation, mice were fed with irradiated Global

2018 rodent diet with 4100 ppm Uniprim® (Envigo).

J-Lat cell surface marker staining: For each staining, 1×10^6 J-Lat cells were washed once with

PBS (Gibco), resuspended in 100 µl PBS, and stained with Zombie dye (Cat. #423105,

BioLegend) according to manufacturer's protocol (1:100 dilution) to enable subsequent

discrimination between live and dead cells. 10 min after incubation with the Zombie dye,

human CD45-PE (clone HI30, Cat. #304039, BioLegend), CD4-PE (clone OKT4, Cat. #317410,

BioLegend), TCR α/β-PE (clone IP26, Cat. #306708, BioLegend), CD27-PE (clone M-T271, Cat.

#356406, BioLegend), CD147-APC (clone HIM6, Cat. #306214, BioLegend), CD29-APC (clone

TS2/16, Cat. #303008, BioLegend), and HLA-ABC-APC/Cy7 (clone W6/32, Cat. #311426,

BioLegend) antibodies were added respectively, and were incubated for another 20-30 min at

room temperature (RT) in the dark. Cells were then washed with 2 ml of cell staining buffer

(Cat. #420201, BioLegend), resuspended in 300 µl PBS and measured using an LSR II flow

cytometer (BD Biosciences). Respective Isotype stained samples were used as control. At least

10,000 events were recorded for each sample.

J-Lat cell transplantation into mice: If not otherwise stated, J-Lat cell transplantation was

performed as follows: mice were irradiated with 175 cGy radiation dose using a RS2000 X-Ray

Biological Irradiator (RAD Source Technologies, Inc.) 3 hours prior to cell transplantation. Mice

were transplanted with 10×10^6 J-Lat cells in a volume of 200 µl by tail vein injection. J-Lat cell

transplanted mice as well as control mice (left untransplanted) were sacrificed and analyzed

within 25 days of cell transplantation.

Engraftment analysis of J-Lat cells in mice by flow cytometry: Transplanted mice were

sacrificed by cervical dislocation 3 weeks post injection (or as indicated in figure legends). In

one experiment, BM, brain, gut, intraepithelial lymphocytes (IEL) from gut tissue, heart, spleen,

lung, lymph nodes, and PB were harvested. In subsequent experiments, selected tissues were

harvested including BM, spleen, lung, and PB. Single cell suspensions from the BM, spleen, and

PB were prepared as described previously by Beyer et al.(46). Brain, gut, heart, lymph node,

and lung specimens were processed as follows: after harvest, all samples were stored in PBS on

ice. Brain and lymph node specimens were washed twice in PBS, brain specimens additionally

cut into small fragments (3-4 mm^2), followed by mashing the tissue fragments with a pestle and

passaging the cell suspensions through 70 µm cell strainers. Gut, heart, and lung specimens

were washed twice with PBS and then cut into 3-4 mm^2 pieces in a petri dish in 1 ml digestion

solution consisting of 1 mg/ml DE Collagenase (Cat. #011-1040, VitaCyte) and 100 U/ml DNase I

(Cat. #D5025-15KU, Sigma-Aldrich) final concentration in HBSS (Gibco). Fragments were

transferred to 50 ml falcon tubes, digestion solution was added to a final volume of up to 5 ml,

and samples were incubated at 37°C for 30 min (gut, heart) or 50 min (lung). IEL from gut tissue

were isolated as supernatant after digestion incubation. Afterwards, up to 5 ml of stop solution,

consisting of 0.5% BSA (Cat. #A2153, MilliPoreSigma) and 2 mM EDTA (Cat. #E0306, Teknova)

final concentration in PBS, was added to each sample to end the enzymatic digestion reaction.

Single cell suspensions were prepared by passage through a 70 µm cell strainer and washed

with PBS. For the analysis of J-Lat cell engraftment, single cells from harvested mouse tissues

179 were stained with human CD147-APC or human CD147-APC combined with human CD29-APC,

180 and additional Pacific Blue conjugate, mouse-specific lineage markers including CD45-Pacific

181 Blue (clone 30-F11, Cat. #103126, BioLegend), TER-119-Pacific Blue (clone TER-119, Cat.

182 #116232, BioLegend), and H-2K^D-Pacific Blue antibodies (clone SF1-1.1, Cat. #116616,

183 BioLegend). Cells were incubated for 30 min at RT in the dark. Zombie dye staining detected in

184 the APC-Cy7 channel was used for discrimination of live and dead cells. Following staining, cells

185 were washed and run on an LSR II flow cytometer, where at least 10,000 events were recorded

186 per sample. Flow cytometry data were analyzed using FlowJo software (FlowJo, Inc.).

187 **Treatment of J-Lat engrafted mice with latency reversal agent TNF-α:** Mice engrafted with J-

188 Lat cells were treated with recombinant human TNF-α (Cat. #PHC3011, Gibco), a potent and

189 well-known LRA. Briefly, NSG mice were transplanted with J-Lat cells and 3 weeks post-

190 transplantation, mice received TNF-α (diluted in 200 µl PBS) intravenously at a dose of

191 20 µg/mouse(47). NSG mice injected with J-Lat cells and treated with PBS were used as vehicle

192 control group.

193 After 24h of TNF-α treatment, mice were sacrificed to determine viral reactivation in PB, BM,

194 lung and spleen tissues. Latency reversal of the HIV provirus was analyzed by comparing GFP

195 expression of J-Lat cells in tissues of TNF-α treated mice vs the vehicle control group. GFP

196 expression of engrafted J-Lat cells was determined by flow cytometry as described above.

197 **Data analysis:** All data were analyzed using GraphPad Prism version 8.2 and are presented as

198 mean ± the standard error of the mean (SEM). Applied statistical tests and analyses are

199 described in figure legends. A P value of ≤0.05 was considered as statistically significant.

200 Individual P values are indicated in figures.

Results

The human cell surface proteins CD147 and CD29 enable discrimination of J-Lat cells from mouse cells. Our model involves transplantation of J-Lat cells (immortalized human T cells harboring a latent HIV provirus with a GFP reporter reflecting viral transcriptional activity) into irradiated adult NSG mice(41). We therefore searched for cell surface markers that identify engrafted J-Lat cells in a background of mouse cells, exhibiting three key features: 1) universal expression across J-Lat cells, 2) high per-cell expression on J-Lat cells, and 3) absence of expression on the surface of mouse cells.

Our candidate panel included seven cell surface proteins commonly known to be expressed in human CD4+ T cells: CD45, CD4, TCR α/β, CD27, CD147, CD29, and HLA-ABC. Four of these cell surface proteins were found to be expressed universally among the J-Lat population: CD45, CD147, CD29 and HLA-ABC (**Figure 1A**). The mean fluorescence intensity (MFI) of the four proteins was measured, reflecting the relative abundance of each protein on the cell surface. CD147 exhibited the highest MFI, followed by CD29 (**Figure 1B**). We next tested if the CD29 and CD147 antibodies (both APC conjugated to achieve maximum signal to noise ratio in the APC-channel) showed detectable binding to mouse cells obtained from different tissues. We harvested bone marrow (BM), lung, brain, gut, intraepithelial lymphocytes (IEL) from gut tissue, lymph nodes, heart and spleen from an untransplanted control mouse. In addition, based on Hoggatt et al.(48), who reported that blood parameters such as cell composition significantly varied depending on sampling manner and site, we performed peripheral blood (PB) harvest in three different ways: by retro-orbital bleed (r.-o.), tail vein bleed, or heart bleed. Human CD29

and CD147 staining was combined with antibodies targeting the mouse-specific lineage markers

CD45, TER-119, and H-2K^d for a positive identification of mouse cells. While 100% of cultured J-

Lat cells expressed CD29 and CD147 and were negative for mouse-specific markers (**Figure 1C**),

the frequency of CD147$^+$/CD29$^+$ cells was negligible or absent across mouse tissues (**Figure 1D**

and E). Most tissues harbored no CD147$^+$/CD29$^+$ cells, while 0.06% CD147$^+$/CD29$^+$ cells were

detected in the lung and 1.37% CD147$^+$/CD29$^+$ cells were detected in PB upon heart bleed

(**Figure 1E**). For both samples, lung and PB heart bleed, these percentages represent a single

event detected to be positive for CD147/CD29 (**Figure 1D and E**). Thus, we concluded that

among the tested cell surface proteins, human CD147 combined with CD29 function as a

specific marker pair for the identification of J-Lat cells engrafted in mouse tissues.

Establishment and optimization of J-Lat cell engraftment in humanized mice. In the context of

establishing a humanized mouse model, we examined and optimized several parameters

simultaneously to achieve a highly reproducible and robust *in vivo* platform (**Figure 2**).

In the following pilot experiments, we focused on J-Lat cell engraftment in the BM (**Figure 3**). To

measure the kinetics of J-Lat cell engraftment, we harvested the BM at different time points

post J-Lat cell injection (**Figure 3A**). We examined the effect of varying cell doses (**Figure 3B**)

and irradiation of recipient mice (**Figure 3C**) on J-Lat cell engraftment levels, and compared

engraftment levels in NOD.Cg-*Prkdc^{scid} Il2rg^{tm1Wj}I* Tg(CMV-IL3,CSF2,KITLG)1Eav/MloySzJ (NSG-

3GS) and NSG mice (**Figure 3D**). We detected J-Lat cells in the BM 18 days post cell injection

(pci), and detection levels reached a significant increase 25 days pci compared to 1 and 3 days

pci with an average of 16.3±8.6% engrafted J-Lat cells (**Figure 3A**). Using a J-Lat cell injection

dose of 10 x 10^6 J-Lat cells compared to 5 x 10^6 J-Lat cells per mouse led to a higher but not

significant increase in engraftment levels in the BM of nonirradiated NSG-3GS mice (**Figure 3B**).

Nonirradiated NSG-3GS recipient mice harbored lower frequencies of engrafted J-Lat cells in

the BM compared to irradiated NSG-3GS recipient mice (**Figure 3C**). Again, this difference was

not significant but showed a slight increase in J-Lat cell engraftment levels when recipient mice

were irradiated. Finally, when testing engraftment levels in different mouse strains, we found

significantly higher numbers of J-Lat cells in NSG mice (**Figure 3D**). J-Lat cell engraftment levels

in the BM 3 weeks pci of 10 x 10^6 J-Lat cells were 2.4 fold higher in NSG mice compared to NSG-

3GS mice (**Figure 3D**). Based on our results, we concluded that irradiation of recipient NSG mice

and a cell dose of 10 x 10^6 J-Lat cells per mouse administered through intravenous injection

resulted in efficient and reproducible cell engraftment levels which peaked 3 weeks following

cell transplantation.

J-Lat cells engraft successfully in several tissues in transplanted NSG mice. Using the

determined experimental conditions, we investigated J-Lat cell engraftment in several tissues,

as listed above, that we harvested from 5 irradiated NSG mice 3 weeks pci (**Figure 4**). To

analyze the expression of J-Lat and mouse markers on cells obtained from different mouse

tissues, we followed the gating strategy as described before (see **Figure 1C and Figure 4A and**

B). J-Lat cell engraftment levels were evaluated based on the frequency, MFI, and the absolute

number of CD147$^+$/CD29$^+$ cells in the respective processed tissues (**Figure 4C-E and**

Supplementary Figure 1). We found engrafted J-Lat cells in several tissues including the BM,

lung, spleen, and PB (**Figure 4A and B**). The mean frequency of engrafted J-Lat cells varied

across tissues from approximately 30% in the BM to 0.7% in the spleen (**Figure 4C**). While the

MFI of the engrafted J-Lat cells was comparable within and across tissues (**Figure 4D**), the

absolute number of engrafted J-Lat cells varied drastically. We detected the highest J-Lat cell

amount in the BM, followed by the amount in the lung and then the spleen (**Figure 4E**). We did

not detect engrafted J-Lat cells in the gut, IEL from the gut or heart tissues (**Figure 4C-E**).

Processing brain and lymph node tissues yielded only low overall cell numbers that were

evaluable (**Supplementary Figure 1A and B**). While J-Lat cells were identified in processed brain

tissues at high frequency (**Supplementary Figure 1C**), the actual number of engrafted J-Lat cells

was on average 5 cells (**Supplementary Figure 1E**). We thus decided to exclude the brain in

subsequent experiments due to the low overall cell numbers. We did not detect engrafted J-Lat

cells in the lymph node (**Supplementary Figure 1B-E**). When focusing on engraftment in the PB,

we found that heart and r.-o. bleed yielded a substantial higher frequency of engrafted J-Lat

cells compared to tail vein bleed (**Figure 4C**). However, PB harvest by r.-o. bleeding showed the

highest MFI, and more importantly, the highest absolute number of engrafted J-Lat cells (**Figure**

4D and E). Taken together, we found different engraftment levels based on the sample

collection method as observed by Hoggatt et al.(48). Thus, due to considerable engraftment

levels and practicality of collection, we decided to harvest PB in subsequent experiments via r.-

o. bleeding.

Based on these results and parameter thresholds applied in concert (frequency > 0.005; MFI >

15,000; absolute number > 10), we focused in subsequent experiments on J-Lat engraftment in

the BM, lung, spleen and PB via r.-o. bleed and established a suitable engraftment and sample

collection protocol to drive the model.

J-Lat cells maintain low GFP expression upon engraftment in selected tissues. Once we

confirmed reproducible and robust engraftment in several tissues, we wanted to investigate

GFP expression levels of engrafted J-Lat cells. As described before, the GFP reporter integrated

into the latent provirus in J-Lat cells is under the control of the proviral LTR and thus reflects

viral transcriptional activity. High GFP expression levels would thus indicate spontaneous viral

reactivation *in vivo* and render the model inapplicable for testing LRAs. We measured J-Lat cell

engraftment levels and GFP background signal of engrafted J-Lat cells in the selected tissues

(BM, lung, spleen, PB r.-o.) of 10 NSG mice 3 weeks pci (**Figure 5**). We applied our previously

described gating strategy to identify engrafted J-Lat cells based on CD147/CD29 expression and

measured GFP signal within this compartment (**Figure 5A-C**). Engraftment of J-Lat cells was

detected in the BM, lung, spleen, and PB r.-o. as observed before (**Figure 5D**). Across these

tissues, GFP expression levels of engrafted J-Lat cells remained low and frequencies did not

exceed an average of 4% GFP-positive cells (**Figure 5E**). The results showed that while these

mouse tissues harbored substantial levels of HIV latently-infected cells *in vivo,* engraftment of J-

Lat cells in the tested tissues did not lead to spontaneous reactivation of viral transcription.

TNF-α treatment induces GFP expression in tissue engrafted J-Lat cells reflecting reactivation

of latent provirus *in vivo*. Finally, we sought to determine if HIV LRA administration would

result in viral reactivation *in vivo*. Viral reactivation was measured based on the frequency of

GFP expressing cells within engrafted J-Lat cells following LRA treatment (**Figure 6A**).

The proviruses within the J-Lat family of clones were selected to be responsive to TNF-α

stimulation, resulting in viral LTR-driven GFP expression(41). Previous studies have shown, that

among the established clones, J-Lat 11.1 cells displayed greatest viral reactivation upon TNF-α

treatment in cell culture settings(43). 24h *in vitro* treatment of J-Lat cells with 20 ng/µl TNF-α

resulted in a significant increase of GFP-positive cells compared to mock-treated (0.5% DMSO)

and untreated cells, which were largely GFP-negative (**Figure 6B**). Therefore, we performed *in*

vivo reactivation experiments using 20 µg of TNF-α as an LRA (with PBS as a vehicle

control)(47). 24h following TNF-α tail vein injection, effects on tissue engraftment and

reactivation of latent provirus were estimated in a cross-sectional manner at necropsy,

comparing 9 animals treated with TNF-α versus 10 vehicle control animals. J-Lat cell

engraftment and GFP expression of J-Lat cells were analyzed as described in **Figure 5A**. Analyses

of all tissues demonstrated no significant effect of TNF-α treatment on J-Lat cell engraftment

levels within the respective compartment (**Figure 6C**). However, TNF-α treatment did lead to

significant increases in the frequency of GFP-positive cells as well as GFP MFI among engrafted

J-Lat cells (**Figure 6D-F**). Comparing the two populations (TNF-α vs vehicle control)

demonstrated a GFP expression fold-change of 3.2 in the BM, 17.8 in the spleen, 11.7 in the

lung, and 12.7 in the PB, illustrating potent and significant viral reactivation.

Discussion

The development of an HIV cure will be accelerated by the deployment of a convenient and

cost-effective animal model that enables the determination of an agent's therapeutic efficacy

as it permeates through tissue-specific barriers and the circulatory, respiratory and excretory

systems. We present a novel humanized mouse model of HIV latency that aims to address this

gap. In this study, we established a reliable method to discriminate between J-Lat and mouse

cells (**Figure 1** and **Figure 4**) and demonstrated robust engraftment of these cells in multiple

tissues sites (**Figure 4** and **Figure 5**). Significant viral reactivation was observed in TNF-α-treated

animals with respect to vehicle control-treated animals (**Figure 6D-F**). Importantly, although *in*

vivo treatment with TNF-α induced GFP expression and thus viral reactivation by 3.2 (BM), 17.8

(spleen), 11.7 (lung), and 12.7 (PB) -fold compared to vehicle control treated animals (**Figure**

6E), TNF-α treatment *in vitro* induced a 25.3-fold increase in GFP expressing cells (**Figure 6B**).

Moreover, the observed variability in TNF-α-mediated viral reactivation across tissue sites

demonstrates that the efficacy of latency reversal approaches can differ dramatically between

distinct anatomical niches due to pharmacokinetic factors and drug penetration efficiencies.

These data emphasize the importance of testing the efficacies of promising and potent

regimens in an *in vivo* system, and reinforce the relevance of the μ-Lat model.

Although not intended to serve as a pathophysiology model, the μ-Lat platform offers key

advantages over existing preclinical models focused on HIV eradication. Firstly, the model is

highly efficient; the timeframe to generate mouse colonies that are ready for administration of

experimental therapies is approximately three weeks. In comparison, even the simplified

humanized mouse model introduced by Kim et al.(31) requires four weeks for robust

engraftment of intraperitoneally injected human PBMCs in NSG mice and an additional 5 weeks

of incubation upon HIV infection. Secondly, the presence of the GFP reporter cassette in the

integrated J-Lat viral genome allows for rapid and convenient assessment of HIV latency

reversal *in vivo*, eliminating the requirement for PCR or culture-based diagnostics to determine

LRA potency. Thirdly, the model is characterized by high frequencies of HIV latently-infected

cells (e.g. in some experiments exceeding 30% engraftment in BM), and these cells are

distributed in a number of relevant anatomical sites that are central to HIV persistence. This

stands in contrast to organoid-based models (e.g. the *SCID-hu thy/liv* mouse model) that

exclusively involves viral colonization within a xenografted tissue(49). Lastly, the model is highly

scalable due to the low costs and limited labor requirements of the method involving the easily

and widely available J-Lat cell line (provided free of charge via the NIH AIDS Reagent Program).

Importantly, the clonal nature of cell lines promotes consistency and high reproducibility across

laboratories, while allowing for an easy transition from the petri-dish stage to an *in vivo*

evaluation system.

Similar to other animal models, there are caveats associated with the μ-Lat model. Although

the GFP reporter present within the J-Lat cell conveniently broadcasts transactivation of the

viral LTR and induction of viral transcription (informative precursor steps that may lead to the

production of viral antigen and release of virus), other assays will need to be applied to the μ-

Lat model to specifically quantify viral production or clearance of infected cells mediated by a

candidate HIV eradication agent. In addition, the efficiency of the model is largely driven by the

administration of an HIV latently-infected cell line, rather than differentiated primary cells. As

presented here, a single latent clone with a single proviral integration site was injected into

mice. Proviral integration site is known to affect viral latency and responsiveness to LRA

administration(4,50–52). However, this issue could be ameliorated by mixing different HIV

latently-infected cell clones at specific ratios, thereby achieving higher integration site

heterogeneity to more accurately represent *in vivo* variability. For instance, there are 11 J-Lat

clones currently available via the NIH AIDS Reagent Program, each of which is characterized by

a distinct proviral integration site. These clones often behave differently from each other as

well when exposed to LRAs *in vitro*, likely reflecting a diversity of molecular mechanisms

reinforcing viral latency across clones(43). This mechanistic diversity could further enhance the

predictive potential of the μ-Lat model.

Beyond concerns regarding integration site heterogeneity, the immortalized nature of the J-Lat

cell line may impact molecular and regulatory pathways that affect HIV latency. However, the

widespread usage of the J-Lat model and its derivative clones to examine viral latency and to

evaluate the efficacy of HIV cure strategies *in vitro* speak to the model's utility(53–62). Ample

data from primary cell-based models of latency and experiments involving *ex vivo*

administration of LRAs to cells obtained from HIV-infected individuals on ART suggest that

differences between applied models and between individuals can have dramatic effects on the

establishment, maintenance, and reversal of HIV latency(43,63,64). Moreover, profiling of HIV

latency in multiple tissue sites has demonstrated that the nature of viral latency may even vary

extensively within a single infected individual(65–67). Therefore, it needs to be considered that

any applied model system or cells from any given individual may introduce biases that impact

generalizability, just as a cell line-based system(26).

Our work described here represents a proof of concept, demonstrating that the engraftment of

a cell line-based model of HIV latency may constitute a useful testbed for HIV cure strategies.

This general approach is highly versatile and should allow for a broad range of infected cell

types to be examined *in vivo*. For example, HIV latency in the myeloid compartment is likely

critical to viral persistence(35,68–71), and multiple reports suggest this compartment may

respond quite differently to curative approaches, as compared to lymphoid reservoirs(72,73). It

may be fruitful to inject the U1 HIV chronically-infected promonocytic cell line(74) into NSG

mice to examine LRA responses in myeloid cells. Building further on the myeloid theme, the

central nervous system (CNS) compartment is an important viral sanctuary site in the setting of

ART(73,75–78), and HIV eradication approaches will almost certainly face unique challenges in

this niche(78,79). As direct injection of human cells into the murine CNS has been used

successfully as an engraftment approach(80–84), site-specific injection of the recently

developed HC69.5 HIV latently-infected microglial cell line(85,86) into the brains of NSG mice

may provide a convenient platform to gauge CNS-focused cure strategies.

Beyond preclinical investigation of LRAs, the μ-Lat framework may provide a convenient model

system to evaluate gene therapy-based HIV eradication approaches. Gene therapy approaches

targeting HIV infection generally fall into two categories: 1) Gene editing can be used to target

or "excise" the HIV provirus directly in infected cells as an eradication approach(87,88) or 2)

Editing can be used to modulate host cells to render them refractory to HIV infection and/or

potentiate antiviral immune responses(89). In the former case, *in vivo* delivery of the gene

therapy modality will likely be necessary to pervasively attack the HIV reservoir within a broad

range of tissue sites in infected individuals. The μ-Lat model is well-suited to investigate gene

delivery in this context, as it is characterized by robust engraftment of HIV latently-infected

cells into diverse anatomical niches. This will allow for efficient assessment of gene therapy

vector dissemination and antiviral function across anatomic sites. The LTR-driven GFP cassette

within the J-Lat integrated provirus may facilitate this assessment; direct targeting of the GFP

sequence may be used to examine vector trafficking and delivery, while specific targeting of the

HIV LTR as a cure approach would be associated with a convenient readout (relative loss of GFP

expression upon induced latency reversal).

In summary, the μ-Lat model is optimized for efficient and scalable evaluation of select HIV

eradication approaches *in vivo*, allowing determination of therapeutic efficacy in addition to

essential safety, tolerability, and pharmacokinetic parameters. Further development and

diversification of the μ-Lat model system may enable convenient testing of HIV eradication

approaches, including antiviral gene therapy strategies, in a range of cell types and tissue sites

in vivo.

Acknowledgments

This study was supported by the National Institutes of Health grants R01 AI150449 (SKP) and R01 MH112457 (SKP), and was additionally supported by a grant from the National Institutes of Health, University of California San Francisco-Gladstone Institute of Virology & Immunology Center for AIDS Research (P30 AI027763). We thank Dr. Roland Schwarzer for valuable input and careful review of the manuscript.

Conflict of interest statement

The authors have declared that no conflict of interest exists.

Author contributions

HSS, PPT, KAR, MSB conceived and performed experiments; PPT, RG, AGG, MOM performed *in vivo* experiments; HSS, PPT collected and analyzed the data; HSS, PPT, MOM, SKP designed the study and wrote the manuscript.

References

1.   Wong JK, Hezareh M, Günthard HF, Havlir D V, Ignacio CC, Spina CA, et al. Recovery of

replication-competent HIV despite prolonged suppression of plasma viremia. Science

(80-) [Internet]. 1997 Nov 14 [cited 2019 Nov 15];278(5341):1291–5. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/9360926

2.   Piot P, Bartos M, Ghys PD, Walker N, Schwartländer B. The global impact of HIV/AIDS.

Nature [Internet]. 2001 Apr 19 [cited 2019 Nov 15];410(6831):968–73. Available from:

http://www.nature.com/articles/35073639

3.   Pitman MC, Lau JSY, McMahon JH, Lewin SR. Barriers and strategies to achieve a cure for

HIV. Lancet HIV [Internet]. 2018 Jun [cited 2019 Nov 15];5(6):e317–28. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0031938416312148

4.   Schwarzer, Gramatica, Greene. Reduce and Control: A Combinatorial Strategy for

Achieving Sustained HIV Remissions in the Absence of Antiretroviral Therapy. Viruses

[Internet]. 2020 Feb 8;12(2):188. Available from: https://www.mdpi.com/1999-

4915/12/2/188

5.   Policicchio BB, Pandrea I, Apetrei C. Animal Models for HIV Cure Research. Front

Immunol [Internet]. 2016 Jan 28 [cited 2019 Nov 15];7(JAN):12. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/26858716

6.   Nixon CC, Mavigner M, Silvestri G, Garcia JV. In vivo models of human immunodeficiency

virus persistence and cure strategies. J Infect Dis. 2017;215(Suppl 3):S142–51.

7.   Denton PW, Søgaard OS, Tolstrup M. Using animal models to overcome temporal, spatial

and combinatorial challenges in HIV persistence research. J Transl Med [Internet]. 2016

Dec   9;14(1):44.   Available   from: http://www.translational-

medicine.com/content/14/1/44

8.   Shytaj IL, Norelli S, Chirullo B, Della Corte A, Collins M, Yalley-Ogunro J, et al. A highly

intensified ART regimen induces long-term viral suppression and restriction of the viral

reservoir in a simian AIDS model. PLoS Pathog. 2012;8(6).

9.   Dinoso JB, Rabi SA, Blankson JN, Gama L, Mankowski JL, Siliciano RF, et al. A Simian

Immunodeficiency Virus-Infected Macaque Model To Study Viral Reservoirs That Persist

during Highly Active Antiretroviral Therapy. J Virol. 2009;83(18):9247–57.

10. North TW, Higgins J, Deere JD, Hayes TL, Villalobos A, Adamson L, et al. Viral Sanctuaries

during Highly Active Antiretroviral Therapy in a Nonhuman Primate Model for AIDS. J

Virol. 2010;84(6):2913–22.

11. Kumar N, Chahroudi A, Silvestri G. Animal models to achieve an HIV cure. Curr Opin HIV

AIDS   [Internet].   2016   Jul;11(4):432–41.   Available   from:

https://linkinghub.elsevier.com/retrieve/pii/S0031938416312148

12. Sayah DM, Sokolskaja E, Berthoux L, Luban J. Cyclophilin A retrotransposition into TRIM5

explains owl monkey resistance to HIV-1. Nature [Internet]. 2004 Jul 7;430(6999):569–

73. Available from: http://www.nature.com/articles/nature02777

13. Mariani R, Chen D, Schröfelbauer B, Navarro F, König R, Bollman B, et al. Species-specific

481   exclusion of APOBEC3G from HIV-1 virions by Vif. Cell. 2003;114(1):21–31.

482 14. Baldauf HM, Pan X, Erikson E, Schmidt S, Daddacha W, Burggraf M, et al. SAMHD1

483   restricts HIV-1 infection in resting CD4 + T cells. Nat Med. 2012;18(11):1682–7.

484 15. Manel N, Littman DR. Hiding in plain sight: How HIV evades innate immune responses.

485   Cell [Internet]. 2011;147(2):271–4. Available from:

486   http://dx.doi.org/10.1016/j.cell.2011.09.010

487 16. Rehwinkel J, Maelfait J, Bridgeman A, Rigby R, Hayward B, Liberatore RA, et al. SAMHD1-

488   dependent retroviral control and escape in mice. EMBO J [Internet]. 2013;32(18):2454–

489   62. Available from: http://dx.doi.org/10.1038/emboj.2013.163

490 17. Behrendt R, Schumann T, Gerbaulet A, Nguyen LA, Schubert N, Alexopoulou D, et al.

491   Mouse SAMHD1 Has Antiretroviral Activity and Suppresses a Spontaneous Cell-Intrinsic

492   Antiviral Response. Cell Rep [Internet]. 2013 Aug;4(4):689–96. Available from:

493   https://linkinghub.elsevier.com/retrieve/pii/S2211124713003999

494 18. Wittmann S, Behrendt R, Eissmann K, Volkmann B, Thomas D, Ebert T, et al.

495   Phosphorylation of murine SAMHD1 regulates its antiretroviral activity. Retrovirology.

496   2015;12(1):1–15.

497 19. Seay K, Qi X, Zheng JH, Zhang C, Chen K, Dutta M, et al. Mice Transgenic for CD4-Specific

498   Human CD4, CCR5 and Cyclin T1 Expression: A New Model for Investigating HIV-1

499   Transmission and Treatment Efficacy. PLoS One. 2013;8(5).

500 20. Rahim MMA, Chrobak P, Hu C, Hanna Z, Jolicoeur P. Adult AIDS-Like Disease in a Novel

Inducible Human Immunodeficiency Virus Type 1 Nef Transgenic Mouse Model: CD4+ T-

Cell Activation Is Nef Dependent and Can Occur in the Absence of Lymphophenia. Vol.

83, Journal of Virology. 2009. p. 11830–46.

21. Hanna Z, Weng X, Kay DG, Poudrier J, Lowell C, Jolicoeur P. The Pathogenicity of Human

Immunodeficiency Virus (HIV) Type 1 Nef in CD4C/HIV Transgenic Mice Is Abolished by

Mutation of Its SH3-Binding Domain, and Disease Development Is Delayed in the Absence

of Hck. Vol. 75, Journal of Virology. 2001. p. 9378–92.

22. Dickie P, Felser J, Eckhaus M, Bryant J, Silver J, Marinos N, et al. HIV-associated

nephropathy in transgenic mice expressing HIV-1 genes. Virology. 1991;185(1):109–19.

23. Maung R, Hoefer MM, Sanchez AB, Sejbuk NE, Medders KE, Desai MK, et al. CCR5

Knockout Prevents Neuronal Injury and Behavioral Impairment Induced in a Transgenic

Mouse Model by a CXCR4-Using HIV-1 Glycoprotein 120. J Immunol [Internet]. 2014 Aug

15;193(4):1895–910.   Available   from:

https://linkinghub.elsevier.com/retrieve/pii/S0031938416312148

24. Marsden MD, Zack JA. Studies of retroviral infection in humanized mice. Virology

[Internet].   2015   May;479–480(1):297–309.   Available   from:

https://linkinghub.elsevier.com/retrieve/pii/S004268221500029X

25. Victor Garcia J. Humanized mice for HIV and AIDS research. Curr Opin Virol. 2016;19:56–

64.

26. Whitney JB, Brad Jones R. In Vitro and In Vivo Models of HIV Latency. In: Zhang L, Lewin

521  SR, editors. HIV Vaccines and Cure◻: The Path Towards Finding an Effective Cure and

522  Vaccine [Internet]. Singapore: Springer Singapore; 2018. p. 241–63. Available from:

523  https://doi.org/10.1007/978-981-13-0484-2_10

524 27. Krishnakumar V, Durairajan SSK, Alagarasu K, Li M, Dash AP. Recent updates on mouse

525  models for human immunodeficiency, influenza, and dengue viral infections. Viruses.

526  2019;11(3).

527 28. Zhou J, Takahashi M, Burnett JC, Li S, Rossi JJ, Li H, et al. HIV Replication and Latency in a

528  Humanized NSG Mouse Model during Suppressive Oral Combinational Antiretroviral

529  Therapy. J Virol. 2018;92(7):e02118-17.

530 29. Lai F, Chen Q. Humanized mouse models for the study of infection and pathogenesis of

531  human viruses. Vol. 10, Viruses. 2018. p. 1–17.

532 30. Mosier DE, Gulizia RJ, Baird SM, Wilson DB. Transfer of a functional human immune

533  system to mice with severe combined immunodeficiency. Nature. 1988;335(6187):256–

534  9.

535 31. Kim KCKCKCKC, Choi BS, Kim KCKCKCKC, Park KH, Lee HJ, Cho YK, et al. A Simple Mouse

536  Model for the Study of Human Immunodeficiency Virus. AIDS Res Hum Retroviruses.

537  2016;32(2):194–202.

538 32. Flerin NC, Bardhi A, Zheng JH, Korom M, Folkvord J, Kovacs C, et al. Establishment of a

539  Novel Humanized Mouse Model To Investigate In Vivo Activation and Depletion of

540  Patient-Derived HIV Latent Reservoirs. Silvestri G, editor. J Virol [Internet]. 2019 Jan

9;93(6):1–18. Available from: http://jvi.asm.org/lookup/doi/10.1128/JVI.02051-18

33. Lapidot T, Pflumio F, Doedens M, Murdoch B, Williams DE, Dick JE. Cytokine stimulation

of multilineage hematopoiesis from immature human cells engrafted in SCID mice.

Science   (80-   )   [Internet].   1992;255(5048):1137–41.   Available   from:

https://science.sciencemag.org/content/255/5048/1137

34. Honeycutt JB, Wahl A, Archin N, Choudhary S, Margolis D, Garcia J V. HIV-1 infection,

response to treatment and establishment of viral latency in a novel humanized T cell-

only mouse (TOM) model. Retrovirology [Internet]. 2013;10(1):1. Available from:

Retrovirology

35. Honeycutt JB, Wahl A, Baker C, Spagnuolo RA, Foster J, Zakharova O, et al. Macrophages

sustain HIV replication in vivo independently of T cells Find the latest version⬛:

Macrophages sustain HIV replication in vivo independently of T cells. J Clin Invest.

2016;126(4):1353–66.

36. McCune J, Namikawa R, Kaneshima H, Shultz L, Lieberman M, Weissman I. The SCID-hu

mouse: murine model for the analysis of human hematolymphoid differentiation and

function. Science (80-) [Internet]. 1988 Sep 23;241(4873):1632–9. Available from:

https://science.sciencemag.org/content/241/4873/1632

37. Namikawa R, Weilbaecher KN, Kaneshima H, Yee EJ, McCune JM. Longterm human

hematopoiesis in the SCID-hu mouse. J Exp Med. 1990;172(4):1055–63.

38. Lan P, Tonomura N, Shimizu A, Wang S, Yang YG. Reconstitution of a functional human

immune system in immunodeficient mice through combined human fetal thymus/liver

and CD34+ cell transplantation. Blood. 2006;108(2):487–92.

39. Melkus MW, Estes JD, Padgett-Thomas A, Gatlin J, Denton PW, Othieno FA, et al.

Humanized mice mount specific adaptive and innate immune responses to EBV and TSST-

1. Nat Med. 2006;12(11):1316–22.

40. McCune JM, Weissman IL. The Ban on US Government Funding Research Using Human

Fetal Tissues: How Does This Fit with the NIH Mission to Advance Medical Science for the

Benefit of the Citizenry? [Internet]. Vol. 13, Stem Cell Reports. 2019 [cited 2019 Nov 19].

p. 777–86. Available from: http://www.ncbi.nlm.nih.gov/pubmed/31722191

41. Jordan A, Bisgrove D, Verdin E. HIV reproducibly establishes a latent infection after acute

infection of T cells in vitro. 2003 [cited 2017 Jun 26];22(8). Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC154479/pdf/cdg188.pdf

42. Stoszko M, De Crignis E, Rokx C, Khalid MM, Lungu C, Palstra RJ, et al. Small Molecule

Inhibitors of BAF; A Promising Family of Compounds in HIV-1 Latency Reversal.

EBioMedicine [Internet]. 2016;3:108–21. Available from:

http://dx.doi.org/10.1016/j.ebiom.2015.11.047

43. Spina CA, Anderson J, Archin NM, Bosque A, Chan J, Famiglietti M, et al. An In-Depth

Comparison of Latent HIV-1 Reactivation in Multiple Cell Model Systems and Resting

CD4+ T Cells from Aviremic Patients. Emerman M, editor. PLoS Pathog [Internet]. 2013

Dec 26;9(12):e1003834. Available from:

http://dx.plos.org/10.1371/journal.ppat.1003834

44.   Spivak AM, Planelles V. Novel Latency Reversal Agents for HIV-1 Cure. Annu Rev Med

583   [Internet]. 2018 Jan 29 [cited 2018 Aug 15];69(1):421–36. Available from:

584   https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5892446/

585   45.   Jordan A, Defechereux P, Verdin E. The site of HIV-1 integration in the human genome

586   determines basal transcriptional activity and response to Tat transactivation. EMBO J.

587   2001;20(7):1726–38.

588   46.   Beyer AI, Muench MO. Comparison of human hematopoietic reconstitution in different

589   strains of immunodeficient mice. Stem Cells Dev. 2017;26(2):102–12.

590   47.   Slørdal L, Muench MO, Warren DJ, Moore MAS. Radioprotection by murine and human

591   tumor-necrosis factor: Dose-dependent effects on hematopoiesis in the mouse. Eur J

592   Haematol. 1989;43(5):428–34.

593   48.   Hoggatt J, Hoggatt AF, Tate TA, Fortman J, Pelus LM. Bleeding the laboratory mouse: Not

594   all methods are equal. Exp Hematol [Internet]. 2016;44(2):132-137.e1. Available from:

595   http://dx.doi.org/10.1016/j.exphem.2015.10.008

596   49.   Namikawa R, Kaneshima H, Lieberman M, Weissman IL, McCune JM. Infection of the

597   SCID-hu mouse by HIV-1. Science (80-). 1988;242(4886):1684–6.

598   50.   Symons J, Cameron PU, Lewin SR. HIV integration sites and implications for maintenance

599   of the reservoir. Curr Opin HIV AIDS [Internet]. 2018 Mar;13(2):152–9. Available from:

600   http://insights.ovid.com/crossref?an=01222929-201803000-00009

51.   Anderson EM, Maldarelli F. The role of integration and clonal expansion in HIV infection:

Live long and prosper 11 Medical and Health Sciences 1108 Medical Microbiology Ben

Berkhout, Alexander Pasternak. Retrovirology [Internet]. 2018;15(1):1–22. Available

from: https://doi.org/10.1186/s12977-018-0448-8

52. Maldarelli F, Wu X, Su L, Simonetti FR, Shao W, Hill S, et al. Specific HIV integration sites

are linked to clonal expansion and persistence of infected cells. Science (80-) [Internet].

2014   Jul   11;345(6193):179–83.   Available   from:

https://www.sciencemag.org/lookup/doi/10.1126/science.1254194

53. Abdel-Mohsen M, Chavez L, Tandon R, Chew GM, Deng X, Danesh A, et al. Human

Galectin-9 Is a Potent Mediator of HIV Transcription and Reactivation. PLoS Pathog

[Internet].   2016;12(6):1–28.   Available   from:

http://dx.doi.org/10.1371/journal.ppat.1005677

54. Jiang G, Mendes EA, Kaiser P, Wong DP, Tang Y, Cai I, et al. Synergistic Reactivation of

Latent HIV Expression by Ingenol-3-Angelate, PEP005, Targeted NF-kB Signaling in

Combination with JQ1 Induced p-TEFb Activation. PLoS Pathog. 2015;11(7):1–27.

55. Darcis G, Kula A, Bouchat S, Fujinaga K, Corazza F, Ait-Ammar A, et al. An In-Depth

Comparison of Latency-Reversing Agent Combinations in Various In Vitro and Ex Vivo

HIV-1 Latency Models Identified Bryostatin-1+JQ1 and Ingenol-B+JQ1 to Potently

Reactivate Viral Gene Expression. PLoS Pathog. 2015;11(7):1–36.

56. Reuse S, Calao M, Kabeya K, Guiguen A, Gatot JS, Quivy V, et al. Synergistic activation of

HIV-1 expression by deacetylase inhibitors and prostratin: Implications for treatment of

latent infection. PLoS One. 2009;4(6).

57.   Brogdon J, Ziani W, Wang X, Veazey RS, Xu H. In vitro effects of the small-molecule

624   protein kinase C agonists on HIV latency reactivation. Sci Rep [Internet].

625   2016;6(November):1–8. Available from: http://dx.doi.org/10.1038/srep39032

626   58.   Chen D, Wang H, Aweya JJ, Chen Y, Chen M, Wu X, et al. HMBA Enhances Prostratin-

627   Induced Activation of Latent HIV-1 via Suppressing the Expression of Negative Feedback

628   Regulator A20/TNFAIP3 in NF- B Signaling. Biomed Res Int. 2016;2016.

629   59.   Rochat MA, Schlaepfer E, Speck RF. Promising Role of Toll-Like Receptor 8 Agonist in

630   Concert with Prostratin for Activation of Silent HIV. J Virol. 2017;91(4):1–21.

631   60.   Klase Z, Yedavalli VSRKRK, Houzet L, Perkins M, Maldarelli F, Brenchley J, et al. Activation

632   of HIV-1 from Latent Infection via Synergy of RUNX1 Inhibitor Ro5-3335 and SAHA. Luban

633   J, editor. PLoS Pathog [Internet]. 2014 Mar 20;10(3):e1003997. Available from:

634   https://dx.plos.org/10.1371/journal.ppat.1003997

635   61.   Bouchat S, Delacourt N, Kula A, Darcis G, Van Driessche B, Corazza F, et al. Sequential

636   treatment with 5-aza-2-deoxycytidine and deacetylase inhibitors reactivates HIV -1 .

637   EMBO Mol Med. 2016;8(2):117–38.

638   62.   Kutsch O, Wolschendorf F, Planelles V. Facts and Fiction: Cellular Models for High

639   Throughput Screening for HIV-1 Reactivating Drugs. Curr HIV Res [Internet]. 2011 Dec

640   1;9(8):568–78.   Available   from:

641   http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1570-

642   162X&volume=9&issue=8&spage=568

643   63.   Bradley T, Ferrari G, Haynes BF, Margolis DM, Browne EP. Single-Cell Analysis of

644   Quiescent HIV Infection Reveals Host Transcriptional Profiles that Regulate Proviral

645   Latency. Cell Rep [Internet]. 2018 Oct;25(1):107-117.e3. Available from:

646   https://linkinghub.elsevier.com/retrieve/pii/S2211124718314505

647   64.   Telwatte S, Morón-López S, Aran D, Kim P, Hsieh C, Joshi S, et al. Heterogeneity in HIV

648   and cellular transcription profiles in cell line models of latent and productive infection:

649   implications for HIV latency. Retrovirology [Internet]. 2019;16(1):32. Available from:

650   https://doi.org/10.1186/s12977-019-0494-x

651   65.   Dahl V, Gisslen M, Hagberg L, Peterson J, Shao W, Spudich S, et al. An example of

652   genetically distinct HIV type 1 variants in cerebrospinal fluid and plasma during

653   suppressive therapy. J Infect Dis. 2014;209(10):1618–22.

654   66.   T. Blackard J. HIV Compartmentalization: A Review on a Clinically Important

655   Phenomenon. Curr HIV Res. 2012;10(2):133–42.

656   67.   Telwatte S, Lee S, Somsouk M, Hatano H, Baker C, Kaiser P, et al. Gut and blood differ in

657   constitutive blocks to HIV transcription, suggesting tissue-specific differences in the

658   mechanisms that govern HIV latency. PLoS Pathog. 2018;14(11):1–28.

659   68.   Thompson KA, Cherry CL, Bell JE, McLean CA. Brain cell reservoirs of latent virus in

660   presymptomatic HIV-infected individuals. Am J Pathol [Internet]. 2011;179(4):1623–9.

661   Available from: http://dx.doi.org/10.1016/j.ajpath.2011.06.039

69.   Castellano P, Prevedel L, Valdebenito S, Eugenin EA. HIV infection and latency induce a

unique metabolic signature in human macrophages. Sci Rep [Internet]. 2019;9(1):1–14.

Available from: http://dx.doi.org/10.1038/s41598-019-39898-5

70.   Wong ME, Jaworowski A, Hearps AC. The HIV Reservoir in Monocytes and Macrophages.

Front   Immunol   [Internet].   2019   Jun   26;10(June).   Available   from:

https://www.frontiersin.org/article/10.3389/fimmu.2019.01435/full

71.   Gama L CJACVRACGSPDFEQSSEBBLMMPKBSMLMJMGFOS. Myeloid and CD4 T Cells

Comprise the Latent Reservoir in Antiretroviral Therapy-Suppressed SIVmac251-Infected

Macaques. Silvestri G, Bennink JR, editors. MBio [Internet]. 2019 Aug 20;10(4):1–19.

Available from: http://mbio.asm.org/lookup/doi/10.1128/mBio.01659-19

72.   Gray LR, On H, Roberts E, Lu HK, Moso MA, Raison JA, et al. Toxicity and in vitro activity

of HIV-1 latency-reversing agents in primary CNS cells. J Neurovirol [Internet]. 2016 Aug

4;22(4):455–63. Available from: http://link.springer.com/10.1007/s13365-015-0413-4

73.   Hellmuth J, Valcour V, Spudich S. CNS reservoirs for HIV: implications for eradication. J

virus   Erad   [Internet].   2015;1(2):67–71.   Available   from:

http://www.ncbi.nlm.nih.gov/pubmed/26430703%0Ahttp://www.pubmedcentral.nih.go

v/articlerender.fcgi?artid=PMC4586130

74.   Folks TM, Justement J, Kinter A, Dinarello CA, Fauci AS, Folks TM, et al. Cytokine-Induced

Expression of HIV-1 in a Chronically Infected Promonocyte Cell Line. Science (80-)

[Internet]. 1987;238(4828):800–2. Available from: https://www.jstor.org/stable/1700363

75.   Joseph SB, Arrildt KT, Sturdevant CB, Swanstrom R. HIV-1 target cells in the CNS. J

Neurovirol [Internet]. 2015 Jun 19;21(3):276–89. Available from:

http://link.springer.com/10.1007/s13365-014-0287-x

76.   Barat C, Proust A, Deshiere A, Leboeuf M, Drouin J, Tremblay MJ. Astrocytes sustain long-

term productive HIV-1 infection without establishment of reactivable viral latency. Glia

[Internet]. 2018 Jul [cited 2018 Aug 15];66(7):1363–81. Available from:

https://onlinelibrary.wiley.com/doi/abs/10.1002/glia.23310

77.   Edén A, Fuchs D, Hagberg L, Nilsson S, Spudich S, Svennerholm B, et al. HIV-1 Viral Escape

in Cerebrospinal Fluid of Subjects on Suppressive Antiretroviral Treatment. J Infect Dis.

2010;202(12):1819–25.

78.   Churchill M, Nath A. Where does HIV hide? A focus on the central nervous system. Curr

Opin HIV AIDS [Internet]. 2013 May;8(3):165–9. Available from:

http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=01222929

-201305000-00003

79.   Nath A, Clements JE. Eradication of HIV from the brain: Reasons for pause. Aids.

2011;25(5):577–80.

80.   Tyor WR, Power C, Gendelman HE, Markham RB. A model of human immunodeficiency

virus encephalitis in scid mice. Proc Natl Acad Sci U S A. 1993;90(18):8658–62.

81.   Potula R, Poluektova L, Knipe B, Chrastil J, Heilman D, Dou H, et al. Inhibition of

indoleamine 2,3-dioxygenase (IDO) enhances elimination of virus-infected macrophages

in an animal model of HIV-1 encephalitis. Blood. 2005;106(7):2382–90.

82.   Poluektova LY, Munn DH, Persidsky Y, Gendelman HE. Generation of Cytotoxic T Cells

Against Virus-Infected Human Brain Macrophages in a Murine Model of HIV-1

Encephalitis. J Immunol [Internet]. 2002 Apr 15;168(8):3941–9. Available from:

http://www.jimmunol.org/lookup/doi/10.4049/jimmunol.168.8.3941

83.   Poluektova L, Gorantla S, Faraci J, Birusingh K, Dou H, Gendelman HE. Neuroregulatory

Events Follow Adaptive Immune-Mediated Elimination of HIV-1-Infected Macrophages:

Studies in a Murine Model of Viral Encephalitis. J Immunol [Internet]. 2004 Jun

15;172(12):7610–7.   Available   from:

http://www.jimmunol.org/lookup/doi/10.4049/jimmunol.172.12.7610

84.   Persidsky Y, Limoges J, McComb R, Bock P, Baldwin T, Tyor W, et al. Human

immunodeficiency virus encephalitis in SCID mice. Am J Pathol. 1996;149(3):1027–53.

85.   Garcia-Mesa Y, Jay TR, Checkley MA, Luttge B, Dobrowolski C, Valadkhan S, et al.

Immortalization of primary microglia: a new platform to study HIV regulation in the

central nervous system. J Neurovirol [Internet]. 2017 [cited 2018 Sep 5];23(1):47–66.

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5329090/

86.   Alvarez-Carbonell D, Garcia-Mesa Y, Milne S, Das B, Dobrowolski C, Rojas R, et al. Toll-like

receptor 3 activation selectively reverses HIV latency in microglial cells. Retrovirology

[Internet].   2017;14(1):9.   Available   from:

http://retrovirology.biomedcentral.com/articles/10.1186/s12977-017-0335-8

87.   Yin C, Zhang T, Qu X, Zhang Y, Putatunda R, Xiao X, et al. In Vivo Excision of HIV-1 Provirus

by saCas9 and Multiplex Single-Guide RNAs in Animal Models. Mol Ther [Internet]. 2017

May   3   [cited   2018   Aug   16];25(5):1168–86.   Available   from:

http://www.sciencedirect.com/science/article/pii/S1525001617301107

88.   Dash PK, Kaminski R, Bella R, Su H, Mathews S, Ahooyi TM, et al. Sequential LASER ART

and CRISPR Treatments Eliminate HIV-1 in a Subset of Infected Humanized Mice. Nat

Commun   [Internet].   2019   Dec   2;10(1):1–20.   Available   from:

http://dx.doi.org/10.1038/s41467-019-10366-y

89.   Tebas P, Stein D, Tang WW, Frank I, Wang SQ, Lee G, et al. Gene Editing of CCR5 in

Autologous CD4 T Cells of Persons Infected with HIV. N Engl J Med [Internet]. 2014 Mar 6

[cited   2019   Nov   15];370(10):901–10.   Available   from:

http://www.ncbi.nlm.nih.gov/pubmed/24597865

Figure Legends

Figure 1: The cell surface proteins CD147 and CD29 are abundantly and specifically expressed in J-Lat 11.1 cells. 1×10^6 J-Lat cells were stained with antibodies targeting select cell surface proteins to (A) measure the frequency of marker-positive cells and (B) mean fluorescence intensity (MFI) of proteins on the cell surface using flow cytometry. The MFI of marker signal positive cells was normalized to signal positive cells of respective isotype control stained samples. A representative experiment is summarized in the bar graphs. (C) Gating strategy to identify human J-Lat cells upon multicolor staining with CD147/CD29-APC (J-Lat cell marker) and CD45/TER-119/H-2Kd-Pacific Blue (mouse cell marker) *in vitro*. (D) Representative flow cytometry plots show multicolor staining of single cell suspensions prepared from different mouse tissues obtained from an untransplanted control animal. Mouse cells were stained, measured, and analyzed in the same way as *in vitro* J-Lat cells to evaluate specificity and background signal of human CD29 and CD147 antibodies. (E) Single cell suspensions were prepared from 8 mouse tissues and three different harvest methods for peripheral blood (PB) to determine the applicability of the multicolor staining for subsequent engraftment studies. BM = bone marrow, IEL = intraepithelial lymphocytes, LN = lymph node, PB = peripheral blood, r.-o. = retro-orbital.

Figure 2: Schematic representation of parameters that were examined and optimized to establish the μ-Lat model. J-Lat cell engraftment kinetics, J-Lat cell dosage, irradiation of recipient mice, and testing of J-Lat cell engraftment in different mouse strains were investigated.

Figure 3: Optimal engraftment levels of J-Lat cells are achieved 3 weeks post injection in

irradiated NSG mice using 10 x 10^6 J-Lat cells for transplantation. The parameters described in

Figure 2 were examined and optimized concomitantly by focusing on engraftment levels in the

BM. (A) <u>J-Lat cell engraftment kinetics</u>: Nonirradiated NSG-3GS mice were transplanted with 5 x

10^6 J-Lat cells and the BM harvested at indicated time points post cell injection. (B) <u>J-Lat cell</u>

<u>injection dosage</u>: J-Lat cell engraftment levels in the BM of nonirradiated NSG-3GS were

compared using different initial injection cell doses of 5 x 10^6 J-Lat cells (BM harvested 18 days

post cell injection) or 10 x 10^6 J-Lat cells (BM harvested 14 days post cell injection). (C)

<u>Irradiation of recipient mice</u>: NSG-3GS recipient mice were nonirradiated (nonIRR) or irradiated

(IRR) before transplantation of 10 x 10^6 J-Lat cells, and engraftment levels were determined two

weeks post cell transplantation. (D) <u>Mouse strain selection</u>: 10 x 10^6 J-Lat cells were injected

either into irradiated NSG-3GS or irradiated NSG mice and engraftment levels in the BM were

measured 3 weeks post cell injection and 24h post TNF-α treatment. Each dot represents an

individual animal for the respective condition. Each plot represents an independent experiment

with n ≥ 3. P values in (A) were determined using the nonparametric Kruskal-Wallis test with

the uncorrected Dunn's test for multiple comparisons. P values in (B-D) were determined using

an unpaired, two-tailed t-test. A standard P<0.05 significance threshold was used.

Figure 4: J-Lat cells engraft successfully in several tissues in transplanted NSG mice. (A) Gating

strategy to identify human J-Lat cells in harvested mouse tissues exemplified here with BM. (B)

Representative flow cytometry plots are shown for engraftment levels across tissue sites

observed at necropsy and engraftment levels in PB (n = 5) using three different harvest

approaches: r.-o. bleeding, tail vein bleeding, and heart bleeding. Bar graphs summarize (C)

frequency, (D) MFI, and (E) number of engrafted J-Lat cells in the respective tissue. Each data

point represents an individual animal. Colors indicate tissues harvested from the same animal.

Error bars show the standard error of the mean (SEM).

Figure 5: Engrafted J-Lat cells exhibit low basal levels of GFP expression 3 weeks post cell

transplantation. (A) Gating strategy to identify human J-Lat cells and GFP background signal of

engrafted J-Lat cells in harvested mouse tissues illustrated here with a representative BM

sample. Representative flow cytometry plots are shown for (B) engraftment levels and (C) GFP

background signal across selected tissues observed at necropsy (n = 10). Bar graphs summarize

(D) cell frequency and (E) GFP expression levels of engrafted J-lat cells. Each data point

represents an individual animal. Colors indicate tissues harvested from the same animal. Error

bars show SEM.

Figure 6: TNF-α treatment reactivates latent HIV *in vivo* in the μ-Lat model. (A) Schematic

representation of the procedure to test LRAs *in vivo* using the μ-Lat model: (1) Mice receiving

cells of interest for transplantation are irradiated 3 hours prior to cell injection. (2) Each mouse

receives 10 x 10^6 J-Lat cells via intravenous injection. (3) Three weeks post injection (21 days)

mice are treated for 24h with LRAs of interest, followed by (4) necropsy, tissue harvest and

processing, and preparation of single-cell suspensions. (5) Single-cell suspensions are stained

for J-Lat and mouse cell markers to assess engraftment levels via flow cytometry. Reactivation

of latent provirus following LRA treatment is assessed by measuring GFP expression via flow

cytometry (the J-Lat provirus contains an LTR-driven GFP reporter). (B) J-Lat cells were treated

in vitro for 24h with 0.5% DMSO (vehicle control), 20 ng/µl TNF-α, 20 nM PMA/1 µM Ionomycin

(positive control) or left untreated (negative control) to assess viral reactivation based on GFP

expression (n = 5). (C) J-Lat engraftment levels were measured 24h post tail vein injection of 20

µg TNF-α (n = 9) or vehicle control (PBS, n = 10) in BM, spleen, lung and PB of mice. (D)

Representative flow cytometry plots demonstrating the effect of vehicle control (top panel) or

TNF-α treatment (bottom panel) on viral reactivation (GFP expression) in engrafted J-Lat cells.

The induction of GFP expression among engrafted J-Lat cells across tissues is summarized as (E)

frequency of GFP+ cells and (F) GFP MFI. One-way ANOVA and uncorrected Fisher's LSD for

multiple comparisons was used to analyze (B) *in vitro* reactivation data, and a mixed effects

model and uncorrected Fisher's LSD for multiple comparisons was used to analyze (C) *in vivo*

engraftment and (E-F) viral reactivation data. Colors indicate specific animals and tissues

obtained from the same animal. Error bars show SEM.

Figure 1

Figure 2

Model optimization

Engraftment kinetics

Cell injection 0

Tissue harvest ?

Cell dosage

low high

Irradiation

Mouse strain

Tissue harvest

Figure 3

Figure 4

A BM

Viable cells — Single cells — Scatter plot — Engrafted J-Lat cells

B Lung, Gut, Gut_IEL, Heart, Spleen, PB – heart bleed, PB – r.-o. bleed, PB – tail vein bleed

J-Lat cell marker (vertical axis)
Mouse cell marker (horizontal axis)

C % engrafted J-Lat cells

D MFI CD147/CD29 (J-Lat marker)

E # engrafted J-Lat cells

Figure 5

Figure 6