Ecology of domestic dogs *Canis familiaris* as an emerging reservoir of Guinea worm *Dracunculus*

medinensis infection

Robbie A. McDonald[1]*, Jared K. Wilson-Aggarwal[1], George J.F. Swan[1], Cecily E.D. Goodwin[1],

Tchonfienet Moundai[2], Dieudonné Sankara[3], Gautam Biswas[3], James A. Zingeser[4]

[1]Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, UK.

[2]Guinea Worm Eradication Programme, Ministry of Public Health, N'Djamena, Republic of Chad.

[3]World Health Organisation, Geneva, Switzerland.

[4]The Carter Center, Atlanta, GA, USA.

Email addresses: Robbie McDonald r.mcdonald@exeter.ac.uk, Jared Wilson-Aggarwal

jw695@exeter.ac.uk, George Swan g.swan@exeter.ac.uk, Cecily Goodwin c.goodwin@exeter.ac.uk,

Tchonfienet Moundai tchomcalvin@yahoo.fr, Dieudonné Sankara sankarad@who.int, Gautam Biswas

biswasg@who.int, James Zingeser Jim.Zingeser@cartercenter.org.

***Correspondence author:** Robbie McDonald, Environment and Sustainability Institute, University of

Exeter, Penryn Campus, Penryn, UK. email: r.mcdonald@exeter.ac.uk. Telephone +44 798 9069834.

Short title: Guinea worm in domestic dogs

Keywords: Neglected Tropical Diseases, zoonoses, domestic dogs, Guinea worm, disease eradication,

One Health, epidemiology, disease ecology, dracunculiasis

Statement of authorship: RM, JZ, GB and DS conceived the study. RM, JW-A, GS, TM and JZ

conducted the fieldwork. JW-A, CG and GS analysed samples and data. RM wrote the manuscript.

JW-A, CG, GS, DS and JZ revised the manuscript.

Abstract

Global eradication of human Guinea worm disease (dracunculiasis) has been set back by the

emergence of infections in animals, particularly domestic dogs *Canis familiaris*. The ecology and

epidemiology of this reservoir is unknown. We tracked dogs using GPS, inferred diets using stable

isotope analysis and analysed correlates of infection in Chad, where numbers of Guinea worm

infections are greatest. Dogs had small ranges that varied markedly among villages. Diets consisted

largely of human staples and human faeces. A minority of ponds, mostly <200 m from dog-owning

households, accounted for most dog exposure to potentially unsafe water. The risk of a dog having had

Guinea worm was reduced in dogs living in households providing water for animals but increased with

increasing fish consumption by dogs. Provision of safe water might reduce dog exposure to unsafe

water, while prioritisation of proactive temephos (Abate) application to the small number of ponds to

which dogs have most access is recommended. Fish might have an additional role as transport hosts

for Guinea worm, by concentrating copepods infected with worm larvae.

Author summary

Guinea worm is a parasite that causes profoundly debilitating disease in humans. An eradication

program has been successful in nearly eliminating the disease from people. However, the same worm

has now been found in domestic dogs and the frequency of detecting Guinea worm in dogs has been

increasing. This means that to eradicate Guinea worm, the infection must be eliminated in dogs as well

as in people. However, not much is known about the disease in dogs. This study is the first to

investigate dog ecology in relation to Guinea worm infection. We worked in the worst-affected

country, Chad. We attached GPS collars to dogs to track their ranging and use of water bodies and

analysed their diets using a forensic technique, based on analysing stable isotope compostion of their

whiskers and potential food items. We showed that dogs living in households that provided water to

their animals had a lower risk of having had Guinea worm and that dogs that ate more fish had an

increased risk. These findings suggest there is a classical route for worm transmission in dogs, via

drinking contaminated water, as well as a novel route, potentially by eating fish carrying a source of

infection.

Introduction

Guinea worm *Dracunculus medinensis* is a nematode parasite that causes Guinea worm disease (dracunculiasis) in humans. It was once widespread in Asia and Africa [1] but an eradication campaign has made impressive progress in reducing human cases from 3.5 million per annum in 21 countries in 1986, to only 28 in 2018, in Chad (17), South Sudan (10), and Angola (1) [2]. Alongside the near-absence of human cases, however, infections were detected in 1040 domestic dogs *Canis familiaris* and 25 domestic cats *Felis catus* in Chad, 11 dogs, five cats and one olive baboon *Papio anubis* in Ethiopia, and 18 dogs and two cats in Mali in 2018 [2] Adult Guinea worms emerging from humans and these non-human animals are genetically indistinguishable [3].

Prior to re-emergence in 2010, no human cases of Guinea worm were reported in Chad for ten years [4], suggesting an unknown reservoir of infection in non-human animals and/or undetected infection in humans, though surveillance of the disease in humans was also problematic at that time. Given the current high number of dogs infected relative to human cases and improved human surveillance in Chad, it is clear that non-human animals, particularly dogs, now constitute a contemporary reservoir of infection. Little is known about the epidemiology of Guinea worm in its non-human hosts. Although direct transmission between dogs and humans is not possible, dogs clearly maintain a persistent source of potential infection for humans in their shared environments. Thus, to eradicate Guinea worm, transmission of infection must be interrupted in non-human as well as human hosts [5]. Alongside civil insecurity in the remaining endemic areas, non-human animal infections are now the major impediment to Guinea worm eradication [6].

The pathways by which dogs acquire infection remain unknown, though two main possibilities exist. First, similar to the classical route for humans, dogs may acquire infection from drinking water containing infected copepods as the intermediate hosts. Second, dogs might consume a paratenic or transport host, most likely a fish or amphibian, that has itself eaten infected copepods [7]. In Chad, dogs drink from permanent and transient surface water bodies that often contain large copepod populations. Large-scale human exploitation of fish and frogs also provides the potential for dog consumption of their uncooked remains [7].

Options for interrupting dog infections are broadly similar to those for humans, and include: detection

and containment of hosts, treating water bodies with the organophosphate temephos (Abate), and

limiting exposure by reducing consumption of contaminated water and, potentially, food [8]. Since

different approaches are required to implement such interventions when aimed at free-ranging dogs, it

is important that risk factors are identified, to help inform management. To understand the

epidemiology and control of Guinea worm in dogs, we undertook a study of their ecology in Chad, to

characterise their ranging and access to water, as well as their diets and consumption of aquatic foods.

We show that both a classical route of transmission, relating to exposure to potentially unsafe water,

as well as a novel route, via consumption of fish, potentially as a paratenic host, influence the risk of

dog infection and highlight management implications for proactive water treatment.

Material and methods

Ethics statement

This study was approved by the University of Exeter College of Life and Environmental Sciences

(Penryn Campus) Ethics Committee (2016/1488). The Committee and the project adhered to the

"Guidelines for the treatment of animals in behavioural research and teaching" of the Association for

the Study of Animal Behaviour.

Study area and subject recruitment

Fieldwork was conducted between 24 June and 12 July 2016 in three settlements (Kakale, Magrao and

Largana), each comprised of multiple villages, situated in Guelendeng sous-préfecture, close to the

Chari River, Chad (Figure 1). Settlements were selected because of the high numbers of infections in

dogs and relative ease of access. The chief of each village provided approval and dog owners

individually provided consent. For each household, we recorded the GPS location and conducted a

questionnaire, recording whether dogs were taken hunting (reported by owners as entering the bush to

capture wild game, and classified as hunting or not) and whether water was provided for animals

within the household. For each dog, we recorded sex, age in months, body condition (on a scale of 1 =

emaciated to 9 = severely obese [9] and then classified as poor ≤2 or moderate ≥3), and whether the

dog had history of emergent adult Guinea worms. The Chad Guinea Worm Eradication Programme

(CGWEP) provided additional surveillance data identifying dogs with confirmed infections.

Dog tracking

All dogs, apart from pups that were <6 months old and were too small to be collared, were fitted with

adjustable nylon collars (Ancol, Walsall, UK) equipped with an iGotU GT600 GPS logger (Mobile

Action Technology, Taipei) with a 10 minute fix interval. After deployment for up to 14 days, collars

were retrieved and data downloaded. Locations before 6am the day after deployment and after 6am the

day before retrieval were excluded to avoid interference arising from our activities. Fixes that

suggested speeds greater than 20km/hr were considered spurious and excluded. In one village

(Gomba) in Largana, owners removed collars early, so locations after midnight prior to their removal

of collars were excluded. The dogs' core and total ranges were characterised by 60% kernel density

estimates and 100% minimum convex polygons, respectively.

DigitalGlobe satellite imagery was obtained for February, October and December 2012 for Magrao,

Kakale and Largana, respectively. This imagery was the nearest available in time to when our

fieldwork was conducted at the end of the dry season, and likely reflects the availability of surface

water earlier in the dry season. We sampled the distribution of ponds by drawing a box around all dog

locations for each settlement and visually searching images for ponds. For each dog, we quantified

exposure to potential sources of infection using four variables: 1. The number of GPS fixes in the

vicinity (locations within 100m) of a water body (ponds and the Chari River); 2. the number of

independent visits to the vicinity of water bodies (defined by an interval of 30 minutes between fixes);

3. The number of water bodies in core and total ranges; and 4. the number of different water bodies

visited. For dogs with small ranges and/or those living in households close to ponds or to the Chari

River, this would lead inevitably to relatively high measures of exposure, since these individuals

would, in effect, be constantly 'visiting' the vicinity of water bodies. We considered, nevertheless, that

this was an appropriate measure of variation among dogs in their relative exposure to potentially

unsafe water sources. For each pond, we measured the number of dog visits, the number of dogs that

visited and the number of visits by each dog (defined by an interval of 30 minutes between fixes). To

characterise the ponds most visited by dogs, ponds were ranked by number of visits by all dogs and

the distance of each pond to the nearest household with tracked dogs.

Variation in dog range sizes was analysed using linear models (LMs). Ranges were ln-transformed.

Models contained settlement, sex, age, body condition, use in hunting activity and household water

provision as explanatory variables, and the number of tracking days as a fixed effect to control for

effort. Variation in the time dogs spent in the vicinity of water bodies (the number of fixes within 100

m) was analysed using a general linear model (GLM) with the same explanatory variables, except for

use in hunting activity, plus the log of the number of tracking days as an offset to control for effort. A

generalised additive model (GAM) related the cumulative total of dog visits per water body to the log-

transformed distance from a household with tracked dogs.

Dog diets

Composition of dog diets was determined using stable isotope analysis of dog whiskers and putative

food items. Two whiskers were plucked from dogs in Kakale and Magrao during collar retrieval. The

period over which whiskers had grown, and diet could be inferred, was calculated by dividing whisker

length by growth rate; a subsample of dogs were fed ~50 g baits with 0.8 g (40-80 mg/kg) of

rhodamine B, a food colourant that stained whiskers with a fluorescent band. Growth rate (mm/day)

was calculated by dividing the distance between the whisker base and the distal end of the band by the

interval between bait consumption and plucking.

To identify putative food items, owners were asked what they fed their dog, what their dog ate

yesterday and what they saw other people's dogs eating. The latter question was to identify 'taboo'

items. Foods reported with >2% frequency were sampled, within households where possible.

Otherwise, livestock meats were sampled at a market in Guelendeng, wild animal meats were obtained

from hunters, and fish were sampled from fishers on the Chari. Samples were dried on the day of

collection and stored in ambient conditions. Items were heat sterilised before and after importation

under licence to the UK.

For each dog, one whisker was rinsed in distilled water, scraped to remove surface contaminants, and

dried for 24 hours. Whiskers were cut into ~0.7 mg sections and each was weighed and sealed in a tin

cup for analysis. Food samples were freeze-dried, homogenised and ~0.7 mg was weighed and sealed

in a tin cup. Stable isotope analyses of carbon (δ^{13}C) and nitrogen (δ^{15}N) were conducted using a

standard protocol [10] on a Sercon 2020 elemental analyser isotope ratio mass spectrometer. We

applied a lipid-normalisation model to δ^{13}C values of samples with high lipid content [11,12].

We estimated the relative contributions of putative food sources to dog diets using the SIMMR

package (v.3.0) [13]. Local food source values were used where available. Isotope ratios were

averaged for each dog. Trophic discrimination factors for dogs for δ^{15}N (2.75‰ SD 1.19) and δ^{13}C

(1.64‰ SD 1.43) were obtained using the SIDER package [14]. Concentration dependence values

(mean N/C) were added to the model as dogs have omnivorous diets with high variation in elemental

composition among sources [15].

Guinea worm infection

To explore correlates of infection, we analysed variation in a) whether a dog had ever been reported to

have emergent Guinea worms or not, using a GLM with a binomial error structure, and b) the number

of worms that had emerged from infected dogs, using a GLM with a negative binomial error structure.

Explanatory variables were settlement, age, sex, body condition, household water provision and the

proportion of fish in the diet estimated from stable isotope analysis. The number of GPS fixes

recorded in the vicinity of water bodies, number of different water bodies visited and total range sizes

were also included in model a). To maximise sample size, a staged analysis was conducted; the model

was run first using data for individuals with no missing data and then run again having removed the

variables that appeared in fewer than 50% of the top models, allowing the inclusion of further

individuals with some missing data. The retention of diet data meant these analyses were limited to

Magrao and Kakale. Settlement was always retained in the model to control for underling spatial

variation in risk. Results are expressed as odds of having had Guinea worm and as relative risks, with

95% confidence intervals from bootstrapping 10,000 times with replacement. The same analyses were

conducted on our field survey records of infection and the longer-term data collected by the CGWEP.

For the models of CGWEP data, some (<5%) bootstrapped models did not converge and were

excluded from calculations of confidence intervals.

Model selection adopted an information theoretic approach. A difference in Akaike Information

Criterion (ΔAIC) of <2 was used to select the top model set. Unless otherwise stated, full model-

averaged effect size coefficients with 95% confidence intervals and p-values from full model-averaged

tables are reported. Correlations between explanatory variables were investigated prior to analyses and

correlated variables were precluded from appearing in the same models. Analyses were undertaken in

R (v.3.3.3) and Quantum Geographic Information System (v2.18.1). Locations were projected into the

EPSG (32634) coordinate system. lme4 (v1.1-12) [16] was used to conduct GLMMs, MuMIn

(v1.15.6) [17] for model selection and mgcv (v1.8.12) [18] for GAMs.

Results

Dog tracking

150 dogs were collared and 134 generated usable tracking data (Tables 1-2). Mean deployment was

7.7 days (SE=0.2). Dog ranging was significantly affected by body condition, settlement and the

number of days for which a dog was monitored (Table 3); age and sex appeared in the top model sets

but did not contribute significantly to variation after model averaging. Dogs in poor body condition

had smaller total ranges (Mean = $10.50 \text{ km}^2 \pm$ SE Mean 2.29 km^2) than those in moderate condition

($11.74 \pm 1.91 \text{ km}^2$; Effect size for logged range (95 % CIs) = -0.45 (-0.89, 0.00), p<0.05). The core

and total ranges of dogs in Largana and Magrao were significantly smaller than those of dogs living in

Kakale (Table 2; Largana vs Kakale: Effect size for logged core range: -0.89 (-1.69, -0.10), p<0.05;

total range: -1.52 (-2.08, -0.95), p<0.001; Magrao vs Kakale: Effect size for logged core range: -2.00

(-2.71, -1.30), p<0.001; Effect size for logged total range: -1.47 (-1.97, -0.96), p<0.001). Use in

hunting activity and household provision of water did not appear in any of the top models for variation

in range areas (Table 3).

Using satellite imagery from earlier in the dry season, 342 ponds were identified in the three search

areas. 30 ponds were located using GPS in the field, of which 17 were not detected using satellite

imagery, suggesting that new ponds were forming as the rains commenced. A sample of 359 ponds

was used in our analyses. On average, dogs had 23% of their locations within 100 m of a detected

pond and 2% of their relocation points within 100 m of the Chari River. Dogs in poor body condition

spent more time in the vicinity of ponds than those in moderate condition (Effect size in negative

binomial model = 0.72 (0.28, 1.15), p<0.01; n=110). Water provision for animals was associated with

dogs spending less time in the vicinity of ponds (-1.01 (-1.61, -0.42), p<0.001). Dogs from Kakale and

Magrao spent more time in the vicinity of ponds than dogs from Largana. In Magrao, 37 of 101 (37%)

ponds were visited by at least one dog, while in Kakale, 87 of 202 (43%) ponds were visited. A small

minority of ponds accounted for the great majority of dog activity. For example, in Kakale, 80% of

dog visits occurred at nine (5%) ponds, all of which were <100 m from a household with a tracked dog

and 95% of visits occurred at 17 (8%) ponds, 16 of which were <200 m from a household (Figure 2).

Dog diet

From 92 households, we recorded 346 reports of food items, grouped into seven broad, isotopically-

distinct categories for which we collected 174 samples: C3 plant food (e.g. potatoes, peanuts, rice,

n=9), C4 plant food (e.g. millet, sorghum, maize, n=17), livestock meat (n=25, including samples of 7

cows, 7 sheep, 9 goats, 2 chickens), wild animal meat (n=24, including samples of 17 amphibians, 2

reptiles, 2 mammals, 3 birds), fish (n=88) and human faeces (n=11). Whiskers were obtained from 108

dogs from Kakale and Magrao. Mean whisker length was 50.6 mm (SD 9.1), mean growth rate was

0.42 mm/day (SD 0.21, n=21), hence the period represented by the average whisker was ~120 days.

C4 plant food ("boule", a sorghum or millet porridge), human faeces and C3 plant food comprised the

majority of dog diet and other items were eaten relatively infrequently, but in similar measure (Figure

3).

Guinea worm infection

Infection data were collected for 149 dogs, of which 34 (23%) had a history of Guinea worm

emergence. A model including all explanatory variables was initially run (n=86), however body

condition, age, sex or any spatial variables appeared in fewer than 50% of the top models and so dogs

with missing data for these variables were included in the final model. Using our field data, 13 of 51

(25.5%) dogs in Magrao and 14 of 49 (28.6%) in Kakale had some history of Guinea worm infection.

Dogs in households where water was provided for animals were less likely to have had Guinea worm

(Figure 4; Table 3; Relative risk = 0.33 (0.11, 0.63) p<0.01), whereas dogs with a higher proportion of

fish in their diets were more likely to have had Guinea worm (Figure 4; Table 3; Relative risk = 1.14

(1.00, 1·48), $p < 0.05$; Figure 4). CGWEP surveillance data recorded that 15 of 52 (28.8% dogs in

Magrao and 8 of 50 (16.0%) dogs in Kakale had some history of worm infection. Analyses using

CGWEP data suggested an effect of settlement (Increased risk in Magrao, compared to Kakale,

Relative risk = 2.17 (0.19, 11.44), a similar effect of water provision (Relative risk = 0.40 (0.06, 0.71)

but no effect of fish consumption (Tables S1-3).

The median number of emergent worms per infected dog was 2. Two dogs with exceptionally high

worm burdens (9 and 14 worms) had particular influence on analyses and when they were excluded,

none of the explanatory variables had an effect on the number of worms that had emerged from an

infected dog.

Discussion

Human activity is a key influence upon domestic dog ecology in these settlements and relates to

variation in ranging, diet, access to water and, critically, history of Guinea worm infection. Dogs spent

most of their time close to their human households and their core ranges tended to be less than 1 km^2.

The exception was for dogs in Kakale, whose owners were moving frequently between permanent and

seasonal residences for crop cultivation.

Dog exposure to ponds and their infection history were functions of the provision of water to animals

by the household. We have shown that household provision of water to animals was protective of

Guinea worm infection, and that water provision was negatively related to time spent close to ponds.

There is a question of cause and effect here; the availability of water within the household might

reduce the tendency for dogs to visit ponds but it is also likely that households located further from

standing water might more frequently provide water for their animals. We did not determine the

source of water provided for animals and analysis of whether water comes from safe, borehole sources

or unsafe surface water sources may help refine our understanding of risk. Clearly, however, our

findings highlight the importance of variation in where and how dogs access water in determining

their exposure to Guinea worm infection.

In terms of potential sources of infection, a small number of ponds close to households accounted for

almost all exposure of dogs to surface water, while visits to the Chari River were comparatively rare.

Dog exposure to ponds close to settlements means that such ponds merit targeted investigation and

prioritisation for systematic treatment with temephos (Abate). Dogs acquiring infection by drinking

water contaminated with infected copepods would fit with a classical route for Guinea worm

transmission. The inconsistency of this transmission pathway with very low incidence in humans can

be explained by the relative ease of access to borehole water. Our analyses of satellite imagery reveal

problems in accessing contemporary imaging at an adequate resolution and in reliably identifying

ponds using optical images. Therefore, we suggest that additional techniques, such as radar, aerial and

ground-based surveys, may assist in understanding the distribution of surface water in and around

villages badly affected by Guinea worm.

Dogs relied heavily on anthropogenic foods, principally boule and human faeces. Given the similarity

in the stable isotope ratios of these items, there is uncertainty in the exact contribution they each make,

though together they clearly make up the majority of dog diet [19]. Although we were successful in

determining between-dog variation in fish consumption, we could not distinguish amphibians from

other wild animal meats in these settlements. We had strong *a priori* interest in consumption of

aquatic foods, as a potential route for exposure of dogs to infection. We identified a significant effect

of fish consumption on Guinea worm infection. There was considerable uncertainty associated with

this effect, though over the range of observed values the risk increased markedly. This may highlight

the importance of novel routes for exposure of dogs to infection, perhaps by the consumption of fish,

fish guts or other remains.

It is, of course, possible that dogs are exposed to Guinea worm infection by both classical and novel

routes, or perhaps that novel routes explain occasional introductions and the classical route

"amplifies" infection in non-human hosts, in a similar manner as was once seen with humans.

We were able to conduct exploratory epidemiological analyses using history of adult female Guinea

worm emergence as an indicator of infection. The underlying assumption of these analyses is that

contemporary investigations reflect the ecology of the dog at the time at which it acquired Guinea

worm infection. This may not be unreasonable, given dog longevity and consistency of their

environment. However, our study, undertaken at the beginning of the rainy season, missed the major

environmental variation arising from rains and flooding. We also took emergent adult worms as the

only measure of infection, which omits dogs that are truly infected but have not yet progressed to

worm emergence (i.e. some animals classified as uninfected were genuinely infected) and dogs that

have been exposed (i.e. exhibit the ecological traits of infected animals) but have not progressed to

infection. An effective diagnostic of prepatent infection in dogs could potentially reduce this

classification uncertainty.

It is perhaps remarkable that we have, with even this relatively small-scale study, identified correlates

of Guinea worm infection in these new non-human animal hosts. We hope our work provides

encouragement that the impressive progress towards dracunculiasis eradication might continue apace,

by addressing this new but now demonstrably tractable epidemiological setting in which the Guinea

worm has emerged.

Acknowledgements

Financial and logistic support was provided by The Carter Center (cartercenter.org), Chad Ministry of

Public Health (sante-tchad.org) and World Health Organization (who.int). The Bill and Melinda Gates

Foundation (gatesfoundation.org) provided satellite imagery. With the exception of the named authors

affiliated with the funding organisations, the funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. We thank Stuart Bearhop, Io Blair-

Freese, Chris Cleveland, Melinda Denson, Mark Eberhard, Donald Hopkins, Rich Inger, Aileen Mill,

Ken Neal, Philip Tchindebet Ouakou, Sharon Roy, Ernesto Ruiz-Tiben, Jordan Schermerhorn, Adam

Weiss, Michael Yabsley, Hubert Zirimwabagabo and colleagues at University of Exeter for their

advice and support.

References

1. Watts S. Dracunculiasis in Africa in 1986: its geographic extent, incidence, and at-risk population.

Am J Trop Med Hyg 1986; **37**: 119–25.

2. Centers for Disease Control. Guinea Worm Wrap Up No. 259 February 28[th] 2019.

https://www.cartercenter.org/resources/pdfs/news/health_publications/guinea_worm/wrap-

up/259.pdf

3. Thiele EA, Eberhard ML, Cotton JA, Durrant C, Berg J, Hamm K and Ruiz-Tiben E. Population

genetic analysis of Chadian Guinea worms reveals that human and non-human hosts share common

parasite populations. *PLoS Negl Trop Dis* 2018; **12**: p.e0006747.

4. Centers for Disease Control. Renewed transmission of dracunculiasis. Chad, 2010. *Morbidity and*

Mortality Weekly Reports 2011; **60**: 744-8.

5. International Commission for the Certification of Dracunculiasis Eradication. Report of 12[th]

Meeting 2018; World Health Organization, Geneva.

6. Roberts L. Battle to wipe out debilitating Guinea worm parasite hits 10 year delay. *Nature* 2019;

574: 157–58.

7. Eberhard ML, Ruiz-Tiben E, Hopkins DR, *et al.* The peculiar epidemiology of dracunculiasis in

Chad. *Am J Trop Med Hyg* 2014; **90**: 61–70.

8. Molyneux D, Sankara DP. Guinea worm eradication: Progress and challenges— should we beware

of the dog? *PLoS Negl Trop Dis* 2017; **11**: 4–7.

9. Laflamme DP. Development and Validation of a Body Condition Score System for Dogs. *Canine*

Pract 1997; **22**: 10–5.

10. Inger R, Ruxton GD, Newton J, *et al.* Using daily ration models and stable isotope analysis to

predict biomass depletion by herbivores. *J Appl Ecol* 2006; **43**: 1022–30.

11. McConnaughey T, McRoy CP. Food-Web structure and the fractionation of Carbon isotopes in the

Bering Sea. *Mar Biol* 1979; **53**: 257–62.

12. Kiljunen M, Grey J, Sinisalo T, Harrod C, Immonen H, Jones RI. A revised model for lipid-

normalizing d13C values from aquatic organisms, with implications for isotope mixing models. *J*

Appl Ecol 2006; **43**: 1213–22.

13. Parnell A, Inger R. simmr: A stable isotope mixing model. R package. Version 0.3. 2016.

14. Healy K, Guillerme T, Kelly SBA, Inger R, Bearhop S, Jackson AL. SIDER: an R package for

predicting trophic discrimination factors of consumers based on their ecology and phylogenetic

relatedness. *Ecography* 2017; **41**: 1233-1410.

15. Phillips DL, Koch PL. Incorporating concentration dependence in stable isotope mixing models.

Oecologia 2002; **130**: 114–25.

16. Bates DM, Machler M, Bolker BM, Walker SC. Fitting Linear Mixed-Effects Models using lme4.

J Stat Softw 2014; **67**. DOI:10.18637/jss.v067.i01.

17. Barton K. MuMIn: Multi-Model Inference. R package v. 1.15.6. 2016. https://cran.r-

project.org/package=MuMIn.

18. Wood S. mgcv: GAMs and generalized ridge regression for R. *R news* 2001; **1**: 20–5.

19. Butler JRA, Brown, WY, Du Toit JT. The value and supply of human faeces in the diet of free-

ranging dogs. *Animals* 2018; **8**: 67.

Table 1. **Summary of dog characteristics in three settlements in Chad.** Data are from all collared

dogs (n=150), but some dogs have missing data for individual characteristics. Water provision is the

proportion of households which reported providing water for animals.

Settlement	n dogs	Sex (M:F)	Body condition (Poor:Moderate)	Mean age (months)	Age range (months)	Water provision (%)
Largana	34	16 : 18	22 : 12	32 ± 4	6 - 96	60
Kakale	52	26 : 26	25 : 27	35 ± 3	4 - 144	75
Magrao	64	40 : 24	16 : 42	34 ± 3	5 - 144	100
All	150	82 : 68	63 : 81	34 ± 2	4 - 144	82

360 **Table 2**. **Summary of dog ranging behaviour in three settlements in Chad.** Core range is the 60%

361 kernel density estimate and total range is the 100% minimum convex polygon. The mean and standard

362 error in parentheses are presented for each variable.

Settlement	n dogs	Core range (km^2)	Total range (km^2)	Fixes within 100m of a pond (%)	Unique ponds visited	Ponds in core range	Ponds in total range	Fixes within 100m of the Chari River (%)
Largana	31	0.30(0.09)	4.70(1.33)	11(4)	2(0.3)	2(0.3)	4(0.8)	4(2)
Kakale	48	2.53(0.74)	21.01(3.08)	36(5)	8(0.6)	8(1.3)	25(3.1)	<1(<1)
Magrao	55	0.17(0.04)	4.39(0.63)	18(4)	3(0.3)	1(0.2)	6(0.7)	2(1)
All	134	1.05(0.28)	10.42(1.35)	23(3)	4(0.3)	3(0.6)	12(1.4)	2(<1)

Table 3. Summaries of analyses of variation in dog range sizes, time spent by dogs in the vicinity of

ponds and history of Guinea worm infection in dogs in rural Chad. Core range is the 60% kernel

density estimate and total range is the 100% minimum convex polygon. History of Guinea worm

infection is from our field survey records and records held by the Chad Guinea Worm Eradication

Program. Model outcomes are shown as the top model sets (ΔAICc < 2 from the top model) for all

analyses.

Formula	df	LogLik	AICc	ΔAICc	Weight
Total range size (100 % MCP)					
Days monitored + body condition + settlement	6	-165.34	343.5	0	0.54
Days monitored + body condition + settlement + age	7	-165.01	345.1	1.62	0.24
Days monitored + body condition + settlement + sex	7	-165.06	345.2	1.71	0.23
Core range size (60 % KDE)					
Settlement	4	-204.48	417.3	0	0.33
Settlement + days monitored	5	-203.80	418.2	0.85	0.22
Settlement + body condition	5	-203.91	418.4	1.07	0.19
Settlement + days monitored + body condition	6	-203.17	419.1	1.81	0.13
Settlement + water provision	5	-204.31	419.2	1.85	0.13
Time spent in the vicinity of ponds (n fixes within 100 m of ponds)					
Body condition + settlement + water provision + age	7	-1357.02	2729.0	0	0.54
Body condition + settlement + water provision	6	-1358.29	2729.3	0.29	0.46
Guinea worm infection (reported history of infection or not)					
Our field records (main analysis)					
% fish consumed + water provision	3	-52.34	110.9	0	0.52
% fish consumed + water provision + settlement	4	-52.18	112.8	1.85	0.21
Longer-term CGWEP records (for validation)					
Water provision + settlement	3	-49.89	106.0	0	0.54
Water provision + settlement + % fish consumed	4	-49.70	107.8	1.78	0.22

371
Figure 1: A) Locations of dog study sites on the Chari River, Chad. B) Adult female Guinea

worm *Dracunculus medinensis* being extracted from a domestic dog. C) Typical dog fitted with

GPS tracking collar. The settlements were: Magrao (10°59'44.31"N, 15°29'29.27"E) which

encompasses the associated permanent village of Sawata, Largana (10°45'15.26"N, 16° 1'39.86"E) and

the associated permanent village of Gomba (10°44'46.22"N, 16° 2'46.70"E), and Kakale-Mberi

(10°53'0.79"N, 15°38'8.45"E) and the associated seasonal settlement of Awine (10°48'6.34"N,

15°37'56.61"E). Guelendeng is the major town in the sous-préfecture. Photographs Jared Wilson-

Aggarwal. The satellite image was generated using the Esri world imagery basemap (sources: Esri,

DigitalGlobe, GeoEye, i-cubed, USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP,

swisstopo, and the GIS User Community).

383
Figure 2. Distances from human households of the ponds most exposed to visits by dogs in two

settlements in Chad. A is Kakale and B is Magrao. Dog visits were determined by locating dogs

within 100m of a pond by GPS tracking. Ponds were located from satellite imagery within a search

area enclosing all dog location points for each village and are a sub-sample of all available ponds. All

visits by all dogs were summed for each pond and the cumulative total of all dog visits is plotted

against the distance of the pond to the nearest household with tracked dogs, with the red line from a

fitted generalised additive model. The dotted lines indicate the distance from a household with dogs at

which 80%, 90% and 95% of all dog visits are captured.

Figure 3. Stable isotope analysis of the diets of dogs in two settlements in Chad. Settlements are

A) Kakale and B) Magrao. i) The δ^{15}N and δ^{13}C values for sampled dogs (black dots) and the mean ±

standard deviation of δ^{15}N and δ^{13}C for putative food groups. Trophic discrimination factors, derived

from package SIDER, have been applied to adjust dog isotope ratios downwards for both δ^{15}N and

δ^{13}C. ii) Outputs of Bayesian mixing models, from package SIMMR, of the proportional contributions

of the main food groups to dog diets.

A   **B**

Figure 4. Effects of water provision and of fish consumption on Guinea worm infection history

in dogs in Chad. The odds, with 95% confidence intervals, of having had Guinea worm are shown for

dogs A) living in households in which animals are, and are not, provided with water, and B) with

different modelled proportions of fish in their diets. The effects of fish consumption are presented

separately for dogs living in households, in which animals are (dark grey) and are not (light grey)

provided with water. Odds are from exponentiated model coefficients describing the relationship

between the history of Guinea worm infection from our field survey, averaged across the top model set

and for the different levels of values of predictive factors. Results presented are from dogs in Kakale

but dogs in Magrao show the same relationships.