Ecology of domestic dogs *Canis familiaris* as an emerging reservoir of Guinea worm *Dracunculus*

medinensis infection

Robbie A. McDonald[1]*, Jared K. Wilson-Aggarwal[1], George J.F. Swan[1], Cecily E.D. Goodwin[1],

Tchonfienet Moundai[2], Dieudonné Sankara[3], Gautam Biswas[3], James A. Zingeser[4]

[1]Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, UK.

[2]Guinea Worm Eradication Programme, Ministry of Public Health, N'Djamena, Republic of Chad.

[3]World Health Organisation, Geneva, Switzerland.

[4]The Carter Center, Atlanta, GA, USA.

Email addresses: Robbie McDonald r.mcdonald@exeter.ac.uk, Jared Wilson-Aggarwal

jw695@exeter.ac.uk, George Swan g.swan@exeter.ac.uk, Cecily Goodwin c.goodwin@exeter.ac.uk,

Tchonfienet Moundai tchomcalvin@yahoo.fr, Dieudonné Sankara sankarad@who.int, Gautam Biswas

biswasg@who.int, James Zingeser Jim.Zingeser@cartercenter.org.

***Correspondence author:** Robbie McDonald, Environment and Sustainability Institute, University of

Exeter, Penryn Campus, Penryn, UK. email: r.mcdonald@exeter.ac.uk. Telephone 0798 9069834.

Short title: Guinea worm in domestic dogs

Keywords: Neglected Tropical Diseases, zoonoses, domestic dogs, Guinea worm, disease eradication,

One Health, epidemiology, disease ecology, dracunculiasis

Statement of authorship: RM, JZ, GB and DS conceived the study. RM, JW-A, GS, TM and JZ

conducted the fieldwork. JW-A, CG and GS analysed samples and data. RM wrote the manuscript.

JW-A, CG, GS, DS and JZ revised the manuscript.

Data accessibility statement: Should this manuscript be accepted for publication, all data will be

available at Dryad data repository.

Abstract

Global eradication of human Guinea worm disease has been set back by the emergence of infections in

animals, particularly domestic dogs. The ecology and epidemiology of this reservoir is unknown. We

tracked dogs using GPS, inferred diets using stable isotope analysis and analysed correlates of

infection in Chad, where Guinea worm prevalence is highest. Dogs had small ranges that varied

markedly among villages. Diets consisted largely of human staples and human faeces. A minority of

ponds, mostly <200m from households, accounted for most dog exposure to potentially unsafe water.

Risk of infection was reduced in households providing water for animals but increased with fish

consumption. Provision of safe water might reduce dog exposure to unsafe water while proactive

temephos application to the small number of ponds to which dogs have most access is recommended.

Fish might have an additional role as transport hosts, by concentrating copepods infected with worm

larvae.

Author summary

Guinea worm is a parasite that causes profoundly debilitating disease in humans. An eradication

program has been successful in nearly eliminating the disease from people. However, the same worm

has now been found in domestic dogs and the frequency of detecting Guinea worm in dogs has been

increasing. This means that to eradicate Guinea worm, the infection must be eliminated in dogs as well

as in people. However, not much is known about the disease in dogs. This study is the first to study

dogs in relation to Guinea worm infection. We worked in the worst-affected country, Chad. We

attached GPS collars to dog to track their ranging and use of water bodies and analysed their diets

using a forensic technique, based on analysing stable isotope compostion of their whiskers and

potential food items. We showed that dogs living in households that provided water to their animals

had a lower risk of having had Guinea worm and that dogs that ate more fish had an increased risk.

These findings suggest there is a classical route for worm transmission in dogs, via drinking

contaminated water, as well as a novel route, potentially by eating fish carrying a source of infection.

Introduction

Guinea worm *Dracunculus medinensis* is a nematode parasite that causes Guinea worm disease (dracunculiasis) in humans. It was once widespread in Asia and Africa (Watts 1986) but an eradication campaign has made impressive progress in reducing human cases from 3.5 million per annum in 21 countries in 1986, to only 28 in 2018, in Chad (17), South Sudan (10), and Angola (1) (Centers for Disease Control 2019). Alongside the near-absence of human cases, however, infections were detected in 1040 domestic dogs *Canis familiaris* and 25 domestic cats *Felis catus* in Chad, 11 dogs, five cats and one olive baboon *Papio anubis* in Ethiopia, and 18 dogs and two cats in Mali in 2018 (CDC 2019) Adult Guinea worms emerging from humans and these non-human animals are genetically indistinguishable (Thiele *et al.* 2018).

Prior to re-emergence in 2010, no human cases of Guinea worm were reported in Chad for ten years (CDC 2011), suggesting an unknown reservoir of infection in non-human animals and/or undetected humans. Given the current high number of dogs infected relative to human cases and improved human surveillance in Chad, it is clear that non-human animals, particularly dogs, now constitute a contemporary reservoir of infection. Little is known about the epidemiology of Guinea worm in its non-human hosts. Although direct transmission between dogs and humans is not possible, dogs clearly maintain a persistent source of potential infection for humans in their shared environments. Thus, to eradicate Guinea worm, transmission of infection must be interrupted in non-human as well as human hosts (International Commission for the Certification of Dracunculiasis Eradication 2018). Alongside civil insecurity in the remaining endemic areas, non-human animal infections are now the major impediment to Guinea worm eradication (Roberts 2019).

The pathways by which dogs acquire infection remain unknown, though two main possibilities exist. First, similar to the classical route for humans, dogs may acquire infection from drinking water containing infected copepods as the intermediate hosts. Second, dogs might consume a paratenic or transport host, most likely a fish or amphibian, that has itself eaten infected copepods (Eberhard *et al*. 2014). In Chad, dogs drink from permanent and transient surface water bodies that often contain large

copepod populations. Large-scale human exploitation of fish and frogs also provides the potential for

dog consumption of their uncooked remains (Eberhard *et al*. 2014).

Options for interrupting dog infections are broadly similar to those for humans and include: detection

and containment of hosts, treating water bodies with temephos, and limiting exposure by reducing

consumption of contaminated water and food (Molyneux & Sankara 2017). Since different approaches

are required to implement such interventions in free-ranging dogs, it is important that risk factors are

identified, to help inform management. To understand the epidemiology and control of Guinea worm

in dogs, we undertook a study of their ecology in Chad, to characterise their ranging and access to

water, as well as their diet and consumption of aquatic foods. We show that both a classical route of

transmission, relating to exposure to potentially unsafe water, as well as a novel route, via

consumption of fish, potentially as a paratenic host, influence the risk of dog infection and highlight

management implications for proactive water treatment.

Material and methods

Ethics statement

This study was approved by the University of Exeter College of Life and Environmental Sciences

(Penryn Campus) Ethics Committee (2016/1488). The Committee and the project ahered to the

"Guidelines for the treatment of animals in behavioural research and teaching" of the Association for

the Study of Animal Behaviour.

Study area and subject recruitment

Fieldwork was conducted between 24 June and 12 July 2016 in three settlements (Kakale, Magrao and

Lagarna), each comprised of multiple villages, situated in Guelendeng sous-prefecture, close to the

Chari River, Chad (Figure 1). Settlements were selected because of high incidence of infection in dogs

and relative ease of access. The chief of each village provided approval and dog owners individually

provided consent. For each household, we recorded GPS location and conducted a questionnaire,

recording whether dogs were taken hunting and whether water was provided for animals within the

household. For each dog, we recorded sex, age in months, body condition (on a scale of 1 = emaciated

to 9 = severely obese (Laflamme 1997) and then classified as poor ≤2 or moderate ≥3), and whether

the dog had history of emergent adult Guinea worms. The Chad Guinea Worm Eradication

Programme (CGWEP) provided additional surveillance data identifying dogs with confirmed

infections.

Dog tracking

All dogs, apart from pups, were fitted with adjustable nylon collars (Ancol, Walsall, UK) equipped

with an iGotU GT600 GPS logger (Mobile Action Technology, Taipei) with a 10 minute fix interval.

After deployment for up to 14 days, collars were retrieved and data downloaded. Locations before

6am the day after deployment and after 6am the day before retrieval were excluded to avoid

interference arising from our activities. Fixes that suggested speeds greater than 20km/hr were

considered spurious and excluded. In one village (Gomba) in Largana, owners removed collars early,

so locations after midnight prior to their removal of collars were excluded. The dogs' core and total

ranges were characterised by 60% kernel density estimates and 100% minimum convex polygons,

respectively.

DigitalGlobe satellite imagery was obtained for February, October and December 2012 for Magrao,

Kakale and Largana, respectively. We sampled the distribution of ponds by drawing a box around all

dog locations for each settlement and visually searching images for ponds. For each dog, we

determined the number of visits (locations within 100m) to a pond and to the Chari River, the number

of ponds in core and total ranges and the number of ponds visited. For each pond, we measured the

number of dog visits, the number of dogs that visited and the number of visits by each dog (defined by

an interval of 30 minutes between fixes). To characterise the ponds most visited by dogs, ponds were

ranked by number of visits by all dogs and the distance of each pond to the nearest household with

tracked dogs.

Variation in dog range sizes was analysed using linear mixed models (LMMs). Ranges were log-

transformed. Models contained settlement, sex, age, body condition, hunting activity and water

provision as explanatory variables, and the number of tracking days as a fixed effect to control for

effort. Variation in the time dogs spent close to ponds (the number of fixes within 100m of a pond)

was analysed using a general linear model (GLM) with the same explanatory variables but with

tracking days as an offset to control for effort. A generalised additive model (GAM) related the

cumulative total of dog visits per pond to the log-transformed distance from a household with tracked

dogs.

Dog diets

Composition of dog diets was determined using stable isotope analysis of dog whiskers and putative

food items. Two whiskers were plucked during collar retrieval. The period over which whiskers had

grown, and diet could be inferred, was calculated by dividing whisker length by growth rate; a

subsample of dogs were fed ~50g baits with 0.8g (40-80mg/kg) of rhodamine B, a food colourant that

stained whiskers with a fluorescent band. Growth rate (mm/day) was calculated by dividing the

distance between the whisker base and the distal end of the band by the interval between bait

consumption and plucking.

To identify putative food items, owners were asked what they fed their dog, what their dog ate

yesterday and what they saw other people's dogs eating. The latter question was to identify 'taboo'

items. Foods reported with >2% frequency were sampled, within households where possible.

Otherwise, livestock meats were sampled at a market in Guelendeng, wild animal meats were obtained

from hunters, and fish were sampled from fishers on the Chari. Samples were dried on the day of

collection and stored in ambient conditions. Items were heat sterilised before and after importation

under licence to the UK.

For each dog, one whisker was rinsed in distilled water, scraped to remove surface contaminants, and

dried for 24 hours. Whiskers were cut into ~0.7mg sections and each was weighed and sealed in a tin

cup for analysis. Food samples were freeze-dried, homogenised and ~0·7mg was weighed and sealed

in a tin cup. Stable isotope analyses of carbon (δ^{13}C) and nitrogen (δ^{15}N) were conducted using a

standard protocol (Inger *et al.* 2006) on a Sercon 2020 elemental analyser isotope ratio mass

spectrometer. We applied a lipid-normalisation model to δ^{13}C values of samples with high lipid

content (McConnaughey & McRoy 1979; Kiljunen et al. 2006).

We estimated the relative contributions of putative food sources to dog diets using the SIMMR

package (v.3.0) (Parnell & Inger 2016). Local food source values were used where available. Isotope

ratios were averaged for each dog. Trophic discrimination factors for dogs for δ^{15}N (2.75‰ SD 1.19)

and δ^{13}C (1.64‰ SD 1.43) were obtained using the SIDER package (Healy *et al*. 2017). Concentration

dependence values (mean N/C) were added to the model as dogs have omnivorous diets with high

variation in elemental composition among sources (Phillips & Koch 2002).

Guinea worm infection

To explore correlates of infection, we analysed variation in a) whether a dog had ever been reported to

have emergent Guinea worms or not, using a GLM with a binomial error structure, and b) the number

of worms that had emerged from infected dogs, using a GLM with a negative binomial error structure.

Explanatory variables were settlement, age, sex, body condition, household water provision and the

proportion of fish in the diet. The proportion of time spent at ponds, number of unique ponds visited,

total and core range sizes and hunting activity were also included in model a). To maximise sample

size, a staged analysis was conducted; the model was run first using data for individuals with no

missing data and then run again having removed the variables that did not explain a significant amount

of variation in the response, allowing the inclusion of further individuals with some missing data.

Settlement was always retained in the model to control for underling spatial variation in risk. Results

are expressed as odds of having had Guinea worm and as relative risks, with 95% confidence intervals

from bootstrapping 10,000 times with replacement. The same analyses were conducted on our field

survey records of infection and the longer-term data collected by the CGWEP.

Model selection adopted an information theoretic approach. A difference in Akaike Information

Criterion (ΔAIC) of <2 was used to select the top model set. Unless otherwise stated, full model-

averaged effect size coefficients and 95% confidence intervals are reported. Correlations between

explanatory variables were investigated prior to analyses and correlated variables were precluded from

appearing in the same models. Analyses were undertaken in R (v.3.3.3) and Quantum Geographic

Information System (v2.18.1). Locations were projected into the EPSG (32634) coordinate system.

lme4 (v1.1-12; Bates *et al*. 2014) was used to conduct GLMMs, MuMIn (v1.15.6; Barton 2016) for

model selection and mgcv (v1.8.12; Wood 2001) for GAMs.

Results

Dog tracking

150 dogs were collared and 134 generated usable tracking data (Tables 1-2). Mean deployment was 7.7 days (SE=0.2). Dogs in better body condition had larger total ranges (Effect size (95 % CIs) = -0.45 (-0.89, 0.0), p<0.05). Dogs in Kakale had larger core and total ranges than in Lagarna (core range: -0.89 (-1.69, -0.10), p<0.05; total range: -1.52 (-2.08, -0.95), p<0.001) and Magrao (core range: -2.00 (-2.71, -1.30), p<0.001; total range: -1.47 (-1.97, -0.96), p<0.001). Hunting activity and the provision of water did not appear in any of the top models for range areas (Supplementary Information Table S1).

Using satellite imagery, 359 ponds were identified in the three search areas, though of 30 ponds located using GPS in the field, only 13 were detected using satellite images. On average, dogs had 23% of their locations within 100m of a detected pond and 2% of their relocation points within 100m of the Chari River. Dogs in poor body condition spent more time close to ponds than those in moderate condition (0.72 (0.28, 1.15), p<0.01; n=110). Water provision for animals was associated with dogs spending less time close to ponds (-1.01 (-1.61, -0.42), p<0.05). Dogs from Kakale and Magrao spent more time at ponds than dogs from Lagarna. In Magrao, 37 of 101 (37%) ponds were visited by at least one dog, while in Kakale, 87 of 202 (43%) ponds were visited. A small minority of ponds accounted for the great majority of dog activity. For example, in Kakale, 80% of dog visits occurred at nine (5%) ponds, all of which were <100m from a household with a tracked dog and 95% of visits occurred at 17 (8%) ponds, 16 of which were <200m from a household (Figure 2).

Dog diet

From 92 households, we recorded 346 reports of food items, grouped into seven broad, isotopically-distinct categories for which we collected 174 samples: C3 plant food (e.g. potatoes, peanuts, rice, n=9), C4 plant food (millet, sorghum, maize, n=17), livestock meat (n=25), wild animal meat (including amphibians, n=24), fish (n=88) and human faeces (n=11). Whiskers were obtained from 108 dogs from Kakale and Magrao. Mean whisker length was 50·6mm (SD 9.1), mean growth rate was 0.42mm/day (SD 0.21, n=21), hence the period represented by the average whisker was ~120

days. C4 plant food ("boule", a sorghum or millet porridge), human faeces and C3 plant food

comprised the majority of dog diet and other items were eaten relatively infrequently, but in similar

measure (Figure 3).

Guinea worm infection

Infection data were collected for 149 dogs, of which 34 (23%) had a history of Guinea worm

emergence. A model including all explanatory variables was initially run (n=86), however neither

body condition, age, sex or any spatial variables explained significant variation in the response in a

conditional analysis and so dogs with missing data for these variables were included in the final model

(n=102). Dogs in households where water was provided for animals were less likely to have had

Guinea worm (Relative risk = 0.33 (0.11, 0.63) $p<0.01$; Figure 4), whereas dogs with a higher

proportion of fish in their diets were more likely to have had Guinea worm (Relative risk = 1.14 (1.00,

1·48), $p < 0.05$; Figure 4). Analysis using CGWEP surveillance data suggested an effect of settlement,

a similar effect of water provision but no effect of fish consumption (Tables S1-3). The median

number of emergent worms per infected dog was 2. Two dogs with exceptionally high worm burdens

(9 and 14 worms) had particular influence on analyses and when they were excluded, none of the

explanatory variables had an effect on the number of worms emerging from an infected dog.

Discussion

Human activity is a key influence upon domestic dog ecology in these settlements and relates to

variation in ranging, diet, access to water and, critically, history of Guinea worm infection. Dogs spent

most of their time close to their human households and their core ranges tended to be less than 1km^2.

The exception was for dogs in Kakale whose owners were moving frequently between permanent and

seasonal residences for crop cultivation.

Dog exposure to ponds and their infection history were functions of the provision of water to animals

by the household. We have shown that provision of water to animals was protective of Guinea worm

infection, and that water provision was negatively related to time spent close to ponds. There is a

question of cause and effect here; the availability of water within the household might reduce the

tendency for dogs to visit ponds but it is also possible that households located further from standing

water might more frequently provide water for their animals. We did not determine the source of water

provided for animals and analysis of whether water comes from safe, borehole sources or unsafe

sources may help refine our understanding of risk. Clearly, however, our findings highlight the

importance of variation in where and how dogs access water in determining their exposure to Guinea

worm infection.

In terms of potential sources of infection, a small number of ponds close to households accounted for

almost all exposure of dogs to surface water, while visits to the Chari River were comparatively rare.

Dog exposure to ponds close to settlements means that such ponds merit targeted investigation and

prioritisation for systematic treatment with larvicide. Dogs acquiring infection by drinking water

contaminated with infected copepods would fit with a classical route for Guinea worm transmission.

The inconsistency of this transmission pathway with very low incidence in humans can be explained

by the relative ease of access to borehole water. Our analyses of satellite imagery reveal problems in

accessing contemporary imaging at an adequate resolution and in reliably identifying ponds using

optical images. Therefore, we suggest that additional techniques, such as radar, aerial and ground-

based surveys, may assist in understanding the distribution of surface water in and around villages

badly affected by Guinea worm.

Dogs relied heavily on anthropogenic foods, principally boule and human faeces. Given the similarity

in the stable isotope ratios of these items, there is uncertainty in the exact contribution they each make,

though together they clearly make up the majority of dog diet (Butler et al 2018). Although we were

successful in determining between-dog variation in fish consumption, we could not distinguish

amphibians from other wild animal meats in these settlements. We had strong *a priori* interest in

consumption of aquatic foods, as a potential route for exposure of dogs to infection. We identified a

significant effect of fish consumption on Guinea worm infection. There was considerable uncertainty

associated with this effect, though over the range of observed values the risk increased markedly. This

may highlight the importance of novel routes for exposure of dogs to infection, perhaps by the

consumption of fish, fish guts or other remains.

It is, of course, possible that dogs are exposed to Guinea worm infection by both classical and novel

routes, or perhaps that novel routes explain occasional introductions and the classical route

"amplifies" infection in non-human hosts, in a similar manner as was once seen with humans.

We were able to conduct exploratory epidemiological analyses using history of adult female Guinea

worm emergence as an indicator of infection. The underlying assumption of these analyses is that

contemporary investigations reflect the ecology of the dog at the time at which it acquired Guinea

worm infection. This may not be unreasonable, given dog longevity and consistency of their

environment. However, our study, undertaken at the beginning of the rainy season, missed the major

environmental variation arising from rains and flooding. We also took emergent adult worms as the

only measure of infection, which misses dogs that are truly infected but have not yet progressed to

worm emergence (i.e. some animals classified as uninfected were genuinely infected) and misses some

animals that have been exposed (i.e. exhibit the ecological traits of infected animals) but had not

progressed to infection. An effective diagnostic of prepatent infection in dogs could reduce this

classification uncertainty.

It is perhaps remarkable that we have, with even this relatively small-scale study, identified correlates

of Guinea worm infection in new non-human animal hosts. We hope our work provides

encouragement that the impressive progress towards dracunculiasis eradication might continue apace,

by addressing the new but now demonstrably tractable epidemiological setting in which the Guinea

worm is now emerging.

Acknowledgements

Financial and logistic support was provided by The Carter Center (cartercenter.org), Chad Ministry of

Public Health (sante-tchad.org) and World Health Organization (who.int). The Bill and Melinda Gates

Foundation (gatesfoundation.org) provided satellite imagery. With the exception of the named authors

affiliated with the funding organisations, the funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. We thank Stuart Bearhop, Io Blair-

Freese, Chris Cleveland, Melinda Denson, Mark Eberhard, Donald Hopkins, Rich Inger, Aileen Mill,

Ken Neal, Philip Tchindebet Ouakou, Sharon Roy, Ernesto Ruiz-Tiben, Jordan Schermerhorn, Adam

Weiss, Michael Yabsley, Hubert Zirimwabagabo and colleagues at University of Exeter for their

advice and support.

References

Barton K. MuMIn: Multi-Model Inference. R package v. 1.15.6. 2016. https://cran.r-

project.org/package=MuMIn.

Bates DM, Machler M, Bolker BM, Walker SC. Fitting Linear Mixed-Effects Models using lme4. *J*

Stat Softw 2014; **67**. DOI:10.18637/jss.v067.i01.

Butler JRA, Brown, WY, Du Toit JT. The value and supply of human faeces in the diet of free-

ranging dogs. *Animals* 2018; **8**: 67.

Centers for Disease Control. Guinea Worm Wrap Up No. 259 February 28[th] 2019.

https://www.cartercenter.org/resources/pdfs/news/health_publications/guinea_worm/wrap-

up/259.pdf

Centers for Disease Control. Renewed transmission of dracunculiasis. Chad, 2010. *Morbidity and*

Mortality Weekly Reports 2011; **60**: 744-8.

Eberhard ML, Ruiz-Tiben E, Hopkins DR, *et al.* The peculiar epidemiology of dracunculiasis in Chad.

Am J Trop Med Hyg 2014; **90**: 61–70.

Healy K, Guillerme T, Kelly SBA, Inger R, Bearhop S, Jackson AL. SIDER: an R package for

predicting trophic discrimination factors of consumers based on their ecology and phylogenetic

relatedness. *Ecography* 2017; **41**: 1233-1410.

Inger R, Ruxton GD, Newton J, *et al.* Using daily ration models and stable isotope analysis to predict

biomass depletion by herbivores. *J Appl Ecol* 2006; **43**: 1022–30.

International Commission for the Certification of Dracunculiasis Eradication. Report of 12[th] Meeting

2018; World Health Organization, Geneva.

Kiljunen M, Grey J, Sinisalo T, Harrod C, Immonen H, Jones RI. A revised model for lipid-

normalizing d13C values from aquatic organisms, with implications for isotope mixing models. *J*

Appl Ecol 2006; **43**: 1213–22.

Laflamme DP. Development and Validation of a Body Condition Score System for Dogs. *Canine*

Pract 1997; **22**: 10–5.

McConnaughey T, McRoy CP. Food-Web structure and the fractionation of Carbon isotopes in the

Bering Sea. *Mar Biol* 1979; **53**: 257–62.

Molyneux D, Sankara DP. Guinea worm eradication: Progress and challenges— should we beware of

the dog? *PLoS Negl Trop Dis* 2017; **11**: 4–7.

Parnell A, Inger R. simmr: A stable isotope mixing model. R package. Version 0.3. 2016.

Phillips DL, Koch PL. Incorporating concentration dependence in stable isotope mixing models.

Oecologia 2002; **130**: 114–25.

Roberts L. Battle to wipe out debilitating Guinea worm parasite hits 10 year delay. *Nature* 2019; **574**:

157–58.

Thiele EA, Eberhard ML, Cotton JA, Durrant C, Berg J, Hamm K and Ruiz-Tiben E. Population

genetic analysis of Chadian Guinea worms reveals that human and non-human hosts share common

parasite populations. *PLoS Negl Trop Dis* 2018; **12:** p.e0006747.

Watts S. Dracunculiasis in Africa in 1986: its geographic extent, incidence, and at-risk population. *Am*

J Trop Med Hyg 1986; **37**: 119–25.

Wood S. mgcv: GAMs and generalized ridge regression for R. *R news* 2001; **1**: 20–5.

Table 1. **Summary of dog characteristics in three settlements in Chad.** Data are from all collared

dogs (n=150), but some dogs have missing data for individual characteristics. Water provision is the

proportion of households which reported providing water for animals.

Settlement	n dogs	Sex (M:F)	Body condition (Poor:Moderate)	Mean age (months)	Age range (months)	Water provision (%)
Lagarna	34	16 : 18	22 : 12	32 ± 4	6 - 96	60
Kakale	52	26 : 26	25 : 27	35 ± 3	4 - 144	75
Magrao	64	40 : 24	16 : 42	34 ± 3	5 - 144	100
All	150	82 : 68	63 : 81	34 ± 2	4 - 144	82

334 **Table 2**. **Summary of dog ranging behaviour in three settlements in Chad.** Core range is the 60%

335 kernel density estimate and total range is the 100% minimum convex polygon. The mean and standard

336 error in parentheses are presented for each variable.

Settlement	n dogs	Core range (km²)	Total range (km²)	Fixes within 100m of a pond (%)	Unique ponds visited	Ponds in core range	Ponds in total range	Fixes within 100m of the Chari River (%)
Lagarna	31	0.30(0.09)	4.70(1.33)	11(4)	2(0.3)	2(0.3)	4(0.8)	4(2)
Kakale	48	2.53(0.74)	21.01(3.08)	36(5)	8(0.6)	8(1.3)	25(3.1)	<1(<1)
Magrao	55	0.17(0.04)	4.39(0.63)	18(4)	3(0.3)	1(0.2)	6(0.7)	2(1)
All	134	1.05(0.28)	10.42(1.35)	23(3)	4(0.3)	3(0.6)	12(1.4)	2(<1)

338
Figure 1: A) Locations of dog study settlements on the Chari River, Chad. B) Adult female
Guinea worm *Dracunculus medinensis* being extracted from a domestic dog *Canis familiaris*. C)
Typical dog fitted with GPS tracking collar. The settlements were: Magrao (10°59'44.31"N,
15°29'29.27"E) which encompasses the associated permanent village of Sawata, Largana
(10°45'15.26"N, 16° 1'39.86"E) and the associated permanent village of Gomba (10°44'46.22"N, 16°
2'46.70"E), and Kakale-Mberi (10°53'0.79"N, 15°38'8.45"E) and the associated seasonal settlement of
Awine (10°48'6.34"N, 15°37'56.61"E). Photographs Jared Wilson-Aggarwal. The satellite image was
generated using the Esri world imagery basemap (sources: Esri, DigitalGlobe, GeoEye, i-cubed,
USDA FSA, USGS, AEX, Getmapping, Aerogrid, IGN, IGP, swisstopo, and the GIS User
Community).
349

350
Figure 2. Distances from human households of the ponds most exposed to visits by dogs in two
settlements in Chad. Dog visits were determined by locating dogs within 100m of a pond by GPS
tracking. Ponds were located from satellite imagery within a search area enclosing all dog location
points for each village and are a sub-sample of all available ponds. All visits by all dogs were summed
for each pond and the cumulative total of all dog visits is plotted against the distance of the pond to the
nearest household with tracked dogs. The dotted lines indicate the distance from a household with
dogs at which 80%, 90% and 95% of all dog visits are captured.

Figure 3. Stable isotope analysis of the diets of dogs in two settlements in Chad. Settlements are

A) Kakale-Mberi and B) Magrao. i) The δ^{15}N and δ^{13}C values for sampled dogs (black dots) and the

mean ± standard deviation of δ^{15}N and δ^{13}C for putative food groups. Trophic discrimination factors,

derived from package SIDER, have been applied to adjust dog isotope ratios downwards for both δ^{15}N

and δ^{13}C. ii) Outputs of Bayesian mixing models, from package SIMMR, of the proportional

contributions of the main food groups to dog diets.

Figure 4. Effects of water provision and of fish consumption on Guinea worm infection history in dogs in Chad. The odds, with 95% confidence intervals, of having had Guinea worm are shown for dogs A) living in households in which animals are, and are not, provided with water, and B) with different modelled proportions of fish in their diets. The effects of fish consumption are presented separately for dogs living in households, in which animals are (dark grey) and are not (light grey) provided with water. Odds are from exponentiated model coefficients looking at the relationship between the history of Guinea worm infection from our field survey, averaged across the top model set and for the different levels of values of predictive factors. Results presented are from dogs in Kakale-Mberi, but dogs in Magrao show the same relationships.

Supplementary Information

374   **Table S1.** Top model sets (ΔAICc < 2) for all analyses included in the results.

Formula	df	LogLik	AICc	ΔAICc	Weight
Guinea worm incidence (history of infection or not)					
Exeter field records (main analysis in the paper)					
% fish consumed + water provision	3	-52.34	110.9	0	0.30
% fish consumed + water provision + settlement	4	-52.18	112.8	1.85	0.19
Longer-term CGWEP records (for validation)					
Water provision + settlement	3	-49.89	106.0	0	0.35
Water provision + settlement + % fish consumed	4	-49.70	107.8	1.78	0.14
Amount of time dogs spend near ponds (no. of fixes within 100 m of ponds)					
Body condition + settlement + water provision + age	7	-1357.02	2729	0	0.37
Body condition + settlement + water provision	6	-1358.29	2729.3	0.29	0.32
Total range size (100 % MCP)					
Days monitored + body condition + settlement	3	12.76	-17.3	0	0.23
Days monitored + body condition + settlement + age	3	12.51	-16.8	0.50	0.10
Days monitored + body condition + settlement + sex	4	13.75	-15.5	1.84	0.10
Core range size (60 % KDE)					
Settlement	4	-204.48	417.3	0	0.13
Settlement + days monitored	5	-203.80	418.2	0.85	0.09
Settlement + body condition	5	-203.91	418.4	1.07	0.08
Settlement + days monitored + body condition	6	-203.17	419.1	1.81	0.05
Settlement + water provision	5	-204.31	419.2	1.85	0.05

375   **Table S2.** The odds ratios and relative risks for predictors of Guinea worm infection status derived from the
376   Chad Guinea Worm Eradication Programme monitoring records. Values are derived from full modelling
377   averaging across the top model (see Table 1) and 95% CIs are derived from model bootstrapping. Significant
378   variables are Settlement and Water provision.

Variable	Odds ratio	95% CIs	Relative risk	95% CIs
Settlement: The risk of dogs in Magrao vs. Kakale	4.69	(1.23, 17.94)	3.24	(1.23, 62.3)
Water provision	0.12	(0.02, 0.63)	0.21	(0.03, 0.68)

379   **Table S3.** Numbers and proportions of dogs with and without history of Guinea worm infection in two
380   settlements, from the data used in the final model of Guinea worm infection (n=100 dogs for Field questionnaire
381   records and n=102 for Chad Guinea Worm Eradication Programme monitoring); according to two different
382   sources of infection history: records taken in the field during household questionnaires when owners were asked
383   whether their dog had ever had Guinea worm, and records of infected dogs from CGWEP.

Record source	Settlement					
	Magrao			Kakale		
	Infection history	*No infection history*	*Rate (%)*	*Infection history*	*No infection history*	*Rate (%)*
Field questionnaires	13	38	25.5	14	35	28.6
CGWEP records	15	37	28.8	8	42	16.0