A single gene expressed in fibroblastic reticular cells predicts response to cancer immunotherapy

Daniele Biasci[1,2,✉]**, Simon Tavaré**[2,3,*]**, and James Thaventhiran**[1,2*]

[1]MRC Toxicology Unit, University of Cambridge, Robinson Way, Cambridge, CB2 0RE, United Kingdom
[2]Cancer Research UK Cambridge Institute, Robinson Way, Cambridge CB2 0RE, United Kingdom
[3]Herbert and Florence Irving Institute for Cancer Dynamics, Columbia University, Schermerhorn Hall, Suite 601, 1190 Amsterdam Ave, New York, NY 10027, United States

While the role of T cells in immune response to cancer is well established, the role of B cells remains more controversial (1–3). Recently, it was reported that expression of B cell specific genes inside tumours is associated with response to immune checkpoint blockade (ICB), thus suggesting an unappreciated biological role for B cells in promoting ICB response (4). However, association between B cell gene expression and ICB response was not statistically significant when other immune populations, including CD8+ T cells, were taken into account (4). For this reason, it remains unclear whether B cells are associated with ICB response independently of CD8+ T cell infiltration, a validated predictor of immunotherapy response (5–7) strongly correlated with B cells (8). Two complementary studies did not address this issue, because they described predictive signatures containing genes expressed in CD8+ T cells, as well as other immune cell types (9, 10). Helmink et al. acknowledge this limitation of their work, attributing it to limited sample size, and call for larger studies to clarify the point (4). By conducting the largest gene expression study of ICB response to date, here we show that: 1) pre-treatment expression of B cell genes is indeed associated with ICB response; 2) such association is independent of CD8+ T cells; 3) the association can be completely explained by a single gene expressed in fibroblastic reticular cells (FRCs). We further validate this last finding in three independent cohorts of patients treated with ICB.

We analysed the entire transcriptome of 137 melanoma samples, considering only samples collected before commencing treatment with immune checkpoint inhibitors (see supplementary methods). We excluded tumour samples obtained from lymph node metastasis, in order to minimise the risk of capturing unrelated immune populations from the adjacent lymphoid tissue. We calculated the correlation between expression of each gene at baseline and subsequent treatment response, as defined in the original studies according to the RECIST criteria (11). A distinct group of genes showed significant association with treatment response, namely: *CR2* (CD21), *MS4A1* (CD20), *CD19*, *FCER2* (CD23), *PAX5, BANK1, VPREB3, TCL1A, CLEC17A* and *FDCSP* (Fig. 1A and Table S1). Literature reports that these genes are predominantly expressed in B cells (12), with the exception of *FDCSP*, which was found to be expressed follicular dendritic cells (FDCs) isolated from secondary lymphoid organs, but not in B cells (13, 14). In order to identify the cell populations expressing these genes in tumours, we assessed their expression in single-cell RNA sequencing data (scRNAseq) from human cancer samples (15). First, we confirmed that genes identified by our association study were specifically expressed in tumour-associated B cells (Fig. S8). Second, we observed that *FDCSP* was not expressed in tumour-associated B cells (Fig. S1), but identified a subset of fibroblastic cells (*FAP+ COL1A1+ CD31-*; Fig. S2) expressing markers consistent with FRC identity (*CCL19+ CCL21+ BAFF+*; 16, 17; Figures S3 and S4 and tables S5 and S6). Finally, we confirmed these results in an independent set of melanoma samples profiled at the single-cell level (18; Figures S4 to S7 and S9). We then sought to determine whether expression of these genes was associated with ICB response independently of CD8+ T cell infiltration. To this aim, we repeated the association analysis after taking into account expression markers of T cells, NK cells, cross-presenting dendritic cells (19) and interferon-gamma response (20; Fig. 1D and table S1). We found that only four genes were associated with ICB response independently of all immune markers considered: *MS4A1* (prototypical B cell marker), *CLEC17A* (expressed in dividing B cells in germinal centers; 21), *CR2* (expressed in mature B cells and FDCs; 22) and *FDCSP* (expressed in FRCs but not in B cells). This result suggested that presence of B cells and FRCs in the tumour microenvironment (TME) before treatment was associated with subsequent ICB response independently of CD8+ T cells (Fig. 1D and Table S7). Remarkably, we observed that *FDCSP* remained significantly associated with ICB response even after B cell markers were taken into account (Fig. 1D) and that no other gene remained significant after *FDCSP* expression was included in the model (Fig. 1D). Taken together, these results suggest that expression of *FDCSP* might be an independent predictor of ICB response. We tested this hypothesis in a completely independent cohort of melanoma patients treated with anti-PD1, either alone or in combination with anti-CTLA4 (23). In this validation cohort, patients with higher expression of *FDCSP* at baseline experienced significantly longer overall survival (log-rank $p < 0.0001$) and progression-free survival (log-rank $p < 0.0001$) after commencing treatment with ICB (Figures 1C and 1E). Moreover, we observed a significant association between expression of *FDCSP* at baseline and subsequent RECIST response (Cochran-Armitage test $p < 0.0001$; Fig. 1E). We sought to replicate this observation in two additional independent cohorts of patients treated with ICB in melanoma (4) and clear cell renal cell carcinoma (24) and found similar results (Figures 1F and 1G). On the other hand, testing B cell markers in the validation cohorts produced less consistent results (Fig. S10). Finally, we performed multiple linear regres-

---

* These authors have contributed equally to this work.

sion analysis and found that the association between *FDCSP* and ICB response was independent of CD8+ T cells, CD4+ T cells, B cells, NK cells, myeloid Dendritic cells (mDCs), Macrophages and Plasma cells (Tables S2 and S3).

While CD8+ T cells express the molecular targets of ICB, and their presence in the TME is a validated predictor of ICB response [5–7], the role of B cells remains more controversial [1–3]. Because T and B cells often co-occur in the TME [25], gene expression studies consistently reported strong correlation between T cells, B cells and other immune cell types, especially in melanoma [8]. As a consequence, assessing whether B cells are associated with ICB response independently of CD8+ T cells has been proven challenging [4]. By analysing a larger number of samples compared to previous studies, we were able to demonstrate that B cells predict ICB response independently of CD8+ T cells (Fig. 1D and Table S4). This supports the idea that B cells might actively promote ICB response [3], rather than being irrelevant [1] or detrimental to it [2]. However, our data also shows that such positive association can be completely explained by a single gene expressed outside of the B cell compartment. We have demonstrated that *FDCSP* is transcribed in a subset of fibroblastic cells up-regulating *CCL19, CCL21* and *BAFF* and down-regulating genes associated with canonical fibroblasts, such as extracellular matrix genes (Fig. S4, Table S5, Table S6), an expression pattern consistent with FRC identity [16, 17]. One way to reconcile these observations is to consider that FRCs are required to initiate efficient B and T cell responses [16]. In both secondary lymphoid organs (SLO) and tertiary lymphoid structures (TLS), FRCs form reticular networks that facilitate interactions between B cells, T cells and their cognate antigens [17, 26]. Homing of immune cells into the network requires interaction between chemokines produced by FRCs (CCL19 and CCL21) and their receptor (CCR7) expressed on T cells, B cells and dendritic cells [27, 28; see also Table S1]. Accordingly, depleting FRCs causes loss of T cells, B cells and dendritic cells in both SLO and TLS and decreases the magnitude of B and T cell responses to subsequent viral infection [16, 29]. Perhaps more importantly, differentiation of fibroblasts into FRCs occurs early during TLS formation, precedes B and T cell infiltration and can still be observed in *Rag2-/-* mice [29]. In this context, our finding that a gene expressed in FRCs is associated with ICB response independently of B and T cell infiltration is coherent with the current understanding of TLS neogenesis. Taken together, these observations suggest that the reported association between B cells and ICB response [4], albeit independent of CD8+ T cells, might ultimately be secondary to the presence of FRCs in the TME. The identification of *FDCSP* as a single marker of ICB response should enable even larger studies to test this conclusion.

Bibliography

1. Damsky, W. *et al.* B cell depletion or absence does not impede anti-tumor activity of pd-1 inhibitors. *Journal for immunotherapy of cancer* **7**, 153 (2019).
2. Somasundaram, R. *et al.* Tumor-associated b-cells induce tumor heterogeneity and therapy resistance. *Nature communications* **8**, 1–16 (2017).
3. Hollern, D. P. *et al.* B cells and t follicular helper cells mediate response to checkpoint inhibitors in high mutation burden mouse models of breast cancer. *Cell* **179**, 1191–1206 (2019).
4. Helmink, B. A. *et al.* B cells and tertiary lymphoid structures promote immunotherapy response. *Nature* (2020).
5. Ji, R.-R. *et al.* An immune-active tumor microenvironment favors clinical response to ipilimumab. *Cancer Immunology, Immunotherapy* **61**, 1019–1031 (2012).
6. Ayers, M. *et al.* Ifn-γ–related mrna profile predicts clinical response to pd-1 blockade. *The Journal of clinical investigation* **127**, 2930–2940 (2017).
7. Cristescu, R. *et al.* Pan-tumor genomic biomarkers for pd-1 checkpoint blockade–based immunotherapy. *Science* **362**, eaar3593 (2018).
8. Iglesia, M. D. *et al.* Genomic analysis of immune cell infiltrates across 11 tumor types. *JNCI: Journal of the National Cancer Institute* **108** (2016).
9. Cabrita, R. *et al.* Tertiary lymphoid structures improve immunotherapy and survival in melanoma. *Nature* (2020).
10. Petitprez, F. *et al.* B cells are associated with survival and immunotherapy response in sarcoma. *Nature* (2020).
11. Schwartz, L. H. *et al.* Recist 1.1—update and clarification: From the recist committee. *European journal of cancer* **62**, 132–137 (2016).
12. Kassambara, A. *et al.* Genomicscape: an easy-to-use web tool for gene expression data analysis. application to investigate the molecular events in the differentiation of b cells into plasma cells. *PLoS computational biology* **11**, e1004077 (2015).
13. Marshall, A. J. *et al.* Fdc-sp, a novel secreted protein expressed by follicular dendritic cells. *The Journal of Immunology* **169**, 2381–2389 (2002).
14. Al-Alwan, M. *et al.* Follicular dendritic cell secreted protein (fdc-sp) regulates germinal center and antibody responses. *The Journal of Immunology* **178**, 7859–7867 (2007).
15. Zilionis, R. *et al.* Single-cell transcriptomics of human and mouse lung cancers reveals conserved myeloid populations across individuals and species. *Immunity* **50**, 1317–1334 (2019).
16. Denton, A. E., Roberts, E. W., Linterman, M. A. & Fearon, D. T. Fibroblastic reticular cells of the lymph node are required for retention of resting but not activated cd8+ t cells. *Proceedings of the National Academy of Sciences* **111**, 12139–12144 (2014).
17. Cremasco, V. *et al.* B cell homeostasis and follicle confines are governed by fibroblastic reticular cells. *Nature immunology* **15**, 973–981 (2014).
18. Jerby-Arnon, L. *et al.* A cancer cell program promotes t cell exclusion and resistance to checkpoint blockade. *Cell* **175**, 984–997 (2018).
19. Hartung, E. *et al.* Induction of potent CD8 t cell cytotoxicity by specific targeting of antigen to cross-presenting dendritic cells in vivo via murine or human XCR1. *The Journal of Immunology* **194**, 1069–1079 (2014).
20. Chow, M. T. *et al.* Intratumoral activity of the cxcr3 chemokine system is required for the efficacy of anti-pd-1 therapy. *Immunity* **50**, 1498–1512 (2019).
21. Graham, S. A. *et al.* Prolectin, a glycan-binding receptor on dividing b cells in germinal centers. *Journal of Biological Chemistry* **284**, 18537–18544 (2009).
22. Takahashi, K. *et al.* Mouse complement receptors type 1 (cr1; cd35) and type 2 (cr2; cd21): expression on normal b cell subpopulations and decreased levels during the development of autoimmunity in mrl/lpr mice. *The Journal of Immunology* **159**, 1557–1569 (1997).
23. Gide, T. N. *et al.* Distinct immune cell populations define response to anti-pd-1 monotherapy and anti-pd-1/anti-ctla-4 combined therapy. *Cancer cell* **35**, 238–255 (2019).
24. Miao, D. *et al.* Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. *Science* **359**, 801–806 (2018).
25. Garnelo, M. *et al.* Interaction between tumour-infiltrating b cells and t cells controls the progression of hepatocellular carcinoma. *Gut* **66**, 342–351 (2017).
26. Denton, A. E., Carr, E. J., Magiera, L. P., Watts, A. J. & Fearon, D. T. Embryonic fap+ lymphoid tissue organizer cells generate the reticular network of adult lymph nodes. *Journal of Experimental Medicine* **216**, 2242–2252 (2019).
27. Gunn, M. D. *et al.* Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *The Journal of experimental medicine* **189**, 451–460 (1999).
28. Förster, R. *et al.* Ccr7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* **99**, 23–33 (1999).
29. Nayar, S. *et al.* Immunofibroblasts are pivotal drivers of tertiary lymphoid structure formation and local pathology. *Proceedings of the National Academy of Sciences* **116**, 13490–13497 (2019).

Fig. 1. | Expression of FDCSP is an independent predictor of response to immune checkpoint blockade. A. Volcano plot showing genes significantly associated with response to ICB. **B-C** Kaplan-Meier curves showing different overall survival and progression-free survival in melanoma patients after commencing treatment with ICB. Patients from the first validation cohort (23; see supplementary methods) were stratified in three groups according to pre-treatment expression of *FDCSP*. **D** Heatmap showing association between pre-treatment expression of B cell or FRC genes (rows) and subsequent ICB response after the effect of other immune markers (columns) was taken into account. The number inside each cell represents the Wald test p-value for the underlying effect size estimate obtained using the negative binomial model for RNAseq counts implemented in DESEq2 (see supplementary methods). Nominal p-values were corrected for multiple testing using Benjamini–Yekutieli false discovery rate (FDR) and are also reported in Table S7. A red background highlights effect sizes estimated to be greater than zero with FDR < 0.05. In this study, an effect size greater than zero represents a positive association between gene expression and probability of ICB response. **E-G** Association between pre-treatment expression of *FDCSP* and subsequent ICB response in the first (23), second (4) and third validation cohort (24). See supplementary methods.

Appendix S1: supplementary methods

Bulk RNAseq: discovery cohorts. The RNAseq datasets used in this study as discovery cohorts were downloaded from the following published sources: data for 27 melanoma samples from patients that went on to receive ICB treatment was downloaded from the NIH Genomics Data Common Portal, study TCGA-SKCM [1, 2]; data for 42 pre-ICB melanoma samples was downloaded from dbGaP [3], study phs000452.v2.p1 [4]; data for 109 melanoma samples collected before or during ICB treatment was downloaded from Gene Expression Omnibus (GEO; [5]), series GSE91061 [6]; data for 28 melanoma samples collected before ICB treatment was downloaded from GEO, series GSE78220 [7]; data for 37 melanoma samples collected before or during ICB treatment was downloaded from GEO, series GSE115821 [8]. Clinical annotations for each sample (i.e. type of ICB treatment received, sample collection time point, tumour site, treatment response) were obtained from the original publications. Samples collected after commencing treatment with ICB, samples obtained from lymph node metastasis, and samples for which treatment response status was not available were excluded from subsequent analyses, leaving 137 pre-treatment samples available for this study.

Bulk RNAseq: validation cohorts. The RNAseq datasets used in this study as validation cohorts were downloaded from the following published sources. *First validation cohort*: data for 91 melanoma samples collected before or during ICB treatment was downloaded from the European Nucleotide Archive (ENA; [9]), study PRJEB23709 [10]. *Second validation cohort*: data for 16 melanoma samples collected before ICB treatment was obtained from [11]. *Third validation cohort*: data for 33 clear cell renal cell carcinoma samples collected before ICB treatment was obtained from [12]. Clinical annotations for each sample were obtained from the original publications. Samples collected after commencing treatment with ICB and samples for which treatment response status was not available were excluded from subsequent analyses.

Bulk RNAseq: gene expression quantification. Sequences of all known human transcripts were downloaded from Ensembl release 97 [13]. Transcript quantification was performed using Kallisto ver. 0.43 [14] and gene-level count matrices for use in DESeq2 [15] were calculated using tximport [16] as recommended by DESeq2 authors [17]. All subsequent analyses on gene expression were performed using R 3.5.0 [18]. For differential expression analysis, raw counts were used directly in DESeq2 as recommended [17]. For other downstream analyses (i.e. survival curves, multiple correlation analysis and others) counts data were transformed using the variance stabilizing transformation [19] as implemented in DESEq2 [17] and potential systematic differences between cohorts were corrected using ComBat [20] as implemented in the sva package [21]. The methods described in this section were used to quantify gene expression in all datasets used in this study as discovery cohorts and for the first validation cohort, for which original FASTQ files were available. For the second and third validation cohort [10], we used the gene expression values provided in the original publications [11, 12].

Bulk RNAseq: genes associated with ICB response. In the discovery cohorts, association between gene expression and ICB response was calculated using the DESeq2 model based on the negative binomial distribution [15]. Raw counts for each gene were used directly in the model as recommended by DESeq2 authors [17]. We used the ICB response status for each patient as reported in the original study [1, 4, 6–8] according to the RECIST criteria [22]. RECIST response categories were encoded as a numerical score as follows: PD (progressive disease) $= -1$, SD (stable disease) $= 0$, PR (partial response) $= 0.5$, CR (complete response) $= 1$. When we assessed the use of a different numerical score (i.e. PD$= 0$, SD$=0.33$, PR$= 0.66$, CR$= 1$), we obtained similar results. In order to account for systematic differences between cohorts, a categorical variable encoding the cohort of origin for each sample was included as a covariate in the DESeq2 association model. In order to take into account the effect of known immune markers (Fig. 1D), the association analysis was repeated after their expression level was calculated according to DESeq2 documentation [17] and added to the model as an additional covariate.

Bulk RNAseq: survival curves. Survival curves were plotted using the R package survminer [23]. Patients were stratified in three groups according to the baseline expression of the gene of interest: the high expression group was defined as containing the 25% of samples with highest expression, the low expression group was defined as containing the 25% of samples with lowest expression, and the medium expression group was defined as containing the remaining samples. Statistical significance of the observed difference between groups was assessed using the Logrank test [24].

Single-cell RNAseq: original data sets. Expression values (normalised counts) for 54773 single cells isolated from non-small-cell lung cancer (NSCLC) tumour samples were downloaded from GEO, series GSE127465 [25]. Cell type annotation and a two-dimensional visualisation (SPRING plot, [26]) of single-cell transcriptomes were also obtained from the original publication [25] and used for subsequent analyses. Expression values (transcripts per million) for 7186 single cells isolated from melanoma samples were downloaded from GEO, series GSE115978 [27]. Cell type annotation and a two-dimensional visualisation (tSNE plot, [28]) of single-cell transcriptomes were obtained from the original publication [27] and used for subsequent analyses.

Single-cell RNAseq: differentially expressed genes. Expression values obtained from GSE127465 and GSE115978 were transformed in logarithmic scale ($y = \log_2(x+1)$). Cells annotated as fibroblasts in the original study were selected and used to compare fibroblasts expressing *FDCSP* (*FDCSP+*) against fibroblasts not expressing *FDCSP* (*FDCSP-*). Statistical significance for gene expression differences observed between these two groups was assessed using unpaired two-samples Wilcoxon test (29). Genes were ranked using a numerical score which takes into account both p-value and $\log_2(fc)$: score $= \sqrt{-\log_{10}(\text{p} - \text{value})} * |\log_2(fc)| * \text{sign}(\log_2(fc))$. Results for GSE127465 (non-small cell lung cancer) and GSE115978 (melanoma) were meta-analysed using the RankProduct method (30, 31) and are presented in Tables S5 and S6.

Code availability. The authors declare that the code used for this study is available upon request.

Supplementary methods references.

1. Akbani, R. *et al.* Genomic classification of cutaneous melanoma. *Cell* **161**, 1681–1696 (2015).
2. Grossman, R. L. *et al.* Toward a shared vision for cancer genomic data. *New England Journal of Medicine* **375**, 1109–1112 (2016).
3. Mailman, M. D. *et al.* The ncbi dbgap database of genotypes and phenotypes. *Nature genetics* **39**, 1181–1186 (2007).
4. Van Allen, E. M. *et al.* Genomic correlates of response to ctla-4 blockade in metastatic melanoma. *Science* **350**, 207–211 (2015).
5. Edgar, R., Domrachev, M. & Lash, A. E. Gene expression omnibus: Ncbi gene expression and hybridization array data repository. *Nucleic acids research* **30**, 207–210 (2002).
6. Riaz, N. *et al.* Tumor and microenvironment evolution during immunotherapy with nivolumab. *Cell* **171**, 934–949 (2017).
7. Hugo, W. *et al.* Genomic and transcriptomic features of response to anti-pd-1 therapy in metastatic melanoma. *Cell* **165**, 35–44 (2016).
8. Auslander, N. *et al.* Robust prediction of response to immune checkpoint blockade therapy in metastatic melanoma. *Nature medicine* **24**, 1545–1549 (2018).
9. Leinonen, R. *et al.* The european nucleotide archive. *Nucleic acids research* **39**, D28–D31 (2010).
10. Gide, T. N. *et al.* Distinct immune cell populations define response to anti-pd-1 monotherapy and anti-pd-1/anti-ctla-4 combined therapy. *Cancer cell* **35**, 238–255 (2019).
11. Helmink, B. A. *et al.* B cells and tertiary lymphoid structures promote immunotherapy response. *Nature* (2020).
12. Miao, D. *et al.* Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. *Science* **359**, 801–806 (2018).
13. Cunningham, F. *et al.* Ensembl 2019. *Nucleic Acids Research* **47**, D745–D751 (2018). URL https://doi.org/10.1093/nar/gky1113.
14. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **34**, 525–527 (2016). URL https://doi.org/10.1038/nbt.3519.
15. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15** (2014). URL https://doi.org/10.1186/s13059-014-0550-8.
16. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2015). URL https://doi.org/10.12688/f1000research.7563.1.
17. Love, M. I., Anders, S. & Huber, W. Analyzing rna-seq data with deseq2 (2019). URL http://bioconductor.org/packages/devel/bioc/vignettes/DESeq2/inst/doc/DESeq2.html#tximport.
18. R Core Team. *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing, Vienna, Austria (2018). URL https://www.R-project.org/.
19. Huber, W., von Heydebreck, A., Sültmann, H., Poustka, A. & Vingron, M. Parameter estimation for the calibration and variance stabilization of microarray data. *Statistical applications in genetics and molecular biology* **2** (2003).
20. Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression data using empirical bayes methods. *Biostatistics* **8**, 118–127 (2007).
21. Leek, J. T. *et al.* Sva: surrogate variable analysis. 2015. *R package version* **3**, 25–27.
22. Schwartz, L. H. *et al.* Recist 1.1—update and clarification: From the recist committee. *European journal of cancer* **62**, 132–137 (2016).
23. Kassambara, A., Kosinski, M., Biecek, P. & Fabian, S. Package 'survminer'. *Drawing Survival Curves using 'ggplot2'.(R package version 0.3. 1.)* (2017).
24. Peto, R. & Peto, J. Asymptotically efficient rank invariant test procedures. *Journal of the Royal Statistical Society: Series A (General)* **135**, 185–198 (1972).
25. Zilionis, R. *et al.* Single-cell transcriptomics of human and mouse lung cancers reveals conserved myeloid populations across individuals and species. *Immunity* **50**, 1317–1334 (2019).
26. Weinreb, C., Wolock, S. & Klein, A. M. Spring: a kinetic interface for visualizing high dimensional single-cell expression data. *Bioinformatics* **34**, 1246–1248 (2018).
27. Jerby-Arnon, L. *et al.* A cancer cell program promotes t cell exclusion and resistance to checkpoint blockade. *Cell* **175**, 984–997 (2018).
28. Maaten, L. v. d. & Hinton, G. Visualizing data using t-sne. *Journal of machine learning research* **9**, 2579–2605 (2008).
29. Mann, H. B. & Whitney, D. R. On a test of whether one of two random variables is stochastically larger than the other. *The annals of mathematical statistics* 50–60 (1947).
30. Breitling, R., Armengaud, P., Amtmann, A. & Herzyk, P. Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS letters* **573**, 83–92 (2004).
31. Hong, F. & Breitling, R. A comparison of meta-analysis methods for detecting differentially expressed genes in microarray experiments. *Bioinformatics* **24**, 374–382 (2008).

Appendix S2: supplementary figures

(A)

(B)

(C)

(D)

Fig. S1. | **FDCSP is not expressed in B cells infiltrating lung cancer tumours.** Two-dimensional visualizations (SPRING plots) of single-cell RNA sequencing data obtained from lung cancer tumours and published in (25). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). Expression of *FDCSP* is observed in a group of cells indicated by the black arrow in (**A**) well distinct from cells expressing prototypical B cell markers (**B-D**).

Fig. S2. | **FDCSP is expressed in a subset of fibroblasts in lung cancer tumours.** Two-dimensional visualizations (SPRING plots) of single-cell RNA sequencing data obtained from lung cancer tumours and published in (25). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). **A.** Expression of *FDCSP*. The black arrow indicates a group of transcriptionally related cells containing *FDCSP+* cells. **B.** Expression of *FAP*. **C.** Expression of *COL1A1*. **D.** Expression of *PECAM1/CD31*.

Fig. S3. | FDCSP+ cells express markers of FRCs in lung cancer tumours. Two-dimensional visualizations (SPRING plots) of single-cell RNA sequencing data obtained from lung cancer tumours and published in (25). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). **A.** Expression of *FDCSP*. The black arrow indicates a group of transcriptionally related cells containing *FDCSP+* cells. **B.** Expression of *CCL19*. **C.** Expression of *CCL21*. **D.** Expression of *TNFSF13B/BAFF*.

Fig. S4. | *FDCSP+ fibroblasts upregulate FRCs markers in lung cancer and melanoma.* Differential gene expression analysis comparing single-cell transcriptomes of *FDCSP+* fibroblasts versus *FDCSP-* fibroblasts obtained from lung cancer and melanoma tumours. The analysis was performed using gene expression data published in (25, 27); see also supplementary methods, Table S5 and Table S6. **A.** Volcano plot representing genes differentially expressed between *FDCSP+* and *FDCSP-* fibroblasts shows that *FDCSP+* fibroblasts significantly upregulate expression of FRCs markers (*CCL19, CCL21, TNFSF13B/BAFF*) in lung cancer. Genes considered significantly up-regulated ($\log_2(fc) > \max(\log_2(fc))/4$ and FDR < 0.05) are marked in red; dotted lines represent chosen $\log_2(fc)$ and FDR cutoffs. **B.** Volcano plot representing genes differentially expressed between *FDCSP+* and *FDCSP-* fibroblasts and confirming significant up-regulation of FRCs markers in melanoma. **C.** Violin plots comparing distribution of gene expression values in *FDCSP+* and *FDCSP-* fibroblasts in lung cancer. The plots shows that *FDCSP+* fibroblasts significantly upregulate cytokines relevant to FRCs function while down-regulating genes involved in secretion of the extracellular matrix (*TIMP3, FN1*). Statistical significance was assessed using unpaired two-samples Wilcoxon test. A red diamond represents the median value of each distribution. **D.** Comparing *FDCSP+* and *FDCSP-* obtained from melanoma tumours leads to similar results (see panel C).

(A)

(B)

(C)

(D)

Fig. S5. | **FDCSP is not expressed in B cells infiltrating melanoma.** Two-dimensional visualizations (tSNE plots) of single-cell RNA sequencing data obtained from melanoma tumours and published in (27). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). Expression of *FDCSP* is observed in a group of cells indicated by a black arrow in (**A**) well distinct from cells expressing prototypical B cell markers (**B-D**).

Fig. S6. | **FDCSP is expressed in a subset of fibroblasts in melanoma.** Two-dimensional visualizations (tSNE plots) of single-cell RNA sequencing data obtained from melanoma tumours and published in (27). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). **A.** Expression of *FDCSP*. The black arrow indicates a group of transcriptionally related cells containing *FDCSP+* cells. **B.** Expression of *FAP*. **C.** Expression of *COL1A1*. **D.** Expression of *PECAM1/CD31*.

Fig. S7. | FDCSP+ cells express markers of FRCs in melanoma. Two-dimensional visualizations (tSNE plots) of single-cell RNA sequencing data obtained from melanoma tumours and published in (27). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). **A.** Expression of *FDCSP*. The black arrow indicates a group of transcriptionally related cells containing *FDCSP+* cells. **B.** Expression of *CCL19*. **C.** Expression of *CCL21*. **D.** Expression of *TNFSF13B/BAFF*.

Fig. S8. | Expression of ICB response associated genes in single cells obtained from lung cancer tumours. Two-dimensional visualizations (SPRING plots) of single-cell RNA sequencing data obtained from lung cancer tumours and published in (25). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The expression level color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). Panels B and C are included in the figure in order to further confirm the identity of cells annotated as "B cells" and "Plasma cells" in panel A. **A.** Cell types annotation as reported in the original study (25). **B.** Expression of *MS4A1/CD20*. **D.** Expression of *CD79A* is detectable in B cells and Plasma cells as expected. **D.** Expression of *TNFRSF17/BCMA* is detectable in Plasma cells but not in B cells as expected. **E.** Expression of *CD19*. **F.** Expression of *PAX5*. **G.** Expression of *BANK1*. **H.** Expression of *VPREB3*. **I.** Expression of *CR2*. **J.** Expression of *FCER2*. **K.** Expression of *TCL1A*, also detectable in Plasmacytoid dendritic cells (pDC, see panel A). **L.** Expression of *CLEC17A*.

Fig. S9. | Expression of ICB response associated genes in single cells obtained from melanoma tumours. Two-dimensional visualizations (tSNE plots) of single-cell RNA sequencing data obtained from melanoma tumours and published in (27). Each dot represents the transcriptional profile of a single cell. Cells closely associated in each plot are more likely to transcribe similar genes and might thus belong to the same cell type. Expression values and coordinates for each dot were obtained from the original study. The expression level color scale indicates expression level for a particular gene ranging from 0 (minimum expression value observed in the dataset for that gene, usually corresponding to non detectable expression) to 100 (maximum expression value observed in the dataset for that gene). Panels B and C are included in the figure in order to further confirm the identity of cells annotated as "B cells" and "Plasma cells" in panel A. **A.** Cell types annotation as reported in the original study (27). **B.** Expression of *MS4A1/CD20*. **D.** Expression of *CD79A* **D.** Expression of *TNFRSF17/BCMA*. A small cluster of Plasma cells might be visible but was not clearly annotated in this dataset (see panel A). **E.** Expression of *CD19*. **F.** Expression of *PAX5*. **G.** Expression of *BANK1*. **H.** Expression of *VPREB3*. **I.** Expression of *CR2*. **J.** Expression of *FCER2*. **K.** Expression of *TCL1A*. A small cluster of pDC might be visible but was not clearly annotated in this dataset (see panel A). **L.** Expression of *CLEC17A*.

Fig. S10. | Predicting response in ICB treated patients using pre-treatment expression of FRCs or B cells markers. **A-B.** Kaplan-Meier curves showing different overall survival and progression-free survival in melanoma patients after commencing treatment with ICB. Patients from (10) were stratified in three groups according to pre-treatment expression of *FDCSP*. **C-D.** Association between pre-treatment expression of *FDCSP* and subsequent ICB response in melanoma patients from (10; Panel C) and in clear cell renal cell carcinoma patients from (12; Panel D). Patients were stratified in three groups according to pre-treatment expression of *FDCSP* maintaining the same proportions used in panel A and D. **E-H.** Corresponding results obtained using pre-treatment expression of *MS4A1/CD20*. **I-L.** Corresponding results obtained using pre-treatment expression of *CD19*. **M-P.** Corresponding results obtained using pre-treatment expression of *CR2*.

Appendix S3: supplementary tables

Table S1 | Top 50 genes for which pre-treatment expression is associated with subsequent ICB response in melanoma. Genes are ordered by the column *score* which takes into account both p-value and $\log_2(fc)$: score $= \sqrt{-\log_{10}(\text{p}-\text{value}) * |\log_2(fc)|} * \text{sign}(\log_2(fc))$. The complete list is provided in a separate spreadsheet file (icb_resp_baseline.xlsx).

	gene	$\log_2(fc)$	$\log_2(fc)$ se	p-value	padj	stat	description	score
1	CR2	2.5	0.27	3.3E-21	4.9E-17	9.5	complement C3d receptor 2	7.1
2	FCER2	2.2	0.32	1.3E-13	6.3E-10	7.4	Fc fragment of IgE receptor II	5.3
3	CLEC17A	2	0.28	1.9E-14	1.5E-10	7.7	C-type lectin domain containing 17A	5.3
4	PAX5	2.2	0.33	8.2E-13	3.1E-09	7.2	paired box 5	5.1
5	FDCSP	2.2	0.36	6.7E-11	1.7E-07	6.5	follicular dendritic cell secreted protein	4.7
6	CD19	1.8	0.3	1.2E-11	3.5E-08	6.8	CD19 molecule	4.5
7	TCL1A	1.9	0.34	4.2E-10	6.3E-07	6.2	T cell leukemia/lymphoma 1A	4.2
8	VPREB3	1.7	0.31	1.7E-10	2.9E-07	6.4	V-set pre-B cell surrogate light chain 3	4.1
9	BANK1	1.6	0.28	1E-10	2E-07	6.5	B cell scaffold protein with ankyrin repeats 1	4
10	MS4A1	1.7	0.32	1.2E-09	1.6E-06	6.1	membrane spanning 4-domains A1	3.9
11	TLR10	1.4	0.24	9E-11	1.9E-07	6.5	toll like receptor 10	3.8
12	BLK	1.3	0.25	3.1E-09	3.6E-06	5.9	BLK proto-oncogene, Src family tyrosine kinase	3.3
13	MUC7	1.7	0.44	6.6E-07	0.00038	5	mucin 7, secreted	3.3
14	PSCA	1.5	0.31	4.1E-08	3.8E-05	5.5	prostate stem cell antigen	3.3
15	SPIB	1.3	0.26	9.9E-09	9.9E-06	5.7	Spi-B transcription factor	3.3
16	CYP4F11	1.4	0.31	5.1E-08	4.5E-05	5.4	cytochrome P450 family 4 subfamily F member 11	3.2
17	FAM129C	1.2	0.23	5.3E-09	5.7E-06	5.8	family with sequence similarity 129 member C	3.2
18	CNR2	1.2	0.25	6.6E-08	5.5E-05	5.4	cannabinoid receptor 2	2.9
19	CUX2	1.3	0.3	3.2E-07	0.0002	5.1	cut like homeobox 2	2.9
20	FCRL1	1.3	0.33	1E-06	0.00053	4.9	Fc receptor like 1	2.8
21	CCR7	1.2	0.25	2E-07	0.00014	5.2	C-C motif chemokine receptor 7	2.8
22	P2RX5	1.1	0.24	2.2E-07	0.00015	5.2	purinergic receptor P2X 5	2.7
23	GPX2	1.1	0.29	3E-06	0.0014	4.7	glutathione peroxidase 2	2.5
24	TSPAN8	1.2	0.38	1.1E-05	0.0035	4.4	tetraspanin 8	2.5
25	TNFRSF13C	0.98	0.22	8.7E-07	0.00048	4.9	TNF receptor superfamily member 13C	2.4
26	CCL21	1.2	0.39	1.6E-05	0.0048	4.3	C-C motif chemokine ligand 21	2.4
27	ABAT	0.9	0.21	1.2E-06	0.00061	4.9	4-aminobutyrate aminotransferase	2.3
28	RGS13	1.1	0.34	1.8E-05	0.0051	4.3	regulator of G protein signaling 13	2.2
29	GP1BA	0.88	0.21	2.1E-06	0.00096	4.7	glycoprotein Ib platelet subunit alpha	2.2
30	STAP1	1	0.31	1.4E-05	0.0044	4.3	signal transducing adaptor family member 1	2.2
31	CD79A	1	0.37	2.7E-05	0.0068	4.2	CD79a molecule	2.2
32	SLC27A2	1	0.32	1.8E-05	0.0051	4.3	solute carrier family 27 member 2	2.2
33	C4A	0.88	0.23	4.9E-06	0.002	4.6	complement C4A (Rodgers blood group)	2.2
34	TREML2	0.93	0.26	9.8E-06	0.0033	4.4	triggering receptor expressed on myeloid cells like 2	2.2
35	APOC4-APOC2	1.1	0.46	5.1E-05	0.011	4.1	APOC4-APOC2 readthrough (NMD candidate)	2.2
36	DNASE1L3	0.96	0.3	1.8E-05	0.0051	4.3	deoxyribonuclease 1 like 3	2.1
37	LRMP	0.78	0.2	5.5E-06	0.0021	4.5	lymphoid restricted membrane protein	2
38	AMDHD1	0.83	0.25	1.7E-05	0.0051	4.3	amidohydrolase domain containing 1	2
39	CPS1	0.88	0.31	3.5E-05	0.0085	4.1	carbamoyl-phosphate synthase 1	2
40	PLAC8	0.84	0.27	2.5E-05	0.0066	4.2	placenta associated 8	2
41	CCR6	0.96	0.46	9.7E-05	0.017	3.9	C-C motif chemokine receptor 6	2
42	SELL	0.8	0.25	2E-05	0.0054	4.3	selectin L	1.9
43	C4B	0.78	0.25	2.5E-05	0.0066	4.2	complement C4B (Chido blood group)	1.9
44	MUC3A	0.77	0.27	4.1E-05	0.0092	4.1	mucin 3A, cell surface associated	1.8
45	VIPR1	0.78	0.32	7.4E-05	0.014	4	vasoactive intestinal peptide receptor 1	1.8
46	IGHV1-58	0.89	0.62	0.00026	0.035	3.7	immunoglobulin heavy variable 1-58	1.8
47	SLC22A1	0.79	0.37	0.00011	0.018	3.9	solute carrier family 22 member 1	1.8
48	CFHR4	0.9	0.77	0.00036	0.039	3.6	complement factor H related 4	1.8
49	ABCG5	0.79	0.37	0.00012	0.019	3.9	ATP binding cassette subfamily G member 5	1.8
50	OIT3	0.83	0.47	0.00018	0.027	3.7	oncoprotein induced transcript 3	1.8

Table S2 | Multiple linear regression analysis of ICB response. Expression values of cell type specific markers were summarised by calculating their geometric mean. The resulting summarised values were used as independent variables in a multiple linear regression model aimed at explaining ICB response. The table reports the regression coefficient estimates obtained by fitting the model to all pre-treatment melanoma samples available for this study (see Supplementary methods). The probability of observing estimates different from zero merely by chance was calculated by leveraging on the null distribution of the t-test statistic ($Pr(>|t|)$). The results show that expression of *FDCSP* is significantly associated with ICB response independently of genes expressed in major immune cell types. Asterisks indicate the level of statistical significance: *** indicates p-value ≤ 0.001, ** indicates p-value ≤ 0.01, * indicates p-value ≤ 0.05.

	Cell type	Markers	Estimate	Std. Error	t value	$Pr(>|t|)$	
1	CD8+ T cells	CD8A, CD8B, TRAC	0.032	0.078	0.40	0.69	
2	CD4+ T cells	CD4, TRAC	0.057	0.110	0.52	0.60	
3	NK cells	GNLY, KLRF1	-0.089	0.092	-0.96	0.34	
4	B cells	MS4A1, CD19	0.044	0.049	0.90	0.37	
5	Macrophages	CD163, MARCO	-0.014	0.049	-0.29	0.77	
6	Dendritic cells	CD1A, CD1B, CD1E, LILRA4	-0.053	0.075	-0.70	0.48	
7	FDCSP+ cells	FDCSP	0.071	0.027	2.60	0.0092	**

Table S3 | Multiple linear regression analysis of ICB response, including plasma cell markers. expression of *FDCSP* is significantly associated with ICB response independently of genes expressed in major immune cell types, including Plasma cells. Asterisks indicate the level of statistical significance: *** indicates p-value ≤ 0.001, ** indicates p-value ≤ 0.01, * indicates p-value ≤ 0.05.

	Cell type	Markers	Estimate	Std. Error	t value	$Pr(>|t|)$	
1	CD8+ T cells	CD8A, CD8B, TRAC	0.077	0.082	0.93	0.35	
2	CD4+ T cells	CD4, TRAC	0.049	0.110	0.45	0.65	
3	NK cells	GNLY, KLRF1	-0.074	0.092	-0.80	0.42	
4	B cells	MS4A1, CD19	0.078	0.053	1.50	0.14	
5	Macrophages	CD163, MARCO	-0.019	0.049	-0.40	0.69	
6	Dendritic cells	CD1A, CD1B, CD1E, LILRA4	-0.059	0.075	-0.78	0.44	
7	Plasma cells	TNFRSF17, MZB1	-0.093	0.056	-1.70	0.098	
8	FDCSP+ cells	FDCSP	0.067	0.027	2.50	0.014	*

Table S4 | Multiple linear regression analysis of ICB response, excluding *FDCSP*. Expression of B cell markers is significantly associated with ICB response independently of CD8+ T cells and other immune cell types. However, such association becomes statistically non-significant when expression of *FDCSP* is taken into account in the multiple regression model (Table S2 and Table S3). Asterisks indicate the level of statistical significance: *** indicates p-value ≤ 0.001, ** indicates p-value ≤ 0.01, * indicates p-value ≤ 0.05.

	Cell type	Markers	Estimate	Std. Error	t value	$Pr(>|t|)$	
1	CD8+ T cells	CD8A, CD8B, TRAC	0.096	0.083	1.20	0.25	
2	CD4+ T cells	CD4, TRAC	0.044	0.110	0.40	0.69	
3	NK cells	GNLY, KLRF1	-0.045	0.093	-0.48	0.63	
4	B cells	MS4A1, CD19	0.120	0.051	2.20	0.027	*
5	Macrophages	CD163, MARCO	-0.018	0.049	-0.37	0.71	
6	Dendritic cells	CD1A, CD1B, CD1E, LILRA4	-0.073	0.076	-0.97	0.33	
7	Plasma cells	TNFRSF17, MZB1	-0.110	0.056	-1.90	0.064	

Table S5 | Top 50 genes up-regulated in *FDCSP+* fibroblasts compared to *FDCSP-* fibroblasts detected by using single-cell RNAseq data. Differential gene expression analysis comparing single-cell transcriptomes of *FDCSP+* fibroblasts versus *FDCSP-* fibroblasts obtained from lung cancer (NSCLC) and melanoma tumours (MEL). The analysis was performed using gene expression data published in (25, 27). Results from both datasets were meta-analysed using the RankProduct method (RP; see supplementary methods). The complete list is provided in a separate spreadsheet file (FDCSP_pos_vs_FDCSP_neg_fibroblasts.xlsx).

	gene	NSCLC_$\log_2(fc)$	NSCLC_pvalue	MEL_$\log_2(fc)$	MEL_pvalue	description	RP
1	CCL21	0.91	2.5E-16	12	8.5E-13	C-C motif chemokine ligand 21	1.4
2	FDCSP	1.7	3.2E-127	7.5	4.7E-17	follicular dendritic cell secreted protein	1.4
3	CCL19	1.2	7.2E-11	8.4	1.3E-08	C-C motif chemokine ligand 19	3.9
4	SLCO2B1	0.33	6.9E-10	2.9	2.3E-11	solute carrier organic anion transporter family member 2B1	10
5	TNFSF13B	0.71	7.5E-09	4.2	3.5E-06	TNF superfamily member 13b	11
6	CADM3	0.49	1.3E-15	2	9.1E-09	cell adhesion molecule 3	12
7	C3	1.8	7.3E-06	4.3	0.01	complement C3	18
8	C7	0.83	0.00043	4.7	0.00025	complement C7	19
9	CP	0.22	9E-07	3.4	2.2E-12	ceruloplasmin	20
10	UBD	0.06	2.4E-10	7	6.2E-13	ubiquitin D	25
11	ADH1B	0.48	0.0069	4.6	2.6E-06	alcohol dehydrogenase 1B (class I), beta polypeptide	32
12	PAPLN	0.51	3.3E-09	2	0.016	papilin, proteoglycan like sulfated glycoprotein	39
13	CXCL13	0.23	3.4E-06	2.4	2.3E-07	C-X-C motif chemokine ligand 13	41
14	VCAM1	0.56	0.0023	4	0.00055	vascular cell adhesion molecule 1	42
15	C10orf10	0.58	0.068	6.3	8.3E-05		42
16	CD74	0.37	0.036	6.5	7.5E-05	CD74 molecule	45
17	RBP5	0.18	4E-08	3.3	0.00011	retinol binding protein 5	46
18	CXCL14	0.77	0.0049	4.4	0.023	C-X-C motif chemokine ligand 14	54
19	PTGDS	0.25	0.22	8.9	2.8E-06	prostaglandin D2 synthase	60
20	DAAM1	0.38	0.0072	2.7	0.0001	dishevelled associated activator of morphogenesis 1	70
21	ZFP36L2	0.68	0.0014	2.1	0.014	ZFP36 ring finger protein like 2	71
22	NFIB	0.66	0.0018	1.9	0.0098	nuclear factor I B	73
23	CSF2RB	0.23	5.2E-06	1.7	3.2E-05	colony stimulating factor 2 receptor beta common subunit	74
24	RAMP3	0.02	0.043	6.4	4.7E-17	receptor activity modifying protein 3	76
25	CSMD1	0.2	0.0031	1.4	1.6E-10	CUB and Sushi multiple domains 1	79
26	IL6ST	0.54	0.022	2.6	0.001	interleukin 6 signal transducer	86
27	SLC22A3	0.04	0.0034	5.1	2.7E-08	solute carrier family 22 member 3	88
28	NFKBIA	0.45	0.1	4.7	0.0017	NFKB inhibitor alpha	88
29	C1R	0.89	0.0022	2.1	0.27	complement C1r	103
30	BIRC3	0.28	0.025	1.4	4.1E-08	baculoviral IAP repeat containing 3	105
31	SERPINB9	0.25	0.0011	3.6	0.016	serpin family B member 9	109
32	CTSH	0.08	0.08	6.4	3.6E-05	cathepsin H	110
33	IGFBP7	1.1	0.0097	1.6	0.18	insulin like growth factor binding protein 7	112
34	CTSS	0.39	5.5E-05	1.7	0.055	cathepsin S	114
35	CLK1	0.57	0.0013	1.7	0.068	CDC like kinase 1	120
36	ZBTB16	0.17	0.04	2.8	9.7E-06	zinc finger and BTB domain containing 16	120
37	RPL13	0.98	0.00015	0.35	0.24	ribosomal protein L13	124
38	LPAR1	0.31	0.00015	2.1	0.041	lysophosphatidic acid receptor 1	129
39	SOD2	0.62	0.12	3.5	0.014	superoxide dismutase 2	130
40	KLRK1	0.1	5.9E-10	0.97	0.00031	killer cell lectin like receptor K1	135
41	XIST	0.37	0.0089	2.1	0.013	X inactive specific transcript	136
42	IL33	0.29	1.3E-05	1.4	0.041	interleukin 33	138
43	IER3	0.51	0.032	2.5	0.027	immediate early response 3	139
44	HLA-DRB1	0.36	2.5E-06	1.3	0.21	major histocompatibility complex, class II, DR beta 1	139
45	CYP1B1	0.66	0.00034	1	0.23	cytochrome P450 family 1 subfamily B member 1	140
46	ARHGAP15	0.05	0.37	4	5.4E-10	Rho GTPase activating protein 15	144
47	JUN	0.92	0.026	1.9	0.17	Jun proto-oncogene, AP-1 transcription factor subunit	148
48	FKBP1A	0.36	0.05	2.1	0.0016	FKBP prolyl isomerase 1A	152
49	CLU	0.21	0.29	5.5	0.001	clusterin	154
50	IRF8	0.08	0.019	2.4	8.4E-07	interferon regulatory factor 8	154
51	CXCL12	0.26	0.12	3.6	0.0021	C-X-C motif chemokine ligand 12	158
52	C1S	0.58	0.068	2.7	0.054	complement C1s	158
53	TMEM176B	0.28	0.25	4.2	0.0017	transmembrane protein 176B	161
54	NAP1L1	0.69	0.0041	1.2	0.13	nucleosome assembly protein 1 like 1	164
55	ABHD2	0.6	2.3E-05	0.43	0.37	abhydrolase domain containing 2	168
56	KEL	0	0.88	3.3	9.4E-14	Kell metallo-endopeptidase (Kell blood group)	183
57	LAMA3	0.05	0.0031	2.1	1.4E-06	laminin subunit alpha 3	189
58	STRA6	0.08	0.038	1.6	6.2E-08	stimulated by retinoic acid 6	195
59	RBP1	0.07	0.25	4.6	0.00017	retinol binding protein 1	196
60	ABCC3	0.04	0.14	3.2	5.2E-07	ATP binding cassette subfamily C member 3	198

Table S6 | Top 50 genes down-regulated in *FDCSP*+ fibroblasts compared to *FDCSP*- fibroblasts detected by using single-cell RNAseq data. Differential gene expression analysis comparing single-cell transcriptomes of *FDCSP*+ fibroblasts versus *FDCSP*- fibroblasts obtained from lung cancer (NSCLC) and melanoma tumours (MEL). The analysis was performed using gene expression data published in (25, 27). Results from both datasets were meta-analysed using the RankProduct method (RP; see supplementary methods). The complete list is provided in a separate spreadsheet file (FDCSP_pos_vs_FDCSP_neg_fibroblasts.xlsx).

	gene	NSCLC_$\log_2(fc)$	NSCLC_pvalue	MEL_$\log_2(fc)$	MEL_pvalue	description	RP
1	FN1	-1.1	0.0024	-5.8	0.001	fibronectin 1	14226
2	CTHRC1	-0.79	0.0042	-4.9	0.0032	collagen triple helix repeat containing 1	14222
3	TIMP3	-0.87	0.019	-3.2	0.0017	TIMP metallopeptidase inhibitor 3	14218
4	COL1A2	-1.1	0.011	-3.1	0.0032	collagen type I alpha 2 chain	14212
5	COL1A1	-1.6	0.011	-3	0.0056	collagen type I alpha 1 chain	14210
6	COL5A1	-0.87	0.0061	-2.9	0.0053	collagen type V alpha 1 chain	14206
7	ARF4	-0.43	0.0095	-3.2	0.0077	ADP ribosylation factor 4	14204
8	LGALS1	-0.46	0.13	-5.8	0.00057	galectin 1	14203
9	COL5A2	-0.81	0.019	-2.9	0.0073	collagen type V alpha 2 chain	14199
10	PRSS23	-0.34	0.084	-3.8	0.0044	serine protease 23	14194
11	POSTN	-0.6	0.11	-3.6	0.021	periostin	14187
12	MMP2	-0.48	0.14	-3.1	0.011	matrix metallopeptidase 2	14180
13	COL8A1	-0.35	0.12	-3.3	0.0094	collagen type VIII alpha 1 chain	14176
14	VCAN	-0.42	0.12	-3.1	0.011	versican	14176
15	SPON2	-0.32	0.038	-2.8	0.011	spondin 2	14174
16	CYB5R3	-0.38	0.024	-2.5	0.014	cytochrome b5 reductase 3	14161
17	ACTB	-0.4	0.078	-2.2	0.0053	actin beta	14160
18	ITGB5	-0.37	0.0093	-2.2	0.0094	integrin subunit beta 5	14158
19	CAV1	-0.2	0.15	-3.9	0.0032	caveolin 1	14154
20	ITGA11	-0.3	0.035	-2.4	0.0094	integrin subunit alpha 11	14153
21	FBN1	-0.53	0.012	-2.4	0.015	fibrillin 1	14152
22	CHPF	-0.17	0.067	-2.7	0.0044	chondroitin polymerizing factor	14151
23	FKBP9	-0.27	0.087	-2.5	0.006	FKBP prolyl isomerase 9	14150
24	ECM1	-0.14	0.089	-3.4	0.0027	extracellular matrix protein 1	14145
25	ANXA5	-0.38	0.055	-2.1	0.0084	annexin A5	14145
26	LOX	-0.22	0.045	-3	0.023	lysyl oxidase	14137
27	RAB31	-0.45	0.033	-2.1	0.015	RAB31, member RAS oncogene family	14122
28	ITGB1	-0.69	0.017	-2.3	0.024	integrin subunit beta 1	14122
29	CNN2	-0.52	0.0089	-2	0.014	calponin 2	14116
30	TPM1	-0.29	0.12	-2.7	0.027	tropomyosin 1	14108
31	CTGF	-0.32	0.16	-3.2	0.05		14105
32	INHBA	-0.41	0.097	-2.3	0.024	inhibin subunit beta A	14102
33	S100A11	-0.23	0.34	-4.3	0.001	S100 calcium binding protein A11	14101
34	GPNMB	-0.14	0.16	-3.4	0.0032	glycoprotein nmb	14100
35	PXDN	-0.4	0.051	-2	0.016	peroxidasin	14099
36	THBS2	-0.47	0.091	-2.6	0.044	thrombospondin 2	14098
37	SFRP2	-0.29	0.36	-4.1	0.043	secreted frizzled related protein 2	14096
38	CAPN2	-0.15	0.076	-2.6	0.021	calpain 2	14091
39	EFEMP2	-0.12	0.13	-3.8	0.0046	EGF containing fibulin extracellular matrix protein 2	14085
40	TPM4	-0.42	0.18	-2	0.014	tropomyosin 4	14083
41	COPS8	-0.18	0.19	-2.4	0.014	COP9 signalosome subunit 8	14071
42	FHL2	-0.12	0.15	-2.8	0.0094	four and a half LIM domains 2	14065
43	CAPZB	-0.29	0.19	-1.9	0.013	capping actin protein of muscle Z-line subunit beta	14065
44	NOP10	-0.12	0.11	-2.7	0.019	NOP10 ribonucleoprotein	14062
45	CERCAM	-0.13	0.22	-3.5	0.0044	cerebral endothelial cell adhesion molecule	14061
46	FSCN1	-0.22	0.12	-1.6	0.0051	fascin actin-bundling protein 1	14057
47	NDUFB10	-0.12	0.12	-2.6	0.014	NADH:ubiquinone oxidoreductase subunit B10	14054
48	HSPG2	-0.23	0.065	-1.6	0.0098	heparan sulfate proteoglycan 2	14053
49	SULF1	-0.48	0.04	-2.2	0.052	sulfatase 1	14044
50	PLAT	-0.26	0.037	-1.9	0.027	plasminogen activator, tissue type	14042
51	SPARC	-1.1	0.0011	-1.8	0.033	secreted protein acidic and cysteine rich	14036
52	CD55	-0.15	0.11	-2.1	0.022	CD55 molecule (Cromer blood group)	14034
53	CALM2	-0.25	0.28	-2.5	0.044	calmodulin 2	14020
54	VDAC2	-0.14	0.15	-2.6	0.032	voltage dependent anion channel 2	14019
55	REXO2	-0.21	0.15	-2	0.029	RNA exonuclease 2	14019
56	HTRA1	-0.15	0.31	-2.9	0.0044	HtrA serine peptidase 1	14010
57	MFAP2	-0.13	0.22	-3.1	0.016	microfibril associated protein 2	14010
58	MFAP5	-0.12	0.22	-3.8	0.021	microfibril associated protein 5	14009
59	GSTP1	-0.19	0.27	-2.3	0.024	glutathione S-transferase pi 1	14004
60	ASPN	-0.33	0.076	-2.5	0.097	asporin	14002

Table S7 | Expression of FDCSP is an independent predictor of response to immune checkpoint blockade. FDR values used to generate Fig. 1D

	TRAC	TRBC1	TRBC2	CD8A	GZMA	GZMB	PRF1	GNLY	KLRF1	NKG7	CCL5	IFNG	CXCL9	CXCL10	CXCL11	IDO1	CLEC9A	XCR1	MS4A1	CD19	PAX5	BANK1	FDCSP
FCER2	1.5E-02	1.6E-05	5.0E-04	7.6E-08	1.2E-07	3.3E-05	3.0E-05	7.3E-05	1.3E-03	1.4E-05	1.1E-08	4.3E-10	1.1E-08	5.4E-11	9.9E-12	1.5E-06	2.7E-05	1.8E-06	1.0E+00	1.0E+00	1.0E+00	1.0E+00	3.5E-01
TCL1A	3.0E-01	1.3E-01	5.1E-01	5.3E-02	4.7E-04	3.3E-03	1.6E-03	1.2E-01	1.8E-03	1.9E-04	4.3E-03	7.8E-03	2.9E-05	3.6E-07	3.9E-02	1.0E-02	4.1E-04	5.5E-01	1.0E+00	1.0E+00	1.0E+00	1.0E+00	1.0E+00
VPREB3	2.0E-01	3.2E-01	1.9E-01	5.1E-04	1.1E-05	2.8E-03	1.9E-04	1.5E-02	1.5E-03	7.8E-04	6.6E-05	8.1E-05	6.9E-07	1.6E-08	7.4E-03	1.3E-02	2.3E-04	9.3E-02	1.0E+00	1.0E+00	1.0E+00	1.0E+00	1.0E+00
PAX5	8.4E-01	6.1E-01	5.0E-01	3.5E-02	7.5E-04	4.3E-03	1.0E-02	3.6E-01	3.6E-03	1.7E-01	8.2E-05	1.9E-02	2.0E-07	2.2E-09	1.9E-02	1.9E-02	1.8E-03	1.0E+00	1.0E+00	1.0E+00	0.0E+00	1.0E+00	1.0E+00
CLEC17A	1.5E-03	1.1E-03	1.3E-03	1.4E-05	5.6E-07	2.6E-04	2.9E-05	1.5E-03	3.9E-05	9.7E-05	8.1E-07	1.6E-07	3.7E-10	4.9E-12	6.7E-04	3.9E-05	2.2E-06	5.6E-01	4.7E-01	8.3E-01	1.9E-01	3.2E-01	3.3E-01
CR2	3.7E-07	2.0E-06	1.4E-06	5.0E-09	1.4E-11	4.0E-10	4.0E-09	1.1E-04	6.6E-06	3.6E-08	1.7E-07	6.9E-14	3.7E-17	1.0E-18	3.4E-09	1.9E-04	5.6E-06	7.8E-01	5.4E-01	7.8E-01	3.3E-02	5.3E-02	3.3E-01
MS4A1	4.0E-04	3.5E-04	4.7E-04	2.1E-06	1.1E-08	1.0E-03	4.1E-04	8.2E-03	5.7E-03	1.1E-04	1.1E-07	3.7E-07	9.0E-10	4.7E-12	1.9E-07	8.1E-02	1.2E-05	1.0E+00	0.0E+00	1.0E+00	1.0E+00	1.0E+00	1.0E+00
CD19	5.3E-02	5.4E-02	1.0E-01	2.6E-04	1.8E-06	1.0E-03	5.3E-04	2.9E-01	2.9E-04	4.7E-04	9.1E-07	1.3E-07	1.9E-09	4.9E-12	1.0E-03	1.5E-03	2.1E-04	1.0E+00	1.0E+00	0.0E+00	1.0E+00	1.0E+00	1.0E+00
BANK1	1.4E-01	1.1E-01	1.8E-01	7.0E-04	2.1E-05	7.8E-03	5.5E-03	7.5E-01	1.5E-02	8.4E-03	2.4E-05	4.4E-06	3.7E-08	6.2E-10	5.5E-03	5.4E-02	2.1E-04	1.0E+00	1.0E+00	1.0E+00	1.0E+00	0.0E+00	1.0E+00
FDCSP	6.1E-05	2.5E-06	1.4E-05	1.2E-06	5.4E-09	1.1E-04	2.0E-07	8.1E-07	1.5E-02	9.6E-07	3.7E-07	4.0E-06	9.5E-07	1.1E-08	6.6E-10	1.6E-03	4.8E-04	3.9E-05	1.0E+00	1.8E-03	5.6E-03	5.3E-04	0.0E+00

 Biasci *et al.* | A single gene expressed in fibroblastic reticular cells predicts response to immunotherapy