Title: Evidence for immediate enhancement of hippocampal memory encoding by network-
targeted theta-burst stimulation during concurrent fMRI

Abbreviated title: Immediate impact of theta-burst stimulation

Authors: Molly S. Hermiller*[1-4], Yu Fen Chen[5], Todd B. Parrish[1,5,6], and Joel L. Voss[1-4]

Affiliations:
[1] Interdepartmental Neuroscience Program, Northwestern University
[2] Department of Medical Social Sciences, Feinberg School of Medicine, Northwestern University
[3] Ken & Ruth Davee Department of Neurology, Feinberg School of Medicine, Northwestern University
[4] Department of Psychiatry and Behavioral Sciences, Feinberg School of Medicine, Northwestern
University
[5] Department of Radiology, Feinberg School of Medicine, Northwestern University
[6] Department of Biomedical Engineering, McCormick School of Engineering and Applied Science,
Northwestern University

*Correspondence to: Molly S. Hermiller (molly.hermiller@northwestern.edu)

Number of figures: 6
Number of tables: 1
Number of words, Abstract: 248
Number of words, Introduction: 649
Number of words, Discussion: 1497

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: This research was supported by R01-MH106512 and R01-MH111790 from the

National Institute of Mental Health and by T32-NS047987 and F31-NS111892 from the National Institute

of Neurological Disorders and Stroke. The content is solely the responsibility of the authors and does not

necessarily represent the official view of the National Institutes of Health. This research was supported in

part through the computational resources and staff contributions provided for Quest, the high-

performance computing facility at Northwestern University, which is jointly supported by the Office of the

Provost, the Office for Research, and Northwestern University Information Technology. Neuroimaging

was performed at the Northwestern University Center for Translational Imaging, supported by

Northwestern University Department of Radiology. We thank Rachael A. Young and Stephen

VanHaerents for contributing to data collection. We thank Plochman, Inc. for providing premium mustard.

Abstract: The hippocampus supports episodic memory via interaction with a distributed brain network. Previous experiments using network-targeted noninvasive brain stimulation have identified episodic memory enhancements and modulation of activity within the hippocampal network. However, mechanistic insights were limited because these effects were measured long after stimulation and therefore could have reflected various neuroplastic aftereffects with extended timecourses. In this experiment with human subjects of both sexes, we tested for immediate stimulation impact on encoding-related activity of the hippocampus and immediately adjacent medial-temporal cortex by delivering theta-burst transcranial magnetic stimulation (TBS) concurrent with fMRI, as an immediate impact of stimulation would suggest an influence on neural activity. We reasoned that TBS would be particularly effective for influencing the hippocampus because rhythmic neural activity in the theta band is associated with hippocampal memory processing. First, we demonstrated that it is possible to obtain robust fMRI correlates of task-related activity during concurrent TBS. We then identified immediate effects of TBS on encoding of visual scenes. Brief volleys of TBS targeting the hippocampal network increased activity of the targeted (left) hippocampus during scene encoding and increased subsequent recollection. Stimulation did not influence activity during an intermixed numerical task with no memory demand. Control conditions using beta-band and out-of-network stimulation also did not influence hippocampal activity or recollection. TBS targeting the hippocampal network therefore immediately impacted hippocampal memory processing. This suggests direct, beneficial influence of stimulation on hippocampal neural activity related to memory and supports the role of theta-band activity in human episodic memory.

Significance Statement: Can noninvasive stimulation directly impact function of indirect, deep-brain targets such as the hippocampus? We tested this by targeting an accessible region of the hippocampal network via transcranial magnetic stimulation during concurrent fMRI. We reasoned that theta-burst stimulation would be particularly effective for impacting hippocampal function, as this stimulation rhythm should resonate with the endogenous theta-nested-gamma activity prominent in hippocampus. Indeed, theta-burst stimulation targeting the hippocampal network immediately impacted hippocampal activity during encoding, improving memory formation as indicated by enhanced later recollection. Rhythm- and

58   location-control stimulation conditions had no such effects. These findings suggest a direct influence of

59   noninvasive stimulation on hippocampal neural activity and highlight that the theta-burst rhythm is

60   relatively privileged in its ability to influence hippocampal memory function.

Introduction

The hippocampus exhibits theta-band (~4-8 Hz) oscillatory neural activity that is thought to provide a temporal framework for episodic memory (Buzsaki, 2002; Lisman and Jensen, 2013; Herweg et al., 2020). Episodic memory involves hippocampal interaction with a distributed network (Squire and Zola-Morgan, 1991; Eichenbaum et al., 2007; Ranganath and Ritchey, 2012) that exhibits interregional synchrony of memory-related activity preferentially in the theta band (Fell et al., 2001; Buzsaki and Draguhn, 2004; Foster et al., 2013; Lisman and Jensen, 2013; Staudigl and Hanslmayr, 2013). Although the functional significance of hippocampal theta activity has been tested via stimulation in rodents (Shirvalkar et al., 2010; Zutshi et al., 2018), such direct functional tests present major challenges for human experimentation.

Stimulation using a theta-rhythmic pattern, as in theta-burst stimulation (TBS; high-frequency stimulation delivered in a theta rhythm), should be capable of testing the role of theta in episodic memory. This is because TBS mimics the endogenous theta rhythm thought to support hippocampal memory processing and hippocampal network synchronization, and therefore may optimally influence this network's function via activity entrainment (Buzsaki, 2002; Thut et al., 2011b; Romei et al., 2016). Direct stimulation of the hippocampus is not possible in human subjects without neurosurgery, but noninvasive stimulation can directly test network functional properties (Fox et al., 2012). Multiple experiments using transcranial magnetic stimulation (TMS) have shown that it is possible to influence episodic memory by targeting the hippocampal network (Hebscher and Voss, 2020). In those experiments, network-targeted TMS increased hippocampal network fMRI connectivity, memory-related fMRI activity, and/or improved memory performance for hours to weeks post-stimulation e.g., (Wang et al., 2014; Kim et al., 2018; Tambini et al., 2018; Freedberg et al., 2019; Hermiller et al., 2019b; Warren et al., 2019). Consistent with the hypothesized importance of theta for memory, one study found that network-targeted TBS had greater impact on memory accuracy and memory-related hippocampal fMRI connectivity than did a non-theta (20-Hz) control frequency (Hermiller et al., 2019a). However, a weakness of previous noninvasive stimulation experiments with respect to mechanistic interpretation is that these studies measured long-lasting aftereffects of stimulation (ranging from minutes to weeks), which could be mediated by a variety of indirect neuroplasticity mechanisms (Thickbroom, 2007).

Evidence for preferred influence of TBS on memory-related neural activity would require immediate

assessment of stimulation impact.

To address this, we delivered TBS to a hippocampal-network-targeted (HNT) location in the

parietal cortex during concurrent fMRI while subjects performed a memory task. We developed custom

fMRI parameters that allowed TBS as well as control-frequency (12.5 Hz) stimulation during scanning.

These scanning parameters were suitable for our focus on the effects of TBS on hippocampus and

adjacent cortex of the medial temporal lobe (MTL), as high-quality fMRI signals were obtained from these

areas.

Human subjects studied complex visual scenes that were each immediately preceded by either

HNT TBS or by one of several control stimulation conditions (Fig. 1). We hypothesized that HNT TBS

delivered in the seconds immediately preceding individual stimuli would increase fMRI activity evoked by

scenes in the hippocampus, reflecting the effects of successful entrainment of the hippocampal theta

rhythm by TBS. Based on robust evidence that increased hippocampal activity during encoding is

associated with later recollection (Paller and Wagner, 2002; Kim, 2011; Rugg et al., 2012) and previous

findings that hippocampal network-targeted stimulation enhances recollection (Hebscher and Voss,

2020), we predicted increased recollection for scenes encoded with HNT TBS. We included various

control conditions, including controls for the cognitive task (memory versus non-memory), stimulation

rhythm (TBS versus non-theta), stimulation location (HNT versus out-of-network), and hemisphere

(targeted left versus non-targeted right hippocampus). Because the experiment involved a randomized

trial-based stimulation design, immediate and selective effects of HNT TBS on hippocampal memory-

related activity would indicate that this type of stimulation impacted hippocampal neural activity, rather

than longer-lasting neuroplasticity processes that would carry over to affect other trials, and would

thereby support the role of theta in human memory formation.

Materials and Methods

Overview

Following a baseline session, subjects completed a two-day experiment in which they attempted to

remember complex visual scenes that were each immediately preceded by different stimulation

conditions, performed during fMRI scanning. The main condition of interest was TBS delivered to a

hippocampal-network targeted (HNT) location in the parietal cortex immediately before the onset of

scenes. Several control conditions were used to test specificity. Subjects also received stimulation

immediately before the presentation of numeric judgments, interleaved randomly with the scenes

throughout the task (Fig. 1A). We expected no effect of stimulation on hippocampal activity for this

condition, as hippocampal activity is generally not evoked by numeric judgments (Stark and Squire,

2001) and therefore would not increase via direct effects of stimulation on hippocampal neural activity.

Furthermore, the same scene and number conditions were administered in three control stimulation

conditions: (i) a different stimulation rhythm (beta; 12.5 Hz) (Fig. 1B) applied to the same HNT location,

(ii) TBS applied to a control location in the supplementary motor area (SMA) outside the hippocampal

network (Fig. 1C), and (iii) beta stimulation of the SMA location. None of these control conditions were

expected to influence hippocampal activity. Finally, stimulation was not delivered for a subset of scene

and number trials, providing a no-stimulation ("off") control condition. Scene and number trials with and

without stimulation were intermixed throughout scanning sessions, guarding against confounding

influences such as stimulation-induced fMRI artifact and stimulation carry-over effects across trials. All

conditions were administered in each subject using a within-subjects counterbalanced design over two

experimental sessions (Fig. 1D). As indicated above, our *a priori* hypotheses were that the HNT TBS

encoding condition would be unique among all stimulation conditions in producing: (1) increased

likelihood of later recollection during the memory test, and (2) increased activity in the left (targeted)

hippocampus during encoding of scenes that would be later recollected during the memory test.

<u>Subjects</u>

Adult human subjects passed standard MRI and TMS safety screenings (Rossi et al., 2009), reported no

present use of psychoactive drugs, and were free of known neurological and psychiatric conditions.

Datasets from 16 subjects were included in all reported analyses (8 females, 8 males; ages 20-35 years,

average age=27.6, SD=4.3). Data from two additional subjects were collected but excluded from all

reported analyses due to poor behavioral performance (overall miss rate > 50%). In addition, data

collection was attempted from three additional subjects but failed due to technical malfunction (n=2) or

attrition (n=1). Subjects gave written informed consent approved by the Northwestern University

Institutional Review Board and were paid for participation. The sample size of N=16 was chosen to

match or exceed previous experiments that demonstrated memory improvement for stimuli encoded

following short volleys of TMS (i.e., <2 s TMS immediately before stimulus onset) (Kohler et al., 2004;

Demeter et al., 2016).

<u>Baseline session</u>

Subjects completed a baseline session to determine stimulation locations and intensity prior to two

experimental sessions, performed on different days (described below).

Baseline MRI to determine stimulation locations

Resting-state fMRI and structural MRI were collected using a 3T Siemens PRISMA scanner with

a 64-channel head/neck coil. Baseline resting-state functional images were acquired using a blood-

oxygenation-level-dependent (BOLD) contrast sensitive gradient-echo echo-planar imaging (EPI)

pulse sequence (270 frames; TE 20 ms; TR 2000.0 ms; flip angle 80°; voxel resolution 1.7 mm isotropic;

70 ascending axial slices; 210x203 mm FOV; scan duration 9 min). During the resting-state scan

subjects were instructed to lie as still as possible, to keep their eyes open and focused on a fixation cross

presented in the center of the screen, and to let their minds wander. Structural images were acquired

using a T1-weighted MPRAGE sequence (176 frames; TE 1.69 ms; TR 2170 ms; TI 1100 ms; flip angle

7°; voxel resolution 1.0 mm isotropic; 256x256 mm FOV; GRAPPA acceleration of a factor of 2; scan

duration 6.36 min).

Baseline scans were submitted to resting-state fMRI connectivity analysis to determine

stimulation locations. All fMRI analyses used AFNI (Cox, 1996) and were visualized with the BrainNet

Viewer Matlab (The MathWorks, Inc. Natick, MA, USA) toolbox (Xia et al., 2013) on a smoothed Colin27

template. Anatomical scans were skull-stripped (*3dSkullStrip*) and co-registered to standardized space

using the Colin27 template (*auto_tlrc*). Preprocessing of the functional volumes included outlier

suppression (*3dDespike*), slice timing and motion correction (*3dvolreg*), and co-registration to the

anatomical scan (*align_epi_anat*). The transformations were applied simultaneously in a single

resampling step (*3dAllineate*). Motion parameters were calculated for each volume as the Euclidean

norm of the first difference of six motion estimates (three translation and three rotation). Volumes with

excessive motion (>0.2 mm), as well as the previous volume, were censored. On average, 0.42%

(SD=1.11, range=0-4.44%) of the resting-state volumes were censored. Data were spatially smoothed

using a 4-mm full-width-at-half-maximum (FWHM) isotropic Gaussian kernel (*3dmerge*) and signal

intensity was normalized by the mean of each voxel. EPI masks were created that included only voxels in

the brain that were not excluded due to instability by *3dAutomask*. Bandpass filtering (0.01-0.1 Hz),

motion censoring, and nuisance time series (estimates of motion parameters and their derivatives) were

detrended from each voxel simultaneously (*3dDeconvolve, 3dTproject*) to yield a residual time series

used in connectivity analyses.

Seed-based resting-state fMRI connectivity was used to determine subject-specific stimulation

locations used in the subsequent concurrent TMS-fMRI experimental sessions. For each subject, a 2 mm

seed in the left hippocampus (MNI: -30 -18 -18) was used in a seed-based functional connectivity

analysis (*3dTcorr1D*) to identify a left lateral parietal cortex location with robust fMRI connectivity to the

left hippocampal seed (mean z(r)=0.38, SD=0.05; average MNI: -53 -41 27). This was the stimulation

location used for hippocampal-network-targeted (HNT) stimulation (Fig. 1C). The control out-of-network

stimulation location was anatomically determined in the left supplementary motor area (SMA; average

MNI: -36 -3 67), a region outside of the targeted hippocampal network. Both the HNT and SMA locations

allowed the TMS coil to be positioned in the scanner without blocking the subjects' view of the screen.

Due to coil displacement during scanning, the actual achieved stimulation locations deviated from these

intended targets (see below and Fig. 1C).

Baseline stimulation intensity determination

TMS was delivered with a MagPro X100 stimulator using a MagPro MRi-B91 air-cooled butterfly

coil and MRI-compatible TMS setup (MagVenture A/S, Farum, Denmark). Resting motor threshold (RMT)

was found during the baseline session in order to determine the stimulation intensity used during the

experimental sessions (see below). Subjects sat at the entrance of the MRI bore with their arms resting

comfortably during RMT determination. The MRi-B91 TMS coil was used to determine RMT as the

minimum percentage of stimulator output (% SO) necessary to generate a visible contraction of the right

thumb (*abductor pollicis brevis)* for five out of ten consecutive single pulses. Pulses were biphasic, as

were pulses delivered during experimental sessions. RMT values ranged between 45.0-85.0% SO

(mean=61.6, SD=10.8).

<u>Experimental Sessions</u>

Following the baseline session, subjects returned for two experimental TMS/fMRI sessions on separate

days to complete a 2x2 crossover design. One stimulation pattern (TBS or beta) was used during each

session. Within each session, there were two study phases that differed in stimulation location (HNT or

SMA). The order of these conditions (TBS or beta session; HNT-then-SMA or SMA-then-HNT within

each session) was counterbalanced across subjects. Memory for scenes encoded during both study

phases was tested at the end of the experimental session, after subjects finished MRI scanning.

Experiment Design

There were two study phases during each of the two experimental sessions. Each study phase

lasted ~70 min and comprised 144 trials (288 trials total for the session). Each trial began with a white

fixation cross presented in the center of the screen, during which ~ 2 s stimulation was delivered (see

TMS/fMRI acquisition methods for exact timing). Immediately following stimulation, a visual stimulus was

presented for 2 s. The study item was followed by a white fixation cross that remained on the screen until

the next trial for a randomly varied duration between 11-19.5 s. Different visual stimuli were presented

during each study phase. Complex visual scenes (50% of trials; 144 scenes total for the session) were

randomly intermixed with numeric stimuli (50% of trials; 144 numbers total for the session). During the

scene presentation, subjects were instructed to imagine visiting the depicted location and to rate via

button press whether they would like to visit the location (right hand button) or not (left hand button).

Scenes were chosen from the SUN397 dataset (Xiao et al., 2016) based on the following criteria:

complex outdoor natural scenes (e.g., mountains, beaches, forests, waterfalls, deserts) without

prominent humans, animals, or man-made objects; color image; image did not include text. Subjects

were told that memory would be tested for all scenes following the study phase (i.e., intentional

encoding). Numeric stimuli were randomly selected from the integers 1-864 and presented in white font

for 2 s. Subjects used a button response to indicate if the number was even (right hand button) or odd

(left hand button). Visual scene stimuli were randomly assigned to either a study trial stimulation

condition or to serve as a lure during memory testing (see below) for each subject. Stimuli were

presented in the center of an MRI-compatible LCD screen (Nordic Neuro Lab, Bergen, Norway)

positioned at the subjects' feet, on a gray background, viewed via a mirror attached to the head coil.

Responses were made with hand-held fiber optic button boxes (Current Designs, Inc., Philadelphia, PA,

USA). Subjects were told that they could make their responses during the white fixation cross following

each stimulus and that response times were not important (i.e., self-paced responses).

Stimulation was delivered for ~2 s (see TMS/fMRI acquisition methods for exact timing)

immediately preceding stimulus onset for 66% of scene and numeric trials (i.e., stimulation presence on),

with no stimulation for the remaining trials (i.e., stimulation presence off). Long inter-trial intervals (11-

19.5 s) were used to reduce stimulation carry-over effects (Huang et al., 2005). During one study phase

stimulation targeted the hippocampal network via left parietal cortex (HNT), and during the other study

phase stimulation targeted the SMA. Subjects came out of the scanner for a break of ~10 min between

study phases for TMS coil repositioning. The order of these conditions was counterbalanced across

subjects.

Prior to getting in the scanner for the study phases, MRI-navigated TMS software (Localite

GmbH, St. Augustin, Germany) was used to physically mark the individualized stimulation locations on

the participant's scalp. A conformable MRI-compatible marker was affixed to the scalp at the intended

stimulation location (12.7 mm x 12.7 mm re-sealable plastic bag filled with yellow-mustard MRI contrast

agent; Plochman, Inc., Manteno, IL, USA). The markers were used to position the TMS coil against the

subject's head in the scanner and coil location was recorded via MRI anatomical scans during each

study phase (see below).

One experimental session used TBS and the other used beta TMS, administered in

counterbalanced order across subjects. For both stimulation patterns, 30 TMS pulses were delivered

during the 2 s prior to stimulus onset per trial, delivered at the same intensity for each subject (80%

RMT). For TBS, pulses were delivered as 50 Hz triplets at 5 Hz. For beta stimulation, pulses were

delivered individually at 12.5 Hz. TMS pulses were synchronized with the MRI scan and with visual

stimulus onset (see below). To acclimate subjects to the stimulation protocols and to ensure that

stimulation did not cause scalp/facial twitches, a train of stimulation was applied once the subject was

positioned inside the scanner and the TMS coil was positioned at the targeted location before scanning

in each study phase. Stimulation intensity was lowered during one or both sessions due to technical

limitations for 5 subjects. On average, TBS was delivered at 78.8% RMT (SD=1.9, range=75.0-80.0) and

beta stimulation was delivered at 78.5% RMT (SD=2.2, range=74.1-80.0). The experimental sessions

were scheduled at least two days apart, with an average of 27 days between sessions (range=3-84

days). For 1 subject, a session was discarded due to technical difficulties and the subject returned for a

third "replacement" session. The replacement session was performed with the same location and pattern

order as the discarded session, but with different visual stimuli. There were no relationships between the

delay from the first to second sessions and the effect of the TBS and beta stimulation conditions on

recollection memory reported below, as indicated by non-significant across-subject Pearson correlations

of these variables (P's>0.14).

At the end of each experimental session, memory was tested for the scenes that were presented

during both of the study phases for that session (one study phase with HNT stimulation and one with

SMA stimulation). After completing both study phases, subjects rested out of the scanner for ~15 min

before taking the memory test, which was not scanned. The 144 scenes presented during the study

phases were presented one at a time intermixed randomly with 144 novel lures that were not presented

during study phases, in randomized order. Subjects were instructed to respond with (i) "Remember" if

they specifically recalled details about the scene, (ii) "Familiar" if they recognized the scene but could not

recollect details, and (iii) "New" if the scene was a lure (Yonelinas, 2002; Eichenbaum et al., 2007).

Assessment of remember versus familiar responses was a key aspect of the study design, as

recollection is highly related to increased hippocampal activity during encoding (Paller and Wagner,

2002; Kim, 2011; Rugg et al., 2012) and our previous experiments using hippocampal network-targeted

stimulation have identified increased recollection rather than nonspecific aspects of memory performance

(Hebscher and Voss, 2020). Thus, we hypothesized that the effects of HNT TBS would be specific to

remember responses. Trials were self paced, with the scene remaining on the screen until a response

was registered. The duration of the test phases was 20.8 min on average (range=14-29, SD=4.48).

Simultaneous TMS/fMRI acquisition

MRI was performed during study phases using a 3T Siemens PRISMA scanner with a single-

channel transmitter/receiver head coil. Fast low-angle shot (FLASH) anatomical scans were collected

between study phases to localize the actual location of the TMS coil relative to markers placed on the

scalp, including a T1 sagittal (50 slices; TE 2.42 ms; TR 311.0 ms; flip angle 80°; 1.0 mm in-plane

resolution; 4.0 mm thick sagittal slices with 0 mm gap; 50% phase oversampling; 256x256 mm FOV;

scan duration 44 sec) and a T1 oblique axial (40 slices; TE 2.42 ms; TR 249.0 ms; flip angle 80°; 1.0 mm

in-plane resolution; 4.0 mm thick axial slices with 0 mm gap; 60% phase oversampling; 256x256 mm

FOV; scan duration 36 sec). These anatomical scans were later used to localize the TMS coil targeting

displacement (see below).

We developed two fMRI scan sequences to interface with TMS pulses for the TBS and beta-

patterned stimulation conditions. Task-based functional images were acquired using a BOLD contrast

sensitive gradient echo EPI pulse sequence that contained custom programmed temporal gaps

interleaved between slice acquisitions. Rather than delivering stimulation during slice acquisition, which

causes TMS-induced artifact that requires volumes to be discarded (Bestmann et al., 2008; Siebner et

al., 2009), TMS was delivered between MRI slice acquisitions during the inserted temporal gaps. This

TMS-fMRI method did not cause artifact beyond that associated with the physical presence of the TMS

coil, which produces stable artifact near the coil (see Results).

For both stimulation patterns, 30 TMS pulses were delivered during the imaging volume

immediately prior to visual stimulus onset for conditions that involved stimulation. Pulses were delivered

over a duration of 2000 ms in the TBS condition (Fig. 2A) and 2400 ms in the beta-patterned stimulation

condition (Fig. 2B), for a total of 5760 pulses aggregate over the entire experimental session. For TBS,

107-ms temporal gaps were inserted after every two EPI slices (93 ms). During this temporal gap, a 50

Hz triplet burst (pulse every 20 ms) was delivered, with one triplet burst delivered every 200 ms during

such temporal gaps (665 frames; TE 20 ms; TR 2230.0 ms; 2442 Hz/pixel bandwidth; flip angle 90°;

voxel resolution 3.0 mm isotropic; 22 interleaved 3.0 mm thick axial slices angled to AC-PC alignment

and centered on the longitudinal axis of the temporal lobes; 50% phase oversampling in the phase-

encoding direction; 192x192 mm FOV; scan duration 24.83 min; 72 stimulated trials per scan) (Fig. 2A).

For beta-patterned stimulation, 34-ms temporal gaps were inserted after each slice (46 ms), in which a

single TMS pulse could be delivered, such that one pulse was delivered every 80 ms during such

temporal gaps (270 frames; TE 20 ms; TR 2440.0 ms; 2442 Hz/pixel bandwidth; flip angle 90°; voxel

resolution 3. mm isotropic; 30 interleaved 3.0 mm thick axial slices angled to AC-PC alignment and

centered on the longitudinal axis of the temporal lobes; 50% phase oversampling in the phase-encoding

direction; 192x192 mm FOV; scan duration 28.38 min; 72 stimulated trials per scan) (Fig. 2B). The scans

were programmed such that the last TMS pulse would occur at the end of the TR (i.e., all pulses during

the final 2000 ms of the 2230 ms TR for the TBS scan and during the last 2400 ms of the 2440 ms TR of

the beta-patterned stimulation scan). Phase oversampling in the phase-encoding direction was used in

both scans to shift any Nyquist ghosting induced by the presence of the TMS coil outside the brain.

The limited coverage in both the theta- and beta-patterned task-based scans precluded whole-

brain imaging but did adequately cover MTL regions of interest when the imaging volume was centered

at the MTL (Fig. 2AB). Notably regions imaged directly under/around the TMS coil typically exhibit

irreparable TMS-induced artifacts (Bestmann et al., 2008; Siebner et al., 2009), but the parietal cortex

and SMA stimulation locations did not fall within our limited coverage. Our hypothesis-driven regions of

interest were instead in downstream regions distant from the stimulation location. We confirmed image

quality by subject-level visual inspection, as well as validating signal quality at the group-level (see

Results). Two task-based scans using the same parameters were acquired per the two study phases

during each experimental session (~70 min per phase, 144 trials per phase). Participants wore air-

conduction earplugs during the scans to attenuate both scanner and TMS noise.

A PC in the MRI control room received transistor-transistor logic (TTL) pulses from the MR

scanner and, based on the experiment code, then sent TTL pulses to the TMS device to trigger

stimulation at appropriate times. TTL pulses were sent per EPI slice acquisition to the experiment control

PC, which in turn triggered the TMS device to deliver the programmed stimulation sequence. Thus,

stimulation delivery was trial-specific and time-locked to the slice-based MR trigger. During the study

phases, experiment events (e.g., pulse signals from the MRI and TMS, stimulator settings, participant

responses, task stimuli, etc.) were monitored and recorded in output files created by Presentation

(Neurobehavioral Systems, Inc. Berkeley, CA, USA), MagVenture (MagVenture A/S, Farum, Denmark),

and LabChart (ADInstruments, Inc. Colorado Springs, CO, USA), as well as by the experimenter in the

MRI control room. These records were used by the experimenter to update copies of raw output files with

trial-specific deviations (i.e., trials were discarded if only part of the stimulation train was delivered due to

coil over-heating; records reflected that the trial condition changed from "ON" to "OFF" if stimulation

failed entirely during the 2-s pre-stimulus period).

TMS coil displacement during fMRI

We used FLASH anatomical MRI scans (see above) collected before and after the study phase

fMRI scans to evaluate the actual location of the TMS coil during the experiment relative to its intended

location to account for possible displacement. The scans were uploaded into the MRI-navigated TMS

software (Localite GmbH, St. Augustin, Germany) and aligned to the subject's high-resolution anatomical

scan collected during the baseline session. We utilized contrast-agent markers on the TMS coil (vitamin

E capsules and oil-filled tubing) and on the scalp (mustard packets, described above) to identify the

position, orientation, and rotation of the TMS coil inside the scanner (i.e., 4D Matrix of coordinates)

relative to the target that was identified based on resting-state fMRI for each subject (see above). The

matrices were transformed with a displacement vector to estimate the cortical coordinates directly under

the TMS coil. The across-participant mean MNI coordinates of the achieved HNT location was -50, -52,

32 (SD=4.8, 6.7, 6.8) for the TBS session and -49, -51, 33 (SD=4.8, 7.7, 7.3) for the beta stimulation

session. The SMA location was -31, -16, 66 (SD=5.4, 9.0, 3.7) for the TBS session and -32, -13, 63

(SD=6.0, 9.3, 5.5) for the beta stimulation session. For each participant and condition, the deviation in

achieved versus intended stimulation locations was calculated as the Euclidean distance. For the TBS

and beta-patterned sessions, there was an average deviation of 8.47 mm (SD=4.04) for the HNT

condition and a deviation of 9.02 mm (SD=4.12) for the SMA condition. The amount of deviation did not

significantly vary between the two locations (P>0.7). There were no relationships between location

deviation across sessions (i.e., TBS versus beta) and the differential effect of stimulation on the

recollection memory measures described below, as indicated by non-significant across-subject Pearson

correlations of these variables (P's>0.36).

To confirm that the HNT and SMA conditions differentially targeted the hippocampal network as

intended despite the in-scanner coil displacement, we analyzed resting-state fMRI connectivity of the

achieved stimulation location (considering displacement) with the hippocampal network. Using the high-

resolution resting-state fMRI scan collected at baseline, we calculated the hippocampal network as

regions with robust connectivity to the left hippocampus (defined as 2-mm spherical segments centered

at MNI coordinates: -23, -10, -21; -26, -14, -20; -30, -18, -18; -31 -22 -14; -30, -26, -12; see below for

description of these locations in the task-based fMRI analysis). The hippocampal functional connectivity

map for each subject was created by correlating (Pearson's *r*) the spatially averaged time series of these

hippocampal coordinates with every voxel's time series (*3dTcorr*). A Fisher's *z* transformation was

applied to yield a normally distributed correlation map for each subject (*3dcalc*). Group-level voxel-wise

analysis of these connectivity maps using one-sample one-tailed *t*-tests (*3dttest++*) identified clusters of

contiguous voxels with robust connectivity to the hippocampal seed mask (300+ voxels with *z(r)*

significantly greater than 0; t-threshold=5.2; P<0.0001). These clusters were saved as a hippocampal

network mask (*3dclust*; 6,695 voxels total). This network included locations expected based on the

previous literature on typical hippocampal fMRI connectivity (Kahn et al., 2008), including medial and

lateral temporal, lateral parietal, ventral medial prefrontal cortex areas. For each subject, we then

assessed resting-state connectivity between the achieved stimulation location to every voxel in the

hippocampal network mask (*3dTcorr*) and spatially averaged the correlation value to obtain one overall

network connectivity value for each stimulation condition for every subject (*3dmaskave*). A 2x2 repeated

measures ANOVA testing the effects of stimulation location and pattern on connectivity to the

hippocampal network indicated significant variation by location ($F_{1,12}$=4.83, P=0.04, η^2_p=0.24), such that

connectivity was significantly greater for the HNT locations (mean=0.15, SD=0.12) relative to the SMA

locations (mean=0.06, SD=0.07) (Fig. 1C). Thus, differential targeting of the hippocampal network by the

HNT and SMA locations was successful despite coil displacement during fMRI scanning.

To confirm that the SMA location stimulated belonged to a distinct network and therefore was a

suitable out-of-network control condition, we assessed the connectivity of the achieved SMA locations

using the same connectivity analysis method that was used to define the left hippocampal network but
with the achieved SMA location in each subject used as the connectivity seed rather than the left
hippocampus (300+ voxels with $z(r)$ significantly greater than 0; t-threshold=5.2; P<0.0001; 30,275 total
voxels identified). As expected, this network included canonical somatomotor regions including the
majority of M1/S1, supplementary motor and somatosensory areas. The hippocampal network and SMA
network had minimal overlap (7 voxels overlapping between networks out of 36,963 total voxels in both
networks), confirming that the SMA location was suitable for its intended purpose as an out-of-network
control.

Subject-level task-based fMRI processing

Anatomical scans were skull-stripped (*3dSkullStrip*) and co-registered to the Colin27 template
(*auto_tlrc*). Preprocessing of the functional volumes included outlier suppression (*3dDespike*), slice
timing and motion correction (*3dvolreg*), and co-registration to the anatomical scan (*align_epi_anat*). The
transformations were applied simultaneously in a single resampling step (*3dAllineate*). Motion
parameters were calculated for each volume as the Euclidean norm of the first difference of six motion
estimates (three translation and three rotation). Volumes with excessive motion (>0.3 mm), as well as the
previous volume, were flagged for censoring during the regression analyses, which is a typical threshold
for task-based fMRI analysis. On average, 2.8% (SD=5.0) of the TBS HNT condition, 3.6% (SD=6.1) of
the TBS SMA condition, 5.0% (SD=8.0) of the beta HNT condition, and 5.9% (SD=10.2) of the beta SMA
condition time series were motion censored. There was no significant difference in the amount of
censoring across conditions (all pairwise comparison *Ps*>0.10). Data were spatially smoothed using a 6-
mm full-width-at-half-maximum (FWHM) isotropic Gaussian kernel (*3dmerge*) and signal intensity was
normalized by the mean of each voxel. Task-based masks were created that consisted only of voxels in
the brain with stable signal across the scanning sessions (*3dAutomask*).

Two general linear models (GLMs) incorporating hemodynamic response deconvolution were
applied to the preprocessed data to estimate voxel-wise event-related activity regression coefficients for
each trial type, separately for each stimulation condition (HNT TBS, SMA TBS, HNT beta, and SMA
beta) (*3dDeconvolve*). GLMs were constructed separately per condition because differences in the scan

parameters required for TBS versus beta stimulation precluded concatenation of all conditions into one

GLM. In each GLM, trials were separated based on experiment condition (scenes with TMS ON, scenes

with TMS OFF, numbers with TMS ON, and numbers with TMS OFF). In a second GLM, the scene trials

were further sorted by subsequent memory performance (Remember, Familiar, or New responses during

the test). Time points with motion spikes and time series outliers were censored. Polynomial trends and

motion estimates and their derivatives were included as nuisance regressors of no interest. Condition-

specific activity estimates used the duration-modulated gamma function. Each condition of interest (HNT

TBS, SMA TBS, HNT beta, and SMA beta) was modeled, with each event beginning at the stimulus

(scene or number) onset. Restricted Maximum Likelihood (REML) estimation methods were used to

generate voxel-wise parameter estimates and measures of variability for each trial type for each

stimulation condition (*3dREMLfit*). Parameter estimates from each subject were later analyzed at the

group-level (see below)

<u>Data analysis</u>

Memory performance

Performance on the scene recognition test was computed as the rate of hits ("Remember" and

"Familiar" responses for studied scenes) and correct rejections ("New" response for novel lures) for each

subject separately for every stimulation condition. To evaluate stimulation effects on hippocampal-

dependent recollection, we calculated the proportion of hits that were recollected ("Remember"

responses) for every stimulation condition. Accuracy was also calculated, as described in the Results

section.

Group-level task-based fMRI analyses

Voxel-wise analyses were performed in order to confirm that our scan parameters provided

sensitivity to expected fMRI correlates of cognitive processing (i.e., scenes but not numbers should

evoke activity in parahippocampal, fusiform, and occipital regions (Stern et al., 1996; Stark and Squire,

2001) and stimulation sensations (i.e., sound emitted by stimulation should evoke activity in auditory

cortex). The first contrast compared BOLD activity evoked by task stimuli (scenes versus numbers)

regardless of stimulation presence, location, or pattern; and the second contrasted activation due to

stimulation presence (on versus off) regardless of stimuli type or stimulation location or pattern. Subject-

level GLMs were used to estimate voxel-wise event-related activity regression coefficients for each trial

type (i.e., scenes, numbers, TMS ON, and TMS OFF) and REML estimation methods were used to

generate voxel-wise parameter estimates and measures of variability for each subject (*3dDeconvolve,*

3dREMLfit; see above). GLM maps were analyzed at the group-level using generalized least squares

with a local estimate of random effects variance (*3dMEMA*) to identify regions of significant difference

($P<0.001$, t-threshold=4.07) for scenes versus numbers and for stimulation on versus off. Data from

theta-patterned and beta sessions were analyzed separately due to the differences in scan parameters.

To test whether differences in parameters between the TBS and beta stimulation scans did not

significantly affect the signal quality, the Signal-to-Fluctuation-Noise Ratio (SFNR) summary value

(Friedman and Glover, 2006) was assessed and compared between the two fMRI sequences. The mean

signal was divided by the standard deviation of the residuals (*3dTstat, 3dcalc)* and then averaged within

the limited task coverage mask over the whole session (*3dmaskave*) to yield one TSNR value per

stimulation condition per subject.

To measure the effect of stimulation on activity during the memory task, performance on the

retrieval task was used to back-sort fMRI data to analyze the effects of stimulation on encoding-related

activity that predicted subsequent recollection (trials that were later endorsed with "Remember"

responses for the main analysis, and trials later endorsed with "Familiar" and "New responses in a

control analysis). Subject-level GLMs estimated voxel-wise event-related activity regression coefficients

for each trial type (i.e., remembered scenes with TMS ON, remembered scenes with TMS OFF, scenes

not-recollected with TMS ON, scenes not-recollected with TMS OFF, numbers with TMS ON, and

numbers with TMS OFF) for each stimulation condition (*3dDeconvolve, 3dREMLfit*; see above*). For

scenes with "Remember" responses, there were 16 trials on average (range=6-33) per condition (trial

counts did not vary by condition; Ps>0.2). Trials with remembered scenes with TMS OFF were collapsed

across all study sessions to create the "combined off" condition (average number of trials=27, range=15-

46). For scenes with non-recollected responses ("Familiar" and "New" responses), there were 33 trials on

average per stimulation condition (range=16-64) and a mean of 17 trials in the "combined off" condition

(range=8-26). Similarly, activity for numeric judgment trials was estimated, but without respect to test

performance (i.e., all trials), separately for each stimulation condition, and collapsed across all study

sessions for the "combined off" condition.

The influence of stimulation conditions on fMRI activity estimates for studied scenes and numeric

judgment trials were tested at the group level using a region-of-interest (ROI) approach. 6-mm radius

spherical regions of interest (ROIs) were defined along the hippocampal longitudinal axis in each

hemisphere. The middle ROI in the left hemisphere was placed in the body of the hippocampus,

centered at the coordinate that was targeted via its connectivity with parietal cortex as measured during

the baseline session resting-state fMRI scan (centroid MNI coordinate: -30 -18 -18). However, because

the coil was displaced, this location was not accurately targeted and was instead used simply because of

its anatomical location in the body of the hippocampus. Two spheres were placed anterior to this location

(centroid MNI coordinates: -23 -10 -21; -26 -14 -20), and two posterior (centroid MNI coordinates: -31 -22

-14; -30, -26, -12), in 4mm increments along the longitudinal axis. These coordinates were mirrored into

the right hemisphere (centroid MNI coordinates: 23 -10 -21; 26 -24 -20; 30 -18 -18; 31 -22 -14; 30 -26 -

12. The two most anterior spheres encompassed the head of the hippocampus and the middle and two

posterior spheres fell in the body of the hippocampus. These spherical ROIs encompassed hippocampal

as well as small portions of immediately adjacent entorhinal/perirhinal and parahippocampal cortex,

which was necessary due to the relatively large voxel size. Spherical ROIs were used in lieu of more

anatomically precise methods (e.g., hippocampal subfield identification) due to the limited spatial

resolution imposed by the scanning parameters that are possible with the single-channel MRI head coil,

which was necessary to accommodate the TMS coil. The hippocampal tail was likewise not measured,

as very few voxels of this size can capture the hippocampal tail without also including substantial

portions of adjacent white matter and ventricle.

The primary analysis collapsed these ROIs within each hemisphere to create separate left and

right aggregate hippocampus ROIs, with the main hypothesis being that HNT TBS would increase

activity evoked by scenes only in the left hippocampus. A follow-up exploratory analysis used rmANOVA

to determine whether stimulation conditions had differential impact along the longitudinal axis. The goal

was to identify locations along the hippocampal long axis that responded to stimulation conditions, taking

into account potential functional distinctions along the anterior-posterior axis (Aggleton and Brown, 1999;

Ranganath and Ritchey, 2012; Poppenk et al., 2013) as in our previous experiments investigating the

variation in effects of TMS on hippocampal activity along its longitudinal axis (Wang et al., 2014;

Nilakantan et al., 2019).

Voxel-wise analysis with an intentionally liberal threshold was used to evaluate whether there was

any evidence for effects of stimulation in areas other than in the hippocampal ROIs, which was not

expected. This analysis compared activity estimates for scenes with stimulation that were later

recollected between stimulation conditions. Each subject's voxel-wise regression coefficients for

remembered scenes with TMS ON for every stimulation condition (HNT TBS, SMA TBS, HNT beta, SMA

beta) was entered into repeated measures ANOVA (*3dANOVA3*) to assess voxels for a significant

interaction between stimulation location and pattern. Clusters of significant interaction were identified

using a liberal threshold (two-tailed $P<0.05$ voxel-wise threshold, F-statistic=4.54, >40 contiguous supra-

threshold voxels) (*3dClust*).

<u>Statistical Analysis</u>

Statistical analysis was performed using AFNI and Matlab. Group-level analysis of multiple

conditions was performed using repeated-measures analysis of variance (rmANOVA), with partial eta

squared (η^2_p) reported as the effect size. The within-subject factors corresponding to each test are

provided in the Results. Greenhouse-Geisser correction was applied if the assumption of sphericity was

not met (significant Mauchly's test at $P<0.05$), with the corrected p-values and degrees of freedom

reported ($F_{(GG)}$). Follow-up tests avoided making all possible pairwise comparisons among the many

stimulation control conditions used in the experiment by focusing on the *a priori* hypothesis that the HNT

TBS condition would be unique among all stimulation conditions in that it alone would increase

hippocampal fMRI activity and recollection relative to all other control conditions. HNT TBS ON was

therefore the condition of interest whereas all other conditions were controls. For these follow-up

comparisons, we tested whether each individual stimulation condition was significantly greater than the

aggregate of all control conditions (other than the individual stimulation condition being tested) using

one-tailed Students t-tests (t) or Wilcoxon signed rank tests (z) if the assumption of normality was

violated (significant Shapiro-Wilk test at P<0.05), and then corrected for the number of individual

stimulation conditions that were tested using the Bonferroni method. Bonferroni corrected P values are

indicated as "P_B" throughout the manuscript. Effect size measures were calculated as Cohen's *d* (*d*) for

the t-tests and as rank-biserial correlation (*r*) for the Wilcoxon tests. This follow-up assessment method

therefore indicated whether any stimulation condition significantly increased hippocampal fMRI activity or

recollection relative to the control conditions, and we hypothesized that this would only be the case for

HNT TBS ON. For comprehensiveness, an exploratory analysis of all possible pairwise comparisons was

used to assess whether there was evidence that any condition increased fMRI activity or recollection

relative to any other condition using one-tailed Students t-tests (t) or Wilcoxon signed rank tests (z) if the

assumption of normality was violated (significant Shapiro-Wilk test at P<0.05), with no correction for

multiple comparisons and no reporting of effect sizes for the exploratory analysis.

Code and Data Accessibility

Raw data are freely available on the Northwestern University Neuroimaging Data Archive

(https://nunda.northwestern.edu/). Dataset identifiers will be provided with publication to permit

unrestricted access to raw data. Custom code and scripts to replicate analyses will also be available via

this archive.

Results

Validation of concurrent TMS-fMRI

The fMRI scanning methods used for TBS and beta stimulation differed in a number of critical

parameters, including parameters of the scan sequence as well as the timing of scan acquisition relative

to interleaved TMS pulses (Fig. 2AB). To test whether these differences affected signal quality, the

Signal-to-Fluctuation-Noise Ratio (SFNR) summary value (Friedman and Glover, 2006) was calculated

for each subject for the TBS and beta stimulation sessions. We expected SFNR values to be ~120, as

this was the approximate SFNR value obtained when we performed the same scans on an additional

subject who had the TMS coil at the same approximate locations but without any pulses delivered

(average SNFR across both scans = 120.3). Furthermore, another group using the same scanner and

head coil models as in this experiment reported similar SFNR values for their interleaved TMS/fMRI

scans (Moisa et al., 2009). The average SFNR value for the TBS sessions was 122.5 (SD=18.3,

range=94.8-160.2) and 123.4 for the beta sessions (SD=14.2, range=92.2-144.8), with no significant

difference between sessions (P=0.75). Therefore, scan stability and thus, sensitivity, did not vary by scan

sequence or stimulation pattern.

We performed two voxel-wise fMRI analyses to confirm expected neural activity correlates of

cognitive processing within the task. That is, scenes but not numbers should evoke activity in

parahippocampal, fusiform, and occipital regions (Stern et al., 1996; Stark and Squire, 2001) and

stimulation sensations such as TMS sounds should evoke activity in auditory cortex. The contrast of fMRI

activity evoked by scenes versus numbers, calculated across stimulation presence (on and off), and

location (HNT and SMA) identified significantly greater activity (P<0.001, t-threshold=4.07) of bilateral

occipital, fusiform, and posterior parahippocampal cortex as well as hippocampus for both stimulation

patterns (Fig. 3A). Contrasts between TBS and beta stimulation identified no voxels with significant

differences even at a liberal threshold (P<0.01 uncorrected). The contrast of stimulation "on" versus "off",

calculated across stimulation location (HNT and SMA), identified significantly greater activity for "on"

(P<0.001, t-threshold=4.07) in bilateral auditory cortex for both TBS and beta stimulation (Fig. 3B).

Again, the direct contrast identified no voxels with significantly different activity for TBS versus beta

stimulation at a liberal threshold (P<0.01 uncorrected). Notably, although auditory-related activity due to

stimulation was identified robustly, the limited imaging volume did not permit identification of likely

somatosensory activation. Collectively, these analyses indicate that fMRI sensitivity was sufficient for

identifying typical neural signals of scene viewing and auditory stimulation despite concurrent TMS and

that there was no obvious variation in sensitivity for TBS versus beta scan parameters.

<u>Effects of stimulation on memory encoding</u>

Subjects demonstrated above-chance discrimination of old/studied scenes from new scenes

overall across all stimulation conditions and irrespective of whether old scenes were endorsed with

"Remember" or "Familiar" responses (mean d'=0.81, SD=0.35; t_{15}=9.15, P<0.001 vs. chance value of 0).

For all stimulation conditions, the proportion of old scenes endorsed with "Remember" responses

(mean=0.30, SD=0.10) was greater than the proportion of new scenes endorsed with "Remember"

responses (mean=0.07, SD=0.05; t_{15}=21.18, P<0.001, d=2.29). Likewise, the proportion of old scenes

endorsed with "Familiar" responses (mean 0.34, SD=0.12) was greater than the proportion of new

scenes endorsed with "Familiar" responses (mean=0.28, SD=0.13; t_{15}=6.26, P<0.001, d=0.97). As

expected, "Remember" responses were significantly more likely to be hits to old items (mean=0.84,

SD=0.10) than were "Familiar" responses (mean=0.59, SD=0.10; t_{15}=6.94, P<0.001, d=1.71). Thus,

subjects performed the recognition task (Fig. 4A) accurately and the "Remember" responses were

characteristic of high-confidence, high-accuracy recognition memory, as has been linked to the

hippocampus in multiple previous experiments (Ranganath et al., 2004).

Our *a priori* hypothesis was that HNT TBS during encoding would increase subsequent

recollection of scenes, as quantified as the proportion of hits endorsed with "Remember" responses,

relative to all control stimulation conditions. Because of their different rhythms, TBS and beta could

produce differential effects on sensory (auditory and somatosensory) entrainment of brain activity

unrelated to the hypothesized effects of stimulation on hippocampal activity, and they were furthermore

administered during different experimental sessions/days (Fig. 1). To test for their differential effects on

recollection while accounting for these nonspecific differences, we compared the effects of TBS when

applied targeting the hippocampal network (HNT) relative to the control SMA location versus the effects

of beta when applied targeting HNT relative to SMA. We submitted the HNT minus SMA difference

scores for the proportion of recollected hits to 2x2 rmANOVA with factors TMS presence (on, off) and

rhythm (TBS, beta). There was a main effect of TMS presence ($F_{1,15}$=5.13, P=0.04, η^2_p=0.23) qualified by

significant interaction between TMS presence and rhythm ($F_{1,15}$=6.53, P=0.02, η^2_p=0.31). As indicated in

Figure 4B, TBS increased recollection when applied to HNT versus SMA to a greater extent than did

beta, which had almost no differential effect when applied to HNT versus SMA. Essentially the same

pattern was evident when performing this comparison without subtracting the HNT and SMA conditions.

Recollection varied significantly by TMS presence (on versus off), rhythm (TBS versus beta), and

location (HNT versus SMA) (3-way rmANOVA interaction $F_{1,11}$=6.63, P=0.02, η^2_p=0.44). Thus, TBS and

beta rhythms had different impacts on recollection when applied targeting the hippocampal network

versus the control network.

The follow-up test followed our *a priori* hypothesis that HNT TBS would be the only stimulation

condition to increase recollection relative to other conditions. We tested whether the proportion of

recollection hits was greater for HNT TBS versus the average of all control conditions (HNT TBS OFF,

SMA TBS ON, SMA TBS OFF, HNT BETA ON, HNT BETA OFF, SMA BETA ON, SMA BETA OFF),

and, for comparison, tested whether recollection hits were significantly greater for each of the control

conditions versus the average of all other control conditions. Consistent with our hypothesis, HNT TBS

was the only stimulation condition with significantly greater recollection relative to other conditions

(t_{15}=3.59, P=0.001, P_B<0.01 corrected for 8 tests, d=0.90; P_B = 1 for each other control condition versus

the average of its respective other controls) (Fig. 4C).

An exploratory analysis of all possible pairwise comparisons among stimulation conditions was

also performed, although the number of comparisons is high (56 total) and this analysis does not account

for possible nonspecific differences among conditions that are relatively controlled in the primary

analysis. HNT TBS was the only condition for which recollection was significantly greater than all of the

other individual conditions (uncorrected P = 0.007 versus HNT TBS OFF, P = 0.03 versus HNT beta ON,

P = 0.008 versus HNT beta OFF, P = 0.0005 versus SMA TBS ON, P = 0.03 versus SMA TBS OFF, P =

0.03 versus SMA beta ON, and P = 0.02 versus SMA beta OFF). In contrast, no control condition was

associated with greater recollection than any one other condition (range of all uncorrected P-values =

0.08-0.99).

The experiment design used trial-specific stimulation with randomized condition order and long

inter-trial intervals to ensure that the effects of stimulation were specific, as any effects of stimulation that

outlasted the trial (i.e., due to longer-lasting neuroplasticity) would not have been condition specific. To

more directly assess whether effects of HNT TBS on recollection outlasted the trial during which it was

applied, we tested whether recollection of scenes with no stimulation (OFF) varied based on whether the

scene immediately followed a trial with HNT TBS (ON) versus a trial from the same study session with no

stimulation (OFF). There was no difference in the proportion of recollected hits for these two categories

of non-stimulated trials (t_{15}=0.66, P=0.52; two-tailed paired t-test). Thus, there was no evidence that HNT

TBS continued to affect recollection on the trial after which it was applied.

Although our hypotheses concerned the effects of stimulation on recollection, we also examined

effects on overall hit rates. Overall hit rate (old items endorsed with "Remember" and "Familiar"

responses) did not vary by stimulation pattern, location, or presence (all 3-way rmANOVA main effects

and interactions P-values>0.1) (Table 1). Notably however, although effects of stimulation on hit rates

were not identified, overall memory strength could have been affected by stimulation, which could

influence both the hit rate and the false alarm rate. The format of the experiment precluded calculation of

false alarm rates separately for each stimulation condition because study phases for both stimulation

locations (HNT and SMA) were tested at the end of each experimental session using the same set of

novel lures (see Fig. 1D). Therefore, lures could be segregated based on stimulation pattern (TBS versus

beta) but not based on stimulation location. The false alarm rate was significantly lower (t_{15}=2.54,

P=0.02, d=0.64) for TBS (mean=0.31, SD=0.17) than beta stimulation (mean=0.38, SD=0.12). Likewise,

the accuracy of "Remember" responses (proportion made to old items) was significantly greater for TBS

(mean=0.88, SD=0.09) than for beta (mean=0.81, SD=0.13; t_{15}=2.44, P=0.03, d=0.62; two-tailed paired t-

test), whereas the accuracy of "Familiar" responses did not differ significantly for TBS (mean=0.61,

SD=0.11) compared to beta (mean=0.57, SD=0.13; t_{15}=1.89, P=0.08; two-tailed paired t-test). Although

we could not test for differences based on stimulation location, these findings indicate that TBS

increased overall memory strength and recollection accuracy to a greater extent than beta stimulation,

which is consistent with the primary findings that recollection was improved selectively for HNT TBS.

<u>Effects of stimulation on recollection-related hippocampal activity during encoding</u>

As outlined above, we hypothesized that HNT TBS would be unique among stimulation conditions

in increasing hippocampal fMRI activity evoked by later-recollected scenes, and that this would occur

selectively for the left hippocampus that was targeted based on its fMRI connectivity to the HNT

stimulation location and based on the relatively lateralized projections from lateral parietal cortex to

ipsilateral medial temporal lobe (Mesulam et al., 1977; Mufson and Pandya, 1984). Further, we

hypothesized that these effects would be specific to scenes, with no impact of stimulation conditions on

activity evoked by numeric-judgment trials that were randomly intermixed with scene trials during the

encoding periods.

We first evaluated the effects of stimulation on activity evoked by later-recollected scenes for the

left hippocampus (Fig. 5A). Activity varied significantly by stimulation condition ($F_{4,75}$=3.40, P=0.01,

η^2=0.18, 1x5 rmANOVA for the HNT TBS, SMA TBS, HNT beta, SMA beta, and "off" conditions). The

right hippocampus showed a numerically similar trend of increased activity for HNT TBS, but activity did

not vary significantly by stimulation condition for right hippocampus ($F_{(GG)2.3,42.2}$=0.76, P=0.49).

The follow-up test of left hippocampal activity followed our *a priori* hypothesis that HNT TBS

would be the only stimulation condition to increase activity relative to other stimulation conditions. We

tested whether activity evoked by later-recollection scenes was greater for HNT TBS versus the average

of all control conditions (SMA TBS, HNT beta, SMA beta, and off), and also tested whether activity was

significantly greater for each of the control conditions versus the average of all other control conditions.

Consistent with our hypothesis, HNT TBS was the only stimulation condition with significantly greater

activity relative to control conditions (z_{15}=2.61, P=0.005, P_B=0.03 corrected for 5 tests; r=0.65; P_B for

each control condition relative to the average of its respective other controls range = 0.67-1; Fig. 5B).

An exploratory analysis of all possible pairwise comparisons among stimulation conditions was

also performed. HNT TBS was the only condition for which left hippocampal activity evoked by later-

recollected scenes was significantly greater than for all of the other individual conditions (uncorrected P =

0.03 versus HNT beta, P = 0.002 versus SMA TBS, P = 0.02 versus SMA beta, and P = 0.03 versus

combined OFF). In contrast, no control condition was associated with greater activity than any other one

condition (range of uncorrected P-values = 0.07-0.93).

As shown in Fig. 5C, numeric judgments did not evoke robust activity in left or right hippocampus,

relative to activity evoked by scenes (Fig. 5B), as anticipated. This activity did not vary significantly by

stimulation condition for either left ($F_{4,75}$=1.04, P>0.4) or right ($F_{4,75}$=0.81, P>0.5) hippocampus. Focusing

specifically on the HNT TBS condition of interest and left hippocampus, the effect of stimulation relative

to the average of all control conditions was significantly greater for scene-evoked recollection-related

activity (difference between HNT TBS bar and adjacent white bar in Fig. 5B) than it was for activity

evoked by numeric judgments (difference between HNT TBS bar and adjacent white bar in Fig. 5C)

(z_{15}=2.30, P=0.01, r=0.58), indicating the impact of HNT TBS on activity evoked by later-recollected

scenes was greater than the impact of HNT TBS on activity evoked by numeric judgments. Collectively,

these findings indicate that HNT TBS selectively increased left hippocampal activity evoked by later-

recollected scenes during encoding.

<u>Exploratory analysis of variation in stimulation effects along the hippocampal longitudinal axis</u>

707   Based on evidence for anatomical and functional specialization along the hippocampal

longitudinal axis (Ranganath et al., 2004; Poppenk et al., 2013) as well as our own previous evidence

that network-targeted TMS can affect fMRI activity of specific hippocampal long-axis segments (Wang et

al., 2014; Nilakantan et al., 2019), we performed an exploratory analysis to test for variation in the effects

of HNT TBS along the longitudinal axis using five segments extending from anterior head to posterior

body (Fig. 6A) (described in Methods). For left hippocampal segments, activity varied significantly by the

five stimulation conditions for the two most anterior segments ($F_{(GG)1.9,36.0}$=4.14, P=0.03, η^2_p=0.28;

$F_{(GG)1.7,32.4}$=3.80, P=0.04, η^2_p=0.25; all other Ps>0.1) (Fig. 6B). Follow-up pairwise comparisons were

made among conditions using averaged activity of these two anterior segments, not corrected for

multiple comparisons in this exploratory analysis. Activity evoked by later-recollected scenes was greater

for HNT TBS relative to all control conditions (range of uncorrected P-values = 0.00009-0.03). Activity

was also significantly greater for OFF relative to SMA TBS (t_{15}=2.47, P=0.01, d=0.62), but no other

control condition significantly differed from another (range of uncorrected P-values = 0.07-0.99).

720   As expected based on findings from the primary analysis (Fig. 5), the exploratory analysis

performed for right (non-targeted) MTL segments yielded a numerically similar pattern of greater activity

following HNT TBS relative to other conditions, but there was no significant variation by stimulation

condition for any longitudinal segment (Ps>0.3) (Fig. 6B). Furthermore, as expected, MTL activity evoked

by numeric judgments was minimal along the longitudinal axis and did not vary significantly by

stimulation condition or segment in either the left or right hemisphere (Fig. 6C; main effect of condition

and interaction of condition by longitudinal segment in the left and right hemispheres Ps>0.3).

Collectively, this exploratory analysis provides preliminary evidence that effects of HNT TBS on activity

evoked by later-recollected scenes were most pronounced for relatively anterior hippocampus in the

targeted hemisphere.

 <u>Voxel-wise fMRI analysis to evaluate non-hippocampal stimulation effects</u>

To complement the ROI-based approach that was used to assess the hypothesized effects of

stimulation on hippocampal activity evoked by later-recollected scenes, we performed a voxel-wise

analysis with an intentionally liberal statistical threshold to determine if there was any evidence for effects

of stimulation elsewhere within the FOV (Material and Methods). The rationale for using an exceptionally

liberal statistical threshold is that this provides high sensitivity to the any potential effects of stimulation,

with the caveat that any results would require further testing via a more rigorous approach. The voxel-

wise rmANOVA assessing the interaction of stimulation location (HNT, SMA) with rhythm (TBS, beta) on

activity evoked by later-recalled scenes did not identify supra-threshold activity anywhere outside of the

MTL. The only supra-threshold activity identified by this test was for the left anterior MTL (82-voxel

cluster, peak MNI = -20, 4, -23), and this location overlaps considerably with the anterior-left

hippocampal regions of interest where activity evoked by later-recollected scenes was most robustly

influenced by HNT TBS (Figs. 6A). This indicates that this liberal voxel-wise test had the sensitivity to

identify hypothesized effects of stimulation on activity (as it identified such effects in a similar area as in

the ROI-based analysis), yet there were no robust differential effects of stimulation location and rhythm

outside of the hippocampal regions of interest. The same test performed for activity evoked by numeric

digits did not yield any supra-threshold clusters for either the main effects or interaction.

<u>No effects of stimulation on hippocampal activity during relatively unsuccessful scene encoding</u>

In order to determine whether stimulation effects on fMRI activity evoked by scenes were specific

to those scenes that were later recollected, we performed the same set of fMRI analyses previously

described for the later-recollected scenes but for the scenes that were later endorsed with "Familiar" and

"New" (miss) responses. As indicated above, "Familiar" responses were significantly less accurate than

"Remember" responses, and "New" responses indicated memory failure. Thus, memory encoding was

far less successful for this combined Familiar/New condition (44% "Familiar" responses on average,

SD=6.49%) than it was for the "Remember" condition.

Stimulation condition did affect activity for this condition in the left hippocampus ($F_{4,75}=0.34$,

P>0.8) or right hippocampus ($F_{4,75}=1.45$, P>0.2). Considering the hippocampal longitudinal axis

exploratory analysis, there were also no main effects of stimulation condition or interaction of stimulation

condition by the five segments for either the left or right hippocampus (main effects and interaction of

condition by longitudinal segment in the left and right hemispheres Ps>0.2). The liberal whole-brain

voxel-wise analysis did not find any supra-threshold clusters with an interaction of stimulation location by

stimulation rhythm. Thus, the effects of HNT TBS on activity evoked by scenes were specific to the

scenes that were later-recollected, with no effects of stimulation on activity for scenes comprising this

low-memory condition.

Discussion

TBS targeting the network of the left hippocampus (HNT) during encoding selectively increased fMRI

activity evoked by later-recollected scenes in the targeted (left) hippocampus and increased subsequent

recollection memory. These increases were rhythm- and location-specific, as they were not observed for

HNT beta stimulation nor for either stimulation rhythm (TBS or beta) targeting an out-of-network control

location (SMA). Further, effects were cognitively specific, as no condition (including HNT TBS) influenced

hippocampal activity for scenes that were not later remembered (later-familiar and later-miss trials) nor

did any condition influence hippocampal activity during numeric judgments, which do not typically evoke

MTL activity (Stark and Squire, 2001). Effects were specific to the left hippocampus, which was targeted

via its connectivity to the stimulated left parietal area. A numerically similar but non-significant pattern

was observed for right hippocampus, which could be due to commissural connectivity and is consistent

with previous findings of relatively (but not completely) lateralized effects of network-targeted TMS on

hippocampal activity (Wang et al., 2014; Nilakantan et al., 2019). Finally, HNT TBS only influenced the

recollective aspect of scene memory, which supports the conclusion that hippocampal function was

affected, as recollection is associated with hippocampal activity during encoding (Kim, 2011; Rugg et al.,

2012).

Notably, the experiment design constrains mechanistic interpretation. Stimulation was delivered

as 2-sec volleys immediately before stimulus onset and with conditions intermixed across encoding trials,

with intentionally long inter-stimulus intervals to avoid cumulative effects of stimulation across trials

(Huang et al., 2005; Demeter et al., 2016). Any effects of HNT TBS versus control conditions must

therefore have been immediate and trial specific, and indeed a control analysis provided no evidence

that HNT TBS continued to affect encoding even on the subsequent trial. The experiment design and

pattern of findings thus suggests that HNT TBS influenced hippocampal neural activity, as opposed to

neuroplasticity and/or neuromodulatory mechanisms that can support persistent/long-lasting effects of

stimulation (Cirillo et al., 2017). Notably, the design and findings also indicate that HNT TBS effects on

fMRI activity were unlikely to have reflected TMS-related fMRI artefact, as different trial types were

intermixed and all received the same stimulation parameters, yet effects were specific to later-recollected

scenes.

The premise of the experiment was that TBS mimics endogenous theta-band neural activity

characteristic of the hippocampus and hippocampal network synchrony (Buzsaki, 2002; Buzsaki and

Draguhn, 2004; Lisman and Jensen, 2013) and therefore might optimally influence hippocampal memory

processing via entrainment of neural activity (Thut et al., 2011a; Hanslmayr et al., 2019). Thus, if

targeting hippocampal network theta activity via noninvasive TBS were successful, we expected that it

would cause population synchrony of theta and therefore increase the evoked fMRI BOLD response to a

stimulus that evokes processing by the affected region(s). Indeed, increased hippocampal theta activity

predicts successful memory formation, particularly for complex associative information (Rutishauser et

al., 2010; Fell et al., 2011; Herweg et al., 2020), including theta activity in the pre-stimulus interval

(Guderian et al., 2009; Fell et al., 2011; Merkow et al., 2014) when TBS was applied in our experiment.

Although our findings are consistent with this prediction, a weakness is that theta cannot be measured

with fMRI and we can only infer an impact based on fMRI activity and memory performance. Further, it is

also possible that recollection improvement due to HNT TBS was mediated by a mechanism other than

the corresponding increase in hippocampal activity, as fMRI provides a limited window on brain activity.

Full confirmation of our interpretation requires additional evidence, including direct measurement of

stimulation effects on hippocampal theta.

The findings inform mechanisms by which brain stimulation influences hippocampal activity.

Direct electrical stimulation of the hippocampus and its immediate entorhinal inputs via depth electrodes

in human neurosurgical cases typically disrupts memory [(Coleshill et al., 2004; Jacobs et al., 2016;

Goyal et al., 2018b), but see (Suthana et al., 2012; Jun et al., 2019)]. In contrast, direct/neurosurgical

electrical stimulation of hippocampal network areas has yielded modest evidence for memory

enhancement. Stimulation of lateral temporal cortex enhanced memory in four patients (Kucewicz et al.,

2018), and closed-loop stimulation of approximately the same location caused relative enhancement

compared to disruptive effects for other brain regions (Ezzyat et al., 2018). Of relevance to theta,

memory enhancement was achieved in four patients receiving TBS of the fornix (Miller et al., 2015), and

TBS with microstimulation of entorhinal cortex enhanced memory in several cases in which white matter

(rather than gray matter) was targeted (Titiz et al., 2017). However, demonstrations of memory

enhancement by invasive TBS (Miller et al., 2015; Titiz et al., 2017) did not include non-theta control

conditions, and therefore do not permit strong conclusions regarding the specific role of theta.

TMS targeting the hippocampal network can generate long-lasting and robust recollection

improvement and corresponding changes in hippocampal and hippocampal network activity (Hebscher

and Voss, 2020), with greater aftereffects using TBS (Hermiller et al., 2019a). The current findings

provide novel mechanistic insights by showing that hippocampal activity is immediately sensitive to

stimulation applied to its network and mimicking its endogenous theta activity. This suggests an impact of

network-targeted stimulation on hippocampal neural activity and supports the interpretation (Hebscher

and Voss, 2020) that effects of noninvasive stimulation targeting this network are due to the impact of

stimulation on the hippocampus.

Although sampling of activity elsewhere in the brain was limited by the fMRI methods, we found

no evidence for effects of HNT TBS among locations sampled outside of the left MTL. This suggests that

the hippocampus may be unique in its ability to be immediately impacted by stimulation. In contrast,

previous experiments using longer stimulation trains and/or multiple days of stimulation found effects

distributed throughout the network (Hebscher and Voss, 2020). Interestingly, in those studies, although

stimulation was delivered at a parietal network target, the most robust effects were on fMRI activity of the

hippocampus and network locations with highest hippocampal connectivity (Hebscher and Voss, 2020). It

is possible that brief TBS trains affect activity in only those areas most sensitive to this stimulation

pattern (hippocampus) whereas more extensive network locations are recruited with increasing

stimulation.

Functionally distinct theta bands have recently been reported along the human longitudinal

hippocampal axis, with slower theta (~3-5 Hz) more prevalent in the anterior and higher theta (~6-8 Hz)

more prevalent in the posterior hippocampus (Lega et al., 2016; Goyal et al., 2018a). TBS was delivered

at 5 Hz and the exploratory analysis suggested that it had relatively greater impact on the anterior

hippocampus. This further suggests that TBS might have affected neural activity in the matching

frequency band, if 5-Hz theta is more pronounced in anterior hippocampus. Formal testing of this

possibility is a direction for future experiments.

The reported effects of HNT TBS are highly unlikely to have resulted from the impact of

stimulation on the left parietal area where HNT stimulation was applied. This parietal location has been

associated with confidence and meta-memory judgments during retrieval but not with successful memory

encoding (Uncapher and Wagner, 2009; Kim, 2011). Stimulation in this experiment was applied at

encoding, not during retrieval. Moreover, because effects were condition- and trial-specific, with no

impact of HNT TBS even on the very next trial after it was applied, it is incredibly unlikely that stimulation

had any lingering impact during the delayed test period. Because hippocampus is robustly implicated in

encoding that produces later recollection whereas the stimulated area of parietal cortex is not, the

experiment affords greater confidence that the effects of HNT TBS on recollection memory were related

to the corresponding increases in hippocampal activity during encoding produced by HNT TBS (Tambini

et al., 2018).

A limitation of the current experiment is that the memory test was given after both study phases

on a given experiment session, and those study phases differed in the stimulation location (HNT versus

SMA), keeping stimulation rhythm constant (TBS or beta). Thus, we could not compare effects of location

on memory accuracy, as novel foils were not segregated by location. Nonetheless, we did find reduced

false alarms to novel foils for TBS versus beta as well as an increase in hit rates for HNT TBS versus

TBS to the SMA location. Furthermore, the accuracy of recollection, but not familiarity, responses

increased for TBS versus beta. These effects on accuracy are consistent with the main findings that HNT

TBS specifically impacted recollection, but future experiments will be necessary to directly test whether

TBS increases accuracy only when applied to target the hippocampus versus to any location.

Hippocampal network dysfunction is related to memory impairments in many disorders (Andrews-

Hanna et al., 2007; Buckner et al., 2008; Dickerson and Eichenbaum, 2010; Small et al., 2011). The

current findings suggest that memory processing by the core structure of this network can be

immediately and beneficially influenced via TMS of an appropriate location and rhythm. Given the

immediate impact and relatively precise locus of stimulation-related activity changes, concurrent TBS

with fMRI could be a powerful tool for testing hypotheses about normal hippocampal function as well as

dysfunction in memory disorders.

References:

876
Aggleton JP, Brown MW (1999) Episodic memory, amnesia, and the hippocampal-anterior thalamic axis.
Behav Brain Sci 22:425-444; discussion 444-489.
Andrews-Hanna JR, Snyder AZ, Vincent JL, Lustig C, Head D, Raichle ME, Buckner RL (2007)
Disruption of large-scale brain systems in advanced aging. Neuron 56:924-935.
Bestmann S, Ruff CC, Blankenburg F, Weiskopf N, Driver J, Rothwell JC (2008) Mapping causal
interregional influences with concurrent TMS-fMRI. Exp Brain Res 191:383-402.
Buckner RL, Andrews-Hanna JR, Schacter DL (2008) The brain's default network: anatomy, function,
and relevance to disease. Ann N Y Acad Sci 1124:1-38.
Buzsaki G (2002) Theta oscillations in the hippocampus. Neuron 33:325-340.
Buzsaki G, Draguhn A (2004) Neuronal oscillations in cortical networks. Science 304:1926-1929.
Cirillo G, Di Pino G, Capone F, Ranieri F, Florio L, Todisco V, Tedeschi G, Funke K, Di Lazzaro V (2017)
Neurobiological after-effects of non-invasive brain stimulation. Brain Stimul 10:1-18.
Coleshill SG, Binnie CD, Morris RG, Alarcon G, van Emde Boas W, Velis DN, Simmons A, Polkey CE,
van Veelen CW, van Rijen PC (2004) Material-specific recognition memory deficits elicited by
unilateral hippocampal electrical stimulation. J Neurosci 24:1612-1616.
Cox RW (1996) AFNI: software for analysis and visualization of functional magnetic resonance
neuroimages. Comput Biomed Res 29:162-173.
Demeter E, Mirdamadi JL, Meehan SK, Taylor SF (2016) Short theta burst stimulation to left frontal
cortex prior to encoding enhances subsequent recognition memory. Cogn Affect Behav Neurosci
16:724-735.
Dickerson BC, Eichenbaum H (2010) The episodic memory system: neurocircuitry and disorders.
Neuropsychopharmacology 35:86-104.
Eichenbaum H, Yonelinas AP, Ranganath C (2007) The medial temporal lobe and recognition memory.
Annu Rev Neurosci 30:123-152.
Ezzyat Y et al. (2018) Closed-loop stimulation of temporal cortex rescues functional networks and
improves memory. Nat Commun 9:365.
Fell J, Klaver P, Lehnertz K, Grunwald T, Schaller C, Elger CE, Fernandez G (2001) Human memory
formation is accompanied by rhinal-hippocampal coupling and decoupling. Nat Neurosci 4:1259-
1264.
Fell J, Ludowig E, Staresina BP, Wagner T, Kranz T, Elger CE, Axmacher N (2011) Medial temporal
theta/alpha power enhancement precedes successful memory encoding: evidence based on
intracranial EEG. J Neurosci 31:5392-5397.
Foster BL, Kaveh A, Dastjerdi M, Miller KJ, Parvizi J (2013) Human retrosplenial cortex displays transient
theta phase locking with medial temporal cortex prior to activation during autobiographical
memory retrieval. J Neurosci 33:10439-10446.
Fox MD, Halko MA, Eldaief MC, Pascual-Leone A (2012) Measuring and manipulating brain connectivity
with resting state functional connectivity magnetic resonance imaging (fcMRI) and transcranial
magnetic stimulation (TMS). Neuroimage 62:2232-2243.
Freedberg M, Reeves JA, Toader AC, Hermiller MS, Voss JL, Wassermann EM (2019) Persistent
Enhancement of Hippocampal Network Connectivity by Parietal rTMS Is Reproducible. eNeuro 6.
Friedman L, Glover GH (2006) Report on a multicenter fMRI quality assurance protocol. J Magn Reson
Imaging 23:827-839.
Goyal A, Miller J, Qasim A, Watrous AJ, Stein JM, Inman CS, Gross RE, Willie JT, Lega B, Lin J, Sharan
A, Wu C, Sperling MR, Sheth SA, McKhann GM, Smith EH, Schevon C, Jacobs J (2018a)
Functionally distinct high and low theta oscillations in the human hippocampus. bioRxiv.
Goyal A, Miller J, Watrous AJ, Lee SA, Coffey T, Sperling MR, Sharan A, Worrell G, Berry B, Lega B,
Jobst BC, Davis KA, Inman C, Sheth SA, Wanda PA, Ezzyat Y, Das SR, Stein J, Gorniak R,
Jacobs J (2018b) Electrical Stimulation in Hippocampus and Entorhinal Cortex Impairs Spatial
and Temporal Memory. J Neurosci 38:4471-4481.
Guderian S, Schott BH, Richardson-Klavehn A, Duzel E (2009) Medial temporal theta state before an
event predicts episodic encoding success in humans. Proc Natl Acad Sci U S A 106:5365-5370.
Hanslmayr S, Axmacher N, Inman CS (2019) Modulating Human Memory via Entrainment of Brain
Oscillations. Trends Neurosci 42:485-499.

Hebscher M, Voss JL (2020) Testing network properties of episodic memory using non-invasive brain
stimulation. Curr Opin Behav Sci 32:35-42.
Hermiller MS, VanHaerents S, Raij T, Voss JL (2019a) Frequency-specific noninvasive modulation of
memory retrieval and its relationship with hippocampal network connectivity. Hippocampus
29:595-609.
Hermiller MS, Karp E, Nilakantan AS, Voss JL (2019b) Episodic memory improvements due to
noninvasive stimulation targeting the cortical-hippocampal network: A replication and extension
experiment. Brain Behav 9:e01393.
Herweg NA, Solomon EA, Kahana MJ (2020) Theta Oscillations in Human Memory. Trends Cogn Sci
24:208-227.
Huang YZ, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC (2005) Theta burst stimulation of the human
motor cortex. Neuron 45:201-206.
Jacobs J, Miller J, Lee SA, Coffey T, Watrous AJ, Sperling MR, Sharan A, Worrell G, Berry B, Lega B,
Jobst BC, Davis K, Gross RE, Sheth SA, Ezzyat Y, Das SR, Stein J, Gorniak R, Kahana MJ,
Rizzuto DS (2016) Direct Electrical Stimulation of the Human Entorhinal Region and
Hippocampus Impairs Memory. Neuron 92:983-990.
Jun S, Kim JS, Chung CK (2019) Direct Stimulation of Human Hippocampus During Verbal Associative
Encoding Enhances Subsequent Memory Recollection. Front Hum Neurosci 13:23.
Kahn I, Andrews-Hanna JR, Vincent JL, Snyder AZ, Buckner RL (2008) Distinct cortical anatomy linked
to subregions of the medial temporal lobe revealed by intrinsic functional connectivity. J
Neurophysiol 100:129-139.
Kim H (2011) Neural activity that predicts subsequent memory and forgetting: a meta-analysis of 74 fMRI
studies. Neuroimage 54:2446-2461.
Kim S, Nilakantan AS, Hermiller MS, Palumbo RT, VanHaerents S, Voss JL (2018) Selective and
coherent activity increases due to stimulation indicate functional distinctions between episodic
memory networks. Sci Adv 4:eaar2768.
Kohler S, Paus T, Buckner RL, Milner B (2004) Effects of left inferior prefrontal stimulation on episodic
memory formation: a two-stage fMRI-rTMS study. J Cogn Neurosci 16:178-188.
Kucewicz MT, Berry BM, Miller LR, Khadjevand F, Ezzyat Y, Stein JM, Kremen V, Brinkmann BH,
Wanda P, Sperling MR, Gorniak R, Davis KA, Jobst BC, Gross RE, Lega B, Van Gompel J, Stead
SM, Rizzuto DS, Kahana MJ, Worrell GA (2018) Evidence for verbal memory enhancement with
electrical brain stimulation in the lateral temporal cortex. Brain 141:971-978.
Lega B, Burke J, Jacobs J, Kahana MJ (2016) Slow-Theta-to-Gamma Phase-Amplitude Coupling in
Human Hippocampus Supports the Formation of New Episodic Memories. Cereb Cortex 26:268-
278.
Lisman JE, Jensen O (2013) The theta-gamma neural code. Neuron 77:1002-1016.
Merkow MB, Burke JF, Stein JM, Kahana MJ (2014) Prestimulus theta in the human hippocampus
predicts subsequent recognition but not recall. Hippocampus 24:1562-1569.
Mesulam MM, Van Hoesen GW, Pandya DN, Geschwind N (1977) Limbic and sensory connections of
the inferior parietal lobule (area PG) in the rhesus monkey: a study with a new method for
horseradish peroxidase histochemistry. Brain Res 136:393-414.
Miller JP, Sweet JA, Bailey CM, Munyon CN, Luders HO, Fastenau PS (2015) Visual-spatial memory
may be enhanced with theta burst deep brain stimulation of the fornix: a preliminary investigation
with four cases. Brain 138:1833-1842.
Moisa M, Pohmann R, Ewald L, Thielscher A (2009) New coil positioning method for interleaved
transcranial magnetic stimulation (TMS)/functional MRI (fMRI) and its validation in a motor cortex
study. J Magn Reson Imaging 29:189-197.
Mufson EJ, Pandya DN (1984) Some observations on the course and composition of the cingulum
bundle in the rhesus monkey. J Comp Neurol 225:31-43.
Nilakantan AS, Mesulam MM, Weintraub S, Karp EL, VanHaerents S, Voss JL (2019) Network-targeted
stimulation engages neurobehavioral hallmarks of age-related memory decline. Neurology
92:e2349-e2354.
Paller KA, Wagner AD (2002) Observing the transformation of experience into memory. Trends Cogn Sci
6:93-102.

Poppenk J, Evensmoen HR, Moscovitch M, Nadel L (2013) Long-axis specialization of the human hippocampus. Trends Cogn Sci 17:230-240.

Ranganath C, Ritchey M (2012) Two cortical systems for memory-guided behaviour. Nat Rev Neurosci 13:713-726.

Ranganath C, Yonelinas AP, Cohen MX, Dy CJ, Tom SM, D'Esposito M (2004) Dissociable correlates of recollection and familiarity within the medial temporal lobes. Neuropsychologia 42:2-13.

Romei V, Thut G, Silvanto J (2016) Information-Based Approaches of Noninvasive Transcranial Brain Stimulation. Trends Neurosci 39:782-795.

Rossi S, Hallett M, Rossini PM, Pascual-Leone A, Group SoTC (2009) Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. Clin Neurophysiol 120.

Rugg MD, Vilberg KL, Mattson JT, Yu SS, Johnson JD, Suzuki M (2012) Item memory, context memory and the hippocampus: fMRI evidence. Neuropsychologia 50:3070-3079.

Rutishauser U, Ross IB, Mamelak AN, Schuman EM (2010) Human memory strength is predicted by theta-frequency phase-locking of single neurons. Nature 464:903-907.

Shirvalkar PR, Rapp PR, Shapiro ML (2010) Bidirectional changes to hippocampal theta-gamma comodulation predict memory for recent spatial episodes. Proc Natl Acad Sci U S A 107:7054-7059.

Siebner HR et al. (2009) Consensus paper: combining transcranial stimulation with neuroimaging. Brain Stimul 2:58-80.

Small SA, Schobel SA, Buxton RB, Witter MP, Barnes CA (2011) A pathophysiological framework of hippocampal dysfunction in ageing and disease. Nat Rev Neurosci 12:585-601.

Squire LR, Zola-Morgan S (1991) The medial temporal lobe memory system. Science 253:1380-1386.

Stark CE, Squire LR (2001) When zero is not zero: the problem of ambiguous baseline conditions in fMRI. Proc Natl Acad Sci U S A 98:12760-12766.

Staudigl T, Hanslmayr S (2013) Theta oscillations at encoding mediate the context-dependent nature of human episodic memory. Curr Biol 23:1101-1106.

Stern CE, Corkin S, Gonzalez RG, Guimaraes AR, Baker JR, Jennings PJ, Carr CA, Sugiura RM, Vedantham V, Rosen BR (1996) The hippocampal formation participates in novel picture encoding: evidence from functional magnetic resonance imaging. Proc Natl Acad Sci U S A 93:8660-8665.

Suthana N, Haneef Z, Stern J, Mukamel R, Behnke E, Knowlton B, Fried I (2012) Memory enhancement and deep-brain stimulation of the entorhinal area. N Engl J Med 366:502-510.

Tambini A, Nee DE, D'Esposito M (2018) Hippocampal-targeted Theta-burst Stimulation Enhances Associative Memory Formation. J Cogn Neurosci 30:1452-1472.

Thickbroom GW (2007) Transcranial magnetic stimulation and synaptic plasticity: experimental framework and human models. Exp Brain Res 180:583-593.

Thut G, Schyns PG, Gross J (2011a) Entrainment of perceptually relevant brain oscillations by non-invasive rhythmic stimulation of the human brain. Front Psychol 2:170.

Thut G, Veniero D, Romei V, Miniussi C, Schyns P, Gross J (2011b) Rhythmic TMS causes local entrainment of natural oscillatory signatures. Curr Biol 21:1176-1185.

Titiz AS, Hill MRH, Mankin EA, Aghajan ZM, Eliashiv D, Tchemodanov N, Maoz U, Stern J, Tran ME, Schuette P, Behnke E, Suthana NA, Fried I (2017) Theta-burst microstimulation in the human entorhinal area improves memory specificity. Elife 6.

Uncapher MR, Wagner AD (2009) Posterior parietal cortex and episodic encoding: insights from fMRI subsequent memory effects and dual-attention theory. Neurobiol Learn Mem 91:139-154.

Wang JX, Rogers LM, Gross EZ, Ryals AJ, Dokucu ME, Brandstatt KL, Hermiller MS, Voss JL (2014) Targeted enhancement of cortical-hippocampal brain networks and associative memory. Science 345:1054-1057.

Warren KN, Hermiller MS, Nilakantan AS, Voss JL (2019) Stimulating the hippocampal posterior-medial network enhances task-dependent connectivity and memory. Elife 8.

Xia M, Wang J, He Y (2013) BrainNet Viewer: a network visualization tool for human brain connectomics. PLoS One 8:e68910.

Xiao JX, Ehinger KA, Hays J, Torralba A, Oliva A (2016) SUN Database: Exploring a Large Collection of Scene Categories. International Journal of Computer Vision 119:3-22.

039   Yonelinas AP (2002) The nature of recollection and familiarity: A review of 30 years of research. Journal
040   of Memory and Language 46:441-517.
041   Zutshi I, Brandon MP, Fu ML, Donegan ML, Leutgeb JK, Leutgeb S (2018) Hippocampal Neural Circuits
042   Respond to Optogenetic Pacing of Theta Frequencies by Generating Accelerated Oscillation
043   Frequencies. Curr Biol 28:1179-1188 e1173.
044

    Figure Legends

Figure 1. Trial-specific stimulation during episodic memory formation. (**A**) Scene-encoding and
numeric-judgment trials were randomly intermixed during each study phase, with ~2 s of stimulation
delivered immediately before stimulus onset for a subset of trials (ON) and no preceding stimulation for
remaining trials (OFF). Study phases were completed during fMRI scanning with memory test phases
after scanning. There were four stimulation conditions for ON and OFF trials. (**B**) Stimulation was
delivered as either a theta-burst pattern (TBS: 50 Hz triplet pulses delivered at 5 Hz) or at beta (single
pulses delivered at 12.5 Hz). These conditions had the same overall number of pulses during each
stimulation period delivered at the same intensity. (**C**) Stimulation was delivered to the HNT parietal
location (based on its fMRI connectivity with left hippocampus, as depicted by the blue arrow), or a
control out-of-network SMA location. Achieved stimulation locations confirmed via MRI for each condition
and subject are indicated by colorized spheres on a template brain. Bar plots represent mean ±s.e.m.
baseline resting-state fMRI connectivity of the subject-specific stimulation locations with the hippocampal
network, confirming relatively higher connectivity for the HNT than SMA location. *$P<0.05$ main effect of
stimulation location by one-way rmANOVA. (**D**) HNT or SMA locations were targeted for one of the two
study phases in each experimental session. After both study phases were complete, subjects exited the
scanner for a ~15 min break before taking the memory test. A different stimulation pattern (TBS or beta)
was used for each experimental session. Black arrows indicate stimulation conditions with order
counterbalanced across subjects.

Figure 2. Interleaved TMS/fMRI scan sequences. Depiction of one imaging volume in the (**A**) TBS and (**B**) beta scan sequences. Each grey cube represents one MR EPI slice acquisition and each colored line indicates a TMS pulse (purple for TBS; green for beta). The extent of imaging coverage (22 EPI slices for TBS; 30 EPI slices for beta) is shown on a template brain, with EPI slices colorized to match the TMS pulses (purple for the TBS scan; green for the beta scan).

Figure 3. Expected fMRI signals of scene processing and stimulation sensations confirm fMRI data quality during concurrent TMS. Voxel-wise contrasts confirm that the fMRI signal could distinguish the task stimuli (scenes vs. numbers) and the presence of stimulation (ON vs. OFF). (**A**) Group-level contrast of scenes versus numbers, regardless of stimulation location or presence, identified significantly greater activation by scenes in areas that are typically scene-sensitive for both TBS and beta stimulation, including bilateral hippocampus and occipital, fusiform, and posterior parahippocampal cortex. (**B**) Group-level contrast of TMS ON versus OFF, regardless of stimulation location or the stimuli type (scenes and numbers) identified significantly greater activation for TMS ON in the bilateral auditory cortex for both TBS and beta stimulation. Direct contrasts of TBS versus beta did not identify significant differences for either comparison (see main text). Plots show supra-threshold voxels on a template brain. Color bars indicate t-statistic calculated by *3dMEMA*.

Figure 4. Theta-patterned stimulation of the hippocampal network during encoding selectively increased subsequent recollection. (**A**) Subjects discriminated studied scenes (old; depicted here with a yellow border) from randomly intermixed novel lures (new) using "Remember" responses to indicate the experience of hippocampal-dependent recollection, "Familiar" responses to indicate the experience of familiarity, and "New" responses to indicate lures. Example correct responses are underlined. (**B**) To identify stimulation effects on memory while holding sensory entrainment qualities (e.g., somatosensory, auditory) relatively constant, we compared the difference in the proportion of recollected hits for stimulation targeting HNT versus SMA for each of the stimulation patterns (TBS, beta, and off). * P=0.02 interaction between TMS presence and rhythm reported in text. (**C**) The proportion of recollected hits for HNT TBS was greater than for the aggregate of all control conditions. ** P_B=0.008 reported in text. None of the control conditions increased the proportion of recollected hits relative to the aggregate of all other control conditions (nonsignificant values reported in text). White bars next to each condition indicate the average of the control conditions that the corresponding condition of interest (colorized bars) was tested against. Bars and error bars indicate mean ± s.e.m.

Figure 5. HNT TBS selectively increased left hippocampal activity during recollection memory formation. (**A**) Stimulation-evoked activity was measured within hippocampal/MTL units centered along the hippocampal longitudinal axis in each hemisphere, shown as colorized voxels in coronal slices of a template brain. The left hippocampus/MTL was targeted indirectly via its network functional connectivity (Materials and Methods) and the right hippocampus/MTL served as a non-targeted control. (**B**) fMRI activity evoked by later-recollected scenes for each stimulation condition (colorized bars) and the aggregate of all other control conditions (white bards adjacent to each colorized bars), plotted separately for left and right hippocampus. Bars and error bars indicate mean ± s.e.m. estimated fMRI activity (arbitrary units). Activity evoked by later-recollected scenes for HNT TBS was greater than for the aggregate of all control conditions. * P_B=0.03 reported in the text. None of the control conditions increased active relative to the aggregate of all other control conditions (nonsignificant values reported in text). (**C**) Activity evoked by numbers displayed as in panel B. Significant difference of the effect of HNT TBS relative to the average of all controls for scene-evoked recollection-related activity versus activity evoked by numeric judgments is indicated (*P=0.01 as reported in the text).

Figure 6. Exploratory analysis of differential stimulation effects on activity along the hippocampal longitudinal axis. The same fMRI analyses as described in Figure 5 are presented but segregated for longitudinal segments of the hippocampus in each hemisphere. (**A**) The five segments along the longitudinal axis in each hemisphere are shown as colorized spheres on a template brain. (**B**) fMRI activity evoked by scenes later-recollected was extracted from each of the spherical ROIs in the left and right hemisphere for each stimulation condition. Line plots indicate mean estimated fMRI activity (arbitrary units) ±s.e.m. (indicated by the shaded area), colorized for each of the five stimulation conditions. (**C**) fMRI activity evoked by numbers displayed as in panel B. * P<0.05 main effect of stimulation condition by five-way rmANOVA reported in text are outlined. Follow-up pairwise comparisons among stimulation conditions are described in the text.

Table 1. Memory performance. For the test phase in both sessions, the total hit rate (i.e., both

Remember and Familiar responses) and the proportion of hits that were recollected (i.e., Remember

responses; R if hit) are provided for the scenes that were presented during the study phases for the HNT

and SMA location conditions, for trials with stimulation (TMS ON) and without (TMS OFF). The total false

alarm rate (i.e., both Remember and Familiar responses) and the proportion of false alarms that were

endorsed with recollection (R if false alarm) are provided for scenes that were not presented during the

study phase (novel foils). Statistical comparisons of interest are described in the text. Values are given

as across-participant means (SD).

	TBS session		Beta stimulation session	
	HNT	SMA	HNT	SMA
Total hit rate, TMS ON	0.66 (0.10)	0.61 (0.16)	0.67 (0.12)	0.66 (0.16)
Total hit rate, TMS OFF	0.63 (0.13)	0.62 (0.19)	0.62 (0.15)	0.64 (0.13)
Recollection hits, TMS ON	0.56 (0.13)	0.42 (0.15)	0.48 (0.19)	0.46 (0.21)
Recollection hits, TMS OFF	0.48 (0.18)	0.47 (0.20)	0.44 (0.21)	0.43 (0.22)
Total false alarm rate	0.31 (0.17)		0.38 (0.12)	
Recollection false alarms	0.17 (0.10)		0.21 (0.23)	