Social isolation modulates appetite and defensive behavior via a common

oxytocinergic circuit in larval zebrafish

Caroline L. Wee[1,2,3§], **Erin Song**[1§], **Maxim Nikitchenko**[1,4§], Kristian J. Herrera[1], Sandy Wong[1],

Florian Engert[1]*, Sam Kunes[1]*

§Equal-contribution first authors

*Corresponding authors

Affiliations:

[1]Department of Molecular and Cellular Biology and Center for Brain Science, Harvard

University, Cambridge, Massachusetts, USA

[2]Program in Neuroscience, Department of Neurobiology, Harvard Medical School, Boston,

Massachusetts, USA

[3]Present address: Institute of Molecular and Cell Biology, A*STAR, Singapore

[4]Present address: Duke University, Durham, North Carolina, USA

Correspondence:

Florian Engert / Sam Kunes

Department of Molecular and Cellular Biology

Harvard University, 16 Divinity Ave, Biolabs 2073

Cambridge, MA 02138, USA

ABSTRACT

Animal brains have evolved to encode social stimuli and transform these representations into

advantageous behavioral responses. The commonalities and differences of these

representations across species are not well-understood. Here, we show that social isolation

activates an oxytocinergic (OXT), nociceptive circuit in the larval zebrafish hypothalamus and

that chemical cues released from conspecific animals are potent modulators of this circuit's

activity. We delineate an olfactory to subpallial pathway that transmits chemical social cues to

OXT circuitry, where they are transformed into diverse outputs simultaneously regulating

defensive and feeding behaviors. Our data allow us to propose a model through which social

stimuli are integrated within a fundamental neural circuit to mediate diverse adaptive

behaviours.

INTRODUCTION

In mammals, signaling in oxytocinergic (OXT) circuits modulates a wide spectrum of socially driven behaviors, ranging from pair bonding and parental care to responses to stress and pain[1–3]. OXT has also been described as a potent appetite suppressant[4,5]. We and others previously reported that larval zebrafish OXT neurons encode a response to aversive, especially noxious stimuli, and directly drive nocifensive behavior via brainstem premotor targets[6,7]. Moreover, studies in both zebrafish[6–8] and mammals[9,10] suggest that the highly-conserved[11] OXT-expressing neuronal population is anatomically and functionally diverse and might modulate multiple behaviors.

The optically-transparent and genetically-tractable larval zebrafish is a popular model for relating brain activity to behavioural output. Social cue sensing (visual and mechanosensory) and social interactions have been reported at these larval stages, although they tend to manifest as avoidance behaviors, with social attraction only developing after 2 weeks of age[12–15]. One exception is in the domain of olfaction, where olfactory detection and imprinting of chemical kin cues in zebrafish and other teleost fish have been reported as early as 6 **d**ays-**p**ost-**f**ertilization (dpf)[16], with animals already demonstrating a preference for kin chemical cues at these early ages[17,18]. Recent work suggests that Major Histocompatibility Complex (MHC) signaling via specific olfactory sensory neuron subtypes (e.g. crypt cells) could allow for the recognition of kin olfactory signals, and a downstream forebrain region, the subpallium, has been proposed to mediate such affiliative social behaviors[16–21].

Here, in a brain-wide screen[22] for neuronal populations whose activity reflects social context, we show that larval zebrafish oxytocinergic neurons display diverse responses to conspecific chemosensory stimuli and are key effectors for social context modulation of nociceptive and appetite-driven behaviors. We further describe an olfactory bulb to subpallium circuit that is responsive to conspecific cues and sufficient to modulate OXT neuronal activity. Our results reveal a simple algorithm by which neuromodulatory neurons can represent social context to exert flexible control over hard-wired behavioral drives.

RESULTS

Brain-wide activity mapping of social isolation and its rescue by conspecific chemical cues

Using **p**hosphorylated-**E**xtracellular signal-**R**egulated **K**inase (pERK) based whole-brain activity mapping (MAP-Mapping[22]), neural activity in brains of briefly (2 hrs) socially-isolated larvae (7 - 8 dpf) was compared to animals that had been maintained in the presence of similarly-aged conspecifics. We found that isolated fish showed an enhancement of neural activity in specific regions, including the telencephalon, hindbrain, locus coeruleus, area postrema, caudal hypothalamus, preoptic area (PO, homolog of the hypothalamic paraventricular nucleus in mammals) and posterior tuberculum (PT) (Figure 1a-b, Supplementary Movie 1, Supplementary Data 1). Many of these same regions were observed to be activated by noxious or aversive stimuli[6,7] and may thus represent the signature activity pattern of a negative internal state that can be likewise triggered by social deprivation. Further, these regions appear to scale their activity in a monotonic fashion with the negative valence of the stimulus[6] (Figure 1b), and might encode the state of social isolation analogously.

Neurons expressing the peptide oxytocin (OXT) are localized to the PO and PT[6,23,24] regions. Indeed, by measuring pERK activity for individual GFP-labeled OXT neurons (*Tg(oxt:GFP)*) in high resolution confocal microscopic images of dissected brains[6] (Figure 1c-d, Supplementary Figure 1), we found that OXT-positive neurons clustered in both the preoptic area (OXT_{PO}) and posterior tuberculum (OXT_{PT}) display significantly enhanced activity in socially-isolated fish. To further elucidate the nature of this social signal, we compared OXT neuronal activity in animals exposed to the separated visual or chemical cues of a social environment. Water conditioned by prior exposure to conspecific larval fish (see Methods) reduced the elevated $OXT_{PO/PT}$ neuronal activity observed in socially isolated fish, whereas visual exposure to conspecific larval fish (maintained in a separate water enclosure) was significantly less effective in reducing OXT_{PO} neural activity (Figure 1c-d). We also examined the effect of water conditioned with similarly-aged sibling (kin) fish in relation to similarly-aged fish of a different strain background (non-kin). Both 'kin' and 'non-kin' conditioned water were sufficient to reduce OXT_{PO} neuron activity (Figure 1c-d), but the effect of kin water was overall stronger (Figure 1d). Thus, our results show that hypothalamic OXT activity is increased during brief (2 hr) social isolation and that this enhanced activity is significantly (and rapidly; see Figure 2) suppressed by social cues. The OXT_{PO} cluster displays a particular preference to chemical (and non-visual) cues in water conditioned by conspecific larval fish (Figure 1c-f). OXT_{PT} neurons on the other hand showed stronger suppression by visual and non-kin cues (Figure 1c-f). Further, nearby OXT-negative neurons in the PO and PT area were also suppressed by conspecific-

103   conditioned cues, including visual cues, suggesting that they too may play a role in integrating

104   social responses (Supplementary Figure 1).

106   **FIGURE 1**

108   **Figure 1: phospho-ERK based mapping reveals OXT neuron modulation by social context**

(a) **Left:** pERK-based activity mapping comparing the brain activity of fish that have been isolated for 2 hours vs maintained in the presence of conspecifics. Green voxels highlight regions that are significantly more active in isolated fish as compared to fish maintained in groups. This final map was generated by averaging data from 5 independent experiments in which isolated fish are compared to fish kept in groups of 3 (2 experiments), 5 (1 experiment) or 10 (2 experiments) (See **Supplementary Movie 1**; FDR Threshold used = 0.05%. Yellow = Outline of preoptic (PO) and posterior (PT) OXT populations. These masks were used for the quantification of activity in **Right**. Tel = Telencephalon, cH = Caudal Hypothalamus, LC = Locus Coeruleus, AP = Area Postrema, HB = Hindbrain. A = Anterior, L = Left, V = Ventral. Scale bar = 100 µm. **Right:** Social isolation significantly activates PO and PT regions where OXT neurons reside. Here, the mean pERK signal was calculated per fish across the specified ROIs. Data for other stimuli was also included in Wee et al (2019)[6]. Adjusted p values for social isolation: ***p = 0.00084 (PO), *p = 0.03 (PT), p = 0.42 (whole brain), Wilcoxon Signed-Rank test relative to a median of 1, Bonferroni correction. Other ROIs are quantified in **Supplementary Data 1.**

(b) High-resolution imaging of pERK expression in OXT and surrounding neurons, from dissected isolated fish brains **(bottom)** and compared to brains of fish kept in groups of 3 **(top)**. The maximum intensity projection of one example fish from each condition are shown (green = *Tg(oxt:GFP)*-positive neurons, magenta = anti-pERK staining). White arrows indicate examples of OXT neurons with high pERK intensities. Scale bar = 20 µm.

(c) Effect of social isolation and social sensory cues on mean OXT_{PO} and OXT_{PT} neuron activity per fish. pERK intensities shown are normalized to the respective mean PO or PT intensities of control fish (i.e. fish maintained in groups of 3). Social isolation (n = 11 fish) induced a 1.2 fold increase in OXT_{PO} and OXT_{PT} neuron pERK activity relative to fish maintained in groups of 3 (n = 12 fish). Visual cues of conspecifics moderately reduced OXT_{PO} neuron pERK activity (1.1 fold relative to group, n = 12 fish), non-kin and kin water cues produced an even greater reduction in OXT_{PO} neuron pERK activity (1.01 and 1.00 fold relative to group, n = 11 and 12 fish respectively). Notably, all social cues were sufficient to rescue OXT_{PT} activity to control levels, with visual (0.95 fold relative to group) and non-kin (0.89 fold relative to control) cues having a larger effect as compared to kin odor (1.05 fold relative to group).

OXT_{PO} neurons: p= 0.0019** (group vs isolated) / 0.010* (group vs visual) / 0.15 (group vs non-kin water) / 0.78 (group vs kin water) / 0.47 (isolated vs visual) / 0.15 (isolated vs non-kin water) / 0.066 (isolated vs kin water) / 0. 23 (visual vs non-kin water) / 0.053 (visual vs kin water) / 0.60 (kin vs non-kin water). Two-sided Wilcoxon Rank-Sum test.

OXT_{PT} neurons: p= 0.018* (group vs isolated) / 0.623 (group vs visual) / 0.10 (group vs non-kin water) / 0.644 (group vs kin water) / 0.0023** (isolated vs visual) / 0.0039** (isolated vs non-kin water) / 0.030* (isolated vs kin water) / 0. 16 (visual vs non-kin water) / 0.93 (visual vs kin water) / 0.32 (kin vs non-kin water). Two-sided Wilcoxon Rank-Sum test.

(d) Probability distributions and box plots showing the median, inter-quartile range and 95% confidence interval of normalized pERK fluorescence across all neurons per group. pERK fluorescence was divided by the mean pERK fluorescence of all OXT neurons (PO and PT combined) of the control group. Fish were either kept in groups of 3 (gray, n = 835 OXT_{PO} / 158 OXT_{PT} neurons from 12 fish), isolated (pink, n = 803 OXT_{PO} / 128 OXT_{PT} neurons from 11 fish), isolated but exposed to visual cues of conspecifics through a transparent barrier (red, n = 796 OXT_{PO} / 106 OXT_{PT} neurons from 12 fish) or isolated but exposed to non-kin-conditioned water

(blue, n =751 OXT$_{PO}$ / 115 OXT$_{PT}$ neurons from 11 fish) or kin-conditioned water (orange, n= 722
OXT$_{PO}$ / 134 OXT$_{PT}$ neurons from 12 fish).

OXT$_{PO}$ neurons: p = 6.9x10^{-42}*** (group vs isolated) / 1.3x10^{-24}*** (group vs visual) / 8.5x10^{-6}***
(group vs non-kin water) / 0.86 (group vs kin water) / 0.0060** (isolated vs visual) / 1.6x10^{-18}***
(isolated vs non-kin water) /   2.1x10^{-34}*** (isolated vs kin water) / 4.6x10^{-21}*** (visual vs non-kin
water) / 4.6x10^{-21}***(visual vs kin water) / 7.4x10^{-5}*** (kin vs non-kin water). Two-sided Wilcoxon
Rank-Sum test.

OXT$_{PT}$ neurons: p = 1.9x10^{-82}*** (group vs isolated) / 2.3x10^{-4}*** (group vs visual) / 7.8x10^{-8}***(group vs non-kin water) / 0.025* (group vs kin water) / 1.3x10^{-15}*** (isolated vs visual) /
169
1.9x10^{-18}*** (isolated vs non-kin water) / 3.6x10^{-10}*** (isolated vs kin water) / 0.052 (visual vs
non-kin water) / 0.30 (visual vs kin water) / 0.0069** (kin vs non-kin water). Two-sided Wilcoxon
Rank-Sum test.

(e) Spatial distribution of OXT neurons from each category, color coded according to pERK intensity
(most active = yellow, least active = deep blue). 45 OXT neurons were randomly sampled per
fish, with 495 neurons were drawn from this sample, to ensure even representation across all fish
and groups.

(f)   Anterior-posterior (AP) localization of "active cells" sampled from each category. These are cells
that have normalized pERK values surpassing the threshold by which only 10% of control (PO
and PT combined) cells would be considered active. To account for sample size differences,
these "active cells" were randomly drawn from the same set of 495 neurons in **(e).**

In vivo calcium imaging reveals diverse OXT neuronal responses to chemical conspecific cues

We next turned to *in vivo* calcium imaging to acquire a more temporally precise record of OXT

neuronal activity. Imaging was performed on larvae in which *Tg(UAS:GCaMP6s)* was driven

directly in OXT neurons with an *Tg(oxt:Gal4)* driver (Figure 2a). Tethered 8-11 dpf fish were

subjected to either conspecific-conditioned or control (unconditioned) water released in 10

second pulses. We found that control water flow presents a mechanosensory stimulus that

mildly activates OXT neurons, whereas kin-conditioned water triggers an immediate reduction of

OXT neuronal activity (Figure 2a-b). We did not observe any difference in the effectiveness of

kin conditioned water with respect to social isolation time (from 30 min to 4 hours) or age range

(8-11 dpf) (Supplementary Figure 2a-b). We next examined whether OXT neuron calcium

activity displays distinctions with respect to kin and non-kin conspecific water-borne cues[16,19].

Similar to our pERK experiments in Figure 1, OXT neuronal responses were compared between

sibling-conditioned water ('kin water') and water conditioned by larvae of a distinct genetic

background[25] ('non-kin water'). To minimize the effects of familiarity, conspecifics used to

generate conditioned cues were raised apart from experimental fish. Kin water was found to

induce a greater reduction in OXT$_{PO}$ activity compared to non-kin water. We also examined

water conditioned by incubation with adult kin, which is a potential aversive cue since adult

zebrafish consume their own young[26,27]. In contrast to larval-conditioned water, adult

conditioned water induced a substantial increase in $OXT_{PO/PT}$ activity.

FIGURE 2

Figure 2: Calcium imaging confirms OXT neuron modulation by conspecific cues

(a) **Left:** Maximum z-projection of volumetrically-scanned oxytocin neurons. The *Tg(oxt:Gal4;*
UAS:GCaMP6s) transgenic line labels OXT_{PO} neurons strongly, with weaker labeling of the more
ventro-posteriorly located OXT_{PT} cluster. Scale bar = 20 µm. **Right:** Mean OXT neuron
fluorescence integrated over a 60 s period post-stimulus from two different fish, one imaged with
kin and non-kin water **(top)**, and another with kin and adult water **(bottom)**.

(b) **Left:** Mean stimulus-triggered calcium responses (Δf/f) of OXT_{PO} (top) and OXT_{PT} (bottom)
neurons. All fish (29 fish, n = 1033 neurons) were imaged with water or kin water, some
additionally were imaged with either non-kin (8 fish, n = 245 neurons) or adult water (10 fish, n =
374 neurons). Of all neurons imaged, only 10 were from the OXT_{PT} population. **Right:** Mean

218 stimulus-triggered calcium responses of each neuron normalized to its own mean water
219 response. Gray broken line indicates stimulus onset. Shading indicates SEM.
220

221 (c) Spatial distribution and percentages of neurons that show either suppressed (**top**) or enhanced
222 (**bottom**) responses to each cue relative to water. See **Methods** and **Supplementary Figure 2c**
223 for details. For accurate comparison of spatial distributions, we randomly sampled 25 neurons per
224 fish, then 200 neurons were randomly selected from each group.
225

226 (d) Mean stimulus-triggered calcium responses (Δf/f) of neurons classified as being suppressed (top)
227 or enhanced (bottom) by each cue, normalized to their mean water response. Shading indicates
228 SEM.
229

230 (e) Visualization of overlapping OXT neuron representations of conspecific cues. Left: overlap
231 between kin and non-kin water responses. Right: overlap between kin and adult water responses.
232 We did not image all 3 cues simultaneously. Red = enhancement, Blue = suppression, White =
233 no change. Kin cues induced the highest percentage suppression (25-30%) and lowest
234 percentage activation (10-11%), whereas adult cues induced the lowest percentage suppression
235 (14%) and highest percentage activation (40%) of OXT neurons. Approximately one third of
236 neurons suppressed by kin water (blue) were also suppressed by non-kin (40%) and by adult
237 water (37%). In addition, a common subset of neurons activated by kin water (red) were also
238 activated by non-kin (68%) and adult (95%) water. However, 22% and 36% of kin water-
239 suppressed neurons were alternatively activated by non-kin and adult-conditioned water,
240 respectively. See **Supplementary Figure 2d-e** for additional visualization.
241

242 OXT neurons are diverse in their circuit connectivities[8,28–30] and responses to social

243 isolation (Figure 1) and nociceptive input[6]. To further resolve this heterogeneity, we classified

244 individual OXT neurons into populations that showed either reduced (blue) or increased (red)

245 calcium activity in response to water-borne conspecific cues (Figure 2c-e). Across a range of

246 thresholds, the fraction of OXT-positive neurons whose activities were suppressed by larval kin-

247 conditioned water was consistently higher than the fraction of neurons that were activated

248 (Supplementary Figure 2c). Non-kin-conditioned water, in contrast, induced equivalent neuronal

249 fractions with increased or suppressed activities (Supplementary Figure 2c). Water conditioned

250 by adult fish triggered a greater fraction of neurons with enhanced activity (Supplementary

251 Figure 2c). Under all conditions, activated or suppressed neurons were spatially distributed

252 instead of being segregated into distinct areas (Figure 2c). Further, the specific response

253 properties of a neuron to one conspecific cue did not strongly predict how it would respond to

254 the other cues, although we observed that kin water-activated OXT neurons tended to also be

255 commonly activated by other conspecific (non-kin or adult) cues (Figure 2e, Supplementary

256 Figure 2d-e). The response heterogeneity of the OXT population suggests that these neurons

257 differentially encode conspecific stimuli.

FIGURE 3

FIGURE 3: Calcium imaging reveals olfactory bulb encoding of conspecific stimuli

(a) **Left:** Schematic of imaging and cue delivery setup. Two-photon (2P) calcium imaging of the olfactory bulb was performed as each *Tg(HuC:GCaMP6s)* fish was presented with pulses of kin water, non-kin water and adult water (randomized order), alternated with pulses of water. **Right:** Stimulus-triggered average of all olfactory bulb (OB) neuron responses (Δf/f) to kin, non-kin and adult water.

(b) Δf/f traces (**bottom,** raster plot for an example fish) from individual units (see **Methods**) within the OB were correlated with regressors (**top**) fit to each stimulus type.

271   (c) Mean OB Δf/f integrated over a 60 s period post-stimulus from a single fish (same fish shown in
272   **Figure 3b,** more example fish in **Supplementary Figure 3**), shown for each plane (ventral to
273   dorsal) and each cue in a volumetric stack. **Bottom-most panel** shows the anatomy references
274   for each plane. A = Anterior, L = Left. Scale bar = 50 μm.

276   (d) Δf/f traces from individual units were extracted from the OB (see **Methods**). **Top:** Correlation
277   coefficients for each unit (n = 477 units from the same fish in **b** and **c**) with each regressor are
278   color coded (blue to red) and overlaid over the anatomy image per z-plane. **Bottom:** Each unit's
279   response (i.e. correlation coefficient) to water and conspecific stimuli, from the same fish as
280   above, clustered using K-means clustering (as in **e**).

282   (e) K-means clustering (k = 12) was performed on the matrix of correlation coefficients (color coded
283   blue to red) to water and conspecific stimuli (4018 units from 10 fish). Number of units within each
284   cluster and percentage representation of total units are displayed on the right. Clusters specific to
285   kin water and adult water, as well as clusters with mixed selectivities were observed.

287   (f) Mean stimulus-triggered calcium activity (Δf/f) for each cluster, aligned to stimulus onset. Order of
288   clusters are the same as in (e). As OB responses are often sustained across the entire stimulus
289   epoch, negative Δf/f and correlation coefficients in response to water flow are likely due to wash-
290   out of residual olfactants, rather than an active suppression of the signals by water flow.

292   (g) 3D plot displaying the spatial localization of units within each cluster, for a few selected clusters
293   (all clusters in **Supplementary Figure 3c**). XY coordinates for all units per fish were scaled
294   linearly to their minimum and maximum values in each dimension. Adult-responsive clusters tend
295   to be situated laterally (e.g. cluster 7 and 8), whereas clusters that show more general conspecific
296   responses (e.g. clusters 4,5,10) are situated more medially. Kin-specific responses (clusters 1
297   and 2) are more scattered, though they extend more laterally than general conspecific-responsive
298   clusters. See also **Supplementary Figure 3b.** A = Anterior, P = Posterior, D = Dorsal, V =
299   Ventral.

The zebrafish olfactory bulb and downstream subpallium encode conspecific chemical cues

Since water-borne chemical cues might act as odorants, we included the olfactory bulb (OB) in

the areas specifically interrogated for changes in neuronal activity upon exposure to chemical

conspecific cues that modulate OXT neuron activity. We found that each of three conditioned

water types (kin, non-kin and adult kin) generated overlapping but distinct activity signatures

within the OB (Figure 3a-c, Supplementary Figure 3). Of all the cues, adult water had the

strongest and longest-lasting effect (Figure 3a-b, f), as was also noted for OXT neuronal

responses with this stimulus (Figure 2b). Notably, non-kin water at the same concentration

elicited much weaker OB activity than kin water, consistent with the prior observations of its

weaker effect in suppressing OXT neuron activity (Figures 1d and 2).

To characterize the relationships in OB encoding of the different social stimuli, we

extracted individual units[31] and performed k-means clustering (k = 12) based on their correlation

coefficients with stimulus regressors (see Methods). We identified clusters that were exclusive

to specific social cues as well as clusters with mixed selectivity. Consistent with previous

reports[20], kin-selective (as well as adult-selective clusters) tended to be located more laterally,

while non-selective social cue-responsive clusters were situated medially (examples shown in

Figure 3g; all clusters shown in Supplementary Figure 3).

The zebrafish telencephalon contains multiple GABAergic populations that receive

olfactory input[21,32,33] and hence might transform excitatory input from the OB into inhibitory

signals to the OXT population. We focused on the subpallium (SPa) in particular because it is a

known downstream target of the olfactory bulb[21,32,33] and implicated in social behavior[34,35] and

kin odor recognition[21]. To this end, we imaged neuronal activity in the telencephalon in the pan-

neuronal *Tg(HuC:GCaMP6s)* line in which GABAergic neurons were also fluorescently marked

with red fluorescent protein (*Tg(Gad1b:RFP)*; Supplementary Figures 4-5). We also imaged

activity in the preoptic area of fish in which OXT neurons are labeled with mCherry

(Supplementary Figure 5). The SPa can be broadly divided into anterior (aSPa) and posterior

(pSPa) domains; both of these subpallial domains, including their GABA-positive units, were

responsive to conspecific cues (Supplementary Figures 4-5). Using k-means clustering as

above (k = 12), we identified neuronal clusters responsive to distinct social cues, notably with

more kin- and adult kin-responsive clusters compared to non-kin-responsive clusters

(Supplementary Figures 4-5). In contrast to the SPa, larval conspecific cues reduced OXT_{PO}

neuronal activity (Supplementary Figure 5). These observations are consistent with a role of the

SPa as an intermediary in the olfactory modulation of OXT neuronal activity.

A GABAergic subpallial population encodes conspecific chemical cues and suppresses oxytocin

neuron activity

To demonstrate a circuit pathway through which OB-encoded social cues might be conveyed to

the OXT population, we sought Gal4 transgenic lines with which to perturb neuronal activity in

the subpallium. The *Tg(y321:Gal4)* enhancer trap specifically labels the aSPa population and

has been previously implicated in visually-mediated social behavior and social behavior

development in larval zebrafish[34,35]. Co-labeling y321:Gal4-positive and Gad1b-positive cells in

double transgenics revealed that, among y321-labeled neurons, 88% in the aSPA (aSPa[y321])

and 87% in the pSPa (pSPa[y321]) are GABAergic (n = 7 fish, Figure 4a). Moreover, some OXT

neurons extend dendrites into the SPa, establishing potential connectivity with the SPa[y321]

population (Figure 4c). Analysis of activity via pERK staining (similar to Figure 1) revealed

stronger responses of both aSPa[y321] and pSPa[y321] neurons to kin water, as compared to non-kin

water (Supplementary Figure 6a-b). In addition, calcium imaging of the aSPa[y321] neurons'

response to conspecific cues (Figure 4d-f) revealed clusters of aSPa[y321] neurons that are

differentially responsive to social cues, with more clusters responsive to kin and adult water-

borne cues, consistent with results from pan-neuronal and GABAergic neuronal activity imaging

(Supplementary Figures 4-5).

These correlative results suggested a possible mechanism by which kin cues are

transformed by subpallial GABA neurons into an inhibitory signal that suppresses OXT neuronal

activity. To establish a functional relationship between these two populations, we

optogenetically stimulated y321:Gal4-positive neurons via spatially-restricted laser pulse

illumination of red-shifted channelrhodopsin (ReaChR[36]) and concurrently recorded calcium

sensor (GCaMP6s) fluorescence in OXT neurons (Figure 4g). In control experiments, with

sibling fish lacking the ReaChR transgene, laser illumination of the subpallium induced a small

increase in y321:Gal4 neuronal activity (Figure 4g-h, top rows, Supplementary Figure 6c),

indicating that there is some calcium response in the absence of ReaChR. When both the

ReaChR and y321:Gal4 transgenes were present, y321-positive neuronal activity was strongly

induced and phase-locked to the onset of the laser pulse, confirming the efficacy of the

optogenetic protocol.

We then examined OXT neuronal activity after spatially-targeted aSPa illumination in fish

harboring the y321:Gal4, oxt:Gal4, UAS:ReaChR-RFP and UAS:GCaMP6s transgenes. These

animals express ReaChR and GCaMP6 in both the preoptic OXT population and the y321-

positive subpallium neurons. OXT neuronal activity in these animals was compared to control

siblings lacking the y321:Gal4 transgene; hence lacking ReaChR expression in the subpallium

(Figure 4g-h). In these negative control fish, laser stimulation of the subpallium was associated

with sporadic OXT neuronal activity and moderate reductions in OXT neuropil activity (blue

arrows in bottom traces, Figure 4g). We consider it possible that these effects reflect

optogenetic activation of ReaChR in OXT neuron dendrites that extend into the subpallium

(Figure 4c). In contrast, with y321:Gal4 present and ReaChR expressed in subpallial neurons,

optogenetic stimulation of the aSPa resulted in a dramatic and significant reduction in OXT

neuronal activity, specifically in OXT_{PO} cell bodies and the OXT neuropil (OXT axons) (Figure

4g-h, Supplementary Figure 6d). Thus our data strongly suggest that GABAergic aSPa[y321]

neurons transform excitatory input from conspecific cue activation of the olfactory bulb into an

inhibitory signal for OXT neuronal activity.

379
380
381
382

FIGURE 4

FIGURE 4: A genetically-defined GABAergic subpallial population discriminates social cues and suppresses OXT activity

(a) Schematic depicting the hypothesized telencephalic subpallial (SPa) pathway linking olfactory bulb conspecific cue responses to preoptic OXT neurons.

(b) **Top Left:** Maximum projection image showing overlap of *Tg(y321:Gal4;UAS:Kaede)* (green) with GABAergic neurons labeled by *Tg(Gad1b:RFP)* (magenta) in the telencephalon. y321 and Gad1b co-expression occurs in both the anterior subpallium (aSPa, rostral to the anterior commissure) and posterior commissure (pSpa, caudal to the anterior commissure and rostral to the preoptic area). **Top Right:** Lower magnification maximum projection image showing overlap of *Tg(y321:Gal4;UAS:Kaede)* with *Tg(Gad1b:RFP)* (magenta). Overall, 88.2 ± 1.5 % of aSPa^{y321} and 87.3 ± 3.06 % of pSPa^{y321} neurons are GABAergic (n = 7 fish. Tel = Telencephalon, PO = Preoptic area; PT = posterior tuberculum. **Bottom:** y321 and Gad1b overlap shown across different z-planes. All scale bars = 20 μm.

(c) OXT neuron dendrites project into the location where SPa^{y321} neurons reside. **Top:** Maximum projection image; **Bottom:** 3D projection at different rotation angles. Scale bar = 50 μm.

(d) 2P calcium imaging was performed on *Tg(y321:Gal4; UASGCaMP6s)* neurons in the aSPa region on fish exposed to kin, non-kin and adult water cues. See **Supplementary Figures 4-5** for imaging and analysis of the same telencephalic regions in *Tg(HuC:GCaMP6s; Gad1b:RFP fish).* **Top Left:** Maximum projection image showing the aSPa^{y321} region imaged. **Top Right:** Example Δf/f trace of a kin-responsive unit. **Bottom:** Stimulus triggered averages for units that are kin-selective, adult-selective, or responsive to multiple conspecific cues. Shaded region indicates SEM.

(e) K-means clustering (k = 12) was performed on the matrix of correlation coefficients (color coded blue to red) to water and conspecific stimuli (4875 units from 4 fish). Number of units within each cluster and percentage representation of total units are displayed on the right. Clusters specific to kin, non-kin, and adult water, as well as clusters with mixed selectivity were observed.

(f) Mean stimulus-triggered activity (Δf/f) for each cluster, aligned to stimulus onset. Order of clusters are the same as in (e).

(g) **Left:** Maximum projection image showing regions of optogenetic stimulation (aSPa^{y321} neurons) and calcium imaging (OXT neurons in PO or PT, or neuropil bundles in PT outlined in yellow). **Right:** Mean Δf/f traces from either aSPa^{y321} neurons or OXT cell bodies / neuropil bundles after stimulation with a 633 nm laser focused on the aSPa region at 60 s intervals. Black traces depict mean Δf/f of ReaChR-negative controls (top trace) or y321-negative controls (bottom traces). Red traces depict mean Δf/f from an example *Tg(y321:Gal4, oxt:Gal4; UAS:GCaMP6s; UAS:ReaChR-RFP)* fish of aSPa^{y321} neurons (top trace) or OXT$_{PO/PT}$ cell bodies (middle trace) or neuropil (bottom trace) to aSPa^{y321} optogenetic stimulation. Orange lines indicate stimulus onset. Each box represents calcium traces from an individual fish. Shaded region indicates SEM. Blue arrows indicate dips in OXT neuropil activity in the presence or absence of optogenetic y321 activation.

(h) Stimulus-triggered averages showing the mean Δf/f trace 50 s before and after the light stimulus. Responses to each stimulus were first averaged per unit (cell body or neuropil bundle) before being averaged across all units to obtain the Δf/f trace. Shaded region indicates SEM. **Top panel:** The calcium response (Δf/f) of aSPa^{y321} neurons to 633 nm laser stimulation is significantly higher in the presence (red, n = 320 neurons from 15 fish) than absence (black, n = 118 neurons from 4 fish) of ReaChR (*p = 0.014, Two-sided Wilcoxon Rank-Sum Test). **Middle panel:** The calcium response of OXT$_{PO/PT}$ neurons under aSPa^{y321} optogenetic stimulation is significantly lower (***p $=1.2 \times 10^{-5}$, n = 410 neurons from 18 fish) than for negative controls (*Tg(oxt:Gal4; UAS:GCaMP6s; UAS:ReaChR-RFP) fish*) exposed to the same light stimulus (n = 103 neurons from 5 fish). When

we divided these OXT neurons according to their anterior-posterior position, we found that OXT_{PO}
neurons showed significant suppression after $aSPa^{y321}$ optogenetic stimulation compared to
negative controls (***p =0.00025, n = 294 / 91 neurons), but no significant difference was
observed for OXT_{PT} neurons (OXT_{PT}; p = 0.397, n = 116 / 12 neurons). **Bottom panel:** OXT
neuropil calcium responses under y321 optogenetic stimulation are significantly lower than for
negative controls exposed to the same light stimulus (***p = 9.6 x10^{-5}, n = 36 / 10 neuropil
regions).
447

Conspecific cues suppress nociceptive OXT circuits and defensive behavior

Social buffering is a widely-observed phenomenon in which the presence of conspecifics

ameliorates the effects of aversive experience[37,38]. We previously showed that a large fraction of

OXT neurons are activated by noxious stimuli and drive defensive behaviors, specifically

through brainstem premotor targets that trigger vigorous large-angle tail bends[6]. Given that most

of the OXT circuitry appears to be suppressed by water-borne social cues derived from closely-

related conspecifics, we posited that these cues might reduce the nocifensive behavior induced

by the stimulation of TRPA1 receptors. Indeed, increased swim speed triggered by nociceptive

TRPA1 receptor activation was significantly ameliorated by the presence of kin-conditioned

water (Supplementary Figure 7). Similarly, in 8-10 dpf tethered fish, kin water significantly

reduced the frequency of large-angle swims in tethered fish after kin cue delivery and TRPA1

stimulation but not during other epochs (Figure 5a, Supplementary Figure 8a). Correspondingly,

the average calcium responses of individual OXT_{PO} (but not OXT_{PT}) neurons following TRPA1

stimulation was significantly reduced when preceded by kin-conditioned water (Figure 5b).

We next examined individual OXT neuron responses to TRPA1 stimulation in order to

resolve the social cue modulation of their responses. As previously done[6], TRPA1-responsive

OXT neurons were identified by correlating each neuron's calcium activity with TRPA1-stimulus

and motor output regressors (Figure 5c). Motor regressors either reflected movements that

occurred in temporal association with (within 5 seconds of) the TRPA1 stimulus ($motor_{stim}$) or at

times outside of the stimulus period ($motor_{spon}$). For spontaneous swim bouts outside of the

stimulus period, we further generated regressors for small ($motor_{spon-S}$) and large-angle

($motor_{spon-L}$) tail bends to examine whether neurons tuned to large angle vs small angle

movements may be differentially represented by the OXT population.

FIGURE 5

Figure 5: Chemical kin cues modulate OXT neuron nociceptive responses and behavior

(a) **Top:** Schematic showing setup to probe effect of conspecific-conditioned water on TRPA1-
induced nocifensive responses. Fish were tethered with their nose/mouth and tails freed, and
incubated in 15 µM Optovin solution. **Bottom:** Kin water significantly reduced the frequency of
large angle (>50°) but not small (<= 50°) angle tail bends, both after cue delivery and post-TRPA1
stimulation, though overall bout frequency was much higher after TRPA1 stimulation (bottom-
most row). Post-cue delivery: p = 0.011* (all bouts) / 0.0039** (large angle) / 0.93 (small angle).
Post-TRPA1: p = 0.045* (all bouts) / 0.019* (large angle) / 0.86 (small angle). There was no
significant change in bout kinematics during epochs outside of kin cue or TRPA1 delivery (p =
0.65 / 0.55 / 0.71 for total, large and small tail bends respectively). One -sided Wilcoxon Signed-
Rank test, n = 16 fish. More kinematic features are shown in **Supplementary Figure 8a.**

(b) **Top:** Alternating pulses of water or conspecific water were presented, followed by a 100 ms pulse
of UV light after 30 s to activate TRPA1 receptors. Half of the fish had kin water as the first
stimulus.
494

Bottom: Stimulus-triggered averages (Δf/f) for OXT_{PO} (n = 492) or OXT_{PT} (n = 8) neuron calcium
activity in response to a water pulse or kin water delivery. The integrated calcium response (Δf/f)
of OXT_{PO} neurons both post-cue delivery (***p = 1.3×10^{-12}) as well as post-TRPA1 stimulus
(*p=0.0129) was significantly lower with kin water as compared to water flow. There was no
significant effect of kin water on OXT_{PT} neuron activity both post-cue delivery (p = 0.63) as well as
post-TRPA1 stimulus (p = 0.63), one-sided Wilcoxon Signed-Rank test. Black horizontal line
shows the region over which pre- and post-TRPA1 calcium activity and behavior were integrated
to calculate pre-cue, post-cue and post-TRPA1 responses. Gray dashed line = water or kin water
delivery, purple dashed line = UV stimulus onset, Shading indicates SEM.
504

505   (c) Calcium traces (Δf/f) and respective stimulus and motor regressors (see **Results** and **Methods**),
506   as well as their correlation coefficients with each regressor are shown for two example OXT_{PO}
507   neurons.
508

(d) To characterize the diversity of $OXT_{PO/PT}$ neuron responses to TRPA1 stimulation, K-means
clustering (k = 12) was performed on the matrix of correlation coefficients to each of the stimulus
or motor regressors (n = 500 units from 16 fish). Number of units within each cluster and
percentage representation of total units are displayed on the right. Clusters specific to TRPA1
stimulation, as well as motor-correlated clusters were observed. We grouped these clusters into
broader classes and subclasses (i-v) which are indicated by the black lines and boxes. See also
Supplementary Figure 8c-d.
516

517   (e) The percentage of $OXT_{PO/PT}$ neurons within each cluster that were suppressed, activated, or did
518   not show any change when exposed to kin water, either following cue delivery (**left**) or TRPA1
519   stimulation (**right**). Red = enhancement, Blue = suppression, White = no change. The broader
520   classes (i-v) are indicated by black boxes. See also **Supplementary Figure 8b**.
521

(f) Mean stimulus-triggered calcium responses (Δf/f) of all $OXT_{PO/PT}$ neurons in the presence of
water (black) or kin water (orange)**,** as a function of their class / subclass. Note scale bar for class
i_a is different from the others due to intensity of TRPA1 responses. For class i_b, the integrated
calcium response both post-cue delivery (**p = 0.0015) and post-TRPA1 stimulus (***p=1.0×10^{-5})
was significantly lower in the presence of kin water as compared to water flow. For classes ii_{a-b},
the integrated calcium response both post-cue delivery (***p = 6.6×10^{-4} or ***p = 3.8×10^{-7}) was
significantly lower with kin water as compared to water flow, and the response to TRPA1 stimulus
was only significantly lower for class ii_b (p = 0.73 or *p = 0.032 respectively), one -sided Wilcoxon
Signed-Rank test, n = 500 neurons.
531

As expected, OXT neurons displayed diverse activity patterns and varying degrees of

correlation with stimulus and motor regressors. Neurons tuned to the TRPA1 stimulus tended to

be correlated with motor behavior that followed the TRPA1 stimulus (Figure 5c-d,

Supplementary Figure 8d). Given the diverse responses of OXT neurons, we utilized k-means

clustering (k = 12) to classify neurons on the basis of correlated activity with different

regressors. This analysis uncovered 5 broad activity classes of OXT neurons (Figure 5d): (i)

TRPA1-selective neurons, with varying strength of correlation with the stimulus regressors; (ii)

neurons that were both TRPA1-responsive as well as correlated with spontaneous movements

(iii) neurons that were primarily motor-correlated, particularly to large-angle movements; (iv)

unresponsive neurons; and (v) neurons that showed an anti-correlated relationship to the

TRPA1 stimulus.

Individual OXT neurons were also grouped according to whether kin water presentation

enhanced or suppressed their responses to TRPA1 stimulation (as performed in Figure 2, see

also Supplementary Figure 8b). We found that the majority (54%) of OXT neurons displayed

reduced TRPA1-induced activity in the presence of kin water, with only 32% showing enhanced

activity, indicating that kin water exposure triggers widespread negative modulation of OXT

neuronal responses to TRPA1 stimulation. Notably, 43% of OXT neurons already showed a

reduction in activity post kin-water delivery and prior to TRPA1 stimulation, while 30% displayed

increased activity. While activity suppression by kin water occurred throughout the OXT

population (Figure 5e), there were also varying ratios of suppressed versus activated neurons

within each cluster and class, indicating that subsets of OXT neurons with distinct response

characteristics are differentially modulated by kin water (Figure 5e).

From the above analyses we derive two main observations. First, moderately TRPA1-

responsive OXT neurons (class i_b , Figure 5e, 5f; 25% of OXT neurons) displayed greater

suppression by kin water than highly TRPA1-responsive neurons (class i_a, Figure 5e, 5f; ~3% of

OXT neurons; Figure 5e, 5f). However, this relationship did not extend to weakly TRPA1-

responsive OXT neurons, which were not modulated by kin water (class i_c, Figure 5e, 5f;~8% of

OXT neurons). The average calcium responses of the moderately TRPA1-responsive (class i_b)

neurons displayed a significant divergence from baseline activity before TRPA1 activation

(Figure 5f), suggesting that these OXT neurons respond to both water flow and TRPA1

stimulation, both of which could be significantly suppressed by kin cues. A second observation

is that OXT neurons whose activity was strongly correlated with both the TRPA1 stimulus and

small or large spontaneous tail movements (class ii, representing ~31% of OXT neurons) were

more strongly suppressed by kin water, even prior to the onset of the TRPA1 stimulus. The fact

that spontaneous tail movements prior to the TRPA1 stimulus were reduced (Figure 5a)

suggests these motor-correlated suppressed neurons could play the role of reducing motor

responses generally in the context of an aversive cue. Overall, our results demonstrate the

widespread suppressive effects of kin water on OXT neuron activity and OXT-triggered

behavior. Both kin water-suppressed and activated neurons were scattered along the AP axis

(Supplementary Figure 8c), and a high proportion were located posteriorly, indicating that at

least some of the suppressed neurons have a parvocellular, hindbrain identity and may directly

drive motor output[6]. Thus, the reduced nocifensive responses of larval fish observed in the

presence of social cues is directly reflected in the suppressed activity of a large subset of

TRPA1-responsive neurons within the OXT circuit.

Larval conspecific water-borne cues enhance appetite via suppression of OXT neuronal activity

Social isolation is not only associated with increases in aversive behavioral responses, but is

also reportedly associated with reduced positive-valence behaviors such as feeding[39]. To further

characterize the diverse behaviors subject to social modulation in larval zebrafish, we used a

quantitative food intake assay to determine whether social isolation diminishes feeding

behavior[40–42] (Figure 6a). In this assay, larval fish are deprived of food for a period of two hours,

which results in increased appetite[40–42]. This appetite modulation protocol was performed on fish

maintained either in isolation or in the presence of conspecifics (Figure 6a). Subsequently,

animals were presented with a large excess of fluorescently-labeled paramecia, which reveals

food ingestion on the basis of gut fluorescence quantitation after a short feeding period. We

found that food-deprived fish maintained in isolation consumed significantly less paramecia than

those maintained in groups of three animals (Figure 6b). When group size was varied from 2 to

5 individuals, the food intake per animal scaled with group size (Figure 6b). Water-borne cues

from closely related larval fish (kin), but not visual contact with kin larvae, were sufficient to

rescue the isolation-induced suppression of food intake (Figure 6c).

Given that social isolation increases neuronal activity in a subset of OXT-positive

neurons, we asked if OXT neurons are required for the social control of appetite. To this end,

OXT neurons were chemogenetically ablated in larvae via OXT neuron-specific expression of

bacterial-nitroreductase (NTR)[43]. Animals were incubated with a group of conspecifics along

with the prodrug metronidazole (MTZ), which results in the death of NTR-expressing cells,

during the period of from 5 to 7 dpf. This treatment resulted in loss of ~80% of NTR-expressing

preoptic OXT neurons. At 8 dpf, these larvae, or MTZ-treated control animals lacking the NTR

gene, were placed in either into small groups or isolated and then assayed for feeding as

described above. We found that OXT neuron ablation enhanced food intake in isolated animals

while the food intake of fish maintained in groups was unchanged (Figure 6d). Consistently,

addition of an OXT receptor antagonist strongly increased food intake in isolated fish (Figure

6e), whereas OXT agonists strongly suppressed food intake of fish kept in a group (Figure 6f).

These results indicate that OXT signaling is both necessary and sufficient to mediate the social
modulation of food intake in larval zebrafish.
606
607 **FIGURE 6**

608
609 **Figure 6: Manipulation of OXT signaling affects appetite in a social-context dependent manner**
610
611   (a) Schematic depiction of social feeding behavior experiments. Larval zebrafish are raised in a
612   group to 7 or 8 dpf. They are then isolated in single dishes or placed into dishes with a group of
613   conspecific fish of the same age. After 2 hrs, fluorescently-labeled paramecia are added for 30
614   min, after which gut fluorescence was quantified microscopically. Example images are shown
615   (inset shows higher (100x) magnification).
616
(b) **Left:** Gut fluorescence in a single experiment demonstrates that isolated fish, on average, ingest
less paramecia than fish maintained in a group of 3 conspecific fish. All gut fluorescence
measurements are normalized to the group mean (**p = 0.0058, n = 21/17 fish, two-sided

Wilcoxon Rank-Sum test). **Right:** Normalized food intake (gut fluorescence normalized to the
average for groups of 3 fish) scales with group size (1 fish, n=36; 2 fish, n=7; 3 fish, n=52; 5 fish,
n=29). Single asterisks depict p<0.05 using one-way ANOVA corrected for multiple comparisons).

(c)   Water-borne but not visual cues rescue reduced feeding of isolated fish. Fish were food-deprived
       and incubated alone or in groups, as in (a). For provision of odor cues, conditioned water was
       added to dishes containing isolated larvae prior to supplying fluorescently labeled paramecia . To
       provide visual cues, ~5 larvae were placed in a large dish in which a smaller transparent dish was
       placed containing a single larva during isolation and feeding with labeled paramecia. **Left:**
       Results within a single experiment. *p = 0.03 (isolated vs group), *p = 0.04 (isolated vs kin water),
       p = 0.17 (isolated vs visual), n = 24/24/22/23 fish, two-sided Wilcoxon Rank-Sum test. **Right:**
       Average of multiple experiments. p = ***2.8283×10^{-4} (single vs group), *p = 0.02 (single vs kin
       water), p = 0.35 (single vs visual), n = 86/86/55/87, Two-sided Wilcoxon Rank-Sum test.

(d)   Cell-specific chemogenetic ablation of OXT neurons specifically rescues the effect of social
       isolation on appetite. OXT neurons were ablated via OXT neuron-specific expression of bacterial-
       nitroreductase using *Tg(oxt:Gal4;UAS:nfsb-mCherry)* fish[6,43]. Animals were incubated with a
       group of conspecifics and the prodrug metronidazole (MTZ) during the period from 5 to 7 dpf.
       This treatment resulted in loss of ~80% of nitroreductase-labeled preoptic OXT cells (unablated =
       20.3 ± 1.1 neurons, ablated = 4.5 ± 0.5 neurons; n = 20 control fish, 29 fish with ablation).

       *p = 0.015 (isolated control vs group control), **p = 0.0017 (isolated control vs single ablated), p =
       0.76 (isolated ablated vs group ablated), n = 65 /59/57/56 fish, two-sided Wilcoxon Rank-Sum
       test. Controls are metronidazole (MTZ)-treated, non-transgene-expressing siblings.

(e)   The Oxytocin receptor antagonist (L-368,899) restores social feeding levels to isolated animals.
       Antagonist (5uM) was added to the incubation water at the start of the 2 hr isolation period (see
       (a)).

       **p = 0.0022 (single vs group), **p = 0.0091 (single control vs single antagonist), p = 0.09 (single
       antagonist vs group antagonist), n=40/50/29/34 fish, two-sided Wilcoxon Rank-Sum test.

(f)   An Oxytocin receptor agonist (WAY267,484) (5 µM) reduces group food ingestion to the level of
       isolated animals. Agonist (5uM) was added to the incubation water at the start of the 2 hr
       isolation period (see (a)). **p = 0.0022 (single vs group), p = 1 (single control vs single agonist),
       **p = 0.0069 (group control vs group agonist), n=40/50/20/21 fish, two-sided Wilcoxon Rank-Sum
       test. Control groups for both **(e)** and **(f)** are the same sets of fish, split up for better visualization.
       All data is from multiple sets of experiments and normalized to food intake in a group of 3 fish.

(g)   Comparison of food intake of *oxt* null mutants (*oxt-/-*) and their heterozygous wild-type siblings
       (*oxt+/-*) in isolated and non-isolated contexts Single-group differences in food consumption are
       abolished -- however, food intake was reduced in groups, rather than enhanced in isolated fish.
       **p = 0.0011 (single vs group), p = 0.65 (single control vs single mutant), **p = 0.0012 (group
       control vs group mutant), n = 60/59/58/54 fish, two-sided Wilcoxon Rank-Sum test.

(h)   Model for how oxytocin neurons integrate information on social state to control appetite and
       nocifensive behaviors. We posit that social chemical cues are olfactory, and that GABAergic
       neurons in the subpallium transform olfactory bulb activation into inhibitory signals that

668   differentially modulate the OXT population. The OXT circuit modulates nocifensive behavior via
669   brainstem premotor neurons[6]. The OXT neurons project extensively to other areas of the
670   hypothalamus and the pituitary; these downstream regions may be involved in mediating effects
671   on appetite[42].
672

Lastly, we compared food intake in OXT homozygous null mutant animals with their

heterozygous siblings. As expected, we found that null mutant animals did not modulate their

food intake on the basis of social environment, whereas the behavior of their heterozygous

siblings was indistinguishable from that of wild type animals. However, mutant animals had low

food intake in both the isolated and the group setting (Figure 6g). Thus, as with acute ablation of

OXT neurons, social modulation of feeding was impaired. However, the generally low food

intake in these fish might reflect long-term deficits in circuit activity that altered the development

or maintenance of neural circuits involved in socially modulated behaviors. In summary, these

results demonstrate that an olfactory pathway including OXT neurons modulates diverse social

behaviors in larval zebrafish.

683
DISCUSSION

Until recently, larval zebrafish had not been thought to exhibit robust social interactions, aside

from simple behaviors such as avoidance of other larvae[12–14]. An emerging body of literature

however has revealed the profound influence of visual, chemosensory and mechanosensory

social cues on larval zebrafish development and behavior[12,15,16,34,44]. Some studies have

reported early kin odor preference and imprinting, suggesting an early experience-dependent

development of positive social behaviors[16,18]. Here, we demonstrate that the presence of

conspecific fish and chemosensory conspecific cues significantly increases appetite and

reduces nociceptive responses in larvae, lending further support for the existence of early pro-

social behaviors in zebrafish.

We further demonstrate that at these larval stages, even brief (~2 hr) social isolation

results in distinct neural signatures that include aversive and nociceptive[6,45] brain circuits in the

preoptic area, posterior tuberculum, caudal hypothalamus and, notably, populations of oxytocin

(OXT)-expressing neurons. These results are consistent with those of a recent study[44] that

described increased activity in the preoptic area as a result of long-term social isolation in older

(juvenile, 21 dpf) zebrafish. Thus, in an intriguing parallel to humans[46], we observe that pain and

social isolation exhibit a shared neural signature in the larval zebrafish brain that extends also to

older animals of the species.

Our analysis of the OXT-expressing population reveals that these neurons are

functionally diverse and modulate distinct behavioral outputs, including feeding and nocifensive

escape responses (see also Wee et al, 2019[6]). Further, we show that OXT neurons receive
social information at least in part via an olfactory bulb to subpallial circuit. A number of studies
have implicated cell populations of the anterior subpallium (aSPa) in distinct visual and olfactory
social responses[17,21,34,35]. Our analysis of the GABAergic y321:Gal4 population provides
convergent evidence that the zebrafish subpallium, thought to be homologous to the
mammalian medial amygdala[17,18,21], is a multi-modal, social cue-responsive, inhibitory circuit
that regulates OXT activity.
We thus propose a circuit mechanism by which the larval zebrafish integrates
conspecific social cues into the modulation of defensive and appetitive behaviors (Figure 6h).
Specifically, our data suggest that OXT neurons and other preoptic populations encode a range
of chemical social information at least partially via olfactory inputs which then modulate
nocifensive and appetitive behavioral outputs via downstream projections to the hindbrain and
other hypothalamic regions. The role of other preoptic populations cannot be overstated -- a
recent study[7] confirmed that numerous preoptic neuropeptidergic populations including OXT are
responsive to aversive stimuli, and that only co-ablation of OXT together with Corticotropin
Releasing Factor (CRF)-expressing neurons was sufficient to significantly attenuate aversive
responses.
OXT neurons displayed a variety of responses to chemical social cues, which could
uniquely signal the presence of adult fish (known predators), genetically-related animals (larval
kin), or distantly-related (larval non-kin) conspecifics. Further, water flow itself elicited a mild
response that may represent an arousing or mildly aversive cue. Overall, kin cues induced the
most widespread inhibition of OXT neurons, while the same OXT neurons can be either excited
or inhibited by other cues, suggesting diverse and valence-specific tuning. The significant
heterogeneity in OXT neuron activity and responses complements a larger body of both
mammalian and fish literature describing heterogeneity in gene expression, projection patterns,
receptor localization and function of OXT neurons[8,47]. We hypothesize that activated OXT
neurons could drive anxiolytic or anti-nociceptive effects, consistent with traditional views of
OXT function[3].
We also note that while differences between kin and non-kin responses are clear in the
olfactory bulb and subpallium, they are more subtle in OXT_{PO} and surrounding preoptic neurons,
and further, we have not demonstrated that such differential modulation results in distinct
behavioral outcomes. However, they are consistent with a larger body of studies which have
reported the capacity for kin recognition and discrimination in zebrafish larvae of comparable
ages[16,18]. Further, our results raise the likelihood that OXT neurons in the PO and PT may

respond differentially to social cues. For example, our pERK results (Figure 1) suggest that
OXT_{PT} neurons may be more strongly suppressed by non-kin and visual rather than kin cues.
Calcium imaging (Figure 2) also revealed differences in OXT_{PO} and OXT_{PT} activities, though the
low numbers of OXT_{PT} neurons recorded makes it difficult to identify conclusive differences.
Finally, y321 optogenetic stimulation was sufficient to suppress OXT_{PO} but not OXT_{PT} neuron
activity (Figure 4), again suggesting that OXT_{PO} and OXT_{PT} may differentially encode or respond
to social stimuli. Intriguingly, a previous study[23] had demonstrated a specific role of OXT_{PT} but
not OXT_{PO} neurons in the development of social preference, again suggesting differences in
their functional roles.

In our study, we did not attempt to identify the chemical components within kin or
conspecific-conditioned water that are sufficient to modulate OXT activity and behavior, though
it is an interesting avenue for follow-up work. Previous studies have suggested the role of bile
acids[48,49] or MHC peptides[20] as odorants that mediate social or kin recognition. Our clustering
analyses have demonstrated complexity in the encoding of kin, non-kin, and adult water stimuli
by olfactory and subpallial neurons which strongly suggests that multiple components, rather
than a single compound, are contributing to kin versus conspecific discrimination.

In humans and other mammals, it is known that social cues, including odors, can
attenuate aversive experience and behaviors, a phenomenon known as "social buffering"[37,38].
Social facilitation of appetite has also been observed in many species[39], and is likely
evolutionarily adaptive. In general, an isolated animal needs to shift its priorities from foraging to
vigilance or escape, since it may be more susceptible to the risk of predation. Accordingly, adult
zebrafish display isolation stress in a group size-dependent manner[50]. We had previously shown
that OXT neurons respond to a range of aversive, particularly noxious stimuli, and are sufficient
to drive motor responses by acting on brainstem targets[6]. We also previously showed that
ablation of OXT neurons or OXT gene deletion had a moderate but significant effect on
nocifensive behavior[6]. By demonstrating that chemical social cues converge on this circuit, and
that kin cues, in particular, predominantly diminish the activity of TRPA1-responsive OXT
neurons, we provide a potential mechanistic understanding of how the OXT circuit can mediate
the phenomenon of "social buffering" in a vertebrate organism[51].

Although we have demonstrated an attenuative effect of kin cues on behavior, including
nocifensive responses, we note that we have not demonstrated the converse -- that isolation
can enhance aversive behavioral output. More detailed analyses of nocifensive behavior in
isolated versus grouped larvae, or of the effect of isolation duration on nocifensive behavior, will
help bolster our hypothesis. We also note that the genetic and pharmacological manipulation

experiments of OXT signaling on food uptake (Figure 6) shown in this paper cannot be

generalized to its social buffering effects, and that additional circuit manipulation experiments

will be necessary to directly tie OXT to social cue-dependent attenuation of nociceptive

responses. Further, given the overlapping sensory and behavioral effects of other

neuropeptides in the preoptic area[7], it is likely that OXT is not the sole mediator of social

buffering behavior, and that manipulation of other preoptic populations (e.g. CRF-expressing

neurons) would be necessary to completely abolish social-dependent influences on behavior.

Notably, the attenuative effect of OXT signaling on larval zebrafish feeding behavior is

consistent with mammalian studies: 1) the insatiable appetite and morbid obesity observed in

Prader Willi Syndrome is likely due to impaired OXT signaling[52,53]; 2) acute inhibition of

paraventricular OXT neurons can promote food intake[54]; 3) lesions of the PVN, as well as

mutations that affected OXT neuron development, have been shown to cause hyperphagia and

obesity[55,56], and; 4) direct administration of OXT has been shown to reduce feeding[4,5]. At the

same time, our data suggests that the role of OXT in feeding may be more complex than

previously appreciated. Notably, while OXT homozygous mutants do not show social state-

dependent modulation of appetite, they also eat less than their heterozygous siblings, implying

that a complete absence of OXT may be detrimental towards feeding.

Such a profound influence of social context on OXT's appetite-suppressing effects may

very well generalize to mammals. For example, a recent study found that inhibiting OXT

signaling enhances sugar intake in a socially dominant mouse regardless of their social context,

whereas in subordinate mice, such inhibition only enhanced appetite when cues from the

dominant mouse were not present[57]. Our results also complement the observations of strong

interactions between social and feeding circuits across evolution[58], and reinforce a role for OXT

in prioritizing different motivated behaviors[4]. However, effects of OXT in larval zebrafish may

occur as part of a coordinated response to both social isolation and noxious contexts, rather

than reproduction or parental care.

There are important distinctions between our findings and the canonical view of OXT

function as suggested by mammalian studies, the most significant of which is that the larval

zebrafish OXT neurons show widespread *activation* by social isolation, rather than by cues

indicating the presence of conspecifics[2,59], and may in fact represent a negative valence state in

larval zebrafish, rather than the rewarding experience it is generally associated with.

We propose three possible reasons for these apparent differences: *first*, representations

within both the mammalian and zebrafish OXT population are diverse and thus the observed

activity patterns in zebrafish may reflect those of specific subpopulations of mammalian OXT

cells (indeed, some zebrafish OXT neurons are activated by conspecific cues, and some
mammalian OXT cells are inhibited by conspecific cues[59]). Interestingly, recent studies in
mammalian models have demonstrated that OXT neurons can also be negatively reinforcing,
and promote fear, stress and anxiety in some situations[1,60]. *Second*, OXT response properties
may have changed over the course of evolution, as more sophisticated social functions were
derived. *Third,* there is a possibility that OXT neuron response properties might reverse over the
course of development, since adult and kin odor generate opposite activity signatures; however,
since the enhancement of preoptic area activation appears to persist at least till juvenile
stages[44], when social preference behaviours have developed, any such reversal would have to
happen closer to adulthood. Intriguingly, a recent preprint[61] has demonstrated that OXT receptor
loss-of-function in fact enhances social development in zebrafish, confirming our findings that
OXT signaling in zebrafish may not necessarily be pro-social. Overall, our results may provide a
broader and more intricate perspective of this highly-conserved peptide's social function and
evolution in vertebrate animals.
In conclusion, our study demonstrates how organizing principles and circuit
implementation strategies underlying social behavior can be elucidated by probing social
context-dependent behaviors in a small and optically-accessible model organism. More broadly,
our dissection of the larval zebrafish OXT circuit provides an entrypoint into understanding how
neuromodulatory systems represent behavioral states such as social isolation, hunger, and
acute nociception, on multiple timescales, and how these representations are then used to
modulate behavioral output in a flexible and context-dependent manner.
827

MATERIALS & METHODS

Fish husbandry and transgenic lines
Larvae and adults were raised in facility water and maintained on a 14:10 hr light:dark cycle at
28°C. All protocols and procedures involving zebrafish were approved by the Harvard
University/Faculty of Arts & Sciences Standing Committee on the Use of Animals in Research
and Teaching (IACUC). Fish were raised at a density of ~40 fish per dish and fed paramecia
from 5 dpf till the day of the experiment. Behavioral experiments were carried out mostly on fish
of the WIK background, although other genotypes (e.g., AB, or mit1fa-/- (nacre) in the AB
background) were also utilized and showed similar behavioral results. mit1fa-/- (nacre) in the AB
background, along with additional transgenes described below, were also used for calcium
imaging and MAP-mapping experiments. Transgenic lines *Tg(oxt:GFP)[62]*,
Tg(HuC:GCaMP6s)[63], *Tg(UAS:GCaMP6s)[64]*, *Tg(UAS:nsfbCherry)[43]* [*nitroreductase is referred*
to as NTR in the text], Tg(UAS:ReaCHR-RFP)[36,42], Tg(Gad1b:loxP-dsRed-loxP-GFP) [referred

to as *Tg(Gad1b:RFP)* in text][65]*, Tg(y321:Gal4)*[34,66]*, Tg(oxt:Gal4)* and oxytocin mutants[6] were
previously published.
843
MAP-mapping
7-8 dpf larvae, that had been continuously fed with an excess of paramecia since 5 dpf, were
either isolated or split into small groups, using 35 mm dishes filled with 3 ml embryo water. For
groups of 10, a larger (10 cm) dish was used to prevent overcrowding. Paramecia was present
within each dish to ensure that the fish were well-fed and had ample stimulation. After 2 hrs,
larvae were quickly funneled through a sieve, which was then quickly dropped into ice-cold 4%
paraformaldehyde, immunostained, imaged and analyzed as described below.

Whole-mount immunostaining
24 hr after fixation (4% paraformaldehyde (PFA) in PBS), fish were washed in PBS + 0.25%
Triton (PBT), incubated in 150 mM Tris-HCl at pH 9 for 15 min at 70°C (antigen retrieval),
washed in PBT, permeabilized in 0.05% Trypsin-EDTA for 45 min on ice, washed in PBT,
blocked in blocking solution (10% Goat Serum, 0.3% Triton in Balanced Salt Solution or 2%
BSA in PBS, 0.3% Triton) for at least an hour and then incubated in primary and secondary
antibodies for up to 3 days at 4°C diluted in blocking solution. In-between primary and
secondary antibodies, fish were washed in PBT and blocked for an hour. If necessary,
pigmented embryos were bleached for 5 min after fixation with a 5%KOH/3%H2O2 solution.

The protocol was similar for dissected brains, except that the brains were dissected in PBS after
24 hr of fixation, and the permeabilization step in Trypsin-EDTA and Tris-HCL antigen retrieval
were omitted. Dissected brains were mounted ventral up on slides in 70% glycerol prior to
imaging. Confocal images of dissected brains were obtained using either a Zeiss LSM 700 or
Olympus FV1000.

Exposure to sensory cues for high-resolution pERK experiments
For generation of conspecific-conditioned water, sibling or non-sibling larvae that had been
continuously fed with an excess of paramecia from 5 dpf were transferred into a new 10 cm petri
dish that did not contain any paramecia, at a concentration of 2 fish per ml. After a 2 hr
incubation, a syringe with an attached 0.45 μm filter was used to very gently suck out the
conditioned water, with great care taken not to disturb or stress the fish in the process.
871
7-8 dpf larvae, that had been continuously fed with an excess of paramecia since 5 dpf, were
either isolated or split into small groups, using 35 mm dishes filled with 3 ml embryo water.
Paramecia was present within each dish to ensure that the fish were well-fed and had ample
stimulation. 700 μl of the filtered conspecific-conditioned water was added to each 35 mm dish
(~1:5 dilution), 30 min before fixation, and embryo water was correspondingly added to controls.
For providing visual access to conspecifics, the 35 mm dishes containing single larvae were
inserted into larger (55 mm) dishes containing ~5 larvae that thus be surrounding but unable to
interact with the single larva. After 2 hrs, larvae were quickly funneled through a sieve, which
was then quickly dropped into 4% paraformaldehyde, dissected in PBS and immunostained as
described in Wee et al, 2019[6].

High resolution pERK analysis
For quantification of pERK intensities of individual OXT neurons, pERK experiments were
performed on dissected *Tg(oxt:GFP)* brains. Cellular-resolution imaging of dissected brains was
obtained using the Zeiss (LSM 700 and LSM 880) or Olympus (FV1000) confocal microscopes.
pERK intensities of individual GFP-positive (OXT) or GFP-negative (non-OXT) neurons were
measured using Fiji[67] and quantified using MATLAB as reported in Wee et al, 2019[6]. In Figure
1c, pERK intensities shown are normalized to the respective mean PO or PT intensities of
control fish (i.e. fish maintained in groups of 3), whereas in Figure 1d, pERK intensities are
normalized to the mean of both PO and PT intensities of control fish. PO and PT were
segmented based on their anterior-posterior positions. For quantification of pERK intensities of
y321 neurons (Supplementary Figure 6), GFP-positive cells were manually segmented.
Raincloud plots[68,69] were produced using open-source software (version 2) in Figure 1 and
Supplementary Figure 1. For sampling of neurons for spatial distribution analysis, a fixed
number of neurons were first randomly sampled without replacement per fish, and then another
round of random sampling was performed to ensure even distribution across treatment groups
(as different treatment groups had different numbers of fish). This ensured a more even
representation of neurons across all fish and treatments.

Calcium imaging and olfactory stimulation
8-11 dpf larval *Tg(oxt:Gal4;UAS:GCaMP6s)* fish in the nacre background were used for calcium
imaging experiments. They were embedded in the center of a 55 cm dish in 1.5% agarose with
their tails and noses freed. Fish were isolated for between 30 min to 4 hours prior to the
experiment, with the bulk between 1-3 hours.
904
Kin or non-kin conditioned water (at a concentration of 1 fish/ml) were generated as described
above. In these calcium imaging experiments, we used WIK or EK fish as non-kin fish, since the
Tg(oxt:Gal4;UAS:GCaMP6s) fish we imaged were of the AB genetic background. For adult
water, 5 adult kin (from the parent tank of larvae) were used to condition 500 ml of water (1
adult fish/100 ml) for 2 hrs, and also subsequently filtered. Kin and non-kin fish used to condition
water were raised apart from experimental fish from 3 dpf, to dissociate genetic from familiarity
effects.
912
Alternating water or chemical stimuli were delivered using a custom-built syringe pump system
controlled by custom Labview software. At specified time intervals, 300 µl of each cue (~30
µl/second) was delivered using a zero-dead-volume multi-channel perfusion pencil (AutoMate
Scientific). Embryo water was also constantly circulated through the dish using a peristaltic
pump (Harvard Apparatus). The interstimulus interval (ISI) was 2 minutes for olfactory bulb
imaging and 5 minutes for OXT neuron imaging. The longer ISI for OXT neuron imaging was
implemented to reduce desensitization and ensure that activity truly returned to baseline before
presenting the subsequent stimulus. In order to reduce experiment time and avert the
possibility of OXT neuron desensitization, we also never compared responses to more than 2
cues (e.g. either kin vs non-kin, or kin vs adult, but not all 3 stimuli). For
Tg(y321:Gal4;UAS:GCaMP6s) imaging experiments, solenoid-controlled (Automate Scientific)
gravity flow[70] (Herrera et al. 2021) was used to deliver the four stimuli (control fish-water, kin,

non-kin, adult water) to each fish. The stimulus delivery cycle was as follows: stimulus was
actively flowed onto the fish for 20 seconds, followed by 45 seconds of no flow, which was
followed by 20 seconds of flow of control water to wash out the chemical, and finally another 45
seconds of no flow.
929
Calcium imaging and behavioral monitoring with TRPA1 stimulation was performed as
previously reported[6] on 8-10 dpf larvae, with a number of core differences: 1) 15 µM instead of
25 µM Optovin (Tocris Biosciences) was used, to reduce background signals (see below) 2)
nostrils are exposed to allow for olfactory stimulation 3) the same UV stimulus intensity used
throughout the experiment, and only 4 stimulations were presented (alternating water and kin
stimulus, equal number of fish with the starting with kin or water stimulus). Fish were isolated for
between 1.5 to 6 hours prior to the experiment, with the bulk between 2-3 hours. Since, in
addition to the tail, the nose was exposed in this current paradigm, Optovin, a colored solution,
was rapidly absorbed into the fish's brain and caused a linear increase in background signal (i.e.
even in non-GCaMP-labeled tissue) over time. Post-hoc subtraction of this background signal
from OXT GCaMP signals restored a flat baseline, allowing us to extract meaningful calcium
signals.
942
943 *Data analysis for calcium imaging*
944 All calcium imaging data was analyzed using custom Fiji and MATLAB software. The general
945 protocol for analysis was:
946
947 1) Image registration to correct for motion artifacts using the TurboReg[71] plugin in Fiji
948
2) Extraction of fluorescence signals from both channels using manually or correlation-based
(see code in Herrera et al. 2021[70]) segmented ROIs in MATLAB or Suite2p[31]. For the analysis of
Tg(y321:Gal4;UAS:GCaMP6s) neurons, manually drawn ROIs were created around the cell
body regions of the anterior subpallium to isolate automatically segmented ROIs that were from
soma. For fish expressing *Tg(HuC:GCaMP6s),* cell segmentation from correlated voxels was
performed and curated using Suite2p software. RFP or mCherry-expressing cells were
classified as having a probability threshold of > 0.4 (Supplementary Figure 4-5)
956
3) Calculation of Δf/f signals from raw traces and alignment to tail traces, as needed, in
MATLAB. Δf/f values were calculated from raw traces using the average fluorescence over the
time period before the first stimulus as the baseline to which all traces were normalized.
Behavior (if any), stimulation and calcium imaging timestamps were aligned and used to extract
stimulus-triggered averages as well as to generate motor and /or stimulus regressors to
correlate with calcium activity. The regressors were convolved with a GCaMP6s kernel based
on its measured response delay (0.48 s) and decay time (3 s, based on Chen et al. (2013)) and
cross-correlated with calcium traces that had been smoothed with a 3-frame zero phase filter.
Bayesian Information Criterion and Aikake information criterion analysis was used to determine
optimal number of clusters for k-means clustering of olfactory bulb neuron correlation
coefficients with different regressors, and the same number of clusters was eventually adopted
across all datasets for simplicity.

In order to determine if an OXT neuron was activated or suppressed, we averaged the calcium
signal over a 60 s interval post-stimulus for each stimulus type. In the case of simultaneous
TRPA1 stimulation, integrated activity was quantified either 5 seconds post-TRPA1 stimulation,
or 30 s prior to TRPA1 stimulation (i.e. post-cue delivery but pre-TRPA1 stimulation). If the
difference between the two integrated calcium signal during olfactory and water stimulation was
more than 0.05, we classified the neurons as being either activated or suppressed by the cue
depending on the sign. Although the threshold of 0.05 is arbitrary, we show that across a range
of thresholds, the relationship between the proportions of suppressed and enhanced neurons
remains consistent for each type of stimulus.

For tethered behavior analysis, bouts were first extracted using the 'findpks' function in
MATLAB. The number of large or small angle bouts within 90 seconds prior to cue delivery (pre-
cue period), 30 seconds from cue delivery (post-cue period) or 5 seconds of TRPA1 stimulus
were subsequently quantified. Other parameters quantified included maximum and minimum
velocities and peak cumulative tail angles (and latencies where applicable) of bouts occurring
prior to or directly after TRPA1 stimulation.

Calcium imaging and Optogenetics (Figure 4)
Optogenetic stimulation and calcium imaging was performed on a commercial 2P microscope
with confocal settings (Zeiss LSM 980 NLO Multiphoton) using a 633 nm laser for ReaChR
activation, and a 488 nm laser for calcium imaging. *Tg(y321:Gal4; oxt:Gal4; UAS:ReaChR-RFP;*
UAS:GCaMP6s) transgenic fish were used to record OXT activity after y321 ReaChR activation.
Fish were embedded in 1.8–2% agarose. ReachR activation was then induced in the
telencephalon periodically for 10 s, and ensuing activity in the preoptic area / posterior
tuberculum was recorded continuously during the 60-120 s interval between stimuli.

Free-swimming TRPA1 stimulation (Supplementary Figure 7)
Fish were singly placed into a 20.6 mm cut-out agarose circular mold illuminated by three quad
blue LEDs (Luxeon Star, 470 nm). To probe the effect of conspecific cues, DMSO or Optovin
solutions were generated either in embryo water or kin-conditioned water. Following 5 min of
habituation in DMSO (effectively also the isolation period), fish were stimulated once every 30 s
with a 100 ms pulse of blue light. The DMSO solution was then exchanged with Optovin, and
the same protocol repeated. Behavior was recorded at 200 fps (Pike F-032, Allied Vision) and
analyzed using custom Python software.

1003

Social feeding experiments
For experiments in which feeding was assessed, larvae that had been continuously fed with an
excess of paramecia from 5 dpf were either isolated or placed in groups of 3, in 35 mm dishes
(3 ml embryo water), in the absence of food. After 2 hours, fluorescent-labeled paramecia was
added followed by a quick fixation after 30 min (full protocol is described in Wee et al, *eLife*,
2019). Sensory cues were generated and presented as described above for pERK experiments.
Fixed larvae were subsequently distributed into 96-well flat-bottom dishes and imaged using the

AxioZoom V16 (Zeiss) and analyzed using Fiji software (3D Objects Counter, custom software
also available on www.github.com/carolinewee)

1013

1014   Pharmacological experiments
1015   For pharmacology experiments, the OXT antagonist L-368,899 hydrochloride (Tocris
1016   Biosciences, 2641, 5 µM) and agonist WAY 267464 dihydrochloride (Tocris Biosciences, 3933,
1017   5 µM) were added to each dish from the beginning of the isolation period (2hrs prior to adding
1018   fluorescently-labeled paramecia).

1019

1020   *OXT neuron ablation / inhibition experiments*
1021   Transgenic larvae expressing *Tg(oxt:Gal4;UAS:nsfbCherry)* or their non-transgenic siblings
1022   were incubated in 2.5 mM Metronidazole (MTZ, Sigma-Aldrich, M3761) from 5-6/6-7dpf. MTZ
1023   was subsequently washed out, and food intake was measured at 7 or 8 dpf. For these
1024   experiments, the MTZ-treated siblings were used as the control group.

1025

Oxytocin mutants
For experiments with *oxt* mutants[6], a heterozygous *oxt* mutant was crossed to a null mutant. All
fish from the same clutch were raised together in the same dish and fed from 5dpf. On the day
of the experiment (7 or 8 dpf), the larvae were distributed into isolated or non-isolated conditions
and fed labeled paramecia after 2 hrs. They were fixed after 30 minutes, transferred into 96 well
plates to keep track of experimental conditions and their gut fluorescence was imaged. After
imaging their genomic DNA was extracted and genotyped and their genotypes were matched to
their experimental conditions (isolated or non-isolated).

1034

Statistics
All error bars show mean ± SEM over fish. Significance was reported as follows: *$p<0.05$,
**$p<0.01$, ***$p<0.001$. Significance was determined using the Wilcoxon Signed-Rank test for
paired data and the Wilcoxon Rank-Sum test for independent samples. One-sided tests were
used in cases where there was a clear hypothesis for the direction of effect. One-way ANOVA
with Bonferroni Correction was used in cases where there were multiple comparisons. Wilcoxon
Signed-Rank test was used in Figure 1a for comparing the distribution of normalized OXT ROI
signals across different behavioral stimuli to the null hypothesis of median 1.

1043

1044   **DATA AVAILABILITY**
1045   All data, code (hardware control and analysis) and resources (transgenic lines/mutants
1046   generated) will be made available by the corresponding author upon request.

1047

1048   **CODE AVAILABILITY**
1049   Live versions of the analysis code are maintained at www.github.com/carolinewee.

1050

1051   **ACKNOWLEDGEMENTS**
1052   We are grateful to Adam Douglass (Univ. of Utah) for continued support, advice, and comments.
1053   We thank the CBS imaging facility, Harvard Center for Biological Imaging, and the NorthWest
1054   Undergraduate Teaching Laboratories at Harvard for the successful completion of many

experiments. J. Miller, K. Hurley and B. Hughes provided invaluable fish care. This work was supported by U01-NS090449 (FE), R24-NS086601 (FE), U19-NS104653 (FE and SK) and Simons Foundation grant SCGB 325207 (FE). CLW was supported by the National Science Scholarship from the Agency for Science, Technology and Research (A*STAR), Singapore.

AUTHOR CONTRIBUTIONS

C.L.W., E.S., and S.K. originally conceived of the project, which was then developed into its final form with M.N and F.E. S.K. & F.E. supervised the project. C.L.W., E.S., and M.N. designed and performed most of the experiments, and analyzed most of the data. K.J.H. and E.S. performed y321 calcium imaging and optogenetic imaging experiments, K.J.H and C.L.W. analyzed data from these experiments. M.N. also developed hardware and software for calcium imaging and behavioral experiments, and analyzed the free-swimming behavioral data. S.W. performed feeding experiments and analyzed data. C.L.W., S.K., and F.E. wrote the manuscript with contribution from all other authors.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES:

1. Rash, J. A., Aguirre-Camacho, A. & Campbell, T. S. Oxytocin and pain: a systematic review and synthesis of findings. *Clin. J. Pain* **30**, 453–462 (2014).

2. Lieberwirth, C. & Wang, Z. Social bonding: regulation by neuropeptides. *Front. Neurosci.* **8**, 171 (2014).

3. Jezova, D., Skultetyova, I., Tokarev, D. I., Bakos, P. & Vigas, M. Vasopressin and Oxytocin in Stress. *Ann. N. Y. Acad. Sci.* **771**, 192–203 (1995).

4. Leng, G. *et al.* Oxytocin and appetite. *Prog. Brain Res.* **170**, 137–151 (2008).

5. Sabatier, N., Leng, G. & Menzies, J. Oxytocin, feeding, and satiety. *Front. Endocrinol.* **4**, 35 (2013).

6. Wee, C. L. *et al.* Zebrafish oxytocin neurons drive nocifensive behavior via brainstem premotor targets. *Nat. Neurosci.* **22**, 1477–1492 (2019).

7. Lovett-Barron, M. *et al.* Multiple overlapping hypothalamus-brainstem circuits drive rapid threat avoidance. *bioRxiv* 745075 (2019) doi:10.1101/745075.

8. Herget, U., Gutierrez-Triana, J. A., Salazar Thula, O., Knerr, B. & Ryu, S. Single-Cell Reconstruction of Oxytocinergic Neurons Reveals Separate Hypophysiotropic and Encephalotropic Subtypes in Larval Zebrafish. *eneuro* **4**, ENEURO.0278–16.2016 (2017).

9. Stoop, R. Neuromodulation by oxytocin and vasopressin. *Neuron* **76**, 142–159 (2012).

10. Knobloch, H. S. & Grinevich, V. Evolution of oxytocin pathways in the brain of vertebrates. *Front. Behav. Neurosci.* **8**, 31 (2014).

11. Theofanopoulou, C., Gedman, G., Cahill, J. A., Boeckx, C. & Jarvis, E. D. Universal nomenclature for oxytocin-vasotocin ligand and receptor families. *Nature* **592**, 747–755 (2021).

12. Groneberg, A. H. *et al.* Early-Life Social Experience Shapes Social Avoidance Reactions in Larval Zebrafish. *Curr. Biol.* **30**, 4009–4021.e4 (2020).

13.   Dreosti, E., Lopes, G., Kampff, A. R. & Wilson, S. W. Development of social behavior in

young zebrafish. *Front. Neural Circuits* **9**, 39 (2015).

14.   Harpaz, R., Nguyen, M. N., Bahl, A. & Engert, F. Precise visuomotor transformations

underlying collective behavior in larval zebrafish. *bioRxiv* 2021.05.24.445521 (2021)

doi:10.1101/2021.05.24.445521.

15.   Anneser, L. *et al.* The neuropeptide Pth2 dynamically senses others via mechanosensation.

Nature **588**, 653–657 (2020).

16.   Gerlach, G., Hodgins-Davis, A., Avolio, C. & Schunter, C. Kin recognition in zebrafish: a 24-

hour window for olfactory imprinting. *Proc. Biol. Sci.* **275**, 2165–2170 (2008).

17.   Gerlach, G. & Wullimann, M. F. Neural pathways of olfactory kin imprinting and kin

recognition in zebrafish. *Cell Tissue Res.* **383**, 273–287 (2021).

18.   Gerlach, G. *et al.* Behavioural and neuronal basis of olfactory imprinting and kin recognition

in larval fish. *J. Exp. Biol.* **222**, (2019).

19.   Biechl, D., Tietje, K., Gerlach, G. & Wullimann, M. F. Crypt cells are involved in kin

recognition in larval zebrafish. *Sci. Rep.* **6**, 24590 (2016).

20.   Hinz, C. *et al.* Olfactory imprinting is triggered by MHC peptide ligands. *Sci. Rep.* **3**, 2800

(2013).

21.   Biechl, D. *et al.* Identification of accessory olfactory system and medial amygdala in the

zebrafish. *Sci. Rep.* **7**, 44295 (2017).

22.   Randlett, O. *et al.* Whole-brain activity mapping onto a zebrafish brain atlas. *Nat. Methods*

12, 1039–1046 (2015).

23.   Wircer, E. *et al.* Homeodomain protein Otp affects developmental neuropeptide switching in

oxytocin neurons associated with a long-term effect on social behavior. *Elife* **6**, (2017).

24.   Herget, U., Wolf, A., Wullimann, M. F. & Ryu, S. Molecular neuroanatomy and

chemoarchitecture of the neurosecretory preoptic-hypothalamic area in zebrafish larvae. *J.*

Comp. Neurol. **522**, 1542–1564 (2014).

1144 25. Rauch, G.-J., Granato, M. & Haffter, P. A polymorphic zebrafish line for genetic mapping

1145   using SSLPs on high-percentage agarose gels. *Tech. Tips Online* **2**, 148–150 (1997).

1146 26. Manica, A. Filial cannibalism in teleost fish. *Biol. Rev. Camb. Philos. Soc.* **77**, 261–277

1147   (2002).

1148 27. Mehlis, M., Bakker, T. C. M., Engqvist, L. & Frommen, J. G. To eat or not to eat: egg-based

1149   assessment of paternity triggers fine-tuned decisions about filial cannibalism. *Proc. Biol.*

1150   *Sci.* **277**, 2627–2635 (2010).

1151 28. Saito, D., Komatsuda, M. & Urano, A. Functional organization of preoptic vasotocin and

1152   isotocin neurons in the brain of rainbow trout: central and neurohypophysial projections of

1153   single neurons. *Neuroscience* **124**, 973–984 (2004).

1154 29. Swanson, L. W. & Sawchenko, P. E. Paraventricular nucleus: a site for the integration of

1155   neuroendocrine and autonomic mechanisms. *Neuroendocrinology* **31**, 410–417 (1980).

1156 30. Sawchenko, P. E. & Swanson, L. W. Immunohistochemical identification of neurons in the

1157   paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal cord

1158   in the rat. *J. Comp. Neurol.* **205**, 260–272 (1982).

1159 31. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy.

1160   *bioRxiv* 061507 (2017) doi:10.1101/061507.

1161 32. Miyasaka, N. *et al.* Olfactory projectome in the zebrafish forebrain revealed by genetic

1162   single-neuron labelling. *Nat. Commun.* **5**, 3639 (2014).

1163 33. Miyasaka, N. *et al.* From the olfactory bulb to higher brain centers: genetic visualization of

1164   secondary olfactory pathways in zebrafish. *J. Neurosci.* **29**, 4756–4767 (2009).

1165 34. Stednitz, S. J. *et al.* Forebrain Control of Behaviorally Driven Social Orienting in Zebrafish.

1166   *Curr. Biol.* **28**, 2445–2451.e3 (2018).

1167 35. Bruckner, J. J. *et al.* The microbiota promotes social behavior by neuro-immune modulation

1168   of neurite complexity. *bioRxiv* 2020.05.01.071373 (2020) doi:10.1101/2020.05.01.071373.

1169 36. Lin, J. Y., Knutsen, P. M., Muller, A., Kleinfeld, D. & Tsien, R. Y. ReaChR: a red-shifted

variant of channelrhodopsin enables deep transcranial optogenetic excitation. *Nat.*

Neurosci. **16**, 1499–1508 (2013).

37. Kikusui, T., Winslow, J. T. & Mori, Y. Social buffering: relief from stress and anxiety. *Philos.*

Trans. R. Soc. Lond. B Biol. Sci. **361**, 2215–2228 (2006).

38. Kiyokawa, Y. Social Odors: Alarm Pheromones and Social Buffering. *Curr. Top. Behav.*

Neurosci. (2015) doi:10.1007/7854_2015_406.

39. Higgs, S. & Thomas, J. Social influences on eating. *Current Opinion in Behavioral Sciences*

9, 1–6 (2016).

40. Shimada, Y., Hirano, M., Nishimura, Y. & Tanaka, T. A high-throughput fluorescence-based

assay system for appetite-regulating gene and drug screening. *PLoS One* **7**, e52549

(2012).

41. Jordi, J. *et al.* A high-throughput assay for quantifying appetite and digestive dynamics. *Am.*

J. Physiol. Regul. Integr. Comp. Physiol. ajpregu.00225.2015 (2015).

42. Wee, C. L. *et al.* A bidirectional network for appetite control in larval zebrafish. *Elife* **8**,

(2019).

43. Davison, J. M. *et al.* Transactivation from Gal4-VP16 transgenic insertions for tissue-

specific cell labeling and ablation in zebrafish. *Dev. Biol.* **304**, 811–824 (2007).

44. Tunbak, H., Vazquez-Prada, M., Ryan, T. M., Kampff, A. R. & Dreosti, E. Whole-brain

mapping of socially isolated zebrafish reveals that lonely fish are not loners. *Elife* **9**, (2020).

45. Lovett-Barron, M. *et al.* Multiple convergent hypothalamus–brainstem circuits drive

defensive behavior. *Nat. Neurosci.* (2020) doi:10.1038/s41593-020-0655-1.

46. Eisenberger, N. I., Lieberman, M. D. & Williams, K. D. Does Rejection Hurt? An fMRI Study

of Social Exclusion. *Science* **302**, 290–292 (2003).

47. Althammer, F. & Grinevich, V. Diversity of oxytocin neurons: beyond magno- and

parvocellular cell types? *J. Neuroendocrinol.* e12549 (2017).

48. Cong, X. *et al.* Zebrafish olfactory receptors ORAs differentially detect bile acids and bile

salts. *J. Biol. Chem.* **294**, 6762–6771 (2019).

49. Kermen, F. *et al.* Stimulus-specific behavioral responses of zebrafish to a large range of

odors exhibit individual variability. *BMC Biol.* **18**, 66 (2020).

50. White, L. J., Thomson, J. S., Pounder, K. C., Coleman, R. C. & Sneddon, L. U. The impact

of social context on behaviour and the recovery from welfare challenges in zebrafish, Danio

rerio. *Anim. Behav.* **132**, 189–199 (2017).

51. Smith, A. S. & Wang, Z. Hypothalamic oxytocin mediates social buffering of the stress

response. *Biol. Psychiatry* **76**, 281–288 (2014).

52. Swaab, D. F. Neuropeptides in hypothalamic neuronal disorders. *Int. Rev. Cytol.* **240**, 305–

375 (2004).

53. Francis, S. M. *et al.* Oxytocin and vasopressin systems in genetic syndromes and

neurodevelopmental disorders. *Brain Res.* **1580**, 199–218 (2014).

54. Atasoy, D., Betley, J. N., Su, H. H. & Sternson, S. M. Deconstruction of a neural circuit for

hunger. *Nature* **488**, 172–177 (2012).

55. Xi, D., Gandhi, N., Lai, M. & Kublaoui, B. M. Ablation of Sim1 neurons causes obesity

through hyperphagia and reduced energy expenditure. *PLoS One* **7**, e36453 (2012).

56. Tolson, K. P. *et al.* Inducible neuronal inactivation of Sim1 in adult mice causes

hyperphagic obesity. *Endocrinology* **155**, 2436–2444 (2014).

57. Olszewski, P. K., Allen, K. & Levine, A. S. Effect of oxytocin receptor blockade on appetite

for sugar is modified by social context. *Appetite* **86**, 81–87 (2015).

58. Fischer, E. K. & O'Connell, L. A. Modification of feeding circuits in the evolution of social

behavior. *J. Exp. Biol.* **220**, 92–102 (2017).

59. Resendez, S. L. *et al.* Social stimuli induce activation of oxytocin neurons within the

paraventricular nucleus of the hypothalamus to promote social behavior in male mice. *J.*

Neurosci. (2020) doi:10.1523/JNEUROSCI.1515-18.2020.

60. Grillon, C. *et al.* Oxytocin increases anxiety to unpredictable threat. *Mol. Psychiatry* **18**,

958–960 (2013).

61. Gemmer, A. *et al.* Oxytocin receptors influence the development and maintenance of social

behavior in zebrafish (Danio rerio). *bioRxiv* 2021.06.26.449566 (2021)

doi:10.1101/2021.06.26.449566.

62. Coffey, C. M. *et al.* Novel oxytocin gene expression in the hindbrain is induced by alcohol

exposure: transgenic zebrafish enable visualization of sensitive neurons. *PLoS One* **8**,

e53991 (2013).

63. Kim, D. H. *et al.* Pan-neuronal calcium imaging with cellular resolution in freely swimming

zebrafish. *Nat. Methods* **14**, 1107–1114 (2017).

64. Muto, A. *et al.* Activation of the hypothalamic feeding centre upon visual prey detection.

Nat. Commun. **8**, 15029 (2017).

65. Satou, C. *et al.* Transgenic tools to characterize neuronal properties of discrete populations

of zebrafish neurons. *Development* **140**, 3927–3931 (2013).

66. Marquart, G. D. *et al.* A 3D Searchable Database of Transgenic Zebrafish Gal4 and Cre

Lines for Functional Neuroanatomy Studies. *Front. Neural Circuits* **9**, 78 (2015).

67. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

68. Whitaker, K. *et al. RainCloudPlots/RainCloudPlots: WellcomeOpenResearch*. (2019).

doi:10.5281/zenodo.3368186.

69. Allen, M. *et al.* Raincloud plots: a multi-platform tool for robust data visualization. *Wellcome*

Open Res. **4**, 63 (2021).

70. Herrera, K. J., Panier, T., Guggiana-Nilo, D. & Engert, F. Larval Zebrafish Use Olfactory

Detection of Sodium and Chloride to Avoid Salt Water. *Curr. Biol.* **31**, 782–793.e3 (2021).

71. Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration

based on intensity. *IEEE Trans. Image Process.* **7**, 27–41 (1998).