Swine Growth Promotion with Antibiotics or Alternatives

can Increase Antibiotic Resistance Gene Mobility Potential

Johanna Muurinen,[a]* Jacob Richert,[b] Carmen Wickware,[a] Brian Richert, [a] and Timothy A.

Johnson[a]*

[a]Department of Animal Sciences, Purdue University, West Lafayette, Indiana, USA

[b]Department of Animal Sciences & Industry, Kansas State University, Manhattan, Kansas, USA

Running Title: Growth Promoters Mobilize Antimicrobial Resistance

*Address correspondence to Johanna Muurinen and Timothy Johnson,

johanna.muurinen@onehealth.fi and john2185@purdue.edu, respectively.

Abstract

Even though the use of antibiotics for food-producing animals may contribute to the emergence of antimicrobial resistance, antibiotics are still used as growth promoters. Due to consumer and regulatory pressures, the use of alternatives to antibiotics as growth promoters is increasing, thus more information is needed on their capability to disseminate antimicrobial resistance compared to antibiotics. We investigated the impacts of carbadox (antibiotic), copper sulfate and zinc oxide (metals) and mushroom powder (natural product) on the pig fecal resistome and microbiome. Antibiotic resistance gene (ARG) and mobile genetic element (MGE) abundances were measured using a high-throughput qPCR array with 382 primer pairs. Bacterial community composition was determined by 16S rRNA gene sequencing. More ARGs co-occurred with MGEs in the growth promoter group samples than in the control group samples. Community composition could not be linked to resistome in the growth promoter group samples, indicating a potential decoupling of ARGs and phylogeny. Additionally, machine-learning methods aided in defining the community and resistome differences in response to treatments. Since increased ARG mobility potential was the primary response to the dietary additives used in this study, we suggest that ARG mobility should be considered when designing antimicrobial use policies and antimicrobial resistance surveillances.

Introduction

Antibiotics have been used in pork production to prevent diseases and to increase productivity since the 1940's[1]. Due to the established connection of antibiotic use in animals and the emergence of antibiotic resistance in pathogenic bacteria[2], the regulation and prohibition of antibiotic growth promoters is increasing (prohibition in the EU in 2006[1], restrictions in 2017 in the U.S.[3], and restrictions in 2020 in China[4]). Additionally, a report to the Secretary General of the United Nations suggested that antibiotic growth promoters should be completely phased out

from livestock production[5]. Despite the increasing regulation, most of the world's pork is

produced in countries that still allow growth promotion with antibiotics. In 2020, the U.S. was

the third largest producer of pork worldwide after China and the EU, and the second largest

exporter of pork meat after the EU. Currently, the livestock industries in the U.S. as well as in

many other countries are looking for alternative feed additives to replace antibiotics as growth

promoters due to increasing restrictions and consumer concerns towards antibiotic use in animal

agriculture. Because of these reasons, the markets are now favorable to so-called alternatives to

antibiotics as growth promoters and the interest towards them is increasing rapidly[6,7].

Among the antibiotics that are still allowed for growth promotion in the U.S., carbadox is

used in pigs mainly to control dysentery and bacterial enteritis. Carbadox can cause short-term

but also long-lasting alterations in the microbiome[8] and can promote the mobility of antibiotic

resistance genes (ARGs) through transduction[9]. Still, little is known about the impact of

carbadox or especially alternative feed additives on the abundance of ARGs in pig feces.

Numerous growth promoters that are alternatives to antibiotics are currently under study for their

efficacy to promote animal growth and gastrointestinal health. Among natural products, of

special interest are medicinal mushrooms. *Cordyceps militaris* and *Ophiocordyceps sinensis*

(formerly *C. sinensis*) produce many bioactive compounds, such as Cordycepin and Beta-

Glucan, which have antimicrobial effects and can enhance the immune system[10]. Zinc (Zn) and

copper (Cu) are important trace elements for all organisms, and livestock animals commonly

receive feed supplements that ensure their required concentration in the feed. Higher

concentrations of Zn and Cu can be used for controlling various bacterial infections in livestock

animals, such as diarrheal diseases[7] and also for growth promotion[7,11,12]. However, use of Zn

oxide has shown to select zoonotic methicillin resistant *Staphylococcus aureus* (MRSA) and

multiresistant *E. coli*[13–16]. Consumption of Cu-supplemented animal feed can increase the

prevalence of erythromycin resistance in gram-positive bacteria[17] and various plasmid-mediated

resistance genes in microbiomes[16,18]. Thus, it seems that some growth promoters that are

alternatives to antibiotics are actually antimicrobials and may select ARGs similarly as the

antibiotics they are meant to replace.

On the other hand, when quantifying the antibiotic resistome, it is important to consider

that changes in resistance gene abundance may simply be due to a change in the underlying

community composition, since community composition can explain resistance gene

composition[19,20]. It is thought that growth promoters alter the gut microbial community

composition, and that the microbiota composition could be manipulated to promote animal

growth by inducing populations favorable to growth, immunity, and gut health[21]. Therefore, it is

important to characterize the bacterial community composition to determine if the populations

linked to induced animal growth carry ARGs and mobile genetic elements (MGEs). However,

making connections between taxonomic data and resistance gene data can be challenging, since

microbiome datasets are compositional and rarely meet assumptions of normality that many

statistical tests require[22,23]. In addition, it has been pointed out that different normalization

strategies could influence the results derived from sequence data[24,25].

In order to examine the influence of carbadox, cordyceps mushroom powder and

pharmacological concentrations of Zn and Cu on the community composition and resistome[26],

we took fecal samples from pigs that were administered these growth promoters, extracted DNA,

analyzed 16S rRNA gene sequences and quantified genes related to resistance and horizontal

gene transfer with 382 primer pairs using a high-throughput qPCR array[27,28]. We also compared

two normalization strategies for taxonomic data using statistical analyses that are suitable for

compositional datasets. Our results suggest that inclusion of ARGs into MGEs led to decoupling

of bacterial community composition and resistome composition in response to growth

promotion, highlighting the importance of MGEs in shaping the resistomes under the influence

of different types of antimicrobial agents.

Results

Samples and data quality control

Samples were collected from pigs that had been assigned into non-treatment control group (NTC = no antibiotic or alternative growth promoters), or one of the growth promoter groups (AB = carbadox, M = mushroom powder mixture of *C. militaris* and *O. sinesis*, ZnCu = Zn oxide and Cu sulfate). Each treatment group consisted of six pens and each pen had seven pigs. The DNA for 16S rRNA gene sequencing was extracted from combined fecal samples of one medium weight female and male per pen. After quality filtering, a total of 741,785 sequences were obtained. For 22 samples sequences per sample ranged from 11,392 to 67,072. Two samples were discarded due to low number of sequences, resulting in five samples in the NTC and AB groups. The data analysis for 16S rRNA gene sequencing was completed using two different methods: rarefication and subsampling or total sum scaling (TSS). The TSS normalized 16S rRNA gene sequence data had 132 operational taxonomical units (OTUs), while rarefied and subsampled 16S rRNA gene sequence data 127 OTUs. The same DNA samples were used for qPCR array analysis to examine how the treatments altered the resistome. One hundred and thirty-six assays out of 382 assays (Supplementary Table S1) targeting antibiotic resistance genes (ARGs) or mobile genetic elements (MGEs) were positive. See materials and methods for qPCR data processing and Ct value adjustment for four assays that had unspecific amplification.

The profiles of most abundant genera and genes were similar in different treatment groups

To take into account the influence of normalization of 16S rRNA gene sequence reads on the results, we used two different normalization methods. Only small differences were discovered between TSS normalized and rarefied and subsampled OTUs among the most abundant genera (Fig. 1A and B). The TSS normalized OTUs and rarefied and subsampled

OTUs correlated significantly (ρ = 0.97, $p < 0.05$) (Supplementary Fig. S1), indicating the

agreement of the overall community composition of the two normalization methods. *Prevotella*

was the most abundant genera in all treatments and all samples had several short chain fatty acid

producers (Fig. 1A and B). Very few OTUs were found only in one treatment group with both

normalization methods and a majority of the OTUs were found in all treatment groups, however

more OTUs were found in all treatment groups using TSS normalization than rarefied and

subsampled normalization (Fig. 1D and E).

The different treatment groups also had similar resistome profiles (Fig. 1C). The NTC

group had the highest number of positive assays (110), the AB group had 106 positive assays,

The M group had 100 positive assays and the ZnCu group had 103 positive assays. Twenty-eight

genes were detected in only one treatment group, but most genes were found in all treatment

groups (Fig. 1F). Out of the positive assays, 108 targeted ARGs and 28 MGEs. Among the

detected ARGs, 63 conferred resistance to aminoglycosides, MLSBs or tetracyclines and the

most common resistance mechanism was antibiotic deactivation (Fig. Supplementary S2A and

B). Among MGEs, most of the positive assays targeted insertion sequences (12) or transposases

(7) (Fig. Supplementary S2C).

Growth promoters favored different genera, ARGs and MGEs

Generalized linear models (GLMs) with False discovery rate control[29] for *p*-value

adjustment were used for testing which genera, ARGs and MGEs differed in abundance between

treatments. Mostly only minor differences in abundance of different genera were found between

treatment groups (Supplementary Fig. S3A and B). Compared to all other growth promoters,

carbadox favored unclassified *Veillonellaceae* in feces and decreased the abundance of

Streptococcus, whereas Zn and Cu decreased the abundance of *Bifidobacterium* and

Campylobacter (adjusted *p*-value < 0.05, gamma distribution GLMs [TSS OTUs] and

negative binomial GLMs [rarefied and subsampled OTUs]) (Fig. 2A and B, Supplementary

Table S2 and S3). Mushroom powder decreased the abundance of fecal *Roseburia* and favored

Campylobacter compared to other treatments (adjusted *p*-value < 0.05, gamma distribution

GLMs [TSS OTUs] and negative binomial GLMs [rarefied and subsampled OTUs]) (Fig. 2A and

B, Supplementary Table S2 and Supplementary S3). *Streptococcus* was more abundant in M and

ZnCu group samples compared to other groups (adjusted *p*-value < 0.05, gamma distribution

GLMs [TSS OTUs] and negative binomial GLMs [rarefied and subsampled OTUs]) (Fig. 2A and

B, Supplementary Table S2 and S3). More ARGs and MGEs were differentially abundant in

NTC group and M group comparison than when NTC group was compared to ZnCu or AB

groups (Supplementary Fig. S3C). Compared to all other growth promoters, carbadox and Zn

and Cu increased the relative abundance of *vat(E),* mushroom powder increased the relative

abundance of *tetW*, and Zn and Cu favored *tetM* (adjusted *p*-value < 0.05, gamma distribution

GLMs) (Fig. 2C, Supplementary Table S4). Interestingly, carbadox treatment decreased the

relative abundance of *tetW, tet(32)*, *erm(B)*, *ermT* and *tetM* compared to other treatments or to

NTC group (adjusted *p*-value < 0.05, gamma distribution GLMs) (Fig. 2C, Supplementary

Table S4).

Both OTU normalization methods yielded similar results for differential OTU abundance

in the treatment groups (Fig. 2A and B). The exceptions were that *Phascolarctobacterium* was

among the differentially abundant genera when TSS normalization was used, whereas

Faecalibacterium was differentially abundant when rarefying and subsampling was used (Fig.

2A and B) and that more genera were differentially abundant when TSS normalization was used

(Supplementary Table S2 and S3).

Growth promoters had modest influences on overall bacterial community composition

and resistome

With both TSS normalized OTUs and rarefied and subsampled OTUs, the treatment

group explained 24% of the variability in community composition (PERMANOVA, $R^2 = 0.239$,

$p < 0.05$). With ARGs and MGEs, the treatment group explained 27% of the variability

(PERMANOVA, $R^2 = 0.267$, $p < 0.05$). The number of sequences per sample was also included

as a variable in the PERMANOVA model with both normalization methods, but the library size

did not have influence on the community variability in our samples ($p > 0.05$).

167   The different growth promoters altered the community composition and resistome only

slightly, since in all non-metric multidimensional scaling (NMDS) ordinations the samples

belonging to different treatment groups clustered close to each other with mostly overlapping

centroid confidence intervals (Fig. 3A, B and D). In ordinations of TSS normalized and rarefied

and subsampled OTU-tables, the ZnCu group tended to cluster further away from the other

samples indicating more dissimilar community composition (Fig. 3A and B), however there were

no significant differences between treatment groups among pairwise PERMANOVA

comparisons (False discovery rate control adjusted[29] p-value > 0.05). In the ordination of the

ARG and MGE data, there was no separation between ZnCu group and the other groups, instead

the M group clustered slightly separately (Fig. 2D) and in the pairwise comparisons using the

ARG and MGE data, M vs. AB group explained 32% of the variance (PERMANOVA, $R^2 =$

0.321, False discovery rate control adjusted[29] p-value < 0.05), but in all the other pairwise

comparisons the differences were nonsignificant (False discovery rate control adjusted[29] p-value

> 0.05).

Taxonomic variation did not explain resistome variation in growth promoter group

samples

183   The OTU ordinations were correlated against each other and against the ARG and MGE

NMDS ordination using Procrustes analysis to determine if the taxonomic variation explains

resistome variation. The two OTU NMDS ordinations had reasonably high (0.7) and significant

($p < 0.05$) correlation and the Procrustes residual error remained lower than 0.25 in all samples

except one (Fig. 3C). The correlation between the rarefied and subsampled OTU ordination

(which did not include all the data due to subsampling) and the ARG and MGE ordination was

moderately high (0.6, $p < 0.05$) and there was no pattern in Procrustes residual errors across

different samples (Fig. 3F). Contrariwise, the correlation between the TSS normalized OTU

ordination (that included all quality filtered data) and the ARG and MGE ordination was

nonsignificant ($p > 0.05$) and Procrustes residual errors were high in most samples not belonging

to NTC group, in which the residual errors were mostly equal or less than the first quantile

residual value (Fig. 3E). The Procrustes analyses performed on TSS normalized OTUs and

resistome implies that taxonomic variation explained resistance variation in NTC samples with

less error. However, growth promoters altered the resistome, resulting higher residual errors in

samples belonging to AB, M and ZnCu groups and thus the overall correlation between TSS

normalized OTU ordination and ARG and MGE ordination was not significant (Procrustes

analysis, $p > 0.05$).

The links between bacterial community structure and resistome were further examined

with Mantel's test using Spearman's rank correlation. The correlation coefficients between the

TSS normalized OTUs and ARGs and MGEs distance matrices as well as the rarefied and

subsampled OTUs and ARGs and MGEs distance matrices were low ($\rho = 0.25$ and $\rho = 0.23$,

respectively, $p < 0.05$), which suggests that the phylogenetic composition did not govern the

resistome composition when all samples were included in the analysis. We also used Mantel's

test for all treatment groups individually. With the NTC group and both OTU normalization

methods, the bacterial community distance matrix correlated significantly with the distance

matrix obtained from ARGs and MGEs ($\rho = 0.66$ [TSS OTUs] and $\rho = 0.62$ [rarefied and

subsampled OTUs], $p < 0.05$). With all growth promoter groups and both OTU normalization

methods, the correlations between OTU and ARG and MGE distance matrixes were

nonsignificant, giving more evidence that the growth promoters shaped the resistance

composition and thus taxonomic variation did not explain the resistance variation in the growth

promoter group samples. It should be denoted that we had only five or six samples in each

treatment group; however, the results were consistent, since the correlations between taxonomic

structure and resistome were reasonably high and significant in NTC group and nonsignificant in

all other groups.

Growth promoters increased the co-occurrences of ARGs and MGEs

A correlation matrix between ARG and MGE relative abundances was visualized using

network analysis to examine if the growth promoters selected resistance genes into mobile

genetic elements. The network of NTC group was simpler compared to growth promoter group

networks as the number of correlating ARGs and MGEs increased in response to growth

promotion (Fig. 4). There were no integrons co-occurring with resistance genes in the NTC

group network, but integrons were present in all the growth promoter group networks (Fig. 4)

and co-occurred with many ARGs. AB and M group networks had more aminoglycoside

resistance genes than other networks and M group also had the most multidrug resistance genes,

but the least tetracycline resistance genes, whereas AB group network had the most vancomycin

resistance genes (Fig. 4).

Transposase gene for insertion sequence-like element IS1216 was the best predictor for

resistance and community composition

We used machine-learning approaches to identify drivers of changes in community

composition. First, a cluster analysis was performed on a combined data table of TSS normalized

OTUs and ARGs and MGEs using *t*-distributed stochastic neighbor embedding (t-SNE)

algorithm[30] and HDBSCAN algorithm[31]. Then, a classification random forests model[32,33] was

used for identifying the predictors for the clustering pattern. Three clusters were identified after

dimension reduction. Most of the samples belonged to cluster 2, while cluster 1 had two ZnCu

samples, one M group sample and one AB group sample, and cluster 3 contained only three

ZnCu samples (Fig. 5A). According to the partial dependence plot that shows the most important

predictors found by the classification random forests model, clusters 1 and 3 were separated from

cluster 2 due to higher abundance of transposase gene linked to IS1216 element and from each

other because the abundance of *tetO* was lower in cluster 3 (Fig. 5B). Five of the nine best

predictors for clustering pattern were MGEs and ARGs and four were bacterial genera (Fig. 5B).

Interestingly, only one of the ZnCu group samples belonged to cluster 2, while all the other

ZnCu samples clustered to clusters 1 and 3. Thus, growth promotion with Zn and Cu might cause

more alterations in the community composition and resistome than the other growth promoters

examined in this study.

Discussion

Our study explored the influences of antibiotic and alternative growth promoters on the

pig fecal resistome, ARG mobility, and bacterial community composition. Resistome structure

and taxonomic structure correlated in the NTC group samples but there was no correlation

between resistome and phylogeny in the growth promotion groups. Some have reported that

community composition predicts resistance profile[19,20,34], while others have reported that under

the presence of a selective pressure, phylogeny and resistome can become uncoupled in the

swine feces and manure[35]. Our network analysis revealed that the growth promoters in this study

increased linkages between ARGs and MGEs, i.e. more ARGs correlated with MGEs in the

resistome of growth promoter group pigs than in the resistome of NTC group pigs. Thus, it

seems plausible that resistance gene mobility would have increased in response to growth

promotion, since subinhibitory antimicrobial concentrations and bacterial stress response

mechanisms can induce horizontal gene transfer[36,37].

Besides the inclusion of ARGs into MGEs, another possible scenario is that bacteria

carrying MGEs with multiple ARGs and other embedded features would have tolerated the

growth promoters better than bacteria carrying MGEs with only few ARGs, possibly due to more

variability in stress response mechanisms[16,36,37]. Despite the mechanism, more ARGs co-

occurring with MGEs in response to growth promotion could indicate more persistent resistance

gene collection[38] and potential for ARGs to be mobilized by the MGEs in the future, and this

way the ARGs in the animal microbiomes could eventually be transmitted to human

microbiomes[2,36].

Despite the changes in ARG mobility potential, the abundances of nearly all ARGs and

MGEs were on similar levels in the non-treatment control group and the growth promoter

groups, which indicates that it is unlikely that changing the substance that is used for growth

promotion would reduce antimicrobial resistance in the pig fecal microbiome, at least in a short

period of time (33 days in this study). Gut bacteria, especially gram-negative species, are known

to carry many ARGs and MGEs and the gastro-intestinal tract is suspected to be a major hotspot

for horizontal gene transfer[36,39]. These observations have also been made with individuals

without antibiotic exposure[40,41].

Interestingly, some ARGs were less abundant in the AB group than in the NTC group and

alternative growth promoter groups. A potential explanation for this could be that carbadox is a

broad acting antibiotic and suppresses most bacterial populations[8]. Overall, we did not observe

large shifts among genera as a result of growth promotion feed additives. *Bifidobacteria*, which

have been previously linked to lower antibiotic resistance level[42,43], were somewhat more

abundant in the NTC and M group samples; however, in this study the ARG abundances in

samples belonging to different groups were similar. It is possible that the shifts in the community

composition caused by growth promoters were small with high variability and thus the changes

in the community structure are difficult to capture with community-wide molecular approaches.

Although the changes in community composition and resistome in response to treatments

were modest and mostly nonsignificant, we observed shifts with machine learning methods. The

differences we observed did not precisely follow the experimental grouping (but largely isolated

the ZnCu group), and therefore they were not found with commonly used ordination methods or

differential abundance analysis. The transposase gene linked to IS1216 element was the driver

for the clustering result and has been previously associated to Gram-positive bacteria[35,44], mainly

Enterococcus faecium[45]. However, *Enterococcus* was not among the most abundant or

differentially abundant genera, or among the predictors of the clustering result in random forest

analysis. Even though we cannot demonstrate all the factors explaining the clustering result, the

observation that zinc and copper treatment perhaps altered the microbiome and resistome more

than the other treatments would not have been possible without the use of a machine learning

approach for statistical analyses. Thus, machine learning is an important tool in determining the

complex relationship between resistome and bacterial community compositions.

Considering the two approaches to normalize OTU counts between samples, the rarefied

and subsampled OTU data and TSS normalized OTU data mostly agreed; however, in Procrustes

analysis, the rarefied and subsampled OTU data correlated significantly with the resistome data

when all samples were used, although the same analysis using the TSS normalized data as input

showed that the taxonomic data did not explain resistance. It is important to acknowledge that

rarefying and subsampling captures the most abundant OTUs for each sample[25] and discards rare

OTUs, as well as slightly adjusting OTU relative abundances in different samples. If the sample

size and differences in OTU abundances are small before the procedure, the shifts might change

the outcome if rarefied OTU data is used in comparison with data obtained using a different

method. Thus, researchers making connections between taxonomic data and resistome

observations should use multiple methods to confirm their findings.

In this experiment, we observed a decoupling of phylogeny and resistance gene

composition, possibly through increased co-occurrences between ARGs and MGEs. This

indicates that the studied feed-additives can increase and maintain resistance gene mobility

potential. On the other hand, under this experimental design, the withdrawal of antibiotics did

not decrease the abundance of antibiotic resistance, however, we did not observe enrichment of

ARGs in response to growth promotion either. Thus, we suggest that the mobility potential of

ARGs should be considered to include in antimicrobial resistance surveillances to establish the

impact of changes in agricultural practices on antimicrobial resistance.

Materials and methods

Animal experiment statement

318 All procedures involving animal use were approved by the Purdue University Animal

319 Care and Use Committee (protocol #1303000841), and animal care and use standards were based

320 upon the Guide for the Care and Use of Agricultural Animals in Research and Teaching[46].

Samples and DNA

The pig fecal samples were obtained from growth promoter experiment where 210

weanling pigs ((Duroc × (York × Landrace)) avg. 19 d of age and 5.8 kg were used in a 33-day

trial. The experiment had 7 pigs in each pen and 6 pens per each treatment. Feed amendment

treatments were: 1) non-treatment control (NTC); 2) antibiotic growth promoter (carbadox, 55

ppm) (AB); 3) mushroom powder (mixture of *C. militaris* and *O. sinesis*, 300 ppm) (M); 4)

carbadox and mushroom powder mixture (results are not included in this study); 5) copper

sulfate (125 ppm) and zinc oxide (3000 ppm d 0-7, 2000 ppm d 7-35) (ZnCu). After 33 days,

fecal samples were taken from 1 median weight female and male per pen. Samples from the

same pen were pooled, and DNA was extracted using the DNeasyPowerLyzer PowerSoil DNA

Isolation Kit (Qiagen) according to the manufacturer's protocol. Extracted DNA was stored at

−20 °C before 16S sequencing and qPCR array.

16S sequencing and quantitative PCR array

The 16S rRNA gene library was constructed as described[47]. Briefly, the V4 region of the

bacterial 16S rRNA gene was amplified with the 515R (GTGCCAGCMGCCGCGGTAA) /

806R (GGACTACHVGGGTWTCTAAT) primers. 16S rRNA gene libraries were also prepared

for a known mock community (20 Strain Even Mix Genomic Material; ATCC® MSA–1002TM)

and a no-template control (water). The amplified DNA from one 96-well plate was normalized

using a SequalPrep Normalization Plate (Invitrogen), and pooled into a single library. Library

concentrations were determined using the KAPA Library Quantification Kit (Roche) and the

average fragment length was determined using a high sensitivity kit with the Bioanalyzer

(Agilent). The pooled samples, mock community, and water were sequenced with Illumina

MiSeq v2 (500 cycles). Sequences were demultiplexed using oligonucleotide bar code sequences

and Illumina software.

Quantitative PCR reactions and raw data processing were conducted using WaferGen

SmartChip Real-time PCR system as reported[48]. The qPCR array reactions were performed using

384 previously described and validated primer sets (assays) (Supplementary Table S1)[28]. One

sample from M group (2_M) was not included in the qPCR array analysis due to technical error.

Samples from pigs that received both mushroom powder and carbadox were also excluded and

therefore the results from 16S rRNA amplicon sequencing are not presented in this study.

16S sequence analysis and qPCR array data processing

The 16S rRNA amplicon sequences were analyzed using mothur (v 1.39.3)[47]: contigs

were made from paired forward and reverse raw reads, aligned to reference sequences (SILVA

database release 132)[49], screened and filtered to remove low quality reads (ambiguous bases

allowed = 0, maximum read length = 275, homopolymers allowed = 8), classified with reference

to known taxonomic classifications (RDP training set 16)[50] and clustered into OTUs. The

sequences clustered into 137 different OTUs at the 3% dissimilarity level. One NTC group

sample (11_NTC) and one AB group sample (16_AB) were discarded due to low number of

obtained sequences. 16S sequences were normalized using two different methods: total sum

scaling (TSS) using R and rarefying and subsampling using mothur. To produce the rarefied and

subsampled OTU table, the data were subsampled to 7,500 reads per sample according to

rarefication curves (Supplementary Fig. S4). The rarefied and subsampled OTU table and only

quality filtered OTU table (for TSS) were imported into R. After removing the results of samples

from animals that received both mushroom powder and carbadox, 132 different OTUs remained

in the TSS normalized OTU table and 127 different OTUs in the rarefied and subsampled OTU

table. The TSS normalization was carried out in R by dividing each OTU read count by the total

number of reads in that sample and all the NA observations (zero sequences) were replaced with

1.490935e-07, which was 100-fold lower than the lowest observed relative abundance.

In the qPCR array, assays "16S old 1_1","blaOXY-1_1118", "cmlV_911", "czcA_1536",

"fabK_1520", "intI1F165_clinical_359" and "tetPA_1507" were positive in the negative control,

however the Ct-values in the negative control were mostly higher than in experimental samples

(Supplementary Table S5). Assay "tetPA_1507" was detected only in the negative control and

the Ct-values of assay "czcA_1536" were removed from the results since they were lower in the

negative control than in samples. Assay "16S old 1_1" results were removed and not used in

normalization since DNA amplification was more efficient in assay "16S new 2_2". The Ct-

values of the remaining four assays that were positive in the negative control were adjusted as

follows: The Ct values of each of these assays in each sample were subtracted from Ct value of

the assay in the negative control. The resulting numbers were then subtracted from 27, which

was the Ct value was used as the cutoff between true positive values and primer-dimer

amplification. Next, all the Ct values that were higher than 27 were set to "NA". After this, all

the assays that were undetected in all the samples were removed, resulting in 136 assays out of

382 targeting to AGRs or MGEs being included in the data table. The ΔCt values, ΔΔCt values

and relative gene abundances were calculated from these Ct values as previously described[27].

Genes under the detection limit were given a ΔCt value of 20, which was higher than any

observed ΔCt (17.4).

 Statistical analyses

R version 3.5.1 (2018-07-02) was used for data exploration, visualization and for all

statistical analyses. Analysis of differential abundances of taxa and ARGs and MGEs were

carried out using gamma distribution GLMs with TSS normalized OTUs and ARGs and MGEs

(relative abundances) and negative binomial GLMs with rarefied and subsampled OTUs

(abundance). Gamma distribution model was selected because the relative abundance values

(TSS normalized OTUs and ARGs and MGEs) did not fit to normal distribution but followed the

gamma distribution. Negative binomial models were selected for rarefied and subsampled OTUs

because of the overdispersion in the abundance data. With both model types, p-values were

obtained with Tukey's post-hoc test and adjusted with False discovery rate control[29] using *glht*

function in the multcomp package[51].

The community and resistome compositions were analyzed using vegan package[52].

PERMANOVA was used to examine the shifts in community composition and resistome

between treatment groups with function *adonis* and 9,999 permutations. Nonmetric

multidimensional scalings (NMDS) were completed using the Bray–Curtis dissimilarity index

with function *metaMDS*. Procrustes analysis with the *protest* function was used to examine the

agreement of ordinations of TSS normalized OTUs and rarefied and subsampled OTUs as well

as ordination of ARGs and MGEs and both OTU ordinations separately. Mantel's test and

Spearman's rank correlation was used to analyze the links between microbial community

structure and resistome: first, Bray-Curtis dissimilarity indexes were calculated for TSS

normalized OTUs, rarefied and subsampled OTUs and for ARGs and MGEs with function

vegdist. Then the *mantel* function was applied for the dissimilarity indexes of TSS normalized

OTUs and ARGs and MGEs as well as rarefied and subsampled OTUs and ARGs and MGEs.

Mantel's tests between both OTU dissimilarity matrices and dissimilarity matrix of ARGs and

MGEs were also run for all the treatment groups separately.

To examine if the treatments selected resistance genes into mobile genetic elements, a

correlation matrix between ARG and MGE relative abundances was visualized using network

analysis with Gephi[53]. Spearman's rank correlations between ARGs and MGEs within treatment

groups and their *p*-values used in network analysis were obtained with package psych[54] using

False discovery rate control[29]. Only ARG-MGE-pairs that were detected at least in three samples

with a strong positive correlation ($\rho > 0.8$, adjusted *p*-value < 0.05) were included.

The factors influencing the shifts in taxonomic structure and resistome were analyzed

with machine learning algorithms as previously described[55]. First, dimension reduction was

executed on a combined data table of TSS normalized OTUs and ARGs and MGEs using t-SNE

algorithm (29) and the R package Rtsne[56], with 50,000 iterations and "perplexity" set to 5. Then,

clusters in the two-dimensional data were identified using HDBSCAN algorithm[31] in the package

dbscan[57]. The minimum number of members in clusters ("minPts") was set to 3. Classification

random forest model[32] was used with partial dependence plot function in edarf package[58] for

identifying the most important predictors for the clustering pattern. The forests were grown to

10,000 trees using the ranger package[59] and the best predictors were screened using Gini index

by adding predictors one at a time in the order of decreasing importance[60]. The final model was

then selected according to the highest Cohen's Kappa (comparison of observed accuracy and

expected accuracy).

Availability of materials and data

Raw reads from 16S rRNA gene amplicon sequencing are deposited under BioProject

accession number PRJNA605462 at NCBI. The R code, mothur commands and all datasets used

in statistical analyses are available at https://github.com/sjmuurine/ZnCu.

References

1. Dibner, J. J. & Richards, J. D. Antibiotic growth promoters in agriculture: history and mode of action. *Poultry Science* **84**, 634–643 (2005).

2. Marshall, B. M. & Levy, S. B. Food animals and antimicrobials: Impacts on human health. *Clinical Microbiology Reviews* **24**, 718–733 (2011).

3. Centner, T. J. Recent government regulations in the United States seek to ensure the effectiveness of antibiotics by limiting their agricultural use. *Environ Int* **94**, 1–7 (2016).

4. Hu, Y. J. & Cowling, B. J. Reducing antibiotic use in livestock, China. *Bulletin of the World Health Organization* 98, 360-361 (2020).

5. Interagency Coordination Group on Antimicrobial Resistance (IACG). No time to wait: securing the future from drug-resistant infections. (2019).

6. Thacker, P. A. Alternatives to antibiotics as growth promoters for use in swine production: a review. *J Anim Sci Biotechnol* **4**, 35–35 (2013).

7. Willing, B. P. *et al.* Bacterial resistance to antibiotic alternatives: A wolf in sheep's clothing? *Animal Frontiers* **8**, 39–47 (2018).

8. Looft, T., Allen, H. K., Casey, T. A., Alt, D. P. & Stanton, T. B. Carbadox has both temporary and lasting effects on the swine gut microbiota. *Frontiers in Microbiology* **5**, 1–1 (2014).

9. Johnson, T. A. *et al.* The in-feed antibiotic carbadox induces phage gene transcription in the swine gut microbiome. *mBio* **8**, 1–14 (2017).

10. Richert, J. *et al.* Effects of Cordyceps Mushroom Powder on Nursery Pig Performance. *Kansas Agricultural Experiment Station Research Reports* **4**, (2018).

11. Di Giancamillo, A. *et al.* Copper sulphate forms in piglet diets: Microbiota, intestinal morphology and enteric nervous system glial cells. *Animal Science Journal* **89**, 616–624 (2018).

457   12.   Jacela JY, DeRouchey JM, Tokach MD, et al. Feed additives for swine: Fact sheets –

458   high dietary levels of copper and zinc for young pigs, and phytase.

459   13.   Slifierz, M. J., Friendship, R. & Weese, J. S. Zinc Oxide Therapy Increases Prevalence

460   and Persistence of Methicillin-Resistant Staphylococcus aureus in Pigs: A Randomized

461   Controlled Trial. *Zoonoses and Public Health* **62**, 301–308 (2015).

462   14.   Bednorz, C. *et al.* The broader context of antibiotic resistance: zinc feed supplementation

463   of piglets increases the proportion of multi-resistant Escherichia coli in vivo. *International*

464   *journal of medical microbiology□: IJMM* **303**, 396–403 (2013).

465   15.   Ciesinski, L. *et al.* High dietary zinc feeding promotes persistence of multi-resistant E.

466   coli in the swine gut. *PLoS ONE* **13**, 1–18 (2018).

467   16.   Poole, K. At the Nexus of Antibiotics and Metals: The Impact of Cu and Zn on Antibiotic

468   Activity and Resistance. *Trends in Microbiology* **25**, 820–832 (2017).

469   17.   Van Noten, N., Gorissen, L. & De Smet, S. Assistance in the update of the systematic

470   literature review (SLR): 'Influence of copper on antibiotic resistance of gut microbiota on pigs

471   (including piglets)'. *EFSA Supporting publication* **2016:EN–10**, (2016).

472   18.   Fang, L. *et al.* Co-spread of metal and antibiotic resistance within ST3-IncHI2 plasmids

473   from E. coli isolates of food-producing animals. *Scientific Reports* **6**, 25312 (2016).

474   19.   Munk, P. *et al.* Abundance and diversity of the faecal resistome in slaughter pigs and

475   broilers in nine European countries. *Nature Microbiology* **3**, (2018).

476   20.   Forsberg, K. J. *et al.* Bacterial phylogeny structures soil resistomes across habitats.

477   *Nature* **509**, 612–6 (2014).

478   21.   Ward, T. L. *et al.* Antibiotics and host-tailored probiotics similarly modulate effects on

479   the developing avian microbiome, mycobiome, and host gene expression. *mBio* **10**, 1–17

480   (2019).

22.   Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. & Egozcue, J. J. Microbiome

Datasets Are Compositional: And This Is Not Optional. *Frontiers in Microbiology* **8**, 2224

(2017).

23.   Zuur, A. F., Ieno, E. N. & Elphick, C. S. A protocol for data exploration to avoid

common statistical problems. *Methods in Ecology and Evolution* **1**, 3–14 (2010).

24.   McMurdie, P. J. & Holmes, S. Waste Not, Want Not: Why Rarefying Microbiome Data

Is Inadmissible. *PLoS Computational Biology* **10**, (2014).

25.   Weiss, S. *et al.* Normalization and microbial differential abundance strategies depend

upon data characteristics. *Microbiome* **5**, 27–27 (2017).

26.   Wright, G. D. The antibiotic resistome: the nexus of chemical and genetic diversity.

Nature Reviews Microbiology **5**, 175–186 (2007).

27.   Muurinen, J., Karkman, A. & Virta, M. High Throughput Method for Analyzing

Antibiotic Resistance Genes in Wastewater Treatment Plants. in *Antimicrobial Resistance in*

Wastewater Treatment Processes 253–262 (John Wiley & Sons, Inc., 2017).

doi:10.1002/9781119192428.ch14.

28.   Stedtfeld, R. D. *et al.* Primer set 2.0 for highly parallel qPCR array targeting antibiotic

resistance genes and mobile genetic elements. *FEMS Microbiology Ecology* **94**, (2018).

29.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *Journal of the Royal statistical society: series B*

(Methodological) **57**, 289–300 (1995).

30.   Van Der Maaten, L. Accelerating t-SNE using tree-based algorithms. *The Journal of*

Machine Learning Research **15**, 3221–3245 (2014).

31.   Campello, R. J. G. B., Moulavi, D. & Sander, J. Density-Based Clustering Based on

Hierarchical Density Estimates. in *Advances in Knowledge Discovery and Data Mining* (eds.

Pei, J., Tseng, V. S., Cao, L., Motoda, H. & Xu, G.) 160–172 (Springer Berlin Heidelberg, 2013).

32.   Breiman, L. Random Forests. *Machine Learning* **45**, 5–32 (2001).

33.   Liaw, A. & Wiener, M. Classification and regression by randomForest. *R news* **2**, 18–22 (2002).

34.   Su, J. Q. *et al.* Antibiotic Resistome and Its Association with Bacterial Communities during Sewage Sludge Composting. *Environmental Science and Technology* **49**, 7356–7363 (2015).

35.   Johnson, T. A. *et al.* Clusters of Antibiotic Resistance Genes Enriched Together Stay. **7**, 1–11 (2016).

36.   Davies, J. & Davies, D. Origins and Evolution of Antibiotic Resistance. *Microbiology and Molecular Biology Reviews* **74**, 417–433 (2010).

37.   Poole, K. Bacterial stress responses as determinants of antimicrobial resistance. *Journal of Antimicrobial Chemotherapy* **67**, 2069–2089 (2012).

38.   Enne, V. I. *et al.* Assessment of the fitness impacts on Escherichia coli of acquisition of antibiotic resistance genes encoded by different types of genetic element. *Journal of Antimicrobial Chemotherapy* **56**, 544–551 (2005).

39.   Netherwood, T. *et al.* Gene transfer in the gastrointestinal tract. *Applied and environmental microbiology* **65**, 5139–41 (1999).

40.   Moore, A. M. *et al.* Gut resistome development in healthy twin pairs in the first year of life. *Microbiome* **3**, 27 (2015).

41.   Gibson, M. K. *et al.* Developmental dynamics of the preterm infant gut microbiota and antibiotic resistome. *Nature Microbiology* **1**, 16024 (2016).

42.   Pärnänen, K. *et al.* Maternal gut and breast milk microbiota affect infant gut antibiotic resistome and mobile genetic elements. *Nature Communications* **9**, 3891 (2018).

43.   Taft, D. H. *et al.* Bifidobacterial Dominance of the Gut in Early Life and Acquisition of

Antimicrobial Resistance. *mSphere* **3**, 1–24 (2018).

44.   Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile Genetic Elements

Associated with Antimicrobial Resistance. *Clin. Microbiol. Rev.* **31**, e00088-17 (2018).

45.   Clewell, D. B. *et al.* Extrachromosomal and mobile elements in enterococci:

transmission, maintenance, and epidemiology. in *Enterococci: From commensals to leading*

causes of drug resistant infection [Internet] (Massachusetts Eye and Ear Infirmary, 2014).

46.   Chapter 11: Swine in *Guide for the Care and Use of Agricultural Animals in Research*

and Teaching. 143–155 (Federation of Animal Science Societies 2010).

47.   Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D.

Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing

Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Appl. Environ.*

Microbiol. **79**, 5112 (2013).

48.   Wang, F. *et al.* High throughput profiling of antibiotic resistance genes in urban park

soils with reclaimed water irrigation. *Environmental science & technology* (2014)

doi:10.1021/es502615e.

49.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res* **41**, D590–D596 (2013).

50.   Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput rRNA

analysis. *Nucleic Acids Research* **42**, D633–D642 (2013).

51.   Hothorn T. et al. Package 'multcomp.' Simultaneous inference in general parametric

models. https://CRAN.R-project.org/package=multcomp (2016).

52.   Oksanen, J. et al. vegan: Community Ecology Package. https://CRAN.R-

project.org/package=vegan (2019).

53.   Bastian M., Heymann S. & Jacomy M. Gephi: an open source software for exploring and

manipulating networks. *International AAAI Conference on Weblogs and Social Media.*
http://www.aaai.org/ocs/index.php/ICWSM/09/paper/view/154 (2009).

54.   Revelle, W. psych: Procedures for Personality and Psychological

 Research. https://CRAN.R-project.org/package=psych (2019).

55.   Ruuskanen M. O., St Pierre K. A., St Louis V. L., Aris-Brosou S. & Poulain A. J.

Physicochemical drivers of microbial community structure in sediments of lake hazen, nunavut,

canada. *Front Microbiol* **9**:1138–1138. doi:10.3389/fmicb.2018.01138 (2018).

56.   Krijthe J. & van der Maaten L. Rtsne: T-distributed stochastic neighbor embedding using

a barnes-hut implementation. https://cran.r-project.org/web/packages/Rtsne/index.html (2018).

57.   Hahsler M., Piekenbrock M., Arya S. & Mount D. dbscan: Fast Density-Based Clustering

with R. *Journal of Statistical Software* **91**:1-30.. https://CRAN.R-project.org/package=dbscan

(2019).

58.   Jones Z. & Linder F. edarf: Exploratory Data Analysis using Random Forests. *Journal of*

Open Source Software **1**:92. https://CRAN.R-project.org/package=edarf (2017).

59.   Wright M. N. & Ziegler A. ranger: A fast implementation of random forests for high

dimensional data in C++ and R. *Journal of Statistical Software*, **77**:1-17. https://CRAN.R-

project.org/package=ranger (2017).

60.   Menze B. H., et al. A comparison of random forest and its Gini importance with standard

chemometric methods for the feature selection and classification of spectral data. *BMC*

Bioinformatics **10**:213–213. doi:10.1186/1471-2105-10-213 (2009).

Acknowledgements

We thank Ms. Olivia Consoli for excellent technical assistance in laboratory analysis and

Purdue University for internal financial support. Information Technology at Purdue is

acknowledged for providing computational resources.

Contributions

J. M. and C. W. analyzed the data (bioinformatics and statistical analyses) and J. M. wrote the manuscript. J. R. was responsible for the practical implementation of the pig feeding trial and fecal sampling. B. R., J. R. and T. A. J. designed the animal experiment and sampling. O.C. performed laboratory work under the supervision of T. A. J. and C. W. All authors contributed to manuscript revisions and have read and approved the final version of the manuscript.

Corresponding authors

Correspondence to Johanna Muurinen and Timothy A. Johnson, johanna.muurinen@onehealth.fi and john2185@purdue.edu, respectively.

Completing interests statement

The authors declare no competing interests.

Figures and Figure legends

Fig. 1. Comparison of most abundant genera, ARGs and MGEs in different treatment groups.

Samples on x-axis are grouped according to the treatments and color-coded. Sample names are

as follows: NTC= Non-treatment control, AB= Carbadox (antibiotic), M= Mushroom powder

and ZnCu= zinc oxide and copper sulfate. The number in front of the group code denotes the

number of the pen. (A) Stacked bar plot showing 16 most abundant genera in OTUs normalized

using total sum scaling (TSS). (B) Stacked bar plot showing 16 most abundant genera in rarefied

and subsampled OTUs. (C) Most abundant genes related to antibiotic resistance and mobile

genetic elements (n=27). Each row represents results of each primer set (assay) (Supplementary

Table S1) displayed on the y-axis. Assays are grouped according to the antibiotic group which

the   target   genes   confer   resistance.   MLSB   is   abbreviation   for

Macrolide−Lincosamide−Streptogramin B resistance and MGE for mobile genetic elements. One

sample from mushroom powder group was left out from the qPCR array analysis due to a

technical error. (D) Venn diagram showing the OTUs that are shared between samples belonging

into different treatment groups when TSS normalization was used. (E) Venn diagram showing

the OTUs that are shared between samples belonging into different treatment groups when

Rarefying and subsampling was used. (F) Venn diagram showing the ARGs and MGEs that are

shared between samples belonging into different treatment groups.

609
Fig. 2. Boxplots showing the most abundant genera, ARGs and MGEs with statistically

significant differences between treatment groups labeled on right. NTC= Non-treatment control,

AB= Carbadox (antibiotic), M= Mushroom powder and ZnCu= zinc oxide and copper sulfate.

The asterisks "*", "**" and "***" denote statistical significance levels at $p < 0.05$, $p < 0.01$ and

$p < 0.001$, respectively. (A) Most abundant genera in TSS normalized OTUs (n=14), (B) Most

abundant genera in rarefied and subsampled OTUs (n=14), (C) Most abundant genes related to

resistance and transfer (n=10). See Supplementary Fig. S3A, B and C for all differentially

abundant genera and ARGs and MGEs and Supplementary Table S2, S3 and S4 for fold changes

of the differently abundant genera and ARGs and MGEs.

Fig. 3. NMDS ordinations of TSS normalized OTUs, rarefied and subsampled OTUs and ARGs

& MGEs and Procrustes errors between them. Sample names are as follows: NTC= Non-

treatment control, AB= Carbadox (antibiotic), M= Mushroom powder and ZnCu= zinc oxide and

copper sulfate. The number in the sample name denotes the number of the pen. (A) NMDS

ordination of TSS normalized OTUs. (B) NMDS ordination of rarefied and subsampled (RS)

OTUs. (C) Procrustes errors between NMDS ordinations of TSS normalized OTUs and rarefied

and subsampled OTUs. (D) NMDS ordination of ARGs and MGEs. (E) Procrustes errors

between NMDS ordinations of TSS normalized OTUs and ARGs & MGEs. (F) Procrustes errors

between NMDS ordinations of rarefied and subsampled OTUs and ARGs & MGEs. The

Procrustes residual error line plots (C, E and F) allow residual error size comparisons. The bars

show the difference in the community structures between the two normalization methods (C) as

well as the differences in community structure and resistome structure in samples belonging to

different treatment groups (E and F). Horizontal lines denote the median (solid), 25% and 75%

quantiles (dashed).

Fig. 4. Network analysis showing co-occurrence patterns between ARGs and MGEs within the samples in different treatment groups. Nodes of the MGEs are triangles, and circle resistance gene nodes are colored according to the antibiotic they confer resistance. Edges between resistance gene nodes and mobile genetic element nodes have the color of the resistance gene

node. Nodes have equal sizes, edges have equal weights, and distance between the nodes is
irrelevant.

Fig. 5. t-SNE analysis of a dataset containing ARGs, MGEs and TSS normalized OTUs (all

relative abundances) and the distribution of the pens where experimental pigs were kept. (A) The

clustering pattern of the samples. (B) Partial dependence plot of cluster numbers and the most

important predictors in the order of decreasing importance. The partial dependence plot shows

the effect of each predictor on the model outcome one by one, while the other predictors are

fixed to their average value.

Supplementary information legends

Supplementary Table S1. List of the used primer sets.

Supplementary Fig. S1. Correlation between rarefied and subsampled OTUs and TSS normalized

OTUs.

Supplementary Fig. S2 Composition of positive assays grouped by (A) antibiotic group the

targeted gene confers resistance, (B) resistance mechanism and (C) mobile genetic element

group.

Supplementary Fig. S3. Differentially abundant genera and genes. Samples on x-axis are

grouped according to the treatments. Sample names are as follows: NTC= Non-treatment control,

AB= Carbadox (antibiotic), M= Mushroom powder and ZnCu= zinc oxide and copper sulfate.

The number in front of the group code denotes the number of the pen. Each row represents

abundance of each genus or the results of each primer set (assay) (Supplementary Table S1)

displayed on the y-axis. Only genera and genes with statistically significant differences between

treatment groups are shown (A) Differentially abundant genera with TSS normalization. See

Supplementary Table S2 for fold changes. (B) Differentially abundant genera with Rarefying and

subsampling. See Supplementary Table S3 for fold changes. (C) Differentially abundant ARGs

and MGEs. See Supplementary Table S4 for fold changes.

Supplementary Table S2. Pairwise comparisons of gamma distribution GLMs of relative

abundances of each genera between treatment groups. TSS normalized OTU table was used as

the input.

Supplementary Table S3. Pairwise comparisons of negative binomial GLMs of abundances of

each genera between treatment groups. Rarefied and subsampled OTU table was used as the

input.

Supplementary Table S4. Pairwise comparisons of gamma distribution GLMs of relative

abundances of each ARG or MGE between treatment groups.

Supplementary Fig. S4. Rarefaction curves. OTU collection curves determined from sequence

analysis. Each line represents one sample. Vertical line shows the subsampling cutoff: 7500

sequences

Supplementary Table S5. Assays that had unspecific amplification. Ct values in the negative

control and mean Ct-values in samples.

688