A New World begomovirus infecting Grapevine

Humberto Debat[1,2]*[Ω], Diego Zavallo[3][Ω], Sabrina Moyano[4], Facundo Luna[4], Sebastian Asurmendi[3], Sebastian Gomez-Talquenca[4]*

[1] Instituto de Patología Vegetal, Centro de Investigaciones Agropecuarias, Instituto Nacional de Tecnología Agropecuaria (IPAVE-CIAP-INTA), 11 de setiembre 4755, Córdoba, Argentina, X5020ICA.

[2] Unidad de Fitopatología y Modelización Agrícola, Consejo Nacional de Investigaciones Científicas y Técnicas (UFYMA-CONICET), 11 de setiembre 4755, Córdoba, Argentina, X5020ICA.

[3] Instituto de Agrobiotecnología y Biología Molecular (IABIMO), Instituto Nacional de Tecnología Agropecuaria (INTA), Consejo Nacional de investigaciones Científicas y Tecnológicas (CONICET), Los Reseros y Nicolas Repeto, Hurlingham, Buenos Aires, Argentina, 1686.

[4] Estación Experimental Agropecuaria Mendoza, Instituto Nacional de Tecnología Agropecuaria (EEA-Mendoza-INTA), San Martín 3853, Luján de Cuyo, Mendoza, Argentina, 5534.

[Ω] These authors contributed equally.

* Correspondence:

Humberto Debat, debat.humberto@inta.gob.ar

Sebastian Gomez-Talquenca, gomez.talquenca@inta.gob.ar

ORCID ID:

HD 0000-0003-3056-3739

DZ 0000-0002-9021-2175

SA 0000-0001-9516-5948

SGT 0000-0001-9265-8333

Keywords: Geminiviridae, Begomovirus, Grapevine, New World Begomovirus, virus discovery, Vitis vinifera, Torrontes

Abstract

Grapevine (*Vitis* L.), a deciduous woody vine, is a highly valuable agricultural crop. Grapevine, as other crops, is vulnerable to infectious pathogens. Several of them, including viruses, are a major threat to viticulture. Geminiviruses (family *Geminiviridae*) are insect transmitted, small non-enveloped viruses, with circular single-stranded DNA genomes, which are encapsidated in quasi-icosahedral geminated virions. There are only four geminiviruses associated to grapevine: two members of genus *Grablovirus* and two unassigned species. Here we present evidence of a novel begomovirus (genus *Begomovirus*) infecting grapevines from Argentina. Two circular ssDNA virus sequences were assembled from high-throughput sequencing data from *Vitis vinifera* cv Torrontes from Mendoza province, Argentina. Structural and functional annotation indicated that the virus sequences corresponded to complete DNA components A and B of a novel New World bipartite begomovirus. Genetic distance and evolutionary analyses support that the detected sequences correspond to a new virus, the first begomovirus reported to infect grapevine, a tentative prototype member of a novel species which we propose the name "grapevine begomovirus A" (GBVA).

Annotated Sequence Record

Grapevine (*Vitis* L.), a deciduous woody vine, is a highly valuable agricultural crop. Viticulture oriented to wine industry, consumption of fresh fruits, or raisins represents a multi-billion dollar industry. Grapevine, as other crops, is vulnerable to infectious pathogens. Several of them, including viruses, are a major threat to viticulture. Given its economical importance, grapevine has been the focus of intensive virology studies worldwide. As of today, grapevine has been reported to be linked to more than 80 viruses and thus represent the crop naturally hosting the largest number of viruses ever studied [1]. Even though Argentina is one of the top ten global wine and grapevine producer, the study of pathogens in general, and viruses in particular associated to this crop remain scarce. Only 13 viruses have been identified in grapevine in Argentina [2, 3], the latter represents the recent detection of Grapevine red blotch-associated virus (GRBV, *Grablovirus*) in *V. vinifera* cv. Flame Seedless from Mendoza, one of the very few geminiviruses linked to grapevine.

Geminiviruses (family *Geminiviridae*) are insect transmitted, small non-enveloped viruses, with circular single-stranded DNA genomes (2.5-5.2 kbps), which are encapsidated in quasi-icosahedral geminated virions of 22-38 nm size [4]. Geminiviruses are significant crop pathogens and cause important damages to economically relevant vegetables in many tropical and subtropical areas of the world [4]. There are only four geminiviruses associated to grapevine: the aforementioned GRBV [5, 6] and another grablovirus found in non-cultivated grapevine (*Vitis sp.*) from North America and named 'Wild Vitis virus 1' (WVV1) [7], Temperate fruit decay-associated virus (TFDaV, unassigned) in Brazil infecting also apple and pear [8] and grapevine geminivirus A (GGVA, unassigned) [9]. These four viruses have monopartite DNA genomes. To our knowledge, no begomoviruses (genus *Begomovirus*) have been linked to grapevine. Begomoviruses have mono or bipartite genomes and are transmitted by whiteflies (*Bemisia tabaci*). Genus *Begomovirus* currently harboring 409 species is the largest genus of plant infecting viruses, some of them causing severe diseases in vegetables and ornamental crops [4].

In order to gain knowledge about grapevine viruses in Argentina, we started surveying *V. vinifera* samples from several regions and cultivars, representing the most significant wine cultivars from the main viticulture areas of the country. A pooled sample consisting of *V. vinifera* cv Torrontes from Mendoza province and four criollas varieties were subjected to high-throughput sequencing (HTS). To this end, total RNAs from cambial scrapings of grapevine samples were purified using the Spectrum Plant RNA Miniprep Kit (Sigma-Aldrich, USA). Equal quantities were pooled and subjected to HTS in a Illumina Novaseq 6000 instrument generating 171,490,780 paired reads (150 nt long). These reads were trimmed and filtered with the Trimmomatic tool, as implemented in http://www.usadellab.org/cms/?page=trimmomatic, and the resulting 154,115,794 reads were mapped with Bowtie2 available at http://bowtie-bio.sourceforge.net/bowtie2/ using standard parameters against the *V. vinifera* genome (assembly 12X) available at https://www.ncbi.nlm.nih.gov/nuccore/CAAP00000000.3. The resulting 26,594,330 unaligned paired reads were assembled *de novo* with Trinity v2.9.0 release with standard parameters. The obtained transcripts were subjected to bulk local Blastx searches (E-value $<1e^{-5}$) against a complete NCBI Refseq of virus proteins database retrieved at ftp://ftp.ncbi.nlm.nih.gov/refseq/release/viral/viral.1.protein.faa.gz. Two contigs of 400 and 320 nt long obtained significant hits to the replication associated protein (REP) of

Sida mosaic Bolivia virus 1 (SiMBoV1, E-value = 3e-87, Identity 91.73%, GenBank accession number YP_004207827.1), and to the nuclear shuttle protein (NSP) of Sida mosaic Bolivia virus 2 (SiMBoV2, E-value = 2e-63, Identity 91.51%, GenBank accession number YP_004207829.1), respectively. To further extend these transcripts the filtered reads were iteratively mapped to these contigs, dubbed putDNA-A and putDNA-B, using the Geneious v8.1.9 platform (Biommaters, USA) map-to-reference tool with low sensitivity parameters as described elsewhere [10,11]. Eventually, both sequences were widely extended and supported by more than two thousand reads each (putDNA-A: 2,680 nt long, supported by 2,121 reads, mean coverage = 97X and putDNA-B: 2,645 nt long, supported by 2,395 reads, mean coverage = 2,348X). Interestingly, read mapping indicated a circular nature of both sequences, which was confirmed by identifying overlapping reads at all positions, and thus a continuum of virus reads supported circularity at all sequence positions.

Blastn analysis of sequences putDNA-A and putDNA-B using default parameters showed the greatest nucleotide sequence identity to DNA component A of an isolate of SiMBoV2 of *Sida micrantha* from Bolivia (HM585443.1, coverage 100%, E-value 0.0, Identity 84.89%), and to DNA component B of an isolate of SiMBoV2 of chia (*Salvia hispanica*) from Argentina (KJ742422.1, coverage 100%, E-value 0.0, Identity 78.61%), respectively (**Table 1**; **Supplementary Table 1**). A rapid search of the sequences for open reading frames (ORFs) greater than 150 nt in size with the program ORF Finder (https://www.ncbi.nlm.nih.gov/orffinder/) indicated that the sequence architecture of putDNA-A and putDNA-B resembled that of SiMBoV2 New World bipartite begomovirus genome. That is, putDNA-A presented one ORF in the virion sense (V) and four in the complementary sense (C), and putDNA-B presented one ORF in the virion sense (V) and one in the complementary sense (C) (**Figure 1**). ORF AV1 of putDNA-A is predicted to encode a 249 aa coat protein (CP) sharing a 94.8% aa pairwise identity (PI) to that of SiMBoV2, and presenting a Gemini_coat domain (pfam00844, coordinates 13-249 aa, E-value = 1.21e-98) according to the NCBI-CDD conserved domain database available at https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi. To trace back these putative virus sequences to the original samples of our pooled HTS library we designed specific primers targeting the CP CDS of putDNA-A. Total RNA was purified from cambium scrapings of each of the grapevine samples employed for the HTS pooled library using the same conditions as described above. The extracted RNA was reverse transcribed using random primers. The resulting cDNAs were amplified by PCR with primers Fw-BEG-DNA-A: TAAGCGGGATCCCTCATGGC and Rv-BEG-DNA-A: TCGTACTTCCCAGCTTCCTGGT, which we designed using PrimerBLAST (https://www.ncbi.nlm.nih.gov/tools/primer-blast/), to avoid *V. vinifera* endogenous targets. PCR reactions included Taq Polymerase (Promega) and used a program consisting of 35 cycles of 30 s at 94 °C, 45 s at 58 °C, 1 min at 72 °C, and a final elongation step of 7 min at 72 °C. The PCR reactions were run on standard agarose gel electrophoresis. Only a single sample corresponding to RNA of a *V. vinifera* cv Torrontes from Mendoza province, Argentina, sampled 2019-May-05 showed PCR products of the expected size (ca. 633 nt) on the gels. The PCR product was cloned and subjected to enzymatic restriction using *HinfI*, which based on our assembled genome would generate two fragments of 154 nt and 479 nt (**Supplementary Figure 4**). PCR products of three clones (1-3) were assessed and the restriction pattern was in line with the single expected cut. Thus, we hypothesized that the circular assembled sequences correspond to genome component A and B a new virus in *V. vinifera* cv

Torrontes that we tentatively dubbed grapevine begomovirus A (GBVA). To entertain this hypothesis we advanced in the structural and functional annotation of the proposed virus sequences. When comparing DNA-A and B of GBVA by MUSCLE sequence alignments we detected a common region (CR), 201 nt long sharing a 95.5% PI. This bipartite begomovirus typical region suggests that the DNA components correspond to the same virus. Further inspection of this CR showed three repeated iteron like regions "TGGTACTCA" (two in the virion and one in the complementary strand) upstream of a TATA box and the canonical nonanucleotide sequence "TAATATT/AC" that marks the origin of virion-strand DNA replication, which was designated as position 1 at the nick region. DNA foldings of the CR using the MFOLD server implemented at http://unafold.rna.albany.edu/?q=mfold/DNA-Folding-Form suggested that, as expected, the nonanucleotide sequence is accessible to the REP protein in the loop of a well-supported stem-loop (**Supplementary Figure 1**). Besides the aforementioned AV1 CDS, DNA-A presented four ORFs in the complementary sense (AC1-AC4) (**Figure 1**). AC1 (coordinates 2,550-1,474 nt) is predicted to encode a 358 aa replication-associated protein (REP) sharing a 86% aa PI to that of SiMBoV2. The N-terminal region presented an Origin of replication-binding domain, RBD-like (SSF55464, coordinates 4-121 aa, E-value = 1.13e-57) as indicated by the SUPERFAMILY HMM search platform implemented at http://supfam.org/SUPERFAMILY/cgi-bin/hmm.cgi. In addition, the REP protein presented a Gemini_AL1 domain (pfam00799, coordinates 7-119 aa, E-value = 3.71e-64), a Gemini_AL1_M domain (pfam08283, coordinates 126-231 aa, E-value = 5.43e-46), followed by a REP associated domain (PD000736, coordinates 248-334 aa, E-value = 3e-17) according to the ProDom tool available at http://prodom.prabi.fr/. The overlapping ORF AC2 (coordinates 1,553-1,164 nt) encodes a 129 aa transcriptional regulator protein (TrAP/AL2) sharing a 86.8% aa PI to that of SiMBoV2, and presented a Gemini_AL2 domain (pfam01440, coordinates 1-124 aa, E-value = 1.67e-42). ORF AC3 (coordinates 1,417-1,019 nt) encodes a 132 aa replication enhancer protein (REn) sharing a 85.6% aa PI to that of SiMBoV2, and presented a Gemini_AL3 domain (pfam01407, coordinates 2-121 aa, E-value = 5.19e-50). Lastly, ORF AC4 (coordinates 2,399-2,136 nt), completely embedded within ORF AC1, encodes a 132 aa AC4 protein (REn) sharing a 77% aa PI to that of SiMBoV2, with a Gemini_C4 domain (pfam01492, coordinates 3-86 aa, E-value = 3.46e-35). On the other hand, the 2,645 nt long DNA-B, with two ORFS, encoded a nuclear shuttle protein (NSP) in virion ORF BV1 (505-1,275 nt coordinates) and a movement protein in ORF BC1 (coordinates 2,260-1,379 nt). The 256 aa NSP shared a 84% aa PI to that of SiMBoV2, and had a Gemini_coat domain (pfam00844, coordinates 7-256 aa, E-value = 1.70e-90). The 293 aa movement protein shared a 93.2% aa PI to that of SiMBoV2, and presented a Gemini_BL1 domain (pfam00845, coordinates 13-292 aa, E-value = 0) (**Figure 1**).

Given the clear evidence based on genomic architecture and structural annotation that the identified virus sequences corresponded to a New World bipartite begomovirus, we advanced with further genetic distance and evolutionary insights. These analyses were oriented to assess whether we could determine if the virus corresponded to a new strain of a previously reported virus such as SiMBoV2, or to a tentative member of a novel species of begomovirus. Firstly, we retrieved the 50 closest hits based on Blastn searches with standard parameters using as query complete DNA components A and B, and additionally African cassava mosaic virus (ACMV, NC_001467- NC_001468). As observed in **Table 1** and **Supplementary Table 1**, no virus sequences presented sequence identity higher than 84.89% (to DNA-A), or 78.61% (to DNA-B).

Nevertheless, the current criteria for begomoviruses recommended by the *Geminiviridae* Study Group of the International Committee on Taxonomy of Viruses indicates a threshold of <91% nt sequence identity for the complete DNA-A as a requisite for species demarcation [12], in the context of identities calculated from true pairwise sequence identities excluding gaps. Thus, as recommended, we employed the Sequence Demarcation Tool (SDT) v. 1.0 [13] with MUSCLE [14] alignment option to generate pairwise sequence comparisons. Our results confirm that the identified virus sequences are well below this <91% nt sequence identity threshold, being SiMBoV2 (HM585443) the highest identity virus with 85.9% PI for DNA-A, and 80.1% PI with SiMBoV2 (KJ742422) DNA-B (**Supplementary Figure 2**, **Supplementary Table 2**). In this scenario, we propose that the detected sequences correspond to a new virus, a tentative member of a novel species, which we suggest the name "grapevine begomovirus A" (GBVA). To complement these findings we generated phylogenetic insights based on multiple genome alignments of both DNA-A and B components using MAFFT v7.017 https://mafft.cbrc.jp/alignment/server/ (1PAM/k2 scoring matrix), using E-INS-i as the best-fit model. The aligned genomes were used as input for maximum likelihood phylogenetic trees using FastTree 2.1.5 as implemented in http://www.microbesonline.org/fasttree/ (best-fit model = JTT-Jones-Taylor-Thorton with single rate of evolution for each site = CAT) computing local support values with the Shimodaira-Hasegawa test (SH) and 1,000 tree resamples, the trees were rooted at ACMV. Our results suggest that GBVA unequivocally clusters within genus *Begomovirus,* in a separated clade with affinity to SiMBoV2 and other Sida sp., Solanaceae, Salvia and Okra viruses (**Figure 2**, **Supplementary Figure 3**). It is worth mentioning that GBVA highest identity viruses correspond to isolates of viruses detected in the region, mostly Brazil and Argentina (**Table 1, Supplementary Table 1)** and that there is an apparent clustering of viruses according to geographic distribution on phylogenetic trees **(Figure 2)**. This is in line with a suggested evolutionary path of divergence through isolation [15] as a consequence of the incapacity of whiteflies, the most plausible vectors, to fly long distances.

All in all, our results suggest that a novel New World bipartite begomovirus is associated to grapevine. As of today, we have no additional information on the infectivity, host range, geographical distribution or biology of GBVA. Nevertheless, we believe that this report, the associated sequences, and the newly developed detection specific-primers would aid in further characterizing the genetic variability of this virus which would eventually redound in an assessment of the prevalence of this agent in the region and worldwide. While the only plant presenting GBVA did not show any obvious symptoms, it is too premature to assume this virus to be asymptomatic. Preliminary assessment with primers Fw-BEG-DNA-A and Rv-BEG-DNA-A failed to detect GBVA in 38 additional grapevine samples from diverse cultivars. To our knowledge, this is the first report of a begomovirus linked to grapevine, which highlights the remarkable diversity of viruses able to infect this economically important crop.

Data availability.

The genome sequences reported here are available at GenBank: GBVA-DNA-A (MT075773), GBVA-DNA-B (MT075774) and as Supplementary Material of this article.

Acknowledgements

We thank P. Schubert for helpful discussions and insightful comments.

Funding

This work was supported in part by project ANPCyT PID: 2018-0021, and by ANPCyT PICT 2015-1532 and PICT 2016-0429. The funders had no role in study design and analysis, decision to publish, or preparation of the manuscript.

Compliance with ethical standards

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

References

1. Dolja VV, Meng B, Martelli GP (2017) Evolutionary aspects of grapevine virology. Grapevine viruses: molecular biology, diagnostics and management. Springer, Cham, pp 659–688. ISBN 978-3-319-57706-7

2. Luna F, Debat H, Moyano S, Zavallo D, Asurmendi S, Gomez-Talquenca S (2019) First report of grapevine red blotch virus infecting grapevine in Argentina. J Plant Pathol 101:1239–1239. https://doi.org/10.1007/s42161-019-00298-3

3. Debat H, Zavallo D, Luna F, Moyano S, Asurmendi S, Gomez-Talquenca S (2019) First report of grapevine virus E infecting grapevine in Argentina. J Plant Pathol 101:1221–1222. https://doi.org/10.1007/s42161-019-00282-x

4. Zerbini FM, Briddon RW, Idris A, Martin DP, Moriones E, Navas-Castillo J, Rivera-Bustamante R, Roumagnac P, Varsani A, and ICTV Report Consortium (2017) ICTV Virus Taxonomy Profile: Geminiviridae. J Gen Virol 98:131–133. https://doi.org/10.1099/jgv.0.000738

5. Perry KL, McLane H, Hyder MZ et al (2016) Grapevine red blotch-associated virus is present in free-living *Vitis* sp. proximal to cultivated grapevines. Phytopathology 106:663–670. https://doi.org/10.1094/PHYTO-01-16-0035-R

6. Krenz B, Thompson JR, McLane H et al (2014) Grapevine red blotch-associated virus is widespread in the United States. Phytopathology 102:1232–1240. https://doi.org/10.1094/PHYTO-02-14-0053-R

7. Perry KL, McLane H, Thompson JR et al (2018) A novel grablovirus from non-cultivated grapevine (Vitis sp.) in North America. Arch Virol. 163:259–262. https://doi.org/10.1007/s00705-017-3567-y

8. Basso MF, da Silva JCF, Fajardo TVM et al (2015) A novel, highly divergent ssDNA virus identified in Brazil infecting apple, pear and grapevine. Virus Res 210:27–33. https://doi.org/10.1016/j.virusres.2015.07.005

9. Al Rwahnih M, Alabi OJ, Westrick NM et al (2017) Description of a novel monopartite geminivirus and its defective subviral genome in grapevine. Phytopathology 107:240–251. https://doi.org/10.1094/PHYTO-07-16-0282-R

10. Debat HJ, Ng TFF (2019) Complete genome sequence of a divergent strain of Tibetan frog hepatitis B virus associated with a concave-eared torrent frog (Odorrana tormota). Arch Virol 164:1727–1732. https://doi.org/10.1007/s00705-019-04227-8

11. Debat HJ, Bejerman N (2019) Novel bird's-foot trefoil RNA viruses provide insights into a clade of legume-associated enamoviruses and rhabdoviruses. Arch Virol 164:1419–1426. https://doi.org/10.1007/s00705-019-04193-1

12. Brown JK, Zerbini FM, Navas-Castillo J, Moriones E, Ramos-Sobrinho R, Silva JC, Fiallo-Olivé E, Briddon RW, Hernández-Zepeda C, Idris A, Malathi VG, Martin DP, Rivera-Bustamante R, Ueda S, Varsani A (2015) Revision of Begomovirus taxonomy based on pairwise sequence comparisons. Arch Virol 160:1593–1619. https://doi.org/10.1007/s00705-015-2398-y

13. Muhire BM, Varsani A, Martin DP (2014) SDT: a virus classification tool based on pairwise sequence alignment and identity calculation. PLOS One 9:e108277. https://doi.org/10.1371/journal.pone.0108277

14. Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinform 5:1–19. https://doi.org/10.1186/1471-2105-5-113

15. Rocha CS, Castillo-Urquiza GP, Lima AT et al (2013) Brazilian begomovirus populations are highly recombinant, rapidly evolving, and segregated based on geographical location. J Virol 87:5784–5799. https://doi.org/10.1128/JVI.00155-13

Figure Legends

Figure 1. Molecular characterization of Grapevine begomovirus A (GBVA). Genome graph depicting predicted circular DNA components A and B of GBVA. Start and end coordinates of each gene product are indicated. Arrowed rectangles represent each ORF. Curved yellow rectangles represent each predicted proteins and conserved domains are shown in pink. In the lower panel the circular track indicates GC contant variation (GC blue, AT green). Abbreviations: AV1/BV1, virion strand ORF 1 of DNA-A or DNA-B; AC1-4/BC1, complementary strand ORF1-4 of DNA-A or DNA-B; Ren, replication enhancer protein; TrAP, transcriptional regulator protein; CRA/CRB, common region of DNA-A or DNA-B.

Figure 2. Phylogenetic insights of Grapevine begomovirus A (GBVA) based on MAFFT alignments and maximum likelihood trees generated with DNA-A of GBVA and reported begomoviruses. The scale bar indicates the number of substitutions per site. Node labels indicate FastTree support values. The tree is rooted at ACMV. Genbank Accession numbers are indicated following the virus names. Viruses corresponding to the same species are demarcated with colored rectangles. Standard ICTV virus abbreviations are shown on the right.

Supplementary Figure 1. (**A**) MUSCLE sequence alignments of DNA-A and B of GBVA common regions (CR), sharing a 95.5% PI. The CR presented three repeated iteron like regions "TGGTACTCA" (shown in green) upstream of a TATA box (in red) and the canonical nonanucleotide sequence "TAATATT/AC" (in purple) flanked by complementary regions (in red). (**B**) DNA folding of a sub region of the CR using the MFOLD server showing a well-supported stem-loop flanking the typical nonanucleotide sequence in the loop. The bases are colored by base pair probability from blue to red. Nonanuclotide sequence is demarcated with a colored rectangle. Nick region is indicated with a dotted line.

Supplementary Figure 2. Pairwise sequence comparisons plots of GBVA DNA-A (**A**) or DNA-B (**B**) and reported begomovirus including the 50 closest Blastn hits using as query the complete genome complement and ACMV. The plots were generated with the Sequence Demarcation Tool (SDT) v. 1.0 with MUSCLE alignment option. SDT matrix scores are available in **Supplementary Table 2**.

Supplementary Figure 3. Phylogenetic insights of Grapevine begomovirus A (GBVA) based on MAFFT alignments and maximum likelihood trees generated with DNA-A of GBVA and reported begomoviruses. The scale bar indicates the number of substitutions per site. Node labels indicate FastTree support values. The tree is rooted at ACMV. Genbank Accession numbers are indicated following the virus names. Viruses corresponding to the same species are demarcated with colored rectangles. Standard ICTV virus abbreviations are shown on the right.

Supplementary Figure 4. PCR/Restriction assay confirmation of virus sequence. Specific primers were designed to detect GBVA DNA-A, targeting the AV1 CDS (Fw-BEG-DNA-A and Rv-BEG-DNA-A). The only sample which generated an amplification product (ca. 633nt) corresponded to *V. vinifera* cv Torrontes from Mendoza province. The PCR product was cloned re-amplified and subjected to enzymatic restriction using HinfI, which based on our assembled genome would generate two fragments of 154 nt and 479 nt. PCR products of three clones (1-3) were assessed and the restriction pattern was in line with the expected cut.