Mapping human laryngeal motor cortex during vocalization

Nicole Eichert[1], Daniel Papp[1], Rogier B. Mars[1,2] & Kate E. Watkins[3]

[1] Wellcome Centre for Integrative Neuroimaging, Centre for Functional MRI of the Brain

(FMRIB), Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of

Oxford, Oxford, United Kingdom

[2] Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen,

Nijmegen, The Netherlands

[3] Wellcome Centre for Integrative Neuroimaging, Department of Experimental Psychology,

University of Oxford, Oxford, United Kingdom

Abbreviated title: human laryngeal motor cortex

Correspondence:

Nicole Eichert

Wellcome Centre for Integrative Neuroimaging

Nuffield Department of Clinical Neurosciences

John Radcliffe Hospital

University of Oxford

Oxford OX3 9DU

United Kingdom

Direct Line: +44 (0) 1865 271363

E-mail: nicole.eichert@psy.ox.ac.uk

<u>Abstract</u>

The representations of the articulators involved in human speech production are organized

somatotopically in primary motor cortex. The neural representation of the larynx, however,

remains debated. Both a dorsal and a ventral larynx representation have been previously

described. It is unknown, however, whether both representations are located in primary motor

cortex. Here, we mapped the motor representations of the human larynx using fMRI and

characterized the cortical microstructure underlying the activated regions. We isolated brain

activity related to laryngeal activity during vocalization while controlling for breathing. We also

mapped the articulators (the lips and tongue) and the hand area. We found two separate

activations during vocalization – a dorsal and a ventral larynx representation. Structural and

quantitative neuroimaging revealed that myelin content and cortical thickness underlying the

dorsal, but not the ventral larynx representation, are similar to those of other primary motor

representations. This finding confirms that the dorsal larynx representation is located in

primary motor cortex and that the ventral one is not. We further speculate that the location

of the ventral larynx representation is in premotor cortex, as seen in other primates. It remains

unclear, however, whether and how these two representations differentially contribute to

laryngeal motor control.

<u>Keywords</u>

- Functional MRI

- Larynx

- Myelin

- Quantitative MRI

- Speech

1. Introduction

The voluntary control of highly complex speech movements is regarded as one aspect of behavior that is unique to humans (Fitch 2017). Speech production requires the fine coordination of a large number of muscles to control the supralaryngeal articulators, respiration, and the vocal folds in the larynx during voice production (Jürgens 2002). In addition to its role as main source of vocal sound production, the larynx is implicated in various biological functions such as protection of the airways and swallowing (Ludlow 2005). Several pairs of intrinsic and extrinsic muscles connect the laryngeal cartilages with each other and to the skeleton. During vocalization, these muscles are controlled in a complex fashion, so that the tension in the vocal folds allows them to be set into vibration as air from the lungs passes through them. Several lines of evidence showed that the ventral part of the precentral gyrus and the central sulcus in the human brain are involved in speech motor control (Bohland and Guenther 2006; Ackermann et al. 2014). The question as to which brain areas specifically control laryngeal activity during vocalization, however, remains debated.

Nearly 100 years ago, direct cortical stimulation of the ventral portion of the precentral gyrus in the human brain was shown to elicit vocalization (Foerster 1936; Penfield and Boldrey 1937). In more recent times, functional brain imaging studies show highly inconsistent results when mapping laryngeal activity during vocalization (reviewed in Belyk & Brown, 2017). Several studies report activity evoked by vocalization in both a ventral portion of the precentral gyrus located close to the Sylvian fissure and in a more dorsal portion of central sulcus and precentral gyrus (Terumitsu et al. 2006; Galgano and Froud 2008; Olthoff et al. 2008; Grabski et al. 2012). Some studies report vocalization-evoked activity in the dorsal location only (Sörös et al. 2006;

Brown et al. 2008; Kleber et al. 2013; Belyk and Brown 2014; Belyk et al. 2018). Only a few

studies specifically isolated laryngeal activity during voice production by contrasting it with

supralaryngeal articulation (Sörös et al. 2006; Terumitsu et al. 2006; Brown et al. 2009; Grabski

et al. 2012), but the results were inconsistent across studies.

Another line of evidence comes from direct neurophysiological recordings from the cortical

surface using implanted high-density electrode arrays in patients being prepared for epilepsy

surgery. These studies also show activity in both a dorsal and a ventral region during speech

production (Bouchard et al. 2013; Chang et al. 2013; Toyoda et al. 2014; Breshears et al. 2015;

Dichter et al. 2018). Interpretation of electrocorticography (ECoG) studies, however, is

inherently limited due to the typically small sample size and potential neurological

abnormalities in the patients' brains.

In addition to these inconsistent results, an additional confound in several studies mentioned

above is breathing. Voice production and respiration functionally interact during speech

production and volitional expiration has been shown to activate motor areas that are located

close to the putative dorsal laryngeal representation (Ramsay et al. 1993; Evans et al. 1999;

McKay et al. 2003). Several previous studies, however, did not control for breathing (e.g. Brown

et al., 2008; Sörös et al., 2006). Those studies that specifically studied vocalization and

breathing found largely overlapping activity during both conditions and a contrast showed no

difference in motor cortex indicating that the activity was not specific to larynx activity (Loucks

et al. 2007; Simonyan et al. 2009).

In addition to a lack of control for confounds such as breathing and supralaryngeal articulation,

most neuroimaging studies mentioned above did not assess individual differences in the

activation patterns. It is common to report group-level cluster-corrected results following

volumetric nonlinear image registration of task-activation maps. Interpreting group-level

results, however, might obscure subject-specific features and inter-individual variability, which

limits sensitivity and functional resolution (Bennett and Miller 2010; Nieto-Castañón and

Fedorenko 2012; Bouchard et al. 2013; Woo et al. 2014). This lack of individual detail might

have caused a failure to detect one of the cortical larynx representations in previous studies.

Moreover, averaging of small sample sizes with high variability, as often performed in ECoG

studies, can show two distinct larynx representations, even when individual patients show

activity in only one of the regions (Bouchard et al. 2013).

Comparisons of human brains and those of other primates indicate strong species differences

in the neural organization underlying laryngeal control during vocalization (Ackermann et al.

2014; Simonyan 2014; Kumar et al. 2016). Most notably, the location of the proposed human

dorsal larynx representation in primary motor cortex is more dorsal-posterior than the non-

human primate homolog, which is in a ventral premotor cortex area (Leyton and Sherrington

1917; Hast et al. 1974; Jürgens 1974; Simonyan and Jürgens 2002; Coudé et al. 2011).

These various findings have led to the proposal of an evolutionary 'duplication and migration'

hypothesis that the larynx motor cortex comprises two structures located dorsally and

ventrally in the human brain (Belyk and Brown, 2017; Jarvis, 2019; reviewed in Mars et al.,

2018). This theory proposes that the dorsal larynx representation is unique to humans and that

it evolved in primary motor cortex due to our especially high demands on laryngeal motor

control. The ancestral primate larynx representation in premotor cortex is presumed to be

homologous to the human ventral larynx representation, which migrated posteriorly,

potentially into human primary motor cortex.

The cortical areas underlying the larynx representations, however, are currently unknown and

it has not been tested if the ventral larynx representation is in primary motor or in premotor

cortex. A myeloarchitectonic approach, which has also become available for neuroimaging,

enables us to describe some properties of cortex (Kuehn et al. 2017). Primary motor cortex,

for example, is characterized by higher cortical myelin content and higher cortical thickness

compared to adjacent premotor cortex and with somatosensory cortex located on the caudal

bank of the central sulcus and postcentral gyrus (Fischl and Dale 2000; Glasser and van Essen

2011; Lutti et al. 2014). Describing the anatomical parcels underlying the larynx

representations can inform evolutionary hypotheses and provide clues about their functional

relevance.

This study sought to determine the anatomical location of larynx-related neural activity in

individual subjects and to characterize the cortical structure underlying these representations.

Our experimental design aimed to isolate brain activity related to voice production by

controlling for breathing-related movements and movements of articulators. In one task, we

identified the (supralaryngeal) articulation and the (laryngeal) vocalization component of

speech during syllable production using a factorial design described in a previous study

(Murphy et al. 1997). We refer to the latter 'vocalization' component as an index for laryngeal

activity during voice production, while other studies have referred to it as 'phonation' or

'voicing'. In a second task, we localized the separate neural representations of lip, tongue and

larynx using highly controlled basic movements, while breathing movements were matched

across conditions.

In order to characterize the microstructural properties underlying the larynx representation,

we compared their myelin content and cortical thickness derived from structural and

quantitative MRI measurements. These quantifications give an indication of the type of cortex

underlying the activated regions, to inform our knowledge about the organization of the

human larynx motor cortex.

2. Materials and Methods

Subjects

20 subjects (12 females, $18 - 40$ years (27.4 ± 5.6, mean $\pm$ SD), 5 self-reported left-handers)

took part in the study. All subjects were self-reported native English speakers; two were raised

bilingually and three were fluent in a second language. All reported normal hearing, normal or

corrected-to-normal vision, no neurological impairments and no history or diagnoses of speech

disorders. The study was approved by the Central University Research Ethics Committee

(CUREC, R55787/RE001) in accordance with the regulatory standards of the Code of Ethics of

the World Medical Association (Declaration of Helsinki). All subjects gave informed consent to

their participation and were monetarily compensated for their participation.

Experimental design and task

We used two tasks to map the motor representations of the articulators and the larynx. Task
1 engaged the speech motor system in an ecologically valid way that required the participants
to utter a short syllable sequence in different conditions. Task 2 required the participants to
perform basic movements that are commonly used in other localizer studies.

Subjects practiced all tasks outside the scanner to be sure they understood the task
requirements. Articulator movements and vocalizations as well as breathing were
demonstrated by the experimenter and practiced until the subjects performed them as
required.

Task 1 - syllable production task

Subjects were instructed to produce the utterance "/la leɪ li la leɪ li/" in four different
conditions: speaking (overt speech), supralaryngeal articulation only (silent mouthing), overt
vowel sound production of vowel /i/ six times but no articulation (vowel production), and
thinking (covert speech). The vowel /i/ was chosen because it is a natural and familiar sound
that requires laryngeal movement during vocalization, but involves minimal movement of the
jaw muscles, lips and pharyngeal part of the tongue (Grabski et al. 2012). Subjects were
instructed not to whisper during the 'silent mouthing' or 'covert speech' conditions as this
would involve laryngeal motor activity.

For all conditions, the breathing pattern was explicitly instructed using the fixation symbol on

the screen (Figure 1A). Subjects were instructed to inhale for 1.5 s (fixation was a square) and

exhale for 4 s (fixation was a cross); the utterance in each condition was performed once during

each 4-s exhalation. Each condition was performed in blocks lasting 22 s, which corresponded

to four repetitions of the breathing cycle. Each block was followed by a rest period of 8 s with

normal breathing (i.e. not explicitly instructed). This rest period allowed the subjects to relax

their breathing patterns and to maintain a comfortable respiratory state. The four conditions

were presented in a fixed pseudo-random order following a balanced Latin-square design

wherein each condition was repeated five times; each condition followed and preceded each

of the other conditions once and the same condition was not presented consecutively.

Subjects were instructed to keep their mouth slightly open during the whole session and to

keep the jaw relaxed.

Task 2 - basic functional localizer

There were four task conditions: tongue retraction, lip protrusion, production of the vowel

sound /i/, and a 'breathing only' condition. The three task conditions for basic speech

movements were contrasted with the 'breathing only' condition in which breathing was

explicitly instructed as above. Articulator movements and vowel production were repeated

during 4 s of exhalation at a rate of approximately 1 - 2 repetitions/s as described for task 1.

Subjects were instructed to keep the movement rate constant throughout the scan for all

movement types. Breathing instructions, task timing and randomization of the blocks were the

same as described for task 1 above, except that each condition was repeated four times during

the scan run. Note that each 22-s block was followed by an 8-s period of rest with normal

breathing, as above.

Hand localizer

During the scanning session, subjects performed the basic localizer first and then the syllable

production task. In between these two tasks involving vocalizations, the subjects performed a

phonological and semantic judgement task (Devlin et al. 2003). Participants had to indicate a

yes/no response by pressing a button with the right index or the middle finger every 3 s. The

task data was used here only to localize the hand representation in the left hemisphere; the

language task data are not reported.

MRI Data Acquisition

MRI data were obtained at the Oxford Centre for Human Brain Activity (OHBA) using a 3-T

Siemens Prisma scanner with a 32-channel head coil. Two structural images of the whole brain

were acquired; a T1w image (MPRAGE sequence; 1 mm^3 isotropic resolution, TR = 1900 ms, TE

= 3.97 ms, TI = 905 ms, 8° flip angle, bandwidth = 200 Hz/pixel, echo spacing = 9.2 ms, FOV =

192 × 192 × 174 mm^3) and a T2w image (SPACE turbo-spin-echo sequence; 1 mm^3 isotropic

resolution, TR = 3200 ms, central TE = 451 ms, variable flip angle, bandwidth = 751 Hz/pixel,

echo spacing = 3.34 ms, echo train duration = 919 ms, Turbo Factor = 282, FOV = 256 × 256 ×

176 mm^3, GRAPPA acceleration factor 2).

For task fMRI, whole-head T2*-weighted echo-planar images (TE = 30 ms) were acquired using

multiband sequence (factor 6, TR = 0.8) at 2.4 mm^3 isotropic resolution. Task fMRI parameters

were adopted from the ABCD study (https://biobank.ctsu.ox.ac.uk/crystal/docs/brain_mri.pdf,

Casey et al., 2018). Two task fMRI scans were conducted lasting 8 min (600 volumes, task 2)

and 10 min (750 volumes, task 1). In between the two tasks, subjects performed a phonological

and semantic judgement task for 9 min, which did not involve vocalization.

Furthermore, a multiparameter mapping (MPM) protocol was acquired (Weiskopf et al. 2013;

Lutti et al. 2014). Proton density-weighted (MPM$_{PDw}$), magnetization transfer-weighted

(MPM$_{MTw}$) and T1-weighted (MPM$_{T1w}$) images were acquired using a tailored pulse sequence

(1 mm^3 isotropic resolution, FOV = 256 × 224 × 176 mm^3, TR = 25 ms, bandwidth = 488

Hz/pixels, first TE/echo spacing = 2.3/2.3 ms, 6° flip angle (MPM$_{PDw}$, MPM$_{MTw}$) or 21° (MPM$_{T1w}$),

slab rotation = 30°, and number of echoes = 8/6/8 (MPM$_{PDw}$/MPM$_{MTw}$/MPM$_{T1w}$ respectively),

GRAPPA acceleration factor 2 × 2, 40 reference lines in each phase encoded direction.

Quantitative R1 (= 1 / T1) maps were estimated from the MPM$_{PDw}$ and MPM$_{T1w}$ images as

demonstrated in Weiskopf et al. (2013), which was extended by including a correction for radio

frequency transmit field inhomogeneities (Lutti et al. 2010) and imperfect spoiling (Preibisch

and Deichmann 2009). Regression of the log-signal from the signal decay over echoes across

all three MPM contrasts was used to calculate a map of R2* (= 1 / T2*) (Weiskopf et al. 2014).

The transmit field map was calculated using a 3D echo-planar imaging (EPI) spin-echo

(SE)/stimulated echo (STE) method (Lutti et al., 2012, 2010; FOV = 256 × 192 × 192 mm^3, matrix

= 64 × 64 × 48 mm^3, TE = 39.06, mixing time = 33.8 ms, TR = 500 ms, nominal α varying from

115° to 65° in steps of 5°, acquisition time 4 min 24 s) and was corrected for off-resonance

effects using a standard B0 field map (double gradient echo FLASH, 3 × 3 × 2 mm^3 resolution,

whole-brain coverage). The MPM parameter maps took approximately 20 minutes to acquire.

In addition to the MRI scans mentioned here, participants underwent a diffusion-weighted

scan and a resting-state scan (data not reported here). The total duration of the entire scanning

session was approximately 1.5 h, preceded by approximately 45 min of briefing and task

practice.

Physiological recordings were carried out throughout scanning using a respiratory belt to

measure the breathing pattern and a pulse oximeter to monitor the heart rhythm during the

scan. Physiological data were recorded using a Biopac MP150 (Biopac, Goleta, CA, USA) at a

sampling frequency of 500 Hz. Subjects wore ear-plugs and MRI-compatible head phones

(OptoActive-II, Optoacoustics Ltd, Moshav Mazor, Israel), which reduced scanner noise using

electrodynamic noise-cancelling. At the beginning of the scanning session, the headphones

were calibrated and noise-cancelling performance was further monitored throughout the

session. Prior to each functional scan the attenuation algorithm 'learned' the scanner noise for

16 s. During the two tasks involving vocalizations, subjects were audio-recorded using an MRI-

compatible microphone with noise cancelling (Dual Channel FOMRI-III, Optoacoustics Ltd,

Moshav Mazor, Israel) at a sampling rate of 22,050 Hz.

 Behavioral data analysis

The subjects' vocal behavior for the tasks was manually assessed using the audio recordings.

The breathing patterns during the task blocks recorded using the Biopac were inspected

visually to verify that subjects complied with the breathing instruction. Individual breathing

traces were cropped into segments of 30 s, which consist of 22 s of instructed breathing during

the task block and 8 s of subsequent rest period with normal breathing (Figure 1C).

Structural MRI analysis

T1w and T2w scans were pre-processed using the HCP-pipeline (Glasser et al. 2013) cloned

from the 'OxfordStructural' - fork (https://github.com/lennartverhagen/Pipelines). The

processing pipeline includes anatomical surface reconstruction using FreeSurfer

and automatic assignment of neuroanatomical labels (Fischl 2012; Jenkinson et al. 2012). The

T2w image was registered to the T1w image using FSL's FLIRT using spline interpolation

(Jenkinson et al. 2002).

The image of the T1w scan was divided by the image of the T2w scan to create a T1w/T2w-

ratio image. The T1w/T2w-ratio was mapped onto the native midthickness surface and then

resampled to the 164k standard (fs_LR) surface mesh using adaptive barycentric

interpolation (approx. 164.000 vertices per hemisphere) using Workbench Command

(www.humanconnectome.org/software/connectome-workbench.html). Mapping was

performed with the `-volume-to-surface-mapping` command using the `-myelin-style` option.

This T1w/T2w map has been shown empirically to correlate with cortical myelin content

(Glasser and van Essen 2011; Glasser et al. 2014). In addition to T1w/T2w myelin maps, the

HCP-pipeline provides automatic generation of cortical thickness surface maps.

MPM parameter maps were reconstructed and pre-processed using the hMRI-toolbox

(Tabelow et al. 2019) embedded in the Statistical Parametric Mapping framework

(SPM12, Wellcome Trust Centre for Neuroimaging, London, UK). For one subject, MPM data

were excluded due to motion-induced blurring. MPM_{MT}, MPM_{R1} and MPM_{R2*} maps were

registered to the MPRAGE T1w scan using FLIRT rigid-body transformation and spline

interpolation and then mapped to the surface using the same steps as for the T1w/T2w myelin

map. The three MPM parameter maps have been shown to correlate to different degrees with

myelin content in white matter, subcortical structures and grey matter (Draganski et al. 2011;

Callaghan et al. 2014; Lutti et al. 2014; Bagnato et al. 2018). Finally, one step of surface-based

smoothing (FWHM = 4 mm) was applied to the five surface maps of interest - T1w/T2w map,

three MPM parameter maps and cortical thickness map.

fMRI data analysis

Functional MRI data processing was carried out using FEAT (FMRI Expert Analysis Tool) Version

6.00, part of FSL (FMRIB's Software Library, www.fmrib.ox.ac.uk/fsl). The following pre-

statistics processing was applied: motion correction using MCFLIRT (Jenkinson et al. 2002);

non-brain tissue removal using BET (Smith 2002); spatial smoothing using a Gaussian kernel

(FWHM = 5 mm); grand-mean intensity normalization of the entire 4D dataset by a single

multiplicative factor; low-frequency drifts were removed using a temporal high-pass filter with

a cut-off of 90 s for all three tasks. Motion corrected images were unwarped using a fieldmap

and PRELUDE and FUGUE software running in FSL (Jenkinson 2003). Registration to the high

resolution structural scan and standard 2 mm MNI-152 template was carried out using FLIRT

(Jenkinson and Smith 2001; Jenkinson et al. 2002). Registration from high resolution structural

to standard space was then further refined using FNIRT nonlinear registration (Andersson et

al. 2007).

Time-series statistical analysis was based on a general linear model (GLM) implemented in

FILM with local autocorrelation correction (Woolrich et al. 2001). Standard motion correction

parameters and individual volumes that were motion outliers, determined using

fsl_motion_outliers, were included as separate regressors at the first level for each subject.

For the syllable production task (task 1), the conditions were analyzed in a factorial model that

allowed the (supralaryngeal) articulation and the (laryngeal) vocalization component of the

task to be separated (Figure 1B). Brain activity associated with the control of articulation was

defined as ('overt speech' *minus* 'vowel production') *plus* ('silent mouthing' *minus* 'covert

speech') and the main contrast for vocalization was derived by the contrast ('overt speech'

minus 'silent mouthing') *plus* ('vowel production' *minus* 'covert speech'). The 8-s periods of

rest with normal breathing between condition blocks served as baseline and were not

modelled as a separate explanatory variable in the GLM; they were not included in contrast to

any task condition in the analysis of this factorial design.

For the basic localizer task (task 2), activity during each condition for basic speech movements

was assessed relative to the 'breathing only' condition with instructed breathing. The periods

of rest with normal breathing in between conditions served as baseline (i.e. they were not

modelled in the GLM as described for task 1). For the hand localizer task, we derived a contrast

of all task conditions involving button presses relative to the rest blocks (note, the breathing

instruction was not used in the hand localizer task).

All contrasts reported in the results for both tasks were assessed relative to conditions with

matched breathing. In addition, we report results of all individual task conditions relative to

the resting condition with normal breathing in a supplementary figure (Figure S1).

Volumetric group average activation maps

To obtain volumetric group average maps, each individual's statistical maps were transformed

to standard space (MNI152) using a nonlinear registration. Volumetric group mean activation

maps were obtained using mixed effects in FLAME (FMRIB's Local Analysis of Mixed Effects,

(Woolrich et al. 2004, Stage 1 only), where subjects are treated as random effects. Group-level

z-statistic images were thresholded using a voxel-wise threshold of $z > 3.5$ ($p < 0.00025$,

uncorrected). Note that the corrected threshold for a whole-brain analysis of these data is $z >$

5. Rather than use a small volume correction and ROI masking, we chose to display results at

the lower threshold to better visualize the activity in the whole brain. The figures of the

volumetric group-level results (Figure 2, 3) are focused, i.e. centered, on the voxel of maximal

z-value within the right hemisphere central somatomotor strip. For the vocalization contrast

(Figure 2) and the vowel production condition (Figure 3), we generated two separate figures

focused on the right dorsal and the ventral larynx representation.

Surface group count maps

In addition to volume-based analysis, task MRI activations were assessed using surface-based

analyses. Surface space permits a better visualization of cortical brain activations.

Furthermore, this allowed us to perform surface-based quantifications of cortical

microstructures.

Each subject's *z*-statistic images were thresholded voxel-wise at $p < 0.05$ (corrected using a *z*-

value determined based on data smoothness and the RESEL count). This thresholded map was

projected onto the individual's native midthickness surface using the '-ribbon-enclosed' option

in wb_command `-volume-to-surface-mapping`. To derive group-level surface count maps, all

individual subject thresholded maps were resampled to the 32k standard (fs_LR) mesh based

on the FreeSurfer registration, binarized and then summed at each vertex (Barch et al. 2013).

The group count maps are shown on an inflated average brain surface, thresholded at $n > 4$

subjects. They give an indication of the between-subject variability in the areas activated by

task, providing complementary information to maps of activity averaged across participants.

Individual surface activation maxima

Surface-based activations were studied further using an ROI-based approach to isolate and

focus on activations in the central somatomotor strip. The two tasks provided complementary

results for further analysis with Task 1 serving as a robust localizer for the larynx

representations, and Task 2 providing an accurate localizer for lip and tongue representations.

In order to assess intra-individual spatial variability of the motor representations, we derived

the location of individual activation maxima in both hemispheres from selected task contrasts:

For the larynx, we used the main contrast for vocalization during the syllable production task

(task 1). For lip and tongue, we used the task contrasts from the basic localizer task (task 2)

and for the hand, we used the hand localizer task (left hemisphere only).

Different ROI masks were used for the different motor representations based on individual

anatomy. In short, we used an ROI of the whole central sulcus for hand, lip and tongue, a more

limited portion of the central sulcus ROI for the dorsal larynx representation and a manually

defined ROI for the ventral larynx. ROI definitions are described in more detail in the

supplementary material.

Individual volumetric ROIs were linearly transformed from FreeSurfer's anatomical to

functional space of the respective task fMRI scan. Within the ROI, the voxel of maximal

intensity was determined from the uncorrected z-statistics image. It should be noted that for

some subjects this local maximum did not achieve the corrected voxel-wise significance

threshold, which was used to create the surface count maps (left hemisphere: hand n = 3,

dorsal larynx n = 6, ventral larynx n = 5; right hemisphere: dorsal larynx n = 5, ventral larynx n

= 4). Using a lower uncorrected threshold is justified given our goal to visualize and assess

spatial variability of the activation maxima. Activation maxima were manually inspected in the

subject's native volume space to confirm that the systematic approach described below

captured task-related activations. The activation maxima were mapped to the individual's

native midthickness surface, resampled to the 32k standard (fs_LR) surface mesh using the

FreeSurfer registration, smoothed (FWHM = 1 mm), and binarized to form a small circular

patch.

Given that the two tasks had some conditions in common, we were able to examine within-

subject reliability of the location of activation maxima for larynx and tongue as described above

compared with those derived based on the vowel condition and the main contrast for

articulation. We computed the geodesic distance on the individual's native midthickness

surface for pairs of maxima (tongue condition and main contrast for articulation; main contrast

for vocalization and vowel condition).

Cortical surface features at activation peaks

We described the cortical microstructure at the individual activation peaks based on different

surface measures. In order to assess cortical myelin content, we used the T1w/T2w map and

the MPM parameter maps (MPM_{MT}, MPM_{R1} and MPM_{R2*}). We extracted the mean value for

each individual subject's surface map within the area defined by the circular patch around the

vertex of maximal activation for hand, lip, tongue, dorsal larynx and ventral larynx

representation, i.e. at the peaks shown in Figure 4A. These measures were obtained from the

subject's native surface mesh prior to resampling to the standard mesh. In order to account

for the different ranges in intensities and to allow direct comparison of the maps, we computed

z-scores based on the distribution of individual values in each map and each ROI. We assessed

the differences in z-scores across the different ROIs using a linear mixed effects analyses as

implemented in R's *lmer* function (Bates and Sarkar (2007), Core Team and Foundation for

Statistical Computing). The model included fixed effects for ROI (dorsal larynx, lip, tongue,

ventral larynx), hemisphere (left, right) and map (T1w/T2w, MPM_{MT}, MPM_{R1}, MPM_{R2*}) and

random effects for subject. A Shapiro-Wilk test revealed a normal distribution of the data at a

significance level of $p > 0.001$.

In addition to cortical myelin content, we also quantified cortical thickness values at the same

individual surface peaks. To increase sensitivity of the quantification described above, we

excluded activation maxima that were located in a part of cortex with lower cortical thickness

and thus likely to be activity in somatosensory rather than motor cortex (Fischl and Dale 2000).

We used a heuristic lower threshold of 2 mm to exclude sensory activation maxima and then

re-ran the statistical evaluation of myelin-values described above. A linear mixed effects

analysis for the fixed effects of ROI and hemisphere and random subject effects was performed

to assess cortical thickness values after exclusion of the sensory activation maxima.

In order to further characterize the differences in myelin content across the ROIs, we used an

additional quantification: We computed the pair-wise Manhattan distance across the ROIs

based on a vector of the four raw (i.e. non-z-transformed) values in each subject after excluding

the sensory activation maxima. The differences across ROIs were visualized in form of a

dissimilarity matrix, where we averaged Manhattan distances within each ROI first across

subjects and then across hemispheres.

A linear discriminant analysis (LDA) was run to explore which combination of z-transformed

surface features (T1w/T2w, MPM$_{MT}$, MPM$_{R1}$, MPM$_{R2*}$), best discriminated the ventral from the

dorsal larynx representation after excluding sensory activation maxima. The parameters of the

LDA were estimated using a singular value decomposition with no shrinkage.

<u>3. Results</u>

We acquired fMRI data in 20 subjects during performance of two tasks: (1) a syllable

production task required subjects to produce "/la leɪ li la leɪ li/" overtly, mouthed silently, and

covertly, and to produce the vowel /i/, (2) a basic localizer task required subjects to make small

repetitive movements of the lips, tongue and larynx (vowel production); an additional task was

used as functional localizer for movement of the right hand.

Auditory recordings

All subjects vocalized as instructed during the conditions that involved vocalizations in both

tasks. In all other conditions, subjects remained silent, as instructed.

Breathing during vocalization tasks

We recorded the breathing traces using a breath belt during both vocalization tasks to confirm

that the breathing patterns were comparable across different conditions (Figure 1C). As

expected for all conditions in both tasks, four breathing cycles were visible in the first 22 s,

during which breathing was instructed. Note that in one subject, where an extension of the

breathing belt was used, the breathing trace differed in overall shape, but the breathing cycles

were still visible. Examination of these figures shows that the shape of the trace and variability

were similar across all conditions, but some differences were observed. For example, in the

vowel production condition of the basic localizer task, exhalation was more gradual and less

rapid than in the other four conditions. All four conditions in the syllable production task

showed a more gradual exhalation pattern than in the basic localizer task.

Figure 1 – Task paradigm and breathing patterns. **A**: Timing of the task paradigm for an example block of the

'overt speech' condition from the syllable production task (task 1). The utterance was repeated four times

while breathing was instructed followed by a resting period with normal breathing. Square – inhale, + -

exhale. **B**: Factorial design used in the syllable production task. The four task conditions can be described

according to orthogonal main factors for laryngeal activity during vocalization and supralaryngeal

articulation. Breathing was instructed during all conditions as shown in A. **C**: Breathing traces during syllable

production task (left two panels) and basic localizer task (right two panels). Black line: group mean after

normalizing all individual breathing traces to the same amplitude; grey lines: average traces for individual

subjects (n = 20). The dashed vertical line indicates the end of the 22-s task block and the beginning of the

8-s resting period with normal breathing.

529
530

Syllable production task (task 1)

Syllable production task - Volumetric results

We localized the cortical activations for movement control of (supralaryngeal) articulation and

(laryngeal) vocalization during syllable production (Figure 2A). MNI coordinates of activation

maxima in the somatomotor strip are reported in the supplemental material (Table S1). The

main contrasts for vocalization and articulation were defined based on a factorial design that

combined the data from all four task conditions in the syllable production task (Figure 1B). The

main contrast for articulation showed activity in the mid-portion of the central gyrus. Note that

we also found spurious activity in prefrontal white matter, which we presumed was induced

by task-correlated movement.

The main contrast for vocalization showed activity in two somatomotor regions in both

hemispheres: one located within a dorsal region of central sulcus, and a second located in a

more anterior-ventral region. Portions of the superior temporal gyrus were also activated

during vocalization. This is presumed to reflect auditory perception of self-generated

vocalizations. The dorsal and ventral activations were separate and did not appear to be

connected.

Vocalization and articulation also activated parts of cerebellum and SMA in a somatotopic

fashion. Group-level volumetric results for cerebellum and SMA in both functional tasks are

described in the supplementary material (Figure S2). In SMA, one single representation for

laryngeal activity during vocalization was observed, while in cerebellum, two distinct

representations were found.

Syllable production task - Surface results

Group count maps of the syllable production task projected to the surface showed that several

vertices in the mid-portion of the central sulcus were commonly activated during articulation

in 19 out of 20 subjects (Figure 2B). The group count map for vocalization showed separate

dorsal and ventral regions robustly activated across the group. In the main contrast for

vocalization, there was greater variability in the exact location of activity above threshold and

the areas activated in individuals were smaller than for the main contrast for articulation. This

resulted in less overlap of activated vertices for the vocalization group maps. For some

subjects, vocalization-induced activity did not reach significance in relevant brain regions,

which also resulted in lower values in the count map.

Figure 2 – Syllable production task. **A**: Whole-brain group activation maps showing areas activated during

articulation and vocalization (voxel-wise threshold $z > 3.5$, $n = 20$). For the main contrast for vocalization

(blue), both a dorsal and a ventral representation are shown in separate panels. Black arrow: dorsal larynx

representation; white arrow: ventral larynx representation. **B**: Surface group count maps of the same

contrasts. Individual surface maps were thresholded at $p < 0.05$ (corrected voxel-wise), binarized and

resampled onto the fs_LR surface. These were summed across the group of 20 subjects and thresholded at

$n > 4$. Note: in both analyses, vocalization also resulted in activation of the superior temporal gyrus,

presumably reflecting auditory stimulation due to hearing oneself vocalizing.

Basic localizer task (task 2)

Basic localizer task - Volumetric fMRI results

Brain activation during movement of the tongue and the lips and for laryngeal activity during

vowel production was assessed by contrasting basic speech movements with a 'breathing only'

condition, which was matched for controlled breathing. We found distinct activation peaks

that followed the predicted somatotopic organization in the mid-portion of the central sulcus

with the tongue representation more ventral than the lip representation in both hemispheres

(Figure 3A) (Penfield and Rasmussen 1950; Grabski et al. 2012; Carey et al. 2017). The location

of the activity during tongue movement overlapped with the result of the main contrast for

articulation in the syllable production task; as previously, we found spurious activity in the

prefrontal lobe white matter that we presume is movement related.

Vowel production induced activity bilaterally in a central dorsal region, a ventral region anterior

to the central sulcus, and in superior temporal cortex. Within the dorsal region, activation is

found both in the central sulcus and on the precentral gyrus. As for the syllable production

task, activity in superior temporal cortex was presumed due to hearing oneself. In the group

average map, and at the statistical threshold used, the dorsal and ventral larynx

representations appeared to be connected, with residual activity along the ventral central

sulcus.

Basic localizer task - Surface results

Group count maps of the basic localizer task projected to the surface revealed the same

somatotopic activity pattern as seen in the volumetric results for group average activity (Figure

3B). The tongue condition showed highly consistent activation in the mid-portion of the central

sulcus during articulation (maximum overlap was achieved for all 20 subjects). The result for

tongue movement was highly similar to the result of the main contrast for articulation, which

is in line with the volumetric results. Complete overlap of activated vertices was not achieved

for the lip condition, but the activated region was still highly consistent across subjects

(maximum: 15 subjects).

For the vowel production condition, the group count maps show a dorsal and a ventral cluster,

similar to the pattern seen during the main contrast for vocalization (task 1). The dorsal cluster

extended from the central sulcus to the precentral gyrus and in the right hemisphere a distinct

dorsal gyral and a dorsal sulcal activation can be observed. We presumed that this dorsal gyral

activity, but not the sulcal activity, is a residual breathing-related effect, because it overlaps

with activity from the 'breathing only' condition (Figure S1). The ventral activation cluster

appeared to extend into a part of cortex where we expected the tongue representation, and

may reflect residual tongue activity during vowel production. In contrast to the volumetric

results for the group averaged activity, the count maps based on individually thresholded

activation maps show the dorsal and ventral representation activated by vowel production to

be clearly separate.

As described above for the main contrast for vocalization, the values in the group count maps

for vowel production indicate that the location of this activity is less consistent than for the

other conditions; the area activated during vocalization is also smaller, both dorsally and

ventrally, than for the articulators. The highest overlap was achieved in the right dorsal larynx

representation in 11 subjects.

Figure 3 – Basic localizer task. **A**: Whole-brain group activation maps showing areas activated during lip and

tongue movement and during vowel production (vowel-wise threshold $z > 3.5$, $n = 20$). For the vowel

production condition (turquoise), both a dorsal and a ventral representation are shown in separate panels.

See legend to Figure 2 for details. **B**: Surface group count maps of the same conditions. See legend to Figure

2 for details.

Individual surface activation maxima

In order to characterize the variability of brain activity across subjects, we derived individual

activation peaks for the hand movement (based on the hand-localizer, only in the left

hemisphere), lip and tongue movement (based on the basic localizer task) and vocalization

(based on the syllable production task). The main contrast for vocalization was used to localize

the dorsal and a ventral larynx representation. Figure 4A shows the spatial distribution of

individual activation peaks on an inflated brain surface. Overall, the location of the peaks for

the different movement types is highly consistent with similar cross-subject variability. For the

ventral larynx representation, particularly in the left hemisphere, however, the location of the

maxima appears to be more variable.

Within-subject reliability of the activation maxima across the two tasks was compared for

'vocalization' and 'vowel production' for both dorsal and ventral larynx representations as well

as for 'articulation' and 'tongue movement'. Reliability was high with a median distance across

subjects of less than 10 mm for the three ROIs in both hemispheres.

Cortical surface maps of microstructural features

We derived whole-brain average surface maps for different measures related to cortical

microstructure (Figure 4B, C). Overall, the T1w/T2w map and the three MPM parameter maps

(MPM_{MT}, MPM_{R1}, MPM_{R2*}) show a similar pattern of myelin content across the cortex (Figure

4C). In all four maps, the central sulcus as well as precentral and postcentral gyrus are

characteristically high in myelin, which is considered to be a defining feature of primary motor

and sensory cortex (Glasser and van Essen 2011). The location of the ventral boundary of the

somatomotor cortex, indicated by a steep gradient of myelin values, slightly varies across the

four maps, but this boundary is consistently more ventrally located in the right compared with

the left hemisphere.

The four maps that correlate with cortical myelin (T1w/T2w, MPM_{MT}, MPM_{R1} and MPM_{R2*}) are

sensitive to different biophysical properties of the myelin, but it should be noted that their

sensitivity profiles are not completely independent (Callaghan et al. 2016). Therefore, some

dissimilarities regarding the distribution of myelin along the cortex can be observed across the

maps. MPM_{MT} and MPM_{R1} have a stronger signal in the frontal lobe compared to the T1w/T2w

and the MPM_{R2*} map. In addition to myelin, the R2* signal is influenced by cortical properties

such as iron content and calcium (Wu et al. 2009; Bagnato et al. 2018). For the T1w/Tw2 map,

the underlying biophysical model is less well understood. The high R2* values in the ventral

temporal lobe are likely a susceptibility artifact caused by signal loss at the air-tissue boundary.

In the cortical thickness map, a prominent strip of low values (i.e. thinner cortex) can be

observed at the posterior bank of the central sulcus, indicating the location of primary sensory

cortex (Fischl and Dale 2000) (Figure 4B). Cortical thickness values are high (i.e. thicker cortex)

in the anterior bank of the central sulcus and in the precentral gyrus, which is indicating the

location of primary motor cortex. In the cortical thickness map, the ventral boundary of

primary sensory cortex can be determined by a sharp gradient at the level of the subcentral

gyrus. The location of this boundary is comparable with the location described above in the

other maps. The same hemispheric difference can be observed with the somatomotor

boundary being located further ventrally in the right hemisphere.

The whole brain maps demonstrate that these cortical surface measures are informative about

the cortical microstructure underlying our functional activation maxima, which we describe in

the next section. The result of the reconstructed MPM parameter maps *per se* is not a primary

result of the paper.

Figure 4 – **A**: Individual activation maxima derived for hand movement (only left hemisphere), lip movement,

tongue movement and vocalization (i.e. larynx activity during voice production) (n = 20). For vocalization,

activation maxima in two separate brain regions were derived to localize the dorsal and ventral larynx

representations. **B**: Average cortical thickness map (n = 20). **C**: Average T1w/T2w myelin map (n = 20) and

MPM parameter maps (n = 19).

Microstructural properties at activation maxima

Next, we determined the intensity value of the microstructural surface maps at the individual

subjects' activation maxima for the different movement types (Figure 5). Figure 5A shows z-

transformed intensity values for T1w/T2w, MPM_{MT}, MPM_{R1} and MPM_{R2*} maps at hand, lip,

tongue as well as dorsal and ventral larynx maxima, i.e. at the peaks presented in Figure 4A.

Mixed effects analyses with the factors ROI (excluding the hand ROI), hemisphere and map

demonstrated that there were highly significant effects of ROI ($F(3, 600) = 60.69$, $p < 0.001$)

and hemisphere ($F(1, 600) = 43.54$, $p < 0.001$) but no significant effect of map ($F(3, 600) < 1$,

n.s.). The significant effect of ROI reflects that values for the ventral larynx are lower than in

the other ROIs (post-hoc pairwise t-tests with Tukey adjustment, $p < 0.001$), but the values for

dorsal larynx representation, lip and tongue were not significantly different. The main effect of

hemisphere reflects higher values in the right hemisphere ($p < 0.001$). The interaction between

ROI and hemisphere was significant ($F(3, 600) = 3.93$, $p = 0.009$) indicating that the difference

between the ventral larynx representation and the other motor representations was greater

on the right hemisphere than on the left.

Cortical thickness values show a high inter-individual variability for hand, dorsal larynx, tongue

and lip ROIs (Figure 5B). This effect can be attributed to the fact that some maxima are located

in the depth of the central sulcus, where values are low, while others are located on the

anterior bank of the central sulcus, where values are high. Variability in the right hemisphere

is lower, which is consistent with the location of the maxima predominantly on the anterior

bank (Figure 4A). This result potentially indicates that some of derived maximal activation

peaks represent activity evoked by somatosensation rather than by motor control. If maxima

with a cortical thickness level below a cut-off value of 2 mm (Fischl and Dale 2000) are

excluded, the comparison of cortical thickness values shows a marginally nonsignificant effect

of ROI ($F(3, 120) = 2.53$, $p < 0.061$), but not of hemisphere ($F(1, 120) < 1$). The marginal effect

of ROI is driven by the ventral larynx representation having lower cortical thickness values

compared to the other ROIs ($p < 0.068$).

To increase sensitivity in the quantification of myelin values described above, we re-ran the

linear mixed model while excluding the sensory activation maxima as defined by a cortical

thickness below 2 mm. The significant main effects of ROI ($F(3, 480) = 73.62$, $p < 0.001$) and

hemisphere ($F(1, 480) = 20.81$, $p < 0.001$) were stronger than reported above. As reported

above, the main effect of ROI is due to decreased values in the ventral larynx representation

and the hemispheric effect due to increased values in the right hemisphere. The interaction

effect of ROI and hemisphere ($F(3, 480) = 2.26$, $p = 0.08$)) was no longer significant, indicating

that the difference between ventral larynx representation and the other motor

representations did not differ between hemispheres.

To assess effects based on the non-transformed values of the four maps (T1w/T2w, MPM_{MT},

MPM_{R1}, MPM_{R2*}) rather than the z-scores, we also derived a 'dissimilarity matrix' based on the

pair-wise Manhattan distances between ROIs after excluding the sensory activation maxima

(Figure 5C). The values were averaged across hemispheres, because no significant interaction

effect was found. The dissimilarity matrix reflects the effect of the ventral larynx

representation being most dissimilar from the other motor representations based on these

quantitative measures of cortical myelin.

Figure 5 – Measures of cortical microstructure derived at individual activation maxima (*n* = 19). **A**:

Microstructural values that correlate with myelin (T1w/T2w, MPM$_{MT}$, MPM$_{R1}$, MPM$_{R2*}$) derived at activation

maxima (for individual maxima see Figure 4A). **B**: Individual values of cortical thickness. The dashed grey line

indicates the threshold that was chosen to determine the exclusion of activation maxima that are presumed

to be sensory due to their location in cortex < 2 mm thick. **C**: Dissimilarity matrix of the myelin values across

ROIs. The matrix shows a quantitative comparison of the values shown in **A** after excluding sensory activation

maxima. Pair-wise dissimilarity is defined as the Manhattan distance between each pair of ROIs.

A linear discriminant analysis (LDA) revealed that the following equation can be used to

discriminate the ventral larynx representation from the dorsal larynx representation:

$$y = -0.28 \cdot T1w/T2w - 2.82 \cdot MPM_{MT} - 3.03 \cdot MPM_{R1} - 0.47 \cdot MPM_{R2*} - 1.19$$

When inputting the *z*-transformed values of the surface maps at a specific location, a resulting

value of y > 0 indicates that the profile of values is more similar to the ventral larynx

representation than to the dorsal larynx representation. The formula indicates that MPM_{MT}

and MPM_{R1} are the most informative measures to discriminate the two larynx representations

(largest weighting). The mean performance accuracy of the classifier is 0.94 (while 0.5 indicates

chance level performance).

Taken together, these quantifications show that the ventral larynx representation is located in

cortex that has lower myelin content and lower cortical thickness compared to primary motor

cortex, where the other movement representations including the dorsal larynx representation,

are located.

4. Discussion

The goal of this study was to characterize the cortical microstructure underlying the laryngeal

representations in the human brain. We localized brain activity evoked by voluntary control of

laryngeal movements during vocalization using a novel paradigm. We showed that even when

controlling for breathing, vocalization elicits brain activity in two separate parts of the

somatomotor cortex: a dorsal region in the central sulcus and a ventral region close to the

Sylvian fissure. On an individual level, the laryngeal activations and the activations during

movement of hand, lips, and tongue show a consistent somatotopic arrangement.

Characterization of cortical microstructure based on structural and quantitative MRI shows

that the dorsal larynx representation has a similar profile to other movement representations

primary motor cortex, while the ventral larynx representation has lower myelin content and

cortical thickness. These results suggest that the dorsal larynx representation is the primary

locus of laryngeal motor control in primary motor cortex, while the ventral larynx

representation relates more to secondary activations in non-primary motor cortex.

Our experimental task was designed to separate laryngeal activity from that evoked by

supralaryngeal articulation and breathing. We were able to dissociate the supralaryngeal and

laryngeal components of syllable production using a factorial design and orthogonal task

contrasts (Murphy et al. 1997). Covert speech is known to engage similar somatomotor brain

areas to overt speech production (Kleber et al. 2007). For our purpose, however, using a 'covert

speech' condition was essential to allow us to construct a fully factorial model aimed at

isolating activity related to the execution of movements involved in articulation and

vocalization while controlling for other processes such as selection and planning of movement

sequences. The result of a factorial design is statistically more robust than a subtraction design,

because the main contrasts include data from all task conditions improving the mathematical

estimates and associated statistics. The main contrast for vocalization showed some residual

activity of the tongue representation. Using the neutral vowel schwa or glottal stops instead

(Gick 2002; Loucks et al. 2007; Brown et al. 2008; Grabski et al. 2012; Belyk et al. 2018),

however, might have caused pharynx activity that would have been more difficult to dissociate.

In addition to the syllable production task, we used a basic localizer task that involved

movement of the lips, and tongue, and vowel production. Results of the basic localizer task

were consistent with the syllable production task, and within-subject variability across the

comparable contrasts was low. The areas activated for the main contrast of articulation overlap

with the result from the tongue localizer, which is expected given that the syllables produced

mainly rely on tongue movement. The areas activated for the main contrast for vocalization

overlap with the result from the vowel production condition, but some differences could be

observed. Residual tongue activity resulting from producing the vowel /i/ is more apparent in

the vowel production condition, suggesting that this was better controlled for in the factorial

design.

When studying vocalizations, control for breathing is essential given that human vocal speech

sounds are mostly produced during exhalation. Several previous neuroimaging studies,

however, did not control for breathing. Some previous studies used an instructed exhalation

condition for comparison with vocalization (Loucks et al. 2007; Galgano et al. 2019), but a

difference in activation of regions in motor cortex was not found consistently. One likely

explanation is that explicitly instructing subjects to exhale might engage laryngeal muscles in

such a way that no difference to laryngeal activity during vocalization can be observed. In this

study, the breathing pattern and rate were matched for all task conditions. Inhalation and

exhalation were explicitly instructed on a screen and monitored using a breath belt. Although

slight deviations in the shape of the breathing trace can be observed, the overall breathing

traces were comparable across subjects and conditions. The deviations in breathing patterns

observed in the basic localizer task suggest that breathing is less well controlled for compared

with breathing during the syllable production task, where subjects produced the same

utterance in different conditions.

Our results suggest that controversial findings within the neuroimaging literature can largely

be explained by differences in experimental design. Several studies focused on the dorsal

region as location of the 'laryngeal motor cortex' or 'larynx phonation area' (Sörös et al. 2006;

Brown et al. 2008; Kleber et al. 2013; Belyk and Brown 2014; Belyk et al. 2018). Brown et al.

(2008) suggested the presence of a 'dorsolateral' larynx representation on the crown of the

precentral gyrus (x = 50, y = -2, z = 37) and a 'ventromedial' larynx representation in the central

sulcus (x = 44, y = -8, z = 34). Our results suggest, however, that activity in the gyral portion of

the dorsal region is a residual breathing-related signal and not specific for laryngeal activity

during vocalization. When contrasting the 'breathing only' condition, during which breathing

was explicitly instructed, to the resting baseline with normal breathing, we found activity in a

dorsal gyral region (x = 55, y = 0, z = 43), which is presumably related to voluntary control of

the diaphragm during instructed inhalation and exhalation (Figure S1B) (Ramsay et al. 1993;

Olthoff et al. 2008). Similarly, we found the dorsal gyral activation when contrasting the 'covert

speech' condition, which involved instructed breathing, but with no overt movement, to the

resting baseline with normal breathing (Figure S1A). We also find this dorsal gyral activity

during vowel production, but only when we assess it relative to the resting baseline with

normal breathing (Figure S1B). When we assess vowel production relative to the 'breathing

only' activity, only the sulcal portion of the dorsal activity, which we also found in the main

contrast for vocalization, remains (x = 43, y = -13, z = 38). The activation that Brown et al.

(2008) refer to as 'ventromedial' larynx area, overlaps most closely with the region that we

focus in this paper as dorsal (sulcal) larynx representation.

In addition to activation of the dorsal larynx representation, multiple neuroimaging studies

reported activation in a region that we refer to as ventral larynx representation (Terumitsu et

al. 2006; Olthoff et al. 2008; Grabski et al. 2012). Strong evidence for the involvement of a

ventral larynx representation in laryngeal motor control also comes from electrical recording

from the cortical surface during vocalization and microsimulation studies (Galgano and Froud

2008; Bouchard et al. 2013; Chang et al. 2013; Breshears et al. 2015; Dichter et al. 2018).

Electrocorticography demonstrated the presence of both a dorsal and a ventral larynx

representation much more consistently than the fMRI literature (reviewed in Conant et al.,

2014). Direct electrical stimulation of the dorsal larynx representation causes vertical larynx

movement that correlates with stimulation magnitude and evokes vocalization (Dichter et al.

2018); stimulation of the ventral larynx representation causes speech arrest (Chang et al.

2017).

In addition to our novel experimental design, this study focuses on investigating individual

differences in laryngeal activation patterns. The task count maps demonstrate that laryngeal

activity is much less consistent across subjects compared with other movement types. In some

subjects the dorsal or ventral activity did not reach the individual significance threshold. The

location of the ventral larynx representation, in particular in the left hemisphere, show a much

wider spatial spread across the cortex indicating large intra-individual variability. A recent

anatomical study based on the dataset presented here, showed that inter-individual

morphological variability in the ventral end of the central sulcus is higher than in the rest of

the central sulcus (Eichert et al. 2020). It was shown that variability in sulcal morphology can

explain, in part, spatial variability in functional activation peaks. As a result of the variability

and the low *z*-statistical values, group level fMRI analysis in previous studies might have failed

to detect one of the larynx representations or the task paradigm was not optimized to localize

the larynx representations.

The existence of two separate brain regions correlating with laryngeal activity in the central

sulcus raises the question as to whether both of them are involved in motor control of the

larynx. Given that the larynx area in the macaque is located in a ventral premotor area (Hast

et al. 1974; Jürgens 1974; Simonyan and Jürgens 2002; Coudé et al. 2011), an evolutionary

'migration and duplication' hypothesis has been proposed (Belyk and Brown, 2017; Jarvis,

2019; reviewed in Mars et al., 2018). The theory states that the human ventral larynx

representation migrated posteriorly during evolution, possibly into a different

cytoarchitectonic area. The dorsal larynx representation is thought to have evolved as human

novelty in primary motor cortex.

Additional support for a duplication theory also comes from genetic profiling analyses

comparing gene expressions in song-learning birds and humans (Pfenning et al. 2014;

Chakraborty and Jarvis 2015). Gene expression in the avian vocal nuclei, that are involved in

vocal learning, are similar to the expression profiles in both the dorsal and the ventral larynx

representation of the human brain. In the context of this more general brain pathway

duplication theory, it has been recently suggested that there was an additional duplication in

human vocal premotor cortex, leading to the emergence of a pre-larynx motor cortex (preLMC)

just anterior to the ventral larynx representation (Jarvis 2019). The proposed genetic

mechanisms for the evolutionary brain pathway duplication theory in laryngeal motor control,

however, remain to be verified experimentally.

An alternative interpretation of the non-human primate literature would suggest that the

species differences are quantitative rather than qualitative. While traditional electrical

mapping studies failed to find a focal larynx representation in primary motor cortex in non-

human primates (Bailey et al. 1950; Hast et al. 1974; Simonyan and Jürgens 2002), neuronal

activity correlating with laryngeal movement was observed, for example, during swallowing in

macaques (Martin et al. 1997, 1999). Furthermore, tract-tracing studies showed that the

premotor larynx representation in the macaque has direct connections to primary motor

cortex (Künzle 1978; Simonyan and Jürgens 2002). These findings might indicate the presence

of a non-human homology of the dorsal larynx representation, which is experimentally more

challenging to localize due to its smaller spatial extent.

The present study is the first to characterize the microstructural properties of cortex

underlying the dorsal and ventral larynx representation. The use of high-resolution

quantitative neuroimaging allows us to characterize cortical architecture noninvasively

(Weiskopf et al. 2013; Lutti et al. 2014). Measures from T1w/T2w maps and MPM parameter

maps (MPM_{MT}, MPM_{R1}, MPM_{R2*}) are sensitive to different microstructural properties of the

cortex, but all of them have been shown to correlate with myelin to varying degrees (Glasser

and van Essen 2011; Weiskopf et al. 2013; Callaghan et al. 2014, 2015; Bagnato et al. 2018).

Age-related effects can confound measures of cortical microstructure (Gennatas et al. 2017;

Grydeland et al. 2019), but the effects in the regions studied are expected to be minimal.

The cytoarchitectonic area underlying the larynx representations, remains to be determined

histologically. Our quantification, however, suggests strongly that the dorsal larynx

representation is located in primary motor cortex, which is characterized by high myelin

content and thicker cortex (Fischl and Dale 2000; Glasser and van Essen 2011). The ventral

larynx representation has lower myelin content and thinner cortex indicating that it is located

in a different cortical territory. Based on a cytoarchitectonic map of the human brain

(Brodmann 1905), we suggest that the human ventral larynx representation is located in

Brodmann area (BA) 6 (premotor cortex). This interpretation is consistent with the idea that

the primate brain contains multiple laryngeal representations in different cortical areas: The

human ventral larynx representation is homologous to the premotor macaque larynx

representation (Hast et al. 1974; Jürgens 1974; Simonyan and Jürgens 2003; Coudé et al. 2011)

and the human dorsal larynx representation is homologous to a macaque larynx

representation in primary motor cortex, which receives projections from the premotor larynx

representation (Künzle 1978; Simonyan and Jürgens 2002).

Many individual maxima for the ventral larynx representation were located in the subcentral

part of cortex. Brodmann considered this a postcentral and therefore somatosensory cortical

region (BA 43), subjacent and anterior to primary somatosensory cortex (BA 3, 1 and 2)

(Brodmann 1905). A magnetoencephalography study showed that air-puff stimulation of the

larynx evokes activity in a ventral region (Miyaji et al. 2014). Using fMRI, BA 43 was also found

to be activated by movements of the tympanic membrane associated with changes in pressure

in the oropharyngeal cavity such as those that occur during gustation and swallowing and in

vocalization (Job et al. 2011). These results suggest that the ventral larynx activity could be a

sensory rather than a motor representation.

The question regarding the distinct functional contributions of the two larynx representations

during voice productions remains unanswered. Based on our results, we formulate a

hypothesis regarding the causal role of the two laryngeal representations during voice

production: We propose that, as for vocalizations in non-human primates, the ventral larynx

representation is involved in basic control of the vocal folds so that a vocal sound can be

produced. Rapid adduction (tensioning) and abduction (relaxation) of the vocal folds at the

onset and offset of a vocalization is primarily modulated by the intrinsic laryngeal muscles

(Jürgens 1974; Simonyan and Jürgens 2003). Fine motor control, which is required for pitch

modulations in human speech and singing, however, also relies on the vertical movement of

the larynx within the trachea, which is mediated by the extrinsic muscles. We suggest that pitch

control is facilitated by the dorsal larynx representation, which is located in primary motor

cortex (Kleber et al. 2013; Dichter et al. 2018; Finkel et al. 2019). Activity in both intrinsic and

extrinsic laryngeal muscles, however, are tightly coupled and might not be controlled by

distinct brain areas (Belyk and Brown 2014). Our hypothesis is in line with an evolutionary

theory suggesting that the dorsal larynx representation is unique to the human brain (Belyk

and Brown 2017). Such a functional dissociation of dorsal and ventral larynx representation

during vocalization, however, still needs to be tested directly using, for example, non-invasive

brain stimulation.

In sum, we used neuroimaging to localize neural activity related to laryngeal motor control

during vocalization, while controlling for confounding factors such as breathing and

supralaryngeal articulation. We found two activated regions for laryngeal activity during

vocalization, which are in anatomically distinct brain areas. Quantification of cortical

microstructure suggests that the dorsal representation, but not the ventral representation, is

located in primary motor cortex. It remains open, whether and how these two representations

differentially contribute to laryngeal motor control.

<u>Acknowledgements</u>

N.E. is a Wellcome Trust Doctoral student in Neuroscience at the University of Oxford

[203730/Z/16/Z]. The project was supported by the NIHR Oxford Health Biomedical Research

Centre. The Wellcome Centre for Integrative Neuroimaging is supported by core funding from

the Wellcome Trust [203139/Z/16/Z]. The work of R.B.M. is supported by the Biotechnology

and Biological Sciences Research Council (BBSRC) UK [BB/N019814/1] and the Netherlands

Organization for Scientific Research NWO [452-13-015]. The Authors would like to thank

Martina Callaghan from University College London for the MPM sequence.

<u>Code Availability</u>

Upon acceptance of the manuscript, all processing code will be made available from the

Wellcome   Centre   for   Integrative   Neuroimaging's   GitLab   at

git.fmrib.ox.ac.uk/neichert/project_larynx. FSL tools, are available from fsl.fmrib.ox.ac.uk.

Connectome Workbench is available at www.humanconnectome.org/software/connectome-

workbench.html.

Supplementary Material

Surface count maps of all individual task contrasts

1000

1001

1002

Figure S1 – Related to Figure 2 and Figure 3. Surface count maps of all individual task contrasts. **A**: Syllable production task. Upper two rows: Four task conditions - overt speech, silent mouthing (articulation only), vowel production (vocalization only), covert speech (thinking) when compared to the resting baseline with normal breathing. Third row: Two orthogonal main contrasts for the articulation and vocalization component (these are also shown in Figure 2B). **B**: Basic localizer task. Left panel: Task conditions compared to the resting baseline with normal breathing. The 'breathing only' condition is a contrast between the breathing condition (where breathing was instructed) and the resting baseline with normal breathing. A lower threshold of $z = 1.96$ was used to threshold individual z-statistical images for this contrast due to generally lower activation. Right panel: Movement conditions compared to the breathing condition with explicit breathing instruction (these are also shown in Figure 3B).

Task activations in supplementary motor area

The two main contrasts for articulation and vocalization - at a lower voxel-wise threshold of $z > 2$ - revealed activity bilaterally in SMA (Figure S2). A somatotopic arrangement was observed with vocalization activating a more anterior and more dorsal part of cortex. The representations of the effectors during the basic localizer task also show a clear somatotopy: In dorsal-to-ventral direction we first find the representation of the larynx, which is also most anterior, then of the lip and then of the tongue. Only one single representation of the larynx was found in SMA in both the main contrast for vocalization and in the vowel condition.

The location of the representation of the speech effectors is in line with previous accounts in the literature (Picard and Strick 1996). It has been suggested that the vertical line crossing the anterior commissure, the VAC, is a landmark for a division of SMA proper and pre-SMA (Picard

1028   and Strick 1996; Rizzolatti et al. 1996). Our results thus suggest that lip and tongue

1029   representations are located just posterior to VAC in an anterior region of SMA proper.

1030   Laryngeal activity during vocalization, however, activates a part of cortex anterior to VAC,

1031   presumably in preSMA.

1032

1033

1034   *Task activations in cerebellum*

1035

1036   Movement of the articulators and laryngeal activity also evoke activation of the cerebellum in

1037   a somatotopic fashion, which is mirroring the order observed for motor cortex (Figure S2).

1038   Most ventrally, a representation of the larynx is observed, which activates during vowel

1039   production and vocalization. This is followed dorsally by a representation of the lips and the

1040   tongue and then by a second representation of the larynx.

1041

According to a probabilistic atlas of the human cerebellum (Diedrichsen et al. 2009), all the

activations observed are located in the anterior cerebellar lobule VI. This finding is consistent

with previous neuroimaging studies that found activation of the anterior-superior aspect of

cerebellum during speech movements (Petersen et al. 1989; Fiez and Raichle 1997). A previous

resting-state functional connectivity study demonstrated that cerebellar representations

mirror the topography in cerebral motor cortex (Buckner et al. 2011). It is therefore presumed

that the ventral-most representation of the larynx in cerebellum is related to the dorsal larynx

representation in motor cortex and vice versa.

1050

Figure S2 – Related to Figure 2 and Figure 3. Whole-brain group activation maps showing areas activated

during syllable production task (left panels) and basic localizer task (right panels) (vowel-wise threshold $z >$

3.5, n = 20). Coordinates are given for the voxel of maximal activation in the left hemisphere. **A**: Activity on

the medial brain surface centered on the voxel of maximal activity in the left supplementary motor area

(SMA) (note that for the main contrast for vocalization the threshold was lowered to $z >$ 2). **B**: Cerebellar

activity in the same task contrasts and shown in the same colors as in **A**. For the main contrast for

vocalization and for the vowel production contrast, two separate activations are shown in different slices.

1058

1059

1060   *Regions-of-interest to derive maxima during task activation*

1061

1062   An example of the volumetric ROIs in an individual is shown in Figure S3. The central sulcus ROI

1063   used for the hand, lip and tongue was defined using FreeSurfer's automatic volumetric labelling

1064   based on the Destrieux Atlas.

1065

For the larynx, we identified two activation maxima in separate ROIs: One for the dorsal and

one for the ventral larynx representation. The dorsal larynx ROI was a portion of the same

central sulcus ROI used above from z-coordinates in MNI space of 50 - 30. The limits were

determined empirically, so that ROI did not capture the ventral larynx representation or an

unrelated supra-dorsal activation in the trunk area, which was observed in some individuals

(Foerster 1931).

1072

The ventral larynx representation lay outside the central sulcus and was located ventrally in

the subcentral part of cortex. Due to the high intra-individual morphological variability in this

region (Eichert et al. 2020), the ventral larynx ROI was derived manually based on individual

anatomy in surface space. A liberal surface ROI was drawn on each individual's midthickness

surface covering the ventral part of the central sulcus and adjacent gyri (Figure S3A). Anteriorly,

the ROI was delineated by the inferior portion of the precentral sulcus and posteriorly the ROI

spanned the postcentral gyrus. If present, the lateral portion of the ascending sulcus in the

subcentral gyrus was included within the ROI. The dorsal limit of the ROI was defined by a

horizontal plane across the gyrus at the level of the usual location of the posterior ramus of

the inferior precentral sulcus. The ventral larynx surface ROI was converted into a volumetric

ROI covering the underlying cortical ribbon using wb_command. We checked that the ventral

larynx ROI did not overlap with subjacent auditory cortex in the temporal lobe or inferior

frontal cortex.

In some subjects, the main contrast for vocalization in the syllable production task had

additional activity related to articulation of the tongue. To remove this, we transformed the

coordinates for each individual's maximal voxel from the tongue contrast (from the basic

localizer task) to the functional space of the syllable production task (task 1) using rigid-body

transformation and then derived a spherical ROI (7 voxels diameter) around it. This sphere was

used to mask the *z*-statistic image of the main contrast for vocalization prior to localizing the

maxima for laryngeal activity in the dorsal and ventral ROIs described above.

1094

1095

Figure S3 – Related to Methods. **A**: Example of a manually-drawn surface ROI for the ventral larynx

representation. **B**: Volumetric ROIs overlaid onto an individual's functional scan. The central sulcus ROI (red)

was used to derive the voxel of maximal activation for the hand, lip and tongue. The intersection of the

central sulcus ROI and a dorsal mask (dark blue) was used to derive the voxel of maximal activation for the

dorsal larynx representation (black arrow, pink voxel). The ventral mask was projected from surface to

volume space (bright blue) to derive the voxel of maximal activation in the ventral larynx representation

(white arrow, pink voxel). A spherical ROI around the voxel of maximal activation in the tongue contrast was

masked out from the dorsal and ventral mask (yellow).

1104

1105

1106 *Group-level task activation maxima*

1107

1108 Table S1: Group-level task activation maxima. Reported are MNI coordinates of maxima in somatomotor

1109 cortex for the main contrasts reported in the manuscript (LH, RH: left and right hemisphere). For the main

1110 contrast for vocalization and vowel production, a dorsal and a ventral maxima are reported separately.

1111

task	main contrast	LH			RH		
		x	y	z	x	y	z
hand localizer	hand	-38	-21	57			
syllable production	articulation	-50	-12	32	54	-2	32
	vocalization (dorsal)	-40	-16	36	43	-13	38
	vocalization (ventral)	-58	-2	18	60	-2	16
basic localizer	lip	-48	-10	40	50	-8	36
	tongue	-52	-11	28	56	-4	30
	vowel (dorsal)	-42	-16	40	50	-6	40
	vowel (ventral)	-58	-2	20	58	-4	18

1112

1113 **References**

1114

Ackermann H, Hage S, Ziegler W. 2014. Brain mechanisms of acoustic communication in

humans and nonhuman primates: An evolutionary perspective. Behavioral and Brain

Sciences. 37:529–604.

Andersson JLR, Jenkinson M, Smith SM. 2007. Non-linear optimisation FMRIB technial eport

TR07JA1.

Bagnato F, Hametner S, Boyd E, Endmayr V, Shi Y, Ikonomidou V, Chen G, Pawate S,

Lassmann H, Smith S, Brian Welch E. 2018. Untangling the R2* contrast in multiple

sclerosis: A combined MRI-histology study at 7.0 Tesla. PLoS ONE. 13:e0193839.

Bailey P, Bonin G von, McCulloch WS. 1950. The Isocortex of the Chimpanzee. 440.

Barch DM, Burgess GC, Harms MP, Petersen SE, Schlaggar BL, Corbetta M, Glasser MF, Curtiss

S, Dixit S, Feldt C, Nolan D, Bryant E, Hartley T, Footer O, Bjork JM, Poldrack R, Smith S,

Johansen-Berg H, Snyder AZ, Van Essen DC. 2013. Function in the human connectome:

Task-fMRI and individual differences in behavior. NeuroImage. 80:169–189.

Bates DM, Sarkar D. 2007. Bates, Douglas M., and Deepayan Sarkar. "lme4: linear mixed-

effects models using S4 classes. R package version 099875-6.

Belyk M, Brown S. 2014. Somatotopy of the extrinsic laryngeal muscles in the human

sensorimotor cortex. Behavioural Brain Research. 270:364–371.

Belyk M, Brown S. 2017. The origins of the vocal brain in humans. Neuroscience and

Biobehavioral Reviews. 77:177–193.

Belyk M, Lee YS, Brown S. 2018. How does human motor cortex regulate vocal pitch in

singers ? Royal Society Open Science. 5:172208.

Bennett CM, Miller MB. 2010. How reliable are the results from functional magnetic

resonance imaging? Annals of the New York Academy of Sciences.

Bohland JW, Guenther FH. 2006. An fMRI investigation of syllable sequence production.

NeuroImage. 32:821–841.

Bouchard KE, Mesgarani N, Johnson K, Chang EF. 2013. Functional organization of human

sensorimotor cortex for speech articulation. Nature. 495:327–332.

Breshears JD, Molinaro AM, Chang EF. 2015. A probabilistic map of the human ventral

sensorimotor cortex using electrical stimulation. Journal of Neurosurgery. 123:340–349.

Brodmann K. 1905. Beitrage zur histologischen Lokalisation der Grosshirnrinde: III.

Mitteilung. Die Rindenfelder der niederen Affen. J Psychol Neurol. 4:177–206.

Brown S, Laird AR, Pfordresher PQ, Thelen SM, Turkeltaub P, Liotti M. 2009. The somatotopy

of speech: Phonation and articulation in the human motor cortex. Brain and Cognition.

70:31–41.

Brown S, Ngan E, Liotti M. 2008. A larynx area in the human motor cortex. Cerebral Cortex.

18:837–845.

Buckner RL, Krienen FM, Castellanos A, Diaz JC, Yeo BTT, Thomas Yeo BT. 2011. The

organization of the human cerebellum estimated by intrinsic functional connectivity.

Journal of Neurophysiology. 106:2322–2345.

Callaghan MF, Freund P, Draganski B, Anderson E, Cappelletti M, Chowdhury R, Diedrichsen J,

FitzGerald THB, Smittenaar P, Helms G, Lutti A, Weiskopf N. 2014. Widespread age-

related differences in the human brain microstructure revealed by quantitative

magnetic resonance imaging. Neurobiology of Aging. 35:1862–1872.

Callaghan MF, Helms G, Lutti A, Mohammadi S, Weiskopf N. 2015. A general linear

relaxometry model of R1 using imaging data. Magnetic Resonance in Medicine.

73:1309–1314.

Callaghan MF, Mohammadi S, Weiskopf N. 2016. Synthetic quantitative MRI through

relaxometry modelling. NMR in Biomedicine. 29:1729–1738.

Carey D, Krishnan S, Callaghan MF, Sereno MI, Dick F. 2017. Functional and quantitative MRI

mapping of somatomotor representations of human supralaryngeal vocal tract. Cerebral

Cortex. 27:265–278.

Casey BJ, Cannonier T, Conley MI, Cohen AO, Barch DM, Heitzeg MM, Soules ME, Teslovich T,

Dellarco D V., Garavan H, Orr CA, Wager TD, Banich MT, Speer NK, Sutherland MT,

Riedel MC, Dick AS, Bjork JM, Thomas KM, Chaarani B, Mejia MH, Hagler DJ, Daniela

Cornejo M, Sicat CS, Harms MP, Dosenbach NUF, Rosenberg M, Earl E, Bartsch H, Watts

R, Polimeni JR, Kuperman JM, Fair DA, Dale AM. 2018. The Adolescent Brain Cognitive

Development (ABCD) study: Imaging acquisition across 21 sites. Developmental

Cognitive Neuroscience. 32:43–54.

Chakraborty M, Jarvis ED. 2015. Brain evolution by brain pathway duplication. Philosophical

transactions of the Royal Society of London Series B, Biological sciences. 370:20150056-.

Chang EF, Breshears JD, Raygor KP, Lau D, Molinaro AM, Berger MS. 2017. Stereotactic

probability and variability of speech arrest and anomia sites during stimulation mapping

of the language dominant hemisphere. Journal of Neurosurgery. 126:114–121.

Chang EF, Niziolek CA, Knight RT, Nagarajan SS, Houde JF. 2013. Human cortical sensorimotor

network underlying feedback control of vocal pitch. Proceedings of the National

Academy of Sciences. 110:2653–2658.

Conant D, Bouchard KE, Chang EF. 2014. Speech map in the human ventral sensory-motor

cortex. Current Opinion in Neurobiology. 24:63–67.

Coudé G, Ferrari PF, Rodà F, Maranesi M, Borelli E, Veroni V, Monti F, Rozzi S, Fogassi L. 2011.

Neurons controlling voluntary vocalization in the macaque ventral premotor cortex.

PLoS ONE. 6:1–10.

Devlin JT, Matthews PM, Rushworth MFS. 2003. Semantic Processing in the Left Inferior

Prefrontal Cortex: A Combined Functional Magnetic Resonance Imaging and

Transcranial Magnetic Stimulation Study. Journal of Cognitive Neuroscience. 15:71–84.

Dichter BK, Breshears JD, Leonard MK, Chang EF. 2018. The Control of Vocal Pitch in Human

Laryngeal Motor Cortex. Cell. 174:21-31.e9.

Diedrichsen J, Balsters JH, Flavell J, Cussans E, Ramnani N. 2009. A probabilistic MR atlas of

the human cerebellum. NeuroImage. 46:39–46.

Draganski B, Ashburner J, Hutton C, Kherif F, Frackowiak RSJ, Helms G, Weiskopf N. 2011.

Regional specificity of MRI contrast parameter changes in normal ageing revealed by

voxel-based quantification (VBQ). NeuroImage. 55:1423–1434.

Eichert N, Watkins KE, Mars RB, Petrides M. 2020. Morphological and functional variability in

central and subcentral motor cortex of the human brain. bioRxiv. 2020.03.17.995035.

Evans KC, Shea SA, Saykin AJ. 1999. Functional MRI localisation of central nervous system

regions associated with volitional inspiration in humans. Journal of Physiology. 520:383–

392.

Fiez JA, Raichle ME. 1997. Linguistic Processing. In: International Review of Neurobiology. p.

233–254.

Finkel S, Veit R, Lotze M, Friberg A, Vuust P, Soekadar S, Birbaumer N, Kleber B. 2019.

Intermittent theta burst stimulation over right somatosensory larynx cortex enhances

vocal pitch-regulation in nonsingers. Human Brain Mapping. 1–14.

Fischl B. 2012. FreeSurfer. NeuroImage.

Fischl B, Dale AM. 2000. Measuring the thickness of the human cerebral cortex from

magnetic resonance images. Proceedings of the National Academy of Sciences of the

United States of America. 97:11050–11055.

Fitch WT. 2017. Empirical approaches to the study of language evolution. Psychonomic

Bulletin and Review. 24:3–33.

Foerster O. 1931. THE CEREBRAL CORTEX IN MAN. Lancet. 309–312.

Foerster O. 1936. Symptomatologie der Erkrankungen des Grosshirns. Motorische Felder und

Bahnen. Sensible corticale Felder, Handbuch der Neurologie 6. Berlin: Springer.

Galgano J, Froud K. 2008. Evidence of the voice-related cortical potential: An

electroencephalographic study. NeuroImage. 41:1313–1323.

Galgano J, Pantazatos S, Allen K, Yanagihara T, Hirsch J. 2019. Functional connectivity of PAG

with core limbic system and laryngeal cortico-motor structures during human

phonation. Brain Research. 1707:184–189.

Gennatas ED, Avants BB, Wolf DH, Satterthwaite TD, Ruparel K, Ciric R, Hakonarson H, Gur

RE, Gur RC. 2017. Age-related effects and sex differences in gray matter density,

volume, mass, and cortical thickness from childhood to young adulthood. Journal of

Neuroscience. 37:5065–5073.

Gick B. 2002. An X-ray investigation of pharyngeal constriction in American English Schwa.

Phonetica. 59:38–48.

Glasser MF, Goyal MS, Preuss TM, Raichle ME, Van Essen DC. 2014. Trends and properties of

human cerebral cortex: Correlations with cortical myelin content. NeuroImage.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S,

Webster M, Polimeni JR, Van Essen DC, Jenkinson M. 2013. The minimal preprocessing

pipelines for the Human Connectome Project. NeuroImage. 80:105–124.

Glasser MF, van Essen DC. 2011. Mapping human cortical areas in vivo based on myelin

content as revealed by T1- and T2-weighted MRI. Journal of Neuroscience. 31:11597–

11616.

Grabski K, Lamalle L, Vilain C, Schwartz JL, Vallée N, Tropres I, Baciu M, Le Bas JF, Sato M.

2012. Functional MRI assessment of orofacial articulators: Neural correlates of lip, jaw,

larynx, and tongue movements. Human Brain Mapping. 33:2306–2321.

Grydeland H, Vértes PE, Váša F, Romero-Garcia R, Whitaker K, Alexander-Bloch AF, Bjørnerud

A, Patel AX, Sederevičius D, Tamnes CK, Westlye LT, White SR, Walhovd KB, Fjell AM,

Bullmore ET. 2019. Waves of maturation and senescence in micro-structural MRI

markers of human cortical myelination over the lifespan. Cerebral Cortex. 29:1369–

1381.

Hast MH, Fischer JM, Wetzel AB, Thompson VE. 1974. Cortical motor representation of the

laryngeal muscles in Macaca mulatta. Brain Research. 73:229–240.

Jarvis ED. 2019. Evolution of vocal learning and spoken language. Science. 366:50–54.

Jenkinson M. 2003. Fast, automated, N-dimensional phase-unwrapping algorithm. Magnetic

Resonance in Medicine. 49:193–197.

Jenkinson M, Bannister P, Brady M, Smith S. 2002. Improved optimization for the robust and

accurate linear registration and motion correction of brain images. NeuroImage.

17:825–841.

Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. 2012. FSL. NeuroImage.

62:782–790.

Jenkinson M, Smith S. 2001. A global optimisation method for robust affine registration of

brain images. Medical Image Analysis. 5:143–156.

Job A, Paucod JC, O'Beirne GA, Delon-Martin C. 2011. Cortical representation of tympanic

membrane movements due to pressure variation: An fMRI study. Human Brain

Mapping. 32:744–749.

Jürgens U. 1974. On the elicitability of vocalization from the cortical larynx area. Brain

Research. 81:564–566.

Jürgens U. 2002. Neural pathways underlying vocal control. Neuroscience and Biobehavioral

Reviews.

Kleber B, Birbaumer N, Veit R, Trevorrow T, Lotze M. 2007. Overt and imagined singing of an

Italian aria. NeuroImage. 36:889–900.

Kleber B, Zeitouni AG, Friberg A, Zatorre RJ. 2013. Experience-Dependent Modulation of

Feedback Integration during Singing: Role of the Right Anterior Insula. Journal of

Neuroscience. 33:6070–6080.

Kuehn E, Dinse J, Jakobsen E, Long X, Schäfer A, Bazin PL, Villringer A, Sereno MI, Margulies

DS. 2017. Body Topography Parcellates Human Sensory and Motor Cortex. Cerebral

Cortex. 27:3790–3805.

Kumar V, Croxson PL, Simonyan K. 2016. Structural Organization of the Laryngeal Motor

Cortical Network and Its Implication for Evolution of Speech Production. Journal of

Neuroscience. 36:4170–4181.

Künzle H. 1978. Cortico-cortical efferents of primary motor and somatosensory regions of the

cerebral cortex in Macaca fascicularis. Neuroscience. 3:25–39.

Leyton ASF, Sherrington CS. 1917. Observations on the Excitable Cortex of the Chimpanzee,

Orang-Utan, and Gorilla. Quarterly Journal of Experimental Physiology. 11:135–222.

Loucks TMJ, Poletto CJ, Simonyan K, Reynolds CL, Ludlow CL. 2007. Human brain activation

during phonation and exhalation: Common volitional control for two upper airway

functions. NeuroImage. 36:131–143.

Ludlow CL. 2005. Central nervous system control of the laryngeal muscles in humans.

Respiratory Physiology and Neurobiology.

Lutti A, Dick F, Sereno MI, Weiskopf N. 2014. Using high-resolution quantitative mapping of

R1 as an index of cortical myelination. NeuroImage. 93:176–188.

Lutti A, Hutton C, Finsterbusch J, Helms G, Weiskopf N. 2010. Optimization and validation of

methods for mapping of the radiofrequency transmit field at 3T. Magnetic Resonance in

Medicine. 64:229–238.

Lutti A, Stadler J, Josephs O, Windischberger C, Speck O, Bernarding J, Hutton C, Weiskopf N.

2012. Robust and fast whole brain mapping of the RF transmit field B1 at 7T. PLoS ONE.

7:1–7.

Mars RB, Eichert N, Jbabdi S, Verhagen L, Rushworth MFS. 2018. Connectivity and the search

for specializations in the language-capable brain. Current Opinion in Behavioral

Sciences. 21:19–26.

Martin RE, Kemppainen P, Masuda Y, Yao D, Murray GM, Sessle BJ. 1999. Features of

cortically evoked swallowing in the awake primate (Macaca fascicularis). Journal of

Neurophysiology. 82:1529–1541.

Martin RE, Murray GM, Kemppainen P, Masuda Y, Sessle BJ. 1997. Functional properties of

neurons in the primate tongue primary motor cortex during swallowing. Journal of

Neurophysiology. 78:1516–1530.

McKay LC, Evans KC, Frackowiak RSJ, Corfield DR. 2003. Neural correlates of voluntary

breathing in humans. Journal of Applied Physiology. 95:1170–1178.

Miyaji H, Hironaga N, Umezaki T, Hagiwara K, Shigeto H, Sawatsubashi M, Tobimatsu S,

Komune S. 2014. Neuromagnetic detection of the laryngeal area: Sensory-evoked fields

to air-puff stimulation. NeuroImage. 88:162–169.

Murphy K, Corfield DR, Guz A, Fink GR, Wise RJ, Harrison J, Adams L. 1997. Cerebral areas

associated with motor control of speech in humans. Journal of applied physiology

(Bethesda, Md : 1985). 83:1438–1447.

Nieto-Castañón A, Fedorenko E. 2012. Subject-specific functional localizers increase

sensitivity and functional resolution of multi-subject analyses. NeuroImage. 63:1646–

1669.

Olthoff A, Baudewig J, Kruse E, Dechent P. 2008. Cortical Sensorimotor Control in

Vocalization: A Functional Magnetic Resonance Imaging Study. The Laryngoscope.

118:2091–2096.

Penfield W, Boldrey E. 1937. Somatic motor and sensory representation in the cerebral

cortex of man as studied by electrical stimulation. Brain. 60:389–443.

Penfield W, Rasmussen T. 1950. The Cerebral Cortex of Man: Clinical Study of Localization of

Function. Academic Medicine. 25:375.

Petersen SE, Fox PT, Posner MI, Mintun M, Raichle ME. 1989. Positron emission tomographic

studies of the processing of single words. Journal of Cognitive Neuroscience. 1:153–170.

Pfenning AR, Hara E, Whitney O, Rivas M V, Wang R, Roulhac PL, Howard JT, Wirthlin M,

Lovell P V, Ganapathy G, Mountcastle J, Moseley MA, Thompson JW, Soderblom EJ, Iriki

A, Kato M, Gilbert MTP, Zhang G, Bakken T, Bongaarts A, Bernard A, Lein E, Mello C V,

Hartemink AJ, Jarvis ED. 2014. Convergent transcriptional specializations in the brains of

humans and song-learning birds. Science. 346:1256846–1256846.

Picard N, Strick PL. 1996. Motor areas of the medial wall: A review of their location and

functional activation, Cerebral Cortex. Narnia.

Preibisch C, Deichmann R. 2009. Influence of RF spoiling on the stability and accuracy of T1

mapping based on spoiled FLASH with varying flip angles. Magnetic Resonance in

Medicine. 61:125–135.

Ramsay SC, Adams L, Murphy K, Corfield DR, Grootoonk S, Bailey DL, Frackowiak RS, Guz A.

1993. Regional cerebral blood flow during volitional expiration in man: A comparison

with volitional inspiration. The Journal of Physiology. 461:85–101.

Rizzolatti G, Luppino G, Matelli M. 1996. The classic supplementary motor area is formed by

two independent areas. Advances in neurology. 70:45–56.

Simonyan K. 2014. The laryngeal motor cortex: Its organization and connectivity. Current

Opinion in Neurobiology. 28:15–21.

Simonyan K, Jürgens U. 2002. Cortico-cortical projections of the motorcortical larynx area in

the rhesus monkey. Brain Research. 949:23–31.

Simonyan K, Jürgens U. 2003. Efferent subcortical projections of the laryngeal motorcortex in

the rhesus monkey. Brain Research. 974:43–59.

Simonyan K, Ostuni J, Ludlow CL, Horwitz B. 2009. Functional But Not Structural Networks of

the Human Laryngeal Motor Cortex Show Left Hemispheric Lateralization during Syllable

But Not Breathing Production. Journal of Neuroscience. 29:14912–14923.

Smith SM. 2002. Fast robust automated brain extraction. Human Brain Mapping. 17:143–

155.

Sörös P, Sokoloff LG, Bose A, McIntosh AR, Graham SJ, Stuss DT. 2006. Clustered functional

MRI of overt speech production. NeuroImage. 32:376–387.

Tabelow K, Balteau E, Ashburner J, Callaghan MF, Draganski B, Helms G, Kherif F, Leutritz T,

Lutti A, Phillips C, Reimer E, Ruthotto L, Seif M, Weiskopf N, Ziegler G, Mohammadi S.

2019. hMRI – A toolbox for quantitative MRI in neuroscience and clinical research.

NeuroImage. 194:191–210.

Terumitsu M, Fujii Y, Suzuki K, Kwee IL, Nakada T. 2006. Human primary motor cortex shows

hemispheric specialization for speech. Neuroreport. 17:1091–1095.

Toyoda G, Brown EC, Matsuzaki N, Kojima K, Nishida M, Asano E. 2014. Electrocorticographic

1353  correlates of overt articulation of 44 English phonemes: Intracranial recording in

1354  children with focal epilepsy. Clinical Neurophysiology. 125:1129–1137.

1355 Weiskopf N, Callaghan MF, Josephs O, Lutti A, Mohammadi S. 2014. Estimating the apparent

1356  transverse relaxation time (R2*) from images with different contrasts (ESTATICS)

1357  reduces motion artifacts. Frontiers in Neuroscience. 8.

1358 Weiskopf N, Suckling J, Williams G, Correia M. MM, Inkster B, Tait R, Ooi C, Bullmore T. ET,

1359  Lutti A. 2013. Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: A

1360  multi-center validation. Frontiers in Neuroscience. 7:95.

1361 Woo CW, Krishnan A, Wager TD. 2014. Cluster-extent based thresholding in fMRI analyses:

1362  Pitfalls and recommendations. NeuroImage. 91:412–419.

1363 Woolrich MW, Behrens TEJ, Beckmann CF, Jenkinson M, Smith SM. 2004. Multilevel linear

1364  modelling for FMRI group analysis using Bayesian inference. NeuroImage. 21:1732–

1365  1747.

1366 Woolrich MW, Ripley BD, Brady M, Smith SM. 2001. Temporal autocorrelation in univariate

1367  linear modeling of FMRI data. NeuroImage. 14:1370–1386.

1368 Wu Z, Mittal S, Kish K, Yu Y, Hu J, Haacke EM. 2009. Identification of calcification with MRI

1369  using susceptibility-weighted imaging: A case study. Journal of Magnetic Resonance

1370  Imaging. 29:177–182.

1371