Molecular epidemiology of third-generation cephalosporin-resistant Enterobacteriaceae

from Southeast Queensland, Australia

Stewart AG[1,2], Price EP[3,4], Schabacker K[3,4], Birikmen M[3,4], Harris PNA[2,5], Choong K[1], Subedi S[1,6]

and Sarovich DS[3,4#]

[1]Department of Infectious Diseases, Sunshine Coast University Hospital, Birtinya, QLD,

Australia

[2]Centre for Clinical Research, University of Queensland, Herston, QLD, Australia

[3]GeneCology Research Centre, University of the Sunshine Coast, Sippy Downs, QLD, Australia

[4]Sunshine Coast Health Institute, Birtinya, QLD, Australia

[5]Pathology Queensland, Central Microbiology, Royal Brisbane and Women's Hospital,

Herston, QLD, Australia

[6]Department of Microbiology, Pathology Queensland, Sunshine Coast University Hospital,

Birtinya, QLD Australia

Keywords: Antibiotic, antimicrobial, resistance, cephalosporin, ESBL, whole-genome

sequencing, real-time PCR, Enterobacteriaceae

Running title: Population genomics of 3GC-R Enterobacteriaceae

#Corresponding author

Derek S. Sarovich

dsarovich@usc.edu.au

+61754563748

University of the Sunshine Coast, Locked Bag 4, Maroochydore DC, QLD, 4558, Australia

Synopsis

Third-generation cephalosporin-resistant (3GC-R) Enterobacteriaceae represent a major threat to human health. Here, we captured 288 3GC-R Enterobacteriaceae clinical isolates from 264 patients presenting at a regional Australian hospital over a 14-month period. Alongside routine mass spectrometry speciation and antibiotic sensitivity testing, isolates were examined using rapid (~40 min) real-time PCR assays targeting the most common extended spectrum β-lactamases (ESBLs; CTX-M-1 and CTX-M-9 groups, plus TEM, SHV, and an internal 16S ribosomal DNA control). AmpC CMY β-lactamase prevalence was also examined. *Escherichia coli* (80.2%) and *Klebsiella pneumoniae* (17.0%) were dominant, with *Klebsiella oxytoca*, *Klebsiella aerogenes* and *Enterobacter cloacae* infrequently identified. Ceftriaxone and cefoxitin resistance were identified in 97.0% and 24.5% of *E. coli* and *K. pneumoniae* isolates, respectively. Consistent with global findings in Enterobacteriaceae, most (98.3%) isolates harbored at least one β-lactamase gene, with 144 (50%) encoding $bla_{CTX-M-1}$ group, 92 (31.9%) $bla_{CTX-M-9}$ group, 48 (16.7%) bla_{SHV}, 133 (46.2%) bla_{TEM}, and 34 (11.8%) bla_{CMY}. A subset of isolates (n=98) were subjected to whole-genome sequencing (WGS) to identify the presence of cryptic resistance determinants, and to verify genotyping accuracy. WGS of β-lactamase negative or carbapenem-resistant isolates identified uncommon ESBLs and carbapenemases, including bla_{NDM} and bla_{IMP}, and confirmed all PCR-positive genotypes. We demonstrate that our PCR assays enable the rapid and cost-effective identification of ESBLs in the hospital setting, which has important infection control and therapeutic implications.

Introduction

Third-generation cephalosporin-resistant (3GC-R) Enterobacteriaceae represent a significant threat to human health due to their ability to rapidly transmit antimicrobial resistance (AMR) determinants within and among bacterial populations (1). Both infection and colonization with these globally-disseminated organisms are associated with poor clinical outcomes and death (2). 3GC-R Enterobacteriaceae prevalence rates vary considerably among hospitals and countries, with particularly high rates in India, Asia, and the Middle East (3). Australia has historically observed a relatively low frequency of 3GC-R Enterobacteriaceae. However, this pattern may be changing, with an extended spectrum β-lactamase (ESBL) phenotype detected in 13.3% of *E. coli* and 9.8% of *K. pneumoniae* in blood culture isolates in 2018 (4), up from 2013 rates of 7.6% and 6.3%, respectively. Moreover, ceftriaxone resistance was seen in 13.4% of *E. coli* and 9.4% of *K. pneumoniae*, with 86.3% and 82.6% containing CTX-M-type ESBL genes, respectively (4).

AMR in 3GC-R Enterobacteriaceae is generally conferred by the presence of an ESBL or the over-expression of a chromosomal or plasmid-borne AmpC β-lactamase (5). CTX-M-type ESBLs are currently the most common AMR determinants found in 3GC-R Enterobacteriaceae isolated in Australian hospitals, with $bla_{CTX-M-1}$ and $bla_{CTX-M-9}$ groups collectively comprising 86.2% of identified CTX-M-type enzymes (4). Detection of ESBL subtypes is not standard clinical practice; instead, their presence is typically inferred from antimicrobial susceptibility patterns and phenotypic confirmatory tests, which are laborious, costly (~$AUD25/isolate), and time-consuming (>24h) to perform (6). Use of molecular-based diagnostics for AMR determinant detection in the clinical setting would improve patient outcomes by enabling more rapid, directed, and personalized treatment strategies, thereby improving antibiotic stewardship measures and reducing AMR burden. Their utility

has already been demonstrated in a number of clinical settings, including for bloodstream

infections, pneumonia, and infection control screening (7-9).

In the current study, we prospectively collected all 3GC-R Enterobacteriaceae identified over

a 14-month period at the Sunshine Coast University Hospital, Queensland, Australia, a 450-

bed, regional teaching and tertiary hospital. Using large-scale comparative genomics, we

developed a novel quadriplex and two singleplex real-time PCR assays to rapidly detect the

most commonly identified β-lactamase (including ESBL) genes in 3GC-R Enterobacteriaceae.

In association with WGS, we used our diagnostic real-time assays to determine the

molecular epidemiology of the β-lactamase genes in our isolate set, and compared their

performance with both genotypic and phenotypic typing methods to validate assay

accuracy, sensitivity and specificity. Finally, we examined the population structure of our

3GC-R Enterobacteriaceae population to identify potential cases of nosocomial spread or

outbreak clusters.

Materials and Methods

Ethics. This study was approved by The Prince Charles Hospital Human Resources Ethics

Committee (HREC/18/QPCH/110). A waiver of informed consent was granted due to the low

to negligible risk associated with this study.

Isolate collection. Clinical specimens from 264 adult and pediatric patients (including ICU

patients [n=48]) that were culture-positive for 3GC-R Enterobacteriaceae were collected

prospectively from July 2017 through September 2018 at the Sunshine Coast University

Hospital microbiology laboratory (n=288). Multiple isolates were collected from 23 patients,

either due to re-infection over the course of the study (n=9), multiple samples taken during

initial admission (n=11), or collected during the course of their hospital stay (n=3). Samples

were inoculated onto ChromID^{TM} ESBL agar (bioMérieux) and ChromID^{TM} Carba/OXA agar

(bioMérieux) and examined for colored colonies at 18-24 hours. An ESBL screening disc test

was performed on isolates growing on ESBL agar. Isolates were speciated using the VITEK® 2

MALDI-TOF MS with the GN ID card (bioMérieux, Murrarie, QLD, Australia). Isolate

information is detailed in Table S1.

Susceptibility testing. Antibiotic susceptibility testing and minimum inhibitory concentration

(MIC) values were determined using VITEK® 2 or Etest (bioMérieux) according to the

manufacturer's instructions. EUCAST criteria (v.10.0) were used for susceptibility and

resistance interpretation.

Culture DNA preparation. For PCR, crude DNA was extracted from all isolates using 5%

chelex-100 resin (Bio-Rad Laboratories, Gladesville, NSW, Australia). For a subset of samples

(*n*=50), a pinhead-sized amount of culture collected using a sterile 10 µL pipette tip was

placed directly into 4 µL of PCR mastermix to facilitate rapid turnaround time. For WGS, DNA

was extracted using the Quick-DNA miniprep kit (Zymo Research, Irvine, CA, USA).

PCR development and β-lactamase detection. A quadriplex real-time PCR assay was

developed to detect the two major CTX-M ESBL groups ($bla_{CTX-M-1}$ and $bla_{CTX-M-9}$) and bla_{SHV}

(both ESBL and non-ESBL); a 16S rDNA probe-based assay (10) was included in this

quadriplex to verify DNA integrity. Additionally, two singleplex assays were designed to

amplify ESBL and non-ESBL bla_{TEM} and bla_{CMY} β-lactamase genes (Figure 1). To identify

conserved regions for PCR assay design, all publicly available genetic sequences for $bla_{CTX-M-1}$,

$bla_{CTX-M-9}$, bla_{CMY}, bla_{SHV}, and bla_{TEM} were downloaded from the NCBI Nucleotide database (as

of March 2019) and aligned using Clustal Omega v1.2.2. Aligned sequences were visualized

in Geneious v11.0.4. Candidate primers and probes were identified in Primer Express v3.0

(Applied Biosystems, Foster City, California, USA) or Primer3 v0.4.0 (11) followed by primer

dimer and heterodimer assessment using NetPrimer

(http://www.premierbiosoft.com/netprimer/).

Black Hole Quencher probes were used to enable multiplexing in a single PCR well and

robust target identification. For the quadriplex assay, the following primers and probes (5' to

3'; Macrogen Inc., Geumcheon-gu, Seoul, Republic of Korea) were designed: CTX-1 (CTX1_F:

CGAMACGTTCCGTCTCGAC, CTX1_R: CTTTCATCCAYGTCACCAGCTG, CTX1_P1: FAM-

CGTGATACCACTTCACCTCGGG-BHQ1), CTX-9 (CTX9_F: CGTCSCGCTCATCGATACC, CTX9_R:

CAGCTGCTTTTGCGTTTCACTC, CTX9_P1: HEX-CGCTTTCCAATGTGCAGTACC-BHQ1), and SHV

(SHV_F: CCARCGTCTGAGCGCC, SHV_R: TATCGGCGATAAACCAGCCC, SHV_P1: Cy5-

CAGCACGGAGCGGATCA-BHQ2). The previously described 16S assay (10) incorporated a

different fluorophore on the probe (Texas Red-CACGAGCTGACGACARCCATGCA-BHQ2) to

accommodate the quadriplex assay format. Two singleplex assays targeting TEM and CMY

were also designed: TEM (TEM_F: TGTGGYGCGGTATTATCCCGT, TEM_R:

TGCATAATTCTCTTACTGTCAWGCCA, TEM_P1: Cy5.5-CACCAGTCACAGAAAAGCATCT-BHQ2),

CMY (CMY_F: TGGCGTATTGGYGATATGTA, CMY_R: TTATGCACCCATGAGGCTTT, CMY_Pr:

FAM-TGGGAGATGCTGAACTGGCC-BHQ1). Coincidentally, the reverse primer sequence for

our CMY assay was identical to one designed previously (12).

The quadriplex PCR was carried out using 0.1 µM 16S probe and primers, 0.2 µM CTX-1 and

CTX-9 probes and primers, 0.3 µM SHV primers, and 0.4 µM SHV probe. For the TEM

singleplex, 0.2 µM TEM primers and probe were used, and for the CMY singleplex, 0.2 uM

primers and 0.4 probe were used. All reactions contained 1X Sso Advanced Universal Probes

Supermix (Bio-Rad), primers, probe/s, 1 µL DNA template, and molecular grade H_2O, to a 5

µL total volume. PCRs were performed on a CFX96 Touch Real-Time PCR Detection System

using white Hard Shell 96-well PCR plates and Microseal 'C' Adhesive seals (Bio-Rad) or 0.2

mL white 8-tube PCR strips and optical caps (Bio-Rad). For bacterial cultures, thermocycling

conditions comprised an initial 2 min denaturation at 95 °C, followed by 45 cycles of 95 °C

for 1 sec and 60 °C for 3 sec (total run time=39 min).

LoD and LoQ determination. Three previously characterized strains (13) that represent the

PCR assays developed in this study – MER85 (*K. pneumoniae*), MER110 (*E. coli*), and MER132

(*E. coli*) – were subject to total DNA extraction using the Gram-positive bacteria extraction

protocol of the Qiagen DNeasy kit (Qiagen, Chadstone Centre, VIC, Australia). DNA quantities

were measured on the Qubit 4 fluorimeter (Thermo Fisher Scientific, Seventeen Mile Rocks,

Qld, Australia) using the Qubit dsDNA BR kit, and normalized to 40 ng/µL in molecular grade

H_2O. LoD and LoQ values were determined across 1:10 serial dilutions (range: 40 ng to 0.4

fg) using MER85 (*bla*$_{TEM}$ singleplex), MER110 (*bla*$_{CMY}$ singleplex), and a 50:50 mix of MER85

($bla_{CTX-M-1}$ group, bla_{SHV}) and MER132 ($bla_{CTX-M-9}$ group) for the quadriplex assay. For PCR

thermocycling, denaturation and annealing times were increased to 5 sec and 10 sec,

respectively, to ensure assay robustness under very low DNA input conditions. Genome

equivalents were calculated by assuming *E. coli* and *K. pneumoniae* genome sizes of 4.984

Mbp and 5.638 Mbp, respectively.

Assay specificity. The quadriplex and singleplex assay amplicons were first assessed for

specificity *in silico* using the microbial nucleotide discontiguous BLAST tool and the Complete

genomes database (http://blast.ncbi.nlm.nih.gov/; performed March 2019). Next, assay

specificity was examined by laboratory testing against a range of both pathogenic and non-

pathogenic microbial species found humans. This diversity panel comprised *Achromobacter*

spp. (*n*=4), *Bordetella bronchiseptica* (*n*=1), *Candida albicans* (*n*=1), *Cupriavidus* sp. (*n*=2),

Enterobacter aerogenes (*n*=1), *Enterobacter cloacae* (*n*=3), *Enterococcus faecalis* (*n*=1),

Klebsiella pneumoniae (*n*=2), *Klebsiella oxytoca* (*n*=1), *Lactobacillus* spp. (*n*=2), *Prevotella*

spp. (*n*=6), *Pseudomonas aeruginosa* (*n*=10), *Staphylococcus aureus* (*n*=4), *Staphylococcus*

epidermidis (*n*=1), *Stenotrophomonas maltophilia* (*n*=5), and *Veillonella* spp. (*n*=4).

Whole-genome sequencing (WGS). WGS was performed at the Australian Centre of

Ecogenomics (University of Queensland, St Lucia, QLD, Australia) on the NextSeq 500

(Illumina, San Diego, CA, USA) with Nextera DNA Flex libraries (Illumina).

Genomic analysis. All genomic data were lightly quality-filtered using Trimmomatic v0.39

(14) using parameters described elsewhere (15). WGS data were screened for multiple

species using Kraken 2 v2.0.8 (16). Single species were assembled using MGAP v1.1

(https://github.com/dsarov/MGAP---Microbial-Genome-Assembler-Pipeline), whereas WGS

data containing multiple species were assembled with metaSPAdes (17). To determine

plasmid copy number and identity, plasmids were assembled using the plasmid module of

SPAdes(18) with plasmid homology determined using BLAST(19). Assemblies were binned

with MaxBin v2.0 (20) and speciated with Kraken 2 v2.0.8 (16). Antibiotic resistance genes

were identified using the Resistance Gene Identifier (21). Phylogenomic analysis was

performed using SPANDx v3.2.1 (22). Samples identified as containing multiple strains

according to the method described by Aziz and colleagues (23) were excluded from

phylogenetic analysis (n=7); samples able to be split into single species using MaxBin were

included in phylogenetic analysis (n=4). *E. coli* O157:H7 Sakai (NCBI Assembly ID: ASM886v2)

(24) or *K. pneumoniae* HS11286 (NCBI Assembly ID: ASM24018v2) (25) were used as read

mapping reference genomes for species-specific analyses. To achieve the highest possible

resolution, error-corrected assemblies of strains with matching multilocus sequence types

(STs) were used as reference genomes for transmission analyses. ST profiles were

determined using SRST2 v0.2.0 (26). Phylogenetic trees were reconstructed using PAUP

v4.4.7 (27) and visualized in iTOL v5.5(28).

WGS data. Raw sequence reads generated in this study are available on the NCBI Sequence

Read Archive database under BioProject accession number PRJNA606985.

Results and Discussion

ESBL enzymes have established themselves as major contributors of AMR in Gram-negative bacteria and are increasing in prevalence, particularly in the community setting. Over 14 months, we collected a total of 288 isolates from all diagnosed cases of 3GC-R Enterobacteriaceae in a regional hospital in Southeast Queensland, Australia. Our patient cohort was predominantly female (64.0%) and aged over 65 (60.9%). ICU patient specimens comprised 54 (18.8%) of the total collection, and were obtained from 48 ICU patients over a 12-month period. Five ICU patients had multiple isolates collected over the course of their stay in the ICU. Three patients isolated single species with identical AMR profiles, two patients isolated multiple species: one with identical AMR profiles and one with differing profiles (Table S1). *E. coli* and *K. pneumoniae* were dominant, comprising 80.2% and 17.0% 3GC-R strains, respectively (Table 1 and S1). The remaining isolates were identified as *E. cloacae* (1.4%; *n*=4), *K. oxytoca* (1.0%; *n*=3), and *K. aerogenes* (0.3%; *n*=1). Resistance to ceftriaxone, ceftazidime or cefepime was common and observed in 96.2%, 93.4%, and 86.8% of isolates, respectively. In contrast, cefoxitin resistance was noted in only 81 (28.1%) isolates (Table 1). Meropenem resistance was rare, with only six resistant strains (2.1%) identified. The double disc diffusion method for ESBL phenotypic testing was negative in 32 (11.1%) isolates, indicating a lack of β-lactamase inhibition in the presence clavulanic acid. The remaining 256 (88.9%) isolates were positive for this phenotypic test.

ESBL and β-lactamase assay performance. Preliminary validation of our singleplex (bla_{TEM}, bla_{CMY}) and quadriplex ($bla_{CTX-M-1}$ group, $bla_{CTX-M-9}$ group, bla_{SHV}, 16S rDNA) PCR assays was performed to determine their accuracy and specificity. Following *in silico* verification, assays were tested on 12 previously characterized control strains (Table S2) (29) that represented a range of Enterobacteriaceae containing the gene targets of interest. All expected targets were identified in this control dataset; no false positives or false negatives were detected,

although both the bla_{TEM} and 16S rDNA assays yielded late (>30 cycles) amplification in no-

template control wells (Figure 1). This result was expected due to residual *E. coli* DNA

presence in the recombinant Sso Advanced *Taq* polymerase, which encodes both bla_{TEM} and

16S rRNA genes.

We found that all genotypes could be accurately determined for the control strains by

directly inoculating PCR wells with a very small amount of bacterial culture, thereby

obviating the need for DNA extraction and leading to faster time-to-diagnosis. However, this

technique requires skill to ensure that the PCR is not inhibited by excess DNA and cellular

constituents, and can lead to greater PCR failures if not performed correctly. In addition, we

recommend using 10 μL rather than 5 μL total PCR volume as, in our hands, the lower

reaction volume led to a greater failure rate. Inclusion of the 16S rDNA assay enables true

negatives to be differentiated from PCR failures. This method provides an attractive option

where rapid diagnosis is paramount, or where DNA extraction is not feasible.

Next, we screened our PCR assays across a diversity panel of non-Enterobacteriaceae human

pathogens or colonizers (*n*=48 strains). Our assays showed 100% sensitivity and specificity

for all targets, with no amplification detected in these isolates, except for the 16S target in

bacterial strains; as with the control dataset, the bla_{TEM} PCR was inhibited by the presence of

non-bla_{TEM}-encoding DNA and therefore did not amplify in any diversity panel strains.

LoQ and LoD determination of the ESBL and β-lactamase assays. Using serial dilutions of

previously characterized DNA samples ranging from 40 ng to 0.4 fg, the LoQ values for each

assay were identified (Figure 2) as follows: bla_{CMY}, 400 fg (~73 genome equivalents [GEs]);

bla_{TEM}, 400 fg (~65 GEs); 16S rDNA, 40 fg (~3 GEs); $bla_{CTX-M-1}$ group, 4 pg (~344 GEs); $bla_{CTX-M-9}$

group, 4 pg (~344 GEs); and bla_{SHV}, 4 pg (~344 GEs). LoD values for each assay were: bla_{CMY}, 4

fg (~7 GEs); $bla_{CTX-M-1}$ group, 4 pg (~344 GEs); $bla_{CTX-M-9}$ group, 400 fg (~34 GEs); and bla_{SHV},

400 fg (~34 GEs). Due to low levels of recombinant *E. coli* DNA present in the Sso Advanced

Taq polymerase, which results in late amplification in no-template control wells, LoD values

for the bla_{TEM} and 16S rDNA assays could not be determined. As expected, these results

indicate inferior performance of the quadriplex assay when compared with the singleplex

assays. Although not examined in this study, increased LoQ and LoD values may be achieved

for the $bla_{CTX-M-1}$ group, $bla_{CTX-M-9}$ group, and bla_{SHV} assays by omitting the 16 rDNA assay, or

by running these assays in singleplex format. Although we have defined the limits of our

assays using pure bacterial templates, it remains to be determined what detection limits

would be achieved from clinical specimens.

ESBL and β-lactamase prevalence in clinical 3GC-R Enterobacteriaceae isolates. Following

specificity confirmation, the PCR assays were tested across our clinical set (*n*=288). All assays

performed well, with 100% concordance between PCR genotypes and WGS of 98 isolates,

and with no false negatives identified (Table S3). Overall, a β-lactamase genotype was

identified in 277 (96.2%) isolates. Consistent with previously reported rates (29), $bla_{CTX-M-1}$

group was most common, being identified in 144/288 (50%) isolates, including 105/231

(45.5%) *E. coli* and 36/49 (73.5%) *K. pneumoniae*. The $bla_{CTX-M-9}$ group was next frequent,

being found in 93/288 (32.3%) isolates, including 89/231 (38.5%) *E. coli* and 4/49 (8.2%) *K.*

pneumoniae. The presence of both $bla_{CTX-M-1}$ group and $bla_{CTX-M-9}$ group enzymes was

identified in five (1.7%) *E. coli* and one (2.0%) *K. pneumoniae* isolate. Of the eight isolates

belonging to other species, $bla_{CTX-M-1}$ and $bla_{CTX-M-9}$ group genes were present in 4/8 (50%)

and 0/8 (0%) of these isolates, respectively. The acquired *ampC* gene, bla_{CMY}, was identified

in 30/231 (13%) of *E. coli*, 3/49 (6.1%) of *K. pneumoniae*, and 1/8 (13%) other

Enterobacteriaceae species. bla_{CMY} and $bla_{CTX-M-1}$ were both identified in seven isolates (five

E. coli, one *K. pneumoniae*, and one *E. cloacae*) and bla_{CMY} and $bla_{CTX-M-9}$ were present in two

E. coli isolates only (Table 1).

Seven strains (five *E. coli* and two *K. pneumoniae*) had bla_{SHV} only; nine *E. coli* had bla_{TEM}

only, and seven (five *K. pneumoniae*, one *E. coli*, and one *E. cloacae*) had both bla_{SHV} and

bla_{TEM}. Although most SHV (e.g. bla_{SHV-1}) and TEM (e.g. bla_{TEM-1}, bla_{TEM-2}) enzymes do not

typically cause 3GC-R, their presence in strains that lack CTX-M-type ESBLs should be

investigated, as these enzymes can occasionally confer 3GC-R through enzyme active site

mutation or gene overexpression (5). Of the 14 strains encoding bla_{SHV} but not $bla_{CTX-M-1}$

group, $bla_{CTX-M-9}$ group, or bla_{CMY} enzymes, five had accompanying WGS data, which

permitted assessment of bla_{SHV} ESBL status. In all cases, the bla_{SHV} variant was an ESBL (two

bla_{SHV-2}, one bla_{SHV-64}, one bla_{SHV-66}, and one $bla_{SHV-134}$) (30). In contrast, none of the four

strains encoding bla_{TEM} but not $bla_{CTX-M-1}$ group, $bla_{CTX-M-9}$ group, bla_{CMY}, or ESBL-encoding

bla_{SHV} enzymes were ESBLs, with all being bla_{TEM-1}. Three of these were negative with the

double disc diffusion method, indicating probable chromosomal mutations underpinning this

phenotype. The basis for the ESBL phenotype in the fourth strain could not be elucidated.

Correlation between bla_{CMY} and cefoxitin resistance. Of the 81 cefoxitin-resistant isolates,

bla_{CMY} was detected in only 31 (38.3%) cases. Although imperfect, cefoxitin resistance is

often used as a marker for AmpC presence (32). Correlation is also dependent on the MIC,

with higher MICs more reliably predicting AmpC presence (32). Our findings concur with this

prior work, with 61.7% of cefoxitin-resistant isolates failing to amplify bla_{CMY}. AmpC has

previously been reported as an underappreciated mechanism of 3GC-R due to a lack of

standardized testing (33). As such, molecular methods that can detect other common and

emerging β-lactamase gene families (e.g. bla_{DHA}, bla_{OXY}, bla_{MIR}) may prove useful for infection

control management and treatment of multidrug-resistant infections. This is particularly true

for piperacillin-tazobactam, which may not be effective against AmpC producers (29, 34).

WGS identifies uncommon ESBL genes. Eleven (3.8%) PCR-negative isolates that otherwise

successfully amplified their 16S rDNA internal control (seven *E. coli*, one *K. pneumoniae*, one

K. oxytoca, one *K. aerogenes* and one *E. cloacae*) were subjected to WGS to determine the

molecular basis of 3GC-R. Six were found to carry uncommon β-lactamase enzymes, with

three *E. coli* encoding bla_{DHA-1}, one *K. pneumoniae/K. oxytoca* mixture harboring $bla_{OXY-2-4}$,

one *K. oxytoca* carrying $bla_{OXY-2-1}$, and one *E. cloacae* carrying bla_{ACT-17}. The remaining five

isolates had no discernable 3GC-R determinants, although misclassification via automated

susceptibility testing or upregulation of chromosomal β-lactamases are both possible causes

for this discrepancy.

Six isolates (three *E. cloacae*, two *E. coli* and one *K. pneumoniae*) exhibiting meropenem

resistance alongside 3GC-R were also subjected to WGS to determine the basis of resistance

towards this carbapenem. As expected, meropenem resistance was mostly conferred by

carbapenemases, with 4/6 (66.6%) isolates possessing one of these enzymes. bla_{NDM-1} or

bla_{IMP-4} were identified in two *E. cloacae* isolates, bla_{NDM-5} was identified in an *E. coli* strain,

and an OXA-48-like enzyme, $bla_{OXA-232}$, was identified in a *K. pneumoniae* isolate. No

carbapenemase gene could be identified in the remaining two isolates (one *E. coli* and one *E.*

cloacae). Although not investigated in this study, chromosomally-encoded porin loss or

efflux pump overexpression may form the basis of meropenem resistance in these strains.

Monitoring patient-to-patient spread in an ICU. Fifty-four (18.8%) 3GC-R isolates obtained

during this study originated from ICU-admitted patients. Due to their dominance, only *E. coli*

and *K. pneumoniae* genomes were analyzed for potential transmission events. Among the *E.*

coli population (Figure 3), isolates SCHI0016.S.60 and SCHI0016.S.62, and SCHI0016.S.265

and SCHI0016.S.46, were separated by 10 SNPs and 2 SNPs, respectively; however, both

pairs came from individual patients, reflecting expected levels of within-host evolution of a

single infecting strain. *E. coli* isolates SCHI0016.S.42 and SCHI0016.S.294 were very similar,

being separated by 13 SNPs. Although both strains were isolated from ICU-admitted

patients, SCHI0016.S.294 was isolated 306 days after SCHI0016.S.42 with no other linking

epidemiology identified. Additionally, no similar strains were isolated from ICU patients over

this time period suggesting that these do not represent transmission. *E. coli* isolates

SCHI0016.S.172 and SCHI0016.S.274 differed by ~300 SNPs, with no epidemiological or

temporal correlation. ~300 SNPs separated SCHI0016.S.112 and SCHI0016.S.229, again with

no demonstrable epidemiological link. Among *K. pneumoniae* (Figure 4), the closest isolates,

SCHI0016.S.15 and SCHI0016.S.261, differed by 361 SNPs, with no epidemiological link.

Additionally, no horizontal transfer of plasmids carrying ESBL determinants was observed. As

standard practice, all ICU patients, and all patients in other wards infected or colonized with

ESBL-producing *E. coli* and *K. pneumoniae*, have single rooms with an assigned nurse (1:1

ratio), with contact precautions exercised. In addition, hand hygiene rates in our hospital ICU

are generally above 80% (35). Taken together, our results demonstrate no evidence of

transmission in the ICU, suggesting that the implemented hygiene measures are effective.

Population structure of E. coli and K. pneumoniae genomes. We subjected an additional 36

strains to WGS (total *n*=98) to capture a snapshot of genomic diversity and population

structure (Figures 3 and 4). *In silico* MLST of the 79 *E. coli* strains found that ST131 was most

common (*n*=11; 13.9%). This finding is consistent with a recent study of the population

structure of Australian, New Zealand, and Singaporean multidrug-resistant *E. coli* isolates

(29). The highly successful ST131 clone has now spread worldwide, where it frequently

causes urinary tract and bloodstream infections and accounts for 12–27% of all community-

acquired *E. coli* infections (36, 37). Among our ST131 isolates, we found high frequencies of

fluoroquinolone resistance-conferring variants (GyrA$_{S83L}$, GyrA$_{D87N}$, ParC$_{S80I}$ and ParC$_{E84V}$)

alongside frequent carriage of plasmid-borne CTX-M-14 (CTX-M-9 group) and CTX-M-15

(CTX-M-1 group) ESBLs. We also identified three strains belonging to the emerging multi-

drug resistant pandemic clone, ST1193. These isolates also possessed the same three

fluoroquinolone-resistant SNPs as ST131. It has been proposed that ST1193 strains evolved

fluoroquinolone resistance relatively recently through multiple recombination events with

other *E. coli* strains (38).

Phylogenomic analysis of our 10 *K. pneumoniae* genomes identified several distinct lineages,

which comprised two ST14 strains and one each of ST17, ST307, ST309, ST395, ST656, ST697,

ST873, and ST969. Of note, the globally disseminated ST258 clone, which is responsible for a

large number of hospital deaths (39), was not identified in our dataset. However, we

identified another globally disseminated clone, ST395, which has also been retrieved from

clinical cases in Europe and Asia (40). Concerningly, this ST395 isolate, SCHI0016.S.137,

carried both the ESBL genes *bla*$_{SHV-11}$ and *bla*$_{CTX-M-15}$ and the OXA-48-like carbapenemase,

bla$_{OXA-232}$, the latter of which confers low-level meropenem resistance. Additionally, we

observed one *K. pneumoniae* ST307, which is an emerging multi-drug resistant, globally

disseminated clone postulated to have emerged in the mid-1990s(41). Although reports of

this clone from Australian populations in uncommon it has been previously observed(41).

This study had several recognized limitations. First, isolates were collected from a

nosocomial setting; no community sources were sampled. It is well-established that

differences in AMR patterns and their associated molecular mechanisms exist between

hospital and community settings. Second, the absence of certain high-risk populations within

our hospital (e.g. solid organ or hematopoietic stem cell transplantation) limits the

generalizability of our findings to other settings. Third, our PCR assays can only identify the

presence or change in copy number of the most common ESBL and *ampC* genes. Although

these loci make up the majority (>90%) of β-lactamase enzymes present in 3GC-R

Enterobacteriaceae, they do not capture all potential ESBLs, leaving less common ESBL and

ampC genes unidentified. In addition, PCR assays for bla_{TEM} or bla_{SHV} cannot discriminate

between ESBLs and non-ESBLs, with additional assays or concurrent WGS required to

confirm ESBL status. Our assays also cannot detect chromosomal changes leading to AMR

(e.g. porin mutations or efflux pump up-regulation), making these assays less useful for 3GC-

R bacteria that frequently encode these AMR determinants (e.g. *Pseudomonas aeruginosa*).

Fourth, our PCR was unable to distinguish chromosomal vs. low-copy plasmid-borne β-

lactamase genes, which has implications for AMR transmissibility between and among

bacterial populations. Finally, ~40% of samples were obtained from rectal swabs, which have

limitations for detection of multidrug-resistant organisms due to the high microbial burden

coupled with difficulties in sample collection. It is therefore likely that we did not capture all

3GC-R isolates during our study period, and thus may have missed potential transmission

events.

406
Conclusion

Taken together, our findings identified a predominance of CTX-M (and to a lesser extent,

CMY) β-lactamases among 3GC-R Enterobacteriaceae in the Sunshine Coast region. These

findings are consistent with global findings of Gram-negative bacterial AMR determinants,

particularly among *E. coli*. Our study demonstrates the likely absence of 3GC-R

Enterobacteriaceae transmission in our ICU setting, indicating effective infection control

processes. We show that rapid molecular assays, such as our multiplex PCR, can detect

common β-lactamase genes from bacterial culture within 40 minutes. These early molecular

results would permit more informed selection of antibiotic therapy up to 1-2 days prior to

3GC-R confirmation with traditional culture methodologies. Further research is required to

determine whether these molecular assays can be used directly on clinical specimens, and

whether the implementation of molecular assays and more advanced technologies (e.g.

WGS) is associated with improved patient outcomes in the hospital setting.

Acknowledgements and funding

This study was funded by the Wishlist Sunshine Coast Health Foundation (awarded to AGS,

KC, SS, and DSS). DSS and EPP were funded by Advance Queensland (grants AQRF13016-

17RD2 and AQIRF0362018, respectively). We thank Dr Delaney Burnard (University of

Queensland Centre for Clinical Research) and the Sunshine Coast University Hospital

pathology laboratory for laboratory assistance, and Rhys T. White (University of Queensland)

for helpful feedback on this manuscript.

Figures

Figure 1: Singleplex and quadriplex PCR performance on third-generation cephalosporin-

resistant Enterobacteriaceae. 1: *Enterobacter cloacae* SCHI0016.S.104 containing CMY. 2:

Escherichia coli SCHI0016.S.205 containing TEM. 3A: *E. coli* SCHI0016.S.187 positive for CTX-

M group 1. 3B: *E. coli* SCHI0016.S.60 positive for both CTX-M groups 1 and 9. 3C: *E. coli*

SCHI0046.S.46. 3D: *E. coli* with only 16s rDNA amplification with no CTX-M or SHV detected.

Note that both 16S rDNA and the TEM assays have late amplification in the no-template

controls due to low levels of recombinant *E. coli* DNA present in the Sso Advanced *Taq*

polymerase.

Figure 2. Limits of Detection (LoD) and Quantitation (LoQ) for the extended spectrum β-

lactamase real-time PCR assays developed in this study. The bla_{CMY} and bla_{TEM} assays were

examined as singleplex PCRs, whereas the 16S rDNA, $bla_{CTX-M-1}$ group, $bla_{CTX-M-9}$ group, and

bla_{SHV} assays were examined in quadriplex format.

Figure 3. Whole-genome phylogeny of 79 third-generation cephalosporin-resistant

Escherichia coli isolates and their β-lactamase gene complement. Sequence type (ST)131 is

indicated by the gray-shaded box; strains isolated from intensive care unit patients are

452 highlighted in green.

Figure 4. Whole-genome phylogeny and β-lactamase enzyme presence in *Klebsiella*

pneumoniae. Green-shaded isolates were isolated from the intensive care unit.

Transparency declarations

Dr. Harris reports research grants from Shionogi, Sandoz, and MSD and speaker's fees from Pfizer, outside the submitted work. No conflicts are declared for remaining authors

462
References

1. Gupta V, Ye G, Olesky M, Lawrence K, Murray J, Yu K. 2019. National
prevalence estimates for resistant Enterobacteriaceae and Acinetobacter
species in hospitalized patients in the United States. Int J Infect Dis
doi:10.1016/j.ijid.2019.06.017.
2. Biehl LM, Schmidt-Hieber M, Liss B, Cornely OA, Vehreschild MJ. 2016.
Colonization and infection with extended spectrum beta-lactamase
producing Enterobacteriaceae in high-risk patients - Review of the
literature from a clinical perspective. Crit Rev Microbiol 42:1-16.
3. Anonymous. 2014. Antimicrobial resistance - global report on
surveillance 2014.
https://apps.who.int/iris/bitstream/handle/10665/112642/978924156
4748_eng.pdf;jsessionid=519244EE0EEF520027CE4098504150B4?seque
nce=1. Accessed
4. Anonymous. 2018. Gram-negative Sepsis Outcome Program 2018 Report.
Australian Group on Antimicrobial Resistance
5. Paterson DL, Bonomo RA. 2005. Extended-spectrum beta-lactamases: a
clinical update. Clin Microbiol Rev 18:657-86.
6. Consortium BaB. 2015. Technology overview | BD MAX Enteric Bacterial
Panel for identifying pathogens in contagious gastroenteritis | Advice |
NICE, *on* National Institute for Health and Care Excellence.
https://www.nice.org.uk/advice/mib32/chapter/Technology-overview.
Accessed 14-2-2020.
7. MacVane SH, Nolte FS. 2016. Benefits of Adding a Rapid PCR-Based Blood
Culture Identification Panel to an Established Antimicrobial Stewardship
Program. J Clin Microbiol 54:2455-63.
8. Echavarria M, Marcone DN, Querci M, Seoane A, Ypas M, Videla C, O'Farrell
C, Vidaurreta S, Ekstrom J, Carballal G. 2018. Clinical impact of rapid
molecular detection of respiratory pathogens in patients with acute
respiratory infection. J Clin Virol 108:90-95.
9. Souverein D, Euser SM, van der Reijden WA, Herpers BL, Kluytmans J,
Rossen JWA, Den Boer JW. 2017. Clinical sensitivity and specificity of the
Check-Points Check-Direct ESBL Screen for BD MAX, a real-time PCR for
direct ESBL detection from rectal swabs. J Antimicrob Chemother
72:2512-2518.
10. Yang S, Lin S, Kelen GD, Quinn TC, Dick JD, Gaydos CA, Rothman RE. 2002.
Quantitative multiprobe PCR assay for simultaneous detection and
identification to species level of bacterial pathogens. J Clin Microbiol
40:3449-54.
11. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M,
Rozen SG. 2012. Primer3--new capabilities and interfaces. Nucleic Acids
Res 40:e115.
12. Bae IK, Kang HK, Jang IH, Lee W, Kim K, Kim JO, Jeong SH, Lee K. 2015.
Detection of Carbapenemases in Clinical Enterobacteriaceae Isolates
Using the VITEK AST-N202 Card. Infect Chemother 47:167-74.
13. Harris PNA, Tambyah PA, Lye DC, Mo Y, Lee TH, Yilmaz M, Alenazi TH,
Arabi Y, Falcone M, Bassetti M, Righi E, Rogers BA, Kanj S, Bhally H, Iredell

J, Mendelson M, Boyles TH, Looke D, Miyakis S, Walls G, Al Khamis M, Zikri
A, Crowe A, Ingram P, Daneman N, Griffin P, Athan E, Lorenc P, Baker P,
Roberts L, Beatson SA, Peleg AY, Harris-Brown T, Paterson DL,
Investigators MT, the Australasian Society for Infectious Disease Clinical
Research N. 2018. Effect of Piperacillin-Tazobactam vs Meropenem on 30-
Day Mortality for Patients With E coli or Klebsiella pneumoniae
Bloodstream Infection and Ceftriaxone Resistance: A Randomized Clinical
Trial. JAMA 320:984-994.

14. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics 30:2114-20.

15. Price EP, Viberg LT, Kidd TJ, Bell SC, Currie BJ, Sarovich DS. 2018.
Transcriptomic analysis of longitudinal Burkholderia pseudomallei
infecting the cystic fibrosis lung. Microb Genom 4.

16. Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with
Kraken 2. Genome Biol 20:257.

17. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a
new versatile metagenomic assembler. Genome Res 27:824-834.

18. Antipov D, Hartwick N, Shen M, Raiko M, Lapidus A, Pevzner PA. 2016.
plasmidSPAdes: assembling plasmids from whole genome sequencing
data. Bioinformatics 32:3380-3387.

19. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local
alignment search tool. J Mol Biol 215:403-10.

20. Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets.
Bioinformatics 32:605-7.

21. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA,
Dave BM, Pereira S, Sharma AN, Doshi S, Courtot M, Lo R, Williams LE,
Frye JG, Elsayegh T, Sardar D, Westman EL, Pawlowski AC, Johnson TA,
Brinkman FS, Wright GD, McArthur AG. 2017. CARD 2017: expansion and
model-centric curation of the comprehensive antibiotic resistance
database. Nucleic Acids Res 45:D566-D573.

22. Sarovich DS, Price EP. 2014. SPANDx: A genomics pipeline for
comparative analysis of large haploid whole genome re-sequencing
datasets. BMC Res Notes 7:618.

23. Aziz A, Currie BJ, Mayo M, Sarovich DS, Price EP. 2020. Comparative
genomics confirms a rare melioidosis human-to-human transmission
event and reveals incorrect phylogenomic reconstruction due to
polyclonality. Microb Genom doi:10.1099/mgen.0.000326.

24. Hayashi T, Makino K, Ohnishi M, Kurokawa K, Ishii K, Yokoyama K, Han
CG, Ohtsubo E, Nakayama K, Murata T, Tanaka M, Tobe T, Iida T, Takami
H, Honda T, Sasakawa C, Ogasawara N, Yasunaga T, Kuhara S, Shiba T,
Hattori M, Shinagawa H. 2001. Complete genome sequence of
enterohemorrhagic Escherichia coli O157:H7 and genomic comparison
with a laboratory strain K-12. DNA Res 8:11-22.

25. Liu P, Li P, Jiang X, Bi D, Xie Y, Tai C, Deng Z, Rajakumar K, Ou HY. 2012.
Complete genome sequence of Klebsiella pneumoniae subsp. pneumoniae
HS11286, a multidrug-resistant strain isolated from human sputum. J
Bacteriol 194:1841-2.

26. Inouye M, Dashnow H, Raven LA, Schultz MB, Pope BJ, Tomita T, Zobel J,
Holt KE. 2014. SRST2: Rapid genomic surveillance for public health and
hospital microbiology labs. Genome Med 6:90.
27. Wilgenbusch JC, Swofford D. 2003. Inferring evolutionary trees with
PAUP*. Curr Protoc Bioinformatics Chapter 6:Unit 6 4.
28. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates
and new developments. Nucleic Acids Res 47:W256-W259.
29. Harris PNA, Ben Zakour NL, Roberts LW, Wailan AM, Zowawi HM,
Tambyah PA, Lye DC, Jureen R, Lee TH, Yin M, Izharuddin E, Looke D,
Runnegar N, Rogers B, Bhally H, Crowe A, Schembri MA, Beatson SA,
Paterson DL, investigators MT. 2018. Whole genome analysis of
cephalosporin-resistant Escherichia coli from bloodstream infections in
Australia, New Zealand and Singapore: high prevalence of CMY-2
producers and ST131 carrying blaCTX-M-15 and blaCTX-M-27. J
Antimicrob Chemother 73:634-642.
30. Liakopoulos A, Mevius D, Ceccarelli D. 2016. A Review of SHV Extended-
Spectrum beta-Lactamases: Neglected Yet Ubiquitous. Front Microbiol
7:1374.
31. Orencia MC, Yoon JS, Ness JE, Stemmer WP, Stevens RC. 2001. Predicting
the emergence of antibiotic resistance by directed evolution and
structural analysis. Nat Struct Biol 8:238-42.
32. Jacoby GA. 2009. AmpC beta-lactamases. Clin Microbiol Rev 22:161-82,
Table of Contents.
33. Meini S, Tascini C, Cei M, Sozio E, Rossolini GM. 2019. AmpC beta-
lactamase-producing Enterobacterales: what a clinician should know.
Infection 47:363-375.
34. Cheng L, Nelson BC, Mehta M, Seval N, Park S, Giddins MJ, Shi Q, Whittier
S, Gomez-Simmonds A, Uhlemann AC. 2017. Piperacillin-Tazobactam
versus Other Antibacterial Agents for Treatment of Bloodstream
Infections Due to AmpC beta-Lactamase-Producing Enterobacteriaceae.
Antimicrob Agents Chemother 61.
35. Grayson ML, Stewardson AJ, Russo PL, Ryan KE, Olsen KL, Havers SM,
Greig S, Cruickshank M, Hand Hygiene A, the National Hand Hygiene I.
2018. Effects of the Australian National Hand Hygiene Initiative after 8
years on infection control practices, health-care worker education, and
clinical outcomes: a longitudinal study. Lancet Infect Dis 18:1269-1277.
36. Giedraitiene A, Vitkauskiene A, Pavilonis A, Patamsyte V, Genel N, Decre D,
Arlet G. 2017. Prevalence of O25b-ST131 clone among Escherichia coli
strains producing CTX-M-15, CTX-M-14 and CTX-M-92 beta-lactamases.
Infect Dis (Lond) 49:106-112.
37. Nicolas-Chanoine MH, Bertrand X, Madec JY. 2014. Escherichia coli ST131,
an intriguing clonal group. Clin Microbiol Rev 27:543-74.
38. Tchesnokova V, Radey M, Chattopadhyay S, Larson L, Weaver JL, Kisiela D,
Sokurenko EV. 2019. Pandemic fluoroquinolone resistant Escherichia coli
clone ST1193 emerged via simultaneous homologous recombinations in
11 gene loci. Proc Natl Acad Sci U S A 116:14740-14748.
39. David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, Abudahab K,
Goater R, Giani T, Errico G, Aspbury M, Sjunnebo S, Eu SWG, Group ES, Feil
EJ, Rossolini GM, Aanensen DM, Grundmann H. 2019. Epidemic of

607  carbapenem-resistant Klebsiella pneumoniae in Europe is driven by
608  nosocomial spread. Nat Microbiol doi:10.1038/s41564-019-0492-8.
609 40. Lee CR, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. 2016. Global
610  Dissemination of Carbapenemase-Producing Klebsiella pneumoniae:
611  Epidemiology, Genetic Context, Treatment Options, and Detection
612  Methods. Front Microbiol 7:895.
613 41. Wyres KL, Hawkey J, Hetland MAK, Fostervold A, Wick RR, Judd LM,
614  Hamidian M, Howden BP, Lohr IH, Holt KE. 2019. Emergence and rapid
615  global dissemination of CTX-M-15-associated Klebsiella pneumoniae
616  strain ST307. J Antimicrob Chemother 74:577-581.

621 **Tables**

623 **Table 1:** Summary of antimicrobial resistance mechanisms among 288 clinical Enterobacteriaceae isolates retrieved from 264 patients admitted to a
624 Southeast Queensland hospital over a 14-month period.

Organism	E. coli (n=231)	K. pneumoniae (n=49)	K. aerogenes (n=1)	K. oxytoca (n=3)	E. cloacae (n=4)
Phenotypic antimicrobial resistance[a]					
Ceftriaxone	223 (96.5%)	46 (93.9%)	1 (100%)	3 (100%)	4 (100%)
Ceftazidime	215 (93.1%)	47 (95.9%)	1 (100%)	2 (66.7%)	4 (100%)
Cefepime	201 (87.0%)	43 (87.8%)	1 (100%)	2 (66.7%)	3 (75.0%)
Cefoxitin	64 (27.7%)	12 (24.5%)	1 (100%)	0 (0%)	4 (100%)
Piperacillin-tazobactam	65 (28.1%)	31 (63.3%)	1 (100%)	3 (100%)	4 (100%)
Ticarcillin-clavulanate	186 (80.5%)	46 (93. 9%)	0 (0%)	3 (100%)	3 (75.0%)
Meropenem	3 (1.3%)	2 (4.1%)	0 (0%)	0 (0%)	3 (75.0%)
ESBL phenotypic test[b]					
Positive	204 (88.3%)	46 (93.9%)	0 (0%)	2 (66.7%)	4 (100%)
Negative	27 (11.7%)	3 (6.1%)	1 (100%)	1 (33.3%)	0 (0%)
ESBL and _ampC_ gene prevalence					
$bla_{CTX-M-1}$	104 (45.0%)	36 (73.5%)	0 (0%)	2 (66.7%)	2 (50.0%)
$bla_{CTX-M-9}$	89 (38.5%)	4 (8.2%)	0 (0%)	0 (0%)	0 (0%)
bla_{SHV}	9 (3.9%)	38 (77.6%)	0 (0%)	0 (0%)	1 (25.0%)
bla_{TEM}	93 (40.3%)	33 (67.3%)	0 (0%)	2 (66.7%)	2 (50.0%)
bla_{CMY}	30 (13.0%)	3 (6.1%)	0 (0%)	0 (0%)	1 (25.0%)
PCR-negative[c]	7 (3.0%)	1 (4%)	1 (100%)	1 (33.3%)	1 (25.0%)

625 [a]Antimicrobial susceptibility testing performed by automated method (VITEK 2®) and EUCAST breakpoints applied
626 [b]ESBL phenotypic testing performed by disc diffusion method
627 [c]DNA integrity was confirmed by 16S rDNA PCR (10), which forms part of the quadriplex assay. Three PCR negative _E. coli_ strains encoded bla_{DHA-1}, one _E._
628 _cloacae_ encoded bla_{ACT-17}, one _K. oxytoca_ encoded a $bla_{OXY-2-1}$, a second _K. oxytoca_ (included as a mixture with _K. pneumoniae_) encoded $bla_{OXY-2-4}$, and 5/11
629 isolates had no β-lactamases identified despite whole-genome sequencing.