Title

DISPERSE – A trait database to assess the dispersal potential of aquatic macroinvertebrates

Authors

Romain Sarremejane[1*], Núria Cid[2,3*], Rachel Stubbington[1], Thibault Datry[2], Maria Alp[2], Miguel

Cañedo-Argüelles[3], Adolfo Cordero-Rivera[4], Zoltán Csabai[5,6], Cayetano Gutiérrez-Cánovas[7,8], Jani

Heino[9], Maxence Forcellini[2], Andrés Millán[10], Amael Paillex[11], Petr Pařil[6], Marek Polášek[6], José

Manuel Tierno de Figueroa[12], Philippe Usseglio-Polatera[13], Carmen Zamora-Muñoz[12] & Núria Bonada[3]

Affiliations

[1] School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK

[2] INRAE, UR RiverLy, centre de Lyon-Villeurbanne, 5 rue de la Doua CS70077, 69626 Villeurbanne

Cedex, France

[3] Grup de Recerca Freshwater Ecology, Hydrology and Management (FEHM), Departament de

Biologia Evolutiva, Ecologia i Ciències Ambientals, Facultat de Biologia, Institut de Recerca de la

Biodiversitat (IRBio), Universitat de Barcelona (UB), Diagonal 643, 08028 Barcelona, Catalonia, Spain

[4] ECOEVO Lab, E.E. Forestal, Univesidade de Vigo, Campus A Xunqueira, 36005 Pontevedra, Spain

[5] Department of Hydrobiology, University of Pécs, Ifjúság útja 6, H7624 Pécs, Hungary

[6] Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2, 61137 Brno,

Czech Republic

[7] Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of

Minho, Braga, Portugal

[8] Institute of Science and Innovation for Bio-Sustainability (IB-S), University of Minho, Braga, Portugal

[9] Finnish Environment Institute, Freshwater Centre, Paavo Havaksen Tie 3, FI-90570 Oulu, Finland

[10] Department of Ecology and Hydrology, Biology Faculty, Murcia University. Campus de Espinardo,

30100. Murcia, Spain

[11] Department of Aquatic Ecology, Eawag, Swiss Federal Institute of Aquatic Sciences,

Überlandstrasse 133, CH-8600 Dübendorf, Switzerland

[12] Departamento de Zoología, Facultad de Ciencias, Universidad de Granada, Avenida Fuente Nueva,

s/n, 18071 Granada, Spain

[13] Université de Lorraine, CNRS, UMR 7360, LIEC, Laboratoire Interdisciplinaire des Environnements

Continentaux, F-57070 Metz, France

*These authors contributed equally to this work.

Abstract

Dispersal is an essential process in population and community dynamics, but is difficult to measure in

the field. In freshwater ecosystems, information on biological traits related to organisms'

morphology, life history and behaviour provides useful dispersal proxies, but information remains

scattered or unpublished for many taxa. We compiled information on multiple dispersal-related

biological traits of European aquatic macroinvertebrates in a unique resource, the DISPERSE

database. DISPERSE includes 39 trait categories grouped into nine dispersal-related traits for 480

taxa, including Annelida, Mollusca, Platyhelminthes, and Arthropoda such as Crustacea and Insecta,

generally at the genus level. Information within DISPERSE can be used to address fundamental

questions in metapopulation ecology, metacommunity ecology, macroecology and evolutionary

ecology research. Information on dispersal proxies can be applied to improve predictions of

ecological responses to global change, and to inform improvements to biomonitoring and

conservation management strategies. The diverse sources used in DISPERSE complement existing

trait databases by providing new information on dispersal traits, most of which would not otherwise

be accessible to the scientific community.

Background & Summary

Dispersal is a fundamental ecological process that affects the organization of biological diversity at multiple temporal and spatial scales[1,2]. Dispersal strongly influences metapopulation and metacommunity dynamics through the movement of individuals and species, respectively[3]. A better understanding of dispersal processes can inform biodiversity management practices[4,5]. However, dispersal is difficult to measure directly, particularly for small organisms, including most invertebrates[6]. Typically, dispersal is measured for single species[7,8] or combinations of few species within one taxonomic group[9–11], using methods based on mark and recapture, stable isotopes, or population genetics[5,12]. Such methods can directly assess dispersal events but are expensive, time-consuming, and thus impractical for studies conducted at the community level or at large spatial scales. In this context, taxon-specific biological traits represent a cost-effective alternative that may serve as proxies for dispersal[5,6,13,14]. These traits interact with landscape structure to determine patterns of effective dispersal[15,16].

Aquatic macroinvertebrates inhabiting freshwater ecosystems include taxa with diverse dispersal modes and abilities (Figure 1). For species with complex life cycles, such as some insects, this diversity is enhanced by distinct dispersal strategies among life stages. For example, juveniles of many aquatic insects disperse actively and/or passively in water whereas adults fly over land[17]. In any case, dispersal is affected by multiple traits relating to the morphology[6,12], life history and behaviour[2] of different life stages.

We compiled and harmonized information on dispersal-related traits of aquatic macroinvertebrates from across Europe, including both aquatic and aerial (i.e. flying) stages. Although information on some dispersal-related traits such as body size, reproduction, locomotion and dispersal mode is available in online databases for European[18–20] and North American taxa[21], other relevant information is scattered across published literature and unpublished data. Using the expertise of 19 experts, we built a comprehensive database containing nine dispersal-related traits divided into 39

75 trait categories for 480 European taxa. Dispersal-related traits were selected and their trait

76 categories fuzzy-coded[22] following a structure comparable to existing databases[23]. Our aim was to

77 provide a single resource facilitating the inclusion of dispersal in ecological research, and to create

78 the basis for a global dispersal database.

Methods

Dispersal-related trait selection criteria

We defined dispersal as the unidirectional movement of individuals from one location to another[1],

assuming that population-level dispersal rates depend on both the number of dispersers/propagules

and dispersers' ability to move across the landscape[11,24].

We selected nine dispersal-related morphological, behavioural and life-history traits (Table 1).

Selected morphological traits were maximum body size, female wing length and wing pair type, the

latter two relating only to flying adult insects. Body size influences organisms' dispersal[6], especially

for active dispersers[25], with larger animals more capable of active dispersal over longer distances

(e.g. flying adult dragonflies[6]). Wing morphology, and in particular wing length, is related to the

dispersal of flying adult insects[6,26]. Female wing length was selected because females connect and

sustain populations through oviposition, thus representing adult insects' flying and recolonization

capacity[27]. Females with larger wings are likely to oviposit farther from their source population[6,10,28].

We also described wing morphology as insect wing types, i.e. one or two pairs of wings, and the

presence of halters, elytra or hemielytra, or small hind wings[12] (Figure 1). Selected life-history traits

were adult life span, life-cycle duration, annual number of reproductive cycles and lifelong fecundity.

Adult life span and life-cycle duration respectively reflect the adult (i.e. reproductive) and total life

duration, with longer-lived animals typically having more dispersal opportunities[13]. The annual

number of reproductive cycles and lifelong fecundity assess dispersal capacity based on potential

propagule production, with multiple reproductive cycles and abundant eggs increasing the expected

number of dispersal events[6]. Dispersal behaviour was represented by a taxon's predominant

dispersal mode (passive and/or active, aquatic and/or aerial), and by propensity to drift, which

indicates the frequency of flow-mediated passive downstream dispersal.

Data acquisition and compilation

A taxa list was generated based on existing trait databases[20] and modified to reflect current

taxonomic classifications. Trait information was sourced primarily from the literature using Google

Scholar searches of keywords including trait names, synonyms and taxon names, and by searching in

existing databases[18,21]. Altogether, >300 peer-reviewed articles and book chapters were consulted

(Supplementary File 1). When no European studies were available, we considered information from

other continents only if experts evaluated traits as comparable across regions. When published

information was lacking, traits were coded based on authors' expert knowledge and direct

measurements. Specifically, for 139 species in 69 genera of Coleoptera and Heteroptera, female wing

lengths were characterized using measurements of 538 individuals in experts' reference collections,

sampled in Finland, Hungary and Greece. The number of species measured within a genus varied

between 1 and 10 in proportion to the number of species within each genus in Europe. For example,

for the most specious genera, common and rare species as well as northern latitude and

Mediterranean species were included.

Fuzzy-coding approach and taxonomic resolution

Traits were coded using a fuzzy approach in which a value given to each trait category indicates if the

taxon has no (0), weak (1), moderate (2) or strong (3) affinity with the category[22]. This method can

incorporate intra-taxon variability when trait profiles differ among e.g. species within a genus, early

and late instars, or within one species in different environments[29]. Most traits were coded at genus

level, but some Diptera and Annelida were coded at family, sub-families or tribe level because of

their complex taxonomy, identification difficulties and the scarcity of reliable information about their

traits.

Data Records

DISPERSE can be downloaded as an Excel spreadsheet from the Intermittent River Biodiversity Analysis and Synthesis webpage[30] or Figshare at: https://figshare.com/s/4f074fb4bdbbe8cf149e.

The database comprises three sheets: DataKey, Data and Reference list. The "Datakey" sheet summarizes the content of each column in the "Data" sheet. The "Data" sheet includes the fuzzy-coded trait categories and cites the sources used to code each trait. The four first columns list the taxa and their taxonomy (group, family, genus [or lowest taxonomic resolution achieved including e.g. tribes and sub-families] and genus synonyms) to allow users to sort and compile information. Sources are cited in chronological order by the surname of the first author and the year of publication. Expert evaluations are reported as "Unpublished" followed by the name of the expert providing the information. Direct measurements are reported as "Direct measurement from" followed by the expert's name. The "Reference list" sheet contains the references cited in the "Data" sheet, organized in alphabetical order and then by date (also available in Supplementary File 1).

In total, the database contains nine dispersal-related traits divided into 39 trait categories for 480 taxa. Most (78%) taxa are insects, principally Coleoptera and Trichoptera, as these are, together with Diptera, the most diverse orders in freshwater ecosystems[31]. DISPERSE provides complete trait information for 61% of taxa, with 1-2 traits being incomplete for the 39% remaining taxa (Table 1, Figure 2). The traits with the highest percentage of information across taxa were wing pair type and maximum body size, followed by dispersal strategy, life-cycle duration, potential number of reproductive cycles per year, and female wing length (Table 1). The percentage of completed information was lower for two life-history traits: adult lifespan and lifelong fecundity (Table 1).

Technical Validation

Most of the trait information (88%) originated from published literature (Supplementary File 1) and the rest was coded based on expert knowledge (9%) and direct measurements (3%) (Table 1). The database states information sources for each trait and taxon, allowing users evaluate data quality.

Most traits were coded using multiple sources representing multiple species within a genus. When

only one study was available, we supplemented this information with expert knowledge, to ensure

that trait codes represented potential variability in the taxon.

Using insects as an example, we performed a fuzzy correspondence analysis (FCA)[22] to

visualize variability in trait composition among taxa (Figure 3). Insect orders were clearly

distinguished based on their dispersal-related traits, with 32% of the variation explained by the first

two FCA axes. Wing pair type and lifelong fecundity had the highest correlation coefficient (0.87 and

0.63, respectively) with axis A1. Female wing length (0.73) and maximum body size (0.55) best

correlated with axis A2 (Figures 3 and 4). For example, Coleoptera typically produce few eggs and

have intermediate maximum body sizes and wing lengths, Odonata produce an intermediate number

of eggs and have long wings, and Ephemeroptera produce many eggs and have short wings (Figures 1

and 4).

The database currently represents a Europe-wide resource, which can be updated and expanded as

new information becomes available, to include more taxa and traits from across and beyond Europe.

For example, additional information could be collected on other measures of wing morphology[10,14]

and functionality or descriptors of exogenous dispersal vectors such as wind and animals[32]. DISPERSE

lays the foundations for a global dispersal trait database, the lack of which is recognized as limiting

research progress across multiple disciplines[33].

Usage Notes

DISPERSE is the first publicly available database describing the dispersal traits of aquatic

macroinvertebrates including information on both aquatic and aerial (i.e. flying) life stages. It

provides good coverage of macroinvertebrates at the genus level, which is generally considered as

adequate to capture biodiversity dynamics[34–37]. It will promote incorporation of dispersal proxies into

fundamental and applied population and community ecology in aquatic ecosystems[5]. In particular,

metacommunity ecology may benefit from the use of dispersal traits[15,38], which enable classification

of taxa according to their dispersal potential in more detail. Such classification, used in combination

with, for example, spatial distance measurements[39,40], may advance our understanding of the effects

of regional dispersal processes on community assembly and biodiversity patterns. Improved

knowledge of taxon-specific dispersal abilities may also inform the design of more effective

management practices. For example, recognizing dispersal abilities in biomonitoring methods could

inform enhancements to catchment-scale management strategies that support ecosystems adapting

to global change[41,42]. DISPERSE could also inform freshwater conservation strategies by establishing

different priorities depending on organisms' dispersal capacities in relation to spatial connectivity[43].

DISPERSE could also improve species distribution models (SDMs), in which dispersal has

rarely been considered due to insufficient data [13], limiting the accuracy of model predictions[44].

Recent trait-based approaches have begun to integrate dispersal into SDMs, and DISPERSE could

increase model accuracy[45,46]. Including dispersal in SDMs is especially relevant to assessments of

biodiversity loss and species vulnerability to climate change[45,47,48]. DISPERSE could also advance

understanding of eco-evolutionary relationships and biogeographical phenomena. In an evolutionary

context, groups with lower dispersal abilities should be genetically and taxonomically richer due to

long-term isolation[49,50]. From a biogeographical perspective, regions affected by glaciations should

have species with greater dispersal abilities, enabling postglacial recolonization[51].

By capturing different dispersal-related biological traits, DISPERSE provides information on

organisms' potential ability to move between localities as well as on recruitment and reproduction[15].

Traits also facilitate comparison of taxa with different dispersal strategies, which could inform studies

conducted at large spatial scales, independent of taxonomy[52].

Users should note that dispersal-related traits included in DISPERSE represent an indirect

measure of dispersal, not actual dispersal. Therefore, the database is not intended to substitute

population-level studies related to dispersal, but to act as a repository that collates and summarizes

information from such studies. As freshwater biodiversity declines at unprecedented rates[53,54],

collecting, harmonizing and sharing dispersal-related data on freshwater organisms will underpin

evidence-informed initiatives that seek to support the resilience of ecosystems adapting to global

change.

Acknowledgements

The study was supported by the COST Action CA15113 Science and Management of Intermittent

Rivers and Ephemeral Streams (http://www.smires.eu/), funded by COST (European Cooperation in

Science and Technology). NC was supported by the French research program Make Our Planet Great

Again. MC-A was supported by the MECODISPER project (CTM2017-89295-P) funded by the Spanish

Ministerio de Economía, Industria y Competitividad (MINECO) - Agencia Estatal de Investigación (AEI)

and cofunded by the European Regional Development Fund (ERDF). AC-R acknowledges funding by

MINECO-AEI-ERDF (CGL2014-53140-P). CG-C was supported by the EDRF (COMPETE2020 and

PT2020) and the Portuguese Foundation for Science and Technology, through the CBMA strategic

program UID/BIA/04050/2019 (POCI-01-0145-FEDER-007569) and the STREAMECO project

(Biodiversity and ecosystem functioning under climate change: from the gene to the stream,

PTDC/CTA-AMB/31245/2017). PP and MP were supported by the Czech Science Foundation (P505-

20-17305S). ZC was supported by the projects EFOP-3.6.1.-16-2016-00004, 20765-

3/2018/FEKUTSTRAT and TUDFO/47138/2019-ITM.

Author contributions

NB, NC and RSa developed the idea and data collection framework. RSa compiled most of the

dispersal-related trait data and structured the database. All authors contributed to the addition and

checking of information included in the database, and ZC provided direct measurements of several

taxa. AC-R designed Figure 1. MA, MC-A, CG-C and MF helped to finalize the database reference list.

NC wrote the first draft of the manuscript, and all authors contributed to finalizing it. RSt proofread

the manuscript.

Competing interests

226   The authors declare no competing interests.

References

1.   Bohonak, A. J. & Jenkins, D. G. Ecological and evolutionary significance of dispersal by

freshwater invertebrates. *Ecol. Lett.* **6**, 783–796 (2003).

2.   Clobert, J., Baguette, M., Benton, T. G. & Bullock, J. M. *Dispersal ecology and evolution.*

(Oxford Univ. Press, 2012).

3.   Heino, J. *et al.* Metacommunity organisation, spatial extent and dispersal in aquatic systems:

Patterns, processes and prospects. *Freshw. Biol.* **60**, 845–869 (2015).

4.   Barton, P. S. *et al.* Guidelines for using movement science to inform biodiversity policy.

Environ. Manage. **56**, 791–801 (2015).

5.   Heino, J. *et al.* Integrating dispersal proxies in ecological and environmental research in the

freshwater realm. *Environ. Rev.* **25**, 334–349 (2017).

6.   Rundle, S. D., Bilton, D. T. & Foggo, A. By wind, wings or water: body size, dispersal and range

size in aquatic invertebrates. in *Body Size: The Structure and Function of Aquatic Ecosystems*

(eds. Hildrew, A. G., Raffaelli, D. G. & Edmonds-Brown, R.) 186–209 (Cambridge Univ. Press,

2007).

7.   Macneale, K. H., Peckarsky, B. L. & Likens, G. E. Stable isotopes identify dispersal patterns of

stonefly populations living along stream corridors. *Freshw. Biol.* **50**, 1117–1130 (2005).

8.   Troast, D., Suhling, F., Jinguji, H., Sahlén, G. & Ware, J. A global population genetic study of

Pantala flavescens. PLoS One **11**, 1–13 (2016).

9.   French, S. K. & McCauley, S. J. The movement responses of three libellulid dragonfly species to

open and closed landscape cover. *Insect Conserv. Divers.* (2019). doi:10.1111/icad.12355

10.   Arribas, P. *et al.* Dispersal ability rather than ecological tolerance drives differences in range

size between lentic and lotic water beetles (Coleoptera: Hydrophilidae). *J. Biogeogr.* **39**, 984–

994 (2012).

11.   Lancaster, J. & Downes, B. J. Dispersal traits may reflect dispersal distances, but dispersers

may not connect populations demographically. *Oecologia* **184**, 171–182 (2017).

12.   Lancaster, J. & Downes, B. J. Dispersal in the terrestrial environment. in *Aquatic Entomology*

(eds. Lancaster, J. & Downes, B. J.) 137–153 (Oxford Univ. Press, 2013).

13.   Stevens, V. M. *et al.* Dispersal syndromes and the use of life-histories to predict dispersal.

Evol. Appl. **6**, 630–642 (2013).

14.   Outomuro, D. & Johansson, F. Wing morphology and migration status, but not body size,

habitat or Rapoport's rule predict range size in North-American dragonflies (Odonata:

Libellulidae). *Ecography* **42**, 309–320 (2019).

15.   Tonkin, J. D. *et al.* The role of dispersal in river network metacommunities: patterns,

processes, and pathways. *Freshw. Biol.* **63**, 141–163 (2018).

16.   Brown, B. L. & Swan, C. M. Dendritic network structure constrains metacommunity properties

in riverine ecosystems. *J. Anim. Ecol.* **79**, 571–580 (2010).

17.   Wikelski, M. *et al.* Simple rules guide dragonfly migration. *Biol. Lett.* **2**, 325–329 (2006).

18.   Schmidt-Kloiber, A. & Hering, D. www.freshwaterecology.info – An online tool that unifies,

standardises and codifies more than 20,000 European freshwater organisms and their

ecological preferences. *Ecol. Indic.* **53**, 271–282 (2015).

19.   Serra, S. R. Q., Cobo, F., Graça, M. A. S., Dolédec, S. & Feio, M. J. Synthesising the trait

information of European Chironomidae (Insecta: Diptera): towards a new database. *Ecol.*

Indic. **61**, 282–292 (2016).

20.   Tachet, H., Richoux, P., Bournaud, M. & Usseglio-Polatera, P. *Invertébrés d'Eau Douce:*

Systématique, Biologie, Écologie. (CNRS Éditions, 2010).

21.   Vieira, N. K. M. *et al. A Database of Lotic Invertebrate Traits for North America. U.S. Geological*

Survey Data Series 187 (2006).

22.   Chevenet, F., Dolédec, S. & Chessel, D. A fuzzy coding approach for the analysis of long-term

ecological data. *Freshw. Biol.* **31**, 295–309 (1994).

23.   Schmera, D., Podani, J., Heino, J., Erös, T. & Poff, N. L. R. A proposed unified terminology of

species traits in stream ecology. *Freshw. Sci.* **34**, 823–830 (2015).

24.   Lancaster, J., Downes, B. J. & Arnold, A. Lasting effects of maternal behaviour on the

distribution of a dispersive stream insect. *J. Anim. Ecol.* **80**, 1061–1069 (2011).

25.   Jenkins, D. G. *et al.* Does size matter for dispersal distance? *Glob. Ecol. Biogeogr.* **16**, 415–425

(2007).

26.   Harrison, R. G. Dispersal polymorphisms in insects. *Annu. Rev. Ecol. Syst.* **11**, 95–118 (1980).

27.   Graham, E. S., Storey, R. & Smith, B. Dispersal distances of aquatic insects: upstream crawling

by benthic EPT larvae and flight of adult Trichoptera along valley floors. *New Zeal. J. Mar.*

Freshw. Res. **51**, 146–164 (2017).

28.   Hoffsten, P. O. Site-occupancy in relation to flight-morphology in caddisflies. *Freshw. Biol.* **49**,

810–817 (2004).

29.   Bonada, N. & Dolédec, S. Does the Tachet trait database report voltinism variability of aquatic

insects between Mediterranean and Scandinavian regions? *Aquat. Sci.* **80**, 1–11 (2018).

30.   Sarremejane, R. *et al.* DISPERSE: a trait database to assess the dispersal potential of aquatic

macroinvertebrates. *Intermittent River Biodiversity Analysis and Synthesis* (2020). Available at:

http://irbas.inrae.fr/latest-news/disperse-database-sarremejane-et-al-2020.

31.   Lévêque, C., Balian, E. V & Martens, K. An assessment of animal species diversity in

continental waters. *Hydrobiologia* **542**, 39–67 (2005).

32.   Green, A. J. & Figuerola, J. Recent advances in the study of long-distance dispersal of aquatic

invertebrates via birds. *Divers. Distrib.* **11**, 149–156 (2005).

33.   Maasri, A. A global and unified trait database for aquatic macroinvertebrates: the missing

piece in a global approach. *Front. Environ. Sci.* **7**, 1–3 (2019).

34.   Cañedo-Argüelles, M. *et al.* Dispersal strength determines meta-community structure in a

dendritic riverine network. *J. Biogeogr.* **42**, 778–790 (2015).

35.   Datry, T. *et al.* Metacommunity patterns across three Neotropical catchments with varying

environmental harshness. *Freshw. Biol.* **61**, 277–292 (2016).

36.   Swan, C. M. & Brown, B. L. Metacommunity theory meets restoration: isolation may mediate

how ecological communities respond to stream restoration. *Ecol. Appl.* **27**, 2209–2219 (2017).

37.   Sarremejane, R., Mykrä, H., Bonada, N., Aroviita, J. & Muotka, T. Habitat connectivity and

dispersal ability drive the assembly mechanisms of macroinvertebrate communities in river

networks. *Freshw. Biol.* **62**, 1073–1082 (2017).

38.   Jacobson, B. & Peres-Neto, P. R. Quantifying and disentangling dispersal in metacommunities:

How close have we come? How far is there to go? *Landsc. Ecol.* **25**, 495–507 (2010).

39.   Sarremejane, R. *et al.* Do metacommunities vary through time? Intermittent rivers as model

systems. *J. Biogeogr.* **44**, 2752–2763 (2017).

40.   Datry, T., Moya, N., Zubieta, J. & Oberdorff, T. Determinants of local and regional

communities in intermittent and perennial headwaters of the Bolivian Amazon. *Freshw. Biol.*

61, 1335–1349 (2016).

41.   Cid, N. *et al.* A metacommunity approach to improve biological assessments in highly dynamic

freshwater ecosystems. *Bioscience* **70**, 427–438 (2020).

42.   Datry, T., Bonada, N. & Heino, J. Towards understanding the organisation of

metacommunities in highly dynamic ecological systems. *Oikos* **125**, 149–159 (2016).

43.   Hermoso, V., Cattarino, L., Kennard, M. J., Watts, M. & Linke, S. Catchment zoning for

freshwater conservation: refining plans to enhance action on the ground. *J. Appl. Ecol.* **52**,

940–949 (2015).

44.   Thuiller, W. *et al.* A road map for integrating eco-evolutionary processes into biodiversity

models. *Ecol. Lett.* **16**, 94–105 (2013).

45.   Willis, S. G. *et al.* Integrating climate change vulnerability assessments from species

distribution models and trait-based approaches. *Biol. Conserv.* **190**, 167–178 (2015).

46.   Cooper, J. C. & Soberón, J. Creating individual accessible area hypotheses improves stacked

species distribution model performance. *Glob. Ecol. Biogeogr.* **27**, 156–165 (2018).

47.   Markovic, D. *et al.* Europe's freshwater biodiversity under climate change: distribution shifts

and conservation needs. *Divers. Distrib.* **20**, 1097–1107 (2014).

48.   Bush, A. & Hoskins, A. J. Does dispersal capacity matter for freshwater biodiversity under

climate change? *Freshw. Biol.* **62**, 382–396 (2017).

49.   Bohonak, A. J. Dispersal, gene flow, and population structure. *Q. Rev. Biol.* **74**, 21–45 (1999).

50.   Dijkstra, K.-D. B., Monaghan, M. T. & Pauls, S. U. Freshwater biodiversity and aquatic insect

diversification. *Annu. Rev. Entomol.* **59**, 143–163 (2014).

51.   Múrria, C. *et al.* Local environment rather than past climate determines community

composition of mountain stream macroinvertebrates across Europe. *Mol. Ecol.* **26**, 6085–6099

(2017).

52.   Statzner, B. & Bêche, L. A. Can biological invertebrate traits resolve effects of multiple

stressors on running water ecosystems? *Freshw. Biol.* **55**, 80–119 (2010).

53.   Strayer, D. L. & Dudgeon, D. Freshwater biodiversity conservation: recent progress and future

challenges. *J. North Am. Benthol. Soc.* **29**, 344–358 (2010).

54.   Reid, A. J. *et al.* Emerging threats and persistent conservation challenges for freshwater

biodiversity. *Biol. Rev.* **94**, 849–873 (2019).

Figures and Tables

347 **Table 1** Dispersal-related aquatic macroinvertebrate traits included in the DISPERSE database and the

348 relative contribution of different sources of information (i.e. literature, expert knowledge, direct

349 measurement) used to build the database.

Trait	Categories	Taxa completed (%)	Source of information (%)		
			Literature	Expert	Measured
Maximum body size (cm)	< 0.25 ≥ 0.25 – 0.5 ≥ 0.5 – 1 ≥ 1 – 2 ≥ 2 – 4 ≥ 4 – 8 ≥ 8	99	100		
Female wing length (insects only) (mm)	< 5 ≥ 5 – 10 ≥ 10 – 15 ≥ 15 – 20 ≥ 20 – 30 ≥ 30 – 40 ≥ 40 – 50 ≥ 50	95	57	12	31
Wing pair type (insects only)	1 pair + halters 1 pair + elytra or hemielytra 1 pair + small hind wings 2 similar-sized pairs	100	100		
Life-cycle duration	≤ 1 year > 1 year	98	100		
Adult life span	< 1 week ≥ 1 week – 1 month ≥ 1 month – 1 year ≥ 1 year	79	65	35	
Lifelong fecundity	< 100 eggs ≥ 100 – 1000 eggs ≥ 1000 – 3000 eggs ≥ 3000 eggs	75	77	23	
Potential number of reproductive cycles per year	< 1 1 >1	98	100		
Dispersal	Aquatic active	98	100		

Trait	Categories	Taxa	Source of information (%)		
strategy	Aquatic passive				
	Aerial active				
	Aerial passive				
Propensity to drift	Rare/catastrophic				
	Occasional	80	90	10	
	Frequent				
All traits		61	88	9	3

350
351

Figure 1. The dispersal-related trait diversity of aquatic macroinvertebrates. Taxa that disperse in

water include the crustaceans *Potamon* (a) and *Asellus* (red arrow in b), planarians (b), the bivalve

mollusc *Unio* (c), insect larvae such as the Diptera genus *Simulium* (d) and Plecoptera genus *Leuctra*

(e), and adult Coleoptera including dytiscid *Cybister* (f). Such aquatic dispersers may move passively

in the drift (c, d) and/or actively crawl or swim (a, b, e, f). Most adult insects have wings and can fly

overland (f-n). Wings are morphologically diverse and include various types: one wing pair, as in

Diptera such as the syrphid genus *Eristalis* (g); one pair of wings with elytra for Coleoptera including

the genus *Enochrus* (h) or with hemielytra for Heteroptera such as the genus *Hesperocorixa* (i); two

wing pairs including one pair of small hind wings for Ephemoptera including the genus *Ephemera* (j);

and two pairs of similar-sized wings for species such as the Trichoptera genus *Polycentropus* (k), the

Megaloptera genus *Sialis (l)* or the Odonata genera *Ischnura* (m) and *Crocothemis* (n). Wings range in

size from a few mm in some Diptera (g) up to more than 3 cm (l-n), with the Odonata exemplifying

the large morphologies. Taxa vary in the number of eggs produced per individual, ranging from tens

per reproduction cycle for most Coleoptera and Heteroptera such as the genus *Sigara* (o) to several

hundreds in the egg masses of most Ephemeroptera and Trichoptera, such as those of the genus

Hydropsyche (p). Credits: Adolfo Cordero-Rivera (a-g, i, k-n), Jesús Arribas (h), Pere Bonada (j), José

Antonio Carbonell (o) and Maria Alp (p).

Figure 2. Total number of taxa and percentage of the nine traits completed in each insect order and

macroinvertebrate phylum, sub-phylum, class or sub-class. "Other" includes Hydrozoa,

Hymenoptera, Megaloptera and Porifera, for which the database includes only one genus.

Figure 3. Variability in the dispersal-related trait composition of all insect orders with complete trait
profiles along fuzzy correspondence analysis axes A1 and A2. Dots indicate taxa and lines converge to
the centroid of each order to depict within-group dispersion.

Dispersal strategy

Propensity to drift

Lifelong fecundity

Life cycle duration

Adult life span

Maximum body size

Potential number of reproductive cycles per year

Female wing length

Wing pair type

a

b

Figure 4. Trait category locations in the fuzzy correspondence analysis ordination space for each trait: (a) Dispersal strategy = **dis1**: aquatic active, **dis2**:

aquatic passive, **dis3**: aerial active, **dis4**: aerial passive; (b) Propensity to drift = **drift1**: rare/catastrophic, **drift2**: occasional, **drift3**: frequent; (c) Fecundity =

egg1: < 100 eggs, **egg2**: ≥ 100 − 1000 eggs, **egg3**: 1000 − 3000 eggs, **egg4**: ≥ 3000 eggs; (d) Life-cycle duration = **cd1**: ≤ 1 year, **cd2**: > 1 year; (e) Adult life span

= **life1**: < 1 week, **life2**: ≥ 1 week − 1 month, **life3**: ≥ 1 month − 1 year, **life4**: > 1 year; (f) Maximum body size (cm) = **s1**: < 0.25, **s2**: ≥ 0.25 − 0.5, **s3**: ≥ 0.5 − 1,

s4: ≥ 1 − 2; **s5**: ≥ 2 − 4, **s6**: ≥ 4 − 8; (g) Potential number of reproductive cycles per year = **cy1**: < 1, **cy2**: 1, **cy3**: > 1; (h) Female wing length (cm) = **fwl1**: < 5,

fwl2: ≥ 5 − 10, **fwl3**: ≥ 10 − 15, **fwl4**: ≥ 15 − 20, **fwl5**: ≥ 20 − 30, **fwl6**: ≥ 30 − 40, **fwl7**: ≥ 40 − 50, **fwl8**: ≥ 50; (i) Wing pair type = **Wbn2**: 1 pair + halters, **Wbn3**:

1 pair + elytra or hemielytra, **Wbn4**: 1 pair + small hind wings, **Wbn5**: 2 similar-sized pairs.

SUPPORTING INFORMATION

Supplementary File 1. List of references used to build DISPERSE.