**Next-generation phylogeography resolves post-glacial colonization patterns in a widespread carnivore, the red fox (*Vulpes vulpes*), in Europe.**

Allan D. McDevitt[1]*, Ilaria Coscia[1], Samuel S. Browett[1], Aritz Ruiz-González[2], Mark J. Statham[3], Inka Ruczyńska[4], Liam Roberts[1], Joanna Stojak[4], Alain C. Frantz[5], Karin Norén[6], Erik O. Ågren[7], Jane Learmount[8], Mafalda Basto[9], Carlos Fernandes[9], Peter Stuart[10], David G. Tosh[11], Magda Sindicic[12], Tibor Andreanszky[13], Marja Isomursu[14], Marek Panek[15], Andrey Korolev[16], Innokentiy M. Okhlopkov[17], Alexander P. Saveljev[18], Boštjan Pokorny[19], Katarina Flajšman[20], Stephen W. R. Harrison[21], Vladimir Lobkov[22], Duško Ćirović[23], Jacinta Mullins[4], Cino Pertoldi[24], Ettore Randi[24,25], Benjamin N. Sacks[3], Rafał Kowalczyk[4] and Jan M. Wójcik[4]*

[1]School of Science, Engineering and Environment, University of Salford, Salford, United Kingdom

[2]Department of Zoology and Animal Cell Biology, University of the Basque Country, UPV/EHU, Vitoria-Gasteiz, Spain

[3]Mammalian Ecology and Conservation Unit, Center for Veterinary Genetics, and Department of Population Health and Reproduction, School of Veterinary Medicine, University of California, Davis, USA

[4]Mammal Research Institute, Polish Academy of Sciences, Białowieża, Poland

[5]Musée National d'Histoire Naturelle, Luxembourg, Luxembourg

[6]Department of Zoology, Stockholm University, Stockholm, Sweden

[7]Department of Pathology and Wildlife Diseases, National Veterinary Institute, Uppsala, Sweden

[8]National Wildlife Management Centre, Animal and Plant Health Agency, Sand

Hutton, York, United Kingdom

[9]CE3C - Centre for Ecology, Evolution and Environmental Changes, Department of

Animal Biology, Faculty of Sciences, University of Lisbon, Lisbon, Portugal

[10]Biological and Pharmaceutical Sciences Department, Institute of Technology

Tralee, Kerry, Ireland

[11]National Museums of Northern Ireland, Hollywood, Northern Ireland UK

[12]Faculty of Veterinary Medicine, University of Zagreb, Zagreb, Croatia

[13]Croatian Veterinary Institute, Rijeka, Croatia

[14]Finnish Food Authority, Veterinary Bacteriology and Pathology Research Unit,

Oulu, Finland

[15]Polish Hunting Association, Czempiń, Poland

[16]Institute of Biology of Komi Science, Remote Centre of the Ural Branch of the

Russian Academy of Sciences, Syktyvkar, Komi Republic, Russia

[17]Institute of Biological Problems of Cryolithozone, Siberian Branch of Russian

Academy of Sciences, Yakutsk, Russia

[18]Department of Animal Ecology, Russian Research Institute of Game Management

and Fur Farming, Kirov, Russia

[19]Environmental Protection College, Velenje, Slovenia

[20]Slovenian Forestry Institute, Ljubljana, Slovenia

[21]School of Animal Rural & Environmental Sciences, Nottingham Trent University,

Southwell, UK

[22]Odessa I.I. Mechnykov National University, Faculty of Biology, Odessa, Ukraine

[23]Faculty of Biology, University of Belgrade, Belgrade, Serbia

[24]Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark

[25]Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna, Italy

*Correspondence: a.mcdevitt@salford.ac.uk; jwojcik@ibs.bialowieza.pl

**Abstract**

Carnivores tend to exhibit a lack of (or less pronounced) genetic structure at continental scales in both a geographic and temporal sense and this can confound the identification of post-glacial colonization patterns in this group. In this study we used genome-wide data (using Genotyping-by-Sequencing (GBS)) to reconstruct the phylogeographic history of a widespread carnivore, the red fox (*Vulpes vulpes*), by investigating broad-scale patterns of genomic variation, differentiation and admixture amongst contemporary populations in Europe. Using 15,003 single nucleotide polymorphisms (SNPs) from 524 individuals allowed us to identify the importance of refugial regions for the red fox in terms of endemism (e.g. Iberia). In addition, we tested multiple post-glacial re-colonization scenarios of previously glaciated regions during the Last Glacial Maximum using an Approximate Bayesian Computation (ABC) approach that were unresolved from previous studies. This allowed us to identify the role of admixture from multiple source population post-Younger Dryas in the case of Scandinavia and ancient land-bridges in the colonization of the British Isles. A natural colonization of Ireland was deemed more likely than an ancient human-mediated introduction as has previously been proposed and potentially points to an increased mammalian fauna on the island in the early post-glacial period. Using genome-wide data has allowed us to tease apart broad-scale patterns of structure and diversity in a widespread carnivore in Europe that was not evident from using more limited marker sets and provides a foundation for next-generation phylogeographic studies in other non-model species.

**Keywords**

81    Approximate Bayesian Computation; Genotyping-by-Sequencing; Ireland; Mammal;

82    Last Glacial Maximum; Scandinavia; SNP

## Introduction

Over the last 30 years, phylogeographic studies have highlighted the roles of major past climatic and geophysical events in shaping contemporary genetic structure and diversity in a multitude of species (Avise et al., 1987; Pedreschi et al., 2019; Stojak, Borowik, McDevitt, & Wójcik, 2019; Taberlet, Fumagalli, Wust-Saucy, & Cosson, 1998). During the Last Glacial Maximum (LGM; ~27-19 thousand years ago (kyrs BP; Clark et al., 2009)) many terrestrial plant and animal species retreated, and were often restricted, to refugial areas (Sommer & Nadachowski, 2006; Taberlet et al., 1998). In Europe, several overarching patterns have emerged from fossil data and phylogeographic studies, with refugia identified in the three 'classic' Mediterranean (Iberian, Apennine and Balkan) peninsulas (Hewitt, 1999; Sommer & Nadachowski, 2006; Taberlet et al., 1998), but also further north in areas in or adjacent to the Carpathians mountains and the Dordogne region in France as examples of 'cryptic refugia' (McDevitt et al., 2012; Provan & Bennett, 2008; Stojak et al., 2016). Phylogeographic studies of the most widely studied group, the terrestrial mammals, have shown distinct mitochondrial DNA (mtDNA) lineages in small mammals (Searle et al., 2009; Stojak et al., 2016; Vega et al., 2020) and ungulates (Carden et al., 2012; Sommer et al., 2008) that are consistent with contraction and re-expansion from these refugial regions (Hewitt, 1999; Taberlet et al., 1998).

Carnivores appear to be an exception to this general pattern however, with either a lack of, or less pronounced phylogeographic structure shown across continental scales (Frantz et al., 2014; Hofreiter et al., 2004; Korsten et al., 2009; Mucci et al., 2010). One such carnivore, the red fox (*Vulpes vulpes*), is well-represented in the fossil record in Europe (Sommer & Benecke, 2005) and has

numerous records during the LGM in recognized refugial areas such as the

Mediterranean peninsulas, and further north in areas in or adjacent to the Carpathian

mountains, and the Dordogne region in France (Sommer & Benecke, 2005; Sommer

& Nadachowski, 2006). Previous studies using mtDNA on modern and/or ancient

specimens have revealed a general lack of genetic structure on a continental-wide

scale, in both a geographic and temporal sense (Edwards et al., 2012; Kutschera et

al., 2013; Teacher et al., 2011). The lack of phylogeographic structuring in the red

fox and other carnivores has been previously attributed to these species persisting

outside the traditional refugial areas during the LGM, and effectively existing as a

large interbreeding population on a continental scale (Edwards et al., 2012; Teacher

et al., 2011). However, despite the abundant fossil data for red foxes, there is a

distinct lack of fossils from central Europe or further north during the LGM (Sommer

& Benecke, 2004, 2005).

More recent studies (Statham et al., 2014, 2018) identified mtDNA haplotypes

that were unique to particular regions (e.g. Iberia) that potentially indicate long-term

separation from other European populations. The concerns about the use of short

mtDNA sequences is that they may not fully capture the signals of post-glacial

colonization patterns in species with high dispersal capabilities (Keis et al., 2013;

Koblmüller et al., 2016). One solution is the utilisation of microsatellite markers in

conjunction with mtDNA data (Statham et al., 2018). However, several carnivore

species (e.g. badgers *Meles meles* and otters *Lutra lutra*) show a similar lack of

resolution in terms of broad-scale genetic structure across continental Europe using

microsatellites (Frantz et al., 2014; Mucci et al., 2010). For the red fox, several

distinct groups in Europe were identified using microsatellite markers from Bayesian

clustering analysis, with distinction between foxes in Ireland, Britain, Spain, Italy and

Scandinavia being apparent (Statham et al., 2018). The rapid mutation rate of

microsatellites leaves it unclear whether divisions reflect ancient isolation or more

recent population structure, owing to recently limited gene flow. This uncertainty and

lack of resolution in phylogeographic structure outside the 'traditional' (and well-

established) refugia means that inferring post-glacial colonization patterns of

previously glaciated regions during the LGM is challenging with more limited genetic

marker sets. This is particularly evident in the British Isles and Scandinavia where

these regions present differing problems in terms of how and when contemporary

populations of terrestrial species reached these areas in post-glacial periods (Carden

et al., 2012, 2020; Herman et al., 2014).

The island of Ireland has long presented a biogeographical quandary in terms

of how and when terrestrial species colonized it (Carden et al., 2012, 2020; McDevitt

et al., 2011). The area's latitude means that it was covered almost entirely by the ice

sheet during the LGM and it didn't become an island until approximately 15,000

years ago (almost twice as long as Britain; Edwards & Brooks, 2008). Because of

this, humans have been proposed as the primary mechanism of transport for its

mammalian fauna in ancient and modern times (Carden et al., 2012; Frantz et al.,

2014). An estimate of 10.2 kyrs BP was estimated for a split between Irish and

British red foxes using mtDNA data, but with a 95% CI range that incorporated the

possibility of natural colonization before Ireland became an island (Statham et al.,

2018). Haplotype diversity at different mtDNA markers is high in Irish red foxes

(Edwards et al., 2011; Statham et al., 2018) which would appear to contradict a more

recent, human-mediated origin that has been inferred from the only fossil dated to

the Bronze Age (3.8 kyrs BP; Sommer & Benecke, 2005). Indeed, natural

colonization of other species (e.g. stoat *Mustela erminea*) have been proposed over

post-glacial land bridges (Martínková, McDonald, & Searle, 2007) and a recent re-

evaluation of fossil dates has now placed multiple mammalian species on the island

close to the end of the LGM period (Carden et al., 2020). Therefore, a more thorough

investigation of the red fox's colonization history on the island has implications

beyond this species only.

Fennoscandia (present-day Sweden, Norway, Finland and Denmark) was

covered by a huge ice sheet during the LGM and would have been inhospitable to

most species until its complete retreat approximately 10,000 years ago (Patton et al.,

2017). However, some parts of southern Sweden and western Norway would have

had ice free regions during the later Younger Dryas Glaciation (12.9–11.7 kyrs BP)

and several temporary land-bridges connected these countries to Denmark between

approximately 13.1 and 9.2 kyrs BP (Herman et al., 2014; Marková et al., 2020).

Although glacial refugia in Scandinavia during the LGM have been proposed for

several species, including some mammals (Lagerholm et al., 2014; Westergaard et

al., 2019), the red fox does not appear in the fossil records in southern Scandinavia

until after ~9,000 yrs BP (Sommer & Benecke, 2005). Based on evidence from

mtDNA, microsatellites and Y chromosome data, multiple colonization events have

been proposed from the south and east for the red fox (Norén et al., 2015; Wallén et

al., 2018) but the progression of these events remain untested. Several mammalian

species have been proposed to have colonized from the south over temporary land-

bridges and persisted through the Younger Dryas before being supplemented with

further colonization wave(s) from the east (Herman et al., 2014; Marková et al.,

2020). It is unclear however if this is a frequently-used post-glacial colonization route

for many of the mammals that are currently present in the region.

The advent of next-generation sequencing technologies holds great promise

for phylogeographic studies, allowing for thousands of single nucleotide

polymorphisms (SNPs) to be genotyped in non-model organisms and providing a

representation of the organism's entire genome (Garrick et al., 2015; McCormack,

Hird, Zellmer, Carstens, & Brumfield, 2013). The use of reduced-representation

techniques (e.g. genotyping-by-sequencing, GBS) has already demonstrated their

potential in resolving phylogeographic patterns in non-model organisms that are not

fully captured with data with a limited number of genetic markers (Emerson et al.,

2010; Jeffries et al., 2016; Puckett et al., 2016; Marková et al., 2020). Using GBS

data from over 500 individuals, the purpose of the present study was to reconstruct

the phylogeographic history of the red fox in Europe by investigating broad-scale

patterns of genomic variation, differentiation, and admixture amongst contemporary

populations to investigate if contemporary red fox populations are the result of

isolation in particular refugia during the LGM (Statham et al., 2018); or if these

patterns likely emerged in the post-glacial period following the LGM (Edwards et al.,

2012; Teacher, Thomas, & Barnes, 2011). From there, we adopted an Approximate

Bayesian Computation (ABC; (Beaumont, Zhang, & Balding, 2002)) framework to

distinguish between multiple post-glacial re-colonization scenarios of previously

glaciated regions that were not fully resolved in previous studies using mtDNA, Y

chromosomal, and microsatellite markers (Statham et al., 2018; Wallén et al., 2018).

We first focused on the colonization of Ireland and Britain, and whether the red fox

colonized the island of Ireland naturally or was introduced later by humans (Statham

et al., 2018). Finally, we investigated the post-glacial colonization patterns of

northern Europe, with a focus on the mechanism and timing of the colonization of

Scandinavia around and after the Younger Dryas period (Wallén et al., 2018).

**Materials and Methods**

*Laboratory methods*

Red fox tissue samples were obtained from freshly culled (not directly related to this study), roadkill, frozen, ethanol- (70-95%) and DMSO-preserved material from previous studies. No material was collected for the purpose of the present study. Genomic DNA was extracted using the DNeasy Blood and Tissue Kit (Qiagen Ltd.) according to manufacturer's protocols (with the additional treatment of RNase). A total of 30–100 ng/μl of high molecular weight DNA was sent to the Genomic Diversity Facility in Cornell University (USA) where GBS was used for constructing reduced representation libraries (Elshire et al., 2011) using the restriction enzyme EcoT22I (ATGCAT) in six GBS libraries (each consisting of 95 uniquely barcoded individuals and one negative control). Individual ligations were then pooled and purified from adaptor excess using the QIAquick PCR Purification Kit (Qiagen). For library preparation genomic adaptor-ligated fragments were then PCR amplified in 50 μL volumes with 10 μL of DNA fragment pool, 1 × Taq Master Mix (New England Biolabs Inc.) and 12.5 pmol each of the following Illumina primers: 5´-AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT and 5´-CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCT (the underlined parts will hybridize to the two Illumina flowcell oligos). Temperature cycling consisted of 72 ☐C for 5 min, 98 ☐C for 30 s followed by 18 cycles of 98 ☐C for 30 s, 65 ☐C for 10 s, and 72 ☐C for 30 s, with a final extension step at 72 ☐C for 5 min. The EcoT22I GBS libraries (now containing ID tags and Illumina flowcell adaptors) were purified again using the QIAquick PCR Purification Kit (Qiagen). An aliquot was run on the BioAnalyzer™ 2100 to verify fragment sizes.

Library DNA was then quantified on a Nanodrop 2000 (ThermoFisher Scientific) and

subsequently sequenced on an Illumina HiSeq 2000 (Cornell University, Life

Sciences Core Facility).

*Bioinformatics*

The raw Illumina sequence data from 568 individuals were processed through the

GBS analysis pipeline implemented in TASSEL v5.2.31 (Glaubitz et al., 2014). Due

to concerns about the performances of *de novo* approaches to identify SNPs in

reduced representation genomic techniques (particularly for demographic analyses;

Shafer et al., 2017) the reference genome of the domestic dog (CanFam3.1; *Canis*

*lupus familiaris*) was used to align the sequence tags on individual chromosomes

using the BWA-backtrack method in the Burrows-Wheeler alignment tool (Li &

Durbin, 2009).

A total of unique 9,249,177 tags were found, with 6,267,895 tags (67.8%)

uniquely aligned to the dog genome. The rest of the tags were disregarded from

further analyses.  SNPs were initially filtered by removing those with a minor allele

frequency (MAF) of <0.01 and missing data per site of >0.9. This resulted in 144,745

SNPs with a mean individual depth of 15.853 (±SD = 4.629). After the removal of

indels, SNPs with an observed heterozygosity >0.5 (to filter out potential paralogs),

removing SNPs and individuals with call-rates of <0.8, SNPs with a MAF of <0.05,

and SNPs located on the X chromosome, a total of 19,795 bi-allelic SNPs and 524

individuals were retained after filtering in TASSEL v5.2.31 (Glaubitz et al., 2014) and

PLINK v1.07 (Purcell et al., 2007). Because of the potential for linkage disequilibrium

(LD) to bias the results of population genetic analyses, we pruned 3,571 SNPs with

LD of $r^2 > 0.2$ (Schweizer et al., 2016) in a window of 50 SNPs (sliding window with

five SNPs overlapping at a time) in PLINK (Purcell et al., 2007). To filter out SNPs

deviating from Hardy-Weinberg Equilibrium (HWE), individuals were grouped into 29

populations, consisting of seven or more individuals (see below). A least-squares $F_{IS}$

estimator (based on an AMOVA) was implemented in GenoDive v2.0b23 (Meirmans

& Van Tienderen, 2004) using 999 permutations. A total of 869 SNPs which deviated

from HWE ($p < 0.01$) in 15 or more populations (≥ 51.7%) were removed from

subsequent analyses (Van Wyngaarden et al., 2017).

Given that the focus of the study was to investigate demographic and

phylogeographic patterns, loci putatively under selection were identified using

Principal Component Analysis (PCA) and Mahalanobis distance implemented in the

R package *pcadapt* v3.0 (Luu, Bazin, & Blum, 2017). A false discovery rate (FDR) of

α = 0.05 was applied, with *q*-values smaller than α considered as candidate SNPs

under selection. A total of 352 SNPs were removed according to these criteria. Two

datasets were created for subsequent analyses; one containing all 524 individuals

(*524dataset*; Table S1), and the other containing 494 individuals (*494dataset*; Tables

1 and S1), consisting of 29 'populations' with seven or more individuals for

population-level analyses (Fig. 1A-C).

*Individual-based analyses*

Individual-based clustering analysis on the *524dataset* was conducted in

fastSTRUCTURE v1.0 (Raj, Stephens, & Pritchard, 2014). fastSTRUCTURE was

performed using the simple prior with *K* values of 1–30 over five independent runs.

The number of clusters (*K*) was obtained by using the 'chooseK.py' function on each

of these independent runs. Visualization of individual assignments to clusters per

population was initially performed using the 'distruct.py' function, with the final figure

produced using a custom R script. In our second individual-based analysis on the

*524dataset*, we incorporated the capture or sampling location of each individual,

using the spatially-explicit software *TESS3* (Caye, Deist, Martins, Michel, & François,

2016) The default values of the program were implemented and each run was

replicated five times. The optimal value of *K* corresponded to the minimum of the

cross-entropy criterion, across values of *K* = 1–30.

*Population-based analyses*

For the *494datset*, two measures of genomic diversity, allelic richness (AR) and

expected heterozygosity ($H_E$), were calculated in GENODIVE (Meirmans & Van

Tienderen, 2004) and HP-RARE (Kalinowski, 2005). AR and $H_E$ values were

mapped with interpolation using ArcGIS 10.2.1. Geostatistical Analyst. One

population from Siberia (Russia) was excluded from this analysis because it was

geographically distant from all the other populations. Interpolation was carried out

using an Inverse Distance Weight model (IDW, power=1, based on 12 neighbours;

Stojak et al., 2016). Discriminant Analysis of Principal Components (DAPC) as

implemented in the R package *adegenet* v2.0.1 (Jombart & Ahmed, 2011) was

performed on the *494dataset*. DAPC does not make assumptions about Hardy-

Weinberg equilibrium or linkage disequilibrium and provides a graphical

representation of the divergence among pre-defined populations/groups.

*Phylogeographic reconstruction*

Approximate Bayesian Computation was implemented in DIYABC Random Forest

v1.0 (Collin et al., 2021) to further investigate the dynamics of the re-colonization

process of red foxes in Europe. As the method is computationally intensive and has

generally required 100,000 to 1,000,000 simulations to distinguish between

demographic scenarios being tested, several studies have performed their ABC

analyses on subsets of their SNPs randomly selected from the full dataset (usually

~1,000 SNPs; e.g. Huang et al., 2017; Jeffries et al., 2016). However, recent

applications using a tree-based classification method known as 'random forest' in

ABC allow demographic scenarios to be distinguished based on 1,000s to 10,000s of

simulations for each scenario (Fraimout et al., 2017; Kotlík, Marková, Konczal,

Babik, & Searle, 2018; Pudlo et al., 2015; Marková et al., 2020). We followed the

approach of Kotlík et al., (2018) where all SNPs were used, but we chose a subset of

the individuals in each grouping to save on computational time (Table S1).

Based on recent simulation studies, ABC-based methods have received

criticism for their ability to capture the true demographic models under consideration

(Cabrera & Palsboll, 2017; Shafer, Gattepaille, Stewart, & Wolf, 2015). Following

recommendations by Cabrera & Palsboll (2017), scenarios were kept as simple and

different from each other as possible in order to distinguish between the major

demographic events under consideration. In addition, the number of comparable

scenarios was always kept low (Cabrera & Palsboll, 2017). Following on from

outstanding issues in regards to unresolved colonization scenarios (Statham et al.,

2018), we first investigated the colonization history of red foxes in the British Isles.

We grouped individuals into three 'populations' of interest to ascertain the most likely

scenario for the timing and source of existing populations in Ireland and Great Britain

based on the analyses performed in fastSTRUCTURE, DAPC (See Results) and previous studies. These were 'Ireland' (IRE + NIR populations), 'Britain' (UKS + UKN) and 'Europe' (FRA + BEL + SWZ). Three scenarios were incorporated that the data and analysis presented by Statham et al. (2018) could not distinguish between. The first is that Ireland was colonized naturally overland from an unsampled ancestral population (N4) after the LGM (i.e. before Ireland became an island between approximately 19,000 and 15,000 yrs BP; t2; Fig. S1). In this scenario, Britain originated from this ancestral population also but mixes with the European population before it became isolated from the mainland after the flooding of Doggerland around 8 kyrs BP (19–8 kyrs BP; ta). Europe diverged from the unsampled ancestral population at 19–15 kyrs BP with t1>t2. In the second scenario, Ireland's red foxes were founded directly from Britain naturally before it became an island (t2). In the third scenario, Ireland's foxes were transported by humans from Britain after the earliest evidence of human presence on the island at 12,700 yrs BP (Dowd & Carden, 2016) right up to the present day (t3; Fig. S1). Effective populations sizes were allowed to range between 10 and 500,000 individuals for Irish and British populations and between 10 and 1,000,000 for Europe and the unsampled ancestral population. Generation time was assumed to be 2 years (Statham et al., 2018).

For the second ABC-based analysis, we investigated the colonization history of Scandinavia. Wallen et al. (Wallén et al., 2018) proposed that Scandinavia was colonized from multiple sources based on mtDNA and Y chromosome data but did not attempt to date the progression of these events. We attempted to distinguish between five scenarios for the timing and progression of these colonization events.

In the first of these, 'Scandinavia' (FNS + FNN + SWN + SWS + NOR) was the result

of admixture in the east and subsequent colonization (9,000 yrs BP to present; tb)

between 'Central Europe' (PLA + PLC + PLS + GER) and 'Russia' (RUK + RUV)

populations after the region became ice-free after the Younger Dryas Glaciation and

disappearance of land-bridges connecting it to central Europe (Herman et al., 2014).

The Central Europe and Russian populations split at t1 (27,000–19,000 yrs BP). In

the second scenario, the first colonization wave occurred from central Europe over

land-bridges (14,000 to 9,000 yrs BP; ta), with later admixture occurring from Russia

(9,000 yrs BP to present; tb). In the third scenario, the first colonization wave

occurred from Russia (9,000 yrs BP to present; tb), with later admixture from Central

Europe from an eastern route (9,000 yrs BP to present but restricted to after the first

Russian colonization wave; tb >tc). In the fourth scenario, Russia was the result of

admixture between Scandinavia and Central Europe at tb. Finally, the fifth scenario

had all three populations diverging at t1. Effective populations sizes were allowed to

range between 10 and 1,000,000 individuals.

Ten thousand simulated datasets per scenario were used to produce posterior

distributions. Each scenario was considered to be equally probable at the outset. To

check the reliability of the observed summary statistics, a Principal Component

Analysis (PCA) was performed on the summary statistics from the simulated

datasets and compared against the summary statistics from the observed dataset in

order to evaluate how the latter is surrounded by the simulated datasets (Collin et al.,

2021). All simulated datasets were used in each Random Forest training set. We

used five noise variables and ran 1,000 Random Forest trees to select the most

likely scenario and estimate parameters (Collin et al., 2021).

**Results**

*Individual-based analyses*

fastSTRUCTURE identified $K = 7$ as the lower limit of clusters in each of the five independent runs of $K = 1$–30 (Fig. 1C), with the upper limit of $K$ fluctuating from 10-13 between runs. Focusing first on $K = 7$, distinct clusters were identified in each of Ireland ('Ireland') and Great Britain ('Britain'). Iberian populations formed a distinct cluster ('Iberia'), a known glacial refugium. Populations in France, Switzerland, Belgium, Germany, Poland, Slovenia, Croatia, Serbia and the Ukraine formed a distinct cluster ('Central Europe', named for simplicity because of the approximate location of the cluster relative to the other clusters). Several of these populations are in proximity to the Balkan and Carpathian glacial refugia. Localities with small numbers of individuals in Lithuania, Estonia, Belarus and western Russia also belonged to this cluster. These populations in eastern Europe showed evidence of mixed ancestry with individuals from Scandinavia (Figs. 1A and 1C), who formed another distinct cluster ('Scandinavia'). Individuals from European Russia had mixed ancestry between this Scandinavian cluster and individuals from Siberia (another distinct cluster; 'Siberia'). Finally, individuals from central Italy formed a distinct cluster ('Italy'), another well-defined glacial refugium. Individuals in northern Italy showing mixed ancestry with the central European cluster (Figs. 1A and 1C). For the lower values of $K$, the islands of Ireland and Britain were separated from all other samples at $K = 2$. Scandinavia and Russia samples were further separated at $K = 3$, Iberia samples at $K = 4$, Siberia at $K = 5$ and Ireland and Britain each becoming distinct clusters at $K = 6$ (Fig. 2). For $K = 8$, an additional cluster with the European Russian populations was identified. For $K = 9$, another cluster consisting of French, Belgian and Swiss populations was found (with mixed ancestry from Central Europe;

Fig. 2). These populations are in close proximity to the known glacial refugium

located in the Dordogne region. Further admixture was identified within populations

in Central Europe at $K$ = 10–13 (data not shown). The spatially-explicit software

*TESS3* (Caye et al., 2016) failed to resolve the genomic structure within the system,

with the minimum of the cross-entropy criterion decreasing up to $K$ = 30 (Fig. S4).

Further test runs increasing $K$ up to a maximum of 40 failed to resolve the genomic

structure (data not shown).

*Population-based analyses*

Direct estimates of genomic diversity (Table 1) and the IDW interpolation of allelic

richness (AR) and expected heterozygosity ($H_E$) in 28 fox populations showed that

diversity is highest in Central and Eastern Europe and decreases westwards and

northwards (Fig. 2). Genomic diversity was notably lower in the British Isles, with the

Irish populations showing the lowest levels of diversity (Fig. 2). The DAPC revealed

distinct groupings of Iberian samples, Irish samples, British samples, Siberian and

Scandinavia/Russian samples in general agreement with the individual-based

analysis in fastSTRUCTURE (Fig. 1B). Populations in western, central and eastern

Europe were grouped closely together, but the populations in France, Belgium and

Switzerland were more separated from the main European group on the first axis,

aligned with the individual-based Bayesian analyses at $K$ = 9 (Fig. 2). Although the

population in central Italy formed its own genomic cluster in the individual-based

analysis (Fig. 1C), it grouped more closely with the central European group than the

French, Belgian and Swiss populations in this analysis (Fig. 1B).

For the ABC-based analysis, first focusing on the colonization history of

Ireland and Britain, Scenario 1 (Fig. S1) was selected (620/1,000 trees) with the

posterior probability estimated at 0.741. The ancestral population for Ireland and

Britain was estimated to have split from the European mainland at 17.7 kyrs BP

(95% CI: 15.81–18.95 kyrs BP), with the Irish population originating at 16.1 kyrs BP

(95% CI: 15.04–18.1 kyrs BP; Fig. 4; Table S2). For Scandinavia, Scenario 1

(510/1000 trees) was selected with the posterior probability estimated at 0.66. The

initial split of the Central European and Russian populations was estimated at 23.04

kyrs BP (95% CI: 19.34–26.55 kyrs BP). The current Scandinavian originates from

admixture from Russian and Central European populations at 5.36 kyrs BP (95% CI:

1.83–8.75 kyrs BP; Fig. 4; Table S2).

**Discussion**

In this study, we provided a genome-wide assessment of population structure and diversity in the red fox in Europe. By incorporating over 15,000 SNPs and over 500 individuals, we were able to advance previous work by investigating broad-scale patterns of structure and variation to identify putative glacial refugia and post-glacial re-colonization patterns in this widespread species.

*Phylogeographic structure of the red fox in Europe*

Individual- and population-based analyses revealed congruent patterns of genomic structuring at the broad scale of Europe (and Siberia), with certain important nuances being revealed by different approaches (Figs. 1, 2, 3 and S4). Earlier studies had proposed that red foxes may have existed as a single, large panmictic population during the LGM based on a lack of distinct structure at mtDNA markers using modern and/or ancient DNA (Edwards et al., 2012; Teacher et al., 2011). If this was the case, we might have expected to find a more continuous population (excluding the islands potentially) and this is not evidenced here with this greatly expanded dataset in terms of genetic markers, individuals and spatial coverage. In addition, a continuous population over the whole continent at the LGM is not generally congruent with the fossil data and the lack of fossil records beyond the more accepted refugial regions (Sommer & Benecke, 2005; Sommer & Nadachowski, 2006). Our study demonstrates that the observed patterns of genomic variation in contemporary red fox in Europe were mainly shaped by distinct refugial populations, with subsequent post-glacial admixture and isolation when this species had expanded into what is now its current distribution range in Europe (Sommer & Benecke, 2005).

At lower values of $K$, the islands of Ireland and Britain are separated from the rest of the European continent. Scandinavia/Russia and Iberia are then separated from the other continental individuals at $K = 3$ and 4, respectively (Fig. 2). Most of the central European (defined here as those outside of the Mediterranean peninsulas and Scandinavia) and Balkan populations formed a single genomic cluster at $K = 7$ (Figs. 1A and 1C). This and the elevated values of genomic diversity (Fig. 3) are likely reflective of more widespread and connected populations occupying the Balkans and Carpathian refugia during the LGM (as is known from fossil records; Sommer & Nadachowski, 2006) and a subsequent expansion into the rest of central Europe in the post-glacial period. A similar scenario has been proposed for other large mammals (Frantz et al., 2014; Stojak & Tarnowska, 2019). At $K = 7$, French, Belgian and Swiss individuals were grouped with other central European populations but population-level analyses (DAPC; Fig. 1B and 3) showed that these populations were distinct from other populations in close proximity (and they formed their own cluster at $K = 9$ in the individual-based Bayesian analysis; Fig. 2). Fossil records of the red fox are known from the Dordogne region in France during the LGM (Sommer & Nadachowski, 2006) so these populations may stem from a previously isolated refugial population in the area, and now show post-glacial admixture with populations stemming from eastern/Balkan and Iberian refugia (Figs. 1B and 1C). The Iberian populations form a distinct cluster/group at $K = 4$ and above (Figs. 1B, 1C and 2) and this is in line with previous findings using fewer molecular markers identifying this as a glacial refugium. Statham et al. (2018) identified mtDNA haplotypes that were endemic to the region, while microsatellites identified Spanish individuals as being distinct from those in other European populations. A similar pattern was found

previously in badgers, with Iberian populations having many unique mtDNA haplotypes not found elsewhere on the continent (Frantz et al., 2014). The Pyrenees Mountains have remained a formidable barrier for post-glacial re-colonization, and there appears to be little contribution to subsequent northwards expansion when the ice-sheets receded for many terrestrial species (Bilton et al., 1998). Even though the maximum dispersal capabilities of the red fox are up to 1,000 km in Europe (Walton, Samelius, Odden, & Willebrand, 2018), this mirrors the pattern of mountains acting as significant barriers for the species in North America (Sacks, Statham, Perrine, Wisely, & Aubry, 2010). This is in contrast to red foxes in the other Mediterranean refugium, Apennine (Italy). Although red foxes from central Italy are identified as a distinct cluster in fastSTRUCTURE at $K$ = 7 and above, mixed ancestry was identified with neighbouring populations north of the Alps in central Europe and the Balkans (Figs. 1B, 1C and 2). This may rather reflect a more recent divergence of this population given the geography of the region.

Glaciated regions during the LGM such as the British Isles and Scandinavia present differing problems in terms of how contemporary populations of terrestrial species colonized these areas in post-glacial periods. Ireland has existed as an island for approximately 15,000 years (Edwards & Brooks, 2008) and humans have been proposed as the primary mechanism of transport for its mammalian fauna based on existing fossil data and numerous phylogeographic studies (e.g. McDevitt et al., 2011; Carden et al., 2012; Frantz et al., 2014). Statham et al. (2018) proposed a split of approximately 10 kyrs between Irish and British foxes but with confidence intervals that didn't distinguish between natural and human-mediated colonization scenarios. Both Irish and British populations are distinct from mainland European

populations (Figs. 1B, 1C and 2) and have patterns of diversity and structure consistent with colonization and subsequent isolation (Table 1; Figs. 1B and 2). Using an ABC-based approach, a scenario in which Ireland was colonized before humans were known to be present (approximately 15 kyrs BP) was deemed the most likely (Fig. 4; Table S3). Although this conflicts with the current fossil evidence where the oldest known specimen in Ireland is from the Bronze Age (Sommer & Benecke, 2005), this is congruent with previous studies demonstrating high haplotype diversity and the identification of many unique haplotypes at mitochondrial markers on the island in comparison to British and other European populations (Edwards et al., 2012; Statham et al., 2018). The stoat was also proposed to be an early colonizer of Ireland over a post-glacial land bridge (Martínková, McDonald, & Searle, 2007) and several potential prey species (e.g. mountain hare *Lepus timidus* and arctic lemming *Dicrostonyx torquatus*; Woodman et al., 1997) were also present in the early post-glacial period. A recent re-assessment of several Irish mammalian fossils has pushed several species (reindeer *Rangifer tarandus*, grey wolf *Canis lupus* and woolly mammoth *Mammathus primigenius*) to the LGM/post-glacial boundary (Carden et al., 2020). Given that models of glaciation/de-glaciation patterns are reliant on secure and accurate fossil data, the island may have hosted a larger mammalian community in the early post-glacial period and this may have included the red fox also. While humans were still likely an important factor in determining later faunal assemblages on Ireland (Carden et al., 2012; Frantz et al., 2014), the early post-glacial period clearly warrants further investigation on the island based on the results presented here and newly available fossil data (Dowd & Carden, 2016; Carden et al., 2020).

Although glacial refugia in Scandinavia during the LGM have been proposed

for several species including mammals (Lagerholm et al., 2014; Westergaard et al.,

2019), the red fox does not appear in the fossil records in southern Scandinavia until

much later (~9,000 yrs BP; Sommer & Benecke, 2005). Here, we examined five

different hypotheses for the colonization of Scandinavia (Fig. S2). The most likely

scenario for the colonization of Scandinavia is a mixture of foxes from Russian and

central/eastern Europe colonizing from the east (Fig. 4). When the ice retreated from

northern Scandinavia after the Younger Dryas, a lack of geographic barriers led to

later dispersal into the region from the east (Norén et al., 2015; Wallén et al., 2018),

a pattern that is evident in other carnivores also (Dufresnes et al., 2018; Keis et al.,

2013). Several recent studies have pointed to a pattern of post-glacial colonization of

Scandinavia from the south involving individuals from present-day central Europe

first crossing temporary land-bridges prior to the Younger Dryas glaciation. This

pattern is particularly prevalent in small mammals such as the field vole *Microtus*

*agrestis* (Herman et al., 2014) and the bank vole *Myodes glareolus* (Marková et al.

2020). From all the analyses presented here, this scenario is not supported for the

red fox and it is instead apparent that this came later from the east only (Fig. 4). With

Scandinavia being one of the last regions of the continent to be re-colonized, the

higher genomic diversity observed here than some of the more westerly regions

could be due to it being populated from multiple sources from the east (Wallen et al.,

2018; Marková et al., 2020).

Using genome-wide data has allowed us to tease apart broad-scale patterns

of structure and diversity in a widespread carnivore in Europe that was not evident

from more limited marker sets. The use of genomic data allowed us to identify the

importance of refugial regions in terms of both endemism (e.g. Iberia) and sources of

post-glacial re-expansion across the continent (e.g. the Carpathians and Balkans). In

conjunction with ABC-based analyses, we identified patterns of post-glacial

colonization in formerly glaciated regions that contradict previously proposed routes

for the red fox and other similar species. Given the genomic resources now available

(Kukekova et al., 2018), the application of ancient genomics on the extensive fossil

material available for this species (Sommer & Benecke, 2005) should fall into line

with other charismatic carnivores (Loog et al., 2020) to fully understand re-

colonization and temporal patterns that have not been captured in previous studies

of ancient red fox specimens (Edwards et al., 2012; Teacher et al., 2011).

**Acknowledgements**

This study was financed by the National Science Centre, Poland, grant no. DEC-

2012/05/B/NZ8/00976 awarded to JMW, ADM, RK, ER and CP. We thank the

following people for supplying samples: Heikki Ahola, Peter Allason, Evidio Bartolini,

Lucia Burrini, Benoit Combes, Dorothee Ehric, Teresa García Díez, Vaclavas

Gedminas, Christian Gortázar Schmidt, Rebecca Hari, U.A. Kalisnikov, Marta

Kołodziej-Sobocińska, Nikolay Korablev, Antonio Lavazza, Sandro Lovari, Luciano

Palazzi, Giorgia Romeo, Marie-Pierre Ryser-Degiorgis, Ivan Seryodkin, Aleksandr

Sokolov and Pavel Veligurov. ADM thanks Robert Sommer, Norbert Benecke and

Ruth Carden for information on, and access to, their red fox fossil data, and to Petr

Kotlik for advice on ABC analyses. We sincerely thank the three anonymous

reviewers for their comments which significantly improved the manuscript.

## References

Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E., … Saunders, N. C. (1987). Intraspecific Phylogeography: The Mitochondrial DNA Bridge Between Population Genetics and Systematics. *Annual Review of Ecology and Systematics*, *18*(1), 489–522. doi:10.1146/annurev.es.18.110187.002421

Beaumont, M., Zhang, W., & Balding, D. (2002). Approximate Bayesian computation in population genetics. *Genetics*, *162*, 2025–2035.

Bilton, D. T., Mirol, P. M., Mascheretti, S., Fredga, K., Zima, J., & Searle, J. B. (1998). Mediterranean Europe as an area of endemism for small mammals rather than a source for northwards postglacial colonization. *Proceedings of the Royal Society B: Biological Sciences*, *265*(1402), 1219–1226. doi:10.1098/rspb.1998.0423

Cabrera, A. A., & Palsboll, P. J. (2017). Inferring past demographic changes from contemporary genetic data: a simulation-based evaluation of the ABC methods implemented in DIYABC. *Molecular Ecology Resources*, *17*, e94-e110. doi:10.1111/1755-0998.12696

Carden, R. F., McDevitt, A. D., Zachos, F. E., Woodman, P. C., O'Toole, P., Rose, H., … Edwards, C. J. (2012). Phylogeographic, ancient DNA, fossil and morphometric analyses reveal ancient and modern introductions of a large mammal: The complex case of red deer (*Cervus elaphus*) in Ireland. *Quaternary Science Reviews*, *42*, 74–84. doi:10.1016/j.quascirev.2012.02.012

Carden, R. F., Higham, T. F. G., Woodman, P. C. (2020) A reconsideration of the radiocarbon dating of the Marine Isotope Stage 3 fauna from southern Ireland. *Boreas, 49*(3), 674-684.

Caye, K., Deist, T. M., Martins, H., Michel, O., & François, O. (2016). TESS3: fast inference of spatial population structure and genome scans for selection. *Molecular Ecology Resources*, *16*(2), 540–548. doi:10.1111/1755-0998.12471

Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., … McCabe, A. M. (2009). The Last Glacial Maximum. *Science*, *325*(5941), 710-714.

Collin, F-D., Durif, G., Raynal, L., Lombaert, E., Gautier, M., Vitalis, R. Marin, J-M., Estoup, A. (2021) Extending approximate Bayesian computation with supervised machine learning to infer demographic history from genetic polymorphisms using DIYABC Random Forest. *Molecular Ecology Resources,* doi: 10.1111/1755-0998.13413.

Cornuet, J.-M., Ravignie, V., & Estoup, A. (2010). Inference on population history and model checking using DNA sequence and microsatellite data with the software DIYABC (v1.0). *BMC Bioinformatics*, *11*(1), 401. doi:10.1186/1471-2105-11-401

Dowd, M., & Carden, R. F. (2016). First evidence of a Late Upper Palaeolithic human presence in Ireland. *Quaternary Science Reviews*, *139*, 158–163. doi:10.1016/j.quascirev.2016.02.029

Dufresnes, C., Miquel, C., Remollino, N., Biollaz, F., Salamin, N., Taberlet, P., & Fumagalli, L. (2018). Howling from the past: Historical phylogeography and diversity losses in European grey wolves. *Proceedings of the Royal Society B: Biological Sciences, 285*(1884). doi:10.1098/rspb.2018.1148

Edwards, C. J., Soulsbury, C. D., Statham, M. J., Ho, S. Y. W., Wall, D., Dolf, G., … Bradley, D. G. (2012). Temporal genetic variation of the red fox, *Vulpes vulpes*, across western Europe and the British Isles. *Quaternary Science Reviews*, *57*,

95–104. doi:10.1016/j.quascirev.2012.10.010

Edwards, R., & Brooks, A. (2008). The Island of Ireland: Drowning the Myth of an Irish Land-bridge? *Mind the Gap: Postglacial Colonisation of Ireland.*, 19–34.

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell, S. E. (2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. *PLoS ONE 6,* e19379. doi:10.1371/journal.pone.0019379

Emerson, K. J., Merz, C. R., Catchen, J. M., Hohenlohe, P. a, Cresko, W. a, Bradshaw, W. E., & Holzapfel, C. M. (2010). Resolving postglacial phylogeography using high-throughput sequencing. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 16196–16200. doi:10.1073/pnas.1006538107

Fraimout, A., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., … Nielsen, R. (2017). Deciphering the routes of invasion of Drosophila suzukii by Means of ABC Random Forest. *Molecular Biology and Evolution*, *34*(4), 980–996. doi:10.1093/molbev/msx050

Frantz, A. C., McDevitt, A. D., Pope, L. C., Kochan, J., Davison, J., Clements, C. F., … Burke, T. (2014). Revisiting the phylogeography and demography of European badgers (*Meles meles*) based on broad sampling, multiple markers and simulations. *Heredity*, *113*(5), 443–53. doi:10.1038/hdy.2014.45

Garrick, R. C., Bonatelli, I. A. S., Hyseni, C., Morales, A., Pelletier, T. A., Perez, M. F., … Carstens, B. C. (2015). The evolution of phylogeographic data sets. *Molecular Ecology*, *24,* 1164-1171. doi:10.1111/mec.13108

Glaubitz, J. C., Casstevens, T. M., Lu, F., Harriman, J., Elshire, R. J., Sun, Q., & Buckler, E. S. (2014). TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline. *PLoS ONE, 9,* e90346. doi:10.1371/journal.pone.0090346

Herman, J. S., McDevitt, A. D., Kawałko, A., Jaarola, M., Wójcik, J. M., & Searle, J. B. (2014). Land-bridge calibration of molecular clocks and the post-glacial Colonization of Scandinavia by the Eurasian field vole *Microtus agrestis*. *PLoS ONE*, *9*(8), e103949. doi:10.1371/journal.pone.0103949

Hewitt, G. (1999). Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society*, *68*(1–2), 87–112. doi:10.1111/j.1095-8312.1999.tb01160.x

Hofreiter, M., Serre, D., Rohland, N., Rabeder, G., Nagel, D., Conard, N., … Pääbo, S. (2004). Lack of phylogeography in European mammals before the last glaciation. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(35), 12963–12968. doi:10.1073/pnas.0403618101

Huang, Z., Young, N. D., Reagon, M., Hyma, K. E., Olsen, K. M., Jia, Y., & Caicedo, A. L. (2017). All roads lead to weediness: Patterns of genomic divergence reveal extensive recurrent weedy rice origins from South Asian *Oryza*. *Molecular Ecology*, *26*(12), 3151–3167. doi:10.1111/mec.14120

Jeffries, D. L., Copp, G. H., Lawson Handley, L., Olsén, K. H., Sayer, C. D., & Hänfling, B. (2016). Comparing RADseq and microsatellites to infer complex phylogeographic patterns, an empirical perspective in the Crucian carp, *Carassius carassius,* L. *Molecular Ecology*, *25*(13), 2997–3018. doi:10.1111/mec.13613

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics*, *27*(21), 3070–3071. doi:10.1093/bioinformatics/btr521

Kalinowski, S. T. (2005). hp-rare 1.0: a computer program for performing rarefaction on measures of allelic richness. *Molecular Ecology Notes*, *5*(1), 187–189. doi:10.1111/j.1471-8286.2004.00845.x

Keis, M., Remm, J., Ho, S. Y. W., Davison, J., Tammeleht, E., Tumanov, I. L., … Saarma, U. (2013). Complete mitochondrial genomes and a novel spatial genetic method reveal cryptic phylogeographical structure and migration patterns among brown bears in north-western Eurasia. *Journal of Biogeography*, *40*(5), 915–927. doi:10.1111/jbi.12043

Koblmüller, S., Vilà, C., Lorente-Galdos, B., Dabad, M., Ramirez, O., Marques-Bonet, T., … Leonard, J. A. (2016). Whole mitochondrial genomes illuminate ancient intercontinental dispersals of grey wolves (*Canis lupus*). *Journal of Biogeography*, *43*(9), 1728–1738. doi:10.1111/jbi.12765

Korsten, M., Ho, S. Y. W., Davison, J., PÄhn, B., Vulla, E., Roht, M., … Saarma, U. (2009). Sudden expansion of a single brown bear maternal lineage across northern continental Eurasia after the last ice age: A general demographic model for mammals? *Molecular Ecology*, *18*(9), 1963–1979. doi:10.1111/j.1365-294X.2009.04163.x

Kotlík, P., Marková, S., Konczal, M., Babik, W., & Searle, J. B. (2018). Genomics of end-pleistocene population replacement in a small mammal. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1872). doi:10.1098/rspb.2017.2624

Kukekova, A. V., Johnson, J. L., Xiang, X., Feng, S., Liu, S., Rando, H. M., … Zhang, G. (2018). Red fox genome assembly identifies genomic regions associated with tame and aggressive behaviours. *Nature Ecology and Evolution*, *2*(9), 1479–1491. doi:10.1038/s41559-018-0611-6

Kutschera, V. E., Lecomte, N., Janke, A., Selva, N., Sokolov, A. a, Haun, T., … Hailer, F. (2013). A range-wide synthesis and timeline for phylogeographic events in the red fox (*Vulpes vulpes*). *BMC Evolutionary Biology*, *13*(1), 114. doi:10.1186/1471-2148-13-114

Lagerholm, V. K., Sandoval-Castellanos, E., Ehrich, D., Abramson, N. I., Nadachowski, A., Kalthoff, D. C., … Dalén, L. (2014). On the origin of the Norwegian lemming. *Molecular Ecology*, *23*(8), 2060–2071. doi:10.1111/mec.12698

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, *25*(14), 1754–1760. doi:10.1093/bioinformatics/btp324

Loog, L., Thalmann, O., Sinding, M.-H. S., Schuenemann, V. J., Perri, A., Germonpré, M., … Manica, A. (2020). Ancient DNA suggests modern wolves trace their origin to a late Pleistocene expansion from Beringia. *Molecular Ecology*, *29*(9), 1596-1610. doi:10.1111/mec.15329.

Luu, K., Bazin, E., & Blum, M. G. B. (2017). *pcadapt*: an R package to perform genome scans for selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1), 67–77. doi:10.1111/1755-0998.12592

Marková, S., Horniková, M., Lanier, H. C., Henttonen, H., Searle, J. B., Weider, L. J., Kotlik, P. (2020). High genomic diversity in the bank vole at the northern apex of a range expansion: the role of multiple colonizations and end-glacial refigua. *Molecular Ecology*, *29*(9), 1730-1744.

Martínková, N., McDonald, R. a, & Searle, J. B. (2007). Stoats (Mustela erminea) provide evidence of natural overland colonization of Ireland. *Proceedings. Biological Sciences / The Royal Society*, *274*, 1387–1393. doi:10.1098/rspb.2007.0334

McCormack, J. E., Hird, S. M., Zellmer, A. J., Carstens, B. C., & Brumfield, R. T. (2013). Applications of next-generation sequencing to phylogeography and phylogenetics. *Molecular Phylogenetics and Evolution*, *66*(2), 526–538. doi:10.1016/j.ympev.2011.12.007

McDevitt, A. D., Vega, R., Rambau, R. V, Yannic, G., Herman, J. S., Hayden, T. J., & Searle, J. B. (2011). Colonization of Ireland: revisiting "the pygmy shrew syndrome" using mitochondrial, Y chromosomal and microsatellite markers. *Heredity*, *107*(6), 548–57. doi:10.1038/hdy.2011.41

McDevitt, A. D., Zub, K., Kawałko, A., Oliver, M. K., Herman, J. S., & Wójcik, J. M. (2012). Climate and refugial origin influence the mitochondrial lineage distribution of weasels (*Mustela nivalis*) in a phylogeographic suture zone. *Biological Journal of the Linnean Society*, *106*(1), 57–69. doi:10.1111/j.1095-8312.2012.01840.x

Meirmans, P. G., & Van Tienderen, P. H. (2004). GENOTYPE and GENODIVE: Two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes*, *4*(4), 792–794. doi:10.1111/j.1471-8286.2004.00770.x

Mucci, N., Arrendal, J., Ansorge, H., Bailey, M., Bodner, M., Delibes, M., … Randi, E. (2010). Genetic diversity and landscape genetic structure of otter (*Lutra lutra*) populations in Europe. *Conservation Genetics*, *11*(2), 583–599. doi:10.1007/s10592-010-0054-3

Norén, K., Statham, M. J., Ågren, E. O., Isomursu, M., Flagstad, Ø., Eide, N. E., … Sacks, B. N. (2015). Genetic footprints reveal geographic patterns of expansion in Fennoscandian red foxes. *Global Change Biology*, *21*(9), 3299–3312. doi:10.1111/gcb.12922

Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P., … Hall, A. M. (2017). Deglaciation of the Eurasian ice sheet complex. *Quaternary Science Reviews*, *169*, 148–172. doi:10.1016/j.quascirev.2017.05.019

Pedreschi, D., García-Rodríguez, O., Yannic, G., Cantarello, E., Diaz, A., Golicher, D., … Stewart, J. R. (2019). Challenging the European southern refugium hypothesis: Species-specific structures versus general patterns of genetic diversity and differentiation among small mammals. *Global Ecology and Biogeography*, *28*(2), 262–274. doi:10.1111/geb.12828

Pickrell, J. K., & Pritchard, J. K. (2012). Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. *PLoS Genetics*, *8*(11). doi:10.1371/journal.pgen.1002967

Provan, J., & Bennett, K. D. (2008). Phylogeographic insights into cryptic glacial refugia. *Trends in Ecology and Evolution*, *23*(10), 564–571. doi:10.1016/j.tree.2008.06.010

Puckett, E. E., Park, J., Combs, M., Blum, M. J., Bryant, J. E., Caccone, A., … Munshi-South, J. (2016). Global population divergence and admixture of the brown rat (*Rattus norvegicus*). *Proceedings of the Royal Society B: Biological Sciences*, *283*(1841). doi:10.1098/rspb.2016.1762

Pudlo, P., Marin, J. M., Estoup, A., Cornuet, J. M., Gautier, M., & Robert, C. P. (2015). Reliable ABC model choice via random forests. *Bioinformatics*, *32*(6), 859–866. doi:10.1093/bioinformatics/btv684

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., … Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*, *81*(3), 559–575. doi:10.1086/519795

Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: Variational Inference of Population Structure in Large SNP Data Sets. *Genetics*, *197*(2). Retrieved from http://www.genetics.org/content/197/2/573.short

Sacks, B. N., Statham, M. J., Perrine, J. D., Wisely, S. M., & Aubry, K. B. (2010). North American montane red foxes: Expansion, fragmentation, and the origin of the Sacramento Valley red fox. *Conservation Genetics*, *11*(4), 1523–1539. doi:10.1007/s10592-010-0053-4

Schweizer, R. M., vonHoldt, B. M., Harrigan, R., Knowles, J. C., Musiani, M., Coltman, D., … Wayne, R. K. (2016). Genetic subdivision and candidate genes under selection in North American grey wolves. *Molecular Ecology*, *25*(1), 380–402. doi:10.1111/mec.13364

Searle, J. B., Kotlík, P., Rambau, R. V, Marková, S., Herman, J. S., & McDevitt, A. D. (2009). The Celtic fringe of Britain: insights from small mammal phylogeography. *Proceedings. Biological Sciences / The Royal Society*, *276*(1677), 4287–94. doi:10.1098/rspb.2009.1422

Shafer, A. B. A., Gattepaille, L. M., Stewart, R. E. A., & Wolf, J. B. W. (2015). Demographic inferences using short-read genomic data in an approximate Bayesian computation framework: in silico evaluation of power, biases and proof of concept in Atlantic walrus. *Molecular Ecology*, *24*(2), 328–345. doi:10.1111/mec.13034

Shafer, A. B. A., Peart, C. R., Tusso, S., Maayan, I., Brelsford, A., Wheat, C. W., & Wolf, J. B. W. (2017). Bioinformatic processing of RAD-seq data dramatically impacts downstream population genetic inference. *Methods in Ecology and Evolution*, *8*(8), 907–917. doi:10.1111/2041-210X.12700

Sommer, R., & Benecke, N. (2005). Late-Pleistocene and early Holocene history of the canid fauna of Europe (Canidae). *Mammalian Biology*, *70*(4), 227–241. doi:10.1016/j.mambio.2004.12.001

Sommer, R. S., & Nadachowski, A. (2006). Glacial refugia of mammals in Europe: evidence from. *Mammal Review*, *36*(4), 251–265. doi:10.1111/j.1365-2907.2006.00093.x

Sommer, R. S., Zachos, F. E., Street, M., Jöris, O., Skog, A., & Benecke, N. (2008). Late Quaternary distribution dynamics and phylogeography of the red deer (Cervus elaphus) in Europe. *Quaternary Science Reviews*, *27*(7–8), 714–733. doi:10.1016/j.quascirev.2007.11.016

Sommer, Robert, & Benecke, N. (2004). Late- and Post-Glacial history of the Mustelidae in Europe. *Mammal Review*, *34*(4), 249–284. doi:10.1111/j.1365-2907.2004.00043.x

Statham, M. J., Edwards, C. J., Norén, K., Soulsbury, C. D., & Sacks, B. N. (2018). Genetic analysis of European red foxes reveals multiple distinct peripheral populations and central continental admixture. *Quaternary Science Reviews*, *197*, 257–266. doi:10.1016/j.quascirev.2018.08.019

Statham, M. J., Murdoch, J., Janecka, J., Aubry, K. B., Edwards, C. J., Soulsbury, C. D., … Sacks, B. N. (2014). Range-wide multilocus phylogeography of the red fox reveals ancient continental divergence, minimal genomic exchange and distinct demographic histories. *Molecular Ecology*, *23* (19), 4813–4830. doi:10.1111/mec.12898

Stojak, J., Borowik, T., McDevitt, A. D., & Wójcik J. M. (2019). Climatic influences on the genetic structure and distribution of the common vole and field vole. *Mammal Research*, *64*, 19-29. doi: 10.1007/s13364-018-0395-8.

Stojak, J., McDevitt, A. D., Herman, J. S., Kryštufek, B., Uhlíková, J., Purger, J. J., …

Searle, J. B. (2016). Between the Balkans and the Baltic: Phylogeography of a Common Vole Mitochondrial DNA Lineage Limited to Central Europe. *PLOS ONE*, *11*(12), e0168621. doi:10.1371/journal.pone.0168621

Stojak, J., & Tarnowska, E. (2019). Polish suture zone as the goblet of truth in post-glacial history of mammals in Europe. *Mammal Research 64,* 463-475.

Taberlet, P., Fumagalli, L., Wust-Saucy, A. G., & Cosson, J. F. (1998). Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology*, *7*(4), 453–464. doi:10.1046/j.1365-294x.1998.00289.x

Teacher, A. G. F., Thomas, J. a, & Barnes, I. (2011). Modern and ancient red fox (*Vulpes vulpes*) in Europe show an unusual lack of geographical and temporal structuring, and differing responses within the carnivores to historical climatic change. *BMC Evolutionary Biology*, *11*(1), 214. doi:10.1186/1471-2148-11-214

Van Wyngaarden, M., Snelgrove, P. V. R., DiBacco, C., Hamilton, L. C., Rodríguez-Ezpeleta, N., Jeffery, N. W., … Bradbury, I. R. (2017). Identifying patterns of dispersal, connectivity and selection in the sea scallop, *Placopecten magellanicus,* using RADseq-derived SNPs. *Evolutionary Applications, 10*(1), 102–117. doi:10.1111/eva.12432

Vega, R., McDevitt, A. D., Stojak, J., Mishta, A., Wójcik, J. M., Kryštufek, B., & Searle, J. B. (2020). Phylogeographical structure of the pygmy shrew: revisiting the roles of southern and northern refugia in Europe. *Biological Journal of the Linnean Society*. doi:10.1093/biolinnean/blz209

Wallén, J., Statham, M. J., Ågren, E., Isomursu, M., Flagstad, Ø., Bjørneboe-berg, T., … Norén, K. (2018). Multiple recolonization routes towards the north: population history of the Fennoscandian red fox (*Vulpes vulpes*). *Biological Journal of the Linnean Society, 124,* 621–632.

Walton, Z., Samelius, G., Odden, M., & Willebrand, T. (2018). Long-distance dispersal in red foxes *Vulpes vulpes* revealed by GPS tracking. *European Journal of Wildlife Research*, *64*(6). doi:10.1007/s10344-018-1223-9

Westergaard, K. B., Zemp, N., Bruederle, L. P., Stenøien, H. K., Widmer, A., & Fior, S. (2019). Population genomic evidence for plant glacial survival in Scandinavia. *Molecular Ecology*, *28*(4), 818–832. doi:10.1111/mec.14994

Woodman, P., McCarthy, M., & Monaghan, N. (1997). The irish quaternary fauna project. *Quaternary Science Reviews*, *16*(2), 129–159. doi:10.1016/S0277-3791(96)00037-6

**Author contributions**

JMW, ADM, RK, JM, CP and ER conceived and acquired funding for the study. ADM, JMW, RK, MJS, BNS, CP and ER designed the study. MJS, ACF, KN, EÅ, JL, MB, CF, PS, DGT, MS, AG, MI, MP, AK, IMO, AS, BP, KF, VL, SWRH, DC, ER, BNS and RK contributed samples and data towards the study. IR performed the laboratory work. ADM, AR-G and IC generated the final SNP panel. ADM, IC, SSB, LR and JMW designed and performed the data analyses. ADM, JMW, IC, JS and SSB wrote the manuscript, with all authors contributing to edits and discussion.

**Data Accessibility**

Raw SNP data and datasets used for individual and population-level analyses are available on a private peer review link on Dryad (https://datadryad.org/stash/share/lxGNZUG315NeDDYqLZ4srLJGzEqw3U7LLfLztMpaI6E)

**Table 1.** Population-level analyses of 29 pre-defined red fox populations.

Populations are named by country of origin with an accompanying label (Pop ID) for

illustrative purposes (see Figure 1). Approximate geographic co-ordinates are given

in longitude and latitude. The number of individuals in each population is indicated by

*N* and population-level measures of allelic richness (AR) and expected

heterozygosity (He) are shown that were used to calculate the interpolation of

genomic diversity (see Figure 3).

| Population | Pop ID | Longitude | Latitude | *N* | AR | He |
|---|---|---|---|---|---|---|
| Ireland | IRE | 53.3927275 | -9.9262638 | 20 | 1.68 | 0.237 |
| Northern Ireland | NIR | 54.717149 | -6.219987 | 20 | 1.68 | 0.238 |
| England North | UKN | 54.47445462 | -2.680500725 | 17 | 1.73 | 0.254 |
| England South | UKS | 51.6839616 | 0.574959028 | 27 | 1.73 | 0.253 |
| Portugal North | PTN | 40.7115386 | -7.7495801 | 11 | 1.77 | 0.263 |
| Portugal South | PTS | 38.9648718 | -9.2992138 | 18 | 1.76 | 0.26 |
| Spain North | SPN | 42.829585 | -2.700338 | 29 | 1.79 | 0.268 |
| Spain South | SPS | 37.14280344 | -3.647460938 | 8 | 1.71 | 0.254 |
| France | FRA | 48.50605105 | 2.64715577 | 20 | 1.79 | 0.27 |
| Belgium | BEL | 50.589536 | 6.123152 | 19 | 1.81 | 0.274 |
| Switzerland | SWZ | 46.7633251 | 7.1564597 | 19 | 1.79 | 0.271 |
| Italy North | ITN | 45.926389 | 10.251667 | 7 | 1.74 | 0.269 |
| Italy | ITA | 43.32517768 | 11.31591797 | 20 | 1.73 | 0.254 |
| Germany | GER | 52.5200066 | 13.404954 | 28 | 1.81 | 0.273 |
| Slovenia | SOV | 46.31255967 | 15.50934385 | 24 | 1.78 | 0.265 |
| Croatia | CRO | 45.4082114 | 13.6589423 | 19 | 1.81 | 0.274 |
| Poland East | PLA | 53.82505665 | 23.3406601 | 16 | 1.82 | 0.274 |
| Poland South | PLS | 49.7801 | 20.2505 | 14 | 1.81 | 0.275 |
| Poland West | PLC | 52.117714 | 16.779255 | 19 | 1.82 | 0.275 |
| Serbia | SER | 45.83645405 | 20.43457031 | 18 | 1.81 | 0.271 |
| Ukraine | UKR | 47.725824 | 29.952977 | 9 | 1.8 | 0.274 |
| Sweden South | SWS | 57.95217685 | 13.93725559 | 17 | 1.74 | 0.256 |
| Sweden North | SWN | 64.3322949 | 20.28735358 | 16 | 1.77 | 0.263 |
| Norway | NOR | 70.26464 | 30.34619 | 14 | 1.77 | 0.265 |
| Finland North | FNN | 69.9059 | 27.02396 | 8 | 1.76 | 0.269 |
| Finland South | FNS | 60.46667 | 26.91667 | 12 | 1.76 | 0.259 |
| Russia Komi | RUK | 61.8 | 51.516667 | 10 | 1.76 | 0.262 |
| Russia Kirovskaja | RUV | 58.56861045 | 48.86787415 | 25 | 1.78 | 0.265 |
| Siberia | SIB | 135.774568 | 67.898166 | 10 | 1.6 | 0.211 |

**Figure 1.** Approximate locations of the studied populations (A) and the genomic clusters to which they have been assigned based on Discriminant Analysis of Principal Components (DAPC; B) and Bayesian analysis in *fastSTRUCTURE* at $K$ = 7 (C). The proportion of admixture in each population (A) is based on the ancestry coefficients determined in *fastSTRUCTURE* (C).

**Figure 2.** The proportion of admixture in each population based on the ancestry coefficients determined in *fastSTRUCTURE* for $K$ values 2–9.

**Figure 3.** Interpolation of allelic richness and expected heterozygosity in 28 red fox populations (Siberia was excluded in this analysis) in Europe. The interpolated values of both indices are presented in the maps in different colours on a low (blue) to high (red) scale according to the legends. Black circles indicate the population locations.

**Figure 4.** Graphical representation of the most likely post-glacial colonization scenarios for Ireland and Britain (A) and Scandinavia (B) inferred from Approximate Bayesian Computation.

926    Figure 1.

929　Figure 2.

933    Figure 3.

937    Figure 4.