A novel protein complex that regulates active DNA demethylation in *Arabidopsis*

Pan Liu[1,2]†, Wen-Feng Nie[1,3]†,*, Xiansong Xiong[1,2], Yuhua Wang[1], Yuwei Jiang[1,2], Pei Huang[1,2], Xueqiang Lin[1], Guochen Qin[1], Huan Huang[1], Qingfeng Niu[1], Jiamu Du[1], Zhaobo Lang[1], Rosa Lozano-Duran[1], and Jian-Kang Zhu[1,4]*

[1]Shanghai Center for Plant Stress Biology, Center of Excellence in Molecular Plant Sciences, Chinese Academy of Sciences (CAS), Shanghai 201602, China

[2]University of Chinese Academy of Sciences，No.19 (A) Yuquan Road, Shijingshan District, Beijing 100049, China

[3] Department of Horticulture, College of Horticulture and Plant Protection, Yangzhou University, Yangzhou, Jiangsu, 225009, China

[4]Department of Horticulture & Landscape Architecture, Purdue University, West Lafayette, IN 47906, USA

†These authors contributed equally

*Correspondence: jkzhu@psc.ac.cn; wfnie@yzu.edu.cn

Running title: A protein complex regulates active DNA demethylation
Pan Liu, liupan@sibs.ac.cn
Wen-Feng Nie, wfnie@yzu.edu.cn
Xiansong Xiong, xiongxiansong@sibs.ac.cn
Yuhua Wang, wangyuhua@sibs.ac.cn
Yuwei Jiang, ywjiang@sibs.ac.cn
Pei Huang, peihuang@psc.ac.cn
Xueqiang Lin, xueqianglin@foxmail.com
Guochen Qin，qinguochen@sibs.ac.cn
Huan Huang, ahuanghuan@163.com

Qingfeng Niu, qfniu@psc.ac.cn

Jiamu Du , dujm@sustech.edu.cn

Zhaobo Lang, zblang@sibs.ac.cn
Rosa Lozano-Duran, lozano-duran@sibs.ac.cn
Jian-Kang Zhu, jkzhu@sibs.ac.cn

SUMMARY

Active DNA demethylation is critical for altering DNA methylation patterns

and regulating gene expression. The 5-methylcytosine DNA

glycosylase/lyase ROS1 initiates a base excision repair pathway for active

DNA demethylation and is required for the prevention of DNA

hypermethylation at thousands of genomic regions in *Arabidopsis*. How

ROS1 is regulated and targeted to specific genomic regions is not well

understood. Here, we report the discovery of an *Arabidopsis* protein

complex that contains ROS1, regulates *ROS1* gene expression, and likely

targets the ROS1 protein to specific genomic regions. ROS1 physically

interacts with a WD40 domain protein (RWD40), which in turn interacts with

a methyl-DNA binding protein (RMB1) as well as with a zinc finger and

homeobox domain protein (RHD1). RMB1 binds to DNA that is methylated in

any sequence context, and this binding is necessary for its function *in vivo*.

Loss-of-function mutations in *RWD40*, *RMB1*, or *RHD1* cause DNA

hypermethylation at several tested genomic regions independently of the

known ROS1 regulator IDM1. Because the hypermethylated genomic

regions include the DNA methylation monitoring sequence in the *ROS1*

promoter, plants mutated in *RWD40*, *RMB1*, or *RHD1* show increased *ROS1*

expression. Importantly, ROS1 binding to the *ROS1* promoter requires

RWD40, RMB1, and RHD1, suggesting that this complex dictates ROS1

targeting to this locus. Our results demonstrate that ROS1 forms a protein

complex with RWD40, RMB1, and RHD1, and that this novel complex

regulates active DNA demethylation at several endogenous loci in

Arabidopsis.

Key words: DNA methylation, DNA demethylation, ROS1, WD40 domain,

methyl-DNA binding

Introduction

DNA methylation at the fifth position of the cytosine ring is important for gene

regulation, transposon silencing, and imprinting in plants and many other

eukaryotic organisms (Robertson 2005; Slotkin and Martienssen 2007; Law and

Jacobsen 2010; He et al. 2011). DNA methylation occurs in different sequence

contexts, i.e. CG, CHG, and CHH, where H represents A, C, or T. In plants, *de*

novo DNA methylation is mediated by the RNA-directed DNA methylation (RdDM)

pathway, which involves small interfering RNAs (siRNAs) and scaffold RNAs in

addition to an array of proteins (Law and Jacobsen 2010; Zhang et al. 2018). In

Arabidopsis, CG methylation is maintained by DNA METHYLTRANSFERASE 1

(MET1) (Finnegan and Dennis 1993), while CHG methylation is maintained by the

plant-specific DNA methyltransferase CHROMOMETHYLASE 3 (CMT3) (Cao and

Jacobsen 2002). Asymmetric CHH methylation is maintained by DOMAIN

REARRANGED METHYLTRANSFERASE 2 (DRM2) through the RdDM pathway

and by CHROMOMETHYLASE 2 (CMT2) (Haag and Pikaard 2011; Zemach et al.

2013).

Specific DNA methylation patterns are tightly regulated by DNA methylation and

demethylation pathways (Penterman et al. 2007; Hsieh et al. 2009; Zhu 2009;

Law and Jacobsen 2010; Furner and Matzke 2011; Matzke and Mosher 2014).

Passive DNA demethylation results from an absence of DNA methyltransferases

or a reduction in their activity, or from a shortage of methyl donors following DNA

replication (Zhang et al. 2018). In plants, active DNA demethylation is mediated

by 5-methylcytosine DNA glycosylases through a DNA base-excision repair

pathway (Choi et al. 2002; Gehring et al. 2006; Zhu et al. 2007; Zhu 2009). There

are four 5-methylcytosine DNA glycosylases in *Arabidopsis,* namely

REPRESSOR OF SILENCING 1 (ROS1), DEMETER (DME), DEMETER-LIKE 2

(DML2), and DEMETER-LIKE 3 (DML3) (Choi et al. 2002; Gong et al. 2002;

Penterman et al. 2007; Hsieh et al. 2009). Because the 5-methylcytosine DNA

glycosylases can recognize and directly remove the 5-mC base, they are critical

enzymes in the active DNA demethylation pathway and are thus often referred to

as DNA demethylases (Zhang et al. 2018). Research on how these DNA

demethylases are regulated and recruited to their target loci is required to reveal

how distinct genomic DNA methylation patterns are generated and modified

during organismal development and in response to environmental change.

The histone acetyltransferase Increased DNA Methylation 1 (IDM1) is part of the

IDM protein complex, which specifically recognizes certain methylated genomic

regions through the Methyl-CpG-Binding domain (MBD) of its subunits

Methyl-CpG-Binding protein MBD7 and IDM1, and through the DNA-binding

domain of HARBINGER TRANSPOSON-DERIVED PROTEIN HDP2 (Qian et al.

2012; Qian et al. 2014; Lang et al. 2015; Duan et al. 2017). Recent research

showed that ROS1 physically interacts with the histone variant H2A.Z and is

recruited to some genomic regions via SWR1-mediated H2A.Z deposition, which

involves specific histone acetylation marks created by the IDM complex (Nie et al.

2019). Because this mechanism only applies to a subset of target genomic

regions of ROS1, there must be alternative mechanisms for the targeting of

ROS1.

In this study, we found that ROS1 forms a protein complex with RWD40, RMB1,

and RHD1, which are proteins that contain WD40, methyl-DNA binding, and

homeodomain and zinc finger domains, respectively. We show that this novel

protein complex, which we call the RWD40 complex, is required for the prevention

of DNA hypermethylation at several genomic regions in *Arabidopsis*, and that this

function is independent of the IDM-SWR1-H2A.Z pathway. One of these regions

is the *ROS1* promoter, which contains a DNA methylation monitoring sequence

(MEMS) important for DNA methylation homeostasis in *Arabidopsis* (Lei et al.

2015; Williams et al. 2015). Consistent with its role in controlling the DNA

methylation of the *ROS1* promoter, we found that this protein complex negatively

regulates the expression of *ROS1*; importantly, ROS1 binding to the MEMS

requires RWD40, RMB1, and RHD1, suggesting that the RWD40 complex

dictates ROS1 targeting to this locus. Moreover, our results indicate that this

complex plays a role in anti-bacterial resistance, consistent with the known

function of ROS1 in plant-bacteria interactions. Taken together, our results

suggest that RWD40, RMB1, and RHD1 act in concert with ROS1 to control

genomic DNA methylation by regulating *ROS1* gene expression and likely by

dictating the targeting of the ROS1 protein to selected target regions.

RESULTS

A WD40 protein physically interacts with ROS1 and functions in DNA demethylation

To date, two proteins, H2A.Z and MET18 (homolog of yeast MET18, Methionine 18), have been reported to physically interact with ROS1 and to regulate active DNA demethylation in *Arabidopsis* plants (Duan et al. 2015; Nie et al. 2019). H2A.Z is required for ROS1 to target specific genomic regions defined by the IDM and SWR1 protein complexes (Qian et al. 2012; Lang et al. 2015; Nie et al. 2019). MET18 is a critical enzyme in the biosynthesis of the MoCo cofactor required for the methyl-DNA glycosylase/lyase activity of ROS1 (Duan et al. 2015). To identify additional ROS1-interacting proteins, we used yeast two-hybrid (Y2H) assays to determine whether ROS1 interacts with selected proteins found in the anti-FLAG pull-down products from *ROS1-3xFLAG-3xHA* (*ros1-13* mutant expressing *3xFLAG-3xHA*-tagged *ROS1* under its native promoter) *Arabidopsis* plants (Lei et al. 2015) in previous experiments. The results showed that a WD40 domain-containing protein (which we named RWD40 for ROS1-associated WD40 domain-containing protein) interacts with ROS1 (Figure 1A and 1B). The interaction between ROS1 and RWD40 was confirmed by bimolecular fluorescence complementation (BiFC) (Figure 1C). Transient expression assays in *Nicotiana benthamiana* leaves showed that RWD40 is mainly localized in the nucleus (Figure S1A). In the anti-FLAG immunoprecipitate from *ROS1-3xFLAG-3xHA* plants (Lei et al. 2015), but not from Col-0 control plants, we identified multiple RWD40 peptides (Figure 1D). Furthermore, we also identified peptides corresponding to ROS1 in the anti-FLAG immunoprecipitate from

transgenic plants expressing a *3xFLAG-RWD40* fusion driven by the *RWD40*

native promoter in the *rwd40-1* mutant background (Figures1D and S1B-S1D).

These results strongly suggest that ROS1 physically interacts with RWD40 in

Arabidopsis.

To investigate whether RWD40 regulates ROS1-dependent active DNA

demethylation, we used Chop-PCR to determine the DNA methylation level at the

AT5G39160 locus in *rwd40-1* and *rwd40-2* mutant seedlings. Similar to *ros1*

mutant plants (Qian et al. 2012), *rwd40-1* and *rwd40-2* plants showed increased

DNA methylation at the 5' region of AT5G39160 (Figure S1E). Individual locus

bisulfite sequencing confirmed that the DNA methylation levels at the AT5G39160

locus were higher in *rwd40-1*, *rwd40-2*, and *ros1-4* mutant plants than in the

wild-type control (Figure S1F). Interestingly, the methylation level at AT5G39160

was not increased in the *idm1-2* mutant (Figure S1E and S1F). The expression of

RWD40-3xFLAG fusion driven from its native promoter in the *rwd40-1* mutant

reduced the methylation level at this locus (Figure S1E). These results show that

RWD40 interacts with ROS1 and functions in regulating DNA demethylation.

RMB1 and RHD1 interact with RWD40 and function in DNA demethylation

We noticed that the anti-FLAG immunoprecipitates from *3xFLAG-RWD40*

transgenic plants contained peptides corresponding to AT1G63240 and

AT5G42780, in addition to peptides from RWD40 and ROS1 (Figure 1D).

AT1G63240 and AT5G42780 peptides were also detected in the anti-FLAG

immunoprecipitates from *ROS1-3xFLAG-3xHA* transgenic plants (Figure 1D). The

AT1G63240 protein is predicted to have a non-canonical methyl-DNA binding

domain, a CBD domain, and a domain of unknown function, and is hereafter

referred to as RMB1 (for ROS1-associated methyl-DNA binding protein 1) (Figure

S2A). The AT5G42780 protein contains a zinc-finger domain and a homeodomain,

and is hereafter referred to as RHD1 (for ROS1-associated homeodomain protein

1) (Figure S2B). In Y2H assays, RWD40 interacts with RMB1 and RHD1, and

RMB1 interacts with RHD1 (Figure 2A). These interactions as well as the

interaction between RWD40 and ROS1 were confirmed by split luciferase and

BiFC assays in *N. benthamiana* leaves (Figures 2B and S2C). The assays also

show that neither RMB1 nor RHD1 can directly interact with ROS1 (Figures 2A,

2B and S2C).

RMB1 was immunoprecipitated from transgenic plants expressing a

3xFLAG-RMBD1 fusion driven by the RMB1 native promoter in the *rmb1-1* mutant

background (Figure S2A). The immunoprecipitate contained not only RMB1 but

also RWD40, RHD1, and ROS1 (Figure 1D). Similarly, the RHD1

immunoprecipitate from transgenic plants expressing a *3xFLAG-RHD1* fusion

driven by the RHD1 native promoter in the *rhd1-1* background (Figure S2B)

contained not only RHD1 but also RMB1, ROS1, and RWD40 (Figure 1D). These

results show that RWD40, RMB1, and RHD1 proteins are associated with ROS1

in vivo.

Dysfunction of *RMB1* caused DNA hypermethylation at the AT5G39160 locus

(Figures S2D and S2E). The DNA hypermethylation phenotype of the *rmb1-1*

mutant was suppressed in transgenic plants expressing *3xFLAG-RMB1* (Figure

S2D). The *rhd1-1* mutant, however, did not show a DNA hypermethylation

phenotype at the AT5G39160 locus (Figure S2F). We assessed the methylation

levels at another endogenous ROS1 target, AT4G18380, by Chop-PCR. Similar to

ros1-4 mutant plants, *rhd1-1* mutant plants had increased DNA methylation at

AT4G18380 (Figure S2F). Expression of the *3xFLAG-RHD1* transgene

suppressed this hypermethylation phenotype of the *rhd1-1* mutant (Figure S2F).

These results suggest that RMB1 and RHD1 function in regulating locus-specific

demethylation in *Arabidopsis.*

RWD40, RMB1, RHD1, and ROS1 form a protein complex

The interactions of RWD40 with ROS1, RMB1, and RHD1, and between RMB1

and RHD1 in Y2H, BiFC, and split luciferase assays, together with their *in vivo*

association as indicated by immunoprecipitation-mass spectrometry analyses,

suggested that the four proteins may form a protein complex in plant cells. To

assess this possibility, we carried out gel filtration assays with protein extracts

from *3xFLAG-RMB1*, *3xFLAG-RWD40*, *3xFLAG-RHD1*, and

ROS1-3xFLAG-3xHA transgenic plants. The results showed that RMB1, RWD40,

RHD1, and ROS1 were eluted in the same fractions, and indicated that they exist

in a complex with an estimated size of approximately 350 kDa (Figure 3A). This is

close to the predicted total size of 328 kDa if the four proteins are in a 1:1:1:1 ratio.

We co-expressed full-length RWD40, RMB1, and RHD1, and a truncated ROS1

protein (amino acid residues 1 to 100) in Sf9 insect cells using baculovirus vectors.

After conducting histidine affinity purification and gel filtration, we found that the

four proteins were eluted in a single peak fraction (Figure 3B). All the gel bands

(Figure 3B) were extracted and their identities were confirmed by mass

spectrometry (Table S1). Taken together, these results suggest that RWD40,

RMB1, RHD1, and ROS1 form a protein complex *in vivo*, referred to as the

RWD40 complex hereafter.

Identification of the protein domains that mediate the interactions within the

protein complex

ROS1 contains an HhH-GPD domain, an End-III domain, a CXXC domain, an

RRM_DME domain, and an N-terminal domain of unknown function (Figure S3A).

The ROS1 N-terminal region (amino acid residues 41 to 100) has sequence

similarity to the EAR (Ethylene response factor–associated Amphiphilic

Repression) motif (Yang et al. 2018) (Figure S3A), and is thus referred to as the

ELD (EAR-like Domain) domain. To determine which domain in ROS1 may

mediate the interaction between ROS1 and RWD40, we performed Y2H assays

with various deletion mutants of ROS1. We found that the N-terminal half (amino

acid residues 1 to 696) of ROS1 interacts with RWD40, whereas the C-terminal

half (amino acid residues 697 to 1393) does not (Figure S3B). Further truncation

of the N-terminal half of ROS1 showed that the ELD domain (amino acid residues

41 to 100) is sufficient to mediate the interaction with RWD40 (Figure S3B).

We also performed Y2H assays with various deletion mutants of RWD40, and

found that the N-terminal half of RWD40, including two LISH-CTLH domains but

not the WD40 domain, is sufficient to enable the interaction with ROS1 (Figures

1B and S3B). On the other hand, the WD40 domain of RWD40 interacts with

RMB1 in Y2H assays (Figure 2D). The C-terminal region of RMB1 interacts with

the WD40 domain of RWD40 (Figures 2C and 2D), and is thus referred to as the

WID (WD40 interaction domain) domain (Figure 2C). The WD40 domain of

RWD40 is also capable of mediating the interaction between RWD40 and RHD1

(Figures S3C and S3D). The zinc-finger domain of RHD1 is responsible for the

interaction with RWD40, while both the zinc finger domain and homeodomain of

RHD1 are necessary for the interaction with RMB1 (Figures S3C and S3E). In the

Y2H assay, RHD1 also interacts with itself, and this self-interaction is mediated

through the zinc finger domain (Figures S3C and S3D).

RMB1 binds to methylated DNA through the MBD domain

Arabidopsis has 13 canonical MBD proteins (Zemach and Grafi 2007). Our

analysis indicates that *Arabidopsis* also has three non-canonical MBD proteins,

namely RMB1, IDM1, and the protein encoded by AT4G14920 (Figure S4A). The

IDM complex contains two MBD domain proteins, i.e., the histone

acetyltransferase IDM1 and MBD7 (Qian et al. 2012; Lang et al. 2015). The two

MBD domain proteins recognize methylated cytosine and thus ensure that the

IDM histone acetyltransferase complex is directed only to methylated sequences

(Qian et al. 2012; Lang et al. 2015). Many of the amino acid residues in the MBD

domain of RMB1 are not conserved compared to the other MBD proteins (Figure

S4B). We carried out electrophoretic mobility shift assays (EMSA) to determine

whether RMB1 can bind to methylated DNA. The EMSA assays showed that

RMB1 is capable of binding to DNA probes containing CG, CHG, or CHH

methylation, and that the binding was competitively blocked by unlabeled

methylated DNA but to a lower extent by unmethylated DNA of the same

sequence (Figures 4A and 4B). To confirm RMB1 binding to methylated DNA, we

carried out microscale thermophoresis (MST) assays and found that the MBD

domain of RMB1 is capable of binding to DNA methylated in any sequence

context (Figures 4C). We constructed four mutant versions of the MBD domain,

including W22G, Y38F, T49A, and K50T (Figure S4B), and found that the W22G,

Y38F, and T49A mutations decreased the methyl-DNA binding activity, while the

K50T mutation abolished the methyl-DNA binding activity (Figures 4C).

Expression of the wild-type but not of the K50T mutant version of RMB1 under its

native promoter complemented the At5g39160 DNA hypermethylation phenotype

of the *rmb1-1* mutant (Figure S5A). These results show that RMB1 is a novel

methyl-DNA-binding protein, and that its methyl-DNA binding is critical for it to

function in active DNA demethylation.

RWD40, RMB1, and RHD1 regulate *ROS1* expression

To begin to explore the function of the protein complex formed by RWD40, RMB1,

RHD1, and ROS1 in plants, we assessed the DNA methylation levels at the *ROS1*

promoter region, which contains a DNA methylation monitoring sequence that

senses methylation and demethylation activities and that regulates *ROS1*

expression (Lei et al. 2015; Williams et al. 2015). Individual locus bisulfite

sequencing showed that the DNA methylation level at the MEMS was increased in

rwd40-1, *rmb1-1*, *rhd1-1*, and *ros1-4* compared to the wild-type control (Figure

5A). In contrast, the MEMS DNA methylation level was not increased in the

idm1-2 mutant (Figure 5A). Consistent with the changes in DNA methylation

levels at the MEMS, *ROS1* expression was increased in *rwd40*, *rmb1*, and *rhd1*

mutant plants but not in *idm1-2* mutant plants (Figure 5B). In the *ros1-4* mutant,

ROS1 expression measured using the 5' primer pair (Figure 5B) was also

increased, as expected (Lei et al.,2015). The expression of the *3xFLAG-RWD40*

transgene driven by the *RWD40* native promoter in the *rwd40-1* mutant restored

the expression of *ROS1* to the wild-type level (Figure 5B). ChIP-qPCR assays

confirmed that ROS1, RWD40, RMB1, and RHD1 were enriched at the MEMS in

the *ROS1* promoter (Figure 5C), and that *RWD40*, *RMB1*, and *RHD1* were

necessary for ROS1 binding to the MEMS region (Figure 5D). Taken together,

these results indicate that *RWD40*, *RMB1*, and *RHD1* mediate binding of ROS1 to

the MEMS region, hence regulating its methylation level and the subsequent

expression of *ROS1*.

RWD40, RMB1, and RHD1 regulate DNA demethylation in an

IDM1-independent manner

RWD40, *RMB1*, *RHD1*, and *ROS1*, but not *IDM1*, are necessary for the

prevention of DNA hypermethylation at the MEMS in the *ROS1* promoter (Figure

5A). These results suggest that RWD40, RMB1, and RHD1 might regulate

ROS1-mediated DNA demethylation in a manner independent of IDM1. To further

investigate the role of RWD40, RMB1, and RHD1 in the regulation of active DNA

demethylation, we used Chop-PCR to assess the DNA methylation levels in

rwd40-1, *rmb1-1*, and *rhd1-1* single mutants and in the *rwd40 rmb1 rhd1* triple

mutant*,* as well as in *idm1-2*, at several genomic targets of ROS1. Similar to DNA

methylation levels in the *ros1-4* mutant, DNA methylation levels at the

AT1G35140, AT4G00660, AT4G18380, and AT5G55875 loci were increased in

rwd40-1, *rmb1-1*, and *rhd1-1,* but not *idm1-2* mutant plants, relative to the Col-0

control (Figure 5E). Individual locus bisulfite sequencing data confirmed that the

DNA methylation levels at these ROS1 targets were increased in *rwd40-1*,

rmb1-1, *rhd1-1,* and *ros1-4* mutant plants, but not in *idm1-2* mutant plants (Figure

5F). Analysis of the *rwd40 rmb1 rhd1* triple mutant indicated that the *rwd40-1*,

rmb1-1, and *rhd1-1* mutations are not additive in causing the DNA

hypermethylation phenotype (Figures 5F). We also determined the DNA

methylation levels at several genomic regions (including DT-75, DT-76, DT-77,

and DT-78) that are known to be hypermethylated in *idm1* and *ros1* mutants (Qian

et al. 2012)and that are amenable to Chop-PCR assays (Duan et al. 2017). DNA

methylation levels were not increased at these regions in *rwd40-1*, *rmb1-1*, or

rhd1-1 mutant plants (Figure S6A). Together, these results support the inference

that RWD40, RMB1, and RHD1 function in regulating active DNA demethylation

independently of IDM1.

ROS1 and the IDM complex are required for the prevention of silencing of the

35S:SUC2 reporter gene in *Arabidopsis* (Qian et al. 2014; Lang et al. 2015; Duan

et al. 2017; Nie et al. 2019). When we introgressed the *35S:SUC2* reporter gene

into *rwd40-1*, *rmb1-1*, or *rhd1-1* mutant plants through genetic crosses, we found

that the reporter gene was not silenced in any of these mutants (Figures S6B and

S6C). Therefore, unlike IDM1 or ROS1, RWD40, RMB1, and RHD1 are not

required for the prevention of silencing of the *35S:SUC2* reporter gene in

Arabidopsis. We examined the deposition of H2A.Z and H3K18ac in wild-type

35S:SUC2 plants (Nie et al., 2019) at the 5 loci shown in Figure 5F, where DNA

methylation levels were increased in RW40 complex-related mutants, but not in

idm1-2 (Figures 5E and 5F). Unlike the moderate co-existence of H2A.Z and

H3K18ac in the IDM1-dependent AT1G62760 locus, only at the AT4G18380 locus

were H2A.Z and H3K18ac both less deposited; at the AT5G55875 locus, H2A.Z

and H3K18ac were rarely deposited; either H2A.Z or H3K18ac was deposited at

the MEMS, AT1G35140, and AT4G00660 loci (Figure S6D). No consistent change

in histone acetylation or H2A.Z deposition could be found at these 5 selected loci,

which is in line with the notion that RWD40, RMB1, and RHD1 function in an

IDM1-independent manner.

The RWD40 complex prevents the silencing of several endogenous genes

and regulates antibacterial resistance

To test the effect of the dysfunction of the RWD40 complex on genome-wide

transcript levels, we profiled the transcriptomes of *ros1-4* and *rwd40 rmb1 rhd1*

mutant plants by mRNA-seq. There were 3,817 and 2,970 up-regulated genes

and 4,028 and 2,918 down-regulated genes in *rwd40 rmbd1 rhd1* triple-mutant

and *ros1-4* mutant plants ,respectively, compared to wild-type Col-0 plants (Figure

6A and Table S2). Of these, 1,673 up-regulated and 1,941 down-regulated genes

were shared between *rwd40 rmbd1 rhd1* and *ros1-4* (Figure 6B). Several

commonly down-regulated DEGs in *rwd40 rmbd1 rhd1* and *ros1-4* mutants were

confirmed by qRT-PCR (Figure 6C). These results suggest that the RWD40

complex is required for the expression of some endogenous genes.

Exposure to the bacterial pathogen *Pseudomonas syringae* pv. *tomato* DC3000

(*Pto* DC3000) causes moderate but widespread differential DNA methylation in

Arabidopsis (Dowen et al. 2012). A mildly enhanced bacterial growth was

observed in *ros1* plants treated with the bacterial pathogen *Pto* DC3000,

supporting a role of ROS1-dependent DNA demethylation in antibacterial

resistance (Yu et al. 2013). To test whether the RWD40 complex regulates

antibacterial resistance, we first inoculated the RWD40 complex mutants with *Pto*

DC3000 by flood-inoculation assays (Ishiga et al. 2011). We found that similar to

ros1-4 mutant plants, *rwd40-1*, *rmb1-1*, and *rhd1-1* mutant plants were more

susceptible to *Pto* DC3000 than wild-type plants (Figures 6D and 6E). These

results point at a biological relevance of the RWD40 DNA demethylation complex

in *Arabidopsis* antibacterial defense.

DISCUSSION

In this study, we identified RWD40, RMB1, and RHD1 as cellular factors critical

for the regulation of *ROS1* expression and for the prevention of DNA

hypermethylation at several endogenous genomic regions (Figure 5). DNA

methylation patterns are important for organismal development, carcinogenesis,

and many other diseases, as well as for human aging (Jung and Pfeifer 2015;

Klutstein et al. 2016). It is therefore important to understand how DNA methylation

patterns are controlled by active DNA demethylation (Penterman et al. 2007;

Furner and Matzke 2011; Zhang et al. 2018). Although the biochemistry of

ROS1-mediated enzymatic removal of DNA methylation has been extensively

studied, the mechanisms by which the enzymatic machinery is regulated and

recruited to specific target sites are poorly understood (Qian et al. 2012; Lang et

al. 2015; Nie et al. 2019). Our results indicate that RWD40, RMB1, RHD1, and

ROS1 form a RWD40 complex, and that this protein complex contributes to

locus-specific DNA demethylation. Moreover, ROS1 binding to the *ROS1*

promoter requires RWD40, RMB1, and RHD1, suggesting that the RWD40

complex directs ROS1 targeting to this locus to regulate the expression of *ROS1*.

Thus, the activity of the RWD40 complex at target genomic loci is likely

self-restrained by *ROS1* expression level through its demethylation function at the

MEMS site on the *ROS1* gene promoter (Figure 7). These findings provide novel

insights into the regulation and targeting of active DNA demethylation in plants.

In mammals, active DNA demethylation is initiated by the Tet dioxygenases (Wu

and Zhang 2017). Several transcription factors have been shown to interact with

and to target Tet enzymes to genes critical for cell differentiation and

reprogramming (Costa et al. 2013; de la Rica et al. 2013; Wang et al. 2015; Xiong

et al. 2016; Sardina et al. 2018). In *Arabidopsis*, the IDM histone acetyltransferase

complex is required for directing ROS1 to a subset of active DNA demethylation

target regions (Qian et al. 2012; Qian et al. 2014; Lang et al. 2015; Duan et al.

2017; Nie et al. 2019). MBD7 in the IDM complex recognizes dense mCpG sites

and thus ensures that ROS1 is eventually targeted to genomic regions with

densely methylated CpG sequences (Lang et al. 2015). On the other hand, the

SANT/Myb/trihelix DNA-binding motif-containing protein HDP2 in the IDM

complex probably helps direct the complex to possible regulatory sequences, thus

avoiding heavily methylated transposon body regions and some genic regions

that should not be demethylated (Duan et al. 2017). The IDM complex creates

histone acetylation marks that attract the SWR complex, which then deposits the

histone variant H2A.Z. Once deposited at the targeted site, H2A.Z helps to recruit

ROS1 through a direct physical interaction (Nie et al. 2019). The protein complex

identified in this study is similar to the IDM complex in that it also contains a

protein (i.e., RMB1) that recognizes methylated DNA, and a transcription

factor-like protein (i.e., RHD1). RMB1 ensures that the complex is directed to

methylated genomic regions, while RHD1 probably helps to target the complex to

regulatory sequences. Like the IDM complex, the RWD40 protein complex is likely

also important for targeting active DNA demethylation to specific genomic regions

(Figures 5A, 5E, and 5F). Our results suggest that the RWD40 complex targets

active DNA demethylation to genomic regions that are different from those

targeted by the IDM complex. This difference can be explained by the fact that

RMB1 recognizes mC marks in all sequence contexts (Figure 4), whereas MBD7

and IDM1 bind to mCpG sequences only (Qian et al. 2012; Lang et al. 2015)

(Figure S5B and S5C). In addition, HDP2 and RHD1 likely have different

sequence specificities in their interactions with DNA, which probably also

contributes to the different targeting preferences of the two protein complexes.

The lack of DNA hypermethylation at the AT5G39160 locus in *rhd1* mutant plants

(Figure S2F) indicates that RHD1 may be dispensable for some targets of the

RWD40 protein complex, either due to genetic redundancy with another

transcription factor-like protein, or because its function is genomic

region-dependent. It should be noted that H2A.Z deposition and histone

acetylation are not restricted to the IDM1-dependent ROS1-targets, since H2A.Z

or H3K18ac are also deposited at some loci targeted by RWD40 complex (Figure

S6D); this observation suggests that other IDM1-independent acetylation

mechanisms are associated with ROS1 activity and possibly involved in H2A.Z

deposition.

Another important difference between the RWD40 complex and the IDM complex

is that the RWD40 complex contains ROS1 and thus has a direct role in targeting

active DNA demethylation. Furthermore, because the RWD40 complex targets the

MEMS in the *ROS1* promoter, it can regulate *ROS1* gene expression. We propose

that through the regulation of *ROS1* gene expression, the RWD40 complex is

expected to have a broad effect on genomic DNA methylation patterns. The

genome-wide effects of DNA demethylation factors can be difficult to analyze

because DNA methylation changes caused by mutations in these factors are often

quite subtle (Nie et al. 2019). Future studies are needed to determine the

genome-wide effects of *rwd40, rmb1,* and *rhd1* mutations, and to identify the

direct targets of the RWD40 complex at the whole genome level.

The data presented here, together with our previous work (Lang et al. 2015; Nie et

al. 2019), suggest that active DNA demethylation activities are targeted only to

methylated genomic regions by proteins that bind to methyl-DNA. Such targeting

would be biologically relevant, because only methylated genomic regions may

require demethylation. Our findings also suggest that the presence of a

transcription factor-like protein is important for the targeting of active DNA

demethylation. It is tempting to speculate that these transcription factor-like

proteins preferentially bind to regulatory sequences, and thus help target active

DNA demethylation to regulatory sequences in order to prevent harmful silencing

and precisely regulate gene expression.

MATERIALS AND METHODS

Plant materials and growth conditions

Plants were grown under long-day conditions (16-h light/8-h dark) at 22°C. The

T-DNA insertion mutants *rwd40-1* (SALK_068825), *rwd40-2* (SALK_012947),

rmb1-1 (SALK_110885), *rmb1-2* (SALK_100783), *rhd1-1* (SALK_092897), *ros1-4*

(SALK_045303), and *idm1-2* (SALK_062999) were obtained from the *Arabidopsis*

Biological Resource Center (http://www.arabidopsis.org) and genotyped by PCR

with the primers listed in Table S3.

METHODS DETAILS

Mutant plant complementation

For the complementation of mutants, the *RWD40*, *RMB1*, or *RHD1* genomic DNA

with the 2-kb upstream region (as the native promoter region) was amplified from

genomic DNA of Col-0 with the primers listed in Table S3. The amplification

products were cloned into the pCambia1305 vector (with a *3xFLAG* tag or a

3xMYC tag at the C-terminus) by T4 DNA ligase (NEB). The constructs were

transformed into mutants using *Agrobacterium tumefaciens* GV3101 by the

standard floral dip method (Clough and Bent 1998). Unsegregated T3

homozygous complementation lines were then used for further experiments.

Real-time qPCR and Chop-PCR

For real-time qPCR analysis, total RNA was extracted from 0.1 g of 14-day-old

Arabidopsis seedlings with the RNeasy plant kit (Qiagen). 2 μg of the mRNA were

converted into cDNA with the First-Strand cDNA Synthesis Kit (TRNSGENE Bio.,

China) following the manufacturer's instructions, and the cDNAs were used as

templates for real-time PCR with iQ SYBR green supermix (Bio-RAD). For

Chop-PCR, genomic DNA was extracted from 14-day-old seedlings, and 500 ng

of DNA were digested with the indicated methylation-sensitive enzyme in a 20-µL

reaction mixture. After digestion, PCR was performed using 2 µl of the digested

DNA as template in a 20-µl reaction mixture with the primers listed in Table S3.

Undigested DNA was amplified as the loading control.

IP and LC-MS/MS analysis

For immunoprecipitation (IP), about 5 g of floral tissue for each epitope-tagged

transgenic line were used. Dynabeads (10004D, Invitrogen) conjugated with

FLAG antibody (F1804, Sigma) were applied for IP. Affinity purification was

performed as previously described (Law et al. 2010), and the protein samples

were subjected to Liquid Chromatography-Mass Spectrometry (LC-MS) /MS

analysis as previously described (Lang et al. 2015; Qin et al. 2017; Nie et al.

2019).

Protein extraction and western blot detection

About 0.1 g of plant tissue were harvested and ground to a fine power in liquid N_2.

Total protein was extracted by protein lysis buffer (LB: 0.5 mM DTT, 5 mM $MgCl_2$,

50 mM Tris (pH 7.6), 10% glycerol, 150 mM NaCl, 0.1% NP-40, 1 mM PMSF, and

protease inhibitor cocktail (Roche)). The proteins were then detected by western

blot with anti-FLAG (F1804, Sigma) or anti-MYC (05-724, Millipore).

Split luciferase complementation assays

The coding sequences of RWD40, RMB1, RHD1, and ROS1 proteins were cloned

into pCAMBIA-cLUC and/or pCAMBIA-nLUC vectors. *A. tumefaciens* GV3101

carrying different constructs were infiltrated into 4-week-old *N. benthamiana*

leaves. Luciferase activity was detected 48 h post infiltration.

Individual locus bisulfite sequencing

Individual locus bisulfite sequencing

200 ng of genomic DNA extracted from Col-0 wild type and the indicated mutants

were treated with the BisulFlash DNA Modification Kit (Epigentek) according to

the manufacturer's protocol. A 2-µL volume of bisulfite-treated DNA was used for

PCR amplification with the primers listed in Table S3. The PCR products were

cloned into the pMD18-T vector (Takara) according to the supplier's instructions.

At least 15 independent clones were sequenced per sample and analyzed using

the online tool Kismeth (http://katahdin.mssm.edu/kismeth/revpage.pl).

Bimolecular fluorescence complementation (BiFC) assay

The coding sequences of RWD40, RMB1, RHD1, and ROS1 were cloned into the

p2YN or p2YC vector to generate fused split YFP constructs. For protein-protein

interaction analysis, *A. tumefaciens* GV3101 carrying the indicated constructs

were cultured overnight. When the cultures reached an OD_{600} of 1.0, they were

firstly collected by centrifuge and then were resuspended in buffer containing 10

mM $MgCl_2$, 100 µM acetosyringone, and 10 mM MES (pH 5.6). The suspensions

were kept at room temperature for 2 hours and then were infiltrated into *N.*

benthamiana leaves. Fluorescence was examined at 2 days post infiltration.

Gel filtration assays and western blot analysis

Western blotting of gel filtration samples was performed as previously described

(Duan et al. 2017). In brief, 2 g of flower tissue from Col-0 wild type or from the

indicated transgenic plants were harvested and ground into a fine power in liquid

N_2. The fine powder was resuspended in 3 ml of lysis buffer (0.5 mM DTT, 5 mM

$MgCl_2$, 50 mM Tris [pH 7.6], 10% glycerol, 150 mM NaCl, 0.1% NP-40, 1 mM

PMSF), and the suspension was kept at 4°C for 5 min without shaking. The

supernatant was loaded into a HiPrep 16/60 Sephacryl S-300 HR column (GE

Healthcare), and 10 fractions were collected. Each fraction was run on 10%

SDS-PAGE for western blot detection.

Y2H assay

In brief, the corresponding cDNA sequences were cloned into pGADT7-AD or

pGBKT7-BD vectors (Clontech), and the pair of genes to be tested for interaction

were co-transformed into the yeast strain Gold (Clontech). Y2H assays were

performed as previously described (Bai et al. 2013).

Protein expression and purification

Arabidopsis *RWD40*, *RMB1*, and *RHD1* were cloned into the pFastBac1 vector

adding a FLAG tag at their N-terminus. Arabidopsis *ROS1* (encoding amino acids

1-100) was cloned into the pFastBac1 vector adding a double His tag

(6xHis-13aa-10xHis) fused with GFP at its N-terminus. Recombinant

baculoviruses were generated by the Bac-to-Bac system in Sf9 insect cells. For

protein co-expression, the insect cells were grown to a density of 2.0E+06 cells

per mL and were then infected with four separate viruses. The infected cells were

cultured for 60-72□h at 27°C before collection. Collected cell pellets were stored

at -80°C before use. Each cell pellet was resuspended in buffer A (20 mM HEPES

pH 7.5, 150 mM NaCl, 10% v/v glycerol, 50 mM imidazole, and 0.1 mM Tris

(2-carboxyethyl) phosphine hydrochloride (TCEP), supplemented with protease

inhibitor cocktail) and then lysed by sonication. The suspension was centrifuged

at 18,000 × *g* for 30 min at 4°C. The supernatant was incubated with Ni-NTA resin

at 4°C for 1 h. The resin was washed with 20 column volumes of buffer A in a

gravity column. The complex was eluted by buffer B (buffer A with 250 mM

imidazole) and was then loaded onto a Superdex200 10/300 GL column

equilibrated with buffer containing 20 mM HEPES pH 7.5, 100 mM NaCl, and 0.1

mM TCEP. Fractions containing the ROS1 complex were collected and analyzed

with SDS-PAGE and Coomassie brilliant blue staining.

Microscale thermophoresis (MST)

The sequence encoding the MBD domain of RMB1 (amino acids 1-81) was

cloned into a pET-Sumo vector to fuse the target protein with a hexahistidine plus

yeast sumo tag at the N-terminus. The His-Sumo tagged MBD domain of RMB1

recombinant protein was transformed into *Escherichia coli* strain BL21 (DE3).

When the OD_{600} of the cell culture reached 0.7, IPTG was added to a final

concentration of 0.2 mM to induce protein expression. The cells were harvested

by centrifugation, re-suspended in buffer A (500 mM NaCl, 20 mM Tris-HCl, pH

7.5), and then lysed by sonication. The supernatant was loaded into a HisTrap

column (GE Healthcare) and washed with buffer A and buffer B (buffer A plus 50

mM immidazol). The target protein was finally eluted with buffer C (buffer A plus

500 mM immidazol). The eluted protein was dialyzed against buffer A and

digested with Ulp1 protease. The His-Sumo tag was removed by passing the

digestion product through a second step HisTrp column. The flow-through protein

was dialyzed against buffer D (100 mM NaCl, 20 mM HEPES, pH 7.0, 5 mM DTT)

and loaded onto a Heparin column (GE Healthcare). A linear gradient from buffer

D to buffer E (2 M NaCl, 20 mM HEPES, pH 7.0) was applied to elute the target

protein. The eluted protein was further purified using a Superdex G200 column

with buffer E (50 mM NaCl, 20 mM HEPES, pH 7.0, 5 mM DTT) and concentrated

to 10 mg/ml for further use. The mutated forms of the MBD domain of RMB1 were

expressed and purified following the same protocol.

The MST assay was performed using a Monolith NT.115 instrument (NanoTemper

Technologies) (Jerabek-Willemsen et al. 2014; Harris et al. 2018). The purified

MBD domain of RMB1 was half-and-half diluted in 14 steps with the buffer E,

covering a concentration range from about 50 µM to 1.5 nM (Figure 4C). For each

DNA duplex, one DNA strand is labeled with Biotin at the 5' during the synthesis

(Figure 4C). The Biotin labeled strand and the unlabeled complementary strand

were annealed together and further mixed with the diluted protein steps. The

mixture was loaded into the MST capillaries. Three repeats for each

measurement were conducted. The data were analyzed using the NanoTemper

analysis software (NanoTemper Technologies).

Pto DC3000 bacterial infection assays

Pto DC3000 bacterial infection assays were performed according to (Ishiga et al.

2011) with some modifications. Briefly, the indicated mutants were germinated

and grown on 1/2 MS medium. A single colony of *Pto* DC3000 was inoculated

onto new mannitol-glutamate medium agar with rifampicin (50 µg/ml) and

streaked on the plate. The plate was inoculated at 28°C for 36 hours. The

collected bacterial colonies were suspended in an appropriate volume of sterile

distilled water. 40 ml of a *Pto* DC3000 bacterial suspension (4 x 10^6 CFU, colony

forming units) were poured onto the plate containing 3-week-old *Arabidopsis*

seedlings and incubated for 2 minutes at room temperature. The bacterial

suspension was then removed by pipetting and the plates were sealed with 3M

Micropore 2.5 cm surgical tape. The infected seedlings were incubated in

long-day (16-h light / 8-h dark) conditions at 22 °C for three days. The

development of symptoms was evaluated and photographed at 3

days-post-inoculation (dpi). Bacterial growth was evaluated according to (Ishiga et

al. 2011).

mRNA-seq analysis

For mRNA-seq analyses, total RNA was extracted from 2-week-old seedlings with

the RNeasy plant kit (Qiagen), and subjected to RNA deep sequencing. Three

biological replicates were performed per genotype. The libraries were constructed

and sequenced at the Genomics Core Facility of the Shanghai Centre for Plant

Stress Biology, Chinese Academy of Sciences, using an Illumina HiSeq2500.

Low-quality sequences (q < 20) and adapters were trimmed using Trimmomatic

and cutadpte (Bolger et al. 2014). Clean reads were mapped to the *Arabidopsis*

reference genome (TAIR10) using STAR (Dobin et al. 2013). DESeq2 was used to

detect differentially expressed genes according to adjusted p value (padj)

threshold of <0.01 (Love et al. 2014).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Figures S1-S7 and Tables S1-S3, and can be

found with this article online.

AUTHOR CONTRIBUTIONS

J.-K.Z. and P.L. designed the study. P.L., W.-F.N., X.X., Y.W., Y.J., P.H., X.L., G.Q.,

H.H., J.D., and Z.L. performed the experiments. P.L., W.-F.N., and J.-K.Z.

analyzed the data. W.-F.N., R.L.-D., and J.-K.Z. wrote the manuscript.

ACKNOWLEDGEMENTS

This work was supported by the Chinese Academy of Sciences, National Nature

Science Foundation of China (32002046), and Natural Science Foundation of

Jiangsu Province (BK20200948).

References

Bai G, Yang DH, Zhao Y, Ha S, Yang F, Ma J, Gao XS, Wang ZM, Zhu JK (2013)
Interactions between soybean ABA receptors and type 2C protein phosphatases.
Plant Mol Biol 83: 651-664
Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina
sequence data. **Bioinformatics** 30: 2114-2120
Cao X, Jacobsen SE (2002) Locus-specific control of asymmetric and CpNpG

methylation by the DRM and CMT3 methyltransferase genes. **Proc Natl Acad Sci U S**
A 99 Suppl 4: 16491-16498

Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen SE, Fischer
RL (2002) DEMETER, a DNA glycosylase domain protein, is required for endosperm
gene imprinting and seed viability in arabidopsis. **Cell** 110: 33-42

Clough SJ, Bent AF (1998) Floral dip: a simplified method for
Agrobacterium-mediated transformation of Arabidopsis thaliana. **Plant J** 16:
735-743

Costa Y, Ding J, Theunissen TW, Faiola F, Hore TA, Shliaha PV, Fidalgo M, Saunders A,
Lawrence M, Dietmann S, Das S, Levasseur DN, Li Z, Xu M, Reik W, Silva JC, Wang J
(2013) NANOG-dependent function of TET1 and TET2 in establishment of
pluripotency. **Nature** 495: 370-374

de la Rica L, Rodriguez-Ubreva J, Garcia M, Islam AB, Urquiza JM, Hernando H,
Christensen J, Helin K, Gomez-Vaquero C, Ballestar E (2013) PU.1 target genes
undergo Tet2-coupled demethylation and DNMT3b-mediated methylation in
monocyte-to-osteoclast differentiation. **Genome Biol** 14: R99

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. **Bioinformatics** 29:
15-21

Dowen RH, Pelizzola M, Schmitz RJ, Lister R, Dowen JM, Nery JR, Dixon JE, Ecker JR
(2012) Widespread dynamic DNA methylation in response to biotic stress. **Proc Natl**
Acad Sci U S A 109: E2183-2191

Duan CG, Wang X, Tang K, Zhang H, Mangrauthia SK, Lei M, Hsu CC, Hou YJ, Wang C,
Li Y, Tao WA, Zhu JK (2015) MET18 Connects the Cytosolic Iron-Sulfur Cluster
Assembly Pathway to Active DNA Demethylation in Arabidopsis. **PLoS Genet** 11:
e1005559

Duan CG, Wang X, Xie S, Pan L, Miki D, Tang K, Hsu CC, Lei M, Zhong Y, Hou YJ, Wang
Z, Zhang Z, Mangrauthia SK, Xu H, Zhang H, Dilkes B, Tao WA, Zhu JK (2017) A pair of
transposon-derived proteins function in a histone acetyltransferase complex for
active DNA demethylation. **Cell Res** 27: 226-240

Finnegan EJ, Dennis ES (1993) Isolation and identification by sequence homology of a
putative cytosine methyltransferase from Arabidopsis thaliana. **Nucleic Acids Res** 21:
2383-2388

Furner IJ, Matzke M (2011) Methylation and demethylation of the Arabidopsis
genome. **Curr Opin Plant Biol** 14: 137-141

Gehring M, Huh JH, Hsieh TF, Penterman J, Choi Y, Harada JJ, Goldberg RB, Fischer RL
(2006) DEMETER DNA glycosylase establishes MEDEA polycomb gene
self-imprinting by allele-specific demethylation. **Cell** 124: 495-506

Gong Z, Morales-Ruiz T, Ariza RR, Roldan-Arjona T, David L, Zhu JK (2002) ROS1, a

repressor of transcriptional gene silencing in Arabidopsis, encodes a DNA
glycosylase/lyase. **Cell** 111: 803-814

Haag JR, Pikaard CS (2011) Multisubunit RNA polymerases IV and V: purveyors of
non-coding RNA for plant gene silencing. **Nat Rev Mol Cell Biol** 12: 483-492

Harris CJ, Scheibe M, Wongpalee SP, Liu WL, Cornett EM, Vaughan RM, Li XQ, Chen W,
Xue Y, Zhong ZH, Yen LD, Barshop WD, Rayatpisheh S, Gallego-Bartolome J, Groth M,
Wang ZH, Wohlschlegel JA, Du JM, Rothbart SB, Butter F, Jacobsen SE (2018) A DNA
methylation reader complex that enhances gene transcription. **Science** 362: 1182-+

He XJ, Chen T, Zhu JK (2011) Regulation and function of DNA methylation in plants
and animals. **Cell Res** 21: 442-465

Hsieh TF, Ibarra CA, Silva P, Zemach A, Eshed-Williams L, Fischer RL, Zilberman D
(2009) Genome-wide demethylation of Arabidopsis endosperm. **Science** 324:
1451-1454

Ishiga Y, Ishiga T, Uppalapati SR, Mysore KS (2011) Arabidopsis seedling
flood-inoculation technique: a rapid and reliable assay for studying plant-bacterial
interactions. **Plant Methods** 7:

Jerabek-Willemsen M, Andre T, Wanner R, Roth HM, Duhr S, Baaske P, Breitsprecher
D (2014) MicroScale Thermophoresis: Interaction analysis and beyond. **J Mol Struct**
1077: 101-113

Jung M, Pfeifer GP (2015) Aging and DNA methylation. **BMC Biol** 13: 7

Klutstein M, Nejman D, Greenfield R, Cedar H (2016) DNA Methylation in Cancer and
Aging. **Cancer Res** 76: 3446-3450

Lang Z, Lei M, Wang X, Tang K, Miki D, Zhang H, Mangrauthia SK, Liu W, Nie W, Ma G,
Yan J, Duan CG, Hsu CC, Wang C, Tao WA, Gong Z, Zhu JK (2015) The
methyl-CpG-binding protein MBD7 facilitates active DNA demethylation to limit DNA
hyper-methylation and transcriptional gene silencing. **Mol Cell** 57: 971-983

Law JA, Ausin I, Johnson LM, Vashisht AA, Zhu JK, Wohlschlegel JA, Jacobsen SE
(2010) A protein complex required for polymerase V transcripts and RNA- directed
DNA methylation in Arabidopsis. **Curr Biol** 20: 951-956

Law JA, Jacobsen SE (2010) Establishing, maintaining and modifying DNA
methylation patterns in plants and animals. **Nat Rev Genet** 11: 204-220

Lei M, Zhang H, Julian R, Tang K, Xie S, Zhu JK (2015) Regulatory link between DNA
methylation and active demethylation in Arabidopsis. **Proc Natl Acad Sci U S A** 112:
3553-3557

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. **Genome Biol** 15: 550

Matzke MA, Mosher RA (2014) RNA-directed DNA methylation: an epigenetic
pathway of increasing complexity. **Nat Rev Genet** 15: 394-408

Nie WF, Lei M, Zhang M, Tang K, Huang H, Zhang C, Miki D, Liu P, Yang Y, Wang X,

Zhang H, Lang Z, Liu N, Xu X, Yelagandula R, Zhang H, Wang Z, Chai X, Andreucci A, Yu
JQ, Berger F, Lozano-Duran R, Zhu JK (2019) Histone acetylation recruits the SWR1
complex to regulate active DNA demethylation in Arabidopsis. **Proc Natl Acad Sci U S**
A:
Penterman J, Zilberman D, Huh JH, Ballinger T, Henikoff S, Fischer RL (2007) DNA
demethylation in the Arabidopsis genome. **Proc Natl Acad Sci U S A** 104: 6752-6757
Qian W, Miki D, Lei M, Zhu X, Zhang H, Liu Y, Li Y, Lang Z, Wang J, Tang K, Liu R, Zhu JK
(2014) Regulation of active DNA demethylation by an alpha-crystallin domain protein
in Arabidopsis. **Mol Cell** 55: 361-371
Qian W, Miki D, Zhang H, Liu Y, Zhang X, Tang K, Kan Y, La H, Li X, Li S, Zhu X, Shi X,
Zhang K, Pontes O, Chen X, Liu R, Gong Z, Zhu JK (2012) A histone acetyltransferase
regulates active DNA demethylation in Arabidopsis. **Science** 336: 1445-1448
Qin G, Ma J, Chen X, Chu Z, She YM (2017) Methylated-antibody affinity purification to
improve proteomic identification of plant RNA polymerase Pol V complex and the
interacting proteins. **Sci Rep** 7: 42943
Robertson KD (2005) DNA methylation and human disease. **Nat Rev Genet** 6:
597-610
Sardina JL, Collombet S, Tian TV, Gomez A, Di Stefano B, Berenguer C, Brumbaugh J,
Stadhouders R, Segura-Morales C, Gut M, Gut IG, Heath S, Aranda S, Di Croce L,
Hochedlinger K, Thieffry D, Graf T (2018) Transcription Factors Drive Tet2-Mediated
Enhancer Demethylation to Reprogram Cell Fate. **Cell Stem Cell** 23: 905-906
Slotkin RK, Martienssen R (2007) Transposable elements and the epigenetic
regulation of the genome. **Nat Rev Genet** 8: 272-285
Wang D, Xia X, Weiss RE, Refetoff S, Yen PM (2010) Distinct and histone-specific
modifications mediate positive versus negative transcriptional regulation of
TSHalpha promoter. **PLoS One** 5: e9853
Wang Y, Xiao M, Chen X, Chen L, Xu Y, Lv L, Wang P, Yang H, Ma S, Lin H, Jiao B, Ren R,
Ye D, Guan KL, Xiong Y (2015) WT1 recruits TET2 to regulate its target gene
expression and suppress leukemia cell proliferation. **Mol Cell** 57: 662-673
Williams BP, Pignatta D, Henikoff S, Gehring M (2015) Methylation-sensitive
expression of a DNA demethylase gene serves as an epigenetic rheostat. **PLoS Genet**
11: e1005142
Wu X, Zhang Y (2017) TET-mediated active DNA demethylation: mechanism, function
and beyond. **Nat Rev Genet** 18: 517-534
Xiong J, Zhang Z, Chen J, Huang H, Xu Y, Ding X, Zheng Y, Nishinakamura R, Xu GL,
Wang H, Chen S, Gao S, Zhu B (2016) Cooperative Action between SALL4A and TET
Proteins in Stepwise Oxidation of 5-Methylcytosine. **Mol Cell** 64: 913-925
Yang J, Liu Y, Yan H, Tian T, You Q, Zhang L, Xu W, Su Z (2018) PlantEAR: Functional
Analysis Platform for Plant EAR Motif-Containing Proteins. **Front Genet** 9: 590

Yu A, Lepere G, Jay F, Wang J, Bapaume L, Wang Y, Abraham AL, Penterman J, Fischer
RL, Voinnet O, Navarro L (2013) Dynamics and biological relevance of DNA
demethylation in Arabidopsis antibacterial defense. **Proc Natl Acad Sci U S A** 110:
2389-2394
Zemach A, Grafi G (2007) Methyl-CpG-binding domain proteins in plants: interpreters
of DNA methylation. **Trends Plant Sci** 12: 80-85
Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL,
Zilberman D (2013) The Arabidopsis nucleosome remodeler DDM1 allows DNA
methyltransferases to access H1-containing heterochromatin. **Cell** 153: 193-205
Zhang H, Lang Z, Zhu JK (2018) Dynamics and function of DNA methylation in plants.
Nat Rev Mol Cell Biol 19: 489-506
Zhu J, Kapoor A, Sridhar VV, Agius F, Zhu JK (2007) The DNA glycosylase/lyase ROS1
functions in pruning DNA methylation patterns in Arabidopsis. **Curr Biol** 17: 54-59
Zhu JK (2009) Active DNA demethylation mediated by DNA glycosylases. **Annu Rev**
Genet 43: 143-166

FIGURE LEGENDS

Figure 1. RWD40 interacts with ROS1.

(A) Schematic representation of the RWD40 protein (top panel) and of the

truncated forms of RWD40 used in Y2H assays (lower panels). The two

LISH-CTLH-containing regions of RWD40 encompass amino acid residues 7 to

95 and 191 to 279, respectively. The WD40 domain encompasses amino acid

residues 388 to 741. LI-CT indicates the LISH-CTLH domain. aa, amino acid

residue.

(B) RWD40 interacts with ROS1 in Y2H assays. The N-terminal region, including

the two LISH-CTLH domains of RWD40, interacts with ROS1.

(C) RWD40 interacts with ROS1 in BiFC assays.

(D) Proteins detected by LC-MS/MS following immunoprecipitation (IP) using an

anti-FLAG antibody, which immunopurifies ROS1, RWD40, RMB1, and RHD1

from transgenic Arabidopsis lines. Results obtained in one to four independent

biological replicates are shown (Rep1–Rep4).

Figure 2. RMB1 and RHD1 interact with RWD40 but not with ROS1.

(A) Interactions among RWD40, RMB1, RHD1, and ROS1 in Y2H assays. BD,

GAL4-binding domain; AD, GAL4-activation domain.

(B) Interactions among RWD40, RMB1, RHD1, and ROS1 in split luciferase

complementation assays. The indicated proteins were fused to the N-terminal or

the C-terminal part of the luciferase protein (nLuc or cLuc, respectively) and

transiently expressed in *Nicotiana benthamiana* leaves. White circles indicate

regions infiltrated with *Agrobacterium*. The luminescent signal indicates a

protein–protein interaction.

(C) Schematic representation of the truncated forms of RWD40 used in the Y2H

assays. WID, WD40-interaction domain. aa, amino acid residue.

(D) The WID domain of RMB1 interacts with the WD40 domain of RWD40 in Y2H

assays.

Figure 3. ROS1, RWD40, RMB1, and RHD1 form a protein complex.

(A) Western blot analysis of gel filtration eluates. The indicated fractions eluted

from the gel filtration column were probed with anti-MYC or anti-FLAG to detect

the epitope-tagged RWD40, RMB1, RHD1, and ROS1. The arrow indicates the

estimated molecular weight of the protein complex. A 120-ml column of HiPrep

16/60 Sephacryl S-300 HR was used. The elution volumes corresponding to the

indicated fraction No. and indicated molecular weight of standards are shown.

(B) Co-expression of full-length *RWD40*, *RMB1*, *RHD*, and a truncated *ROS1*

protein (amino acid residues 1-100) in an insect cell expression system. RWD40,

RMB1, RHD, and the truncated ROS1 were obtained in a single peak fraction.

The positions of elution markers and their molecular masses (in kDa) are

indicated. a.u., arbitrary units. A 24-ml column of Superdex200 10/300 GL was

used for the Sf9 recombinant proteins.

Figure 4. RMB1 binds to methylated DNA.

(A,B) Electrophoretic mobility shift assay (EMSA) showing that the MBD domain

of RMB1 (residues 1 to 81) binds to methylated oligonucleotides corresponding to

the AT5G39160 locus (A) and the MEMS in the *ROS1* promoter (B). M,

methylated; U, unmethylated.

(C) RMB1 binding to labeled methyl-DNA sequence corresponding to the

AT5G39160 locus assessed by microscale thermophoresis (MST). MST binding

assays were used to quantify the interaction of RMB1 with an unmethylated (Um)

probe or probes methylated (m) in CG, CHG, or CHH contexts. The W22G, Y38F,

T49A, or K50T mutations in conserved residues in the MBD domain abolished or

diminished the binding to methylated DNA. The detailed sequence information in

each probe is displayed. The cytosine is labeled in the CHH context in the forward

strand of the mCHH probe, but not in the complementary strand. Values are

means ± SD (n=3). Curves indicate calculated fits; calculated binding affinities are

displayed.

Figure 5. RWD40, RMB1, and RHD1 function in active DNA methylation.

(A) Bisulfite sequencing data showing DNA methylation in different sequence contexts at the MEMS in the *ROS1* promoter in wild-type Col-0 and the indicated mutants.

(B) Relative expression of *ROS1* in the indicated samples. Values are means ± SD of three biological replicates. *P < 0.05, **P < 0.01, compared with wild-type Col-0 plants; NS, not significantly different compared with wild-type Col-0 plants (2-tailed t test).

(C) Enrichment of RWD40, RMB1, RHD1, and ROS1 at the MEMS in the *ROS1* promoter detected by ChIP-qPCR assays. The ChIP signal was quantified relative to input DNA. The no antibody precipitates served as negative control. *Actin 2* was used as a control region. Values are means ± SD of three technical replicates. Results of two biological replicates are displayed. "Rep" indicates replicate.

(D) Effect of mutations in *RWD40*, *RMB1*, and *RHD1* on ROS1 protein enrichment at the MEMS in the *ROS1* promoter by ChIP-qPCR assays.

(E) Chop-PCR showing that *rwd40-1*, *rmb1-1*, *rhd1-1*, and *ros1-4* mutants displayed an increased methylation phenotype at the indicated loci. Amplification of non-digested DNA served as a control.

(F) Analysis of DNA methylation levels at several DNA demethylation target loci in wild-type Col-0 control plants and the indicated mutants by individual locus bisulfite sequencing analysis.

Figure 6. The RWD40 complex regulates gene expression and antibacterial resistance.

(A) Number of differentially expressed genes identified in *rwd40 rmb1 rhd1* and

ros1-4 mutants by mRNA-seq.

(B) Overlap of up-regulated or down-regulated genes between *rwd40 rmb1 rhd 1*

and *ros1-4* mutants.

(C) Validation of the down-regulation of six genes in *rwd40 rmb1 rhd1* and *ros1-4*

mutants by qRT-PCR. Values are means ± SD of 3 biological replicates relative to

transcript levels in wild-type Col-0 plants. *P < 0.05, compared with Col-0 samples

(2-tailed t test).

(D) Phenotypes of Col-0, *ros1-4, rwd40-1, rmb1-1, rhd1-1*, and *rwd40 rmb1 rhd*

plants inoculated with Pto DC3000. Pictures were taken at 3-d post-inoculation

(dpi).

(E) Bacterial counts in Col-0 and the indicated mutants plants at 3 dpi. **P < 0.01,

compared with Col-0 samples (2-tailed t test).

Figure 7. Schematic model showing that RWD40 may inhibit its own

function by targeting the MEMS at the ROS1 promoter.

The RWD40 complex decreases the DNA methylation level of the MEMS site at

the *ROS1* promoter, causing reduced *ROS1* expression. The reduced *ROS1*

expression would hinder the DNA demethylation function of the RWD40 complex

at target genomic loci.

Figure S1. DNA demethylation phenotypes o*f rwd40-1* a*nd rwd40-2* mutants

and mutant complementation.

(A) RWD40 localization in *Nicotiana benthamiana.*

(B) T-DNA insertion position in *rwd40-1* and *rwd40-2* mutants. Boxes and lines

denote exons and introns, respectively.

(C) Expression of *RWD40* in mutants relative to that in wild-type Col-0 control

plants, as measured by qRT-PCR. Values are means ± SD of three biological

replicates.

(D) Western blot analysis of RWD40-3xFLAG in T2 *rwd40-1* transgenic lines.

(E) DNA methylation levels in *rwd40* mutant alleles and complementation lines at

the AT5G39160 locus as determined by Chop-PCR.

(F) DNA methylation levels at the AT5G39160 locus in *rwd40-1* and *rwd40-2*

mutants as determined by individual bisulfite sequencing.

Figure S2. Characterization of RMB1 and RHD1.

(A) RMB1 domains and mutants. Upper panel, schematic representation of the

RMB1 protein. CBD, C (AT)-rich DNA-binding domain. Middle panel, T-DNA

insertion positions in *rmb1-1* and *rmb1-2* mutants. Boxes and lines denote exons

and introns, respectively. Lower panel, RMB1 transcript levels in the mutants.

(B) Schematic representation of the RMB1 protein (upper panel), T-DNA insertion

position in the *rhd1-1* mutant (middle panel), and RHD1 transcripts levels in the

rhd1-1 mutant by RT-PCR detection (lower panel).

(C) Interactions among RWD40, RMB1, and RHD1 in BiFC assays.

(D) DNA methylation levels in *rmb1* mutant alleles and in complementation lines at

the AT5G39160 locus as determined by Chop-PCR.

(E) DNA methylation levels at the AT5G39160 locus in *rmb1-1* and *rmb1-2*

mutants as determined by individual locus bisulfite sequencing.

(F) DNA methylation levels in wild-type Col-0 and indicated mutants at the

AT5G39160 locus (upper panel), and in *rhd1* mutant alleles and complementation

line at the AT4G18380 locus as determined by Chop-PCR (lower panel).

Figure S3. Interactions among RWD40, RMB1, RHD1, and ROS1 in Y2H

assays.

(A) Schematic representation of the truncated forms of ROS1 used in the Y2H

assays and of the ELM domain identified in N-terminal regions of ROS1.

HhH-GPD, hallmark Helix-hairpin-helix and Gly/Pro-rich loop domain; EndIIII,

endonuclease III; Perm-CXXC, permuted version of a single unit of the zinc

finger-CXXC; RRM_DME, RRM-fold domain present at the C-terminus of

Demeter-like glycoslyases; EAR, ethylene response factor-associated amphiphilic

repression domain; ELD, EAR-like motif.

(B) The ELD domain of ROS1 interacts with RWD40 in Y2H assays.

(C) Schematic representation of the truncated forms of RHD1 used in Y2H

assays.

(D) RHD1 interacts with RHD1 through the zinc-finger domain, and the WD40

domain of RWD40 interacts with the zinc-finger domain of RHD1 in Y2H assays.

(E) RMB1 interacts with full-length RHD1 but does not interact with its zinc-finger

domain or homeodomain.

Figure S4. Analysis of MBD domain-containing proteins in *Arabidopsis.*

(A) Phylogenetic tree of MBD domain-containing proteins in *A. thaliana.*

(B) Alignment of MBD domains in MBD domain-containing proteins in *A. thaliana*.

The amino acid residues targeted for mutations in the MBD domain are indicated

by triangles.

Figure S5. Mutation of the MBD domain reduces the binding of RMB1 to

methylated DNA and abolishes the demethylation function of RMB1.

(A) DNA methylation level at AT5G39160 in the *rmb1-1* mutant and in the *rmbd1-1*

mutant expressing WT (wild-type) or K50T-mutated RMB1 as determined by

Chop-PCR.

(B,C) EMSA of the MBD domain of MBD7 (C) and IDM1 (D) binding to methylated

oligonucleotides corresponding to the AT5G39160 locus. M, methylated. Um,

unmethylated.

Figure S6. Dysfunction of RWD40, RMB1, or RHD1 does not cause silencing

of the *35S:SUC2* transgene.

(A) Chop-PCR showing that *rwd40-1*, *rmb1-1*, and *rhd1-1* mutants do not display

an increased methylation phenotype in IDM1-dependent DNA demethylation

target loci (Qian et al. 2012). Amplification of non-digested DNA served as a

control.

(B) Root phenotype in *rwd40-1*, *rmb1-1*, and *rhd1-1* in the *35S:SUC2* transgenic

background.

(C) Relative expression of the *35S:SUC2* transgene in the indicated mutants.

Values are relative to transcript levels in *35S:SUC2* control plants and are means

± SD of three biological replicates. *P < 0.01, compared with *35S:SUC2* plants;

NS, not significantly different compared with *35S:SUC2* plants (2-tailed t test).

(D) Screenshots of DNA cytosine methylation status in Col-0, *idm1-2*, and *ros1-4*,

and screenshots of H2A.Z deposition and H3K18Ac in wild-type control plants

(Nie et al., 2019) at the tested loci in Figure 5. AT1G62760 is an *IDM1*-dependent

active DNA demethylation target (Qian et al., 2012, Nie et al., 2019), and serves

as a positive control of the co-existence of H2A.Z and H3K18ac.

Figure S7. Images of complete Y2H plates (from Figures 1B, 2A, 2D, and

S3B-E).

Figure 1

D

		ROS1 (AT2G36490)		RWD40 (AT2G25420)		RMB1 (AT1G63240)		RHD1 (AT5G42780)	
		Score	Unique peptides	Score	Unique peptides	Score	Unique peptides	Score	Unique peptides
IP ROS1	Rep.1	5574.0	83	152.0	10	81.0	8	55.0	4
	Rep.2	3643.0	86	139.0	11	148.0	4	78.0	4
IP RWD40	Rep.1	816.0	22	605.0	16	202.0	7	205.0	9
IP RMB1	Rep.1	76.0	4	71.0	3	902.0	20	69.0	3
	Rep.2	1371.0	48	176.0	8	591.0	13	147.0	7
IP RHD1	Rep.1	94.0	7	242.0	9	106.0	4	1085.0	14
IP Col-0 control	Rep.1	0	0	0	0	0	0	0	0
	Rep.2	0	0	0	0	0	0	0	0
	Rep.3	0	0	0	0	0	0	0	0
	Rep.4	0	0	0	0	0	0	0	0

Figure 2

Figure 3

A

Gel filtration

Fraction No. 6 8 10 12 14 16 18 20 22 24 26 28 30

ROS1-MYC

RWD40-FLAG

RMB1-MYC

RHD1-MYC

350kDa

The elution volumes corresponding to indicated fraction No.

Fraction No.	12	14	16
Elution Volumes (ml)	54	58	62

The elution volumes corresponding to indicated molecular weight of standards

Molecular Weight (KDa)	443	350	150
Elution Volumes (ml)	53	58	68

B

Figure 4

A AT5G39160 locus

	-	+	+	+	+	+	+	+	+	+	RMB1-MBD
+	-	-	-	-	-	-	-	-	-	Um probe
-	+	+	+	-	-	-	-	-	-	mCG probe
-	-	-	-	+	+	+	-	-	-	mCHG probe
-	-	-	-	-	-	-	+	+	+	mCHH probe
-	-	M	U	-	M	U	-	M	U	Competitor

←bound
←free

B MEMS in the *ROS1* promoter

	-	+	+	+	+	+	+	+	+	+	RMB1-MBD
+	-	-	-	-	-	-	-	-	-	Um probe
-	+	+	+	-	-	-	-	-	-	mCG probe
-	-	-	-	+	+	+	-	-	-	mCHG probe
-	-	-	-	-	-	-	+	+	+	mCHH probe
-	-	M	U	-	M	U	-	M	U	Competitor

←bound
←free

C

MBD	dsDNA	Kd (μM)
WT	mCG	1.77±0.70
WT	mCHG	3.34±0.27
WT	mCHH	2.86±1.21
WT	Um	>5.27

MBD	dsDNA	Kd (μM)
WT	mCG	1.77±0.70
WT	Um	>5.27
W22G	mCG	13.28±4.67
Y38F	mCG	2.15±0.75
T49A	mCG	3.58±2.07
K50T	mCG	No binding

MBD	dsDNA	Kd (μM)
WT	mCHG	3.34±0.27
WT	unM	>5.27
Y38F	mCHG	5.16±2.00
T49A	mCHG	4.11±2.92
K50T	mCHG	No binding

MBD	dsDNA	Kd (μM)
WT	mCHH	2.86±1.21
WT	unM	>5.27
W22G	mCHH	No binding
Y38F	mCHH	No binding
T49A	mCHH	No binding
K50T	mCHH	No binding

AT5G39160 Um probe F	TTTGGTACGGACTTTTAGTTACAAGAGAGCATTTAAACAAAAATGACCGGTAGATGGAA
AT5G39160 Um probe R	TTCCATCTACCGGTCATTTTTGTTTAAATGCTCTCTTGTAACTAAAAGTCCGTACCAAA
AT5G39160 mCG probe F	TTTGGTA**^mC**GGACTTTTAGTTACAAGAGAGCATTTAAACAAAAATGAC**^mC**GGTAGATGGAA
AT5G39160 mCG probe R	TTCCATCTAC**^mC**GGTCATTTTTGTTTAAATGCTCTCTTGTAACTAAAAGTC**^mC**GTACCAAA
AT5G39160 mCHG probe F	TTTGGTACGGACTTTTAGTTACAAGAGAGCATTTAAACAAAAATGA**^mC**CGGTAGATGGAA
AT5G39160 mCHG probe R	TTCCATCTA**^mC**CGGTCATTTTTGTTTAAATGCTCTCTTGTAACTAAAAGT**^mC**CGTACCAAA
AT5G39160 mCHH probe F	TTTGGTACGGA**^mC**TTTTAGTTA**^mC**AAGAGAG**^mC**ATTTAAA**^mC**AAAAATGACCGGTAGATGGAA
AT5G39160 mCHH probe R	TTCCATCTACCGGTCATTTTTGTTTAAATGCTCTCTTGTAACTAAAAGTCCGTACCAAA

Figure 5

Figure 6

A

	up-regulated genes	down-regulated genes
rwd40 rmb1 rhd1	3,817	4,028
ros1-4	2,970	2,918

B

Up-regulated DEGs

ros1-4 (n=2,970)

1,297   1,673   2,144

rwd40 rmb1 rhd1 (n=3,817)

Down-regulated DEGs

ros1-4 (n=2,918)

977   1,941   2,087

rwd40 rmb1 rhd1 (n=4,028)

C

Col-0
rwd40 rmb1 rhd1
ros1-4

Relative expression

AT2G06002 AT3G30720 AT1G11785 AT1G50040 AT2G03640 AT3G46370

D

Col-0   *ros1-4*   *rwd40-1*

rmb1-1   *rhd1-1*   *rwd40 rmb1 rhd1*

E

$\log_{10}$(CFU/mg)

Col-0 *ros1-4* *rwd40-1* *rmb1-1* *rhd1-1* *rwd40 rmb1 rhd1*

Figure 7

Figure S1

A

pDEX::RWD40-YFP

35S::GFP:RWD40

B

AT2G25420

rwd40-1
(SALK_068825)

rwd40-2
(SALK_012947)

C

RWD40

Relative expression

Col-0 rwd40-1 rwd40-2

1.20
0.80
0.40
0.00

D

Col-0 *rwd40-1/RWD40-3xFLAG*(T2)

#7 #8 #15 #16 #19 #21

Anti-FLAG

β-Actin

E

Col-0 *idm1-2* *ros1-4* *rwd40-1* *rwd40-2* *rwd40-1/ RWD40-3xFLAG*

#8 #15

AT5G39160
HpaII

Undigested DNA

F

AT5G39160 locus

DNA methylation level

100%
80%
60%
40%
20%
0%

CG CHG CHH C

■ Col-0
■ *idm1-2*
■ *ros1-4*
■ *rwd40-1*
■ *rwd40-2*

Figure S2

Figure S3

A

ROS1 (50-65aa) :50-GNKDMSGLDHLSFGDLL-65
NINJA (4-15aa):4 -DN – – – –GLE –LSLG –LSS-15

B

E

C

D

Figure S4

A

B

Figure S5

A

| | Col-0 | ros1-4 | rmb1-1 | K50T | WT |
| | | | | *rmb1-1* | |

AT5G39160
HpaII digested

Undigested
Loading control

B

AT5G39160 locus

-	+	+	+	+	+	MBD7-MBD
+	-	-	-	-	+	Um probe
-	+	+	-	-	-	mCG probe
-	-	-	+	-	-	mCHG probe
-	-	-	-	+	-	mCHH probe
-	-	M	-	-	-	Competitor

←bound

←free

C

AT5G39160 locus

-	+	+	+	+	+	IDM1-MBD
+	-	-	-	-	+	Um probe
-	+	+	-	-	-	mCG probe
-	-	-	+	-	-	mCHG probe
-	-	-	-	+	-	mCHH probe
-	-	M	-	-	-	Competitor

←bound

←free

Figure S6

Figure S7 Full-size images of Y2H shown in this study

Figure 1B

Figure S7 Continued

Figure 2A

Figure S7 Continued

Figure 2D

Figure S7 Continued

Figure S3B

BD/AD	LT	LTH+3AT	LTHA+3AT
ROS1/RWD40_LI-CT-1st			
ROS1/RWD40_2LI-CT			
ROS1_N/RWD40			
ROS1_ELD/RWD40			
ROS1/RWD40_WD40			
ROS1_C/RWD40			
ROS1/Vec			
ROS1_N/Vec			
ROS1_ELD/Vec			
ROS1_C/Vec			
Vec/RWD40			
Vec/RWD40_LI-CT-1st			
Vec/RWD40_WD40			
Vec/Vec			

Figure S7 Continued

Figure S3D

Figure S7 Continued

Figure S3D

BD/AD	LT	LTH+3AT	LTHA+3AT
RHD1/RHD1_P1			
RHD1/RHD1_P2			
RHD1/RHD1_P3			
RHD1/RHD1_P4			
RHD1/RHD1_P5			
RHD1_P3/RHD1			
RWD40/RHD1_P1			
RWD40/RHD1_P2			
RWD40/RHD1_P3			
RWD40/RHD1_P4			
RWD40/RHD1_P5			
RWD40_WD40/RHD1_P2			
Vec/RHD1_P1			
Vec/RHD1_P2			
Vec/RHD1_P3			
Vec/RHD1_P4			
Vec/RHD1_P5			
RWD40/Vec			

Figure S7 Continued

Figure S3E