1 ***In silico* integration of thousands of epigenetic datasets into 707 cell type regulatory**
2 **annotations improves the trans-ethnic portability of polygenic risk scores**
3

4 Tiffany Amariuta*[1-5], Kazuyoshi Ishigaki*[1-3,6], Hiroki Sugishita[7], Tazro Ohta[8,9], Koichi
5 Matsuda[10,11], Yoshinori Murakami[12], Alkes L. Price[3,13,14], Eiryo Kawakami[8,15], Chikashi Terao[6,16,17],
6 Soumya Raychaudhuri[1-5,18]
7
[1]Center for Data Sciences, Harvard Medical School, Boston, Massachusetts, 02115, USA.
[2]Divisions of Genetics and Rheumatology, Department of Medicine, Brigham and Women's Hospital, Harvard
Medical School, Boston, Massachusetts, 02115, USA.
[3]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, Massachusetts,
02142, USA.
[4]Department of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts, 02115, USA.
[5]Graduate School of Arts and Sciences, Harvard University, Cambridge, Massachusetts, 02138, USA.
[6]Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences, Kanagawa,
230-0045 Japan.
[7]Laboratory for Developmental Genetics, RIKEN Center for Integrative Medical Sciences (IMS), Kanagawa, Japan
[8]Medical Sciences Innovation Hub Program, RIKEN, Kanagawa, Japan.
[9]Database Center for Life Science, Joint Support-Center for Data Science Research, Research Organization of
Information and Systems, Shizuoka, Japan.
[10]Laboratory of Genome Technology, Human Genome Center, Institute of Medical Science, The University of
Tokyo, Tokyo, 108-8639 Japan.
[11]Laboratory of Clinical Genome Sequencing, Department of Computational Biology and Medical Sciences,
Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, 108-8639 Japan.
[12]Division of Molecular Pathology, Institute of Medical Science, The University of Tokyo, Tokyo, 108-8639 Japan
[13]Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA.
[14]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA 02115, USA.
[15]Artificial Intelligence Medicine, Graduate School of Medicine, Chiba University, Chiba, Japan.
[16]Clinical Research Center, Shizuoka General Hospital, Shizuoka, 420-8527 Japan.
[17]The Department of Applied Genetics, The School of Pharmaceutical Sciences, University of Shizuoka, Shizuoka,
422-8526 Japan.
[18]Centre for Genetics and Genomics Versus Arthritis, Centre for Musculoskeletal Research, Manchester Academic
Health Science Centre, The University of Manchester, Manchester, UK.
34
35 *indicates equal contributions
36 **Correspondence to Soumya Raychaudhuri, soumya@broadinstitute.org

Abstract

Poor trans-ethnic portability of polygenic risk score (PRS) models is a critical issue that may be

partially due to limited knowledge of causal variants shared among populations. Hence,

leveraging noncoding regulatory annotations that capture genetic variation across populations

has the potential to enhance the trans-ethnic portability of PRS. To this end, we constructed a

unique resource of 707 cell-type-specific IMPACT regulatory annotations by aggregating 5,345

public epigenetic datasets to predict binding patterns of 142 cell-type-regulating transcription

factors across 245 cell types. With this resource, we partitioned the common SNP heritability of

diverse polygenic traits and diseases from 111 GWAS summary statistics of European (EUR,

average N=180K) and East Asian (EAS, average N=157K) origin. For 95 traits, we were able to

identify a single IMPACT annotation most strongly enriched for trait heritability. Across traits,

these annotations captured an average of 43.3% of heritability (se = 13.8%) with the top 5% of

SNPs. Strikingly, we observed highly concordant polygenic trait regulation between

populations: the same regulatory annotations captured statistically indistinguishable SNP

heritability (fitted slope = 0.98, se = 0.04). Since IMPACT annotations capture both large and

consistent proportions of heritability across populations, prioritizing variants in IMPACT

regulatory elements may improve the trans-ethnic portability of PRS. Indeed, we observed that

EUR PRS models more accurately predicted 21 tested phenotypes of EAS individuals when

variants were prioritized by key IMPACT tracks (49.9% mean relative increase in R^2). Notably,

the improvement afforded by IMPACT was greater in the trans-ethnic EUR-to-EAS PRS

application than in the EAS-to-EAS application (47.3% vs 20.9%, P < 1.7e-4). Overall, our study

identifies a crucial role for functional annotations such as IMPACT to improve the trans-ethnic

portability of genetic data, and this has important implications for future risk prediction models

that work across populations.

Introduction

An important challenge for complex trait genetics is that there is no clear framework to transfer

population-specific genetic data, such as GWAS results, to individuals of other ancestries[1–3]. The

importance of this challenge is accentuated by the fact that 80% of all genetic studies have

been performed using individuals of European ancestry, accounting for a minority of the

world's population[4]. This is exacerbated by the fact that population-specific linkage

disequilibrium (LD) between variants confounds inferences about causal cell types and variants

(**Figure 1A**)[5–7]. GWAS have the potential to revolutionize the clinical application and utility of

genetic data to the individual, exemplified by current polygenic risk score (PRS) models[5,8–16].

However, while the utility of PRS models relies on accurate estimation of allelic effect sizes

from GWAS and benefits from genetic similarity between the test cohort and the GWAS cohort,

recent studies have explicitly observed a lack of trans-ethnic portability[2,3,5,8,17,18]. Previous

studies have extensively shown that functional annotations can improve PRS models when

learned and applied to the same population[19,20], by introducing biologically-relevant priors on

causal effect sizes and compensating for inflation of association statistics by LD. However, the

potential for functional annotations to improve trans-ethnic PRS frameworks, where the

influences of population-specific LD are more profound, has not yet been extensively

investigated.

However, designing functional annotations that may improve PRS models is challenging.

While the genetic variation of a complex trait likely regulates diverse biological mechanisms

genome-wide, such functional annotations must strike a balance of specificity, comprehensively

but precisely capturing large regulatory programs. Pinpointing these mechanisms is especially

difficult as genome-wide association studies (GWAS) have identified thousands of genetic

associations with complex phenotypes[8,21–23]. It has been estimated that about 90% of these

associations reside in protein noncoding regions of the genome, making their mechanisms

difficult to interpret[24,25]. Defining the etiology of complex traits and diseases requires

knowledge of phenotyping-driving cell types in which these associated variants act.

Transcription factors (TFs) are poised to orchestrate large polygenic regulatory programs as

genetic variation in their target regions can modulate gene expression, often in cell-type-

specific contexts[26,27]. Genomic annotations marking the precise location of TF-mediated cell

type regulation can be exploited to elucidate the genetic basis of polygenic traits. However,

currently there is no comprehensive catalogue of the binding profiles of the approximately

1,600 human TFs in every known cell type[28]. Moreover, existing TF ChIP-seq datasets are

limited to factors with effective antibodies and suffer from inter-experimental variation, noise,

and genomic bias[29,30].

To overcome these challenges, we previously developed IMPACT, a genome-wide cell-

type-specific regulatory annotation strategy that models the epigenetic pattern around active

TF binding using linear combinations of functional annotations[31]. In rheumatoid arthritis (RA),

IMPACT CD4+ T cell annotations captured substantially more heritability than functional

annotations derived from single experiments, including TF and histone modification ChIP-seq[6].

In this study, we expanded this approach by aggregating 5,345 functional annotations with

IMPACT to create a powerful and generalizable resource of 707 cell-type-specific gene

regulatory annotations (**Web Resources**) based on binding profiles of 142 TFs across 245 cell

types (**Figure 1B,C**). This study builds on our previous study introducing IMPACT, in which we

created only 13 annotations (13 TFs) based on 515 functional annotations. Assuming that causal

variants are largely shared between populations[2,21], we hypothesized that restricting PRS

models to variants within trait-relevant IMPACT annotations, which are more likely to have

regulatory roles and less likely to be confounded by LD, will especially improve their trans-

ethnic portability.

In this study, we identify key IMPACT regulatory annotations that capture genome-wide

polygenic mechanisms underlying a diverse set of complex traits, supported by enrichments of

genetic heritability, multi-ethnic marginal effect size correlation (a mechanism of improved

PRS), and improved trans-ethnic portability of PRS models (**Figure 1D**). Here, we defined and

employed our compendium of 707 IMPACT regulatory annotations to study polygenic traits and

diseases from 111 GWAS summary datasets of European (EUR) and East Asian (EAS) origin.

Assuming shared causal variants between populations, annotations that prioritize shared

regulatory variants must (1) capture disproportionately large amounts of genetic heritability in

both populations, (2) be enriched for multi-ethnic marginal effect size correlation, and (3)

improve the trans-ethnic applicability of population-specific PRS models. Using our

compendium of regulatory annotations, we identified key annotations for each polygenic trait

and demonstrated their utility in each of these three applications toward prioritization of

shared regulatory variants. Overall, this work improves the interpretation and trans-ethnic

portability of genetic data and provides implications for future clinical implementations of risk

prediction models.

Figure 1

Figure 1 legend. Study design to identify regulatory annotations that prioritize regulatory variants in a multi-ethnic setting. A) Population-specific LD confounding and subsequent inflation of GWAS associations complicate the interpretation of summary statistics and transferability to other populations; functional data may help improve trans-ethnic genetic portability. B) Prism of functional data in IMPACT model: 707 genome-wide TF occupancy profiles (green), 5,345 genome-wide epigenomic feature profiles (blue), and fitted weights for these features (pink) to predict TF binding by logistic regression. Using IMPACT annotations, we investigate 111 GWAS summary datasets (yellow) of EUR and EAS origin. C) Compendium of 707 genome-wide cell-type-specific IMPACT regulatory annotations. D) Annotations that prioritize common regulatory variants must I) capture large proportions of heritability in both populations, II) account for consistent effect size estimations between populations and III) improve the trans-ethnic application of PRS.

Results

Building a compendium of *in silico* gene regulatory annotations

To capture genetic heritability of diverse polygenic diseases and quantitative traits, we

constructed a comprehensive compendium of 707 cell type regulatory annotation tracks. To do

this, we applied the IMPACT[31] framework to 707 unique TF-cell type pairs obtained from a total

of 3,181 TF ChIP-seq datasets from NCBI, representing 245 cell types and 142 TFs (**Figure 1B**,

Online Methods, **Web Resources**, **ST1**, **SF1**)[32]. Briefly, IMPACT learns an epigenetic signature of

active TF binding evidenced by ChIP-seq, differentiating bound from unbound TF sequence

motifs using logistic regression. We derive this signature from 5,345 epigenetic and sequence

features, predominantly generated by ENCODE[33] and Roadmap[34] (**Online Methods**, **ST2**); these

data were drawn from diverse cell types, representing the biological range of the 707 candidate

models. IMPACT then probabilistically annotates the genome, e.g. on a scale from 0 to 1,

without using the TF motif, identifying regulatory regions that are similar to those that the TF

binds.

To assess the specificity of our IMPACT annotations, we test whether they (1) accurately

predict binding of the modeled TF, (2) share cell-type-specific characteristics with other tracks

of the same cell type, and (3) score cell-type-specifically expressed genes higher than

nonspecific genes. The 707 models that we defined had a high TF binding prediction accuracy

with mean AUPRC = 0.74 (se = 0.008, **SF2**) using cross-validation. Annotations segregated by cell

type rather than by TF, excluding CTCF, suggesting a single TF may bind to different enhancers

in different cell types (**Figure 2A**). Annotations of the same cell types were more strongly

correlated genome-wide (Pearson r = 0.56, se = 0.06) than annotations of different cell types

(Pearson r = 0.48, se = 0.003, difference of means P < 0.03, **SF2**). Furthermore, the covariance

structure between TF ChIP-seq training datasets is similar to that of corresponding IMPACT

annotations, indicating that the IMPACT model does not introduce spurious correlations among

unrelated ChIP-seq datasets (**SF2**). Lastly, for nine different cell types, we examined cell-type-

specifically expressed genes from Finucane et al[35] and corresponding differential expression t-

statistics. We observed significantly larger IMPACT probabilities at SNPs in and near these genes

(mean = 0.062, se = 0.011) compared to genes that were generally expressed (mean = 0.045, se

= 0.006; difference of means P = 0.024, **Figure 2B**, **SF2**, **Online Methods**), suggesting that

IMPACT annotates relevant cell type regulatory elements.

Partitioning common SNP heritability of 111 GWAS summary statistics in EUR and EAS

We obtained summary statistics from 111 publicly available GWAS summary statistics

for diverse polygenic traits and diseases. Throughout the text, we use five randomly selected

traits to exemplify our results: asthma, RA, prostate cancer (PrCa), mean corpuscular volume

(MCV), and body height. These included 69 from EUR participants[31,36] (average N = 180K,

average heritability z-score = 12.9, 41/69 from UK BioBank)[6,37] and 42 from EAS participants of

BioBank Japan[3,38–40] (average N = 157K, average heritability z-score = 6.6)[22] (**ST3**). All of the

summary statistics used were generated from studies that had a sample size greater than

10,000 individuals and also had a significantly non-zero heritability (z-score > 1.97). There are

29 phenotypes for which we obtained summary statistics in both EUR and EAS. Although 10/29

Figure 2

Figure 2 legend. IMPACT annotates relevant cell type regulatory elements. A) Low-dimensional embedding and clustering of 707 IMPACT annotations using uniform manifold approximation projection (UMAP). Annotations colored by cell type category; TF groups indicated where applicable. B) IMPACT annotates cell type specifically expressed genes with higher scores than nonspecific genes. C) Biologically distinct regulatory modules revealed by cell type-trait associations with significantly nonzero τ^* across 20 of 707 IMPACT regulatory annotations and 5 representative EUR complex traits, color indicates $-\log_{10}$ FDR 5% adjusted P value of τ^*. D) Lead IMPACT annotations capture more heritability than lead cell-type-specific histone modifications across 60 of 69 EUR summary statistics for which a lead IMPACT annotation was identified. * indicates heritability estimate difference of means $P < 0.05$.

traits have a multi-ethnic genetic correlation (R_g) significantly less than 1 ($P < 0.05/29$ tested

traits), overall we observed high R_g for most traits, supporting our assumption that causal

variants are generally shared across populations (**Online Methods**, **SF3**)[41]. At two extremes,

basophil count has a low multi-ethnic R_g of 0.32 (sd = 0.10), while atrial fibrillation has a high

multi-ethnic R_g of 0.98 (sd = 0.11), consistent with previous observations made using *Popcorn*,

but using different parameter estimation strategies (**Online Methods**)[3].

We then partitioned the common SNP (minor allele frequency (MAF) > 5%) heritability

of these 111 datasets using S-LDSC[6] with an adapted baseline-LD model excluding cell-type-

specific annotations[31,36] (**SF3, Online Methods**). Next, we tested each of the traits against each

of the 707 IMPACT annotations, assessing the significance of a non-zero τ^*, which is defined as

the proportionate change in per-SNP heritability associated with a one standard deviation

increase in the value of the annotation (**Online Methods**)[36]. We observed that 95 phenotypes

had at least one significant annotation-trait association ($\tau^* > 0$, two-tailed FDR < 5%, **Ext. Data**

1, Online Methods, **ST4-8**). Here, we highlight associations with EUR summary statistics for the

five exemplary phenotypes mentioned above: asthma, RA, PrCa, MCV, and height (**Figure 2C**).

Consistent with known biology, B and T cells were strongly associated with asthma[42], RA[43], and

MCV[44,45] while other blood cell regulatory annotations predominantly derived from GATA

factors were also associated with MCV. Prostate cancer cell lines were associated with PrCa,

while many cell types including myoblasts[46], fibroblasts[47], and adipocytes[48,49], lung cells, and

endothelial cells were associated with height, perhaps related to musculo-skeletal

developmental pathways.

For each trait, we defined the lead IMPACT regulatory annotation as the annotation

capturing the greatest per-SNP heritability, e.g. the largest while significant τ^* estimate (**ST9**).

With their top 5% of SNPs, lead IMPACT annotations captured an average of 43.3% heritability

(se = 13.8%) across these 95 polygenic traits (**SF4**, **Online Methods**), with more than 25% of

heritability captured for two-thirds of the tested summary statistics (73/111 traits) and more

than 50% captured for 28% (31/111). Returning to our five exemplary phenotypes, with the top

5% of EUR SNPs, IMPACT captured 97.1% (sd = 17.6%) of asthma heritability with the T-bet Th1

annotation, 65.9% (sd = 12.1%) of RA heritability with the B cell TBP annotation, 60.4% (sd =

8.9%) of PrCa heritability with the prostate cancer cell line (LNCAP) TFAP4 annotation, 72.4%

(sd = 6.0%) of MCV heritability with the GATA1 PBMC annotation, and lastly 31.6% (sd = 3.0%)

of height heritability with the lung MXI1 annotation (**Figure 2D**). While the observed association

between lung and height is not intuitive, within the MXI1 gene lies a genome-wide significant

variant associated with height[50]. Moreover, we captured significantly more heritability across

EUR traits using our expanded set of 707 IMPACT annotations (mean = 49.5%, se = 12.0%)

compared to the 13 annotations in our previous study (mean = 32.3%, se = 1.3%, difference of

means P = 0.02).

To demonstrate the value of IMPACT tracks, we compared them to annotations derived

from single experimental assays. For example, since each of the IMPACT tracks was trained on

TF ChIP-seq data, we directly compared the heritability captured by both data types. We

observed that the heritability captured by lead IMPACT annotations (mean τ^* = 3.53, se = 0.91)

was significantly greater than by the analogous TF ChIP-seq used in training (mean τ^* = 1.71, se

= 0.94, difference of means P = 0.02). We also compared IMPACT tracks to histone marks, which

are commonly used to quantify cell type heritability[6]. From 220 publicly available cell-type-

specific histone mark ChIP-seq annotations of EUR SNPs[6], we selected the lead histone mark

track for each of 69 EUR summary statistics. Restricting to the top 5% of SNPs, we observed that

the mean EUR heritability captured by lead IMPACT annotations (49.5%, se = 12.0%) was

significantly greater than by lead histone mark annotations (28.4%, se = 9.0%, difference of

means P = 0.02) (**Figure 2D, ST10**). For example, the lead IMPACT annotation for asthma

captured 1.5x more heritability than the best histone mark annotation (H3K27ac in CD4+ Th2),

capturing 64.2% (sd = 15.5%) of heritability. Similarly, IMPACT captured 1.7x more RA

heritability than H3K4me3 in CD4+ Th17s; IMPACT captured 1.4x more MCV heritability than

H3K4me3 in CD34+ cells; IMPACT captured 2.3x more PrCa heritability than H3K4me3 in CD34+

cells; and IMPACT captured 3.1x more height heritability than H3K4me3 in lung cells. In terms

of τ^*, IMPACT also captured more per-SNP heritability than histone marks: mean τ^* fold

change = 1.38x (**SF5**).

Since some of our IMPACT annotations are similar to each other (**SF2**), we performed

serial conditional analyses in order to identify IMPACT annotations explaining heritability

independently from one another (**Online Methods**). This strategy might identify complex traits

for which several distinct biological mechanisms are independently regulated by genetic

variation. Indeed, we identified 30 EUR phenotypes and 8 EAS phenotypes with multiple

independent IMPACT associations (**SF6, ST11-12**). For example, four annotations were

independently associated with EUR PrCa: prostate (TFAP4), prostate (RUNX2), mesendoderm

(PDX1), and cervix (NFYB). Moreover, for seven EUR traits, three IMPACT annotations were

independently associated: height (adipocytes, fibroblasts, lung), neutrophil count (monocytes,

adipocytes, B cells), osteoporosis (myoblasts, mesenchymal stem cells, cervix), IBD (T cells and

two B cell annotations), platelet count (PBMCs, hematopoietic progenitors, muscle), systolic

blood pressure (endothelial, mesenchymal stem cells, fibroblasts), and white blood cell count (B

cells, adipocytes, hematopoietic progenitors). We found that the heritability z-score, an index

correlated with the power of S-LDSC[6], is strongly predictive of the number of independent

regulatory associations (linear regression P < 5.4e-9), while sample size is not (linear regression

P = 0.19) (**SF7**). Our findings suggest that multiple independent regulatory programs can

contribute to the heritability of complex traits, and we can detect them when phenotypes are

sufficiently heritable and the GWAS provide accurate effect size estimation.

Concordance of polygenic regulation between European and East Asian populations

Previous studies have shown concordance of polygenic effects between EUR and EAS

individuals in RA[1] and between EUR and African American individuals in PrCa[51]. However, to our

knowledge, the extent of these shared effects has not yet been comprehensively investigated

across many functional annotations and in diverse traits. Here, we quantified the SNP

heritability (τ*) of 29 traits in EUR and EAS captured by a set of approximately 100 independent

IMPACT regulatory annotations (**Figure 3B, SF8, Online Methods**). Assuming shared causal

variants in EUR and EAS, IMPACT annotations that best prioritize shared genomic regions

regulating a phenotype presumably also disproportionately capture similar amounts of

heritability in both EUR and EAS (**Figure 1D-I**, **Figure 3A**). Briefly, we selected independent

annotations using an iterative pruning approach: for each trait, we ranked all annotations by τ*

and removed any annotation correlated with Pearson r > 0.5 to the lead annotation and then

repeated. As IMPACT annotations are independent of population-specific factors including LD

and allele frequencies (**SF3**), they are poised to capture the genome-wide distribution of

regulatory variation in a population-independent manner. We observed that τ* estimates

across annotations for EUR and EAS are strikingly similar, with a regression coefficient that is

Figure 3

Figure 3 legend. Multi-ethnic concordance of regulatory elements defined by IMPACT. A) Illustrative concept of concordance versus discordance of τ^* between populations. Concordance implies a similar distribution of causal variants and effects captured by the same annotation. The implications of discordant τ^* are not as straightforward. B) Common per-SNP heritability (τ^*) estimate for sets of independent IMPACT annotations across 29 traits shared between EUR and EAS. Left: color indicates τ^* significance (τ^* greater than 0 at 5% FDR) in both populations (blue), significant in only EUR (green), significant in only EAS (red), significant in neither (gray). Line of best fit through annotations significant in both populations (dark purple line, 95% CI in light purple). Black dotted line is the identity line, y = x. Right: color indicates association to one of five exemplary traits. C) Heterogeneity test at 5% FDR for annotation-trait associations between EUR and EAS. Color indicates significance of difference of means *P* value. D) Heterogeneity test reveals 2.2% of all annotation-trait associations with significantly discordant τ^* estimates between populations.

consistent with identity (slope = 0.98, se = 0.04). For example, we observed a strong Pearson

correlation of τ^* between EUR and EAS for asthma (r = 0.98), RA (r = 0.87), MCV (r = 0.96), PrCa

(r = 0.90), and height (r = 0.96). Furthermore, we found that 97.8% of our τ^* estimates have no

evidence of population heterogeneity (FDR P > 0.05) (**Figure 3C**). Among our five representative

traits, we observed only one instance of heterogeneity, in which the B cell SRF IMPACT

annotation captured RA heritability significantly more in EUR than in EAS (EUR τ^* = 1.20 (se =

0.40), EAS τ^* = -1.06 (se = 0.46), difference of means P < 2.0e-4). Overall, our results suggest

that regulatory variants in EUR and EAS populations are equally enriched within the same

classes of regulatory elements. This does not exclude the possibility of population-specific

variants or causal effect sizes, as evidenced by 10 traits with multi-ethnic genetic correlation

significantly less than 1. Rather, these results suggest that causal biology, including disease-

driving cell types and their regulatory elements, underlying polygenic traits and diseases, is

largely shared between these populations.

Models incorporating IMPACT functional annotations improve the trans-ethnic portability of

polygenic risk scores

PRS models have great clinical potential: previous studies have shown that individuals

with higher PRS have increased risk for disease[8–12]. In the future, polygenic risk assessment may

become as common as screening for known mutations of monogenic disease, especially as it

has been shown that individuals with severely high PRS may be at similar risk to disease as are

carriers of rare monogenic mutations[12]. However, since PRS heavily rely on GWAS with large

sample sizes to accurately estimate effect sizes, there is specific demand for the transferability

of PRS from populations with larger GWAS to populations underrepresented by

GWAS[2,3,5,8,17,18,20]. As we would like to investigate the ability of IMPACT annotations to improve

the trans-ethnic application of PRS, we chose pruning and thresholding (P+T) as our model[3,8].

We elected to use P+T rather than LDpred[5,20] or AnnoPred[19], which compute a posterior effect

size estimate for all SNPs genome-wide based on membership to functional categories. With

P+T, we can partition the genome by IMPACT-prioritized and deprioritized SNPs, whereas the

assumptions of the LDpred and AnnoPred models do not support the removal of variants,

making it difficult to directly assess improvement due to IMPACT prioritization. Moreover,

these models have not been explicitly designed or tested for the trans-ethnic application of PRS

and thus are beyond the scope of our work. We conventionally define PRS as the product of

marginal SNP effect size estimates and imputed allelic dosage (ranging from 0 to 2), summed

over M SNPs in the model. Conventional P+T utilizes marginal effect size estimates and

therefore is susceptible to selecting a tagging variant over the causal one guided by GWAS P

values that are inflated by LD. Therefore, we hypothesized that any observed improvement due

to incorporation of IMPACT annotations could result from prioritization of variants with higher

marginal multi-ethnic effect size correlation (**Figure 1D-II**).

Hence, we tested this hypothesis before assessing PRS performance. We selected 21 of

29 summary statistics shared between EUR and EAS with an identified lead IMPACT association

in both populations. Then, using EUR lead IMPACT annotations for each trait (**ST9**), we

partitioned the genome three ways: (1) the SNPs within the top 5% of the IMPACT annotation,

(2) the SNPs within the bottom 95% of the IMPACT annotation, and (3) the set of all SNPs

genome-wide (with no IMPACT prioritization). We then performed stringent LD pruning ($r^2 < 0.1$

from EUR individuals of phase 3 of 1000 Genomes[52]), guided by the EUR GWAS P value, to

acquire sets of independent SNPs in order to compute a EUR-EAS marginal effect size estimate

correlation (**Online Methods**).

For example, in height, EUR-EAS effect size estimates of SNPs in the top 5% partition

(Pearson r = 0.434, **Figure 4A**) are 11.4-fold more similar than those in the bottom 95%

partition (r = 0.038, **Figure 4B**) and 3.31-fold more similar than the set of all SNPs (r = 0.131).

Meta-analyzed across the 21 traits, the marginal multi-ethnic effect size correlation among the

top 5% of IMPACT SNPs was significantly greater than the set of all SNPs genome-wide, across

the 10 most lenient of 17 GWAS locus P value thresholds examined (all difference of means P <

0.026) (**Figure 4C-D**). Furthermore, this observation was consistent across individual traits (**SF9**).

For comparison, we performed the same analysis using alternative annotations: lead

annotations from 513 cell-type-specifically expressed gene sets (SEG)[35] and 220 cell-type-

specific histone mark annotations (CTS)[6] (**SF10**). Marginal effect size correlation with IMPACT

was comparable to CTS when comparing the top 5% of SNPs to the set of all SNPs (difference of

means P value > 0.05 at 14 of 17 P value thresholds, **SF11**). Compared to SEG, IMPACT-selected

SNPs had a significantly greater correlation at 7 of 17 P value thresholds (all difference of means

P value < 0.02, **SF11**). Overall, our results suggest that we might anticipate improved trans-

ethnic portability of PRS models by prioritizing SNPs in key IMPACT annotations.

Finally, we addressed our hypothesis that IMPACT annotations improve the trans-ethnic

portability of PRS (**Figure 1D-III**). For each of the 21 previously analyzed traits, we built a PRS

using effect size estimates from EUR summary statistics and applied it to predict phenotypes of

EAS individuals from BioBank Japan (BBJ) (**Figure 5A**). Here, we compare two PRS models, both

blind to any EAS genetic or functional information and removing SNPs with LD r^2 > 0.2,

Figure 4

Figure 4 legend. Mechanism by which IMPACT prioritization of shared regulatory variants might improve trans-ethnic PRS performance. A) Estimated effect sizes of variants from genome-wide EUR and EAS height summary statistics in the top 5% of the lead IMPACT annotation for EUR height. Proportions of variants in each quadrant indicated in light blue. B) Estimated effect sizes from genome-wide EUR and EAS height summary statistics of variants in the bottom 95% of the same lead IMPACT annotation for height; mutually exclusive with SNPs in A). C) Meta-analysis of multi-ethnic marginal effect size correlations between populations across 21 traits shared between EUR and EAS cohorts over 17 GWAS P value thresholds (with reference to the EUR GWAS). D) Number of SNPs ($\log_{10}$ scale) at each P value threshold for each partition of the genome corresponding to C).

according to European individuals from phase 3 of 1000 Genomes[52]: (i) standard P+T PRS and

(ii) functionally-informed P+T PRS using a subset of SNPs prioritized by the lead EUR IMPACT

annotation (**Online Methods**). In functionally-informed PRS models, for each trait separately,

we *a priori* selected the subset of top-ranked IMPACT SNPs (top 1%, 5%, 10%, or 50%) which

explained the closest to 50% of total trait heritability (**Online Methods**). For all PRS models, we

report results from the most accurate model across nine EUR GWAS *P* value thresholds.

For each trait, we observed that functionally-informed PRS using IMPACT captured more

phenotypic variance than standard PRS (49.9% mean relative increase in R^2, **Figure 5B**, **SF12**,

ST13-15). The mean phenotypic variance explained across traits by functionally-informed PRS

(R^2 = 2.1%, se = 0.2%) was greater than by standard PRS (R^2 = 1.5%, se = 0.1%). For 20 of 21

traits, e.g. excluding basophil count, IMPACT-informed PRS significantly outperformed standard

PRS (difference of means *P* < 0.01). Using 10,000 bootstraps of the PRS sample cohort, we

found that the IMPACT-informed PRS R^2 estimate was consistently greater than the standard

PRS estimate for the same 20 traits (all bootstrap *P* < 0.004, **ST15**). We observed the largest

improvement for RA from R^2 = 1.4% (sd = 0.33%) in the standard PRS versus R^2 = 4.1% (sd =

0.53%, difference of means *P* < 7.7e-10) in the functionally-informed PRS using the B cell TBP

IMPACT annotation. For asthma, R^2 = 0.37% (sd = 0.10%) in the standard PRS versus R^2 = 0.75%

(sd = 0.14%, *P* < 8.5e-4) in the functionally-informed PRS. For MCV, R^2 = 3.0% (sd = 0.10%) in

the standard PRS versus R^2 = 4.1% (sd = 0.12%, *P* < 1.9e-25) in the functionally-informed PRS.

For PrCa, R^2 = 4.5% (sd = 0.36%) in the standard PRS versus R^2 = 6.4% (sd = 0.45%, *P* < 2.4e-6) in

the functionally-informed PRS. For height, R^2 = 4.2% (sd = 0.10%) in the standard PRS versus R^2

= 5.6% (sd = 0.12%, *P* < 1.2e-37) in the functionally-informed PRS.

For our five representative traits asthma, RA, MCV, PrCa, and height, we further

compared functionally-informed PRS_{EUR} using IMPACT to models using cell-type-specifically

expressed genes (SEG) and cell-type-specific histone modification tracks (CTS)[6,35] (**Figure 5C**,

ST16). Across all of the five traits, models using IMPACT explained significantly greater

phenotypic variance (mean R^2= 4.2%, se = 0.3%) than SEG (0.9%, se = 0.1%, all difference of

means P < 9.9e-11). While IMPACT generally outperformed CTS (R^2= 2.6%, se = 0.2%, difference

of means meta P < 1.2e-8), this observation was only individually consistent with 3 of 5 traits

(difference of means P < 9.3e-8). We performed a similar bootstrap analysis as above, yielding

similar results; for only RA and asthma did IMPACT-PRS not produce consistently greater

R^2estimates than CTS-PRS (**ST16**).

Functionally-informed PRS might to some extent compensate for population-specific LD

differences between populations. Hence, we hypothesized that IMPACT-informed PRS would

improve standard PRS moreso in the trans-ethnic prediction framework, in which EUR PRS

models predict EAS phenotypes, than in a within-population framework, in which EAS PRS

models predict EAS phenotypes. Here, we define within-population PRS as PRS_{EAS} and trans-

ethnic PRS as PRS_{EUR} to avoid confusion. In order to directly compare PRS model improvements

between PRS_{EAS} and PRS_{EUR}, we evaluated prediction accuracy on the same individuals. Briefly,

we partitioned the BBJ cohort to reserve 5,000 individuals for PRS testing, derived GWAS

summary statistics from the remaining individuals, and performed P+T PRS modeling and

prediction as done above (**Figure 5D, SF13-15, ST17-18, Online Methods**). For functionally-

informed PRS_{EAS}, we selected lead IMPACT annotations from S-LDSC results using GWAS

summary statistics, as done above, on the partition of the BBJ cohort excluding the 5,000 PRS

test individuals. We defined improvement as the percent increase in R^2 from standard to

functionally-informed PRS; therefore, differences in PRS performance due to intrinsic factors,

such as GWAS power or genotyping platform, cancel out. In both scenarios, we observed

significant non-zero improvements: averaged across the 21 traits in the trans-ethnic setting

(mean percent increase in R^2 = 47.3%, se = 8.1%, P < 2.7e-9) and in the within-population

setting (mean percent increase in R^2 = 20.9%, se = 6.6%, P < 7.5e-4). Indeed, this revealed a

significantly greater improvement in the trans-ethnic than in the within-population application

(difference of means P < 1.7e-4, **Figure 5E**).

Overall, our results reveal that functional prioritization of SNPs using IMPACT

significantly improves both trans-ethnic and within-population PRS models, but is especially

advantageous for the trans-ethnic application of PRS. In conclusion, our results nominate the

prioritization of SNPs according to functional annotations, especially using IMPACT, as a

potential tentative solution for the lack of trans-ethnic portability of PRS models. While

individuals of European ancestry dominate current genetic studies, population-nonspecific cell-

type-specific IMPACT annotations can help transfer highly powered EUR genetic data to study

still underserved populations.

Discussion

In this study, we created a compendium of 707 cell-type-specific regulatory annotations

(**Web Resources**) capturing disproportionately large amounts of polygenic heritability in 95

complex traits and diseases in EUR and EAS populations. We then proposed a three-step

framework to assess how well prioritization of regulatory variants with functional data can

improve multi-ethnic genetic comparisons. First, we showed that heritability-enriched

Figure 5

Figure 5 legend. Identifying shared regulatory variants with IMPACT annotations to improve the trans-ethnic portability of PRS. A) Study design applying EUR summary statistics-based PRS models to all individuals in the BBJ cohort. (B) Phenotypic variance (R^2) of BBJ individuals explained by EUR PRS using two methods: functionally-informed PRS with IMPACT (pink) and standard PRS (blue). Error bars indicate 95% CI calculated via 1,000 bootstraps. C) Phenotypic variance (R^2) of BBJ individuals across 5 exemplary traits explained by EUR IMPACT annotations relative to lead cell-type-specific histone modification annotations (CTS) and lead cell-type-specifically expressed gene sets (SEG). Error bars indicate 95% CI calculated via 1,000 bootstraps. D) Study design to compare trans-ethnic (EUR to EAS) to within-population (EAS to EAS) improvement afforded by functionally-informed PRS models. For each trait, 5,000 randomly selected individuals from BBJ designated as PRS samples. Remaining BBJ individuals used for GWAS to derive EAS summary statistics-based PRS; no shared individuals between GWAS samples and PRS samples. E) Improvement from standard PRS to functionally-informed PRS compared between trans-ethnic (EUR to EAS) and within-population models (EAS to EAS) using the study design in D).

regulatory elements between EUR and EAS populations capture indistinguishable proportions

of heritability across 29 complex traits. Second, we showed that functional prioritization of

variants selects those with more highly correlated marginal effect sizes between populations;

this might explain the improvement driven by functional prioritization in P+T PRS models which

use marginal effect sizes. Third, we showed that variant prioritization with IMPACT annotations

results in consistently improved PRS prediction accuracy, especially for the trans-ethnic

application; potentially due to overcoming large population-specific influences such as LD, an

important challenge of multi-population models.

Designing genetic models for each complex trait or disease that capture risk for the full

diversity of the human population will be challenging. This necessitates approaches that

effectively transfer predictive genetic information from well studied populations to less well

studied populations. Our work provides insight into the potential clinical implementation of PRS

and broader genetic applications that aim to integrate multi-ethnic data. We argue for the use

of biologically diverse IMPACT annotations to capture relevant genetic signal and compensate,

to some extent, for differences in LD across populations. While we did not assess a PRS model

using meta-analyzed summary statistics from two or more populations in this study, we believe

that this approach could be effective, especially for populations with limited GWAS sample size.

We believe that IMPACT may prioritize phenotype-driving regulatory variation. We have

shown IMPACT to be more effective at capturing genetic variation of complex traits than

commonly used functional annotations such as experimentally-derived cell-type-specific

histone marks or gene sets. We hypothesize the utility of IMPACT comes from 1) cell-type-

specificity of TF binding models which locate key classes of regulatory elements and 2) the

integration of thousands of experimentally-derived annotations, which presumably removes

noise and enriches for biological signal present in each individual annotation. Here, we did not

demonstrate the potential utility of IMPACT to perform functional fine-mapping to reduce

credible sets beyond our previous work[31], due to lack of sufficient gold standards with causal

experimental validation and the limitation to genome-wide significant variants. The specific

application of IMPACT in multi-ethnic fine-mapping needs to be further investigated.

We must consider several important limitations of our work. First, our functional

insights are limited to cell types with publicly available TF ChIP-seq data, lacking ones that are

rarer or more difficult to assay. In the future, the cell-type-specific functional training data for

IMPACT may be replaced by newer experimental strategies to map enhancers. For example,

high-throughput CRISPR screens paired with assays for open chromatin could be used to

precisely redefine the regulatory landscape. Second, we used multi-ethnic data to argue for the

utility of our approach. However, the robustness of multi-ethnic comparisons for a given

phenotype rely on properties surrounding the recruitment of individuals or the exact

genotyping platform used in various biobanks, which may result in cohort-bias that inflates

within-population PRS prediction accuracy. For example, BBJ is a disease ascertainment cohort,

in which each individual has any one of 47 common diseases[53,54]; therefore, BBJ control

samples are not comparable to healthy controls of UKBB. Other biases may arise from clinical

differences in phenotyping. Also, we only considered a single non-EUR population in this study,

while the disparity in trans-ethnic portability and hence resulting benefit from functional

annotations may be greater in other non-EUR populations.

In conclusion, we demonstrated that IMPACT annotations improve the comparison of

genetic data between populations and trans-ethnic portability of PRS models using ancestrally

unmatched data. While a long-term goal of the field must be to diversify GWAS and other

genetic studies in non-European populations, it is imperative that genetic models be developed

that work in multiple populations. Such initiatives will necessitate the use of population-

independent functional annotations, such as IMPACT, in order to capture shared biological

mechanisms regulated by complex genetic variation.

Supplemental Data

See *Supplement.pdf* and *Supplementary_Tables.xlsx*

Online Methods

See *Online_Methods.pdf*

Acknowledgements

This work is supported in part by funding from the National Institutes of Health (NHGRI T32

HG002295, UH2AR067677, 1U01HG009088, U01 HG009379, and 1R01AR063759) and the Doris

Duke Charitable Foundation grant #2013097.

Declaration of Interests

The authors declare no competing financial interests.

Web Resources

1. IMPACT 707 annotations: https://github.com/immunogenomics/IMPACT/IMPACT707

2. IMPACT Github repository: https://github.com/immunogenomics/IMPACT

3. HOMER: http://homer.ucsd.edu/homer/motif/

484  4. S-LDSC: https://github.com/bulik/ldsc

485  5. 1000 Genomes: http://www.internationalgenome.org/

486  6. Cell-type-specifically expressed gene set annotations and LD scores:

487   https://data.broadinstitute.org/alkesgroup/LDSCORE/LDSC_SEG_ldscores/

488  7. Cell-type-specific histone modification ChIP-seq datasets:

489   https://data.broadinstitute.org/alkesgroup/LDSCORE/

490  8. Plink: https://www.cog-genomics.org/plink2

492 **References**

494 1. Kichaev, G. & Pasaniuc, B. Leveraging Functional-Annotation Data in Trans-ethnic Fine-

495   Mapping Studies. *Am. J. Hum. Genet.* **97**, 260–271 (2015).

496 2. Lam, M. *et al.* Comparative genetic architectures of schizophrenia in East Asian and

497   European populations. doi:10.1101/445874

498 3. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health

499   disparities. *Nat. Genet.* **51**, 584–591 (2019).

500 4. Sirugo, G., Williams, S. M. & Tishkoff, S. A. The Missing Diversity in Human Genetic Studies.

501   *Cell* **177**, 26–31 (2019).

502 5. Vilhjálmsson, B. J. *et al.* Modeling Linkage Disequilibrium Increases Accuracy of Polygenic

503   Risk Scores. *Am. J. Hum. Genet.* **97**, 576–592 (2015).

504 6. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

505   association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).

506 7. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity

in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

8.   International Schizophrenia Consortium *et al.* Common polygenic variation contributes to

risk of schizophrenia and bipolar disorder. *Nature* **460**, 748–752 (2009).

9.   Chatterjee, N., Shi, J. & García-Closas, M. Developing and evaluating polygenic risk

prediction models for stratified disease prevention. *Nat. Rev. Genet.* **17**, 392–406 (2016).

10. Stahl, E. A. *et al.* Bayesian inference analyses of the polygenic architecture of rheumatoid

arthritis. *Nat. Genet.* **44**, 483–489 (2012).

11. Chatterjee, N. *et al.* Projecting the performance of risk prediction based on polygenic

analyses of genome-wide association studies. *Nat. Genet.* **45**, 400–5, 405e1–3 (2013).

12. Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify individuals

with risk equivalent to monogenic mutations. *Nat. Genet.* **50**, 1219–1224 (2018).

13. Schumacher, F. R. *et al.* Association analyses of more than 140,000 men identify 63 new

prostate cancer susceptibility loci. *Nat. Genet.* **50**, 928–936 (2018).

14. Sharp, S. A. *et al.* Development and Standardization of an Improved Type 1 Diabetes

Genetic Risk Score for Use in Newborn Screening and Incident Diagnosis. *Diabetes Care* **42**,

200–207 (2019).

15. Kullo, I. J. *et al.* Incorporating a Genetic Risk Score Into Coronary Heart Disease Risk

Estimates: Effect on Low-Density Lipoprotein Cholesterol Levels (the MI-GENES Clinical

Trial). *Circulation* **133**, 1181–1188 (2016).

16. Natarajan, P. *et al.* Polygenic Risk Score Identifies Subgroup With Higher Burden of

Atherosclerosis and Greater Relative Benefit From Statin Therapy in the Primary

Prevention Setting. *Circulation* **135**, 2091–2101 (2017).

17. Márquez-Luna, C., Loh, P.-R., South Asian Type 2 Diabetes (SAT2D) Consortium, SIGMA

Type 2 Diabetes Consortium & Price, A. L. Multiethnic polygenic risk scores improve risk

prediction in diverse populations. *Genet. Epidemiol.* **41**, 811–823 (2017).

18. Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in diverse human

populations. *Nat. Commun.* **10**, 3328 (2019).

19. Hu, Y. *et al.* Leveraging functional annotations in genetic risk prediction for human

complex diseases. *PLoS Comput. Biol.* **13**, e1005589 (2017).

20. Márquez-Luna, C. *et al.* Modeling functional enrichment improves polygenic prediction

accuracy in UK Biobank and 23andMe data sets. *bioRxiv* 375337 (2018).

doi:10.1101/375337

21. Okada, Y. *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery.

Nature **506**, 376–381 (2014).

22. Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population links cell

types to complex human diseases. *Nature Genetics* **50**, 390–400 (2018).

23. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body

mass index in ~700000 individuals of European ancestry. *Hum. Mol. Genet.* **27**, 3641–3649

(2018).

24. Schaub, M. A., Boyle, A. P., Kundaje, A., Batzoglou, S. & Snyder, M. Linking disease

associations with regulatory information in the human genome. *Genome Res.* **22**, 1748–

1759 (2012).

25. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in

regulatory DNA. *Science* **337**, 1190–1195 (2012).

26. Reshef, Y. A. *et al.* Detecting genome-wide directional effects of transcription factor

binding on polygenic disease risk. *Nat. Genet.* **50**, 1483–1493 (2018).

27. Liu, X., Li, Y. I. & Pritchard, J. K. Trans Effects on Gene Expression Can Drive Omnigenic

Inheritance. *Cell* **177**, 1022–1034.e6 (2019).

28. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

29. Teytelman, L., Thurtle, D. M., Rine, J. & van Oudenaarden, A. Highly expressed loci are

vulnerable to misleading ChIP localization of multiple unrelated proteins. *Proc. Natl. Acad.*

Sci. U. S. A. **110**, 18602–18607 (2013).

30. Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution

mapping of DNA binding sites. *Elife* **6**, (2017).

31. Amariuta, T. *et al.* IMPACT: Genomic Annotation of Cell-State-Specific Regulatory Elements

Inferred from the Epigenome of Bound Transcription Factors. *Am. J. Hum. Genet.* **104**,

879–895 (2019).

32. Kawakami, E., Nakaoka, S., Ohta, T. & Kitano, H. Weighted enrichment method for

prediction of transcription regulators from transcriptome and global chromatin

immunoprecipitation data. *Nucleic Acids Res.* **44**, 5010–5021 (2016).

33. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human

genome. *Nature* **489**, 57–74 (2012).

34. Roadmap Epigenomics, C. *et al.* Heravi-428 Moussavi A, Kheradpour P, Zhang Z, Wang J, et

al. Integrative analysis of 111 reference human 429 epigenomes. *Nature* **518**, 317–330

(2015).

35. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies

disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).

36. Gazal, S. *et al.* Linkage disequilibrium–dependent architecture of human complex traits

shows action of negative selection. *Nat. Genet.* **49**, 1421–1427 (2017).

37.   Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association

studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–D1012

(2019).

38.   Akiyama, M. *et al.* Characterizing rare and low-frequency height-associated variants in the

Japanese population. *Nat. Commun.* **10**, 4393 (2019).

39.   Ishigaki, K., Akiyama, M., Kanai, M. & Takahashi, A. Large scale genome-wide association

study in a Japanese population identified 45 novel susceptibility loci for 22 diseases.

bioRxiv (2019).

40.   Akiyama, M. *et al.* Genome-wide association study identifies 112 new loci for body mass

index in the Japanese population. *Nat. Genet.* **49**, 1458–1467 (2017).

41.   Brown, B. C., Ye, C. J., Price, A. L. & Zaitlen, N. Transethnic Genetic-Correlation Estimates

from Summary Statistics. *Am. J. Hum. Genet.* **99**, 76–88 (2016).

42.   Drake, L. Y. *et al.* B cells play key roles in th2-type airway immune responses in mice

exposed to natural airborne allergens. *PLoS One* **10**, e0121660 (2015).

43.   Amariuta, T., Luo, Y., Knevel, R., Okada, Y. & Raychaudhuri, S. Advances in genetics toward

identifying pathogenic cell states of rheumatoid arthritis. *Immunol. Rev.* (2019).

doi:10.1111/imr.12827

44.   Buttari, B., Profumo, E. & Riganò, R. Crosstalk between red blood cells and the immune

system and its impact on atherosclerosis. *Biomed Res. Int.* **2015**, 616834 (2015).

45.   Anderson, H. L., Brodsky, I. E. & Mangalmurti, N. S. The Evolving Erythrocyte: Red Blood

Cells as Modulators of Innate Immunity. *J. Immunol.* **201**, 1343–1351 (2018).

46.   Lui, J. C. & Baron, J. Mechanisms limiting body growth in mammals. *Endocr. Rev.* **32**, 422–

440 (2011).

47. Maier, A. B., van Heemst, D. & Westendorp, R. G. J. Relation between body height and

replicative capacity of human fibroblasts in nonagenarians. *J. Gerontol. A Biol. Sci. Med.*

Sci. **63**, 43–45 (2008).

48. Murphy, R. A. *et al.* Adipose tissue, muscle, and function: potential mediators of

associations between body weight and mortality in older adults with type 2 diabetes.

Diabetes Care **37**, 3213–3219 (2014).

49. Heymsfield, S. B., Gallagher, D., Mayer, L., Beetsch, J. & Pietrobelli, A. Scaling of human

body composition to stature: new insights into body mass index. *Am. J. Clin. Nutr.* **86**, 82–

91 (2007).

50. Kichaev, G. *et al.* Leveraging Polygenic Functional Enrichment to Improve GWAS Power.

Am. J. Hum. Genet. **104**, 65–75 (2019).

51. Gusev, A. *et al.* Atlas of prostate cancer heritability in European and African-American men

pinpoints tissue-specific regulation. *Nat. Commun.* **7**, 10979 (2016).

52. Gibbs, R. A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74

(2015).

53. Nagai, A. *et al.* Overview of the BioBank Japan Project: Study design and profile. *J.*

Epidemiol. **27**, S2–S8 (2017).

54. Hirata, M. *et al.* Cross-sectional analysis of BioBank Japan clinical data: A large cohort of

200,000 patients with 47 common diseases. *J. Epidemiol.* **27**, S9–S21 (2017).

Online Methods

Data

TF ChIP-seq data. We previously collected 3,181 publicly available transcription factor (TF)

chromatin immunoprecipitation (ChIP) datasets derived from human primary cells or cell lines.

We downloaded raw sequencing data in SRA format from NCBI GEO, then converted the data to

FASTQ format using the SRA Toolkit function fastq-dump, used FastQC for quality assessment of

sequencing reads, and finally mapped reads to the human genome (hg19/GRCh37) with

Bowtie2 [v2.2.5] using default parameters. All ChIP-seq datasets were matched to

corresponding control data from which peaks were called with macs [v2.1] with q value < 0.01

under a bimodal model, producing 3,181 bed file-formatted files[1,2]. The 1,542 datasets selected

for use with our IMPACT model framework (see below) are listed with accession codes in **ST1**.

Genome-wide annotation data. We augmented our set of 515 publicly available epigenomic

and sequence feature annotations from our previous study[3] with 116 personally curated

datasets from NCBI, 2,593 ENCODE histone ChIP-seq datasets and 2,121 ENCODE open

chromatin DNase-seq datasets[4], all publicly available at the accessions provided in **ST2**. All files

were collected in 6-column standard bed file format. This augmentation brought the total

number of features to 5,345.

Genome-wide association data. We collected publicly available summary statistics data for 111

genome-wide association studies (GWAS) across separate cohorts of East Asian and European

individuals[5–7]. East Asian GWAS data were collected from Biobank Japan (BBJ) while European

GWAS data were collected from either UKBioBank (UKBB) or the GWAS catalog, referred to as

PASS (publicly available summary statistics) (**ST3**). All GWAS summary statistics were

reformatted to be compatible with S-LDSC (see below) and thus contained the following

information for each SNP (per row): rsID, A1 (reference allele), A2 (alternative allele), GWAS

sample size (effective sample size per SNP, may vary with genotyping), chi-square statistic, *z*-

score. For multi-ethnic genetic correlation and polygenic risk score prediction, all GWAS

summary statistics were reformatted to contain the SNP ID (chr_position_A1_A2),

chromosome, base pair, A1, A2, effect size estimate, effect size estimate standard error, and *P*-

value.

Cell-type-specifically expressed gene set (SEG) and cell-type-specific histone modification

(CTS) annotations. We downloaded 513 publicly available SEG annotations for European SNPs

from phase 3 of 1000 Genomes accompanied by pre-computed LD scores (see **Web**

Resources)[8]. SEG annotations are binary and thus each SNP is designated a 1 or a 0, indicating

that the SNP does or does not lie, respectively, within 100 kb of the gene body of the

corresponding gene set[8]. We downloaded 220 publicly available CTS annotations of peak data

in bed file format, from which we annotated European SNPs from phase 3 of 1000 Genomes[9]

and used S-LDSC to compute LD scores (see **Web Resources**)[7]. These annotations are also

binary, in which case each SNP is designated a 1 or a 0, indicating that the SNP does or does not

like, respectively, within the peak of histone modification.

BioBank Japan data. For PRS analysis, we utilized phenotype and genotype data of the BioBank

Japan Project (BBJ)[10,11]. All of the calculations related to PRS were conducted on the RIKEN

computing server. BBJ is a biobank that collaboratively collects DNA and serum samples from

12 medical institutions in Japan. This project recruited approximately 200,000 patients with the

diagnosis of at least one of 47 diseases. Informed consent was obtained from all participants by

following the protocols approved by their institutional ethical committees. We obtained

approval from the ethics committees of the RIKEN Center for Integrative Medical Sciences and

the Institute of Medical Sciences at the University of Tokyo.

Statistical Methods

IMPACT Model. We implemented our previously defined model to predict TF binding on a motif

site. This model regresses the likelihood (*p*) of a binding event on the epigenomic profile of the

motif site, in a logistic regression framework over *j* epigenomic features as follows:

$$log\left(\frac{p}{1-p}\right) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \ldots + \beta_j X_j.$$

60   We use a weighted average of ridge and lasso regularization terms in the objective function to

61   restrict the magnitude of fit coefficients and enforce sparsity to reduce overfitting, respectively,

62   as follows:

$$argmin_\beta = (||Y - X\beta||^2 + \frac{1}{2}(1 - \alpha)||\beta||^2 + \alpha\,||\beta||).$$

Training IMPACT. We trained an IMPACT model for each unique cell type-TF pair present in our

data collection. Our collection consists of 3,181 TF ChIP-seq profiles, representing 442 TFs, 296

cell types, and 24 tissues. The IMPACT model requires that the assayed TF has a distinct binding

motif and so we removed all ChIP-seq datasets corresponding to a TF that did not have a known

sequence motif in MEME, Jaspar, or Transfac databases. This resulted in 1,542 TF ChIP-seq

profiles across 142 TFs, 245 cell types, 23 tissues, and 728 unique combinations of TFs and cell

types. As we did in our previous study[3], we merged experiments of the same TF-cell type

combination by taking the union of the peaks. We next identified motif sites bound by a TF by

using HOMER [v4.8.3][12] to scan ChIP-seq peaks for motif matches exceeding the empirically

determined motif detection threshold. Similarly, we identified motif sites not bound by a TF by

using HOMER to scan the entire genome for sequence matches. 21 of these models did not

contain sufficient overlap between TF sequence motifs and ChIP-seq peaks which would lead to

underfitting in the logistic regression (fewer than 7), thereby resulting in 707 total possible

IMPACT annotations. We then trained 707 IMPACT models using up to 1,000 TF-bound

sequence motifs (evidenced by ChIP-seq) and 10,000 unbound sequence motifs. For each of

707 TF-cell type pairs, we learned a predictive model of TF binding and annotated SNPs

genome-wide for both EUR and EAS populations, with a mean regulatory probability per

nucleotide of 0.02 (se = 7.5e-4).

Assessing cell type specificity of IMPACT tracks. We acquired lists of specifically expressed

genes in 9 different cell types: T cells, B cells, fibroblasts, monocytes, brain, liver, colon,

prostate, and breast according to differential gene expression *t*-statistics from previous work[8],

specifically labeled as T.4+8int.Th, B.Fo.LN, Cells_Transformed_fibroblasts, Mo.6C+II-.LN,

Brain_Cortex, Liver, Colon_Transverse, Prostate, Breast_Mammary_Tissue, respectively from

either ImmGen or GTEx databases. Large and positive *t*-statistics represent greater specificity of

gene expression in the target cell type, large but negative *t*-statistics represent specifically

repressed genes, and *t*-statistics near 0 represent nonspecific gene expression, representing

commonly expressed genes. For each cell type, we selected the 100 genes with highest *t*-

statistics, e.g. specifically expressed (SE) genes, and 100 genes such that -0.5 < *t*-statistic < 0.5,

e.g. not specifically expressed genes (NS). For each cell type separately, we collected all related

IMPACT annotations from the compendium of 707 total annotations. Then for each annotation

separately, we computed the average IMPACT score over all EUR SNPs from phase 3 of 1000

Genomes within 2kb of each SE or NS gene body. Finally, we computed the average across all

100 SE and 100 NS genes, separately.

Partitioning heritability with S-LDSC. We applied S-LDSC [v1.0.0][7] to partition the common

(MAF > 5%) SNP heritability of 111 polygenic traits and diseases, with significantly non-zero

heritability estimates ($P < 0.05$). We partitioned heritability using a customized version of the

baselineLD model, in which we excluded cell-type-specific regulatory annotations (as we would

be testing the enrichment of such annotations from IMPACT). In total, we used 69 cell-type-

nonspecific baselineLD annotations and added one or more IMPACT annotations to the model

to test for cell-type-specific enrichment. We use three metrics to evaluate how well our

IMPACT annotations capture polygenic heritability: enrichment[7], the proportion of heritability

explained by the top 5% of SNPs[7], and per-annotation standardized effect size, $\tau*$[6]. Briefly,

enrichment is defined as the proportion of common SNP heritability divided by the genome-

wide proportion of SNPs in the annotation, for continuous annotations this is the average

annotation value across SNPs. τ^* represents the average per-SNP heritability of a category of

SNPs, where a single SNP may claim membership to one or more categories. τ^* is defined as the

proportionate change in per-SNP heritability associated with a one standard deviation increase

in the value of the annotation. The sum of the τ^* over categories of SNPs equals the total

estimated heritability of the trait. τ^* has units of heritability and is comparable between traits,

annotations, and populations, because it is normalized for the total heritability (indicative of

the power of the GWAS), the dispersion of the annotation values (annotation size), and the

number of common SNPs (population-specific) considered in the model, respectively. τ, the

precursor of τ^*, is the coefficient estimated in the S-LDSC regression. τ and τ^* are conditionally

dependent on the provided baselineLD annotations. Therefore, the τ^* estimate for an IMPACT

annotation is considered a measure of cell-type-specific or annotation-specific SNP heritability,

as the remaining annotations in the model (baselineLD) are not cell-type-specific. Significance

of τ^* is computed using a *z*-test of how different the τ^* estimate is from 0; the significance of

strictly positive τ^* estimates are reported in our study.

Measuring heritability in top X% of SNPs of a continuous annotation. To partition the

heritability captured by various top echelons of SNPs of a given continuous annotation, we used

the same strategy as in a previous study[6]. By this strategy, the proportion of heritability

explained by a set of SNPs is the sum over all SNPs of the product of the τ^* of each category in

the S-LDSC model, e.g. baselineLD plus IMPACT annotation, and the SNP membership to that

category (1 or 0 in the case of binary annotations, continuous values in the case of continuous

annotations) divided by the same metric for all SNPs genome-wide.

Conditional S-LDSC analysis to identify independent annotation-trait associations. Due to the

redundancy in modeled cell type programs and inherent covariance of IMPACT annotations

(**SF2**), the $\tau*$ associations we find with S-LDSC cannot be independent. To this end, for each of

95 traits across EUR and EAS for which we identified a lead IMPACT annotation, reported in

ST9, we performed a series of conditional analyses using S-LDSC. For each trait with more than

one significant $\tau*$ association, we created S-LDSC models consisting of the 69 baselineLD

annotations, the lead annotation for that trait, and separately, each remaining significant

IMPACT annotation. We kept annotations that retained their $\tau*$ significance when conditioned

on the lead annotation(s), which we also required to retain significance. We iteratively

performed these conditional analyses until we were no longer able to identify independent $\tau*$

associations.

Deming regression of EUR $\tau*$ on EAS $\tau*$. As there is significant correlation among IMPACT

annotations, due to redundancy in cell type regulatory elements, we used an iterative pruning

approach, similar to LD-pruning, to identify independent IMPACT annotations. For each trait,

we ranked all 707 IMPACT annotations by their $\tau*$ significance values. Then, we selected the

lead annotation, removed all annotations correlated with Pearson $r > 0.5$, and selected the next

lead annotation, and so on. This approach produced a set of relatively independent

annotations, for which the assumptions of Deming, or any, regression would not be violated.

For each trait, we ran Deming regression over approximately 100 independent IMPACT

annotations using the R function *deming* within the package *deming*. Across independent

observations for all traits, we tested the null hypothesis that the slope of the Deming

regression, which considers standard errors on both the predictor (EUR τ*) and response

variables (EAS τ*), is equal to 1.

Multi-ethnic and within-population genetic correlation. We computed the genetic correlation

(R_g) between pairs of 29 traits for which we acquired EUR and EAS GWAS using Popcorn

[v.0.9.6][13] with default parameters, including maximum likelihood estimation as opposed to

regression[14]. First, we computed cross-population scores between the two populations using

the *compute* flag with the *popcorn* executable, indicating approximately the correlation

between LD at each SNP using EUR and EAS reference LD panels from phase 3 of 1000

Genomes. Then, we used the *fit* flag with the *popcorn* executable to compute the multi-ethnic

genetic correlation of these 29 traits. R_g estimates computed after restricting to MAF > 5% did

not significantly differ from no MAF restriction. Popcorn computes R_g using either "genetic

impact" (effect sizes normalized by allele frequency) or "genetic effect" (unmodified effect

sizes). We observed no significant heterogeneity between the R_g computed using "genetic

impact" and "effect", although "genetic effect" estimates were consistently but not significantly

larger.

We then computed cross-trait cross-population genetic correlations across 21 traits for

which we observed at least one significant IMPACT annotation association in both EUR and EAS.

Therefore, in total we computed the genetic correlation among 42 traits (21 phenotypes x 2

populations). For pairs of traits with one from EUR and one from EAS, we used Popcorn as

described above with MAF threshold of 5% and "genetic impact". For pairs of traits from the

same population we used LDSC [v.1.0.0]. First we used the *munge_sumstats.py* script to make

the direction of allelic effect consistent in the GWAS summary statistics while also restricting to

well-imputed Hapmap3 SNPs. Then, we used the *ldsc.py* script with the *-rg* flag to compute the

genetic correlation using EUR and EAS reference LD panels from phase 3 of 1000 Genomes

where appropriate.

Multi-ethnic marginal effect size correlation. We acquired GWAS summary statistics for each

of 21 shared traits between EUR and EAS for which there was at least one significant IMPACT

association in each population. Then, we restricted to SNPs shared between EUR and EAS

GWAS summary statistics. Next, we performed stringent iterative LD clumping with PLINK

[v1.90b3][15] using EUR summary statistics (selecting the most significant SNP, then removing all

SNPs in LD with $r^2 > 0.1$ within 1 Mb, then selecting the next most significant SNP, and so on).

This step satisfies the assumption of independence in the Pearson correlation that we will

compute among marginal effect sizes. We selected our initial set of SNPs under three scenarios:

(1) using no functional inference, (2) using the top 5% of SNPs according to the trait's lead EUR

IMPACT annotation, and (3) using the bottom 95% of SNPs according to the trait's lead EUR

IMPACT annotation (mutually exclusive with scenario 2). With our set of independent SNPs for

each trait and under each of three scenarios, we compute a Pearson correlation between the

estimated effect sizes, while further stratifying loci on 17 EUR *P*-values (1, 0.3, 0.1, 0.03, 0.01,

3e-3, 1e-3, 3e-4, 1e-4, 3e-5, 1e-5, 3e-6, 1e-6, 3e-7, 1e-7, 3e-8, 1e-8). For example, stratum with

P = 0.1 includes all SNPs with EUR GWAS P < 0.1.

Polygenic risk score calculation. In this study, we utilized pruning and thresholding (P+T) for the

calculation of PRS. We constructed PRS models from either EUR summary statistics or EAS

summary statistics and evaluated their predictive performance on individual EAS phenotypes.

Here, we define within-population PRS as PRS$_{EAS}$ and trans-ethnic PRS as PRS$_{EUR}$ to avoid

confusion. For PRS$_{EUR}$, we utilized genome-wide summary statistics from EUR as reported in

their publicly available version. For PRS$_{EAS}$, we held out 5,000 individuals for PRS analysis and

conducted GWAS using the remaining individuals to avoid overfitting (see next section). For

each trait separately, we restricted our analysis to variants that exist in both GWAS summary

statistics and post-imputation genotype data of EAS individuals used for PRS analysis

(imputation quality of r^2 > 0.3 in minimac3). A detailed description related to the genotyping

platform and imputation strategy is provided in a previous report[2]. We excluded the MHC

region in this analysis.

We designed PRS models using two strategies: standard PRS and functionally-informed

PRS. For standard PRS$_{EUR}$, we performed conventional LD clumping to acquire sets of

independent SNPs using EUR LD reference panels from phase3 of 1000 Genomes. Similarly for

PRS$_{EAS}$, we utilized EAS LD reference panels from phase3 of 1000 Genomes. We used PLINK

[v1.90b3][15] to remove variants in LD with r^2 > 0.2 with a significance threshold for index SNPs

of P = 0.5. For functionally-informed PRS, we restricted the analysis to variants with high

IMPACT score according to the lead IMPACT annotation before conducting LD clumping. As

before, we define the lead annotation as the one with the largest $\tau*$ estimate that was

significantly greater than 0. When we designed PRS_EUR, we utilized the lead IMPACT annotation

in EUR GWAS summary statistics (EAS summary statistics were not taken into account to avoid

overfitting). Similarly, when we design PRS_EAS, we utilized the lead IMPACT annotation in EAS

GWAS summary statistics for which 5,000 EAS individuals for PRS analysis were removed to

avoid overfitting. We performed LD clumping using variants within a predefined top percentage

of IMPACT scores. This was determined by the percentage that captured the closest to 50% of

total trait heritability; considered percentages included the top 1%, 5%, 10%, and 50%.

We evaluated PRS performance using EAS individuals. First, we used all individuals in the

BBJ cohort for PRS_EUR testing. Second, we compared the improvement afforded by IMPACT in

PRS_EUR relative to PRS_EAS models using 5,000 randomly selected individuals in BBJ; specifically

for case-control GWAS, we randomly selected 1,000 cases and 4,000 controls.

For all models, we built a PRS for each individual *j* in our test set (in all cases, there is no

overlap between GWAS samples and PRS samples) using variant effect size estimates from

GWAS as follows:

$$PRS_j = \sum_i^M A_i \ * \ \beta_i, \qquad \text{(Equation 1)}$$

Where M is the total number of SNPs shared between GWAS summary statistics and post-

imputation genotype data of EAS individuals, *i* is the i^{th} SNP in the model, A_i is the allele dosage

of the trait-increasing allele *i*, and β_i is the estimated effect size of allele *i* from the GWAS. We

calculated PRS using PLINK2.

For QC of quantitative phenotypes, we excluded (1) related samples (PI_HAT > 0.187

estimated by PLINK), (2) samples with age < 18 and age > 85, and (3) samples with measured

values outside three interquartile ranges (IQR) of the upper or lower quartiles. The effect of sex,

age, age^2, the top 10 PCs, and affection status of 47 diseases were removed by linear

regression, and the residuals were further normalized by the rank-based inverse normal

transformation (see Equation 3 below). For QC of case/control phenotypes, we excluded (1)

related samples (PI_HAT > 0.187 estimated by PLINK) and (2) samples with age < 18 and age >

85.

We then regressed our phenotype of interest (Y), a measured quantitative trait or a

diagnosed disease among the PRS samples, on the per-individual PRS as follows:

For diseases,

$Y_j \sim PRS_j + sex + age + Geno\ PC1 + \ldots + Geno\ PC10.$ (Equation 2)

For quantitative traits,

$Normalized\ Y_j \sim PRS_j.$ (Equation 3)

We then report the variance explained; for quantitative traits, this is the variance

explained by a linear model and for diseases, the variance explained is from a logistic model

(Nagelkerke R^2)[14,16,17] which we convert to liability scale pseudo R^2 such that R^2 values are

comparable among both quantitative and case/control phenotypes. We used various GWAS *P*

value thresholds (0.1, 0.03, 0.01, 0.003, 0.001, 3e-4, 1e-4, 3e-5, 1e-5) to assess the predictive

performance of our PRS. For each model, we reported in the text the largest R^2 achieved across

the nine P value thresholds.

To estimate confidence intervals of PRS performance (R^2, as explained above), we

conducted 1,000 bootstraps using the R package *boot*. We also conducted 10,000 bootstraps to

evaluate whether the R^2 difference between two PRS models (functionally-informed - standard)

is significantly greater than 0; we calculated the R^2 difference between two PRS models in each

round of bootstrapping (delta R^2), and assess its distribution in 10,000 bootstraps. If we let N

be the frequency of delta $R^2 < 0$, we define one-sided *P* values for delta $R^2 > 0$ as (N +

1)/10,000.

Genome-wide association studies in BBJ. As described in the previous section, we held out

5,000 randomly selected individuals for the PRS analysis and performed GWAS on the

remaining individuals (sample sizes are provided in **ST13-14**). GWAS was conducted with PLINK2

using the same imputed dosages as used in the PRS analysis. For quantitative traits, normalized

residuals were analyzed by a linear regression model. For diseases, affection status was

analyzed by a logistic regression model using age, sex, and the top 10 genotype PCs as

covariates.

Online Methods References

1.   Kawakami, E., Nakaoka, S., Ohta, T. & Kitano, H. Weighted enrichment method for

prediction of transcription regulators from transcriptome and global chromatin

immunoprecipitation data. *Nucleic Acids Res.* **44**, 5010–5021 (2016).

2.   Ishigaki, K., Akiyama, M., Kanai, M. & Takahashi, A. Large scale genome-wide association

study in a Japanese population identified 45 novel susceptibility loci for 22 diseases.

bioRxiv (2019).

3.   Amariuta, T. *et al.* IMPACT: Genomic Annotation of Cell-State-Specific Regulatory Elements

Inferred from the Epigenome of Bound Transcription Factors. *Am. J. Hum. Genet.* **104**,

879–895 (2019).

4.   ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human

genome. *Nature* **489**, 57–74 (2012).

5.   Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population links cell

types to complex human diseases. *Nat. Genet.* **50**, 390–400 (2018).

6.   Gazal, S. *et al.* Linkage disequilibrium–dependent architecture of human complex traits

shows action of negative selection. *Nat. Genet.* **49**, 1421–1427 (2017).

7.   Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).

8.   Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies

disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).

9.   Gibbs, R. A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74

(2015).

10.   Nagai, A. *et al.* Overview of the BioBank Japan Project: Study design and profile. *J.*

Epidemiol. **27**, S2–S8 (2017).

11.   Hirata, M. *et al.* Cross-sectional analysis of BioBank Japan clinical data: A large cohort of

200,000 patients with 47 common diseases. *J. Epidemiol.* **27**, S9–S21 (2017).

12. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589

(2010).

13. Brown, B. C., Ye, C. J., Price, A. L., Zaitlen, N. & Asian Genetic Epidemiology Network-Type 2

Diabetes. Transethnic genetic correlation estimates from summary statistics.

doi:10.1101/036657

14. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health

disparities. *Nat. Genet.* **51**, 584–591 (2019).

15. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based

linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

16. Lam, M. *et al.* Comparative genetic architectures of schizophrenia in East Asian and

European populations. doi:10.1101/445874

17. Lee, S. H., Goddard, M. E., Wray, N. R. & Visscher, P. M. A better coefficient of

determination for genetic profile analysis. *Genet. Epidemiol.* **36**, 214–224 (2012).