Medium spiny neurons spontaneous activity reveal the discrete segregation of mouse dorsal striatum

J. Alegre-Cortés, M. Sáez, R. Montanari, R. Reig*

Affiliation

Instituto de Neurociencias CSIC-UMH, San Juan de Alicante, Spain

J. A-C and M. S. contributed equally.

*Corresponding Author: Ramón Reig ramon.reig@umh.es

Abstract

Behavioural studies differentiate the rodent dorsal striatum (DS) into lateral and medial regions based on their particular roles; however, anatomical evidence suggest that it is a unified structure. To understand striatal dynamics and basal ganglia functions, it is essential to clarify the circuitry that supports this behavioural-based segregation. Here, we show that the mouse DS is made of two non-overlapping functional circuits divided by a sharp boundary. Combining *in vivo* optopatch-clamp and extracellular recordings of spontaneous activity, we demonstrate different coupling of lateral and medial striatum to the cortex together with an independent integration of the slow and fast oscillations, due to particular cortico-striatal connectivity and local properties of each region. Additionally, distinct properties between striatonigral and striatopallidal neurons were revealed along DS. These results demonstrate that the rodent DS is segregated in two neuronal circuits, in homology with the caudate and putamen nuclei of primates.

Introduction

The basal ganglia are a group of subcortical nuclei involved in a diversity of functions including motor control, learning, decision making and reward (Packard and Knowlton, 2002; Schultz et al., 1997; Yin and Knowlton, 2006). The striatum is the main input structure of the basal ganglia, receiving glutamatergic transmission from the cortex and the thalamus (Alloway et al., 2009; Kincaid et al., 1998; Wilson, 1987). In primates and other mammals, the dorsal part of the striatum is formed by the nuclei caudate and putamen, which are anatomically separated by the internal capsule. In contrast, rodent dorsal striatum (DS) is considered a homogeneous structure due to the absence of anatomical border and the copious presence of overlapping axonal connections (Alloway et al., 2006; Hooks et al., 2018; Hoover et al., 2003). However, some recent studies has divided the DS in two regions, dorsomedial (DMS) and dorsolateral (DLS) striatum, based on their different behavioural roles (Graybiel, 2008; Hauber and Schmidt, 1994).While both regions participate in motor control, DLS is often related with habit, stimulus-response associations and navigation, whereas the DMS mediates action-outcome associations, goal-directed strategy, flexible shifting between behavioural strategies, suggesting a role in higher cognitive behaviours (Faure et al., 2005; Hilário and Costa, 2008; Lerner et al., 2015; Thorn et al., 2010). In order to understand the basal ganglia circuits and their related behaviours, an essential question should be addressed: Is the functional segregation of the DS supported by two different circuits?

Cortex and thalamus project to the striatum, forming organized glutamatergic synapses along its mediolateral axis, and defining multiple striatal subregions (Hintiryan et al., 2016; Hunnicutt et al., 2016). This corticostriatal axonal innervation presents a high degree of convergence and divergence (Flaherty and Graybiel, 1991) and originates in both hemispheres from different subtypes of pyramidal neurons (Cowan and Wilson, 1994; Hooks et al., 2018; Levesque et al., 1996; Reiner et al., 2003; Wilson, 1987). Corticostriatal connections innervate both striatonigral and striatopallidal medium-sized spiny neurons (direct and indirect MSNs respectively) (Doig et al., 2010; Wall et al., 2013), which represent 95% of the neurons in the striatum (Kita and Kitai, 1988), and different types of interneurons (Tepper et al., 2008). The lateral region of the striatum is highly innervated by axons from somatosensory and motor related cortical areas, while medial one receives cortical axons from visual, auditory, associative, limbic (Hintiryan et al., 2016; Hunnicutt et al., 2016) and with lower density from somatosensory cortical regions (Reig and Silberberg, 2014).

In addition to the differences on cortical or thalamic axonal innervation, there are dissimilarities on the composition of the striatal microcircuits. The distribution of parvalbumin (PV) and cholinergic interneurons (ChI) along the mediolateral axis of the DS is not homogeneous (Gerfen et al., 1985; Kita et al., 1990; Matamales et al., 2016; Muñoz-Manchado et al., 2018). Furthermore, dopaminergic projections from the lateral part of the substantia nigra massively innervate the DS (Ikemoto, 2007), with different impact in the activity of ChI in the along DS (Chuhma et al., 2018). All these precise afferent connectivity, microcircuit interactions and neuromodulation regulate the synaptic activity of DLS- and DMS-MSNs.

The activity of every neural circuit is limited by anatomical and functional constrains which will restrict the repertoire of spontaneous and evoked activity patterns, defining the functional connectivity of the brain (Getting, 1989; Luczak et al., 2015). In this work, we describe how MSNs of the lateral and medial regions of the DS integrates the spontaneous brain activity.

As a model to study functional connectivity we used the Slow Wave Oscillation (SWO), which is consider the default activity pattern of the cortex (Sanchez-Vives et al., 2017) and has been previously used as a physiological marker for several neurological disorders (Amiri et al., 2019; Busche et al., 2015; Ruiz-Mejias et al., 2016). This brain state is characterized by periods of high synchronized spontaneous activity (Up states) intermingled with periods of silence (Down states) at ~1Hz (Steriade et al., 1993b). The SWO is originated in the cortex (Sanchez-Vives and McCormick, 2000; Timofeev et al., 2000) and propagates directly to the striatum, modulating the resting state of MSNs (Sáez et al., 2018; Wilson and Kawaguchi, 1996) and interneurons (Reig and Silberberg, 2014).

Because MSNs recorded *in vivo* are known to fire scarcely (Adler et al., 2012; Berke et al., 2004; Wilson, 1993) we performed single and pairs of whole-cell patch-clamp recordings to analyse their subthreshold dynamics, identifying their specific pathways. Our findings show how DS is divided in two non-overlapping circuits, based on the MSNs activity. DLS- and DMS-MSNs differ in the integration of the slow wave and beta oscillations, as well as in the functional coupling to multiple cortical regions. In addition, by means of double *in vivo* patch-clamp recordings, we demonstrate a sequential propagation of the cortical SWO along DS. Finally, we found that the direct and indirect pathways MSNs have particular attributes in the DLS and DMS, displaying differences in their electrophysiological properties and integration of the SWO.

In conclusion, our results demonstrate that DS is divided in two functional circuits, separated by a sharp boundary, each of them with specific properties that are essential to understand the striatum and basal ganglia functions.

Results

MSNs in the DLS and DMS have different electrophysiological properties.
We obtained *in vivo* whole-cell patch-clamp recordings from 223 neurons, located in the dorsal striatum (n=197) and several cortical areas (n=26). All of them displayed SWO with prominent Up and Down states (Fig. 1C, 2A, 4A-B, 4D, 5A-B, 6A, S2), at close frequency of ~0.7 Hz in both brain regions. The particular types of striatal neurons were identified by their electrophysiological properties and morphology (Fig. 1B, see methods). Direct and indirect pathway MSNs were determined by their responses to the light stimulation using the optopatcher (Katz et al., 2019) (Ketzef et al., 2017) (Fig. 7A). All average graphs showed in this study represent the standard deviation unless stated otherwise.
In order to study the circuit attributes that may characterize the dorsal striatum (DS), our initial approach was to compare the electrophysiological properties of striatal MSNs recorded in the lateral (DLS, n=77) and medial (DMS, n=91) regions of the DS during the spontaneous SWO regime. The membrane potential of the Down states was stable during the recordings with no differences between striatal regions (DLS-MSNs= -69.89± 5.79 Hz, DMS-MSNs=-69.867±7.67 Hz, p=0.42). MSNs exhibited a rich subthreshold activity during Up states; however, they displayed a very low rate of spontaneous action potentials with similar frequencies along DS (DLS-MSNs=0.3±0.95 Hz, DMS-MSNs=0.24±0.53 Hz, p=0.58). The input resistance was higher for DLS-MSNs (DLS-MSNs= 310± 25 MΩ, DMS-MSNs=255±26 MΩ, p=0.04), as well as their capacitance (DLS-MSNs= 26.65± 7.56 pF, DMS-MSNs=21.68±2.26 pF, p=0.05). It is expected that changes in resistance and capacitance impact on the neuronal properties of synaptic integration, setting the gain and timing for their synaptic inputs. Thus, these results strongly suggest a particular integration of the spontaneous activity on DLS- and DMS-MSNs.

The spontaneous activity from DLS- and DMS-MSNs have different attributes.

We studied the properties of the SWO recorded in whole-cell from dorsal striatal MSNs. We first computed 13 different features of the SWO (Fig. 1D), from which 11 were statistically different between DLS- and DMS-MSNs. The average voltage in the Up states was higher in the DMS-MSNs, as well as their Up state standard deviation, minimum and maximum voltage during Up state and the total amplitude of the Up states (Fig. 1D, features nº 1-5). The upward and downward Up states transitions slopes and its ratio were also higher in DMS-MSNs (Fig. 1D, features nº 8-10). DLS-MSNs displayed higher number of peaks during the Up states and peak to peak amplitude, as well as longer Up states (Fig. 1D, features nº 11-13) [all features are explained in methods section]. Similar to cortical neurons (Sanchez-Vives and McCormick, 2000), Up states in MSNs are comprised by barrage of excitatory and inhibitory synaptic inputs (Reig and Silberberg, 2014). Therefore, these divergences in the integration of the SWO may reflect changes on the excitatory and inhibitory inputs that DLS- and DMS-MSNs receive. Following our result, we asked whether these differences were sufficient to classify DLS- and DMS-MSNs as two different populations. To do so, we searched for those features of the SWO whose combination maximized the accuracy of classifying MSNs corresponding to DLS or DMS circuits. Using a supervised machine learning technique, named Support Vector Machine (SVM) (Cortes and Vapnik, 1995) with a linear kernel, we performed a Recursive Feature Elimination (RFE) analysis to rank the 13 computed features of the SWO, according to their combined utility to classify DLS- or DMS-MSNs (Fig. 1E). The most relevant features were: first, the number of peaks in the Up states and second, the amplitude inside the Up state, which resulted in an accuracy score of 84.61±0.47% (Fig. 1E-G). When a third feature, the speed of transition from Up to Down states, was added, the accuracy increased to 88.56±0.57% (Fig. 1E-G). The addition of new parameters increased the accuracy up to ~92%, value at which accuracy remained stable (Fig. 1E). In a compromise between accuracy and interpretability, we selected the space formed by the three most relevant features for further analyses (Fig. 1H).

We performed an additional control analysis in which we classified the MSNs recorded from each animal as test set, using the remaining MSNs as training set. The accuracy following this calculation was 88.89% (Fig. S1), excluding the presence of inter-animal bias. Thus, our classifier based on the SWO of the MSNs indicates that DLS and DMS are identified as two distinct functional circuits.

In a second step, to further depict the differences in the SWO between DLS and DMS, we characterized the fast oscillations that are phase locked to the Up state during this brain state. High frequency oscillations are prominent during wakefulness and during the Up states of the SWO (Compte et al., 2008; Steriade et al., 1996). In the cortico-basal ganglia-thalamic loop, high frequency activity has been related with different functions and processes in health and disease (Brown, 2003; Feingold et al., 2015). For instance, exaggerated beta-band oscillations occur in Parkinson's Disease. With the porpoise to study possible differences in the high frequency oscillations of DLS and DMS, we compared the energy in the theta (6-10 Hz), beta (10-20 Hz) and gamma (20-80 Hz) bands of DLS- and DMS-MSNs (Fig. 2A, Fig. S2A-D). To that end, we applied the Noise-Assisted Multivariate Empirical Mode Decomposition (NA-MEMD), extracting the Intrinsic Mode Function (IMF) (see Methods) carrying the information of the different oscillatory activity bands (Fig. 1D, 2A and S2 A-D). The use of NA-MEMD let us obtain an enriched decomposition of the spontaneous activity compared to traditional techniques (Fourier decomposition, wavelet, and others), due to the non-linear properties of neural

oscillations (Averbeck et al., 2006; Cole and Voytek, 2017; Laurent, 1996; Shamir and Sompolinsky, 2004).

Our results show that the energy of the beta band was higher, and statistically different, in DMS-MSNs (Fig. 2B-D). On average, the peak of beta energy occurs in the first third of the Up state in both striatal regions, but it is closer to the beginning of the Up state for DMS-MSNs (Fig. 2C, DMS= -26.81±24.63°, DLS= -13.26±30.01°, p=0.0007). Following this divergence in beta band, we tested whether high frequency oscillation contained enough information to classify DLS- and DMS-MSNs. To that end, we trained an SVM with a linear kernel, as before. We obtained a classification accuracy of 60.19±10 % indicating that the high frequencies do not contribute to discriminate between DS regions. Moreover, the addition of the Beta energy as an extra feature to our previous classifier did not improve the level of discrimination obtained by the SWO features (Fig. S2E). Therefore, we conclude that DLS- and DMS-MSNs can be identified as belonging two different circuits based on the features of the Up states during the SWO.

DLS and DMS are two non-overlapping functional circuits in mouse
The features obtained from the SWO let us classify between DLS and DMS-MSNs. Accordingly, we used them to study the transition from one circuit to the other, exploring the properties of the SWO in dorsocentral striatum (DCS) (Fig. 3A), a hypothetical third region between DLS and DMS [see methods]. We considered three possible scenarios (Fig. 3A, B): If the activity of both circuits is mostly driven by the afferent inputs, we should observe a gradient moving from one circuit to the other (Hypothesis 1), or alternatively a third type of circuit (Hypothesis 2). Whether other factors, either extrinsic or intrinsic to the DS, control the spontaneous activity of these circuits, we should observe a sharp transition from one circuit to the other (Hypothesis 3). To address this question, we recorded MSNs in the DCS [see methods] in whole-cell, extracting the previous 13 features of the SWO (Table S1). Then, we projected the DCS-MSNs into the standardized space used for the classification of DLS- and DMS-MSNs (Fig. 3C), containing the values of number of peaks in the Up state, Up state amplitude and Up to Down transition slopes (Table S1, Fig. S3). In order to study the distribution of the DCS-MSNs along the classification boundary, we used the hypervector computed by the SVM to perform the classification (Fig. 3C, see methods). We projected the data onto this hypervector and compared the distribution of the MSNs recorded from DCS to determine whether they belonged to either DLS or DMS circuits, or whether they had different properties (Fig. 3D). If a sharp functional boundary exists between DLS and DMS, we would have recorded an undetermined amount of MSNs belonging to both circuits (Hypothesis 3). Otherwise, it would appear a new cluster with an undescribed third functional type of MSN, with an unknown distribution along the hypervector (Hypothesis 2) or in between DLS and DMS-MSNs distributions (Hypothesis 1). The distribution of MSNs recorded in the DCS was consistent with a mixture of MSNs from DLS and DMS circuits supporting a sharp transition between DLS and DMS regions [see discussion]. DCS distribution (Fig. 3D, green trace) was significantly different to either the DLS and DMS distributions (p< 0.01, Fig. 3D, red, blue and black traces respectively) and equivalent to the combination of both (p=0.681, Fig. 3D, green and black traces). Thus, we conclude that all recorded MSNs in DS belong to the previously described DMS and DLS functional circuits and they are separated by a sharp functional boundary.

DLS- and DMS-MSNs segregation is explained by their voltage transitions

To understand the main source of differences between the SWO of the DLS- and DMS-MSNs, we started by analysing the first ranked feature, number of peaks during Up states (Fig, 1D-E, feature nº11). We studied whether this difference occurred in cortical regions, from where they can be transmitted to the DS (Fig. 4A). It is known that striatal oscillations depend on the cortical ones (Kasanetz et al., 2002; Wilson, 1993) and in vitro slices containing the striatum are absent of spontaneous oscillations (Planert et al., 2013).

We performed *in vivo* whole-cell recordings in Layer V of FrA (frontal associative cortex), M1 (primary motor cortex), S1 (primary visual cortex) and V1 (primary visual cortex) (Fig. 4B). We found that S1 and M1 have significantly higher average number of peaks per Up state than FrA and V1 (Fig. 4C). This result together with previous anatomical descriptions (Alloway et al., 2006; Hoffer et al., 2003; Hooks et al., 2018) strongly suggest that the membrane dynamics inside the active Up state in the DLS are controlled by the sensory-motor cortical areas.

The main features for the classification of DLS- and DMS-MSNs (Fig. 1 G) were relative to state transitions (Up to Down transition) or intra-state transitions (number of peaks). In order to understand whether these differences may be caused by changes in the depolarizing and hyperpolarizing dynamics of MSNs during the SWO cycle, we studied the temporal properties of the voltage transitions. To analyse the convergence of synchronized hyper/depolarizing events onto MSNs that created detectable changes in the whole-cell recording, we first detected the sharp transitions of the voltage trace and phase locked them to the SWO cycle (Fig. 4D [see methods]). Thus, we obtained a measure of the temporal dynamics of hyper/depolarizing events of the MSNs during the SWO. Then, we computed their ratio for the DLS- and DMS-MSNs. We found differences in the depolarizing/hyperpolarizing ratio (DH ratio) of DLS- and DMS-MSNs aligned to the transitions between Up and Down states (Fig. 4E). The DH ratio was significantly higher, implying more excitation for DMS-MSNs during the transition from the Down to the Up state and higher hyperpolarization values during the transition from the Up to the Down state. This is consistent with the significant differences found for both Up states slopes (Fig. 1D, features 8-9).

In summary, our results show that the main differences between DLS- and DMS-MSNs SWO, selected by our classifier, are consistent with by the number and the temporal distribution of the voltage fluctuations during the Up states, suggesting that they are inherited from specific cortical areas and modulated by the striatal microcircuits.

Sequential propagation of the cortical SWO to the striatum

The SWO has a cortical origin, from where it propagates to the striatum activating MSNs (Kasanetz et al., 2002; Ketzef et al., 2017; Reig and Silberberg, 2016; Sáez et al., 2018; Wilson and Kawaguchi, 1996) and interneurons (Reig and Silberberg, 2014). In order to clarify how cortical SWOs are related with the dorsal striatum, we studied the correlations of the SWO between several cortical areas and MSNs located in DLS and DMS (Fig. 5). As before, we applied the NA-MEMD algorithm to extract the IMF carrying the SWO. Then, we used it to compute the cross-correlation between the whole-cell recordings of the MSNs in DLS or DMS with the pairs of simultaneous local field potentials (LFPs) recorded in FrA, M1, S1 and V1 (Fig. 5A-B). We found that neuronal activity from DLS- and DMS-MSNs had different correlations with FrA, M1 and V1 LFPs (Fig. 5DG). Correlation with FrA and M1 activity was higher for DLS neurons, while correlation with V1 was higher for neurons recorded from DMS. When we represented correlation values with FrA and V1, two clusters were clearly distinguishable, corresponding to

DMS- and DLS-MSNs (Fig. 5E), suggesting specific functional coupling to the cortex for both striatal regions.

Next, we analysed the probability of occurrence of an Up state recorded in a MSN, following the occurrence of a cortical Up state (Fig. 5F). Consistent with the correlation results, the probability that an Up state in FrA is followed by an Up state in DLS was significantly higher than in DMS. Similarly, Up states in V1 had higher probability to be followed by an Up state in DMS than in DLS. In summary, the activity from FrA and sensory-motor cortical regions (M1 and S1) is strongly related with DLS-MSNs, while the activity in V1 does with DMS-MSNs.

These results predicted that, following the known propagation of the SWO across the cortex, with a predominant rostro-caudal direction (Massimini et al., 2004; Ruiz-Mejias et al., 2011), a sequential activation could be expected along the DS. In order to test this hypothesis, we first compared the order of transitions to an Up state recorded in both poles of the cortex by pairs of LFPs. We found an Up state in FrA preceding an Up state in V1 in the 71% of times (p value< 0.0001, data not shown), demonstrating the preference of the rostro-caudal propagation of the SWO in cortex.

Finally, in order to test our prediction, we performed two simultaneous whole-cell patch-clamp recordings in two identified MSNs, located in the DLS and DMS (Fig. 6). A representative example is shown in Fig. 6A-D. In all 6 recorded pairs, the Up state in DLS-MSN preceded the one in DMS-MSN in 71.11± 9.37 % of the times (Fig. 6E, p= 0.0313), with an average delay of 53.24± 32.16 ms. In the rest of the cases (28.62± 8.77 %), in which DMS-MSN preceded the DLS-MSN, the delay was similar (average delay DLS-DMS= 51.91± 29.80 ms).

In conclusion, our results regarding the propagation of the cortical SWO to DS, confirm a stronger functional coupling of FrA to DLS and V1 to DMS. This generates a sequential activation of the DS during the SWO, in which in most of the cases, DLS precedes DMS by tens of milliseconds.

Direct and indirect MSNs have distinct properties in DLS and DMS
In a final step to deconstruct DS circuitry, we studied the spontaneous activity of MSNs corresponding to the direct or indirect pathway. To understand corticostriatal dynamics it is essential to apprehend how both pathways integrate the upstream cortical activity. To answer our question, we analysed the whole-cell recoding of the DS that were optogenetically identified as direct and indirect MSNs through the optopatcher (Fig. 7A, see methods). First, we analysed whether the features that were previously used to describe the SWO
(Fig. 1D) allowed us to differentiate between the MSNs belonging to each pathway in DLS and DMS. Our results show differences in the integration of the SWO between direct and indirect pathways only in DLS. We found three features of the SWO that were significant different when comparing direct and indirect MSNs in DLS: maximum and minimum voltage value of the Up state and the average value of the Up state (Fig. 7C). Then, we asked whether DLS- and DMS-MSNs could be subdivided in two additional populations corresponding with the direct or indirect MSNs in both striatal regions. The result did not yield separation, classification accuracy was 50% for DMS-MSNs and 64% for DLS-MSNs (Fig. 7B). High frequency oscillations were also similar between pathways in both striatal regions (Fig. S4).

Finally, we studied whether the differences in the electrophysiological properties between DLS- and DMS-MSNs, described before (Table 1), were dependent on one of the two pathways. Resistance values, both during Up and Down states, were significantly higher for indirect DLS-

MSNs compared to indirect DMS-MSNs, while no differences were found between direct DLS- and DMS-MSNs (Table 1).

In summary, our results show that DLS- and DMS-MSNs cannot be subdivided by the direct and indirect pathways. However, their differences in resistance and integration of the SWO demonstrate that direct and indirect MSNs have distinct properties in the DLS and DMS.

Discussion

Circuits are limited but not defined by their anatomy; different axonal projections cell types and synaptic dynamics delimit their spontaneous and evoked activity patterns. In this study, we have described two functional circuits in the anatomically homogenous DS, covering the medial and lateral regions and separated by a sharp functional boundary. We discriminated DLS and DMS circuits based on different properties of their spontaneous SWO recorded in MSNs by means of the *in vivo* whole-cell patch-clamp technique. To analyse the oscillatory activity, we used NA-MEMD algorithm (Rehman and Mandic, 2009) together with Hilbert transform (Huang et al., 1998). This algorithm is suited to decompose nonlinear nonstationary signals (Alegre-Cortés et al., 2018; Alegre-Cortés et al., 2016; Hu and Liang, 2014; Mandic et al., 2013). Given the well-known nonlinear properties of neural oscillations (Averbeck et al., 2006; Cole et al., 2017; Laurent, 1996; Shamir and Sompolinsky, 2004), the use of NA-MEMD leads to an increased detail of description when compared with traditional techniques (Alegre-Cortés et al., 2018; Hu and Liang, 2014). In our knowledge is the first time that it is used to analyse membrane oscillations of single neurons.

In humans, the dynamics of the SWO has demonstrated to be highly reproducible within and across subjects, providing information about the general state of the cerebral cortex and suggesting its applicability as a method to unravel functional circuits (Massimini et al., 2004). The analysis of the SWO has been used to detect neurophysiological alterations in humans and in a mouse models of neurological disorders such as; Alzheimer's disease (Busche et al., 2015),Down syndrome (Ruiz-Mejias et al., 2016), or Epilepsy (Amiri et al., 2019), among others. Here, we have extended this concept and proposing a methodology to study the brain functional connectivity.

DLS and DMS are two different functional circuits in mouse
We have shown that DS is divided in two different circuits, demonstrating the presence of a sharp functional transition from one circuit to another in the central region of the DS. This sharp transition is unexpected given the anatomical organization of corticostriatal projections, due to the profusion of axon terminals which provide overlapping inputs from multiple cortical areas to several regions of the DS (Alloway et al., 2006; Hintiryan et al., 2016; Hoffer et al., 2003; Hoover et al., 2003). Hence, an essential question arises: why do we find a sharp boundary instead of a gradient transition between DLS- and DMS-MSNs?
Our results show that the number of peaks during Up states is a critical feature to discriminate between DLS- and DMS-MSNs (Fig. 1E), being higher in DLS-MSNs (Fig. 1D, feature 11). In order to understand this result, we recorded layer V neurons in FrA, M1, S1 and V1, showing larger number of peaks during Up states in FrA, M1, and S1 than in V1. This suggest that, in agreement with previous anatomical studies (Alloway et al., 2006; Hoffer and Alloway, 2001; Hooks et al., 2018), somatosensory-motor and premotor cortical regions exert sharper modulation onto the

DLS-MSNs than onto DMS-MSNs (Fig. 4C), which is more anatomically connected with visual, associative, and limbic prefrontal cortical areas (Hintiryan et al., 2016; Hunnicutt et al., 2016). We also observed a higher number of peaks between M1-S1 with respect to FrA (Fig. 4C). These results suggest that the dynamics of the excitatory/inhibitory inputs during the Up states change between cortical areas and subsequently influence the striatum, opening an interesting question regarding the circuit mechanism underlying the observed cortical Up states differences.

Together with the cortical glutamatergic inputs, several additional factors may contribute to generate a functional boundary. Our classification reached ~90% of accuracy between DLS- and DMS-MSNs when the Up to Down state transition slopes were added (Fig. 3D, E). Moreover, the Down to Up transition and transition ratio were both significantly different between DLS- and DMS-MSNs as well (Fig. 3D). Therefore, Up states slopes are key elements to discriminate between dorsal striatal circuits as well. In order to better understand the slopes changes, we measured the time course of depolarized and hyperpolarized events of the SWO (Fig. 4D) and compared them using the DH ratio (Fig. 4E). Cortical SWO propagates to both types of MSNs but also to FS and ChI (Reig and Silberg, 2014). FS interneurons provide strong feedforward inhibition to MSNs (Gittis et al., 2010; Szydlowski et al., 2013; Tepper et al., 2008). Previous reports demonstrated the presence of a FS gradient (Gerfen et al., 1985; Kita et al., 1990; Monteiro et al., 2018) or diverse types of PV interneurons with different electrophysiological properties along DS (Monteiro et al., 2018; Muñoz-Manchado et al., 2018). Our results show more positive and negative ratio to the DMS-MSNs during the upward and downward SWO transitions slopes respectively, suggesting a different inhibitory recruitment in the DLS compared to DMS. In active cortical slices with SWO, both upward and downward transition slopes are controlled by $GABA_A$ and $GABA_B$ receptor activation (Sanchez-Vives et al., 2010; Perez-Zabalza et al., 2020). Based on these evidences, and beyond the glutamatergic inputs, we hypothesize that PV interneurons may greatly influence the SWO of DLS-MSNs making them different from DMS ones.

In addition, other mechanisms can also contribute to the generation of a functional boundary; for instance, a similar scenario exists for ChIs, which present a functional gradient with higher activity in the medial regions of the striatum (Abudukeyoumu et al., 2019; Matamales et al., 2016). ChIs could modulate differently the SWO in DLS- and DMS-MSNs by disynaptic inhibition (English et al., 2012). Under anaesthesia, dopaminergic neurons of the Substantia Nigra discharge spontaneous action potentials, in tonic or burst firing mode (Aristieta et al., 2016; Brown et al., 2009). This dopaminergic activity could induce different effects on the SWO in DLS- and DMS-MSNs, either directly mediated by distinct dopaminergic projections (Brown et al., 2009) or indirectly by their varied impact on ChIs activity along the mediolateral axis (Chuhma et al., 2018). Finally, it has been recently shown that SOM corticostriatal interneurons modulate the MSNs activity from motor cortical areas (Melzer et al., 2017; Rock et al., 2016). This type of interneuron could regulate the Up states transitions slopes directly, inhibiting the MSNs in their distal dendrites. Future work will have to determine whether FS, ChI, SOM or maybe other types of interneurons and neuromodulators contribute to the generation of the sharp functional boundary between DLS and DMS.

Propagation of the cortical SWO to the striatum
SWO propagates along the cortical network (Sanchez-Vives and McCormick, 2000) and is transmitted to the striatum (Reig and Silberg, 2014; Wilson and Kawaguchi, 1996) as well as

to other subcortical nuclei (Ros et al., 2009; Steriade et al., 1993a). Decortication or disruption of the cortical SWO impairs the striatal ones (Kasanetz et al., 2002; Wilson, 1993) and on the other hand, striatal slices are absent of rhythmic spontaneous activity (Planert et al., 2013), requiring the cerebral cortex to generate it (Vergara et al., 2003). As occurs in human slow-wave sleep (Massimini et al., 2004), anesthetized mice exhibit a predominant pattern of SWO propagation from rostral to caudal brain poles (Ruiz-Mejias et al., 2011). Here, we show how the correlation and the probability of transition of an Up state from FrA is higher to the DLS, while an Up state recorded in V1 has higher probability to propagate to the DMS (Fig. 5D-F). These results suggest a sequential activation of DMS and DLS, similar to the one between both poles of the cortex during the SWO. In order to test this hypothesis, we simultaneously recorded pairs of MSNs in DLS and DMS and we found a similar probability of an Up state to appear in DLS before DMS and to appear in FrA before V1 (Fig. 6). This description of the overall dynamics of corticostriatal transmission is based on functional, rather than anatomical connectivity, supporting a stronger coupling of caudal, sensory related cortical regions to DMS, whereas premotor and sensory-motor frontal regions are more tightly connected to DLS. Consistent with this, the sequential activation of both DS regions during a two-forced-decision-task has been recently reported, in which DMS was first activated during the sensory stimulation period, followed by an increase in DLS activity coinciding with decision making and motor preparation/execution (Peters et al., 2019).

The number of peaks during Up states, the first feature selected to classify between DLS- and DMS-MSNs, were prominent in the DLS-MSNs (Fig. 1D) as well as in S1 and M1 (Fig. 4B-C). These results together with the anatomical descriptions (Alloway et al., 2006; Hoffer et al., 2003; Hooks et al., 2018) strongly suggest that the activity inside the Up state of the DLS-MSNs is controlling by sensory-motor areas. In addition, FrA send copious projections to the DLS (Hintiryan et al., 2016; Hunnicutt et al., 2016) and we found that correlations and probability of propagation of Up states to DLS were higher for FrA than S1 and M1 (Fig. 5D-F). Therefore, we hypothesize that while the activity inside the Up state is governing by sensory-motor areas in DLS-MSNs, their Up states are triggered by more frontal regions, as FrA. In agreement with this, the blockage of synaptic transmission in S1 by TTX application during bilateral whisker stimulation block whisker responses but not the SWO in DLS-MSNs (Reig and Silberberg, 2016), demonstrating that their Up states were not triggered by S1.

High frequency oscillations over the SWO.
High frequency oscillations are usually associated with wakefulness. However, they also occur during the SWO in natural sleep, under anaesthesia (Steriade et al., 1996) and in isolated cortical slices (Compte et al., 2008). Especially relevant regarding the cortico-basal ganglia-thalamic loop are beta oscillations, which have been linked with the control of voluntary movements. Differences in Beta oscillations between caudate and putamen were reported in healthy monkeys at the end of a learned motor task (Feingold et al., 2015). Exaggerated Beta oscillations occur in Parkinson's Disease and in rats under dopamine depletion (Brown, 2003; Mallet et al., 2008; Sharott et al., 2017). Our results did not show differences between DLS- and DMS-MSNs for theta and gamma bands (Fig. S2), however a difference in beta energy was detected (Fig. 2B). Possible candidates to explain the observed higher beta energy in DMS-MSNs are ChI. This type of interneurons are more active in the DMS (Matamales et al., 2016) and can facilitate glutamate release from presynaptic terminals to the MSNs (Abudukeyoumu et al., 2019). Thus, they might promote the fast glutamatergic transmission observed in DMS.

Our work adds further detail to the description of beta oscillation in healthy mice thanks to the use of NA-MEMD together with Hilbert transform (Fig. 2, Fig. S5), providing a new substrate for the study of aberrant beta oscillations in Parkinson disease. This result may indicate that the aberrant oscillations in Parkinson disease may be produced by the misbalance of normal striatal dynamics, as previously suggested (McCarthy et al., 2011).

Integration of the SWO in the direct and indirect MSNs along the dorsal striatum

Cortical inputs innervate the DS targeting both the direct and indirect pathway MSNs (Doig et al., 2010; Kress et al., 2013; Wall et al., 2013). Both types of MSNs are co-activated during action initiation (Cui et al., 2013) and respond to tactile and visual stimulation (Reig and Silberberg, 2014). However, a large number of studies described differences between direct and indirect MSNs in the DLS: dMSNs have higher density of cortical and thalamic afferent synapses (Huerta-Ocampo et al., 2014), bigger response to whisker stimulation (Ketzef et al., 2017; Reig and Silberberg, 2014; Sippy et al., 2015) and based on *in vivo* SWO, it has been estimated that dMSNs receive stronger synaptic input than iMSNs (Filipović et al., 2019), among other differences. Here, we have described differences in the SWO between pathways in the DLS-MSNs (Fig. 7C), which are absent in the DMS-MSNs. Despite these differences, our results show that both DLS and DMS are unified circuits that involve both pathways.

On the other hand, the input resistance between MSNs were different between striatal regions, with bigger values in DLS. This result was weighted by the DLS-iMSNs, which displayed the highest values of resistance and were statistically different than iMSNs in the DMS (Table 1). Interestingly, this variation of resistance between iMSNs occurred over the whole oscillatory cycle; during Up states, when MSNs receive barrages of excitatory and inhibitory inputs, and during Down states, when MSNs are silent, suggesting that changes in resistance are independent of the glutamatergic inputs during the SWO. It is known that dopamine alters the input resistance of MSNs in vivo (Ketzef et al., 2017). Therefore, distinct dopaminergic innervation of the DLS- and DMS-MSNs (Chuhma et al., 2018) could underlie our differences in resistance between striatal regions.

Conclusion

Our novel approach, based on the analysis of the SWO, is a sensible strategy to provide a level of understanding between anatomy and function and it can be extended to the study of any brain circuit, in order to uncover their functional constrains.

Human caudate and putamen have been compared to the rodent DLS and DMS respectively (Balleine and O'Doherty, 2010). Based in the subthreshold activity of the MSNs we have shown how DLS and DMS are two non-overlapping circuits isolated by a sharp functional boundary. This work provides further understanding of the corticostriatal organization and reveals the biological substrate to divide the dorsal striatum in two different circuits. We have shown how DLS- and DMS-MSNs display independent spontaneous regimes during SWO brain state, that can only be explained by a combination of particular cortico-striatal functional connectivity and microcircuit properties. Therefore, our results provide the required understanding to support that this functional segregation is analogous to the anatomical and functional division of the primate striatum in caudate and putamen. Considering the relevance of the mouse striatum as a model for multiple human diseases, our results indicate that research will have to consider the idiosyncrasy of the two regions of the dorsal striatum.

Methods

Ethical permit. All the experimental procedures were conformed to the directive 2010/63/EU of the European Parliament and the RD 53/2013 Spanish regulation on the protection of animals use for scientific purposes, approved by the government of the Autonomous Community of Valencia, under the supervision of the *Consejo Superior de Investigaciones Científicas* and the Miguel Hernandez University Committee for Animal use in Laboratory.

Animal model. D2-Cre (ER44 line, GENSAT) mouse line was crossed with the Channelrhodopsin (ChR2)-YFP reporter mouse line (Ai32, the Jackson laboratory) to induce expression of ChR2 in indirect MSNs and was used to perform optogenetic differentiation of direct (dMSN) and indirect (iMSNs) MSNs (Ketzef et al., 2017), while performing electrophysiological recordings in the striatum (N= 36 mice). C57BL6 mice were used to perform the rest of the electrophysiological recordings of cortical neurons (n= 9).

Electrophysiological Recordings. Adult mice of both sexes (45 animals, 19 males and 26 females), between 12 and 44 weeks of age were used to perform the experiments (Table S2). Anaesthesia was induced by intraperitoneal injection of ketamine (75 mg/kg) and medetomidine (1 mg/kg) diluted in 0.9% NaCl. A maintaining dose of ketamine (30 mg/kg i.m.) was administrated every 2 hours, after changes in the frequency of spontaneous activity recorded by cortical LFP or reflex responses to paw pinches. Tracheotomy was performed to increase mechanical stability during recordings by decreasing breathing related movements. Mice were placed in a stereotaxic device and air enriched with oxygen was delivered through a thin tube placed 1 cm from the tracheal cannula. Core temperature was monitored and maintained at 36.5 ± 0.5°C using a feedback-controlled heating pad (FHC Inc.). Craniotomies were drilled (S210, Camo) at seven sites for patch-clamp and extracellular recordings (from Bregma): AP 0 mm, L 2.5 mm (DMS); AP 0 mm, L 4 mm (DLS); AP 0 mm, L 3,25 mm (DCS); AP 2.7 mm, L 1 mm (FrA); AP 1.5 mm, L 2.0 mm (M1); AP −1.5 mm, L 3.25 mm (S1); AP −3.5 mm from Bregma, L 2.5 mm (V1) (following Paxinos and Franklin, 2001). When additional paired recordings were performed, the craniotomies were drilled at the following coordinates (from Bregma): AP 0 mm, L 1 mm (DMS); AP 0 mm, L 4 mm (DLS). Animals were sacrificed after recordings by injecting an overdose of sodium pentobarbital (200 mg/kg I.P.).

Whole-cell recordings were obtained in the dorsal striatum from DLS, DMS, DCS between 2013-2647 µm depth. The number of striatal neurons was 126: DLS= 77, from which direct MSNs= 49, indirect MSNs= 34; DMS n= 91, from which direct MSNs= 65, indirect MSNs= 32; DCS n= 17, from which dDCS= 10, iDCS= 7. Cortical neurons (n= 26) were recorded from layer V in FrA, M1, S1 and V1 at a depth of 675-926 µm. In detail, the number of recorded neurons was: Frontal association cortex (FrA, n= 8), primary motor cortex (M1, n= 6), primary somatosensory cortex (S1, n= 6), and primary visual cortex (n= 6). Additional paired simultaneous whole-cell recordings of MSNs (n=6 pairs) were obtained from DLS and DMS between 2039-2348 µm of depth, by means of two micromanipulators (Luigs & Neumann, MRE/MLE Mini 25). All of them in a perpendicular penetration angle of ~30º, except for DCS cells, in which the angle was ~20º. The exposed brain was continuously covered with 0.9% NaCl to prevent drying. Signals were amplified using MultiClamp 700B amplifier (Molecular Devices) and digitized at 20 kHz with a CED acquisition board and Spike 2 software (Cambridge Electronic Design).
Borosilicate patch pipettes (1B150F-4, WPI), were pulled with a Flaming/Brown micropipette puller P-1000 (Sutter Instruments) and had an initial resistance of 6–12 MΩ. Pipettes were

back-filled with intracellular solution containing: 125 mM K-gluconate, 10 mM KCl, 10 mM Na-Phosphocreatine, 10 mM HEPES, 4 mM ATP-Mg and 0.3 mM GTP-Na. pH and osmolality were adjusted to ~7.4 and ~280 mOsm/L respectively. Biocytin (0.2-0.4 %, Sigma Aldrich) was then added to the intracellular solution to perform cell reconstruction after the experiment. To perform the analysis, 100 seconds of spontaneous activity (no current injection, no stimulation) were used from the recording. Input resistance (Table 1) was measured as the slope of a linear fit between injected depolarizing and hyperpolarizing current steps and membrane potential. Also, in order to improve the quantification of the described inward membrane rectification at hyperpolarized values of the MSNs membrane potentials, mean resistance in response to the negative and positive steps delivered at Up and Down states was also analysed (Table 1), as in Reig and Silberberg 2014 (Reig and Silberberg, 2014).

Neurons were identified according to their recorded electrical properties and following morphological staining, according to the aspiny dendrites in the case of the ChI and FS interneurons. FS interneurons displayed narrow action potentials, relatively depolarized resting membrane potential, high discharge rate of action potentials, and no apparent inward rectification. Cholinergic interneurons were characterized by their depolarized membrane potential, voltage sag response to current step injections and spontaneous tonic discharge. From the whole population of recorded neurons, 2 of them were identify as FS, 1 in the DLS and other in the DMS, 3 ChI were recorded in the DMS. These 5 striatal interneurons were excluded from the data set. Neurons having a deviation by more than 10 mV from their initial resting membrane potential were excluded from analysis. Only MSNs and cortical neurons were included in the data set.

Extracellular recordings were obtained using unipolar tungsten electrodes with impedance of 1 to 2 MΩ. The electrodes were placed in infragranular layers (~1000 µM depth from the pia) of two cortical regions in each experiment, from FrA, M1, S1, V1 cortex with an angle between 15° and 25°. Recordings were amplified using a differential AC Amplifier model 1700 (A-M Systems) and digitized at 20 KHz with CED and Spike-2, simultaneously with the whole-cell recording.

Optogenetic identification of *in vivo* recorded neurons. In order to identify "on line" the specific type of MSNs belonging to the direct (dMSNs) and indirect (iMSNs) pathways, the optopatcher was used (Katz et al., 2019; Katz et al., 2013; Ketzef et al., 2017) (A-M systems, WA USA). Controlled pulses (SLA-1000-2, two channel universal LED driver, Mightex systems) of blue light (Fibre-coupled LED light source FCS-0470-000, 470 nm, Mightex systems) through Spike 2 software were delivered using an optic fibre (200 µm diameter, handmade) inserted into the patch-pipette, while recording their spontaneous activity (Fig. 7A). One or two serial pulses with 5 light steps of 500 ms each were delivered every 2 s with increasing intensity from 20% to 100 % of full LED power (minimal light intensity was 0.166 mW, maximal intensity was 0.83 mW measured at the tip of the fibre). Power light was measured with an energy meter console (PM100D, Thorlabs). Positive cells responded to light pulses by depolarizing the membrane potential (Fig. 7A, upper trace), responding within 2.69 ± 1.37 ms (ranging from 0.8 to 5 ms) to light pulses by a step-like depolarization of 13.3 ± 8.28 mV (ranging from 4 to 33 mV), therefore were classified as indirect MSNs. Negative cells did not show any depolarization to light pulses (Fig. 7A, bottom trace) and were classified as a putative direct MSNs.

Morphological reconstruction. At the end of each experiment, mice were sacrificed with a lethal dose of sodium pentobarbital and perfused with a solution containing 4% paraformaldehyde in 0.1 M phosphate buffer (PB, pH 7.4). Brains were extracted and stored in PBS solution until the cutting. Before cutting, brains were transferred into PBS containing 30 %

sucrose for at least 48 hours. Coronal slices (25 µm thick) containing the entire striatum from the recorded side (from AP 1.4 mm to AP -1.3 mm, following Paxinos and Franklin, 2001), were obtained using a digital automatic cryotome and collected on gelatine coated slides. Sections were incubated over night with Cy3-conjugated streptavidin (Jackson Immuno Research Laboratories) diluted (1:1000) in 1 % BSA, 0.3 % Triton-X 100 in 0.1 M PBS. Finally, the glass slides were covered with mowiol (Calbiochem) and imaged. Neurons were then reconstructed using a fluorescence microscope (DM 6000B, Leica) and a camera (DC350 FX, Leica) and then processed by ImageJ.

SWO computation using NA-MEMD. We used Noise-assisted Multivariate Empirical Mode Decomposition (NA-MEMD) algorithm (Ur Rehman and Mandic, 2011) together with Hilbert transform (Huang et al., 1998) for the analysis of oscillations both in LFPs and MSNS membrane potential. Because neuronal oscillations are characterized by nonlinear properties, this algorithm is suited to decompose nonlinear nonstationary signals (Alegre-Cortés et al., 2018; Alegre-Cortés et al., 2016; Hu and Liang, 2014; Mandic et al., 2013). The original EMD (Huang et al., 1998) is a data driven algorithm suitable for nonlinear and non-stationary signals that does not rely on any predetermined template. It decomposes a given signal into a subset of oscillatory modes called Intrinsic Mode Functions (IMFs). Each IMF contains the oscillations of the original data in a certain frequency range. Then, Hilbert transform is applied onto each IMF in order to compute its instantaneous frequency and amplitude. The MEMD (Rehman and Mandic, 2009) is a multivariate extension of the original EMD to n-dimensional signals. The MEMD is computed simultaneously in all dimensions of the signal to ensure the same number of IMFs as output. In addition, new dimensions can be added to the data containing White Gaussian Noise (WGN) to increase its performance, as it has been described that WGN addition reduces mode mixing produced by signal intermittence (Wu and Huang, 2009), acting as a quasidyadic filter that enhances time frequency resolution (Flandrin et al., 2004; Ur Rehman and Mandic, 2011). The application of MEMD to the desired signal together with extra White Gaussian Noise dimensions is known as NA-MEMD analysis (Ur Rehman and Mandic, 2011). EMDs algorithms plus Hilbert Transform have been increasingly used in neuroscience during the last years as they produce an enriched Time-Frequency (T-F) Spectrum when compared to traditional T-F linear tools (Alegre-Cortés et al., 2016; Hu and Liang, 2014). In our work, we applied NA-MEMD algorithm to a multivariate signal composed by the intracellular recording, both LFPs and one extra WGN channel. By means of this analysis, we could warrant that the number of IMFs was the same in the intracellular and the LFPs recordings for a direct comparison of the SWO extracted using this method. In order to apply NA-MEMD analysis to our data, we adapted MEMD Matlab package (http://www.commsp.ee.ic.ac.uk/mandic/research/emd.htm). Standard stopping criterion was described elsewhere (Rilling et al., 2003). At last, we extracted the IMF carrying the SWO as the one with maximum correlation with the membrane voltage and visually confirmed it in all the recorded cells (Fig. 1C). Once we isolated the SWO of each recording using NA-MEMD, they were stored for further analysis.

Hilbert transform. We computed the frequency of the SWO, and theta (6-10 Hz), beta (10-20 Hz) and gamma (20-80 Hz) bands as the instantaneous frequency using the Hilbert transform (Huang et al., 1998). For a given time series x(t), its Hilbert transform H(x)(t) is defined as:

$$d(x)(t) = \frac{1}{\pi} C \int_{-\infty}^{\infty} \frac{x(t')}{t - t'} dt'$$

where C indicates the Cauchy principal value. Hilbert transform results in a complex sequence with a real part which is the original data and an imaginary part which is a version of the original data with a 90° phase shift; this analytic signal is useful to calculate instantaneous amplitude and frequency; instantaneous amplitude is the amplitude of H(x)(t), instantaneous frequency is the time rate of change of the instantaneous phase angle.

Extraction of Up and Down states. In order to isolate Up and Down states from the membrane voltage we started by smoothing the trace using a 200 ms window. Then, we extracted the Up states using a threshold consisting of the mean value of the membrane potential (Vm) plus 0.5 standard deviations. Then, we merged the parts of fragmented Up states, detected as an interval shorter than 250 ms between the transition of a prospective Up state to a prospective Down state to the transition from a prospective Down state to a prospective Up state. Finally, detected Up states shorter than 200 ms where discarded.

When studying propagation of Up states, we considered two Up states from two different cortical regions, or a cortical LFP recording and an MSN to be part of the same SWO, thus propagating from one region to the other, when their onsets were closer in time than 500 ms.

Feature extraction of Up states. In order to describe the properties of the SWO of the recorded MSNs, we computed 13 parameters (Fig. 1D). First, we subtracted the IMFs carrying oscillations faster than 50 Hz from the MSNs membrane voltage recordings to eliminate the spikes. Then, we divided the trace into Up and Down states and calculated the different features; the mean of the voltage in the Up state (1), Up state standard deviation (2), minimum voltage in the Up state (3) and maximum voltage in the up state (4). Additionally, we computed the peak to peak distance (12), which is the difference between the maximum and minimum values of the Up state, the maximum (6) and minimum (7) value of the Up state derivative and the number of peaks in the Up state (11), which described the presence of diplets or triplets during the Up state (Fig. 4A, B). To do so, we used the *findpeaks* Matlab function on the smoothed (200 ms smooth) Up state with a minimum peak height of 0.3 standard deviations and a minimum distance between peaks of 160 ms. We also computed the Up state length (13), amplitude (5) and the speed of the transition from the Down to the Up state (8), and from the Up to the Down state (9) as well as the Slope transition ratio (10), which consisted of the magnitude of the Down to Up transition relative to the Up to Down transition. In order to measure the slope speeds, we used a vector consisting in -100 to +200 ms (300 ms total) relative to the transition either to the Up or the Down state. Then, we smoothed (100 ms smooth) the selected region and computed its derivative. We delimited the slope computing the region over a threshold calculated as the mean plus 0.3 standard deviations of the derivative. Once we had delimited the transition, we fitted a lineal function to that region of the recording to compute the slope speed.

Detection of intracellular synaptic events. Intracellular synaptic currents were extracted using sharp deflections of the membrane voltage using a modification of the method presented here (Compte et al., 2008). In brief, we applied a Parks-McClellan low-pass differentiator filter (order 20, cut off at 200 Hz) and computed the first derivative of the intracellular membrane voltage. Then, we extracted the temporal profile of depolarizing and hyperpolarizing events to the recorded cell using a threshold of the first derivative. We empirically determined a 2 standard deviations threshold to extract them. Next, we aligned the detected membrane deflections for both the hyperpolarizing and depolarizing events to the phase of the SWO as computed using NA-MEMD and normalized them to their respective maximum (Fig. 4E) in order to study the profile of the recruitment of excitation and inhibition during SWO. We computed the

depolarizing/hyperpolarizing ratio subtracting the hyperpolarizing normalized value to the depolarizing one at each phase point.

For our whole-cell recordings, we have used a low-chloride intracellular solution (described above), in which the reversal potential for excitation is ~ -5 mV and $GABA_A$ inhibition ~ -70 mV. The average membrane potential during the Down states of the recorded MSNs was -69,2± 6,48 mV for DLS-MSNs and -69,31± 7,68 mV for DMS-MSNs. Around these voltages, the membrane fluctuations are depolarizing and excitatory. However, during the Up states (mean voltage values: -57,24± 6,34 mV for DLS-MSNs and -52,95± 7,08 mV for DMS-MSNs), both depolarized and hyperpolarized events can be detected (Fig. 4D). As previously discussed (Compte et al., 2008), this method does not extract all the EPSPs or IPSPs of the recorded cell, but it provides a major picture of the convergence of coordinated excitatory or inhibitory inputs onto the recorded cell that produced sharp deflections in the membrane voltage. Therefore, this method is sufficient to study the temporal profiles of hyper/depolarizing events onto the recorded cells during SWO.

Lateromedial classification. We classified MSNs from DLS and DMS using a Support Vector Machine (SVM) (Cortes and Vapnik, 1995) with a linear kernel. This supervised classification method estimates a hyperplane to separate both populations. Classification was cross-validated 10 times.

In order to determine which features were more relevant for the classification of DLS- and DMS-MSNs, we used Recursive Feature Elimination. This approach organizes the features relative to their importance to the classification by recursively pruning the least relevant features starting from the initial set.

MSNs distribution along the classification boundary. Support Vector Machines (SVMs) are supervised classifiers that predict the category of a new entry based on their previous training history and are not suitable to predict new categories in novel data. In order to study the spontaneous activity of DCS-MSNs, we examined its distribution in the classification space, relative to the classification boundary instead of directly using the output of a classifier. This approach provides information, not only about whether the new data (DCS-MSNs) creates a new cluster in the classification space, but also about where is it placed along the classification axis used to discriminate between DLS and DMS-MSNs. Hence, it provides a more informative approach to the distribution of DCS-MSNs relative to DLS and DMSMSNs than a direct multi-category classification (i.e. discriminate versus Hypothesis 1 and 2, see results).

To calculate the distribution of the data along the classification boundary, we first obtained the coefficients and intersection values of the SVM used to classify the MSNs belonging to DLS and DMS. Once we obtained these parameters, we used them to project all data points to this hypervector and computed the probability density function of the new distributions (Fig. 3D). Now, the data was distributed along a single hypervector and we could compute the distribution of the MSNs along the classification boundary in a comprehensible manner to study the distribution of the DCS-MSNs.

Statistical analysis. All experimental comparisons were tested using the Wilcoxon Rank-sum test except stated otherwise. Directionality of the SWO in the cortex and in the striatum was tested using Wilcoxon signed rank test. Phase locking of different frequency bands was tested using Rayleigh test. When required, alpha values for multiple comparisons were corrected using Holm-Bonferroni correction. Error bars presented in the graphs represent the standard deviation unless stated otherwise.

Acknowledgements

We thank Gilad Silberberg (Karolinska Institutet) for generous donation of D2-Cre(ChR2)-YFP mouse and for helping us to start-up the laboratory. We also want to thank Gilad Silberberg, Juan Lerma, Sandra Jurado, John F. Wesseling, Maya Ketzef, Cristina García-Frigola, Juan Pérez-Fernández and Roberto de la Torre for critical reading and discussions on this. J. A-C and M. S. are supported by the CSIC-Severo Ochoa Excellence Programme of the Instituto de Neurociencias [SEV2013-0317]; R. M. is supported by La Caixa-Severo Ochoa [2016/00006/001]. The study was supported by MINECO Fellowship [BES-2015-072187], Starting Grant I+D Jovenes Investigadores [BFU2014-60809-IN] and CSIC-Severo Ochoa Excellence Programmes of the Instituto de Neurociencias [SEV-2013-0317 and SEV-2017-0723].

Declaration of interest

The authors declare no competing interests.

Author contributions

R. R. conceived the project and wrote the manuscript. J. A-C. analysed the data and wrote the manuscript. M. S. conducted most of the experiments and collected data. R.M. performed most of the cortical patch-clamp recordings. All authors commented on the manuscript.

Author ORCIDs

Javier Alegre-Cortés   0000-0003-0888-7542

María Sáez   0000-0001-9137-6692

Roberto Montanari   0000-0002-4331-8460

Ramón Reig   0000-0002-6475-4181

References

Abudukeyoumu, N., Hernandez-Flores, T., Garcia-Munoz, M., and Arbuthnott, G.W. (2019). Cholinergic modulation of striatal microcircuits. European Journal of Neuroscience *49*, 604-622.

Adler, A., Katabi, S., Finkes, I., Israel, Z., Prut, Y., and Bergman, H. (2012). Temporal convergence of dynamic cell assemblies in the striato-pallidal network. J Neurosci *32*, 2473-2484.

Alegre-Cortés, J., Soto-Sánchez, C., Albarracín, A.L., Farfán, F.D., Val-Calvo, M., Ferrandez, J.M., and Fernandez, E. (2018). Toward an Improvement of the Analysis of Neural Coding. Frontiers in neuroinformatics *11*, 77.

Alegre-Cortés, J., Soto-Sánchez, C., Pizá, Á., Albarracín, A.L., Farfán, F.D., Felice, C.J., and Fernández, E. (2016). Time–frequency analysis of neuronal populations with instantaneous resolution based on noise-assisted multivariate empirical mode decomposition. Journal of neuroscience methods *267*, 35-44.

Alloway, K.D., Lou, L., Nwabueze-Ogbo, F., and Chakrabarti, S. (2006). Topography of cortical projections to the dorsolateral neostriatum in rats: multiple overlapping sensorimotor pathways. J Comp Neurol *499*, 33-48.

Alloway, K.D., Smith, J.B., Beauchemin, K.J., and Olson, M.L. (2009). Bilateral projections from rat MI whisker cortex to the neostriatum, thalamus, and claustrum: forebrain circuits for modulating whisking behavior. J Comp Neurol *515*, 548-564.

Amiri, M., Frauscher, B., and Gotman, J. (2019). Interictal coupling of HFOs and slow oscillations predicts the seizure-onset pattern in mesiotemporal lobe epilepsy. Epilepsia *60*, 1160-1170.

Aristieta, A., Ruiz-Ortega, J., Miguelez, C., Morera-Herreras, T., and Ugedo, L. (2016). Chronic L-DOPA administration increases the firing rate but does not reverse enhanced slow frequency oscillatory activity and synchronization in substantia nigra pars reticulata neurons from 6-hydroxydopamine-lesioned rats. Neurobiology of disease *89*, 88-100.

Averbeck, B.B., Latham, P.E., and Pouget, A. (2006). Neural correlations, population coding and computation. Nat Rev Neurosci *7*, 358-366.

Balleine, B.W., and O'Doherty, J.P. (2010). Human and rodent homologies in action control: corticostriatal determinants of goal-directed and habitual action. Neuropsychopharmacology *35*, 48-69.

Berke, J.D., Okatan, M., Skurski, J., and Eichenbaum, H.B. (2004). Oscillatory entrainment of striatal neurons in freely moving rats. Neuron *43*, 883-896.

Brown, M.T., Henny, P., Bolam, J.P., and Magill, P.J. (2009). Activity of neurochemically heterogeneous dopaminergic neurons in the substantia nigra during spontaneous and driven changes in brain state. Journal of Neuroscience *29*, 2915-2925.

Brown, P. (2003). Oscillatory nature of human basal ganglia activity: relationship to the pathophysiology of Parkinson's disease. Mov Disord *18*, 357-363.

Busche, M.A., Kekus, M., Adelsberger, H., Noda, T., Forstl, H., Nelken, I., and Konnerth, A. (2015). Rescue of long-range circuit dysfunction in Alzheimer's disease models. Nat Neurosci *18*, 1623-1630.

Chuhma, N., Mingote, S., Yetnikoff, L., Kalmbach, A., Ma, T., Ztaou, S., Sienna, A.-C., Tepler, S., Poulin, J.-F., and Ansorge, M. (2018). Dopamine neuron glutamate cotransmission evokes a delayed excitation in lateral dorsal striatal cholinergic interneurons. Elife *7*, e39786.

Cole, S.R., van der Meij, R., Peterson, E.J., de Hemptinne, C., Starr, P.A., and Voytek, B. (2017). Nonsinusoidal Beta Oscillations Reflect Cortical Pathophysiology in Parkinson's Disease. J Neurosci *37*, 4830-4840.

Cole, S.R., and Voytek, B. (2017). Brain Oscillations and the Importance of Waveform Shape. Trends Cogn Sci *21*, 137-149.

Compte, A., Reig, R., Descalzo, V.F., Harvey, M.A., Puccini, G.D., and Sanchez-Vives, M.V. (2008). Spontaneous high-frequency (10–80 Hz) oscillations during up states in the cerebral cortex in vitro. Journal of Neuroscience *28*, 13828-13844.

Cortes, C., and Vapnik, V. (1995). Support-vector networks. Machine learning *20*, 273-297.

Cowan, R.L., and Wilson, C.J. (1994). Spontaneous firing patterns and axonal projections of single corticostriatal neurons in the rat medial agranular cortex. J Neurophysiol *71*, 17-32.

Cui, G., Jun, S.B., Jin, X., Pham, M.D., Vogel, S.S., Lovinger, D.M., and Costa, R.M. (2013). Concurrent activation of striatal direct and indirect pathways during action initiation. Nature *494*, 238.

Doig, N.M., Moss, J., and Bolam, J.P. (2010). Cortical and thalamic innervation of direct and indirect pathway medium-sized spiny neurons in mouse striatum. Journal of Neuroscience *30*, 14610-14618.

English, D.F., Ibanez-Sandoval, O., Stark, E., Tecuapetla, F., Buzsáki, G., Deisseroth, K., Tepper, J.M., and Koos, T. (2012). GABAergic circuits mediate the reinforcement-related signals of striatal cholinergic interneurons. Nature neuroscience *15*, 123.

Faure, A., Haberland, U., Condé, F., and El Massioui, N. (2005). Lesion to the nigrostriatal dopamine system disrupts stimulus-response habit formation. Journal of Neuroscience *25*, 2771-2780.

Feingold, J., Gibson, D.J., DePasquale, B., and Graybiel, A.M. (2015). Bursts of beta oscillation differentiate postperformance activity in the striatum and motor cortex of monkeys performing movement tasks. Proceedings of the National Academy of Sciences *112*, 13687-13692.

Filipović, M., Ketzef, M., Reig, R., Aertsen, A., Silberberg, G., and Kumar, A. (2019). Direct pathway neurons in mouse dorsolateral striatum in vivo receive stronger synaptic input than indirect pathway neurons. Journal of Neurophysiology *122*, 2294-2303.

Flaherty, A.W., and Graybiel, A. (1991). Corticostriatal transformations in the primate somatosensory system. Projections from physiologically mapped body-part representations. Journal of Neurophysiology *66*, 1249-1263.

Flandrin, P., Rilling, G., and Goncalves, P. (2004). Empirical mode decomposition as a filter bank. IEEE signal processing letters *11*, 112-114.

Gerfen, C.R., Baimbridge, K.G., and Miller, J.J. (1985). The neostriatal mosaic: compartmental distribution of calcium-binding protein and parvalbumin in the basal ganglia of the rat and monkey. Proceedings of the National Academy of Sciences *82*, 8780-8784.

Getting, P.A. (1989). Emerging principles governing the operation of neural networks. Annual review of neuroscience *12*, 185-204.

Gittis, A.H., Nelson, A.B., Thwin, M.T., Palop, J.J., and Kreitzer, A.C. (2010). Distinct roles of GABAergic interneurons in the regulation of striatal output pathways. Journal of Neuroscience *30*, 2223-2234.

Graybiel, A.M. (2008). Habits, rituals, and the evaluative brain. Annu Rev Neurosci *31*, 359-387.

Hauber, W., and Schmidt, W.J. (1994). Differential effects of lesions of the dorsomedial and dorsolateral caudate-putamen on reaction time performance in rats. Behav Brain Res *60*, 211-215.

Hilário, M.R., and Costa, R.M. (2008). High on habits. Frontiers in neuroscience *2*, 30.

Hintiryan, H., Foster, N.N., Bowman, I., Bay, M., Song, M.Y., Gou, L., Yamashita, S., Bienkowski, M.S., Zingg, B., Zhu, M._, et al._ (2016). The mouse cortico-striatal projectome. Nat Neurosci _19_, 1100-1114.

Hoffer, Z.S., and Alloway, K.D. (2001). Organization of corticostriatal projections from the vibrissal representations in the primary motor and somatosensory cortical areas of rodents. Journal of Comparative Neurology _439_, 87-103.

Hoffer, Z.S., Hoover, J.E., and Alloway, K.D. (2003). Sensorimotor corticocortical projections from rat barrel cortex have an anisotropic organization that facilitates integration of inputs from whiskers in the same row. J Comp Neurol _466_, 525-544.

Hooks, B.M., Papale, A.E., Paletzki, R.F., Feroze, M.W., Eastwood, B.S., Couey, J.J., Winnubst, J., Chandrashekar, J., and Gerfen, C.R. (2018). Topographic precision in sensory and motor corticostriatal projections varies across cell type and cortical area. Nat Commun _9_, 3549.

Hoover, J.E., Hoffer, Z.S., and Alloway, K.D. (2003). Projections from primary somatosensory cortex to the neostriatum: the role of somatotopic continuity in corticostriatal convergence. J Neurophysiol _89_, 1576-1587.

Hu, M., and Liang, H. (2014). Search for information-bearing components in neural data. PLoS One _9_, e99793.

Huang, N.E., Shen, Z., Long, S.R., Wu, M.C., Shih, H.H., Zheng, Q., Yen, N.-C., Tung, C.C., and Liu, H.H. (1998). The empirical mode decomposition and the Hilbert spectrum for nonlinear and non-stationary time series analysis. Proceedings of the Royal Society of London Series A: Mathematical, Physical and Engineering Sciences _454_, 903-995.

Huerta-Ocampo, I., Mena-Segovia, J., and Bolam, J.P. (2014). Convergence of cortical and thalamic input to direct and indirect pathway medium spiny neurons in the striatum. Brain Structure and Function _219_, 1787-1800.

Hunnicutt, B.J., Jongbloets, B.C., Birdsong, W.T., Gertz, K.J., Zhong, H., and Mao, T. (2016). A comprehensive excitatory input map of the striatum reveals novel functional organization. Elife _5_, e19103.

Ikemoto, S. (2007). Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens–olfactory tubercle complex. Brain research reviews _56_, 27-78.

Kasanetz, F., Riquelme, L.A., and Murer, M.G. (2002). Disruption of the two-state membrane potential of striatal neurones during cortical desynchronisation in anaesthetised rats. J Physiol _543_, 577-589.

Katz, Y., Sokoletsky, M., and Lampl, I. (2019). In-vivo optogenetics and pharmacology in deep intracellular recordings. Journal of neuroscience methods, 108324.

Katz, Y., Yizhar, O., Staiger, J., and Lampl, I. (2013). Optopatcher—an electrode holder for simultaneous intracellular patch-clamp recording and optical manipulation. Journal of neuroscience methods _214_, 113-117.

Ketzef, M., Spigolon, G., Johansson, Y., Bonito-Oliva, A., Fisone, G., and Silberberg, G. (2017). Dopamine depletion impairs bilateral sensory processing in the striatum in a pathway-dependent manner. Neuron _94_, 855-865. e855.

Kincaid, A.E., Zheng, T., and Wilson, C.J. (1998). Connectivity and convergence of single corticostriatal axons. Journal of Neuroscience *18*, 4722-4731.

Kita, H., and Kitai, S.T. (1988). Glutamate decarboxylase immunoreactive neurons in rat neostriatum: their morphological types and populations. Brain Res *447*, 346-352.

Kita, H., Kosaka, T., and Heizmann, C. (1990). Parvalbumin-immunoreactive neurons in the rat neostriatum: a light and electron microscopic study. Brain research *536*, 1-15.

Kress, G.J., Yamawaki, N., Wokosin, D.L., Wickersham, I.R., Shepherd, G.M., and Surmeier, D.J. (2013). Convergent cortical innervation of striatal projection neurons. Nature neuroscience *16*, 665.

Laurent, G. (1996). Dynamical representation of odors by oscillating and evolving neural assemblies. Trends Neurosci *19*, 489-496.

Lerner, T.N., Shilyansky, C., Davidson, T.J., Evans, K.E., Beier, K.T., Zalocusky, K.A., Crow, A.K., Malenka, R.C., Luo, L., and Tomer, R. (2015). Intact-brain analyses reveal distinct information carried by SNc dopamine subcircuits. Cell *162*, 635-647.

Levesque, M., Charara, A., Gagnon, S., Parent, A., and Deschenes, M. (1996). Corticostriatal projections from layer V cells in rat are collaterals of long-range corticofugal axons. Brain Res *709*, 311-315.

Luczak, A., McNaughton, B.L., and Harris, K.D. (2015). Packet-based communication in the cortex. Nature Reviews Neuroscience *16*, 745-755.

Mallet, N., Pogosyan, A., Sharott, A., Csicsvari, J., Bolam, J.P., Brown, P., and Magill, P.J. (2008). Disrupted dopamine transmission and the emergence of exaggerated beta oscillations in subthalamic nucleus and cerebral cortex. Journal of Neuroscience *28*, 4795-4806.

Mandic, D.P., ur Rehman, N., Wu, Z., and Huang, N.E. (2013). Empirical mode decomposition-based time-frequency analysis of multivariate signals: The power of adaptive data analysis. IEEE signal processing magazine *30*, 74-86.

Massimini, M., Huber, R., Ferrarelli, F., Hill, S., and Tononi, G. (2004). The sleep slow oscillation as a traveling wave. Journal of Neuroscience *24*, 6862-6870.

Matamales, M., Götz, J., and Bertran-Gonzalez, J. (2016). Quantitative imaging of cholinergic interneurons reveals a distinctive spatial organization and a functional gradient across the mouse striatum. PloS one *11*, e0157682.

McCarthy, M., Moore-Kochlacs, C., Gu, X., Boyden, E., Han, X., and Kopell, N. (2011). Striatal origin of the pathologic beta oscillations in Parkinson's disease. Proceedings of the National Academy of Sciences *108*, 11620-11625.

Melzer, S., Gil, M., Koser, D.E., Michael, M., Huang, K.W., and Monyer, H. (2017). Distinct Corticostriatal GABAergic Neurons Modulate Striatal Output Neurons and Motor Activity. Cell Rep *19*, 1045-1055.

Monteiro, P., Barak, B., Zhou, Y., McRae, R., Rodrigues, D., Wickersham, I.R., and Feng, G. (2018). Dichotomous parvalbumin interneuron populations in dorsolateral and dorsomedial striatum. J Physiol *596*, 3695-3707.

Muñoz-Manchado, A.B., Gonzales, C.B., Zeisel, A., Munguba, H., Bekkouche, B., Skene, N.G., Lönnerberg, P., Ryge, J., Harris, K.D., and Linnarsson, S. (2018). Diversity of interneurons in the dorsal striatum revealed by single-cell RNA sequencing and PatchSeq. Cell reports *24*, 2179-2190. e2177.

Packard, M.G., and Knowlton, B.J. (2002). Learning and memory functions of the Basal Ganglia. Annu Rev Neurosci *25*, 563-593.

Perez-Zabalza, M., Reig, R., Manrique, J., Jercog, D., Winograd, M., Parga, N., Sanchez-Vives, M.V. (2020). Modulation of cortical slow oscillatory rhythm by GABAB receptors: an in vitro experimental and computational study. J Physiol. DOI: 10.1101/2019.12.14.866442v2.

Peters, A.J., Steinmetz, N.A., Harris, K.D., and Carandini, M. (2019). Striatal activity reflects cortical activity patterns. bioRxiv, 703710.

Planert, H., Berger, T.K., and Silberberg, G. (2013). Membrane properties of striatal direct and indirect pathway neurons in mouse and rat slices and their modulation by dopamine. PloS one *8*, e57054.

Planert, H., Szydlowski, S.N., Hjorth, J.J., Grillner, S., and Silberberg, G. (2010). Dynamics of synaptic transmission between fast-spiking interneurons and striatal projection neurons of the direct and indirect pathways. Journal of Neuroscience *30*, 3499-3507.

Rehman, N., and Mandic, D.P. (2009). Multivariate empirical mode decomposition. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences *466*, 1291-1302.

Reig, R., and Silberberg, G. (2014). Multisensory integration in the mouse striatum. Neuron *83*, 1200-1212.

Reig, R., and Silberberg, G. (2016). Distinct Corticostriatal and Intracortical Pathways Mediate Bilateral Sensory Responses in the Striatum. Cereb Cortex *26*, 4405-4415.

Reiner, A., Jiao, Y., Del Mar, N., Laverghetta, A.V., and Lei, W.L. (2003). Differential morphology of pyramidal tract-type and intratelencephalically projecting-type corticostriatal neurons and their intrastriatal terminals in rats. J Comp Neurol *457*, 420-440.

Rilling, G., Flandrin, P., and Goncalves, P. (2003). On empirical mode decomposition and its algorithms. Paper presented at: IEEE-EURASIP workshop on nonlinear signal and image processing (NSIP-03, Grado (I)).

Rock, C., Zurita, H., Wilson, C., and Apicella, A.J. (2016). An inhibitory corticostriatal pathway. Elife *5*.

Ros, H., Sachdev, R.N., Yu, Y., Sestan, N., and McCormick, D.A. (2009). Neocortical networks entrain neuronal circuits in cerebellar cortex. J Neurosci *29*, 10309-10320.

Ruiz-Mejias, M., Ciria-Suarez, L., Mattia, M., and Sanchez-Vives, M.V. (2011). Slow and fast rhythms generated in the cerebral cortex of the anesthetized mouse. Journal of neurophysiology *106*, 2910-2921.

Ruiz-Mejias, M., Martinez de Lagran, M., Mattia, M., Castano-Prat, P., Perez-Mendez, L., Ciria-Suarez, L., Gener, T., Sancristobal, B., Garcia-Ojalvo, J., Gruart, A.*, et al.* (2016). Overexpression of Dyrk1A, a Down Syndrome Candidate, Decreases Excitability and Impairs Gamma Oscillations in the Prefrontal Cortex. J Neurosci *36*, 3648-3659.

Sáez, M., Ketzef, M., Alegre-Cortés, J., Reig, R., and Silberberg, G. (2018). A New Micro-holder Device for Local Drug Delivery during In Vivo Whole-cell Recordings. Neuroscience *381*, 115-123.

Sanchez-Vives, M.V., Massimini, M., and Mattia, M. (2017). Shaping the Default Activity Pattern of the Cortical Network. Neuron *94*, 993-1001.

Sanchez-Vives, M.V., Mattia, M., Compte, A., Perez-Zabalza, M., Winograd, M., Descalzo, V.F., and Reig, R. (2010). Inhibitory modulation of cortical up states. Journal of neurophysiology *104*, 1314-1324.

Sanchez-Vives, M.V., and McCormick, D.A. (2000). Cellular and network mechanisms of rhythmic recurrent activity in neocortex. Nature neuroscience *3*, 1027.

Schultz, W., Dayan, P., and Montague, P.R. (1997). A neural substrate of prediction and reward. Science *275*, 1593-1599.

Shamir, M., and Sompolinsky, H. (2004). Nonlinear population codes. Neural Comput *16*, 1105-1136.

Sharott, A., Vinciati, F., Nakamura, K.C., and Magill, P.J. (2017). A Population of Indirect Pathway Striatal Projection Neurons Is Selectively Entrained to Parkinsonian Beta Oscillations. J Neurosci *37*, 9977-9998.

Sippy, T., Lapray, D., Crochet, S., and Petersen, C.C. (2015). Cell-Type-Specific Sensorimotor Processing in Striatal Projection Neurons during Goal-Directed Behavior. Neuron *88*, 298-305.

Steriade, M., Amzica, F., and Contreras, D. (1996). Synchronization of fast (30-40 Hz) spontaneous cortical rhythms during brain activation. J Neurosci *16*, 392-417.

Steriade, M., Contreras, D., Curro Dossi, R., and Nunez, A. (1993a). The slow (< 1 Hz) oscillation in reticular thalamic and thalamocortical neurons: scenario of sleep rhythm generation in interacting thalamic and neocortical networks. J Neurosci *13*, 3284-3299.

Steriade, M., Nunez, A., and Amzica, F. (1993b). A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. J Neurosci *13*, 3252-3265.

Szydlowski, S.N., Dorocic, I.P., Planert, H., Carlén, M., Meletis, K., and Silberberg, G. (2013). Target selectivity of feedforward inhibition by striatal fast-spiking interneurons. Journal of Neuroscience *33*, 1678-1683.

Tepper, J.M., Wilson, C.J., and Koós, T. (2008). Feedforward and feedback inhibition in neostriatal GABAergic spiny neurons. Brain research reviews *58*, 272-281.

Thorn, C.A., Atallah, H., Howe, M., and Graybiel, A.M. (2010). Differential dynamics of activity changes in dorsolateral and dorsomedial striatal loops during learning. Neuron *66*, 781-795.

Timofeev, I., Grenier, F., Bazhenov, M., Sejnowski, T., and Steriade, M. (2000). Origin of slow cortical oscillations in deafferented cortical slabs. Cerebral cortex *10*, 1185-1199.

Ur Rehman, N., and Mandic, D.P. (2011). Filter bank property of multivariate empirical mode decomposition. IEEE transactions on signal processing *59*, 2421-2426.

Vergara, R., Rick, C., Hernández-López, S., Laville, J.A., Guzman, J.N., Galarraga, E., Surmeier, D.J., Bargas, J. (2003). Spontaneous voltage oscillations in striatal projection neurons in a rat corticostriatal slice. J Physiol 553, 169-182.

Wall, N.R., De La Parra, M., Callaway, E.M., and Kreitzer, A.C. (2013). Differential innervation of direct- and indirect-pathway striatal projection neurons. Neuron *79*, 347-360.

Wilson, C.J. (1987). Morphology and synaptic connections of crossed corticostriatal neurons in the rat. J Comp Neurol *263*, 567-580.

Wilson, C.J. (1993). The generation of natural firing patterns in neostriatal neurons. Prog Brain Res *99*, 277-297.

Wilson, C.J., and Kawaguchi, Y. (1996). The origins of two-state spontaneous membrane potential fluctuations of neostriatal spiny neurons. Journal of neuroscience *16*, 2397-2410.

Wu, Z., and Huang, N.E. (2009). Ensemble empirical mode decomposition: a noise-assisted data analysis method. Advances in adaptive data analysis *1*, 1-41.

Yin, H.H., and Knowlton, B.J. (2006). The role of the basal ganglia in habit formation. Nat Rev Neurosci *7*, 464-476.

Figures

Figure 1. Analysis and classification of SWO in dorsal striatum. a, Schematic representation of the in vivo recording setup. b, Morphological reconstruction of DLS-MSN (left) and DMS-MSN (right). Different scales show neuron magnitude and its dendritic spines, confirming that the recorded neuron is a MSN. c, Representative LFP (top) and whole-cell patch-clamp recording of a MSN (bottom), together with SWO extraction by NA-MEMD (orange line). d, Featurization of the SWO of DLS- and DMS-MSNs [See methods]. e, RFE of the computed features (feature number) to determine their order of relevance classifying DLS- and DMS-MSNs. A SVM with a linear kernel was selected as classification algorithm. f, Example ROC curve of the DLS-DMS classification for one of the crossvalidations using the 3 most relevant features, illustrated in D. g, Schematic representation of the data distribution of DLS and DMS groups for the most relevant features (numbers 11, 5, and 9 respectively). White number represent the mean; the radius of the circle represents the variance. h, Subspace of classification based on the RFE results. Dots represent individual MSNs recorded in DLS (red) or DMS (blue). Classification hyperplane obtained after training the SVM with linear kernel in black.

Figure 2. Beta band in membrane voltage of dorsal striatal MSNs. **a,** Example of energy of the beta band (red trace) during the MSN SWO recorded in DMS. **b,** Energy of the beta band of DLS- and DMSMSNs. **c,** Average of phase alignment of beta band to the SWO in DLS- (left) and DMS-MSNs (right) (Raylegh test, p<0.001 in both cases). Asterisk indicates the radial position of the beta peak. Beta peak occurs first in DMS-MSNs (p<0.001). **d,** Representative examples of beta phase locking in DLS- (left) and DMS- MSNs (right). White line represents the beginning of each Up state.

Figure 3. Study of the SWO in the DCS striatum and determination of the functional boundary between DLS and DMS. **a,** Schematic representation of the "boundary question". **b,** Three possible hypotheses about DCS-MSNs SWO. Hyp. 1: There is an intermediate distribution between DLS and DMS SWO (top); Hyp. 2: There is a different type of SWO distribution (middle); Hyp. 3: There is no specific DCS distribution of SWO and the MSNs recorded at DCS are a combination of DLS- and DMS-MSNs (bottom). The black line displays the distribution of the SWO combining both DLS and DMS. **c,** Distribution of DLS-, DMS- and DCS-MSNs in the subspace of classification determined by the RFE. Classification plane in black. Orthogonal hypervector to the classification plane in grey. **d,** Distribution of dorsolateral, dorsomedial and dorsocentral MSNs and the combining DLS-DMS function along the orthogonal hypervector to the hyperplane classification. All comparison between distributions are significant (p<0.01) except for DCS with DMS-DLS (p= 0.681).

Figure 4. Number of peaks and depolarized/hyperpolarized ratio of the SWO in dorsal striatum. **a,** Detection of peaks (black arrows) in the Up state in a DLS-MSN (red) and a DLS-MSN (blue) [See methods]. **b,** Representative examples of SWO in different cortical regions recorded. **c,** Number of peaks per Up state in the different cortical regions. Non-labelled comparisons are not significant. **d,** Extraction of depolarized/hyperpolarized events by computational approach [see methods]. Top: Detection of the events using a threshold in the first derivative of the Vm. Scale bar represents 0.1 dmV/dt. Bottom: Representation of the detected depolarized (green) and hyperpolarized (orange) events from whole-cell recording of a DLS-MSN (middle). **e,** Average of the depolarized/ hyperpolarized ratio aligned to the SWO cycle of DMS- and DLS-MSNs. Comparison of positive (p<0.0072) and negative (p<0.0398) values at peaks.

Figure 5. Integration of the cortical SWO in dorsal striatum. **a,** Example of simultaneous recordings from a DLS-MSN (red) and double LFPs from FrA (purple) and S1 (black). **b,** Example of simultaneous recordings from a DMS-MSN (blue) and double LFP from M1 (green) and V1 (ochre). **c,** Histogram of the membrane potential values of the MSNs in D and E, the back arrow indicates the dissimilar shape of the bimodal distributions. **d,** Correlation coefficient of different cortical regions to DLS- and DMS-MSNs. **e,** Raster plot showing the correlation between all MSNs and the LFP from FrA and V1. **f,** Probability of propagation of the Up state from different cortical regions to DLS- and DMS-MSNs.

Figure 6. Sequential activation of the dorsal striatum during the SWO. **a,** Representative example of simultaneous double *in vivo* whole-cell recordings in DLS (top, red) and DMS (bottom, blue). **b,** Inset of the shaded part in A, showing two aligned voltage traces to their Down states. Notice that DLS-MSN onset is preceding the DMS-MSN. **c,** Raster plot of the same example of a paired recording in DLS (top) and DMS (bottom) MSNs. Each line represents a 1 second time window aligned to the onset of each of the Up states of the DLS-MSN (white line). **d,** Example of the distribution of delays between the onset of the Up state in the DMS- relative to the DLS-MSN. Positive values indicate that the Up state arrive later to the DMS-MSN. Same neuron in A, B, C and D. **e,** Directional probability of DS-MSNs.

Figure 7. Differences between direct and indirect pathway MSNs in dorsal striatum. **a,** Example showing an *in vivo* identification of a MSN using the optopatcher. Responses in D2-ChR2-YFP mice (top trace, ChR2$^+$, green) to light pulses, inducing depolarization in the MSN. Negative cells (bottom trace, ChR2$^-$, black) did not respond to light pulses. Blue squares indicate the intensity of the light pulse stimulation, from 20 to 100 %. **b,** RFE to optimize the classification of dMSNs and iMSNs in DLS (red) or DMS (blue). A SVM with a linear kernel was selected as classification algorithm. **c,** Significant differences in three SWO features were found in DLS (red) between direct (dark) and indirect (light) pathways, but not in DMS (blue).

	Input Resistance (MΩ)	Resistance Down state hyp. (MΩ)	Resistance Down state dep. (MΩ)	Resistance Up state hyp. (MΩ)	Resistance Up state dep. (MΩ)	Capacitance (pF)	Tau (ms)
DLS	**310± 25 ***	**291± 23 ***	**309± 23 ***	304± 23	312± 27	**26.65± 7.56 ***	4.57± 0.23
dMSNs	291± 33	276± 31	279± 28	290± 29	292± 35	32.78± 12.81	4.52± 0.29
iMSNs	**338± 41 #**	**312± 36 #**	**355± 40 #**	324± 37	342± 41	17.58± 2.35	4.63± 0.40
DMS	**255± 26 ***	**241± 25 ***	**259± 27 ***	262± 27	262± 27	**21.68± 2.26 ***	4.61± 0.48
dMSNs	260± 22	249± 28	265± 23	269± 28	267± 23	23.20± 2.21	4.77± 0.19
iMSNs	**241± 18 #**	**223± 18 #**	**248± 16 #**	248± 20	252± 20	18.43± 1.04	4.26± 0.24

Table 1. Intrinsic properties of DLS and DMS of direct and indirect MSNs. Comparisons between DLS- and DMS-MSNs ($p<0.05$, *symbol). Comparisons between direct and indirect MSNs ($p<0.01$, # symbol). All values are means ± standard deviation.

Medium spiny neurons spontaneous activity reveal the discrete segregation of mouse dorsal striatum

J. Alegre-Cortés, M. Sáez, R. Montanari, R. Reig*

Supplementary Information

Figure S1. ROC curve for the classification using one-animal-out strategy.
We independently classified the MSNs of each animal training the classifier using the rest of the dataset to exclude any possible inter-animal bias during training.

Figure S2. Theta and Gamma band in membrane voltage of dorsal striatum.

a, Representative example of the energy of the theta band (orange trace) of a DLS-MSN during SWO regime. **b,** Representative example of the energy of the Gamma band (red trace) of a DLS-MSN during SWO regime. **c,** Energy of the Theta band of DLS- and DMS-MSNs. **d,** Energy of the Gamma band of DLS and DMS-MSNs. Same neuron in A and B.

Figure S3. Z-scoring of the SWO features.

Z-score transformation of the computed features of the SWO that were used for the classification of DLS- and DMS-MSNs in figure 3A. P values are the same as in figure 3A.

Figure S4. Beta, Theta and Gamma band in membrane voltage of direct and indirect pathways in the DLS and DMS. Energy of the Beta, Theta and Gamma band of direct and indirect DLS- and DMS-MSNs.

Figure S5. Example of decomposition of different traces by the NA-MEMD.

a, Example traces of a S1 LFP (left), M1 LFP (center) and a whole-cell recording in DLS (right). **b,** Consecutive IMFs obtained by applying NA-MEMD to the signals in a, starting from the 7th IMF. The IMF classifies the oscillatory activity from the fastest to the slowest.

Tables

Feature	Mean value
1- Average voltage in Up state (mV)	-54.44 ± 6.28
2- Up state standard deviation (mV)	5.32 ± 1.36
3- Min voltage in Up state (mV)	-59.95 ± 6.23
4- Max voltage in Up state (mV)	-43.99 ± 6.20
5- Up state amplitude (mV)	13.62 ± 4.10
6- Max of Up state derivative (dmV/dt)	0.65 ± 0.14
7- Min of Up state derivative (dmV/dt)	-0.512 ± 0.12
8- Down to Up transition slope (mV/s x 10^2)	0.829 ± 0.25
9- Up to Down transition slope (mV/s x 10^2)	0.514 ± 0.13
10- Slope transition ratio (arbitrary units)	1.61 ± 0.36
11- Number of peaks in Up state (units)	1.20 ± 0.15
12- Peak to peak in Up state (mV)	20.74 ± 4.65
13- Up state length (ms)	431.40 ± 71.50

Table S1. Featurization of the SWO of the DCS-MSNs.

The number preceding the feature labels are the same as the ones showed in figure 3. All values display means ± standard deviation.

Animals	45
Males	19
Females	26
Weight (gr)	33.54±1.37
Male weight (gr)	30.04±1.58
Female weight (gr)	37.58±1.94
Age (weeks)	24.66±1.31
Male age (weeks)	22.52±1.79
Female age (weeks)	27.26±1.84
Cells/animal	4.38±0.49
Cells/male	4±0.92
Cells/female	4.69±0.39

Table S2. Data set description.

The three first rows show total values, the rest show means ± standard deviation.