Perturbation of GABA Biosynthesis Links Cell Cycle to Control *Arabidopsis thaliana* Leaf Development

Yaxin Gong, Han Yue, Yu Xiang, Guanghui Yu*

Hubei Provincial Key Laboratory for Protection and Application of Special Plants in Wuling Area of China, Engineering Research Centre for the Protection and Utilization of Bioresource in Ethnic Area of Southern China, College of Life Sciences, South-Central University for Nationalities, Wuhan 430074, China

***Correspondence:**

Guanghui Yu

yuguanghui@scuec.edu.com

Keywords: *Arabidopsis thaliana*; CDKA;1; endoreplication; GABA; Reactive Oxygen Species

Running title: Endoreplication response to GABA deficit

Abstract

To investigate the molecular mechanism underlying increasing leaf area in γ-Aminobutyric acid (GABA) biosynthetic mutants, the first pair of true leaves of GABA biosynthetic mutants was measured. The results showed that the leaf blade area in GABA biosynthetic mutants was larger than that of the wild type to different extents, and the area of the leaf epidermal cells in mutants was larger than that of the wild type. DNA polyploid analysis showed that polyploid cells in GABA biosynthetic mutants were appearing earlier and more abundant than in the wild type. To check the correlation between cell size and endoreplication, the expression of factors involving endocycles, including D-type cyclin gene (*CYCD3;1*, *CYCD3;2*, *CYCD3;3*, and *CYCD4;1*) and kinase *CKDA;1*, were analysed by qRT-PCR. The results showed that *CKDA;1* in GABA biosynthetic mutants was downregulated, and four types of *CYCDs* showed different expression patterns in different GABA biosynthetic mutants. Inconsistent with this result, for *CCS52A* (*CELL CYCLE SWITCH 52A*) (controlling the endocycle entry) in *gad2* and *gad1/gad2* mutants, the expression of *CCS52A2* was significantly higher than that in the wild type. The expression of *SIM* (*SIAMESE*) and *SMR* (*SIAMESE-RELATED*), which inhibit kinase activity, were also upregulated compared with the control. To further study the possible potential relationship between GABA metabolism and endoreplication, we analysed the reactive oxygen species (ROS) levels in guard cells using ROS fluorescent probes. ROS levels were significantly higher in GABA biosynthetic mutants than the control. All results indicated that cyclin, the cyclin-dependent kinase, and its inhibitory protein were

coordinated to participate in endoreplication control at the transcription level in the
leaves of GABA biosynthetic mutant *Arabidopsis*.

Contribution to the field statement

γ-Aminobutyric acid (GABA) metabolic pathway plays a dual role in plant
development. This research investigated the perturbation of GABA biosynthesis on
Arabidopsis leave endoreplication for the first time. In the GABA biosynthetic
mutants, many genes, participating in cell division regulation, are coordinately
transcriptionally expressed to trigger the onset and maintenance of endoreplication,
and this led to the cell expansion and the increase leaf blade area. However, this
initiation of endoreplication links with the decrease of endogenous GABA level and
the increase Reactive oxygen species (ROS). This may be a compensation mechanism
to adapt to abnormal GABA level in plant leaf development. Present evidence
provided hypothesized that the normal GABA level in plant leaf development plays a
brake to inhibit the immature cell expansion and differentiation, and this negative
regulation functions a guarantee mechanism to watchdog the normal leaf development.
In all, this contribution provides an updated perspective on the role of GABA in plant
development.

1   Introduction

γ-Aminobutyric acid (GABA) is a four-C, non-protein component amino acid
commonly found in organisms and is prevalent in bacteria, plants, and vertebrates
(Seifikalhor et al., 2019). The precursor of GABA synthesis is L-glutamic acid (Glu),
which is catalysed by glutamate decarboxylase (GAD) in cytoplasm. The *Arabidopsis*
genome has five genes encoding GAD named GAD1–5. The oxidative metabolism of
GABA occurs in mitochondria and entry is mediated by a GABA permease (Michaeli
et al., 2011). In mitochondria, using α-ketoglutarate or pyruvate as an amino acceptor,
GABA is catalysed by GABA transaminase (GABA-T) to produce glutamic acid (Glu)
or Alanine (Ala) and succinic semialdehyde (SSA), respectively. SSA is oxidized by
SSA dehydrogenase (SSADH) to succinic acid (SucA), and then SucA enters the
tricarboxylic acid (TCA) cycle for further metabolism. This metabolic pathway is
referred to as the GABA shunt. Under hypoxic or high-light conditions, SSA can be
reduced to gamma-hydroxybutyrate (GHB) by SSA reductase (SSR, also known as
GHB dehydrogenase) in the cytoplasm, mitochondria, and chloroplasts (Allan et al.,
2008). Previous studies have reported that unique cytosolic and plastid glyoxylate
reductase isoforms in *Arabidopsis* are known as GLYR1 and atGLYR2, respectively,
and they catalyse the conversion of SSA to GHB and glyoxylic acid to glycolic acid
via an NADPH-dependent reaction (Brikis et al., 2017). The balance of the redox state
was maintained by the accumulation of GHB and the reduction of SSA via the GABA
shunt.

GABA biosynthesis and accumulation from the glutamic acid pathway and its
metabolism are at the junction of N and C metabolism, providing a useful metabolic
substrate for the TCA cycle, electron transport chain, and C skeleton, which are

involved in the balancing of C:N metabolism (Jacoby et al., 2011). It is generally
believed that GABA metabolism is considered to be involved in metabolic signalling,
which plays a dual role in plant development, including both metabolic and signal
regulation (Häusler et al., 2014; Ramesh et al., 2017; Podlešáková et al., 2019). In
addition, aldehyde chemical groups (i.e. H–C=O) produced by GABA biosynthesis
and metabolic pathways in plants have high molecular activity, and aldehydes
accumulated under stressed conditions are highly toxic and can react with DNA, lipids
of oxidative membranes, and modified proteins, or affect the transcription of
stress-related genes, leading to cellular and ontogenetic problems in plants. Previous
studies have reported that GABA metabolism regulates leaf pattern morphogenesis.
The SSADH gene is involved in the formation of the paraxial–abaxial (upper–lower)
leaf model in *Arabidopsis thaliana* (Toyokura et al., 2011; 2012). Mutation of *enf1*
(*enlarged fil expression domain1*) alters the expression pattern of the *FIL* (*YABBY1*,
FILAMENTOUS FLOWER) gene, which is characteristic of meristem and organs in *A.*
thaliana (Sawa et al., 1999), on the abaxial surface of leaf primordium (Sawa et al.,
1999; Siegfried et al., 1999). However, here is a dearth of research on the role of
GABA biosynthesis in the leaf development of *A. thaliana*. In the present study,
GABA biosynthetic mutants, *gad1*, *gad2*, and *gad1/gad2* were examined to explore
the molecular mechanism underlying the GABA negative feedback that regulates leaf
cell endoreplication during leaf development. Our findings will provide evidence for
further understanding the role of GABA in plant development.

2 Materials and methods

2.1 Experimental Materials

A. thaliana Col wild-type seeds, Col ecotype *gad1* mutant, *gad2* mutant, and
gad1/gad2 double mutant were provided by Prof. Barry Shelp (University of Guelph,
Canada). The seeds of the wild type and mutants were sterilized for ~2 h in a sealed
container with chlorine gas, and then inoculated on MS solid medium and
synchronized for 3 d at 4 °C. After synchronization, the plate was taken out and
placed in greenhouse under a photoperiod of 16/8 h light/dark and incubated at 22 °C.

2.2 Measurement of leaf blade area

From the 4th day after *A. thaliana* seedlings being transferred to the greenhouse, their
growth condition was photographed every 24 h with a Brinno time-lapse camera
(TLC100) at a fixed distance to record the leaf blade area.

2.3 Microscopic observation and measurement of cell area

From the 7th day after *A. thaliana* seedlings being transferred to the greenhouse, the first
pair of true leaves was collected daily, the leaf abaxial epidermis located 25% and 75%
from the distance between the tip and the base of the leaf blade was photocopied with
nail polish, and then photographed using an Olympus DP80 microscope, and the area
was calculated with the software that came with the microscope. Under the microscope,
a certain area of the epidermis was confined to count the number of cells, and then the
average cell area was calculated.

2.4 DNA ploidy analysis

Approximately two or three leaves at the young stage or one or two leaves at the
middle and late leaf developmental stages were chopped with a razor blade in nuclei
extraction buffer (CyStain® UV Precise P, Sysmex Partec), and then transferred to the
staining buffer (CyStain® UV Precise P, Sysmex Partec) according to the
manufacturer's instructions. The ploidy level of DNA in leaf cells was determined
using a CyFlow Ploidy Analyser (Sysmex Partec).

2.5   qRT-PCR analysis

Total RNA from *A. thaliana* leaves was extracted by AxyPrep Multisource Total
RNA Miniprep Kit (Axygen Science, Inc). Reverse transcription was performed by
Goldenstar RT6 Gene Synthesis Kit (Tingke Biotechnology Co., Ltd.) with a reverse
transcription reaction system of 20 μL. qRT-PCR analysis was performed by MyGo
Pro qPCR System (IT-IS Life Science Ltd.). The gene *isoprenyl diphosphate delta*
isomerase II (*IPP2*, AT3G02780) was selected as the internal reference gene
according to previous research (Fung-Uceda et al., 2018). The primers of the genes
involved in cell cycle regulation, endocycle initiation, progression, and exit are listed
in the supporting materials (Table S1).

2.6   *In vivo* reactive oxygen species imaging

For GC reactive oxygen species (ROS) staining, H2DCF-DA stock solution (10
mmol/L stock in Dimethyl Sulfoxide) was diluted in deionized water to yield a final
concentration of 6.25 μmol/L with a final dimethyl sulfoxide concentration of
0.0125% (v/v) (Watkins et al., 2017). Epidermal strips were peeled with adhesive tape
and directly stained for 30 min in the above solution. After rinsing with deionized
water for 5 min, images were taken by Olympus DP80 fluorescent microscopy. The
intensity of coloration was quantified using ImageJ software (National Institutes of
Health, USA).

2.7   Statistical analyses

Data are expressed as averages ± standard error (SE). Experiments were conducted
with two or three independent replicates. One-way ANOVA was employed using
SPSS 20.0.

3   Results

3.1   The leaf blade area of the GABA biosynthetic mutant was larger than that
of the wild type

At the early stage of leaf development (4th day after stratification), there was no
significant difference in the area of the first pair of true leaves between the wild-type
and GABA biosynthetic mutants (Figure 1). As time progressed, the area of the
mutant leaves increased more rapidly than that of the wild type. Around the 8th day,
the leaf area of the first pair of true leaves in the wild type was about 1.8 mm^2, and the
first pair of true leaves in *gad1* and *gad2* mutants was about 2.4 mm^2, at 1.3-fold that
of the wild type, and that of the *gad1/gad2* mutants was about 3.2 mm^2 with 1.8-fold
that of the wild type (Figure 1). The leaf area between the GABA biosynthetic
mutants and wild type was significantly different ($P < 0.05$).

**3.2 Cell size in GABA biosynthetic mutants was larger than those of the wild
type**

To determine the reasons for the leaf area in GABA biosynthetic mutants being larger
than that in the wild type, we compared the area of leaf epidermal cells. The average
cell area in the epidermis of the first pair of true leaves of the wild type was about
2700 μm^2 at the 8^{th} day after stratification (Figure 2). The average cell area of the
epidermis of the first pair of leaves of *gad1* and *gad2* mutants was ~3200 μm^2,
1.2-fold that of the wild type (P <0.05). The average cell area of the epidermis of the
first pair of true leaves in *gad1/gad2* mutants was ~3800 μm^2, and was 1.4-fold that of
the wild type with significant differences (P <0.05) (Figure 2).

**3.3 The polyploidy in GABA biosynthetic mutant leaf occurred earlier and
higher than that of the wild type**

The correlation between cell size and DNA ploidy level (Matsunaga et al., 2013)
prompted us to explore whether the cell size of GABA biosynthetic mutants is related
to DNA polyploidy level. On the 7^{th} day of leaf development after stratification, the
cells in the first pair of true leaves in wild-type *A. thaliana* were mostly diploid and
tetraploid, indicating that leaf cells mainly divided at this stage (Figure 3A). However,
leaf cells in *gad1*, *gad2*, and *gad1/gad2* mutants appeared to be in different
proportions of 8-ploid cells except the diploid and tetraploid cells (Figure 3B). The
appearance of 8-ploid cells in leaves is a marker of endoreplication. On the 8^{th} day of
leaf development, 8-ploid cells were not found in wild-type *A. thaliana*, and the
proportion of octoploid cells in three kind of GABA-biosynthetic mutants increased
further (Figure 3B).

To further confirm the relationship between cell enlargement and cell ploidy level, we
compared the cell ploidy level in the middle and late leaf development stages (21^{st} day
after stratification). The proportion of octoploid cells in the wild type was about 18%
(Figure 3C). However, the proportion of the octoploid cells in *gad1*, *gad2*, and
gad1/gad2 mutants was about 30–40%, which was significantly higher than that in the
wild type (Figure 3C).

Overall, at the early stage of leaf development (7–8^{th} day after stratification), the
8-ploid cells were only detected in the GABA biosynthetic mutants (8-ploid cells in
gad1 and *gad2* mutants was 3–5%, and that in *gad1/gad2* double mutants was about
5–9%) (Figure 3D). At the late stage of leaf development (21^{st} day after stratification),
the proportion of 8-ploid cells in wild-type leaves was also detected; however, its
level was significantly lower than that in *gad* mutants (Figure 3D).

**3.4 Type-D cyclin genes involved in endocycle regulation in GABA biosynthetic
mutants**

Considering that endoreplication is a special form of cell division (De Veylder et al.,
2011) and is the reason for cell polyploidy (Matsunaga et al., 2013), type-D relative
gene participation in cell cycle regulation was analysed by qRT-PCR.

CYCD3;1 (At4g34160) is a key component to balance cell proliferation/division and
endoreplication (Dewitte et al., 2007). Downregulation of *CYCD3;1* can lead to

endoreplication, and its overexpression can cause excessive cell proliferation and
inhibit cell differentiation (Dewitte et al., 2003). qRT-PCR confirmed that the
expression of *CYCD3;1* was significantly downregulated in GABA biosynthetic
mutants (Figure 4). In *gad1* and *gad2* single mutants, the relative expression of
CYCD3;1 was about 0.2-fold that of the wild type ($P < 0.01$), and in *gad1/gad2*
double mutants, the relative expression of *CYCD3;1* was about 0.1-fold that of the
wild type ($P < 0.01$) (Figure 4), indicating that blocking GABA synthesis inhibited the
expression of *CYCD3;1*.

CYCD3;2 (At5g67260) and CYCD3;3 (At3g50070) participate in the symmetrical
division of guard mother cells and guard cells (GCs) in the late stomatal lineage
development (Yang et al., 2014). In the *gad1* mutant, the expression of *CYCD3;2* was
significantly downregulated ($P < 0.05$); in the *gad2* and *gad1/gad2* mutants, the
relative expression of *CYCD3;2* was about 0.6-fold that of the wild type with
significant differences ($P < 0.01$) (Figure 4). Similarly, the expression of *CYCD3;3* in
the *gad1* mutant was about 0.3-fold that of the wild type ($P < 0.01$), and in the *gad2*
and *gad1/gad2* mutants, the expression of *CYCD3;3* was 0.7 and 0.5-fold that of the
wild type, respectively ($P < 0.05$) (Figure 4). CYCD4;1 (At5g65420) could activate
the cell cycle of root apical meristem (Masubelele et al., 2005), and the expression of
CYCD*4;1* in the *gad2* mutant was significantly downregulated ($P < 0.05$), and in the
gad1 and *gad1/gad2* mutants the relative expression of *CYCD4;1* decreased to
become the most significantly different to that of the wild type ($P < 0.01$) (Figure 4).

3.5   The expression of *CDKA;1* in GABA biosynthetic mutants was significantly downregulated

CDKA;1 (At3g48750) is one of the core components of cell cycle regulation, mainly
controlling the transition of the G1/S and G2/M mitotic phase (Nowack et al., 2012).
Inhibition of its activity can block mitosis to initiate leaf cell endoreplication (Verkest
et al., 2005). The expression of *CDKA;1* in GABA biosynthetic mutants was
significantly lower than that in the wild type ($P < 0.05$). These results indicated that
blocking GABA synthesis is linked with the decreased expression of *CDKA;1* (Figure
5).

The transcription factor E2Fa (At2g36010) stimulates cell proliferation and delayed
differentiation (Boudolf et al., 2004). In the *gad1* and *gad2* mutants, the expression of
E2Fa was significantly downregulated ($P < 0.05$). In *gad1/gad2*, the level of *E2Fa*
was 0.2-fold that in the wild type ($P < 0.01$) (Figure 5).

3.6   Different expression patterns of *CCS52A2* and *CDC6* were observed in GABA synthetic mutants

CCS52A2 (*CELL CYCLE SWITCH52*, At4g11920) encodes a ubiquitin ligase
regulating cell cycle division phase (M) and a substrate-specific activator of anaphase
promotion complex/cyclosome (Fülöp et al., 2005). Inhibition of kinase activity after
CCS52A2 expression and enhancement of its activity are prerequisites for
endoreplication (Heyman et al., 2017; Umeda et al., 2019). Mutations in the *CCS52*
gene resulted in delayed endoreplication, and its overexpression resulted in increased
DNA ploidy levels (Heyman et al., 2017). In the *gad1* mutant, the relative expression
of *CCS52A2* was not different from that of the wild type; however, in the *gad2* and

gad1/gad2 mutants, the expression of *CCS52A2* was significantly higher than that of
the wild type (P <0.05) (Figure 6).

CDC6 (At2g29680) encodes a homolog of cell division regulatory protein 6
(Castellano et al., 2001), which is a license gene for DNA replication (Fung-Uceda et
al., 2018). CDC6 participates in the initiation of DNA replication and is an important
factor for maintaining endoreplication. The ectopic expression of *CDC6* can increase
the ploidy level caused by endoreplication (Castellano et al., 2001; 2004). In the *gad1*
mutant, the relative expression of *CDC6* was 1.7-fold as much as that in the wild type
(P <0.05) (Figure 6). The expression of *CDC6* in the *gad2* mutant was about 3-fold
that in the wild type (P <0.01), and the expression of *CDC6* in the *gad1/gad2* double
mutant was about 8-fold that in the wild type (P <0.01) (Figure 6).

3.7 **The expression of *SIAMESE* and *SIAMESE-RELATED* in GABA**
biosynthetic mutants was significantly up-regulated

SIAMESE (SIM, At5g04470) is an inhibitory protein of cyclin-dependent kinase and
a regulator of mitotic inhibition and endoreplication (Churchman et al., 2006).
Overexpression of *SIM* results in dwarfing of plants, serrated leaves, and cells with
higher nuclear DNA content (Churchman et al., 2006). The relative expression of the
SIM in *gad1* single mutant is about twice that of the wild type (P <0.05), in the *gad2*
single mutant is about 4-fold that of the wild type (P <0.05), and in the *gad1/gad2*
double mutant is about 7-fold that of the wild type (P <0.01) (Figure 7).

Most components of the *SIAMESE-RELATED* (*SMR*) gene family function in mitosis
inhibition and endocycle promotion (Yi et al., 2014; Kumar et al., 2015; Dubois et al.,
2018). In the chosen genes, the expression of *SMR1* (At3g10525) in *gad1* and *gad2*
mutants was not different to that of the wild type. In the *gad1/gad2* mutants, its
expression is 1.5-fold that of the wild type (P <0.05) (Figure 7). However, the
expression of other components, *SMR2* (At1g08180), *SMR5* (At1g07500), and *SMR8*,
was markedly different to that of the wild type (Figure 7). The expression of *SMR2* in
gad1, *gad2*, and *gad1/2* reached 5-, 6-, and 11-fold of that in the wild type,
respectively (P <0.01). Similarly, the expression of *SMR5* in *gad1*, *gad2*, and *gad1/2*
reached 3-, 5- (P <0.05), and 17-fold that in the wild type (P <0.01), respectively, and
the expression of *SMR8* in *gad1*, *gad2*, and *gad1/2* reached 4-, 4- (P <0.05), and
6-fold that in the wild type (P <0.01), respectively.

3.8 The level of ROS in leaves of GABA biosynthetic mutant was higher than
that of the wild type

The fact that redox regulates cell proliferation and the cell cycle (Schippers et al.,
2016), and GABA could scavenge ROS (Liu et al., 2011; Seifikalhor et al., 2019), led
us to postulate that ROS levels in GABA biosynthetic mutants could be increased.
Imaging of H2DCFDA (DCFH-DA; 2', 7'-Dichlorodihydrofluorescein diacetate) with
the fluorescent probe of ROS showed that the average ROS level intensity in GCs of
gad mutants was higher than that in the wild type (P <0.05) (Figure 8). These results
suggest that GABA synthetic mutants are indeed correlated with ROS accumulation.

4 Discussion

The GABA metabolic pathway plays a dual role during plant development, including
metabolic and signal regulation (Bown & Shelp, 2016; Seifikalhor et al., 2019).
Abnormal GABA biosynthesis and metabolism have important effects on plant
development (Baum et al., 1996; Palanivelu et al., 2003; Renault et al., 2013). The
expression of the *Petunia* GAD gene, which lacked the calmodulin-binding domain in
transgenic tobacco, resulted in abnormal plant development, which was shorter and
more branched than that of normal plants (Baum et al., 1996). In the Arabidopsis
GABA-T (*pop2-1*) mutant, the growth of pollen tubes could not accurately target the
ovule sac in the pistils (Palanivelu et al., 2003). The molecular mechanism was related
to the destruction of the gradient distribution of GABA in the stigma of the pistils
(Palanivelu et al., 2003). Furthermore, the growth retardation in hypocotyl epidermal
cells and root cortex cells is related to the limitation of cell elongation (Renault et al.,
2011), and inadequate expression of *GABA-T* may lead to developmental defects in
roots and hypocotyls and composition change in cell walls (Renault et al., 2013).

In the present study, the molecular mechanism underlying leaf-area enlargement in
GABA biosynthetic mutants was investigated. The growth of plant leaves is largely
limited by the development of the epidermis. Polyploidy in pavement cells is strongly
correlated with cell size (P <0.01) (Melaragno et al., 1993). We confirmed that the
larger leaf area and cell size in GABA biosynthetic mutants (Figures 1, 2) was related
to much higher percentages of polyploid cells in GABA biosynthetic mutant leaves,
which occurred earlier and in higher abundance than that of the wild-type leaf cells
(Figure 3).

Endoreplication is a special kind of cell division (De Veylder et al., 2011), and the
essence of endoreplication is to maintain CDK activity below the threshold that
triggers mitosis (De Veylder et al., 2011). During leaf development of *A. thaliana*, the
decrease in transcription levels of mitotic CDK and cyclin genes resulted in
endoreplication (Beemster et al., 2006). Consistent with these results, the qRT-PCR
analysis confirmed that the type-D cyclin genes (*CYCD3;1*, *CYCD3;2*, *CYCD3;3*, and
CYCD4;1) were differently downregulated in GABA biosynthetic mutants, and the
expression of *CDKA;1* exhibited a decreasing expression trend (Figures 4, 5).

Other factors interrelated with endoreplication regulation, from endoreplication
initiation, progression, and maintenance, and exit (Breuer et al., 2014), synergistically
regulate the occurrence of endoreplication in the leaves of GABA biosynthetic
mutants. *CCS52A* (*CELL CYCLE SWITCH 52A*) plays an important role in cell cycle
exit and endoreplication entry (Lammens et al., 2009; Vlieghe et al., 2005). The
concentration of *CCS52A* in mitotic cells remains below a critical threshold to prevent
immaturely initiating endoreplication (Lammens et al., 2009; Vlieghe et al., 2005). In
the *gad2* and *gad1/gad2* mutants, the relative expression of *CCS52A2* was
significantly higher than that of the wild type (Figure 6). The expression of
cyclin-dependent kinase inhibitor *SIM* (*SIAMESE*) and its *SMR* (*SIAMESE-RELATED*)
was significantly higher than that of the control (Figure 7). These results demonstrated
that the endoreplication regulators, including cyclin, cyclin-dependent kinase, and
kinase inhibitor, at the transcriptional level, all harmoniously contributed to the
regulation of endoreplication in mutants.

It is worth noting that *SMR5*, a member of *SMR* (*SIAMESE-RELATED*), is an
ROS-induced gene, which is more highly expressed in GABA biosynthetic mutants

(3–17-fold that of the control) (Figure 7). ROS is an important signalling molecule
regulating leaf development and is involved in triggering endoreplication (Schippers
et al., 2016). Previous evidence proved that exogenous GABA could scavenge ROS
(Liu et al., 2011), and the GABA level in mutants was much lower than that in the
wild type (Figure S1); thus, we speculated that the perturbation of GABA biosynthesis
may be intrinsically linked with the ROS level. Detection using an ROS fluorescence
probe confirmed that the level of ROS in GCs of GABA biosynthesis mutants was
significantly higher than that in the control (Figure 8). Therefore, we speculated that
blocking the GABA metabolic shunt pathway will lead to the accumulation of ROS
intermediates (Bouché et al., 2003; Fait et al., 2005), which may trigger
endoreplication (Figure 9). In this hypothesis, normal GABA metabolism functioned
as a signal and antioxidant to effectively inhibit the production of ROS. In the GABA
biosynthetic mutants, this inhibition was relieved. The production of ROS promotes
the expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*) expression, of
which, their encoding products inhibited kinase activity to initiate endoreplication
(Figure 9). In contrast, normal GABA metabolism postponed endoreplication, and this
negative regulation functions to prevent premature cell differentiation and to ensure
normal leaf development. However, this conclusion must be made cautiously because,
based on current data, it is difficult to demonstrate a causal relationship between
GABA level and ROS in terms of triggering endoreplication during the development
of Arabidopsis leaves.

Overall, the results of the present study demonstrate that the increase in polyploidy
level in GABA biosynthetic mutant leaves is achieved through endoreplication, which
is reflected in the increase in average cell size (Breuer et al., 2010). In this regulation,
multiple regulators are involved in the initiation, maintenance, and exit of
endoreplication. However, the premature termination of mitosis and the immature
occurrence of endoreplication are perhaps a compensatory mechanism for leaf
immature development, which resulted in increased cell ploidy. This mechanism may
be mediated by ROS signalling, but the complex regulatory mechanism requires
further research.

5   Conclusion

Using GABA biosynthetic mutants, the present study focused on the perturbation of
GABA biosynthesis on endoreplication in Arabidopsis leaf development. The
endoreplication cells that occurred in mutants were earlier and of higher abundance
than those in the wild type. This is the reason to lead to the increase in cell size and
leaf blade area. For transcription-level regulation, qRT-PCR confirmed that many
genes involved in cell cycle regulation were synergistically participating in the
initiation, progression, and maintenance of endoreplication. Among the regulators,
SMR5, encoding protein inhibited CDK activity, was markedly upregulated in GABA
biosynthetic mutants. Owing to the perturbation of GABA biosynthesis, the content of
ROS increased in GABA biosynthetic mutants, which is a potential signal to trigger
endoreplication in mutant leaves. Present evidence indicates that normal GABA
metabolism inhibited endoreplication to prevent immature cell differentiation in leaf
development. This research provides a deeper understanding of the role of GABA in
plant development.

6   Acknowledgments

We apologize if we inadvertently omitted citations of major contributions to this area.
We thank Prof. Barry Shelp (University of Guelph, Canada) for generously providing
the mutants seeds. We would like to thank Editage (www.editage.cn) for English
language editing.

7 Author Contributions

YXG studied the root phenotype in relation to GABA metabolism and wrote the draft
of the manuscript. HY and YX studied the role of ROS in cell-cycle regulation. GHY
studied the role of small molecules in plant development and revised the manuscript.

8 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial
or financial relationships that could be construed as a potential conflict of interest.

9 Funding

This work was supported by the Special Fund for Basic Scientific Research of Central
Colleges, South-Central University for Nationalities (CZP17051), National Natural
Science Foundation of China (31270361), studies on the regulation mechanisms of
agronomic traits in important crops and *A. thaliana* supported by State Administration
of Foreign Experts Affairs in Ministry of Science and Technology of the People's
Republic of China (P193009007) and Fund for Key Laboratory Construction of Hubei
Province (Grant No.2018BFC360).

10 Data Availability Statement

No datasets were generated for this study.

11 References

Allan WL, Simpson JP, Clark SM, Shelp BJ. Gamma-hydroxybutyrate accumulation
in Arabidopsis and tobacco plants is a general response to abiotic stress: putative
regulation by redox balance and glyoxylate reductase isoforms. J Exp Bot 2008;
59: 2555–64.

Baum, G., S Lev-Yadun, S., Fridmann, Y., Arazi, T., Katsnelson, H., Zik, M., and
Fromm, H. (1996). Calmodulin binding to glutamate decarboxylase is required
for regulation of glutamate and GABA metabolism and normal development in
plants. The EMBO J. 15, 2988-2996

Beemster GT, Vercruysse S, De VL, et al. The Arabidopsis leaf as a model system for
investigating the role of cell cycle regulation in organ growth. Journal of Plant
Research, 2006, 119(1):43-50.

Bouché N, Fait A, Bouchez D, Møller SG, Fromm H. Mitochondrial
succinic-semialdehyde dehydrogenase of the gamma-aminobutyrate shunt is
required to restrict levels of reactive oxygen intermediates in plants. Proc Natl
Acad Sci USA 2003; 100:6843-8.

Boudolf V, Vlieghe K, Beemster GT, Magyar Z, Torres Acosta JA, Maes S, Van Der
Schueren E, Inzé D, De Veylder L. (2004). The plant-specific cyclin-dependent
kinase CDKB1;1 and transcription factor E2Fa-DPa control the balance of
mitotically dividing and endoreduplicating cells in Arabidopsis. Plant Cell.
16(10):2683-2692.

Bown AW, Shelp BJ. Plant GABA: Not Just a Metabolite. Trends Plant Sci. 2016,
21(10):811-813.

Breuer C, Braidwood L, Sugimoto K. Endocycling in the path of plant development.
Curr Opin Plant Biol. 2014, 17:78-85.

Breuer C, Ishida T, Sugimoto K. Developmental control of endocycles and cell
growth in plants. Current Opinion in Plant Biology, 2010, 13(6):654-660.

Brikis CJ, Zarei A, Trobacher CP, DeEll JR, Akama K,Mullen RT ,Bozzo GG, Shelp
BJ. Ancient Plant Glyoxylate/Succinic Semialdehyde Reductases: GLYR1s Are
Cytosolic, Whereas GLYR2s Are Localized to Both Mitochondria and Plastids.
Front Plant Sci. 2017; 8: 601.

Castellano MM, Boniotti MB, Caro E, et al. DNA replication licensing affects cell
proliferation or endoreplication in a cell type-specific manner. The Plant cell,
2004, 16(9): 2380-93.

Castellano MM, del Pozo JC, Ramirez-Parra E, Brown S, Gutierrez C. Expression and
stability of Arabidopsis CDC6 are associated with endoreplication. Plant Cell.
2001, 13(12):2671-86.

Churchman ML, Brown ML, Kato N, Kirik V, Hülskamp M, Inzé D, De Veylder L,
Walker JD, Zheng Z, Oppenheimer DG, Gwin T, Churchman J, Larkin JC.
SIAMESE, a plant-specific cell cycle regulator, controls endoreplication onset in
Arabidopsis thaliana. Plant Cell. 2006, 18(11):3145-57.

Coelho, C., Dante, R. Sabelli, P., Sun, Y., Dilkes, B., Gordon-Kamm, W., and Larkins,
B. (2005). Cyclin-dependent kinase inhibitors in maize endosperm and their
potential role in endoreduplication. Plant Physiol. 138, 2323-2336.

Coleman ST, Fang TK, Rovinsky SA, Turano FJ, Moye-Rowley WS. Expression of a
glutamate decarboxylase homologue is required for normal oxidative stress
tolerance in Saccharomyces cerevisiae. J Biol Chem. 2001, 276(1):244-50.

De Veylder L, Larkin JC, Schnittger A. Molecular control and function of
endoreplication in development and physiology. Trends Plant
Sci. 2011,16(11):624-34.

De Veylder L., Beeckman T., Beemster G.T.S., Krols L., Terras F., Landrieu I., van
der Schueren E., Maes S., Naudts M., Inze D. (2001). Functional analysis of
cyclin-dependent kinase inhibitors of *Arabidopsis*. Plant Cell 13:1653-1667

Dewitte W, Riou-Khamlichi C, Scofield S, Healy JM, Jacqmard A, Kilby NJ, Murray
JA. Altered cell cycle distribution, hyperplasia, and inhibited differentiation in
Arabidopsis caused by the D-type cyclin CYCD3. Plant Cell. 2003, 15(1):79-92.

Dewitte W, Scofield S, Alcasabas AA, Maughan SC, Menges M, Braun N, Collins C,
Nieuwland J, Prinsen E, Sundaresan V, Murray JA. Arabidopsis CYCD3 D-type
cyclins link cell proliferation and endocycles and are rate-limiting for cytokinin
responses. Proc Natl Acad Sci USA. 2007, 104(36):14537-42.

Dubois M, Selden K, Bediee A, *et al*. SIAMESE-RELATED1 Is Regulated
Posttranslationally and Participates in Repression of Leaf Growth under
Moderate Drought. Plant physiology, 2018, 176(4): 2834-50.

Fait A, Yellin A, Fromm H. GABA shunt deficiencies and accumulation of reactive
oxygen intermediates: insight from Arabidopsis mutants.FEBS Lett. 2005,
579(2):415-20.

Fülöp K, Tarayre S, Kelemen Z, Horváth G, Kevei Z, Nikovics K, Bakó L, Brown S,
Kondorosi A, Kondorosi E. Arabidopsis anaphase-promoting complexes:
multiple activators and wide range of substrates might keep APC perpetually
busy. Cell Cycle. 2005, 4(8):1084-92.

Fung-Uceda J, Lee K, Seo PJ, Polyn S, De Veylder L, Mas P. The Circadian Clock
Sets the Time of DNA Replication Licensing to Regulate Growth in Arabidopsis.
Dev Cell. 2018，45 (1):101-113.

Häusler RE, Ludewig F, Krueger S. Amino acids--a life between metabolism and
signaling. Plant Sci. 2014, 229:225-237.

Heyman J, Polyn S, Eekhout T, De Veylder L.Tissue-Specific Control of the
Endocycle by the Anaphase Promoting Complex/Cyclosome Inhibitors UVI4
and DEL1. Plant Physiol. 2017,175(1):303-313.

Jacoby RP, Taylor NL, Millar AH. The role of mitochondrial respiration in salinity
tolerance. Trends Plant Sci 2011; 16:614–23.

Jun SE, Okushima Y, Nam J, Umeda M, Kim GT. Kip-related protein 3 is required
for control of endoreduplication in the shoot apical meristem and leaves of
Arabidopsis. Mol Cells. 2013, 35(1):47-53.

Kumar N, Harashima H, Kalve S, Bramsiepe J, Wang K, Sizani BL, Bertrand LL,
Johnson MC, Faulk C, Dale R, Simmons LA, Churchman ML, Sugimoto K,
Kato N, Dasanayake M, Beemster G, Schnittger A, Larkin JC. Functional
Conservation in the SIAMESE-RELATED Family of Cyclin-Dependent Kinase
Inhibitors in Land Plants. Plant Cell. 2015, 27(11):3065-80.

Lammens T, Li J, Leone G, et al. Atypical E2Fs: new players in the E2F transcription
factor family. Trends in Cell Biology, 2009, 19(3):111-118.

Liu CL，Zhao L，Yu GH. (2011). The dominant glutamic acid metabolic influx to
produce γ-amino butyric acid over proline in Nicotiana tabacum leaves under
water stress relates to its significant role of antioxidant activity. Integr. Plant Biol.
53:608-618.

Masubelele NH, Dewitte W, Menges M, Maughan S, Collins C, Huntley R,
Nieuwland J, Scofield S, Murray JA. D-type cyclins activate division in the root
apex to promote seed germination in Arabidopsis. Proc Natl Acad Sci USA.
2005，25; 102(43):15694-9.

Matsunaga, S., Katagiri, Y., Nagashima, Y., Sugiyama, T., Hasegawa, J., Hayashi, K.
and Sakamoto, T. (2013). New insights into the dynamics of plant cell nuclei and
chromosomes. Int. Rev. Cell Mol. Biol. 305, 253-301.

Melaragno J E, Mehrotra B, Coleman A W. Relationship between Endopolyploidy
and Cell Size in Epidermal Tissue of Arabidopsis. Plant Cell, 1993,
5(11):1661-1668

Michaeli S, Fait A, Lagor K, Nunes-Nesi A, Grillich N, Yellin A, et al. A
mitochondrial GABA permease connects the GABA shunt and the TCA cycle,
and is essential for normal carbon metabolism. Plant J 2011; 67:485–498.

Nowack MK, Harashima H, Dissmeyer N, *et al*. Genetic framework of
cyclin-dependent kinase function in *Arabidopsis*. Developmental cell, 2012,
22(5): 1030-40.

Palanivelu R, Brass L, Edlund AF, Preuss D. Pollen tube growth and guidance is
regulated by POP2, an Arabidopsis gene that controls GABA levels. Cell. 2003,
114(1):47-59.

Podlešáková K, Ugena L, Spíchal L, Doležal K, De Diego N. Phytohormones and
polyamines regulate plant stress responses by altering GABA pathway. New
Biotechnol. 2019, 48:53-65.

Ramesh SA, Tyerman SD, Gilliham M, Xu B. γ-Aminobutyric acid (GABA)
signalling in plants. Cell Mol Life Sci. 2017, 74(9):1577-1603.

Renault H, El Amrani A, Berger A, Mouille G, Soubigou-Taconnat L, Bouchereau A,
et al. γ-Aminobutyric acid transaminase deficiency impairs central carbon
metabolism and leads to cell wall defects during salt stress in Arabidopsis roots.
Plant Cell Environ 2013; 36:1009–18.

Renault H, El Amrani A, Palanivelu R, Updegraff EP, Yu A, Renou JP, Preuss D,
Bouchereau A, Deleu C. GABA accumulation causes cell elongation defects and
a decrease in expression of genes encoding secreted and cell wall-related proteins
in Arabidopsis thaliana.Plant Cell Physiol. 2011, 52(5):894-908.

Sawa S, Watanabe K, Goto K, Liu YG, Shibata D, Kanaya E, Morita EH, Okada K.
FILAMENTOUS FLOWER, a meristem and organ identity gene of Arabidopsis,
encodes a protein with a zinc finger and HMG-related domains.Genes Dev. 1999,
13(9):1079-88.

Schippers JH, Foyer CH, Van Dongen JT. Redox regulation in shoot growth, SAM
maintenance and flowering. Current Opinion in Plant Biology, 2016,
29:121-128.

Seifikalhor M, Aliniaeifard S, Hassani B, Niknam V & Lastochkina O. (2019).
Diverse role of γ-aminobutyric acid in dynamic plant cell responses. Plant Cell
Reports. 38(8):847-867.

Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL. Members of
the YABBY gene family specify abaxial cell fate in *Arabidopsis*. Development.
1999, 126(18):4117-28.

Toyokura K, Hayashi M, Nishimura M, and Okada K. Adaxial-abaxial patterning: a
novel function of the GABA shunt. Plant Signal Behav. 2012，7 (7):705-7.

Toyokura K, Watanabe K, Oiwaka A, Kusano M, Tameshige T, Tatematsu K,
Matsumoto N, Tsugeki R, Saito K, Okada K. Succinic semialdehyde
dehydrogenase is involved in the robust patterning of Arabidopsis leaves along
the adaxial-abaxial axis. Plant Cell Physiol. 2011, 52(8):1340-53.

Umeda M, Aki SS, Takahashi N. Gap 2 phase: making the fundamental decision to
divide or not. Curr Opin Plant Biol. 2019, 51:1-6.

Verkest A, Manes CL, Vercruysse S, Maes S, Van Der Schueren E, Beeckman T,
Genschik P, Kuiper M, Inzé D, De Veylder L (2005). The cyclin-dependent
kinase inhibitor KRP2 controls the onset of the endoreduplication cycle during
Arabidopsis leaf development through inhibition of mitotic CDKA; 1 kinase
complexes. Plant Cell. 2005, 17(6):1723-1736.

Vlieghe K, Boudolf V, Beemster G T, et al. (2005). The DP-E2F-like Gene DEL1,
Controls the Endocycle in Arabidopsis thaliana. Current Biology, 15(1):59-63.

Watkins JM, Chapman JM, Muday GK (2017). Abscisic Acid-Induced Reactive
Oxygen Species Are Modulated by Flavonols to Control Stomata Aperture. Plant
Physiol. 175(4):1807-1825.

Yang K, Wang H, Xue S, *et al*. Requirement for A-type cyclin-dependent kinase and
cyclins for the terminal division in the stomatal lineage of *Arabidopsis*. J Exp
Bot, 2014, 65(9): 2449-2461.

Yi D, Alvim Kamei CL, Cools T, Vanderauwera S, Takahashi N, Okushima Y,
Eekhout T, Yoshiyama KO, Larkin J, Van den Daele H, Conklin P, Britt A,
Umeda M, De Veylder L. The Arabidopsis SIAMESE-RELATED
cyclin-dependent kinase inhibitors SMR5 and SMR7 regulate the DNA damage
checkpoint in response to reactive oxygen species. Plant Cell. 2014,
26(1):296-309.

Figure Legends

Figure 1. Leaf area of γ-Aminobutyric acid (GABA) biosynthetic mutants was larger
than that of wild-type Arabidopsis.

Top view of leaves at 8 d after stratification of A. wild type; B. *gad1* mutant, C. *gad2*
mutant and D. *gad1/gad2* mutants; E. Leaf growth curve of GABA mutants and wild
type of Arabidopsis in the early stage (from 4–8 d after stratification). Results are
presented as averages ± SE of three separate experiments (n = 15). Bar in A–D = 860
μm.

Figure 2. Epidermal cell size in the first true leaves of γ-Aminobutyric acid (GABA)
biosynthetic mutants was larger than that of wild-type Arabidopsis at the early growth
stage.

Epidermal cell of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2*
mutants; E. Cell size comparison among GABA mutants and wild type of Arabidopsis
in the early stage (7–9 d after stratification). Results are presented as averages ± SE of
60 cells calculated across three separate experimental replicates. Asterisks represent
significant differences between the mutants and the wild type (P <0.05) and were
determined using one-way ANOVA. Bar in A–D = 55 μm.

Figure 3. Octoploid (8n) cells occurred earlier in biosynthetic mutants than that in the
wild type. Top panel: DNA ploidy analysis in leaf early growth stage (7–8 days after
stratification) between γ-Aminobutyric acid (GABA) biosynthetic mutants and the
wild type. Left bottom panel: DNA ploidy analysis in leaf middle-late stage (21 days
after stratification) between GABA mutants and the control. Right bottom panel: 8C
percentage comparison among mutants and the wild type during the long growth
period.

Figure 4. Expression of *Cyclin D* components in biosynthetic mutants was lower than
that in the wild type at the 8th day after stratification. Results are presented as averages
± SE of triplicate experiments. Asterisks represent significant differences between the
mutants and the wild type (*P <0.05; ** P <0.01) and were determined using
one-way ANOVA.

Figure 5. Expression of *CDA;1* and *E2Fa* in biosynthetic mutants was lower than that
in the wild type at the 8th day after stratification. Results are presented as averages ±
SE of three independent experimental replicates. Asterisks represent significant
differences between the mutants and the wild type (*P <0.05; ** P <0.01) and were
determined using one-way ANOVA.

Figure 6. Expression of *CCS52A* and *CDC6* in biosynthetic mutants was higher than
that in the wild type at the 8th day after stratification. Results are presented as averages
± SE of three independent experimental replicates. Asterisks represent significant
differences between the mutants and the wild type (*P <0.05; ** P <0.01) and were
determined using one-way ANOVA.

Figure 7. Expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*)
components in biosynthetic mutants was higher than that in the wild type at the 8th
day after stratification. Results are presented as averages ± SE of three independent
experimental replicates. Asterisks represent significant differences between the
mutants and the wild type (*P <0.05; ** P <0.01). Statistics were determined using
one-way ANOVA.

Figure 8. Reactive oxygen species (ROS) level in guard cells (GCs) of
γ-Aminobutyric acid (GABA) mutants are higher than that of control A–D. DCF
fluorescence (green) in epidermal GCs of GABA biosynthetic mutants and wild type
Arabidopsis. E. DCF fluorescence was quantified in the mutants and control. Results
are presented as averages ± SE of three separate experiments (n = 16). Asterisks
represent significant differences between the mutant and the wild type (P <0.05) and
were determined using one-way ANOVA. Bar in A–D = 55 μm.

Figure 9. Working hypothesis of γ-Aminobutyric acid (GABA) negatively control the
endoreplication of Arabidopsis leaves.

GABA, as a normal metabolic signal and antioxidant, effectively inhibits reactive
oxygen species (ROS) production. In the GABA biosynthetic mutants, owing to
biosynthetic perturbation, the inhibition of GABA on ROS was relieved. The
production of ROS promotes the expression of *SIAMESE* (*SIM*) and
SIAMESE-RELATED (*SMR*). Both components are inhibitors of Cyclin-dependent
Kinase complexes (e.g. CYC3;1-CDKA;1), and low levels of kinases initiate
endoreplication in Arabidopsis. Other components, such as CDC6, synergistically
respond to low levels of GABA to prime endoreplication. In the wild type, the normal
metabolism of GABA delays endoreplication and prevents premature endoreplication.

Supporting Materials

Table S1. The primers of the genes involved in cell cycle regulation, endocycle
initiation, progression, and exit.

Figure S1. The γ-Aminobutyric acid (GABA) analysis was performed on whole-shoot
samples. Results are presented as averages ± standard error (SE) of three independent

replicates. Significant differences from the control are indicated with asterisks: * P

<0.05, ** $P < 0.01$, by one-way ANOVA.

Figure 1. Leaf area of γ-Aminobutyric acid (GABA) biosynthetic mutants was larger
than that of wild-type Arabidopsis.

Top view of leaves at 8 d after stratification of A. wild type; B. *gad1* mutant, C. *gad2*
mutant and D. *gad1/gad2* mutants; E. Leaf growth curve of GABA mutants and wild
type of Arabidopsis in the early stage (from 4–8 d after stratification). Results are
presented as averages ± SE of three separate experiments (n = 15). Bar in A–D = 860
μm.

Figure 2. Epidermal cell size in the first true leaves of γ-Aminobutyric acid (GABA)
biosynthetic mutants was larger than that of wild-type Arabidopsis at the early growth
stage.

Epidermal cell of A. wild type; B. *gad1* mutant, C. *gad2* mutant and D. *gad1/gad2*
mutants; E. Cell size comparison among GABA mutants and wild type of Arabidopsis
in the early stage (7–9 d after stratification). Results are presented as averages ± SE of
60 cells calculated across three separate experimental replicates. Asterisks represent

significant differences between the mutants and the wild type ($P < 0.05$) and were
determined using one-way ANOVA. Bar in A–D = 55 μm.

Figure 3. Octoploid (8n) cells occurred earlier in biosynthetic mutants than that in the
wild type. Top panel: DNA ploidy analysis in leaf early growth stage (7–8 days after
stratification) between γ-Aminobutyric acid (GABA) biosynthetic mutants and the
wild type. Left bottom panel: DNA ploidy analysis in leaf middle-late stage (21 days
after stratification) between GABA mutants and the control. Right bottom panel: 8C
percentage comparison among mutants and the wild type during the long growth
period.

680   Figure 4. Expression of *Cyclin D* components in biosynthetic mutants was lower than
681   that in the wild type at the 8th day after stratification. Results are presented as averages
682   ± SE of triplicate experiments. Asterisks represent significant differences between the
683   mutants and the wild type (*P <0.05; ** P <0.01) and were determined using
684   one-way ANOVA.

Figure 5. Expression of *CDA;1* and *E2Fa* in biosynthetic mutants was lower than that in the wild type at the 8th day after stratification. Results are presented as averages ± SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (*P <0.05; ** P <0.01) and were determined using one-way ANOVA.

Figure 6. Expression of *CCS52A* and *CDC6* in biosynthetic mutants was higher than that in the wild type at the 8th day after stratification. Results are presented as averages ± SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (*P <0.05; ** P <0.01) and were determined using one-way ANOVA.

Figure 7. Relative expression of *SIM* gene

Figure 7. Expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*) components in biosynthetic mutants was higher than that in the wild type at the 8[th] day after stratification. Results are presented as averages ± SE of three independent experimental replicates. Asterisks represent significant differences between the mutants and the wild type (*P <0.05; ** P <0.01). Statistics were determined using one-way ANOVA.

Figure 8. Reactive oxygen species (ROS) level in guard cells (GCs) of γ-Aminobutyric acid (GABA) mutants are higher than that of control A–D. DCF fluorescence (green) in epidermal GCs of GABA biosynthetic mutants and wild type Arabidopsis. E. DCF fluorescence was quantified in the mutants and control. Results are presented as averages ± SE of three separate experiments (n = 16). Asterisks

723   represent significant differences between the mutant and the wild type ($P < 0.05$) and
724   were determined using one-way ANOVA. Bar in A–D = 55 μm.

Figure 9. Working hypothesis of γ-Aminobutyric acid (GABA) negatively control the endoreplication of Arabidopsis leaves.

GABA, as a normal metabolic signal and antioxidant, effectively inhibits reactive oxygen species (ROS) production. In the GABA biosynthetic mutants, owing to biosynthetic perturbation, the inhibition of GABA on ROS was relieved. The production of ROS promotes the expression of *SIAMESE* (*SIM*) and *SIAMESE-RELATED* (*SMR*). Both components are inhibitors of Cyclin-dependent Kinase complexes (e.g. CYC3;1-CDKA;1), and low levels of kinases initiate endoreplication in Arabidopsis. Other components, such as CDC6, synergistically respond to low levels of GABA to prime endoreplication. In the wild type, the normal metabolism of GABA delays endoreplication and prevents premature endoreplication.

743   **Supporting Materials**

744   Table S1: The primers of the genes involved in cell cycle regulation, endocycle

745   initiation, progression and exit.

Locus tag	Gene name	Primer	Primer sequences
AT3G02780	*IPP2*	IPP2_EXP_F	CATGCGACACACCAACACCA
		IPP2_EXP_R	TGAGGCGAATCAATGGGAGA
AT4G34160	*CYCD3;1*	CYCD3;1_EXP_F	CCTCTCTGTAATCTCCGATTC
		CYCD3;1_EXP_R	AAGGACACCGAGGAGATTAG
AT5G67260	*CYCD3;2*	CYCD3;2_EXP_F	TCTCAGCTTGTTGCTGTGGCTTC
		CYCD3;2_EXP_R	TCTTGCTTCTTCCACTTGGAGGTC
AT3G50070	*CYCD3;3*	CYCD3;3_EXP_F	TCCGATCGGTGTGTTTGATGCG
		CYCD3;3_EXP_R	GCAGACACAACCCACGACTCATTC
AT5G65420	*CYCD4;1*	CYCD4;1_EXP_F	GAAGGAGAAGCAGCATTTGCCAAG
		CYCD4;1_EXP_R	ACTGGTGTACTTCACAAGCCTTCC
AT4G11920	*CCS52A2*	CCS52A2_EXP_F	CGTAGATACCAACAGCCAGGTGTG
		CCS52A2_EXP_R	CGTGTGTGCTCACAAGCTCATTC
AT2G29680	*CDC6*	CDC6_EXP_F	AGGCTCTATGTGTCTGCAGGAG
		CDC6_EXP_R	ACCACTTGACACTCTGGAACTGG
AT2G31270	*CDT1a*	CDT1a_EXP_F	AATCGCTCTTCGGAAAGTGTTTCG
		CDT1a_EXP_R	CCTCTGGAACTTCATCACCCTGAG
AT3G48750	*CDKA;1*	CDKA;1_EXP_F	ACTGGCCAGAGCATTCGGTATC
		CDKA;1_EXP_R	TCGGTACCAGAGAGTAACAACCTC
AT2G36010	*E2Fa*	E2Fa_EXP_F	TAGATCGGGAGGAAGATGCTGTCG

		E2Fa_EXP_R	TTGTCGCCTTTCTCTTTCGTGAAG
AT2G23430	*KRP1*	KRP1_EXP_F	ACGGAGCCGGAGAATTGTTTATG
		KRP1_EXP_R	CGAAACTCCATTATCACCGACGAC
AT3G50630	*KRP2*	KRP2_EXP_F	TAGGAGATTATGGCGGCGGTTAGG
		KRP2_EXP_R	TTTCACCGTCGTCGTCGTAACTC
AT2G32710	*KRP4*	KRP4_EXP_F	AAGCTTCAACAGGACCACAAGGG
		KRP4_EXP_R	GGGTTGTCATGATTTCAGGCCTTC
AT1G49620	*KRP7*	KRP7_EXP_F	GAGGCTCATGAAATCTCCGAAACC
		KRP7_EXP_R	CCGAGTCCATTTCTGCTGTTTCTC
AT5G04470	*SIM*	SIM_EXP_F	AGCCATCAAGATCCGAGCCAAC
		SIM_EXP_R	TTGTGGTCGGAAGAAGTGGGAGTG
AT3G10525	*SMR1*	SMR1_EXP_F	CAAAGAAGGACGAAGGTGATGACG
		SMR1_EXP_R	TGTTCTTGGGATGTGGGTGTGC
AT1G60783	*SMR2*	SMR2_EXP_F	TCACAAGATTCCGGAGGTGGAGAC
		SMR2_EXP_R	ATCTCACGCGGTCGCTTTCTTG
AT1G07500	*SMR5*	SMR5_EXP_F	ACGCCTACACGTGATGATTGCC
		SMR5_EXP_R	TATCCCTTCTTCGGTGGTTCCC
AT1G10690	*SMR8*	SMR8_EXP_F	GCGGTTTCCGTCAGAATTCCAAG
		SMR8_EXP_R	GCACTTCAACGACGGTTTACGC

Figure S1: GABA analysis was performed on whole shoot samples. Avarages ±
standard error (SE) of three independent replicates. Significant differences from
the control are indicated with asterisks: * P <0.05, ** P < 0.01, by one-way
ANOVA.

GABA levels in the leaves of the mutants were determined essentially as
described earlier (Allan and Shelp 2006).

Allan WL, Simpson JP, Clark SM, Shelp BJ. Gamma-hydroxybutyrate
accumulation in Arabidopsis and tobacco plants is a general response to abiotic
stress: putative regulation by redox balance and glyoxylate reductase isoforms. J
Exp Bot 2008; 59: 2555–64.