**Evolutionary and ontogenetic changes of the anatomical organization and modularity in the**

**skull of archosaurs**

**Short title: Evolution of network anatomy in archosaurian skulls**

Hiu Wai Lee[1,2], Borja Esteve-Altava[3*], Arkhat Abzhanov[1,2*]

Affiliations:

[1] Department of Life Sciences, Imperial College London, Silwood Park Campus,

Buckhurst Road, Ascot, Berkshire SL5 7PY, United Kingdom

[2] Natural History Museum, Cromwell Road, London SW7 5BD, United Kingdom

[3] Institute of Evolutionary Biology (UPF-CSIC), Department of Experimental and Health

Sciences, Pompeu Fabra University, Barcelona, Spain.

* Corresponding authors: boresal@gmail.com (B.E.A.), a.abzhanov@imperial.ac.uk (A.A.)

**Abstract**

Comparative anatomy studies of the skull of archosaurs provide insights on the mechanisms of evolution for the morphologically and functionally diverse species of crocodiles and birds. One of the key attributes of skull evolution is the anatomical changes associated with the physical arrangement of cranial bones. Here, we compare the changes in anatomical organization and modularity of the skull of extinct and extant archosaurs using an Anatomical Network Analysis approach. We show that the number of bones, their topological arrangement, and modular organization can discriminate birds from non-avian dinosaurs, and crurotarsans. We could also discriminate extant taxa from extinct species when adult birds were included. By comparing within the same framework, juveniles and adults for crown birds and alligator (*Alligator mississippiensis*), we find that adult and juvenile alligator skulls are topologically similar, whereas juvenile bird skulls have a morphological complexity and anisomerism more similar to those of non-avian dinosaurs and crurotarsans than of their own adult forms. Clade-specific ontogenetic differences in skull organization, such as extensive postnatal fusion of cranial bones in crown birds, can explain this pattern. The fact that juvenile and adult skulls in birds do share a similar anatomical integration suggests the presence of a specific constraint to their ontogenetic growth.

**Keywords**: Comparative Anatomy; Cranium; Anatomical Network Analysis; Birds; Crocodiles; craniofacial evolution, Archosauria

## INTRODUCTION

The skulls of archosaurs are morphologically and functionally diverse, with clade-specific specialized features that set apart crurotarsans (extant crocodilians and their stem lineage) from avemetatarsalians (birds and non-avian dinosaurs)[1-7], as reviewed by Brusatte and colleagues[8]. The evolution and diversification of the skull of archosaurs have been associated with changes in the patterns of phenotypic integration and modularity[9-13]. For more information on integration and modularity in shape, see the review by Klingenberg[14]. Different regions of the skull may act as anatomical modules that can evolve, function, and develop semi-independently from one another. Bones within a same module tend to co-vary in shape and size more with each other than with bones from other such variational modules[15-18]. In addition, the bones of the skull can also modify their physical articulations so that some groups of bones are more structurally integrated than others, and, hence, we can recognize them as distinct anatomical-network modules, which had been defined by Eble as a type of organizational modules[15,19,20]. The relationship between anatomical-network modules and variational modules is not yet fully understood, but it is thought that network anatomy constrain growth patterns and shape variation[21-23].

Changes in the anatomical organization of the skull in archosaurs have been concomitant with a broader evolutionary trend in tetrapods toward a reduction in the number of skull bones due to loses and fusions, a phenomenon known as the Williston's law[24-26]. Understanding how the bones are globally arranged to each other allows us to measure the anatomical complexity and organization of body parts, and explain how structural constraints might have influenced the direction of evolution[25-28]. Werneburg and colleagues compared the skull network-anatomy of a highly derived *Tyrannosaurus rex*, *Alligator mississippiensis* and *Gallus gallus* with that of an opossum, a tuatara, and a turtle[29]. They found that the tyrannosaur has the most modular skull organization among these amniotes, with a modular separation of the snout in upper and lower sub-modules and the presence of a lower adductor chamber module. However, the specific

anatomical changes in the organization of the archosaur skull during their evolutionary transitions

more generally have never been characterized. More recently, Plateau and Foth used anatomical

network analysis to study postnatal ontogenetic changes in the skulls of crown bird and non-avian

theropods[30]. They found that early juvenile crown birds have skulls that are less integrated and

more modular than those of more derived birds, resembling their non-avian theropod ancestors.

Here, we compared the anatomical organization and modularity of the skull of archosaurs using

Anatomical Network Analysis (AnNA) [31] to highlight how skull topology has changed in

evolutionary and developmental scales. We chose AnNA over more conventional methods, such

as geometric morphometrics, to understand how major re-organizations of the skull (i.e., loss and

fusion of bones) affect the overall anatomy regardless of shape. We created network models of

the skull for 21 species of archosaurs, including taxa representing key evolutionary transitions

from early pseudosuchians to crocodiles, from non-avian theropods to modern birds, and from

paleognath birds to neognaths (Fig. 2). Our dataset also includes a representative ornithischian, a

sauropodomorph, and a basal saurischian (Supplementary Information 1) for comparison. To

understand the significance of the ontogenetic transitions in archosaur skulls, we provided our

dataset with juvenile skulls for extant birds and alligator. Network models of the skull were built

by coding individual cranial bones and their articulations with other bones as the nodes and links

of a network, respectively (Fig. 1). Network modules, defined as a group of bones with more

articulations among them than to other bones outside the module, were identified heuristically

using a community detection algorithm. We compared skull architectures using topological

variables (i.e. network parameters) that capture whole-skull anatomical feature (modelling and

analysis of anatomical networks were detailed previously [20,25,31]).

Networks and network modules and their respective complexity, integration, modularity, and

anisomerism could be quantified by these network parameters: density of connections, clustering

coefficient, path length, heterogeneity of connections, and parcellation[20,23,31,32]. Here, complexity

is defined as the relationship of bones in a skull and is associated with how abundant are the

interactions that bones have with each other (i.e. density of connections), how interdependent or

integrated the bones are (i.e. clustering coefficient), and proximity between nodes (i.e. path

length). A more complex network would have higher density, higher clustering coefficient, and

shorter path length. Anisomerism is defined as a deviation among anatomical parts[33] and could be

observed by the specialization of bones and measured by heterogeneity of connections, i.e. how

each bone has a different number of connection[25]. Modularity is measured by parcellation, which

is the number of modules and the consistency in the number of bones per module.

**MATERIAL AND METHODS**

**Sampling**

We sampled extinct and extant species, and for some forms included both adults and juveniles to

account for ontogenetic trends within archosaurs. Namely, adults *Aetosaurus ferratus,*

*Archaeopteryx lithographica, Citipati osmolskae, Coelophysis bauri, Compsognathus longipes,*

*Dakosaurus andiniensis, Desmatosuchus haplocerus, Dibothrosuchus elaphros, Dilophosaurus*

*wetherilli, Eoraptor lunensis, Ichthyornis dispar*, *Plateosaurus engelhardti*, *Psittacosaurus*

*lujiatunensis*, *Riojasuchus tenuisceps*, *Sphenosuchus acutus*, *Velociraptor mongoliensis*, *Gallus*

*gallus*, *Geospiza fortis* and *Nothura maculosa*; and juveniles *Gallus gallus*, *Geospiza fortis*,

*Nothura maculosa* and *Alligator mississippiensis*. Within our sample set, eight species represent

the transition from crurotarsan archosaur ancestor to modern crocodilians and 13 species

represent the transition from non-avian theropods to modern birds as described previously[34–43].

Due to the sample size limitation for extinct taxa, reconstructed and type forms were used to

represent each taxon and intraspecific variation could not be accounted for.

**Phylogenetic Context**

We created a phylogenetic tree (Figure 2) based on the previous studies[34–37,39–44]. The tree was

calibrated using the R package paleotree[45] by the conservative "equal" method[46,47]; branching

events were constrained using the minimum dates for known internal nodes based on fossil data

from Benton and Donoghue[48] (listed in Table S3) and the first and last occurrences of all 21

species from the Paleobiology Database using the paleobioDB package [49] in R. Because there

were two extinct *Nothura* species in the Paleobiology Database, the last occurrence for extant

*Nothura* species was adjusted to 0 (Table S2).

**Network Modelling**

We built anatomical network models for each archosaur skull in our sample set based on detailed

literature descriptions and CT scans of complete skulls (see Supplementary Information 1). Skull

bones were represented as the nodes of the network model and their pair-wise articulations (e.g.

sutures and synchondroses) were represented as links between pairs of nodes (Figure 1). Skull

network models were formalized as binary adjacency matrices, in which a 1 codes for two bones

articulating and a 0 codes for absence of articulation. Bones that were fused together without

trace of a suture in the specimens examined were formalized as a single individual bone.

**Network Analysis**

Following Esteve-Altava et al[28], we quantified the following topological variables for each

network model: the number of nodes (N), the number of links (K), the density of connections (D),

the mean clustering coefficient (C), the mean path length (L), the heterogeneity of connections

(H), the assortativity of connections (A), and the parcellation (P). The morphological

interpretation of these topological variables has been detailed elsewhere[28]. A summary is

provided here. N and K represent the direct count of the number of individual bones and

articulations observed in the skull. D is the number of connections divided by the maximum

number of possible connections (it ranges from 0 to 1); D is a proxy measure for morphological

complexity. C is the average number of neighboring bones that connect to one another in a

network (i.e., actual triangles of nodes compared to the maximum possible): a value close to 1

shows all neighboring bones connect to each other while a value close to 0 shows neighboring

bones do not connect to each other; C is a proxy measure for anatomical integration derived from

co-dependency between bones. L measures average number of links separating two nodes (it

ranges from 1 to N-1); L is a proxy measure of anatomical integration derived from the effective

proximity between bones. H measures how heterogeneous connections are in a network: skulls

composed of bones with a different number of articulations have higher H values. If all bones had

the same number of connections (i.e., H = 0), it means that all bones were connected in the same

way and the skull had a regular shape. A measures whether nodes with the same number of

connections connect to each other (it ranges from -1 to 1); H and A are a proxy measure for

anisomerism or diversification of bones. P measures the number of modules and the uniformity in

the number of bones they group (it ranges from 0 to 1); P is a proxy for the degree of modularity

in the skull. Calculating P requires a given partition of the network into modules (see next below).

Network parameters were quantified in R[50] using the igraph package[51]. Networks visualization

was made using the visNetwork package[52] and Cytoscape[53].

**Principal Component Analysis**

We performed a Principal Component Analysis (PCA) of the eight topological variables with a

singular value decomposition of the centered and scaled measures. On the resulting PCs, we used

a PERMANOVA (10,000 iterations) to test whether topological variables discriminate between:

(1) Avialae and non-Avialae; (2) adults and juveniles; (3) extinct and extant; (4) Crurotarsi and

Avemetatarsalia; (5) Neornithes and non-Neornithes; (6) early flight, can do soaring flight, can do

flapping flight, gliding, and flightless (details in Table S5); (7) Crurotarsi, non-avian Dinosauria,

and Aves; and (8) carnivorous, omnivorous, and herbivorous (dietary information in

Supplementary Information 4). First, we performed the tests listed above for all archosaurs. Then,

we repeated these tests for a sub-sample that included all archosaurs, except for all modern birds.

Next, we repeated these tests for a sub-sample that included all archosaurs, except for adult birds.

**Modularity Analysis**

To find the optimal partition into network modules we used a node-based informed modularity

strategy[54] . This method starting with the local modularity around every individual node, using

cluster_spinglass function in igraph[51], then it returns the modular organization of the network by

merging non-redundant modules and assessing their intersection statistically using combinatorial

theory[55].

**RESULTS**

**Topological discrimination of skull bones**

A Principal Component Analysis (PCA) of the eight topological variables measured in skull

network models discriminates skulls with different anatomical organizations (Figs. S1-S3). When

all sampled skulls are compared together, the first three principal components (PCs) explain 89.4%

of the total variation of the sample. PC1 (57.5%) discriminates skulls by number of their bones

(N), density of connections (D), and degree of modularity (P). PC2 (21.3%) discriminates skulls

by their degree of integration (C) and anisomerism (H). Finally, PC3 (10.6%) discriminates skulls

by whether bones with similar number of articulations connect with each other (A).

PERMANOVA tests confirm that different skull anatomies map onto different regions of the

morphospace. Thus, we can discriminate: Avialae (Aves plus *Ichthyornis* and *Archaeopteryx*)

versus non-Avialae ($F_{1,23} = 4.124$, $p = 0.006699$; Fig. 3B); Neornithes plus toothless archosaurs

versus archosaurs with teeth ($F_{1,23} = 6.99$, $p = 0.0005999$; Fig. 3C); Aves (include all modern

birds) versus Crurotarsi versus non-avian Dinosauria ($F_{2,22} = 3.837$, $p = 0.000699$; Fig. 3D); and

extant and extinct species ($F_{1,23} = 4.304$, $p = 0.0017$; Fig. S1C). However, we find no statistically

significant difference in morphospace occupation between crurotarsans and avemetatarsalians

($F_{1,23} = 1.46$, $p = 0.2002$, Fig. S1D).

When all avians are excluded from the comparison, the first three PCs now explain 80.6% of the

total variation (Figs. S4-6). PC1 (38.6%) discriminates skulls by the density of their inter-bone

connections (D) and effective proximity (L). PC2 (22.6%) discriminates skulls by the number of

bones and their articulations (N and K). Finally, PC3 (19.5%) now discriminates skulls by their

anisomerism (H) and whether bones with the same number of connections connect to each (A).

PERMANOVA tests could not discriminate between Crurotarsi and non-avian Dinosauria ($F_{1,17} =$

$1.235$, $p = 0.3022$; Fig. S4D), and between extant and extinct species ($F_{1,17} = 2.274$, $p = 0.06399$;

Fig. S4C).

When only adult birds are excluded, the first three PCs explain 79.7% of the topological variation

(Figs. S7-9). PC1 (35.8%), PC2 (24.5%), and PC3 (19.5%) discriminate skull similarly as when

all birds are excluded (see above). PERMANOVA tests also could not discriminate between

juvenile birds, crurotarsans, and non-avian dinosaurs ($F_{2,19} = 1.682$, $p = 0.09649$; Fig. S7D), and

between extant and extinct species ($F_{1,20} = 2.119$, $p = 0.06169$; Fig. S7C).

Regardless of the sub-sample compared, we found no statistically significant difference in

morphospace occupation between taxa stratified by flying ability and diet (Fig. S1E, see

Supplementary Information 4 for details). This suggests that at least for the given sample set

changes in cranial network-anatomy (i.e. how bones connect to each other) are independent of

both dietary adaptations and the ability to fly.

**Number of network modules**

The number of network modules identified in archosaur skulls ranged from one (i.e. fully

integrated skull) in adult birds *Nothura maculosa* (the spotted tinamou) and *Geospiza fortis*

(medium ground finch) to eight in the non-avian dinosaur *Citipati* (Table S10). The number of

network modules within the studied taxa decreases during evolution of both major archosaurian

clades: from 6 (*Riojasuchus*) to 4 (*Desmatosuchus*,) and from 6 (*Dibothrosuchus*) to 4

(*Dakosaurus* and all adult crocodilians) modules in Crurotarsi; from 6 (*Coelophysis*) to 4

(*Dilophosaurus* and *Compsognathus*), and from 8 (*Citipati*) to 4 (*Velociraptor, Archaeopteryx,*

*Ichthyornis,* and juvenile modern birds) modules in theropod-juvenile bird transition (Fig. 4A and

4B, Table S10). We found no modular division of the skull in adult *Nothura* and *Geospiza*. This

is most likely because these skulls are highly integrated due to the extensive cranial bone fusion

in adults, which, in turn, results in a network with very few nodes. In general, skull networks are

partitioned into overlapping modules.

**DISCUSSION**

**Occupation of morphospace and evolution of skull architecture**

The two major groups of archosaurs (Crurotarsi and Avemetatarsalia) show an analogous trend

towards a reduction in the number of skull bones (Table S8; Supplementary Information 3), in

line with the Williston's Law, which states that vertebrate skulls tend to become more specialized

with fewer bones as a result of fusions of neighboring bones during evolution[25,56,57].This

reduction in the number of bones and articulations, together with an increase in density, is also

observed within aetosaurs and sphenosuchians (Table S8). Likewise, we observed fusion of

paired bones into new unpaired ones: for example, left and right frontals, parietals, and palatines

are fused through their midline suture in the more derived taxa, such as the crocodilians (Table

S6). Bone fusion in extant species produced skulls that are more densely connected than the

skulls of extinct species (Fig. S1C). It was previously suggested that the more connected skulls

would have more developmental and functional inter-dependences among bones, and, hence, they

would be more evolutionarily constrained[22,23]. Similarly, avian cranium with its strongly

correlated traits has lower evolutionary rates and bird skulls are less diverse overall[12].

Bhullar et al. pointed out that avian kinesis relies on their loosely integrated skulls with less

contact and, thus, skulls with highly overlapping bones would be akinetic[58]. This contradicts our

observations here in that kinetic crown birds have more complex and integrated skulls than the

akinetic crurotarsans and the partially kinetic *Riojasuchus*[59]. The reason could be that Bhullar et

al. factored in how much connective tissue and number of contact points each bone has, but not

the total number of connections possible from the number of bones in these taxa. The total

number of articulations possible is the denominator used to calculate density. More recently,

Werneburg and colleagues showed *Tyrannosaurus*, suspected to have kinesis, also has a higher

density when compared to akinetic *Alligator* but lower density when compared to the more

derived and clearly kinetic *Gallus* skull[29].

When compared with modules identified by Felice et al.[60], the node-based modules, such as the

rostral and neurocranial modules (shown as blue and red modules in Fig. 4), are composed of

elements essentially similar to those described as variational modules (more details in

Supplementary Information 2). The supraoccipital and basioccipital bones were part of the same

topology-defined (Supplementary Information 2, Fig. 4) and shape-defined module in most taxa,

likely due to its functional importance in connecting the vertebral column with the skull[60].

**Crurotarsi**

The aetosaurs, *Aetosaurus* and *Desmatosuchus,* and the sphenosuchians, *Sphenosuchus* and

*Dibothrosuchus,* show an increase in complexity within their lineages. The more derived aetosaur

*Desmatosuchus* has a fused skull roof (parietal fused with supraoccipital, laterosphenoid, prootic

and opisthotic) and toothless premaxilla that are absent in the less derived aetosaur *Aetosaurus*[61–

63]. In contrast, basal and derived sphenosuchian are more topologically similar. Their main

difference is that basipterygoid and epiotic are separate in *Sphenosuchus* but are fused with other

bones in the more derived *Dibothrosuchus*[64,65]. When we compared aetosaurs and sphenosuchians,

we found that sphenosuchians have a skull roof intermediately fused condition between

*Aetosaurus* and *Dibothrosuchus*: interparietal sutures in both sphenosuchians are fused while

supraoccipital, laterosphenoid, opisthotic, and prootic remain separate.

To understand cranial topology in Thalattosuchia, a clade with adaptations specialized for marine

life, we included *Dakosaurus andiniensis*. These adaptations comprise nasal salt glands[66],

hypocercal tail, paddle-like forelimbs, ziphodont dentition, fusion of the inter-premaxillary suture,

a fused vomer, and a short and high snout[67,68]. Despite these adaptations, *Dakosaurus* has a

cranial complexity closer to that of extant crocodilians by similarly having inter-frontal and inter-

parietal fusions[67,68]. In addition to the fused frontals and parietals, both *Crocodylus* and *Alligator*

have a fused palate and a fused pterygoid bones.

In turn, crurotarsans first fuse the skull roof and skull base, followed by the fusion of the face

(more details on Table S6). Interestingly, this resonates with the pattern of sutural fusion in

alligator ontogeny, which cranial (i.e. frontoparietal) has the highest degree of suture closure

followed by skull base (i.e. basioccipital-exoccipital) and then the face (i.e. internasal)[69]

suggesting that the same mechanism may control topological changes in both ontogeny and

evolution.

**Avemetatarsalia**

Avemetatarsalian transition is marked with a faster ontogenetic bone growth in more derived taxa,

indicated by higher degree of vascularization, growth marks, and vascular canal arrangement

(reviewed by Bailleul[70]), more pneumatized skulls (reviewed by Gold[71]), and an increase in

complexity reminiscent of what is observed in crurotarsans. The basal ornithischian *Psittaosaurus*

*lujiatunensis* and basal saurischian *Eoraptor lunensis* are relatively close to each other on the

morphospace (Fig. 3), with the *Psittacosaurus* skull showing slightly more density because of

fused palatines, a trait which is also observed in extant crocodilians and some birds, and its extra

rostral bone as observed in other ceratopsians[72].

The basal sauropodomorph *Plateosaurus engelhardti* has the lowest clustering coefficient (i.e.

lower integration) of archosaurs, suggesting that skulls of sauropodmorphs are less integrated

than those of saurischians[31], accompanied by poorly connected bones (as seen in the network in

Fig. 4C). Poorly connected bones, for example epipterygoid, and some connections, such as the

ectopterygoid-jugal articulation, are later lost in neosauropods [43,73].

Within theropods, the ceratosaurian *Coelophysis* is more derived and has a slightly more complex

and specialized skull than the ceratosaurian *Dilophosaurus*[42]. Their positions on the morphospace

suggest that ceratosaurians occupy a region characterized by a higher mean path length (L), when

compared to other archosaurs (Fig. 3). *Compsognathus* is close to *Riojasuchus* on the

morphospace with a similar mean path length (Figs 3 and S4, Table S8), its facial bones are also

unfused, and it has a similar composition for its facial modules (see facial modules in

*Compsognathus* and nasal modules in *Riojasuchus* on Table S4 and Figure S10). These

observations suggest an ancestral facial topology (see Table S6 and S8 for more details) is

concomitant to the magnitude of shape change reported for compsognathids[34]. *Compsognathus*

possesses an independent postorbital that is absent from *Ichthyornis* to modern birds. It also has

an independent prefrontal that is absent in most Oviraptorsauria and Paraves[74], including *Citipati*,

*Velociraptor*, and from *Ichthyornis* to modern birds. Despite its ancestral features, the back of the

skull and the skull base of *Compsognathus* are fused, similarly to other Paravians and modern

birds.

The oviraptorid *Citipati* has a skull topology that occupies a morphospace within non-avian

theropods, despite its unique vertically-oriented premaxilla and short beak[34,75]. *Citipati* has an

independent epipterygoid that is also present in some non-avian theropods and ancestral

archosaurs, such as *Plateosaurus erlenbergiensis*, but which is absent in extant archosaurs[75–78].

*Citipati* also has fused skull roof (with fused interparietals), skull base, and face, marked with

fused internasal and the avian-like inter-premaxillary sutures.

Like other dromaeosaurids, *Velociraptor*'s eyes are positioned lateral to the rostrum. Its prefrontal

bone is either absent or fused with the lacrimal while it remains separate in other

dromaeosaurids[79–81]. We observed a loss of the prefrontals from *Citipati* to modern birds, but not

in more ancestral archosaurs or crurotarsans. Bones forming the *Velociraptor* basicranium, such

as basioccipital, and basisphenoid are fused with other members of the basicranium (listed in

Table S6). Despite having a similar number of bones and articulations to *Citipati*, the cranial

bones in *Velociraptor* are more integrated with each other and are more likely to connect to bones

with a different number of articulations (i.e. more disparity) (Table S8). Like *Compsognathus* and

other primitive non-avian dinosaurs, *Velociraptor* has an ancestral facial topology with separate

premaxilla, maxilla, and nasal bones.

**Archaeopteryx and Ichthyornis as intermediates between non-avian theropods and modern**

**birds**

The skull of *Archaeopteryx* occupied a region of the morphospace closer to non-avian dinosaurs

and crurotarsans than to juvenile birds (Fig. 3). The distance of *Archaeopteryx* from crown birds

and its proximity in the morphospace to *Velociraptor* and *Citipati* along the PC1 axis (Fig. 3)

may reflect the evolving relationship between cranial topology and endocranial volume. In fact,

*Archaeopteryx* has an endocranial volume which is intermediate between the ancestral non-avian

dinosaurs and crown birds[82,83] and it is within the plesiomorphic range of other non-avian

Paraves[84]. This makes *Archaoepteryx* closer to dromaeosaurid *Velociraptor* than to oviraptor

*Citipati*, for both its skull anatomy and its endocranial volume[84]. Modifications related to the

smaller endocranial volume in *Archaeopteryx* include the unfused bones in the braincase, the

independent reappearance of a separate prefrontal after the loss in Paraves[74], a separate left and

right premaxilla as observed in crocodilian snouts and ancestral dinosaurs, and the presence of

separate postorbitals, which might restrict the fitting for a larger brain[34].

Compared to *Archaeopteryx, Ichthyornis* is phylogenetically closer to modern birds and occupies

a region of the morphospace near the juvenile birds and extant crocodilians when adult birds are

included in the analysis (Fig. 3), but closer to extant crocodilians when all birds or when adult

birds are removed (Figs. S4-9). The proximity between *Ichthyornis* and juvenile birds may be

explained by the similar modular division (as observed in Figs. 4B and 4D; Table S4, Fig. S10),

presence of anatomical features characteristic of modern birds, such as the loss of the postorbital

bones, the fusion of the left and right premaxilla to form the beak, a bicondylar quadrate that form

a joint with the braincase, and the arrangement of the rostrum, jugal, and quadratojugal required

for a functional cranial kinetic system[58,85–88]. The proximity between *Ichthyornis* and extant

crocodilians in terms of complexity (Figs. S4-9, Table S8) may be explained by the fused frontal

and fused parietal, and separate maxilla, nasal, prootic and laterosphenoid (Table S6).

**Paleognath and neognath birds**

Juvenile birds have a skull roof with relatively less fused bones with the interfrontal, interparietal,

and frontoparietal sutures open, and a more fused skull base. Postorbital is already fused in all

juvenile birds (i.e. after hatching). Collectively, juvenile neognaths show a skull anatomy with a

fused cranial base, relatively less fused roof, and unfused face that resembles the anatomy of

ancestral non-avian theropods. Unlike what is observed in non-avian theropods, frontal, parietal,

nasal, premaxilla, and maxilla eventually fuse with the rest of the skull in adult modern birds.

However, in the palatal region not all the sutures are completely closed: the caudal ends of the

vomers remained unfused in adult *Nothura*, which is a characteristic common in Tinamidae[89]. A

similar pattern of suture closure has been described in another paleognath, the emu, in which the

sutures of the base of the skull close first and then the cranial and facial sutures close while

palatal sutures remain open[69]. The only difference is that in *Nothura*, where closure of major

cranial sutures (frontoparietal, interfrontal, and interparietal) happens after the facial sutures

closure.  In summary, when compared with neognaths, the skull of the paleognath *Nothura* is

more homogeneous and complex in both juvenile and adult stages. As the skull grows, its bones

fuse and both its complexity and heterogeneity increase.

Within the neognaths, the skull of *Geospiza fortis* is more complex and more homogenous than

*Gallus gallus* in both juvenile and adult stages: bones in *Geospiza* skull are more likely to connect

with bones with the same number of connections than *Gallus*. These two trajectories illustrate

how the connectivity of each bone diversifies and becomes more specialized within a skull as

sutures fuse together, as predicted by the Williston's law.

As in crurotarsans, major transitions in Avemetatarsalia are associated with the fusion first of the

skull base, then the skull roof, and, finally, with the face (more details on Table S6). This is more

similar to the temporal pattern of sutural closure during ontogeny in the emu (skull base first,

skull roof second, facial third) than to the one observed in the alligator (cranial first, skull base

second, facial third)[69], thus suggesting that the same mechanism for ontogeny may have been co-

opted in Avemetatarsalia evolution.

**Ontogenetic differences in topology between birds and crocodilians**

Our comparisons on network anatomy found that juvenile birds occupy a region of the

morphospace that is closer to the less derived archosaurs and crurotarsans than to that occupied

by adult modern birds (Fig. S1B). Juvenile birds have a degree of anisomerism of skull bones and

skull anatomical complexity closer to that in crurotarsans and non-avian dinosaurs, while their

pattern of integration overlaps with that of adult birds, crurotarsans, and non-avian dinosaurs.

These similarities in complexity and heterogeneity may be explained by the comparably higher

number and symmetrical spatial arrangements of circumorbital ossification centres in early

embryonic stages[74]. For example, both crown avians and *A. mississippiensis* have two ossification

centres that fuse into one for lacrimals[74,90]. Meanwhile, ossification centres that form the

prefrontal and postorbital, fuse in prenatal birds but remain separate in adult non-avian

dinosaurs[74,90,91]. These ossification centres later develop into different, but overlapping, number of

bones and their arrangement in juvenile birds (27 – 34 bones) and adult non-avian theropods (32

– 44 bones) with discrepancies explained by the heterochronic fusion of the ossification centres

(Table S8).

Following postnatal fusions and growth, modern bird skulls become more heterogeneous and

their bones more connected and topologically closer to each other (Figs. 3C and 5; Table S8).

This makes avian skull bones more diverse and functionally integrated. Simultaneously, skull

topology in birds diversifies with ontogeny within their lineage, as shown by the ontogenetic

trajectories of *Gallus*, *Nothura*, and *Geospiza* (Figs. 3C and 5). Thus, bones (1) develop from

ossification centres shared among crurotarsans and avemetatarsalians, (2) interact as modules

with heterogeneity and complexity similar to basal members at juvenile stage, and (3) then fuse

and diversify to produce skulls of adult birds.

The skulls of birds, crocodilians, and dinosaurs develop from ossification centres with

comparable spatial locations in the embryonic head[74]. When both evolutionary and ontogenetic

cranial shape variation was compared among crocodilians, Morris and colleagues showed that at

mid- to late embryonic stages, cranial shapes originated from a conserved region of skull shape

morphospace[92]. They suggested that crocodilian skull morphogenesis at early and late embryonic

stages are controlled by signaling molecules that are important in other amniotes as well, such as

*Bmp4*, *calmodulin*, *Sonic hedgehog* (*Shh*); and *Indian hedgehog*[92–99]. Then, from late prenatal

stages onward, snout of crocodilians narrows[100] and elongates following different ontogenetic

trajectories to give the full spectrum of crocodilian cranial diversity[92].

Another major transformation in archosaurian evolution is the origin of skulls of early and

modern birds from the snouted theropods. This transition involved two significant heterochronic

shifts[34,101]. First, avians evolved highly paedomorphic skull shapes compared to their ancestors by

developmental truncation[34]. This was followed, by a peramorphic shift where primitively paired

premaxillary bones fused and the resulting beak bone elongated to occupy much of the new avian

face [101]. By comparison, the skull of *Alligator* undergoes extensive morphological change and

closing of the interfrontal and interparietal sutures during embryogenesis is followed by the

prolonged postnatal and maturation periods, with the lack of suture closure and even widening of

some sutures[102,103]. Bailleul and colleagues suggested that mechanisms that inhibit suture closure,

rather than bone resorption, cause the alligator sutures to remain open during ontogeny[103].

Nevertheless, juvenile and adult alligators share the same cranial topology featuring similar

module compositions and both occupy a region of morphospace close to *Crocodylus* (Figs. 4D

and S10; Table S4 and S8). Such topological arrangement suggests that conserved molecular,

cellular, and developmental genetic processes underlie skull composition and topology observed

across crocodilians. Likewise, oviraptorid dinosaurs, as represented by *Citipati,* display their own

unique skull shape and ontogenetic transformation[34], while retaining a topology conserved with

other theropods. Combined, this evidence suggests that developmental mechanisms controlling

skull composition and interaction among skull elements are conserved among theropods.

The process of osteogenesis underlies the shape and topology of the bony skull. In chicken

embryo, inhibition of FGF and WNT signaling pathways prevented fusion of the suture that

separates the left and right premaxilla, disconnected the premaxilla-palatine articulation and

changed their shapes giving the distal face a primitive snout-like appearance[101]. The site of bone

fusion in experimental unfused, snout-like chicken premaxillae showed reduced expression of

skeletal markers *Runx2*, *Osteopontin*, and the osteogenic marker *Col I*[101], implying localized

molecular mechanisms regulating suture closure and shape of individual cranial bones. Thus,

changes in gene expression during craniofacial patterning in avians[95,96,98,104–106], non-avian

dinosaurs, and crocodilians[92,101] contribute to the clade-specific differences in skull anatomical

organization resulting from the similar patterns of bone fusion of bones.

Finally, we observe some network modules where some bones within the same modules in

juveniles will later fuse in adult birds, but not in *A. mississippiensis* (Supplementary Information

5; Figs. 4E and S10, Table S4). For example, in *Nothura*, premaxilla, nasal, parasphenoid,

pterygoid, vomer, and maxilla grouped in the same juvenile module will later fuse during

formation of the upper beak in the adult. In *A. mississippiensis*, premaxilla, maxilla, nasal,

lacrimal, prefrontal, jugal, frontal, and ectopterygoid are also in the same juvenile module, but

remain separate structures in adult. These findings suggest that bones within the same module

may be more likely to fuse together in ontogeny but doing so is a lineage-specific feature.

Comparisons of juveniles and adults for extant birds and the alligator revealed ontogenetic

changes linked to the evolution of the skull organization in archosaurs. Whereas the anatomical

organization of the skull of juvenile alligators resembles that of adults, the anatomy of juvenile

modern birds is closer to that of non-avian dinosaurs than to that of adult avians of the same

species in terms of morphological complexity and anisomerism, probably due to the spatial

arrangements of ossification centres at embryonic stages[74,90,91]. More specifically, the differences

in skull organization between crown birds and non-avian dinosaurs could be explained by

postnatal fusion of bones.

**CONCLUSION**

A network-based comparison of the cranial anatomy of archosaurs shows that differences within

and among archosaurian clades are associated with an increase of anatomical complexity, a

reduction in number of bones (as predicted by the Williston's Law), and an increase of

anisomerism marked by bone fusion, for both crurotarsans and avemetatarsalians. Our findings

indicate that the anatomical organization of the skull is controlled by developmental mechanisms

that diversified across and within each lineage: heterotopic changes in craniofacial patterning

genes, heterochronic prenatal fusion of ossification centres[74,90,91], and lineage-specific postnatal

fusion of sutures. Some of these mechanisms have been shown to be conserved in other tetrapods.

For example, heterotopy of craniofacial patterning genes also took place between chick and mice

embryos[95,96,106]. Hu and Marcucio showed that mouse frontonasal ectodermal zone could alter the

development of the avian frontonasal process, suggesting a conserved mechanism for frontonasal

development in vertebrates[96]. Our findings illustrate how a comparative analysis of the

anatomical organization of the skull can reveal both common and disparate patterns and processes

determining skull evolution in vertebrates.

**ACKNOWLEDGEMENT**

We thank Jake Horton for coding the adult and juvenile matrices for *Alligator mississippiensis*

and *Crocodylus moreletii*, Patrick Campbell of Natural History Museum London for providing

reptile specimens, Alfie Gleeson and Digimorph for CT scans of crocodiles, and staff from

Natural History Museum library for literature search. BE-A has received financial support

through the Postdoctoral Junior Leader Fellowship Programme from "la Caixa" Banking

Foundation (LCF/BQ/LI18/11630002) and also thanks the Unidad de Excelencia María de

Maeztu funded by the AEI (CEX2018-000792-M). HWL's Master Thesis that inspired this

project was funded by Imperial College London and Natural History Museum, London.

**AUTHOR CONTRIBUTION**

HWL, BE-A, AA designed the study.

HWL coded network models.

HWL and BE-A wrote the R scripts and performed the analyses.

All authors discussed the results and wrote the manuscript.

**Conflict of Interest:** The authors declare no conflict of interest.

**Data Availability**

Data and R code are available at https://figshare.com/s/80714fb9a06e886cd412.

**REFERENCES**

1.  Gauthier, J. Saurischian monophyly and the origin of birds. *Mem. Calif. Acad. Sci.* **8**, 1–55 (1986).

2.  Benton, M. &Clark, J. Archosaur phylogeny and the relationships of the Crocodylia. in *The Phylogeny and Classification of the Tetrapods: Systematics Association Special* 295–338 (1988).

3.  Sereno, P. C. Basal archosaurs: phylogenetic relationships and functional implications. *Soc. Vertebr. Paleontol. Mem.* **2**, 1–53 (1991).

4.  Juul, L. The phylogeny of basal archosaurs. *Palaeontol. Africana* **31**, 1–38 (1994).

5.  Benton, M. J. Scleromochlus taylori and the origin of dinosaurs and pterosaurs. *Philos. Trans. R. Soc. B Biol. Sci.* **354**, 1423–1446 (1999).

6.  Benton, M. J. Origin and Relationships of Dinosauria. in *The Dinosauria* (eds. Weishampel, D. B., Dodson, P. &Osmólska, H.) 7–19 (University of California Press, 2004). doi:10.1525/california/9780520242098.003.0005

7.  Irmis, R. B., Parker, W. G., Nesbitt, S. J. &Liu, J. Early ornithischian dinosaurs: The triassic record. *Hist. Biol.* **19**, 3–22 (2007).

8.  Brusatte, S. L., Benton, M. J., Lloyd, G. T., Ruta, M. &Wang, S. C. Macroevolutionary patterns in the evolutionary radiation of archosaurs (Tetrapoda: Diapsida). *Earth Environ. Sci. Trans. R. Soc. Edinburgh* **101**, 367–382 (2010).

9.  Sadleir, R. &Makovicky, P. Cranial shape and correlated characters in crocodile evolution. *J Evol. Biol.* **21**, 1578–1596 (2008).

10. Goswami, A., Weisbecker, V. &Sánchez-Villagra, M. Developmental modularity and the marsupial-placental dichotomy. *J Exp Zool. B Mol Dev Evol* **312B**, 186–195 (2009).

11. Hallgrímsson, B. *et al.* Deciphering the palimpsest: Studying the relationship between morphological integration and phenotypic covariation. *Evol Biol* **36**, 355–376 (2009).

12.    Felice, R. N. &Goswami, A. Developmental origins of mosaic evolution in the avian

cranium. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 555–560 (2018).

13.    Felice, R. N., Tobias, J. A., Pigot, A. L. &Goswami, A. Dietary niche and the evolution of

cranial morphology in birds. *Proc. R. Soc. B Biol. Sci.* **286**, (2019).

14.    Klingenberg, C. P. Morphological Integration and Developmental Modularity. *Annu. Rev.*

*Ecol. Evol. Syst.* **39**, 115–132 (2008).

15.    Eble, G. Morphological modularity and macroevolution: conceptual and empirical aspects.

in *Modularity. Understanding the development and evolution of natural complex systems*

(eds. Callebaut, W. &Rasskin☐Gutman, D.) 221–238 (MIT Press, 2005).

16.    Olson, E. &Miller, R. *Morphological Integration*. (University of Chicago Press, 1958).

doi:10.2307/2405966

17.    Wagner, G. P. &Altenberg, L. Complex adaptations and the evolution of evolvability.

*Evolution (N. Y).* **50**, 967–976 (1996).

18.    Wagner, G., Pavlicev, M. &Cheverud, J. The road to modularity. *Nat. Rev. Genet.* **8**, 921–

931 (2007).

19.    Esteve-Altava, B., Marugán-Lobón, J., Botella, H. &Rasskin-Gutman, D. Network Models

in Anatomical Systems. *J. Anthropol. Sci.* **89**, 175–184 (2011).

20.    Esteve-Altava, B. Challenges in identifying and interpreting organizational modules in

morphology. *J. Morphol.* **278**, 960–974 (2017).

21.    Chernoff, B. &Magwene, P. Afterword. in *Morphological Integration: Forty years later.*

319–353 (University of Chicago Press, 1999).

22.    Esteve-Altava, B., Marugán-Lobón, J., Botella, H., Bastir, M. &Rasskin-Gutman, D. Grist

for Riedl's mill: A network model perspective on the integration and modularity of the

human skull. *J. Exp. Zool. Part B Mol. Dev. Evol.* **320**, 489–500 (2013).

23.    Rasskin-Gutman, D. &Esteve-Altava, B. Concept of Burden in Evo-Devo. in *Evolutionary*

*Developmental Biology* (eds. Nuño de la Rosa, L. &Müller, G.) 1–11 (Springer, Cham,

2018). doi:10.1007/978-3-319-33038-9_48-1

24.    Gregory, W. K. 'Williston's law' relating to the evolution of skull bones in the vertebrates.

581            *Am. J. Phys. Anthropol.* **20**, 123–152 (1935).

582     25.    Esteve-Altava, B., Marugán-Lobón, J., Botella, H. &Rasskin-Gutman, D. Structural

583            Constraints in the Evolution of the Tetrapod Skull Complexity: Williston's Law Revisited

584            Using Network Models. *Evol. Biol.* **40**, 209–219 (2013).

585     26.    Esteve-Altava, B., Marugán-Lobón, J., Botella, H. &Rasskin-Gutman, D. Random Loss

586            and Selective Fusion of Bones Originate Morphological Complexity Trends in Tetrapod

587            Skull Networks. *Evol. Biol.* **41**, 52–61 (2014).

588     27.    Esteve-Altava, B., Boughner, J. C., Diogo, R., Villmoare, B. A. &Rasskin-Gutman, D.

589            Anatomical Network Analysis Shows Decoupling of Modular Lability and Complexity in

590            the Evolution of the Primate Skull. *PLoS One* **10**, e0127653 (2015).

591     28.    Esteve-Altava, B. *et al.* Evolutionary parallelisms of pectoral and pelvic network-anatomy

592            from fins to limbs. *Sci. Adv.* **5**, (2019).

593     29.    Werneburg, I., Esteve-Altava, B., Bruno, J., Torres Ladeira, M. &Diogo, R. Unique skull

594            network complexity of Tyrannosaurus rex among land vertebrates. *Sci. Rep.* **9**, 1–14

595            (2019).

596     30.    Plateau, O. &Foth, C. Birds have peramorphic skulls, too: anatomical network analyses

597            reveal oppositional heterochronies in avian skull evolution. *Commun. Biol.* **3**, 1–12 (2020).

598     31.    Rasskin-Gutman, D. &Esteve-Altava, B. Connecting the Dots: Anatomical Network

599            Analysis in Morphological EvoDevo. *Biol. Theory* **9**, 178–193 (2014).

600     32.    Esteve-Altava, B. &Rasskin-Gutman, D. Anatomical Network Analysis in Evo-Devo. in

601            *Evolutionary Developmental Biology* 1–19 (Springer International Publishing, 2018).

602            doi:10.1007/978-3-319-33038-9_57-1

603     33.    Gregory, W. K. Polyisomerism and Anisomerism in Cranial and Dental Evolution among

604            Vertebrates. *Proc. Natl. Acad. Sci.* **20**, 1–9 (1934).

34.    Bhullar, B. A. S. *et al.* Birds have paedomorphic dinosaur skulls. *Nature* **487**, 223–226

(2012).

35.    Brusatte, S., Benton, M., Desojo, J. &Langer, M. The Higher-Level Phylogeny of

Archosauria (Tetrapoda: Diapsida). *J. Syst. Palaeontol.* **8**, 3–47 (2010).

36.    Galton, P. M. &Upchurch, P. Prosauropoda. in *The Dinosauria* (eds. Weishampel, D. B.,

Dodson, P. &Osmólska, H.) 232–258 (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

37.    Hailu, Y. &Dodson, P. Basal Ceratopsia. in *The Dinosauria* (eds. Weishampel, D. B.,

Dodson, P. &Osmólska, H.) 325–334 (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

38.    Holtz, T. R. J. &Osmólska, H. Saurischia. in *The Dinosauria* (eds. Weishampel, D. B.,

Dodson, P. &Osmólska, H.) 21–24 (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

39.    Nesbitt, S. J. The Early Evolution of Archosaurs: Relationships and the Origin of Major

Clades. *Bull. Am. Museum Nat. Hist.* **352**, 1–292 (2011).

40.    Norell, M. A. &Makovicky, P. J. Dromaeosauridae. in *The Dinosauria* (eds. Weishampel,

D. B., Dodson, P. &Osmólska, H.) 196–209 (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

41.    Padian, K. Basal Avialae. in *The Dinosauria* (eds. Weishampel, D. B., Dodson, P.

&Osmólska, H.) 210–231 (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

42.    Tykoski, R. S. &Rowe, T. Ceratosauria. in *The Dinosauria* (eds. Weishampel, D. B.,

Dodson, P. &Osmólska, H.) 47–70 (University of California Press., 2004).

doi:10.1525/california/9780520242098.003.0005

43.    Upchurch, P., Barrett, P. M. &Dodson, P. Sauropoda. in *The Dinosauria* (eds.

Weishampel, D. B., Dodson, P. &Osmólska, H.) 259–322 (University of California Press,

2004). doi:10.1525/california/9780520242098.003.0005

44.    Holtz, T. R. J. &Osmólska, H. Dinosaur distribution and biology. in *The Dinosauria* (eds.

Weishampel, D. B., Dodson, P. &Osmólska, H.) (University of California Press, 2004).

doi:10.1525/california/9780520242098.003.0005

45.    Bapst, D. W. paleotree: an R package for paleontological and phylogenetic analyses of

evolution. (2012).

46.    Brusatte, S. L., Benton, M. J., Ruta, M. &Lloyd, G. T. Superiority, competition, and

opportunism in the evolutionary radiation of dinosaurs. *Science (80-. ).* **321**, 1485–1488

(2008).

47.    Lloyd, G. T., Wang, S. C. &Brusatte, S. L. Identifying heterogeneity in rates of

morphological evolution: Discrete character change in the evolution of lungfish

(Sarcopterygii; Dipnoi). *Evol. Int. J. Org. Evol.* **66**, 330–348 (2012).

48.    Benton, M. J. &Donoghue, P. C. J. Paleontological evidence to date the tree of life. *Mol.*

*Biol. Evol.* **24**, 26–53 (2007).

49.    Varela, S., Hernández, J. G. &Sgarbi, L. F. paleobioDB: Download and Process Data from

the Paleobiology Database. (2019).

50.    R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing. (2018).

51.    Csardi, G. &Nepusz, T. The igraph software package for complex network research,

InterJournal, Complex Systems 1695. (2006).

52.    Almende, B., Thieurmel, B. &Robert, T. visNetwork: Network Visualization using 'vis.js'

Library. (2019).

53.    Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of

biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

54.    Esteve-Altava, B. A Node-based Informed Modularity Strategy to Identify Organizational

Modules in Anatomical Networks. *bioRxiv* 2020.07.06.189175 (2020).

doi:10.1101/2020.07.06.189175

55.    Wang, M., Zhao, Y. &Zhang, B. Efficient Test and Visualization of Multi-Set

Intersections. *Sci. Rep.* **5**, 16923 (2015).

56.    Sidor, C. Simplification as a trend in synapsid cranial evolution. *Evolution (N. Y).* **55**,

1419–1442 (2001).

57.    McShea, D. &Hordijk, W. Complexity by Subtraction. *Evol Biol2* **40**, 504–520 (2013).

58.    Bhullar, B. A. S. *et al.* How to make a bird skull: Major transitions in the evolution of the

avian cranium, paedomorphosis, and the beak as a surrogate hand. *Integr. Comp. Biol.* **56**,

389–403 (2016).

59.    vonBaczko, M. B. &Desojo, J. B. Cranial anatomy and palaeoneurology of the archosaur

riojasuchus tenuisceps from the los colorados formation, La Rioja, Argentina. *PLoS One*

**11**, (2016).

60.    Felice, R. N. *et al.* Evolutionary integration and modularity in the archosaur cranium.

*Integr. Comp. Biol.* (2019). doi:10.1093/icb/icz052

61.    Small, B. J. The Triassic Thecodontian Reptile Desmatosuchus: Osteology and

Relationships. (Texas Tech University, 1985).

62.    Small, B. J. Cranial anatomy of Desmatosuchus Haplocerus (Reptilia: Archosauria:

Stagonolepididae). *Zool. J. Linn. Soc.* **136**, 97–111 (2002).

63.    Schoch, R. R. Osteology of the small archosaur Aetosaurus from the upper Triassic of

Germany. *Neues Jahrb. fur Geol. und Palaeontologie - Abhandlungen* **246**, 1–35 (2007).

64.    Walker, A. D. A revision of Sphenosuchus acutus Haughton, a crocodylomorph reptile

from the Elliot Formation (late Triassic or early Jurassic) of South Africa. *Phil. Trans. R.*

*Soc. Lond. B.* **330**, 1–120 (1990).

65.    Wu, X.-C. &Chatterjee, S. Dibothrosuchus elaphros , a Crocodylomorph from the Lower

Jurassic of China and the Phylogeny of the Sphenosuchina Xiao-Chun Wu and Sankar

Chatterjee. *J. Vertebr. Paleontol.* **13**, 58–89 (1993).

66.    Fernández, M. &Gasparini, Z. Salt glands in the Jurassic metriorhynchid Geosaurus:

684           implications for the evolution of osmoregulation in Mesozoic marine crocodyliforms.

685           *Naturwissenschaften* **95**, 79–84 (2008).

686    67.    Gasparini, Z., Pol, D. &Spalletti, L. A. An unusual marine crocodyliform from the

687           jurassic-cretaceous boundary of Patagonia. *Science (80-. ).* **311**, 70–73 (2006).

688    68.    Pol, D. &Gasparini, Z. Skull anatomy of dakosaurus andiniensis (thalattosuchia:

689           Crocodylomorpha) and the phylogenetic position of thalattosuchia. *J. Syst. Palaeontol.* **7**,

690           163–197 (2009).

691    69.    Bailleul, A. M., Scannella, J. B., Horner, J. R. &Evans, D. C. Fusion patterns in the skulls

692           of modern archosaurs reveal that sutures are ambiguous maturity indicators for the

693           Dinosauria. *PLoS One* **11**, 1–26 (2016).

694    70.    Bailleul, A. M., O'Connor, J. &Schweitzer, M. H. Dinosaur paleohistology: Review,

695           trends and new avenues of investigation. *PeerJ* **2019**, 1–45 (2019).

696    71.    Gold, M. E. L., Brusatte, S. L. &Norell, M. A.  The Cranial Pneumatic Sinuses of the

697           Tyrannosaurid Alioramus (Dinosauria: Theropoda) and the Evolution of Cranial

698           Pneumaticity in Theropod Dinosaurs . *Am. Museum Novit.* **3790**, 1–46 (2013).

699    72.    Sereno, P. C., Xijin, Z. &Lin, T. A new psittacosaur from inner mongolia and the parrot-

700           like structure and function of the psittacosaur skull. *Proc. R. Soc. B Biol. Sci.* **277**, 199–

701           209 (2010).

702    73.    Button, D. J., Barrett, P. M. &Rayfield, E. J. Comparative cranial myology and

703           biomechanics of Plateosaurus and Camarasaurus and evolution of the sauropod feeding

704           apparatus. *Palaeontology* **59**, 887–913 (2016).

705    74.    Smith-Paredes, D. *et al.* Dinosaur ossification centres in embryonic birds uncover

706           developmental evolution of the skull. *Nat. Ecol. Evol.* **2**, 1966–1973 (2018).

707    75.    Norell, M. A., Clark, J. M. &Chiappe, L. M. An embryo of an oviraptorid (Dinosauria:

708           Theropoda) from the Late Cretaceous of Ukhaa Tolgod, Mongolia. *Am. Museum Novit.*

**3315**, 1–17 (2001).

76.   deBeer, G. R. *The Development of the Vertebrate Skull*. (Oxford University Press, 1937).

77.   Gauthier, J., Kluge, A. G. &Rowe, T. Amniote phylogeny and the importance of fossils.

*Cladistics* **4**, 105–209 (1988).

78.   Clark, J. M., Norell, M. A. &Rowe, T. Cranial Anatomy of Citipati osmolskae (Theropoda,

Oviraptorosauria), and a Reinterpretation of the Holotype of Oviraptor philoceratops. *Am.*

*Museum Novit.* **3364**, 1–24 (2002).

79.   Norell, M. A. *et al.* A theropod dinosaur embryo and the affinities of the Flaming Cliffs

dinosaur eggs. *Science (80-. ).* **266**, 779–792 (1994).

80.   Barsbold, R. &Osmolska, H. The skull of Velociraptor (Theropoda) from the Late

Cretaceous of Mongolia. *Acta Palaeontol. Pol.* **442**, 189–219 (1999).

81.   Currie, P. &Dong, Z. New information on Cretaceous troodontids (Dinosauria, Theropoda)

from the People's Republic of China. *Can. J Earth Sci* **38**, 1753–1766 (2001).

82.   Larsson, H. C. E., Sereno, P. C. &Wilson, J. A. Forebrain Enlargemant among Nonavian

Theropod Dinosaurs. *J. Vertebr. Paleontol.* **20**, 615–618 (2000).

83.   Domínguez Alonzo, P., Milner, A. C., Ketcham, R. A., Cookson, M. J. &Rowe, T. B. The

avian nature of the brain and inner ear of Archaeopteryx. *Nature* **430**, 666–669 (2004).

84.   Balanoff, A. M., Bever, G. S., Rowe, T. B. &Norell, M. A. Evolutionary origins of the

avian brain. *Nature* **501**, 93–96 (2013).

85.   Jollie, M. T. The head skeleton of the chicken and remarks on the anatomy of this region

in other birds. *J. Morphol.* **100**, 389–436 (1957).

86.   Bock, W. J. Kinetics of the avian skull. *J. Morphol.* **114**, 1–41 (1964).

87.   Clarke, J. Morphology, phylogenetic taxonomy, and sys- tematics of Icthyornis and

Apatornis (Avialae, Ornithurae). *Bull Am Museum Nat Hist.* **286**, 1–179 (2004).

88.   Field, D. J. *et al.* Complete Ichthyornis skull illuminates mosaic assembly of the avian

head. *Nature* **557**, 96–100 (2018).

89.    Silveira, L. F. &Höfling, E. Cranial osteology in Tinamidae (Birds: Tinamiformes), with

systematic considerations. *Bol. Mus. Para. Emílio Goeldi. Ciências Naturais, Belém* **2**,

15–54 (2007).

90.    Rieppel, O. Studies on skeleton formation in reptiles. v. Patterns of ossification in the

skeleton of Alligator mississippiensis DAUDIN (Reptilia, Crocodylia). *Zool. J. Linn. Soc.*

**109**, 301–325 (1993).

91.    Maxwell, E. &Larson, H. Comparative ossification sequence and skeletal development of

the postcranium of palaeognathous birds (Aves: Palaeognathae). *Zool. J Linn. Soc.* **157**,

169–196 (2009).

92.    Morris, Z. S., Vliet, K. A., Abzhanov, A. &Pierce, S. E. Heterochronic shifts and

conserved embryonic shape underlie crocodylian craniofacial disparity and convergence.

*Proc. R. Soc. B Biol. Sci.* **286**, 20182389 (2019).

93.    Abzhanov, A., Protas, M., Grant, B. R., Grant, P. R. &Tabin, C. J. Bmp4 and

Morphological Variation of Beaks in Darwin's Finches. *Science (80-. ).* **305**, 1462–1465

(2004).

94.    Abzhanov, A. *et al.* The calmodulin pathway and evolution of elongated beak morphology

in Darwin's finches. *Nature* **442**, 563–567 (2006).

95.    Hu, D. &Marcucio, R. S. A SHH-responsive signaling center in the forebrain regulates

craniofacial morphogenesis via the facial ectoderm. *Development* **136**, 107–116 (2009).

96.    Hu, D. &Marcucio, R. S. Unique organization of the frontonasal ectodermal zone in birds

and mammals. *Dev. Biol.* **325**, 200–210 (2009).

97.    Mallarino, R. *et al.* Two developmental modules establish 3D beak-shape variation in

Darwin's finches. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4057–4062 (2011).

98.    Hu, D. &Marcucio, R. S. Neural crest cells pattern the surface cephalic ectoderm during

FEZ formation. *Dev. Dyn.* **241**, 732–740 (2012).

99.    Ahi, E. P. Signalling pathways in trophic skeletal development and morphogenesis:

Insights from studies on teleost fish. *Dev. Biol.* **420**, 11–31 (2016).

100.   Watanabe, A. &Slice, D. E. The utility of cranial ontogeny for phylogenetic inference: A

case study in crocodylians using geometric morphometrics. *J. Evol. Biol.* **27**, 1078–1092

(2014).

101.   Bhullar, B. A. S. *et al.* A molecular mechanism for the origin of a key evolutionary

innovation, the bird beak and palate, revealed by an integrative approach to major

transitions in vertebrate history. *Evolution (N. Y).* **69**, 1665–1677 (2015).

102.   Padian, K., deRicqles, A. J. &Horner, J. R. Dinosaurian growth rates and bird origins.

*Nature* **412**, 405–408 (2001).

103.   Bailleul, A. M. &Horner, J. R. Comparative histology of some craniofacial sutures and

skull-base synchondroses in non-avian dinosaurs and their extant phylogenetic bracket. *J.*

*Anat.* **229**, 252–285 (2016).

104.   Hu, D., Marcucio, R. S. &Helms, J. A. A zone of frontonasal ectoderm regulates

patterning and growth in the face. *Development* **130**, 1749–1758 (2003).

105.   Abzhanov, A., Cordero, D. R., Sen, J., Tabin, C. J. &Helms, J. A. Cross-regulatory

interactions between Fgf8 and Shh in the avian frontonasal prominence. *Congenit. Anom.*

*(Kyoto).* **47**, 136–148 (2007).

106.   Brugmann, S. A. *et al.* Wnt signaling mediates regional specification in the vertebrate face.

*Development* **134**, 3283–3295 (2007).

107.   Prieto-Márquez, A. &Norell, M. A. Redescription of a Nearly Complete Skull of

Plateosaurus (Dinosauria: Sauropodomorpha) from the Late Triassic of Trossingen

(Germany). *Am. Museum Novit.* **3727**, 1–58 (2011).

**SUPPLEMENTARY MATERIALS**

Table S1. Variance distribution across principal components

Table S2. First and last occurrence dates used to calibrate phylogenetic tree

Table S3. Internal nodes used for the phylogenetic tree

Table S4. Composition of modules for each taxon

Table S5. Categories of archosaurs based on capabilities of flight

Table S6. List of major fusion of bones with other bones in archosaurs

Table S7. Variation explained by each parameter

Table S8. Topological network parameters measured for each taxon

Table S9. Network parameters categorized by diet

Table S10. Number of modules.

Fig. S1. First two PC of topological parameters for all taxa.

Fig. S2. Second and third PC of topological parameters for all taxa.

Fig. S3. First and third PC of topological parameters for all taxa.

Fig. S4. First two PC of topological parameters for all taxa excluding avians.

Fig. S5. Second and third PC of topological parameters for all taxa excluding avians.

Fig. S6. First and third PC of topological parameters for all taxa excluding avians.

Fig. S7. First two PC of topological parameters for all taxa excluding adult avians.

Fig. S8. Second and third PC of topological parameters for all taxa excluding adult avians.

Fig. S9. First and third PC of topological parameters for all taxa excluding adult avians.

Fig. S10. Node-based modules of archosaurs based on details listed on Table S4.

Supplementary Information 1 References and notes about the specimens used.

Supplementary Information 2 Comparison between network-modules and variational modules in

archosaurs.

Supplementary Information 3 Comparison of network parameters among Aves, Crurotarsi, and

non-avian Dinosauria.

Supplementary Information 4 Comparison based on diet.

Supplementary Information 5 Comparison of juvenile avian modules with adult avian bones.

Supplementary Information 6 Supplementary Reference

**FIGURE LEGENDS**

**Figure 1. Anatomical network models**. Example of the network models for three archosaurian

skulls: (A) *Aetosaurus* from Schoch (2007)[63]; (B) *Plateosaurus* from Prieto-Marquez & Norell

(2011)[107]; (C) *Gallus* from Digimorph. The pair-wise articulations among the bones of skulls (left)

are formalized as network models (middle) and later analyzed, for example, to identify the skull

anatomical node-based modules (right). See methods for details.

**Figure 2. Phylogenetic framework**. A phylogenetic tree was created based on the evolutionary

relations among taxa as detailed in previous work [34–43]. Bifurcation times were calibrated based

on fossil dates from Benton and Donoghue[48] using the equal method in the paleotree package[45–47].

First and last occurrences were from Paleobiology Database (details listed in Table S2).

Silhouettes were from Phylopic.org. See methods for details.

**Figure 3. Principal components decomposition of topological variables.** (A) Skull distribution

for each taxon (see labels below). (B) Comparison of Avialae versus non-Avialae shows that non-

Avialae occupy part of the Avialae morphospace. (C) Comparison of Neornithes versus non-

Neornithes shows that non-Neornithes overlap with part of the Neornithes morphospace. Orange

dotted arrows show the ontogenetic change in modern birds from juvenile stage to adult stage. (D)

Comparison of Aves, Crurotarsi, and Dinosauria shows that they occupied different morphospace.

Ellipses show a normal distribution confidence interval around groups for comparison. Labels: N,

Number of nodes; K, Number of links; D, Density of Connection; C, Mean clustering coefficient;

H, Heterogeneity of connection; L, Mean path length; A, Assortativity of connection; P,

Parcellation. Aeto, *Aetosaurus;* AllA, adult *Alligator*; AllJ, juvenile *Alligator*; Arcx,

*Archaeopteryx*; Citi, *Citipati*; Coel, *Coelophysis*; Comp, *Compsognathus*; Croc, *Crocodylus*;

Dako, *Dakosaurus*; Desm, *Desmatosuchus*; Dibo, *Dibothrosuchus*; Dilo, *Dilophosaurus*; Eora,

*Eoraptor*; GalA, adult *Gallus*; GalJ, juvenile *Gallus*; GeoA, adult *Geospiza*; GeoJ, juvenile

*Geospiza*; Icht, *Ichthyornis*; NotA, adult *Nothura*; NotJ, juvenile *Nothura*; Plat, *Plateosaurus*;

Psit, *Psittacosaurus*; Rioj, *Riojasuchus*; Sphe, *Sphenosuchus*; Velo, *Velociraptor.* Silhouettes

were from Phylopic.org.

**Figure 4. Visualizations of the module composition changes across phylogeny**. The number of

node-based modules ranged from 1 to 8. (C) shows the difference in module composition among

the ornithischian *Psittacosaurus*, the basal saurischian *Eoraptor*, and the sauropodomorph

*Plateosaurus*. (D) Comparisons of the adult and juvenile stages of extant species. Adult *Nothura*

and *Geospiza* are shaded in grey as one module was identified because of the small number of

nodes and links due to a highly fused skull. Nodes were colored based on their modules.

Composition of each module is listed in Supplementary Table 4.

**Figure 5. Overview of the evolution of archosaurian skull topology:** Modern birds and few

non-avian dinosaurs have more heterogeneous connections than crurotarsans; extant taxa have

fewer bones and articulations than the extinct ones; bones in juvenile modern birds fuse and

produce a more densely connected adult skull. Modules and networks of the following taxa are

shown: (1) *Gallus*, (2) juvenile *Gallus*, (3) *Plateosaurus*, (4) *Dilophosaurus*, (5) *Aetosaurus*, (6)

adult *Alligator*. Morphospace of Aves is significantly different from Crurotarsi and Dinosauria

when adult birds are included. Orange arrows show the ontogenetic changes from juvenile to

adult stages in neornithes. Taxa on the left side of the biplot have higher density and fewer bones,

such as *Gallus* and *Alligator*, than taxa on the right, such as *Aetosaurus* and *Dilophosaurus*.