The costs of competition: high social status males experience accelerated epigenetic aging in wild baboons

Jordan A. Anderson[1]†, Rachel A. Johnston[1]†, Amanda J. Lea[2,3,4], Fernando A. Campos[2,5], Tawni N. Voyles[1], Mercy Y. Akinyi[6], Susan C. Alberts[1,2,6], Elizabeth A. Archie[6,7], Jenny Tung[1,2,6,8,*]

[1]Department of Evolutionary Anthropology, Duke University, Durham, North Carolina 27708, USA
[2]Department of Biology, Duke University, Durham, North Carolina 27708, USA
[3]Lewis-Sigler Institute for Integrative Genomics, Carl Icahn Laboratory, Princeton University, Princeton, NJ 08544, USA
[4]Department of Ecology and Evolution, Princeton University, Princeton, NJ 08544, USA
[5]Department of Anthropology, University of Texas at San Antonio, San Antonio, TX 78249, USA
[6]Institute of Primate Research, National Museums of Kenya, Nairobi 00502, Kenya
[7]Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556, USA
[8]Duke Population Research Institute, Duke University, Durham, NC 27708, USA

†Authors contributed equally to this work
*Correspondence: jt5@duke.edu

Summary

Aging, for virtually all life, is inescapable. However, within species and populations, rates of biological aging (i.e., physical decline with age) vary across individuals. Understanding sources of variation in biological aging is therefore central to understanding the biodemography of natural populations. Here, we constructed a DNA methylation-based predictor of chronological age for a population of wild baboons in which behavioral, ecological, and life history data have been collected for almost 50 years (N = 277 blood samples from 245 individuals, including 30 who were longitudinally sampled). Consistent with findings in humans and model organisms [1-4], DNA methylation patterns exhibit a strong, clock-like association with chronological age, but individuals are often predicted to be somewhat older or younger than their known age. However, the two most robust predictors of lifespan described for this population—cumulative early adversity and social bond strength—do not explain this deviation. Instead, the single most predictive factor is male dominance rank: high-ranking males are predicted to be biologically older than their true chronological age, such that alpha males appear to be nearly a year older than their known age. Longitudinal sampling indicates that males who climb the social hierarchy subsequently look epigenetically "older," likely reflecting the high energetic costs of rank attainment and maintenance in male baboons. Together, our results indicate that environmental effects on survival and epigenetic age can be disjunct, and that achieving high rank for male baboons—the best predictor of reproductive success—imposes physiological costs consistent with a "live fast, die young" life history strategy.

Keywords: Epigenetic age, epigenetic clock, aging, DNA methylation, dominance rank, baboons

Results and discussion

We used a combination of previously published [5] and newly generated reduced-representation bisulfite sequencing (RRBS) data from 245 wild baboons (N = 277 blood samples) living in the Amboseli ecosystem of Kenya [6] to generate a DNA methylation-based age predictor (an "epigenetic clock:" [1, 2]). Starting with a data set of methylation levels for 458,504 CpG sites genome-wide (Figure S1; Table S1), we used elastic net regression to identify a set of 593 CpG sites that accurately predict baboon age to within a median absolute difference (MAD) of 1.1 years (Pearson's r = 0.762, p < 10^{-53}; Table S2; median adult life expectancy in this population is 10.3 years for females and 7.94 for males [7]). Because the clock was significantly more accurate in males (N = 135; MAD = 0.9 years; Pearson's r = 0.86, p < 10^{-40}) than in females (N = 142; MAD = 1.6 years; r = 0.78, p < 10^{-29}; two-sided Wilcoxon test for differences in absolute error by sex: p = 4.35 x 10^{-9}), we separated males and females for all subsequent analyses (Figure 1A and 1B).

Overall, the clock performed favorably relative to other morphological or biomarker predictors of age in this population. The epigenetic clock generally explained more variance in true chronological age, resulted in lower median error, and exhibited less bias than predictions based on body mass index (BMI) or blood cell composition data from flow cytometry or blood smears (traits found to change with age in baboons [8, 9]). Its performance was comparable to molar dentine exposure, a classical marker of age [10] (Figure S2). For 16 males and 14 females, we had two samples collected at different points in time. The predicted ages from these longitudinally collected samples were older for the later-collected samples, as expected (Figure 1C-D; binomial test p = 5.95 x 10^{-5}). Furthermore, the change in epigenetic clock predictions between successive longitudinal samples positively predicted the actual change in age between sample dates (β = 0.312, p = 0.027).

Figure 1. Epigenetic clock age predictions in the Amboseli baboons. Predicted ages are shown relative to true chronological ages for **(A)** females (Pearson's r = 0.78, p < 10^{-29}, N = 142 samples) and **(B)** males (r = 0.86, p < 10^{-40}, N = 135 samples). Solid lines represent the best fit line; dashed lines show the line for y = x. **(C)**

and **(D)** show predictions for individuals with at least two samples in the data set (14 females and 16 males, respectively). In 26 of 30 cases (87%), samples collected later were correctly predicted to be from an older animal.

In addition to differences in overall accuracy, sex differences were also apparent in the slope of the relationship between predicted age and chronological age. Males show a 2.2-fold higher rate of change in predicted age, as a function of chronological age, compared to females (Figure 1A-B; chronological age by sex interaction in a linear model for predicted age: $\beta = 0.448$, $p < 10^{-18}$, N = 277). This result agrees with previous findings showing that male baboons senesce more rapidly than females—a pattern shared with most other primates investigated thus far, including humans [11]. Interestingly, sex differences are not apparent in animals < 8 years, which roughly corresponds to the age at which the majority of males have achieved adult dominance rank and dispersed from their natal group [12-14] (N = 158, chronological age by sex interaction $\beta = -0.038$, $p = 0.808$). Rather, sex difference becomes apparent after baboons have reached full physiological and social adulthood (N = 119, chronological age by sex interaction $\beta = 0.459$, $p < 10^{-6}$ in animals ≥ 8 years), when divergence between male and female life history strategies is most marked [12-14] and when aging rates between the sexes are predicted to diverge [15-17]. This pattern suggests that within each sex, deviations between predicted age and chronological age—commonly interpreted as a measure of "biological age" or accelerated aging—may also be affected by environmental or life history variation, as has been suggested in humans, lab mice, and captive rhesus macaques [3, 4, 18, 19].

To test this hypothesis, we focused on four factors of known importance to fertility and/or survival components of fitness in the Amboseli baboon population. First, we investigated the effects of cumulative early adversity, which is a strong predictor of shortened lifespan in female baboons: females who experience three or more major sources of early adversity have expected adult lifespans that are a decade shorter than those who experience none [20]. Additionally, those females are less capable of raising their own juvenile offspring later in life, suggesting that early adversity compromises their physical condition over the long-term [21]. Following [20, 21], we measured cumulative adversity as a count of major adverse experiences suffered in early life, including low maternal social status, early life drought, a competing younger sibling, maternal loss, and high experienced population density. To maximize our sample size, we omitted early life social connectedness (included in [20] but omitted for the same reason in [21]), because social connectedness data were missing for mothers born relatively early in the long-term study. We predicted that high cumulative early adversity, which is linked to reduced lifespan, would predict increased biological age.

Second, we considered social bond strength in adulthood, which is positively associated with longer adult lifespan in female Amboseli baboons, human populations, and several other wild social mammals [22-24]. We predicted that low social bond strength, due to its relationship with decreased lifespan, would be associated with increased biological age.

Third, we investigated the effects of dominance rank, which is a major determinant of resource access in baboons. High-ranking males sire the most offspring, and high-ranking females experience shorter interbirth intervals, retain higher fertility during droughts, and form stronger social bonds with males [22, 25-27].

Finally, we assessed the effect of BMI, which captures dimensions of both body condition and competitive advantage [8]. In this case, we calculated BMI relative to the expected value for each male's age, which eliminates the correlation between BMI and age and BMI and rank. The predictions for dominance rank and body mass index associations were less clear: improved resource access could conceivably slow biological aging, but increased investment in growth and reproduction (either through higher fertility in females or physical competition for rank in males) could also be energetically costly.

We tested these predictions by modeling the deviation between predicted age and known chronological age (Δ_{age}) as a function of cumulative early adversity, ordinal dominance rank, body mass index (controlling for age), and for females, social bond strength to other females. Social bond strength was not included in the model for males, as this measure was unavailable for a large proportion of males in this data set (53.8%). We also included chronological age as a predictor in the model, as epigenetic age tends to be systematically overpredicted for young individuals and underpredicted for old individuals (Figure 1A-B); including chronological age in the model controls for this compression effect. No predictor variables were strongly linearly correlated (all $R^2 < 0.10$; Table S3).

Surprisingly, despite being two of the strongest known predictors of lifespan in wild baboons, neither cumulative early life adversity nor social bond strength explain variation in Δ_{age} (Table 1). In contrast, high male dominance rank is strongly and significantly associated with larger values of Δ_{age} ($\beta = -0.785$, $p = 7.0 \times 10^{-4}$; Figure 2; Table 1). Alpha males are predicted to be an average of 10.95 months older than their true chronological age—a difference that translates to 11.5% of a male baboon's expected adult lifespan [7].

Dominance rank is strongly age structured in male baboons: males tend to attain their highest rank between 7 and 12 years of age and fall in rank thereafter. Thus, nearly all males in the top four rank positions in our data set were between 7 and 12 years of age at the time they were sampled (however, because not all 7 – 12 year-olds are high-ranking, low rank positions include males of all age groups; Table S1, Figure S3). Our finding that high rank predicts accelerated epigenetic aging therefore implies that males incur the costs of high rank primarily in early to mid-adulthood, and only if they succeed in attaining high rank. Accelerated epigenetic aging is thus a function of absolute rank values, regardless of age, not deviations from the mean rank expected given a male's age (i.e., "rank-for-age," which can be quantified as the residuals of male dominance rank modeled as a quadratic function of chronological age: Figure S3). In support of this interpretation, a model that includes rank-for-age as an additional covariate recapitulates the significant effect of ordinal male rank (p=0.045), but finds no effect of rank-for-age (p=0.819; Table S4). In contrast, we observed no evidence for rank effects on Δ_{age} in females, consistent with overall sex differences in patterns of aging in primates and other mammals [15] and marked sex differences in the effects of rank on other molecular phenotypes in the Amboseli baboons specifically [11, 28].

Covariate	β (Female)	P-value (Female)	β (Male)	P-value (Male)
Intercept	**5.400**	**1.33 x 10⁻¹⁵**	**3.294**	**1.19 x 10⁻⁸**
Cumulative early adversity	-0.050	0.807	-0.005	0.973
Social bond strength	0.382	0.164	—	—
Dominance rank	0.025	0.228	**-0.078**	**7.39 x 10⁻⁴**
Body mass index (age-adjusted)	0.026	0.682	**0.111**	**6.33 x 10⁻³**
Chronological age	**-0.699**	**1.62 x 10⁻²⁸**	**-0.277**	**8.36 x 10⁻⁸**

The header above the table reads: Table 1. Predictors of Δ_{age}[1]

[1]Separate linear models for Δ_{age} were fit for females (N = 66) and for males (N = 93) for whom no data values
were missing; social bond strength was not included in the model for males. Significant results are shown in
bold.

Figure 2. Dominance rank predicts epigenetic aging in male baboons. High rank is associated with
elevated values of Δ_{age} (β = -0.0785, p = 7.39 x 10⁻⁴, N = 105). The y-axis shows relative epigenetic age, a
measure of epigenetic aging similar to Δ_{age} that is based on the sample-specific residuals from the
relationship between predicted age and true chronological age. Positive (negative) values correspond to
predicted ages that are older (younger) than expected for that chronological age. Dominance rank is
measured using ordinal values, such that smaller values indicate higher rank. Dots and error bars represent
the means and standard errors, respectively. Gray values above the x-axis indicate sample sizes for each rank.
172
Previous work has shown that high-ranking male baboons, but not high-ranking
female baboons, up-regulate gene expression in inflammation-related and immune
response pathways [28]. Elevated or chronic inflammation is thought to be one of the
hallmarks of aging, and, in human populations, is one of the strongest predictors of
mortality risk [29-31]. Consistent with these observations, CpG sites in the epigenetic clock
that increase in DNA methylation with age (N = 459 sites) are enriched in or near genes
that are up-regulated in the Amboseli baboons in response to the bacterial endotoxin
lipopolysaccharide (LPS), which is a strong driver of inflammation (Figure S3; Fisher's
exact test: $\log_2$(odds ratio) = 1.88, p = 0.001; gene expression data from [28]). In contrast,
clock sites that decrease in DNA methylation with age (N = 134) are significantly enriched
in or near genes that are down-regulated after LPS exposure (Fisher's exact test: $\log_2$(OR) =

3.28, p = 0.002). Clock sites, especially those for which DNA methylation is positively
correlated with age, are also enriched in genes, CpG islands, promoter regions, and putative
enhancers, compared to the background set of sites we initially considered as candidates
for inclusion in the clock (Figure S4; Fisher's exact tests, all p < 0.005; similar functional
enrichment has been found in a human epigenetic clock [2]). Moreover, clock sites are
enriched in regions previously found to change in methylation levels with age [32] and in
regions showing regulatory activity in high-throughput reporter assays [33]. Together,
these results suggest that accelerated epigenetic aging in males reflects functionally
important changes in DNA methylation levels, concentrated in immune response and
inflammation-related pathways.
In baboon males, dominance rank reflects physical condition and fighting ability.
Rank is therefore dynamic across the life course, such that males in their prime (ages 7 –
12) are most likely to be high-ranking, and the same male is likely to occupy multiple
positions in the social hierarchy during his lifetime [14]. If accelerations in epigenetic age
are tightly coupled to rank, our results predict that across longitudinally collected samples,
a male who becomes higher ranking should look older for age in his second sample relative
to first, whereas a male who loses status should look younger for age in his second sample
relative to first. To assess this possibility, we calculated the residuals of the best fit line
relating chronological age and predicted age ("relative epigenetic age") for 14 males for
whom we had repeated samples collected when they occupied different dominance rank
positions (N = 28 samples, 2 per male). In this data set, the sample collected when males
were higher status typically predicted higher values of relative epigenetic age compared to
the sample collected when they were lower status, consistent with high rank-associated
accelerated biological aging (Figure 3; paired t-test, t = -2.31, p = 0.038). For example, in the
case of one young adult male who we sampled at rank 18 and then rank 1, as he was first
climbing the social hierarchy, the actual time that elapsed between samples was 0.79 years
but he exhibited an increase in *predicted* age of 2.6 years. Moreover, the two males that
showed a decrease in predicted age, despite increasing in chronological age (Figure 1D),
were among those that experienced the greatest drop in social status between samples.
Thus, change in rank between samples for the same male predicted change in Δ_{age},
controlling for chronological age (R^2 = 0.24, p=0.044). Consistent with our cross-sectional
results, we did not observe evidence for a relationship between change in Δ_{age} and rank-
for-age (R^2 = 0.14, p = 0.135).

Figure 3. Male baboons exhibit higher relative epigenetic age when they occupy higher ranks. Relative epigenetic age for males in which multiple samples were collected when they occupied different ordinal rank values. Arrow indicates the temporal direction of rank changes: left-facing arrows represent cases in which the later sample was collected when males were higher-ranking, and right-facing arrows represent cases in the later sample was collected when males were lower-ranking.

Together, our findings indicate that major environmental predictors of lifespan and mortality risk (e.g., social bond strength and early life adversity in this population) do not necessarily predict epigenetic measures of biological age. Although this assumption is widespread in the literature, including for epigenetic clock analyses [34, 35], our results are broadly consistent with empirical results in humans. Specifically, while studies of early life adversity, which also predicts lifespan in human populations, find relatively consistent support for a relationship between early adversity and accelerated epigenetic aging in children and adolescents [36-41], there is little evidence for the long-term effects of early adversity on epigenetic age in adulthood [42-47]. Thus, while DNA methylation may make an important contribution to the biological embedding of early adversity into adulthood [48, 49], it does not seem to do so through affecting the epigenetic clock itself. Social and environmental effects on the clock instead seem to be most influenced by concurrent conditions, lending support to "recency" models for environmental effects on aging that posit that health is more affected by the current environment than past experience [50-52]. Additional longitudinal sampling will be necessary to evaluate whether current conditions alone can explain accelerated epigenetic aging, or whether it also requires integrating the effects of exposures across the life course (the "accumulation" model: [50, 52]). Repeated samples could also help exclude an alternative explanation for our findings: that viability selection against individuals who experienced high early adversity attenuates the true effect of cumulative early adversity on relative epigenetic age.

Finally, our analyses reveal that males who achieve high rank appear epigenetically older than expected given their known chronological age. Although high ranking males also tend to be larger due to increased muscle mass (Pearson's r between rank and BMI = 0.56, p = 6.38 x 10^{-9}), we observed an additional, independent effect of age-adjusted BMI: males who had high BMI relative to their age also looked older for age, controlling for rank and age (β = 0.1107, p = 0.006) (Table 1). These two effects suggest that investment in body

condition, which is a crucial factor in male competitive ability, incurs other physiological
costs that compound to influence biological age. Indeed, previous research on the Amboseli
baboons also points to costs of high rank, including high levels of glucocorticoids in alpha
males [53], increased expression of genes involved in innate immunity and inflammation
[28], and a trend towards elevated mortality risk [54]. These associations may arise
because high-ranking males are both more likely to engage in physical conflict with other
males, and more likely to spend long periods of time in energetically costly mate-guarding
[55, 56]. Alternatively, males who are able to make significant investments in body
condition, while tolerating their accompanying costs, may be able to successfully maintain
high rank. Importantly, since doing so is likely to contribute to higher lifetime reproductive
success for male baboons, the fitness-associated benefits of high rank can exceed the costs,
even if accelerated biological aging increases the risk of mortality.
261

Acknowledgements
We gratefully acknowledge the support provided by the National Science Foundation and
the National Institutes of Health for the majority of the data represented here, currently
through NSF IOS 1456832, NIH R01AG053308, R01AG053330, R01HD088558, and
P01AG031719. R.A.J. is supported by NIH F32HD095616 and J.A.A. by NSF #2018264636.
We thank the members of the Amboseli Baboon Research Project for collecting the data
presented here, especially J. Altmann for her foundational role in establishing the study
population and these data sets; J. Gordon, N. Learn, and K. Pinc for managing the database;
R.S. Mututua, S. Sayialel, and J.K. Warutere for data collection in the field; and T. Wango and
V. Oudu for their assistance in Nairobi. We also thank the Kenya Wildlife Service, University
of Nairobi, the Institute of Primate Research, the National Museums of Kenya, the National
Council for Science, Technology, and Innovation, members of the Amboseli-Longido
pastoralist communities, the Enduimet Wildlife Management Area, Ker & Downey Safaris,
Air Kenya, and Safarilink for their assistance in Kenya. Finally, we thank current and past
members of the Tung, Alberts, Archie, and Altmann labs for their helpful feedback. This
research was approved by IACUCs at Duke University, University of Notre Dame, and
Princeton University and adhered to all the laws and guidelines of Kenya. For a complete
set of acknowledgments of funding sources, logistical assistance, and data collection and
management, please visit http://amboselibaboons.nd.edu/acknowledgements/.

Author Contributions
Conceptualization, R.A.J., J.A.A., J.T., E.A.A., A.J.L.; Investigation, J.A.A., R.A.J., A.J.L., F.A.C.,
M.Y.A., T.N.V., and J.T.; Formal Analysis, J.A.A. and R.A.J.; Writing—Original Draft, R.A.J.,
J.A.A., and J.T.; Writing—Reviewing & Editing, R.A.J., J.A.A., A.J.L., T.N.V., M.Y.A., F.A.C., S.C.A.,
E.A.A., and J.T. Funding Acquisition, J.T., S.C.A., and E.A.A. Supervision, J.T.
287

288 Declaration of Interests
289 The authors declare no competing interests.
290
291

STAR Methods
Study population and biological sample collection
This study focused on a longitudinally monitored population of wild baboons (*Papio*
cynocephalus, the yellow baboon, with some admixture from the closely related anubis
baboon *P. anubis* [57, 58]) in the Amboseli ecosystem of Kenya. This population has been
continuously monitored by the Amboseli Baboon Research Project (ABRP) since 1971 [6].
For the majority of study subjects (N = 242 of 245 individuals), birth dates were therefore
known to within a few days' error; for the remainder, birth dates were known within 3
months' error (Table S1).
All DNA methylation data were generated from blood-derived DNA obtained during
periodic darting efforts, as detailed in [28, 59, 60]. Samples were obtained under approval
from the Institutional Animal Care and Use Committee (IACUC) of Duke University and
adhered to all the laws and guidelines of Kenya. In brief, individually recognized study
subjects were temporarily anesthetized using a Telazol-loaded dart delivered through a
blow gun. Baboons were then safely moved to a new location where blood samples and
morphometric data, including body mass and crown-rump length, were collected. Baboons
were then allowed to recover from anesthesia in a covered holding cage and released to
their group within 2 – 4 hours. Blood samples were stored at -20 C in Kenya until export to
the United States.
311
DNA methylation data
DNA methylation data were generated from blood-extracted DNA collected from
known individuals in the Amboseli study population (N = 277 samples from 245 animals;
14 females and 14 males were each sampled twice, and 1 female and 1 male were each
sampled three times). Here, we analyzed a combined data set that included previously
published reduced representation bisulfite sequencing [61] (RRBS) data from the same
population (N = 36) [5] and new RRBS data from 241 additional individuals.
RRBS libraries were constructed following [62], using ~200 ng baboon DNA plus 0.2
ng unmethylated lambda phage DNA per sample as input. Samples were sequenced to a
mean depth of 17.8 (± 10.5 s.d.) million reads on either the Illumina HiSeq 2000 or HiSeq
4000 platform (Table S1), with an estimated mean bisulfite conversion efficiency (based on
the conversion rate of lambda phage DNA) of 99.8% (minimum = 98.1%). Sequence reads
were trimmed with Trim Galore! [63] to remove adapters and low quality sequence (Phred
score < 20). Trimmed reads were mapped with BSMAP [64] to the baboon genome
(*Panu2.0*) allowing a 10% mismatch rate to account for the degenerate composition of
bisulfite-converted DNA. We used the mapped reads to count the number of methylated
and total reads per CpG site, per sample [64, 65]. Following [5, 32], CpG sites were filtered
to retain sites with a mean methylation level between 0.1 and 0.9 (i.e., to exclude
constitutively hyper- or hypo-methylated sites) and mean coverage $\geq$5x. We also excluded
any CpG sites with missing data for $\geq$15% of individuals in the sample. After filtering, we
retained N = 458,504 CpG sites for downstream analysis. For the remaining missing data
(mean number of missing sites per sample = 1.4% ± 3.5% s.d., equivalent to 6,409 ± 16,024
sites), we imputed methylation levels using a k-nearest neighbors approach in the R
package *impute*, using default parameters [66].
336

Building the epigenetic clock

337
We used the R package *glmnet* [67] to build a DNA methylation clock for baboons.
Specifically, we fit a linear model in which the predictor variables were normalized levels of
DNA methylation at 458,504 candidate clock CpG sites across the genome and the response
variable was chronological age. To account for the excess of CpG sites relative to samples,
glmnet uses an elastic net penalty to shrink predictor coefficients toward 0 [68]. Optimal
alpha parameters were identified by grid searching across a range of alphas from 0
(equivalent to ridge regression) to 1 (equivalent to LASSO) by increments of 0.1. We
defined the optimal alpha as the value that maximized R^2 between predicted and true
chronological age across all samples. We set the regularization parameter lambda to the
value that minimized mean-squared error during n-fold internal cross-validation.
To generate predicted age estimates for a given sample, we used a leave-one-out
cross-validation approach in which all samples but the "test" sample were included for
model training, and the resulting model was used to predict age for the left-out test sample.
Importantly, training samples were scaled independently of the test sample in each leave-
one-out model to avoid bleed-through of information from the test data into the training
data. To do so, we first quantile normalized methylation ratios (the proportion of
methylated counts to total counts for each CpG site) within each sample to a standard
normal distribution. Training samples were then separated from the test sample and the
methylation levels for each CpG site in the training set were quantile normalized across
samples to a standard normal distribution. For predicting age in the test sample, we
compared the methylation value for each site in the test sample to the empirical cumulative
distribution function for the training samples (at the same site) to estimate the quantile in
which the training sample methylation ratio fell. The training sample was then assigned the
same quantile value from the standard normal distribution using the function *qnorm* in R.
362
Comparisons to alternative predictors of aging
To assess the utility of the DNA methylation clock relative to other data types, we
compared its predictive accuracy to clocks based on three other age-related phenotypes:
tooth wear (percent molar dentine exposure [10]), body condition (body mass index: BMI
[8]), and blood cell type composition (blood smear counts and lymphocyte/monocyte
proportions from flow cytometry performed on peripheral blood mononuclear cells, as in
[28, 69]). Leave-one-out model training and prediction were performed for each data type
using linear modeling (i.e., not glmnet, since the number of features was much less than the
number of samples in this case). To compare the relative predictive accuracy of each data
type, we calculated the R^2 between predicted and chronological age, the median absolute
difference between predicted and chronological age, and the bias in age predictions (the
absolute value of 1- slope of the best fit line between predicted and chronological age)
(Figure S2).
Tooth wear. Molar enamel in baboons wears away with age to expose the underlying
dentine layer. Percent dentine exposure (PDE) on the molar occlusal surface has been
shown to be strongly age-correlated in previous work [10]. To assess its predictive power,
we obtained PDE data from tooth casts reported by Galbany and colleagues [10] for the left
upper molars (tooth positions M1, M2, M3) and left lower molars (tooth positions M1, M2,
M3) for 39 males and 34 females in our data set. For each molar position (M1, M2, M3)
within each individual, we calculated PDE as the mean for the upper and lower molars.

Because dentine exposure scales quadratically with respect to age [10], we fit age as a
function of PDE using the following model: $age \sim \sqrt{PDE_{M1}} + \sqrt{PDE_{M2}} + \sqrt{PDE_{M3}}$.
Body mass index. For both male and female baboons in Amboseli, body mass
increases with age until individuals reach peak size, and then tends to decrease with age as
animals lose fat and/or muscle mass [8]. To quantify body condition using body mass, we
calculated body mass index (BMI) values for 139 males and 154 females for whom body
mass and crown-rump length data were available from periodic darting efforts. We
retained only measures taken from animals born into and sampled in wild-feeding study
groups, when sex-skin swellings (in females only) that could affect crown-rump length
measures were absent. BMI was calculated as mass (kilograms) divided by crown-rump
length (meters squared), following [70]. To assess the predictive power of BMI for age, we
built sex-specific piecewise-regression models. Breakpoints for the piecewise-regression
models (to separate "youthful" versus "aged" animals) were set at 8 years old for males and
10 years old for females, following findings from previous work on body mass in the
Amboseli population [8].
Blood cell type composition. The proportions of different cell types in blood change
across the life course, including in baboons [9]. We assessed the predictive power of blood
cell composition for age using two data sets. First, we used data collected from blood smear
counts (N = 134) for five major white blood cell types: basophils, eosinophils, monocytes,
lymphocytes, and neutrophils. Second, we used data on the proportional representation of
five peripheral blood mononuclear cell (PBMC) subsets: cytotoxic T cells, helper T cells, B
cells, monocytes, and natural killer cells, measured using flow cytometry as reported by Lea
and colleagues [28] (N = 53). Cell types were included as individual covariates for leave-
one-out model training.
407
Sources of variance in predicted age
We asked whether factors known to be associated with inter-individual variation in
fertility or survival also predict inter-individual variation in Δ_{age} (predicted age from the
epigenetic clock minus known chronological age). To do so, we fit linear models separately
for males and females, with Δ_{age} as the dependent variable and dominance rank at the time
of sampling, cumulative early adversity, relative BMI (corrected for age), and chronological
age as predictor variables [20]). For females, we also included a measure of social bond
strength to other females as a predictor variable, based on findings that show that socially
isolated females experience higher mortality rates in adulthood [22, 71]. Samples with
missing values for any of the predictor variables were excluded in the model, resulting in a
final analysis set of 66 female samples (from 59 females) and 93 male samples (from 84
males). The chronological ages of samples with complete data relative to samples with
missing data were equivalent for females (t-test, t = 1.95, p = 0.053) but were slighly lower
for males (t-test, t = -3.04, p = 0.003; mean chronological ages are 7.98 and 9.65 years for
complete and missing samples, respectively). Predictor variables were measured as
follows.
Dominance rank. Sex-specific dominance hierarchies were constructed monthly for
every social group in the study population based on the outcomes of dyadic aggressive
encounters. Ordinal dominance ranks were assigned to every adult based on these
hierarchies, such that low numbers represent high rank/social status and high numbers

represent low rank/social status [72]. Although most analyses of data from the Amboseli
baboons have used ordinal ranks as the primary measure of social status, in some analyses
proportional rank (i.e., the proportion of same-sex members of an individual's social group
that he or she dominates) has proven to be a stronger predictor of behavioral, life history,
or physiological outcomes [73]. In this study, we chose to use ordinal ranks, but
proportional and ordinal dominance rank were highly correlated in our dataset (r^2=.70,
p=1.13 x 10^{-58}). Using proportional rank rather than ordinal rank did not qualitatively
affect our analyses. Additionally, to investigate whether the patterns we observed are due
to a consistent effect of rank across all ages, or instead an effect of being high or low rank
relative to the expected (mean) value for a male's age, we also calculated a "rank-for-age"
value. Rank-for-age is defined as the residuals of a model with dominance rank as the
response variable and age and age² as the predictor variables (Fig S4).
Cumulative early adversity. Previous work in Amboseli defined a cumulative early
adversity score as the sum of 6 different adverse conditions that a baboon could experience
during early life [20]. This index strongly predicts adult lifespan in female baboons, and a
modified version of this index also predicts offspring survival [21]. To maximize the sample
size available for analysis, we excluded maternal social connectedness, the source of
adversity with the highest frequency of missing data, leaving us with a cumulative early
adversity score generated from 5 different binary-coded adverse experiences. These
experiences were: (i) early life drought (defined as ≤ 200 mm of rainfall in the first year of
life), which is linked to reduced fertility in females [27, 74]; (ii) having a low ranking
mother (defined as falling within the lowest quartile of ranks for individuals in the dataset),
which predicts rates of maturation [75-77]; (iii) having a close-in-age younger sibling (<1.5
years), which may redirect maternal investment to the sibling [78], (iv) being born into a
large social group, which may increase within-group competition for shared resources [27,
77, 79], and (v) maternal death before the age of 4, which results in a loss of both social and
nutritional resources [77, 80].
Body mass index. BMI was modeled as the residuals from sex-specific piecewise
regression models relating BMI to age. By taking this approach, we asked whether having
relatively high BMI for one's age and sex predicted higher (or lower) Δ_{age}.
Social bond strength. For this analysis, we measured female social bond strength to
other females using the dyadic sociality index (F-DSI) [54]. We did not include this
parameter (male's social bond strength to females) for the male model, because this
measure is unavailable for young males in this dataset. F-DSI was calculated as an
individual's average bond strength with her top three female social partners, in the 365
days prior to the day of sampling, controlling for observer effort. This approach is based on
representative interaction sampling of grooming interactions between females, in which
observers record all grooming interactions in their line of sight while moving through the
group conducting random-ordered, 10-minute long focal animal samples of pre-selected
individuals. Because smaller groups receive more observer effort per individual and per
dyad (and thus record more grooming interactions per individual or dyad), we estimated
observer effort for dyad *d* in year *y* as:

470
$$E_{d,y} = \frac{c_d(s_d)}{f_d}$$

471 where c_d is the number of days the two females in a dyad were coresident in the same

472 social group, s_d is the number of focal samples taken during the dyad's coresidence, and f_d
473 is the average number of females in the group during the dyad's coresidence.
474   F-DSI for each adult female dyad in each year is the z-scored residual, ε, from the
475 model:

476
$$\log(R_{d,y}) = \beta(\log(E_{d,y})) + \varepsilon$$

477 where $R_{d,y}$ is the number of grooming interactions for dyad d in year y divided by the
478 number of days that the two individuals were coresident, and $E_{d,y}$ is observer effort.
479
Analysis of longitudinal samples
To test whether changes in rank predict changes in relative epigenetic age within
individuals, we used data from 5 males from the original dataset and generated additional
RRBS data for 9 samples, resulting in a final set of 14 males who each were sampled at least
twice in the data set, when they occupied different ordinal ranks (mean years elapsed
between samples = 3.92 ± 1.94 s.d.; mean absolute difference in dominance ranks = 6.86 ±
5.07 s.d.). This effort increased our total sample size to N = 286 samples from 248 unique
individuals. To incorporate the new samples into our analysis, we reperformed leave-one-
out age prediction with N-fold internal cross validation at the optimal alpha selected for the
original N = 277 samples (alpha = 0.1). For the 277 samples carried over from the original
analysis, we verified that age predictions were nearly identical between the previous
analysis and the expanded data set (R^2 = 0.98, p = 2.21 x 10^{-239}).
Based on the new age predictions for males in the data set (N = 144), we again
calculated relative epigenetic age as the residual of the best fit line relating predicted age to
chronological age. We then used the 14 males with repeated DNA methylation profiles and
rank measures in this dataset to test whether, within individuals, changes in dominance
rank or rank-for-age explained changes in relative epigenetic age between samples.
497
Epigenetic clock enrichment analyses
To evaluate whether CpG sites included in the epigenetic clock were enriched in
functionally important regions of the baboon genome [32, 81], we used two-sided Fisher's
exact tests to investigate enrichment/depletion of the 593 epigenetic clock sites in (i) gene
bodies and exons, based on the Ensembl annotation *Panu2.0.90*; (ii) CpG islands annotated
in the UCSC Genome Browser; (iii) CpG shores, defined as the 2,000 basepairs flanking CpG
islands (following [32, 81, 82]); and (iv) promoter regions, defined as the 2,000 basepairs
upstream of the 5'-most annotated transcription start site for each protein-coding gene
(following [32, 81]). We also considered (v) putative enhancer regions, which have not
been annotated for the *Panu2.0* assembly. We therefore used ENCODE H3K4me1 ChIP-seq
data from humans [83] and the *liftOver* tool to define likely enhancer coordinates in
Panu2.0.
We also tested for enrichment of clock sites in regions of the genome that have been
identified by previous empirical studies to be of special interest. First, we considered
regions that likely have regulatory activity in blood cells, defined as all 200 base-pair
windows that showed evidence of enhancer activity in a recently performed massively
parallel reporter assay [33]. We used *liftOver* to identify the inferred homologous *Panu2.0*
coordinates for these windows, which were originally defined in the human genome.
Second, we defined age-related differentially methylated regions (age DMRs) in the

Amboseli baboons based on genomic intervals found, in previous analyses, to contain at
least three closely spaced age-associated CpG sites (inter-CpG distance ≤ 1kb), as described
in [32]. Third, we defined lipopolysaccharide (LPS) up-regulated and LPS down-regulated
genes as those genes that were significantly differentially expressed (1% false discovery
rate) between unstimulated Amboseli baboon white blood cells and LPS-stimulated cells
from the same individual, following 10 hours of culture in parallel [28].
523
524   Data and code availabity
525   All sequencing data generated during this study are available in the NCBI Sequence Read
526   Archive (project accession PRJNA607996).

References

1. Hannum, G., Guinney, J., Zhao, L., Zhang, L., Hughes, G., Sadda, S., Klotzle, B., Bibikova, M., Fan, J.-B., and Gao, Y. (2013). Genome-wide methylation profiles reveal quantitative views of human aging rates. Molecular cell *49*, 359-367.

2. Horvath, S. (2013). DNA methylation age of human tissues and cell types. Genome biology *14*, 3156.

3. Petkovich, D.A., Podolskiy, D.I., Lobanov, A.V., Lee, S.-G., Miller, R.A., and Gladyshev, V.N. (2017). Using DNA methylation profiling to evaluate biological age and longevity interventions. Cell metabolism *25*, 954-960. e956.

4. Stubbs, T.M., Bonder, M.J., Stark, A.-K., Krueger, F., von Meyenn, F., Stegle, O., and Reik, W. (2017). Multi-tissue DNA methylation age predictor in mouse. Genome biology *18*, 68.

5. Lea, A.J., Altmann, J., Alberts, S.C., and Tung, J. (2016). Resource base influences genome‐wide DNA methylation levels in wild baboons (Papio cynocephalus). Molecular ecology *25*, 1681-1696.

6. Alberts, S.C., and Altmann, J. (2012). The Amboseli Baboon Research Project: 40 years of continuity and change. In Long-term field studies of primates. (Springer), pp. 261-287.

7. Colchero, F., Rau, R., Jones, O.R., Barthold, J.A., Conde, D.A., Lenart, A., Nemeth, L., Scheuerlein, A., Schoeley, J., and Torres, C. (2016). The emergence of longevous populations. Proceedings of the National Academy of Sciences *113*, E7681-E7690.

8. Altmann, J., Gesquiere, L., Galbany, J., Onyango, P.O., and Alberts, S.C. (2010). Life history context of reproductive aging in a wild primate model. Annals of the New York Academy of Sciences *1204*, 127-138.

9. Jayashankar, L., Brasky, K.M., Ward, J.A., and Attanasio, R. (2003). Lymphocyte modulation in a baboon model of immunosenescence. Clin. Diagn. Lab. Immunol. *10*, 870-875.

10. Galbany, J., Altmann, J., Pérez‐Pérez, A., and Alberts, S.C. (2011). Age and individual foraging behavior predict tooth wear in Amboseli baboons. American Journal of Physical Anthropology *144*, 51-59.

11. Bronikowski, A.M., Altmann, J., Brockman, D.K., Cords, M., Fedigan, L.M., Pusey, A., Stoinski, T., Morris, W.F., Strier, K.B., and Alberts, S.C. (2011). Aging in the natural world: comparative data reveal similar mortality patterns across primates. Science *331*, 1325-1328.

12. Alberts, S.C., and Altmann, J. (1995). Balancing costs and opportunities: dispersal in male baboons. The American Naturalist *145*, 279-306.

13. Alberts, S.C., and Altmann, J. (1995). Preparation and activation: determinants of age at reproductive maturity in male baboons. Behavioral Ecology and Sociobiology *36*, 397-406.

14. Alberts, S.C., Watts, H.E., and Altmann, J. (2003). Queuing and queue-jumping: long-term patterns of reproductive skew in male savannah baboons, Papio cynocephalus. Animal Behaviour *65*, 821-840.

15. Clutton-Brock, T.H., and Isvaran, K. (2007). Sex differences in ageing in natural populations of vertebrates. Proceedings of the Royal Society B: Biological Sciences *274*, 3097-3104.

16. Kirkwood, T.B., and Rose, M.R. (1991). Evolution of senescence: late survival
sacrificed for reproduction. Philosophical Transactions of the Royal Society of
London. Series B: Biological Sciences *332*, 15-24.
17. Williams, G.C. (1957). Pleiotropy, natural selection, and the evolution of senescence.
evolution, 398-411.
18. Maegawa, S., Lu, Y., Tahara, T., Lee, J.T., Madzo, J., Liang, S., Jelinek, J., Colman, R.J.,
and Issa, J.-P.J. (2017). Caloric restriction delays age-related methylation drift.
Nature communications *8*, 539.
19. Ryan, J., Wrigglesworth, J., Loong, J., Fransquet, P.D., and Woods, R.L. (2019). A
systematic review and meta-analysis of environmental, lifestyle and health factors
associated with DNA methylation age. The journals of gerontology. Series A,
Biological sciences and medical sciences.
20. Tung, J., Archie, E.A., Altmann, J., and Alberts, S.C. (2016). Cumulative early life
adversity predicts longevity in wild baboons. Nature communications *7*, 11181.
21. Zipple, M.N., Archie, E.A., Tung, J., Altmann, J., and Alberts, S.C. (2019).
Intergenerational effects of early adversity on survival in wild baboons. eLife *8*,
e47433.
22. Archie, E.A., Tung, J., Clark, M., Altmann, J., and Alberts, S.C. (2014). Social affiliation
matters: both same-sex and opposite-sex relationships predict survival in wild
female baboons. Proceedings of the Royal Society B: Biological Sciences *281*,
20141261.
23. Holt-Lunstad, J., Smith, T.B., and Layton, J.B. (2010). Social relationships and
mortality risk: a meta-analytic review. PLoS Med *7*, e1000316.
24. Snyder-Mackler, N., Burger, J.R., Gaydosh, L., Belsky, D., Noppert, G.A., Campos, F.A.,
Bartolomucci, A., Yang, Y.C., Aiello, A.E., O'Rand, A., et al. Social determinants of
health and survival in humans and other animals. In review.
25. Alberts, S.C., Buchan, J.C., and Altmann, J. (2006). Sexual selection in wild baboons:
from mating opportunities to paternity success. Animal Behaviour *72*, 1177-1196.
26. Gesquiere, L.R., Altmann, J., Archie, E.A., and Alberts, S.C. (2018). Interbirth intervals
in wild baboons: Environmental predictors and hormonal correlates. American
journal of physical anthropology *166*, 107-126.
27. Lea, A.J., Altmann, J., Alberts, S.C., and Tung, J. (2015). Developmental constraints in
a wild primate. The American Naturalist *185*, 809-821.
28. Lea, A.J., Akinyi, M.Y., Nyakundi, R., Mareri, P., Nyundo, F., Kariuki, T., Alberts, S.C.,
Archie, E.A., and Tung, J. (2018). Dominance rank-associated gene expression is
widespread, sex-specific, and a precursor to high social status in wild male baboons.
Proceedings of the National Academy of Sciences *115*, E12163-E12171.
29. Castagné, R., Garès, V., Karimi, M., Chadeau-Hyam, M., Vineis, P., Delpierre, C., Kelly-
Irving, M., and Consortium, L. (2018). Allostatic load and subsequent all-cause
mortality: which biological markers drive the relationship? Findings from a UK birth
cohort. European journal of epidemiology *33*, 441-458.
30. Levine, M.E. (2012). Modeling the rate of senescence: can estimated biological age
predict mortality more accurately than chronological age? Journals of Gerontology
Series A: Biomedical Sciences and Medical Sciences *68*, 667-674.
31. López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The
hallmarks of aging. Cell *153*, 1194-1217.

32. Lea, A.J., Tung, J., and Zhou, X. (2015). A flexible, efficient binomial mixed model for
identifying differential DNA methylation in bisulfite sequencing data. PLoS genetics
11, e1005650.
33. Lea, A.J., Vockley, C.M., Johnston, R.A., Del Carpio, C.A., Barreiro, L.B., Reddy, T.E., and
Tung, J. (2018). Genome-wide quantification of the effects of DNA methylation on
human gene regulation. eLife *7*, e37513.
34. Liu, Z., Chen, B.H., Assimes, T.L., Ferrucci, L., Horvath, S., and Levine, M.E. (2019). The
role of epigenetic aging in education and racial/ethnic mortality disparities among
older US Women. Psychoneuroendocrinology *104*, 18-24.
35. Shalev, I., and Belsky, J. (2016). Early-life stress and reproductive cost: A two-hit
developmental model of accelerated aging? Medical Hypotheses *90*, 41-47.
36. Brody, G.H., Miller, G.E., Yu, T., Beach, S.R., and Chen, E. (2016). Supportive family
environments ameliorate the link between racial discrimination and epigenetic
aging: A replication across two longitudinal cohorts. Psychological Science *27*, 530-
541.
37. Brody, G.H., Yu, T., Chen, E., Beach, S.R., and Miller, G.E. (2016). Family‐centered
prevention ameliorates the longitudinal association between risky family processes
and epigenetic aging. Journal of child psychology and psychiatry *57*, 566-574.
38. Davis, E., Humphreys, K., McEwen, L., Sacchet, M., Camacho, M., MacIsaac, J., Lin, D.,
Kobor, M., and Gotlib, I. (2017). Accelerated DNA methylation age in adolescent
girls: associations with elevated diurnal cortisol and reduced hippocampal volume.
Translational psychiatry *7*, e1223.
39. Jovanovic, T., Vance, L.A., Cross, D., Knight, A.K., Kilaru, V., Michopoulos, V., Klengel,
T., and Smith, A.K. (2017). Exposure to violence accelerates epigenetic aging in
children. Scientific reports *7*, 8962.
40. Marini, S., Davis, K.A., Soare, T.W., Suderman, M.J., Simpkin, A.J., Smith, A.D., Wolf, E.J.,
Relton, C.L., and Dunn, E.C. (2018). Predicting cellular aging following exposure to
adversity: Does accumulation, recency, or developmental timing of exposure
matter? BioRxiv, 355743.
41. Sumner, J.A., Colich, N.L., Uddin, M., Armstrong, D., and McLaughlin, K.A. (2019).
Early experiences of threat, but not deprivation, are associated with accelerated
biological aging in children and adolescents. Biological psychiatry *85*, 268-278.
42. Austin, M.K., Chen, E., Ross, K.M., McEwen, L.M., MacIsaac, J.L., Kobor, M.S., and
Miller, G.E. (2018). Early-life socioeconomic disadvantage, not current, predicts
accelerated epigenetic aging of monocytes. Psychoneuroendocrinology *97*, 131-134.
43. Boks, M.P., van Mierlo, H.C., Rutten, B.P., Radstake, T.R., De Witte, L., Geuze, E.,
Horvath, S., Schalkwyk, L.C., Vinkers, C.H., and Broen, J.C. (2015). Longitudinal
changes of telomere length and epigenetic age related to traumatic stress and post-
traumatic stress disorder. Psychoneuroendocrinology *51*, 506-512.
44. Lawn, R.B., Anderson, E.L., Suderman, M., Simpkin, A.J., Gaunt, T.R., Teschendorff,
A.E., Widschwendter, M., Hardy, R., Kuh, D., and Relton, C.L. (2018). Psychosocial
adversity and socioeconomic position during childhood and epigenetic age: analysis
of two prospective cohort studies. Human molecular genetics *27*, 1301-1308.
45. Simons, R.L., Lei, M.K., Beach, S.R., Philibert, R.A., Cutrona, C.E., Gibbons, F.X., and
Barr, A. (2016). Economic hardship and biological weathering: the epigenetics of

aging in a US sample of black women. Social Science & Medicine *150*, 192-200.

46.   Wolf, E.J., Maniates, H., Nugent, N., Maihofer, A.X., Armstrong, D., Ratanatharathorn,
A., Ashley-Koch, A.E., Garrett, M., Kimbrel, N.A., and Lori, A. (2018). Traumatic stress
and accelerated DNA methylation age: a meta-analysis. Psychoneuroendocrinology
92, 123-134.

47.   Zannas, A.S., Arloth, J., Carrillo-Roa, T., Iurato, S., Röh, S., Ressler, K.J., Nemeroff, C.B.,
Smith, A.K., Bradley, B., and Heim, C. (2015). Lifetime stress accelerates epigenetic
aging in an urban, African American cohort: relevance of glucocorticoid signaling.
Genome biology *16*, 266.

48.   Aristizabal, M.J., Anreiter, I., Halldorsdottir, T., Odgers, C.L., McDade, T.W.,
Goldenberg, A., Mostafavi, S., Kobor, M.S., Binder, E.B., and Sokolowski, M.B. (2019).
Biological embedding of experience: A primer on epigenetics. Proceedings of the
National Academy of Sciences, 201820838.

49.   Hertzman, C. (2012). Putting the concept of biological embedding in historical
perspective. Proceedings of the National Academy of Sciences *109*, 17160-17167.

50.   Ben-Shlomo, Y., and Kuh, D. (2002). A life course approach to chronic disease
epidemiology: conceptual models, empirical challenges and interdisciplinary
perspectives. (Oxford University Press).

51.   Shanahan, L., Copeland, W.E., Costello, E.J., and Angold, A. (2011). Child-, adolescent-
and young adult-onset depressions: differential risk factors in development?
Psychological medicine *41*, 2265-2274.

52.   Shanahan, M.J., and Hofer, S.M. (2011). Molecular genetics, aging, and well-being:
Sensitive period, accumulation, and pathway models. In Handbook of aging and the
social sciences. (Elsevier), pp. 135-147.

53.   Gesquiere, L.R., Learn, N.H., Simao, M.C.M., Onyango, P.O., Alberts, S.C., and Altmann,
J. (2011). Life at the top: rank and stress in wild male baboons. Science *333*, 357-
360.

54.   Campos, F.A., Villavicencio, F., Archie, E.A., Colchero, F., and Alberts, S.C. (2020).
Social relationships, social status, and survival in wild baboons: A tale of two sexes.
Philosophical Transactions of the Royal Society. Series B: Biological Sciences.

55.   Archie, E.A., Altmann, J., and Alberts, S.C. (2012). Social status predicts wound
healing in wild baboons. Proceedings of the National Academy of Sciences *109*,
9017-9022.

56.   Rasmussen, K. (1985). Changes in the activity budgets of yellow baboons (Papio
cynocephalus) during sexual consortships. Behavioral Ecology and Sociobiology *17*,
161-170.

57.   Alberts, S.C., and Altmann, J. (2001). Immigration and hybridization patterns of
yellow and anubis baboons in and around Amboseli, Kenya. American Journal of
Primatology: Official Journal of the American Society of Primatologists *53*, 139-154.

58.   Tung, J., Charpentier, M.J., Garfield, D.A., Altmann, J., and Alberts, S.C. (2008). Genetic
evidence reveals temporal change in hybridization patterns in a wild baboon
population. Molecular Ecology *17*, 1998-2011.

59.   Altmann, J., Alberts, S.C., Haines, S.A., Dubach, J., Muruthi, P., Coote, T., Geffen, E.,
Cheesman, D.J., Mututua, R.S., and Saiyalel, S.N. (1996). Behavior predicts genes
structure in a wild primate group. Proceedings of the National Academy of Sciences
93, 5797-5801.

60. Tung, J., Zhou, X., Alberts, S.C., Stephens, M., and Gilad, Y. (2015). The genetic
architecture of gene expression levels in wild baboons. Elife *4*, e04729.
61. Meissner, A., Gnirke, A., Bell, G.W., Ramsahoye, B., Lander, E.S., and Jaenisch, R.
(2005). Reduced representation bisulfite sequencing for comparative high-
resolution DNA methylation analysis. Nucleic acids research *33*, 5868-5877.
62. Boyle, P., Clement, K., Gu, H., Smith, Z.D., Ziller, M., Fostel, J.L., Holmes, L., Meldrim, J.,
Kelley, F., and Gnirke, A. (2012). Gel-free multiplexed reduced representation
bisulfite sequencing for large-scale DNA methylation profiling. Genome biology *13*,
R92.
63. Krueger, F. (2012). Trim Galore: a wrapper tool around Cutadapt and FastQC to
consistently apply quality and adapter trimming to FastQ files, with some extra
functionality for MspI-digested RRBS-type (Reduced Representation Bisufite-Seq)
libraries. URL http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.
64. Xi, Y., and Li, W. (2009). BSMAP: whole genome bisulfite sequence MAPping
program. BMC bioinformatics *10*, 232.
65. Krueger, F. (2015). Trim Galore!: A wrapper tool around Cutadapt and FastQC to
consistently apply quality and adapter trimming to FastQ files. (0.4).
66. Hastie, T., Tibshirani, R., Narasimhan, B., and Chu, G. (2001). impute: Imputation for
microarray data. Bioinformatics *17*, 520-525.
67. Friedman, J., Hastie, T., and Tibshirani, R. (2009). glmnet: Lasso and elastic-net
regularized generalized linear models. R package version *1*.
68. Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization paths for
generalized linear models via coordinate descent. Journal of statistical software *33*,
1.
69. Snyder-Mackler, N., Sanz, J., Kohn, J.N., Brinkworth, J.F., Morrow, S., Shaver, A.O.,
Grenier, J.-C., Pique-Regi, R., Johnson, Z.P., and Wilson, M.E. (2016). Social status
alters immune regulation and response to infection in macaques. Science *354*, 1041-
1045.
70. Altmann, J., Schoeller, D., Altmann, S.A., Muruthi, P., and Sapolsky, R.M. (1993). Body
size and fatness of free‐living baboons reflect food availability and activity levels.
American Journal of Primatology *30*, 149-161.
71. Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig,
R.M., Seyfarth, R.M., and Cheney, D.L. (2010). Strong and consistent social bonds
enhance the longevity of female baboons. Current Biology *20*, 1359-1361.
72. Hausfater, G., Altmann, J., and Altmann, S. (1982). Long-term consistency of
dominance relations among female baboons (Papio cynocephalus). Science *217*,
752-755.
73. Archie, E.A., Altmann, J., and Alberts, S.C. (2014). Costs of reproduction in a long-
lived female primate: injury risk and wound healing. Behavioral ecology and
sociobiology *68*, 1183-1193.
74. Beehner, J.C., Onderdonk, D.A., Alberts, S.C., and Altmann, J. (2006). The ecology of
conception and pregnancy failure in wild baboons. Behavioral Ecology *17*, 741-750.
75. Altmann, J., and Alberts, S.C. (2003). Intraspecific variability in fertility and offspring
survival in a nonhuman primate: behavioral control of ecological and social sources.
In Offspring: The Biodemography of Fertility and Family Behavior, K.W. Wachter

754   and R.A. Bulatao, eds. (Washington, DC: The National Academies Press).

76. Altmann, J., Hausfater, G., and Altmann, S.A. (1988). Determinants of Reproductive Success in Savannah Baboons, Papio cynocephalus. In Reproductive Success: Studies of Individual Variation in Contrasting Breeding Systems, T.H. Clutton-Brock, ed. (Chicago: The University of Chicago Press).

77. Charpentier, M., Tung, J., Altmann, J., and Alberts, S. (2008). Age at maturity in wild baboons: genetic, environmental and demographic influences. Molecular Ecology *17*, 2026-2040.

78. Altmann, J., Altmann, S.A., and Hausfater, G. (1978). Primate infant's effects on mother's future reproduction. Science *201*, 1028-1030.

79. Altmann, J., and Alberts, S.C. (2003). Variability in reproductive success viewed from a life‐history perspective in baboons. American Journal of Human Biology *15*, 401-409.

80. Lea, A.J., Learn, N.H., Theus, M.J., Altmann, J., and Alberts, S.C. (2014). Complex sources of variance in female dominance rank in a nepotistic society. Animal behaviour *94*, 87-99.

81. Vilgalys, T.P., Rogers, J., Jolly, C.J., Mukherjee, S., and Tung, J. (2018). Evolution of DNA methylation in Papio baboons. Molecular biology and evolution *36*, 527-540.

82. Irizarry, R.A., Ladd-Acosta, C., Wen, B., Wu, Z., Montano, C., Onyango, P., Cui, H., Gabo, K., Rongione, M., and Webster, M. (2009). The human colon cancer methylome shows similar hypo-and hypermethylation at conserved tissue-specific CpG island shores. Nature genetics *41*, 178.

83. Consortium, E.P. (2012). An integrated encyclopedia of DNA elements in the human genome. Nature *489*, 57.