1   Manuscript title:
2   **Specialized DNA structures act as genomic beacons for integration by evolutionarily**
3   **diverse retroviruses**
4

5   Hinissan P. Kohio[1‡], Hannah O. Ajoge[1‡], Macon D. Coleman[1], Emmanuel Ndashimye[1], Richard M.
6   Gibson[1], Eric J. Arts[1] and Stephen D. Barr[1*]
7
8   [1] Western University, Schulich School of Medicine and Dentistry, Department of Microbiology and
9   Immunology, Dental Sciences Building Room 3007, London, Ontario, Canada.
10
* To whom correspondence should be addressed. Tel: 01-519-661-3438; Fax: 01-519-661-3499;
Email: stephen.barr@uwo.ca

‡ The authors wish it to be known that, in their opinion, the first 2 authors should be regarded as joint
first authors

ABSTRACT

Retroviral integration site targeting is not random and plays a critical role in expression and long-term
survival of the integrated provirus. To better understand the genomic environment surrounding
retroviral integration sites, we performed an extensive comparative analysis of new and previously
published integration site data from evolutionarily diverse retroviruses from seven genera, including
different HIV-1 subtypes. We showed that evolutionarily divergent retroviruses exhibited distinct
integration site profiles with strong preferences for non-canonical B-form DNA (non-B DNA). Whereas
all lentiviruses and most retroviruses integrate within or near genes and non-B DNA, MMTV and ERV
integration sites were highly enriched in heterochromatin and transcription-silencing non-B DNA
features (e.g. G4, triplex and Z-DNA). Compared to *in vitro*-derived HIV-1 integration sites, *in vivo*-
derived sites are significantly more enriched in transcriptionally silent regions of the genome and
transcription-silencing non-B DNA features. Integration sites from individuals infected with HIV-1
subtype A, C or D viruses exhibited different preferences for non-B DNA and were more enriched in
transcriptionally active regions of the genome compared to subtype B virus. In addition, we identified
several integration site hotspots shared between different HIV-1 subtypes with specific non-B DNA
sequence motifs present at these hotspots. Together, these data highlight important similarities and
differences in retroviral integration site targeting and provides new insight into how retroviruses
integrate into genomes for long-term survival.

Retrovirus integration site profiles

Distance from integration site

Non-B DNA

Spumavirus (FV)

Deltaretrovirus (HTLV-1)

FIV

SIV

HIV

Lentivirus

Gammaretrovirus (MLV)

Alpharetrovirus (ASLV)

Betaretrovirus (MMTV)

Subtype

Integration in non-B DNA

Integration site hotspots

Graphical Abstract:
Schematic comparing integration site profiles from evolutionarily diverse retroviruses. Upper left, heatmaps showing the fold-enrichment (blue) and fold-depletion (red) of integration sites near non-B DNA features (lower left). Lower right, circa plot showing integration site hotspots shared between HIV-1 subtype A, B, C and D virus.

INTRODUCTION

Retroviruses are divided into seven genera *alpha-*, *beta-*, *gamma-*, *delta-*, and *epsilon-Retroviridae*, *Spumaviridae*, and *Lentiviridae*). Following entry of a retrovirus into cells, the viral RNA genome is converted into double-stranded complimentary DNA (cDNA). In the cytoplasm, the cDNA associates with several viral and host proteins to form a pre-integration complex (PIC). Soon after completion of cDNA synthesis, the viral DNA ends are primed by the enzyme integrase in a process called 3' processing. After docking with a host chromosome, the viral cDNA undergoes a strand transfer reaction resulting in the insertion of the retroviral genome into the chromosome. Host repair enzymes are then thought to complete integration (*1*). Insertion of the retroviral genome results in a persistent life-long infection. Selection of integration sites in the genome by the PIC is not random. The gamma- and deltaretroviruses (e.g. murine leukemia virus (MLV), human T cell leukemia virus Type 1 (HTLV-1)) and foamy virus (FV) favor integration around *transcription start sites* (*TSS*) (*2–4*). Alpharetroviruses (e.g. avian sarcoma leukosis virus (ASLV)) and human endogenous retroviruses (ERVs) show no strong preferences, with integration only slightly favored in transcription units or the 5' end of genes (*5–8*).

Lentiviruses such as human immunodeficiency virus type 1 (HIV-1), simian immunodeficiency virus (isolated from a pig-tailed macaque) (SIV_{mne}) and feline immunodeficiency virus (FIV) strongly favor integration in active transcription units (*9–11*). HIV integrations sites are also associated with regions of high G/C content, high gene density, short introns, high frequencies of short interspersed nuclear elements (*SINEs*) (e.g. Alu repeats), low frequencies of long interspersed nuclear elements (*LINEs*), and characteristic epigenetic modifications (*9*, *12*). Thus far, integration site analyses have only been conducted on HIV-1 subtype B infections. Based on phylogenetic analyses of full-length genomic sequences, HIV-1 isolates are classified into four distinct groups: group M, N, O and P (*13*). HIV-1 M group accounts for the majority of the global pandemic and is subdivided into nine subtypes or clades (A, B, C, D, F, G, H, J and K). Additionally, several circulating recombinant forms (CRF) and unique recombinant forms (URF) have been identified, which are the result of a recombination event between two or more different subtypes. HIV-1 geographical prevalence is extremely diverse. Subtype C represents more than 50% of the infection worldwide and is prevalent in Africa and Asia. Subtype B infection is common in the Americas, Europe, Australia, and part of South Asia, Northern Africa and the Middle East. Subtypes A, D, F, G, H, J and K occur mostly in Sub-Saharan Africa, whereas infections with groups N, O and P have been found in confined regions of West-Central Africa.

Several models, not mutually exclusive, have been proposed to explain integration site selection. In the chromatin accessibility model, the structure of chromatin influences accessibility of target DNA sequences to PICs. *In vivo* target DNA is not expected to be naked but rather wrapped in nucleosomes. Wrapping target DNA in nucleosomes does not reduce integration, but instead creates hotspots for integration at sites of probable DNA distortion (*14*, *15*). Distortion of DNA in several other protein-DNA complexes has also been shown to favour integration in the major grooves facing outwards from the nucleosome core (*16*, *17*). Although chromatin structure can facilitate integration, chromatin accessibility cannot solely explain the differences observed in integration site preferences.

The protein tethering model suggests that a cellular protein, potentially specific for each retroviral genera, act as tethering factors between chromatin and the PIC. The most characterized tethering factor identified to date is lens epithelium-derived growth factor and co-factor p75 (LEDGF/p75) (also known as PSIP1/p75) (18–20). LEDGF/p75 interacts with HIV integrase and tethers the PIC to genomic DNA in transcriptionally active genes marked by specific histone modifications such as H3K20me1, H3K27me1 and H3K36me3. In similar fashion, host protein bromodomain/extraterminal domain proteins (BETs) bound to acetylated histones (e.g. H327ac and H3K9ac) interact with MLV integrase and tethers the PIC to genomic DNA in transcriptionally active promoters, enhancers and super enhancers (21–24). LEDGF/p75 and BET depletion studies demonstrated that integration still occurs but with reduced efficiency and an altered integration site selection profile (20, 22–28). Several cellular proteins have been proposed to facilitate integration including barrier to autointegration factor (BAF), high mobility group A1 (HMGA1), integrase interactor 1 (Ini-1), and heat shock protein 60 (Hsp60) (29–31). Some of these proteins might contribute to PIC function by coating and condensing the viral DNA, thereby assisting the assembly of the viral nucleoprotein complexes. Several cellular chromatin proteins have also been suggested to influence integration such as Ini-1, EED, SUV39H1, and HP1γ (reviewed in references (12, 32, 33)). Cleavage and polyadenylation specificity factor 6 (CPSF6) facilitates HIV-1 PIC import and helps direct the PIC to the nuclear interior to locate gene-dense euchromatin for integration (34–38). Recently, Apolipoprotein B Editing Complex (APOBEC3) was identified as a potential new host factor that also influences integration site selection by promoting a more transcriptionally silent integration site profile (39). Moreover, just as DNA-binding proteins can promote integration, they can also block access of integration complexes, creating regions refractory for integration (16, 40, 41). Binding to different factors or differential binding affinity to a factor by the PIC could modulate integration site selection between lentivirus types or even between different HIV-1 subtypes.

Previous analyses of primary DNA sequences (~20 base pairs (bp)) flanking integration sites only revealed a weak consensus motif (42). We previously assessed a broader window of primary sequence (80 bp) around HIV-1 integration sites discovered that HIV-1 integration sites were highly enriched near specialized genomic features called non-B DNA (43). Non-B DNA form functional secondary structures in our genome formed by specific nucleotide sequences that exhibit non-canonical DNA base pairing. At least 10 non-B DNA conformations are identified including inverted repeats, direct repeats, mirror repeats, short-tandem repeats, guanine-quadruplex (G4), A-phased, cruciform, slipped, triplex and Z-DNA (44, 45). Here we present a comparative analysis of integration site profiles of evolutionarily diverse retroviruses and identify striking similarities and differences among all retroviruses, especially towards non-B DNA.

MATERIAL AND METHODS

Ugandan and Zimbabwean cohort description. Details pertaining to the Uganda study population have been reported previously (46–49). Briefly, women who became HIV infected while participating in the Hormonal Contraception and Risk of HIV Acquisition Study in Uganda were enrolled upon

primary infection with HIV-1 into a subsequent study, the Hormonal Contraception and HIV-1 Genital
Shedding and Disease Progression among Women with Primary HIV Infection (GS) Study. Ethical
approval was obtained from the Institutional Review boards (IRBs) from the Joint Clinical Research
Centre and UNST in Uganda, from University of Zimbabwe, from the University Hospitals of
Cleveland, and recently, from Western University. All adult subjects provided written informed consent
and no child participants were included in the study. Protocol numbers and documentation of these
approvals/renewals are available upon request. Blood and cervical samples were collected every
month for the first six months, then every three months for the first two years, and then every six
months up to 9.5 years. Women who had CD4 lymphocyte counts of 200 cells/ml and/or who
developed severe symptoms of HIV infection (WHO clinical stage IV or advanced stage III disease)
were offered combination antiretroviral therapy (cART) and trimethoprim-sulfamethoxazole (for
prophylaxis against bacterial infections and *Pneumocystis jeroveci* pneumonia).

HIV-1 integration site library. Total genomic DNA was extracted from PBMCs using the QIAmp DNA
mini kit (Qiagen) and processed for integration site analysis and sequenced using the Illumina MiSeq
platform as previously described (*43*, *50*). Samples were sequenced at the London Regional
Genomics Centre/Robarts Research Institute (Western University, Canada) and Case Western
Reserve University (USA).

Computational analysis*.* Fastq sequencing reads were quality trimmed and unique integration sites
identified using our in-house bioinformatics pipeline Barr Lab Integration Site Identification Pipeline
(BLISIP version 2.9) and included the following updates: bedtools (v2.25.0), bioawk (awk version
20110810), bowtie2 (version 2.3.4.1), and restrSiteUtils (v1.2.9) (*39*, *43*). All non-B DNA motifs were
defined according to previously established criteria (*51*). The G-quadruplex algorithm identifies four or
more individual G-runs of at least three nucleotides in length. The algorithm requires at least one
nucleotide between each run and considers up to seven nucleotides as spacer, including guanines
(*51*). Lamina-associated domains (*LADs*) were retrieved from http://dx.doi.org/10.1038/nature06947
(*52*). HIV-1 LTR-containing fastq sequences were identified and filtered by allowing up to a maximum
of five mismatches with the reference NL4-3 LTR sequence and accepted if the LTR sequence had
no match with any region of the human genome (GRCh37/hg19). Matched random control integration
sites (28,800 in total) were generated by matching each experimentally determined site with 50
random sites *in silico* that were constructed to be the same number of bases from the restriction site
as was the experimental site, as previously described (*53*). All genomic sites in each dataset that
hosted two or more sites (i.e. identical sites) were collapsed into one unique site for the analysis.
Sites that could not be unambiguously mapped to a single region in the genome were excluded from
the study. Integration site profile heatmaps were generated using our in-house python program
BLISIP heatmap (BHmap v1.0).

Data and software availability. The sequences reported in this paper will be deposited in the

National Center for Biotechnology Information Sequence Read Archive (NCBI SRA) upon acceptance

for publication.

RESULTS

Integration site dataset acquisition and analyses

Previous integration site analyses of evolutionarily diverse retroviruses identified three general

integration site profiles in which retroviruses can be grouped. For example, HIV-1, SIV$_{mne}$ and FIV

integrate into transcriptionally active genes; MLV, HTLV-1 and FV integrate near *TSS*; ASLV, HTLV

and MMTV show little preference for any genomic feature. The integration site profiles of these

evolutionarily diverse retroviruses with respect to non-B DNA was previously unknown. To determine

integration site profiles of these viruses, including *endogenous retroviruses* (*ERVs*) present within the

human genome, we analyzed ~169,910 integration sites from previously published datasets (**Table S1**).

To allow for a more comparable analysis of integration site profiles, we updated these integration site

profiles using the human genome assembly GRCh37/19. Our update also included an assessment of

genomic features not included in some of the previous retroviral integration site studies. We used our

previously developed in-house bioinformatics pipeline called the Barr Lab Integration Site Pipeline

(BLISIP) to generate integration site profiles from the different retroviral integration site datasets (*39,*

43, 50). BLISIP measures integration site enrichment near the genomic features CpG islands, *DNAseI*

hypersensitivity sites (*DHS*), *ERVs*, heterochromatic DNA regions (e.g. *lamin-associated domains*

(*LADs*) and satellite DNA), *SINEs* and *LINEs* respectively, *low complexity repeats* (*LCRs*), oncogenes,

genes, simple repeats and *TSS*. In addition, BLISIP measures enrichment near the non-B DNA features

A-phased motifs, cruciform motifs, direct repeats, G4 motifs, inverted repeats, mirror repeats, short

tandem repeats, slipped motifs, triplex motifs and Z-DNA motifs. Our analyses focused on unique

integration sites and excluded sites arising from clonal expansion, sites falling in repeat regions that

could not be uniquely identified, and regions that could not be confidently placed on a specific

chromosome (e.g. ChrUn). Enrichment of integration sites within genomic features was determined by

comparing the proportion of sites with either a matched random control (MRC) to account for restriction

site bias in the cloning procedure during library construction, or a random control (RC) for comparison

of datasets that used DNA shearing/fragmentation during library construction (**Table S1**).

Evolutionarily divergent retroviruses exhibit distinct integration site profiles

Integration sites were quantified and placed in five bins based on their distance from each

genomic feature (within the feature, 1-499 bp, 500-4,999 bp, 5,000-49,999 bp, and >49,000 bp).

Heatmaps from each retrovirus showing the fold-enrichment and fold-depletion in each bin were

compared to MRC or RC (**Figure 1A**). Consistent with previous studies (*9–11*), HIV-1, SIV$_{mne}$ and FIV

integration sites are significantly enriched within genes (64%, 84%, 90% respectively) (*P* < 0.0001)

(Figure 1A and 1B, Table S2). Our analysis of HTLV-1, ASLV and MLV also agreed with previous

reports, confirming that these viruses exhibit only modest preferences for integration within *genes* with

54%, 56% and 56% of integration sites found within *genes* respectively **(Figure 1A and 1B, Table S2)**

(*54, 55*)**.** In contrast, MMTV, FV and ERVs showed no preference for integration into genes (47%, 43%

and 32% respectively). No retrovirus showed a preference for integration directly into *TSS*; however,

MLV, HTLV-1 and FV showed significant enrichment of integration sites near *TSS* and CpG islands (*P*

<0.0001) **(Figure 1A, Table S2)**. Repetitive elements, such as *LINEs*, *SINEs*, *ERVs* (e.g.

retrotransposons), satellite DNA, simple repeats (e.g. microsatellites), and *LCRs* account for nearly half

of the human genome sequence. However, no strong preference for integration into these regions was

observed for any of the retroviruses except for HIV-1 and FV which targeted *SINEs*, and FV, MMTV

and HTLV-1 which targeted satellite DNA **(Figure 1A).** Pairwise analyses of the different integration

site profile preferences (fold enrichment and depletion values) showed that HIV-1, MMTV and FIV had

the most similar integration site preferences for genomic features overall, whereas SIV, MLV and ERVs

had the least similarity to the other retroviruses (**Figure 1C**).

MMTV and ERVs strongly favour integration into heterochromatin

The nuclear architecture influences HIV-1 integration site selection and proviral expression (*52,*

56). HIV-1 strongly disfavors integration into heterochromatin positioned in *LADs* at the nuclear

periphery, although some integration does occur in these regions, contributing to the latent reservoir

(*57*). We asked whether other retroviruses similarly disfavor integration into *LADs*. Like HIV-1, most

retroviruses significantly disfavored integration into *LADs* with only 11-28% of sites falling within *LADs*

(Figure 1D, Table S2). In contrast, 48% of MMTV and 44% of ERV integration sites were significantly

enriched in *LADs*.

Evolutionarily divergent retroviruses target non-B DNA for integration

Non-B DNA is a genomic correlate of HIV-1 integration site selection, but its influence on

integration site targeting of other retroviruses was previously unknown (*43*). Analysis of the different

retroviral integration site profiles showed that all retroviruses exhibited enriched integration near non-B

DNA (**Figure 2A and Table S3**). FV, HTLV-1 and SIV$_{mne}$ were the only retroviruses with an enrichment

of sites within 50 bp of multiple non-B DNA features, whereas other retroviruses tended to integrate

more distal (100-500 bp) to the features. HIV-1 integration sites were enriched near all non-B DNA

features except A-phased DNA. SIV$_{mne}$ and FIV sites were enriched near inverted repeats, mirror

repeats and short tandem repeats. HTLV-1 sites were enriched near A-phased, cruciform and inverted

repeats. FV sites were enriched near G4 and Z-DNA. MLV sites were enriched near G4, triplex and Z-

DNA. ASLV sites were enriched in slipped and triplex DNA. MMTV sites were enriched in short-tandem

repeats, G4, slipped, triplex and Z-DNA. ERV sites were enriched near cruciform and triplex DNA.

Pairwise analyses of the different integration site profiles showed that HIV-1 and FV had the most similar

preferences for targeting non-B DNA, whereas SIV$_{mne}$, ERVs and FIV had the least similarity to the

other retroviruses (**Figure 2B**). Taken together, these data show that evolutionarily diverse retroviruses

target non-B DNA for integration and that each virus exhibits distinct preferences for certain non-B DNA

features.

HIV-1 Integration site profiles differ between *in vitro*- and *in vivo*-derived datasets

Despite a large number of integration sites from HIV-1 infected individuals, many of the previous integration site studies have been performed on infections done *in vitro* (*12, 58*). To determine if the HIV-1 integration site profiles from *in vitro*-derived infections differed from those from *in vivo*-derived infections, we analyzed and compared nine previously published HIV-1 integration site datasets from publicly available databases, totalling 13,601 *in vitro*-derived sites and 139,480 *in vivo*-derived sites **(Table S1)**. As expected, integration sites were significantly enriched in genes compared to the random controls in both the *in vitro*- and *in vivo*-derived datasets; however, only 62% of total *in vivo*-derived integration sites were in genes compared to 83% in the *in vitro* dataset ($P < 0.0001$) **(Figure 3A, 3B and Tables S4)**. In contrast with the *in vitro* dataset, integration sites in the *in vivo*-derived dataset were highly enriched near most non-B DNA except A-phased DNA **(Figure 3C and Table S4)**. Notably, *in vivo*-derived sites were significantly more enriched near G4, triplex and Z-DNA than *in vitro*-derived sites **(Figure 3C, 3D and Table S4)**. Together, these data show that there are substantial differences in HIV-1 integration site targeting preferences between *in vitro*-derived and *in vivo*-derived datasets.

Integration site profiles differ in individuals infected with HIV-1 subtype A, B, C or D

Thus far, integration site profiles have been extensively analyzed for HIV-1 subtype B infections, which represents only ~10% of the infections worldwide. We asked if the integration site profiles from individuals infected with HIV-1 non-subtype B virus were similar to those infected with subtype B virus. In the integrase enzyme, the amino acid difference between subtypes is as high as 16% (subtype B vs. C) but typically less than 8% in IN of HIV-1 isolates of a specific subtype. This level of amino acid diversity is the highest among the enzymes encoded by the HIV-1 *pol* gene. The greatest diversity within HIV-1 IN is found in the C-terminal domain (CTD), which is involved in genomic DNA binding. Thus, it is reasonable to propose that we would observe differences in selectivity for integrations in non-B form DNA. Genomic DNA was isolated from peripheral blood mononuclear cells (PBMCs) from a cohort of women in Uganda and Zimbabwe infected with HIV-1 subtype A, C or D and used to generate integration site libraries. Integration site profiles were generated from a total of 48 infected individuals (16 subtype A, 19 subtype C and 13 subtype D) and compared to the integration site profile from 14 individuals infected with subtype B virus generated from previously published datasets **(Table S5)** (*59, 60*)**.**

As previously observed with HIV-1 subtype B infections, integration sites from all HIV-1 subtype viruses were enriched in genes (**Figure 4A and 4B and Table S6**). Notably, subtypes A, C and D had significantly more integration sites in genes compared to subtype B (A:82%, B:63%, C:71%, D:78%) (**Figure 4C**). Integration sites for all subtypes were enriched near *DHS*; however, subtypes A, C and D exhibited a weaker preference compared to subtype B. In addition, subtypes A, C and D had significantly less integration in heterochromatin (e.g. *LADs*) and *SINEs* compared to subtype B, which likely reflects their higher integration into genes. Pairwise analyses of the different integration site profiles showed that subtypes C and D, and B and D, shared the most similarity to each other, and subtype A differed the most from the other subtypes (**Figure 4D**). This relates in part to the sequence distance between IN coding regions of subtypes A, B, C, and D, with B and D being most genetically

similar (**Figure 4E and 4F**). However, it is important to stress that any slight differences in integration
profiles could also relate to discreet single amino acid differences between subtypes, especially in the
CTD domain and/or the LEDGF binding domain.
Analysis of the non-B DNA integration site profiles from the different subtype viruses showed
enriched integration near non-B DNA (**Figure 5A, 5B and Table S7**). Notably, subtypes A and C had
more integration sites near A-phased DNA compared to subtypes B and D. Subtypes A, C and D virus
had significantly less sites near G4 DNA compared to subtype B virus. In addition, subtype C had
significantly less sites near triplex and Z-DNA. Pairwise analysis of the different non-B DNA integration
site profiles showed that the profiles of all subtypes differed substantially from each other, with subtypes
B and D showing slightly more similarity to each other compared to the other subtypes (**Figure 5C**).
Together, these data show that the integration site profiles from individuals infected with HIV-1 subtype
B differs from those infected with non-subtype B virus, especially with respect to their preferences for
non-B DNA. As discussed below, these slight but significant differences in selection for integration at
different non-B DNA features may relate to discreet amino acid differences in the CTD domains of these
IN that evolved in different subtypes. Although there are trends suggesting impact by discreet amino
acid differences, for significance, we need to analyze more integration patterns by more subtype A, C,
and D infections with known IN sequences.
288

Integration hotspots are shared between HIV-1 subtypes
The concept of an HIV-1 integration "hotspot" was introduced to describe areas of the
genome where integrations accumulate more than expected by chance in the absence of any
selection process (*9, 61*). We catalogued and analyzed 1,000 bp windows containing two or more
integration sites from each HIV-1 subtype dataset. This yielded a total of 10,892 hotspots between all
subtypes (A: 41/429, 9.6%; B: 10,762/118,727, 9.1%; C: 74/484, 15.3%; D: 15/323, 4.6%; and MRC:
2/2982, 0.07%) (**Figure 6A**). 16% of the hotspots were shared between subtypes A and C, 14%
between subtypes A and D and 10% between subtypes C and D. Despite the large number of
hotspots identified in the subtype B dataset, less than 0.03% of the hotspots were shared between
subtype B and the other subtypes. Comparison of hotspots revealed one hotspot in the *tbc1d5* gene
that was shared by all subtypes (**Figure 6B and 6C**).
To identify potential local genomic sequences that may serve as 'beacons' for increased
integration, we analyzed a nucleotide window of 100 bp upstream and 100 bp downstream of each
integration site and generated a consensus sequence using DiffLogo (**Figure 6D**) (*62*). Comparison of
the sequence motifs surrounding integration sites located in hotspots with the sequence motifs from
sites not located in hotspots revealed the presence of a slipped DNA motif at the integration site in
each of the HIV-1 subtype datasets. In addition, a G4 motif was located ~40-100 bp downstream from
the integration site in each of the different HIV-1 subtype datasets. Subtypes B and C also had a
triplex motif and subtype B had a short tandem repeat located downstream of the integration site.
Together, these data show that sites located in integration hotspot regions are located in slipped DNA
motifs flanked downstream by other non-B DNA motifs.
310

DISCUSSION

Here we showed that evolutionarily diverse retroviruses from all seven genera exhibit strong preferences for integration within or near non-B DNA features. Whereas all lentiviruses and most retroviruses integrate within or near genes and a preference for non-B DNA, we show that MMTV and ERV integration sites are highly enriched in heterochromatin but with still a preference for non-B DNA. Nonetheless, each retrovirus has distinct preferences for integration at specific types of non-B DNA. Highly distinct preferences for these non-B motifs among different HIV-1 subtypes is less apparent and yet, divergent IN evolution may still result in some differential selection/inhibition of HIV DNA integration in host cellular DNA. Finally, we showed that certain regions of the genome attract more integration sites than others and that the integration sites in these regions are located in slipped DNA motifs and flanked downstream by other non-B DNA motifs.

All retroviruses exhibited enriched integration near non-B DNA with specific non-B DNA features (e.g. slipped, G4, triplex and short-tandem repeat DNA) serving as integration site hotspots. How retroviral PICs recognize non-B DNA for integration is not fully understood. Non-B DNA motifs are enriched near endogenous retroviruses, are near sites of genomic variability, and are targets for preferential homologous recombination (>20-fold) in human cells (*63–65*). It is possible that integrase itself, components of the PIC and/or host proteins can bind non-B DNA directly and promote integration in these regions. We recently showed that PIC-binding host factors APOBEC3 (*39*), LEDGF/p75 and CPSF6 (submitted) influence the distribution of integration sites near non-B DNA features, potentially indicating that a host-directed mechanism is involved. Intriguingly, APOBEC3G binding and deamination hotspots also comprise slipped DNA motifs (*66, 67*). It is possible that viral integrase is also involved in recognizing non-B DNA given that HIV-1 integrase has been shown to bind directly to G4 DNA (*68*).

Analysis of integration site profiles after acute infection showed that HIV-1, FIV and SIV$_{mne}$ had the strongest preference for integrating into transcriptionally active regions of the genome, whereas MMTV, FV and ERVs had the strongest preference for transcriptionally inactive regions. The targeting of transcriptionally active regions of the genome is believed to help maximize provirus expression for establishing infection and promoting its spread (*9*). Over time, especially with the help of antiretroviral drugs, the immune system likely clears these high virus-producing cells leaving behind infected cells that produce little to no virus that escape detection by the immune system. The integration profile of these latently infected cells showed that sites are located in more transcriptionally silent regions of the genome, similar to the profile shown here for ERVs and after acute infection with MMTV and FV. Consistent with the latter observation, latently infected cells have reduced integration in genes and increased integration in *SINEs* and heterochromatin, typically found in gene deserts or transcriptionally silent regions (*57, 60*). Transcriptionally silent regions of the genome are characterized by several features. For example, *LADs* represent a repressive chromatin environment tightly associated with the nuclear periphery (*56, 57*). *SINEs* (e.g. Alu repeats) and other transposed sequences are known to serve as direct silencers of gene expression due to their repressed chromatin marks (histone H3 methylated at Lys 9) (*69, 70*). Moreover, non-B DNA structures such as G4, cruciform, triplex and Z-DNA have been shown to silence expression of adjacent genes (*65, 71, 80, 72–79*). The ability to target

silent regions of the genome for integration could be a long-term survival mechanism for both
retroviruses and their hosts. By integrating into transcriptionally silent regions of the genome, these
viruses can minimize their expression and avoid detection by the immune system. As an extreme
example, endogenous retroviruses, some of which are ancient viruses that have been co-evolving with
their hosts for hundreds of millions of years (e.g. ERVs and FV), are commonly integrated in
transcriptionally silent regions of the genome away from genes (Figures 1 and 2) (*81–83*). The host also
benefits with a lower risk of disease. This is seen with FV which persists in their primate hosts in the
absence of disease, and MMTV which typically does not cause cancer unless it integrates near an
oncogene (*84*). It is currently unknown if integration in transcriptionally silent regions of the genome is
driven by retroviruses, the host or both. Interestingly, host APOBEC3G and APOBEC3F appear to
promote a transcriptionally silent integration site profile of HIV-1, suggesting that the host may indeed
contribute to this silent phenotype (*39*).

Genomic position effects have been shown to influence HIV-1 expression, which can ultimately
impact disease progression (*57, 85–87*). As described above, HIV-1 infection of a host cell results in
preferential integration in actively transcribing genes, due in part to reduced chromatin structure. Thus,
it is not surprising to observe this finding following *in vitro* infections. However, even *in vitro*, there is
evidence of preferential integration near non-B DNA motifs. Within a human host, enhanced HIV-1
integration within genes remains evident but now, there is greater enrichment of those exact integration
sites being near transcription-silencing non-B DNA motifs, suggesting an enrichment in the pool of HIV-
1 proviral DNA within transcriptionally silent sites in the genome. As described above, this may be the
consequence of the immune system consistently clearing away cells propagating HIV-1 while the cells
with silent integrants are slowly enriched. HIV-1 integration sites from these *in vitro* experiments were
typically analyzed from cells infected for 2-3 days, whereas the *in vivo*-derived integration sites were
obtained from chronically infected individuals whose cells contain integrated proviruses that have
undergone more than 2 months of selection in the host. Consistent with previous findings (*57, 60*), HIV
integrants were found enriched within features associated with reduced expression such as *LADs*,
satellite DNA and *SINEs* when compared to *in vitro*-derived sites. Moreover, we identified a sequence-
based integration site bias towards non-B DNA, particularly transcription silencing G4, triplex and Z-
DNA.

Much of our knowledge of HIV-1 integration site selection has come from studies using subtype
B virus. Our analyses showed that HIV-1 non-subtype B viruses have a much stronger preference for
integrating into transcriptionally active regions of the genome compared to subtype B virus. This was
characterized by increased integration in genes and decreased integration in genomic features
associated with transcriptional silencing such as *LADs*, *SINEs*, G4, triplex and Z-DNA. Despite the
enriched proviral integration into non-B DNA among all HIV-1 subtypes, there was some integration site
preferences that differed between each of the HIV-1 subtypes. In the integrase coding region, amino
acid differences between subtypes are as a high as 16% (subtype B vs. C) but typically less than 8%
IN diversity among HIV-1 isolates of one subtype (*13, 88, 89*). This level of amino acid diversity is the
highest among the enzymes encoded by the HIV-1 *pol* gene. The greatest diversity within HIV-1 IN is
found in the CTD, which is involved in genomic DNA binding. Thus, it is reasonable to suspect that HIV-

1 subtypes may target different non-B DNA motifs with differential selectivity. Despite the multiple

substitutions that segregate HIV-1 variants of each subtype, there may be specific amino acid

differences in one subtype versus another that impact preference for integration sites. For example, the

isoleucine found at position 20 in subtype C and D versus the threonine in subtype A and B in the IN

CTD may relate the greater similarity of integration site selection of subtypes C and D (Figure 4E and

4F). Despite the closer genetic relationship between HIV-1 subtype B and D across the genome, similar

genetic distances separating subtypes A, B, C, and D are within the IN coding region. Polymorphisms

in HIV-1 integrase have been reported that retarget integration away from gene dense regions, which

also correlated with increased disease progression and virulence (*87*). This suggests that

characteristics of integrase itself is a driver of integration site selection in retroviruses. We are currently

exploring if subtype-specific polymorphisms in integrase may account for differences in the targeting of

transcriptionally silent regions of the genome and of non-B DNA.

In conclusion, we identified non-B DNA as a feature surrounding integration sites that are

targeted differentially by evolutionarily diverse retroviruses. We have also presented the first

comparative look at the integration site profiles of several HIV-1 non-subtype B viruses and showed

that they differed from HIV-1 subtype B profiles. Together, this data highlights important similarities and

differences in retroviral integration site targeting that can be used in future studies to better understand

the evolution of retroviral integration site targeting and how these viruses integrate into our genomes

for long-term survival.

FUNDING

This work was supported by the Canadian Institutes of Health Research (CIHR) Operating Grants

[IBC-150406 and HBF-143164] to S.D.B.; and in part by CIHR [385787, 377790 to E.J.A]; Canada

Research Chair Tier 1 [230811 to E.J.A.]; National Institute of Allergy and Infectious Diseases-

National Institutes of Health [AI49170 to E.J.A].

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

1.   R. Daniel *et al.*, Evidence that stable retroviral transduction and cell survival following DNA

integration depend on components of the nonhomologous end joining repair pathway. *J Virol*.

78, 8573–8581 (2004).

2.   X. Wu, Y. Li, B. Crise, S. M. Burgess, Transcription start regions in the human genome are

favored targets for MLV integration. *Science*. **300**, 1749–51 (2003).

3.   B. Felice *et al.*, Transcription Factor Binding Sites Are Genetic Determinants of Retroviral
Integration in the Human Genome. *PLoS One*. **4**, e4571 (2009).

4.   G. D. Trobridge *et al.*, Foamy virus vector integration sites in normal human cells. *Proc. Natl.*
Acad. Sci. U. S. A. **103**, 1498–503 (2006).

5.   S. D. Barr, J. Leipzig, P. Shinn, J. R. Ecker, F. D. Bushman, Integration targeting by avian
sarcoma-leukosis virus and human immunodeficiency virus in the chicken genome. *J. Virol.*
79, 12035–12044 (2005).

6.   R. Mitchell *et al.*, Retroviral DNA Integration: ASLV, HIV, and MLV Show Distinct Target Site
Preferences. *PLoS Biol.* **2**, E234 (2004).

7.   A. Narezkina *et al.*, Genome-wide analyses of avain sarcoma virus integration sites. *J.Virol.*
78, 11656–11663 (2004).

8.   T. Brady *et al.*, Integration target site selection by a resurrected human endogenous retrovirus.
Genes Dev. **23**, 633–642 (2009).

9.   A. Schroder *et al.*, HIV-1 Integration in the Human Genome Favors Active Genes and Local
Hotspots. *Cell*. **110**, 521–529 (2002).

10.   B. Crise *et al.*, Simian Immunodeficiency Virus Integration Preference Is Similar to That of
Human Immunodeficiency Virus Type 1. *J. Virol.* **79**, 12199–12204 (2005).

11.   Y. Kang *et al.*, Integration Site Choice of a Feline Immunodeficiency Virus Vector. *J. Virol.* **80**,
8820–8823 (2006).

12.   F. Bushman *et al.*, Genome-wide analysis of retroviral DNA integration. *Nat. Rev. Microbiol.* **3**,
848–58 (2005).

13.   B. Taylor *et al.*, The Challenge of HIV-1 Subtipe Diversity. *N. Engl. J. Med.* **358**, 1590–1602
(2008).

14.   D. Pruss, F. D. Bushman, A. P. Wolffe, Human immunodeficiency virus integrase directs
integration to sites of severe DNA distortion within the nucleosome core. *Proc. Natl. Acad. Sci.*
U. S. A. **91**, 5913–7 (1994).

15.   D. Pruss, R. Reeves, F. D. Bushman, A. P. Wolffe, The influence of DNA and nucleosome
structure on integration events directed by HIV integrase. *J. Biol. Chem.* **269**, 25031–41
(1994).

16.   Y. C. Bor, F. D. Bushman, L. E. Orgel, In vitro integration of human immunodeficiency virus
type 1 cDNA into targets containing protein-induced bends. *Proc. Natl. Acad. Sci. U. S. A.* **92**,
10334–10338 (1995).

17.   H.-P. Muller, H. E. Varmus, DNA bending creates favored sites for retroviral integration: an
explanation for preferred insertion sites in nucleosomes. *EMBO J.* **13**, 4704–4714 (1994).

18.   A. Engelman, P. Cherepanov, The lentiviral integrase binding protein LEDGF/p75 and HIV-1
replication. *PLoS Pathog.* **4**, e1000046 (2008).

19.   E. M. Poeschla, Integrase, LEDGF/p75 and HIV replication. *Cell. Mol. Life Sci.* **65**, 1403–24
(2008).

20.   M. Llano *et al.*, An essential role for LEDGF/p75 in HIV integration. *Science*. **314**, 461–4
(2006).

21.   R. C. Larue *et al.*, Bimodal high-affinity association of Brd4 with murine leukemia virus
integrase and mononucleosomes. *Nucleic Acids Res.* **42**, 4868–4881 (2014).

22.   A. Sharma *et al.*, BET proteins promote efficient murine leukemia virus integration at
transcription start sites. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 12036–41 (2013).

23.   S. S. Gupta *et al.*, Bromo- and extraterminal domain chromatin regulators serve as cofactors
for murine leukemia virus integration. *J. Virol.* **87**, 12721–36 (2013).

24.   S. Aiyer *et al.*, Altering murine leukemia virus integration through disruption of the integrase
and BET protein family interaction. *Nucleic Acids Res.* **42**, 5917–5928 (2014).

25.   M.-C. Shun *et al.*, LEDGF/p75 functions downstream from preintegration complex formation to
effect gene-specific HIV-1 integration. *Genes Dev.* **21**, 1767–78 (2007).

26.   H. M. Marshall *et al.*, Role of PSIP1/LEDGF/p75 in lentiviral infectivity and integration
targeting. *PLoS One.* **2**, e1340 (2007).

27.   A. Ciuffi *et al.*, A role for LEDGF/p75 in targeting HIV DNA integration. *Nat. Med.* **11**, 1287–9
(2005).

28.   L. Vandekerckhove *et al.*, Transient and stable knockdown of the integrase cofactor
LEDGF/p75 reveals its role in the replication cycle of human immunodeficiency virus. *J. Virol.*
80, 1886–96 (2006).

29.   S. P. Goff, Host factors exploited by retroviruses. *Nat. Rev.* **5**, 253–263 (2007).

30.   A. Engelman, The roles of cellular factors in retroviral integration. *Curr. Top. Microbiol.*
Immunol. **281**, 209–38 (2003).

31.   F. D. Bushman *et al.*, Host cell factors in HIV replication: meta-analysis of genome-wide
studies. *PLoS Pathog.* **5**, e1000437 (2009).

32.   W. C. Greene, B. M. Peterlin, Charting HIV's remarkable voyage through the cell: Basic
science as a passport to future therapy. *Nat. Med.* **8**, 673–80 (2002).

33.   Y. Suzuki, R. Craigie, The road to chromatin - nuclear entry of retroviruses. *Nat. Rev.*
Microbiol. **5**, 187–96 (2007).

34.   V. Achuthan *et al.*, Capsid-CPSF6 Interaction Licenses Nuclear HIV-1 Trafficking to Sites of
Viral DNA Integration. *Cell Host Microbe* (2018), doi:10.1016/j.chom.2018.08.002.

35.   C. R. Chin *et al.*, Direct Visualization of HIV-1 Replication Intermediates Shows that Capsid
and CPSF6 Modulate HIV-1 Intra-nuclear Invasion and Integration. *Cell Rep.* **13**, 1717–1731
(2015).

36.   K. Peng *et al.*, Quantitative microscopy of functional HIV post-entry complexes reveals
association of replication with the viral capsid. *Elife.* **3** (2014), doi:10.7554/eLife.04114.

37.   A. Dharan *et al.*, KIF5B and Nup358 Cooperatively Mediate the Nuclear Import of HIV-1 during
Infection. *PLOS Pathog.* **12**, e1005700 (2016).

38.   C. Buffone *et al.*, Nup153 Unlocks the Nuclear Pore Complex for HIV-1 Nuclear Translocation
in Nondividing Cells. *J. Virol.* **92** (2018), doi:10.1128/JVI.00648-18.

39.   H. O. Ajoge *et al.*, Antiretroviral APOBEC3 Cytidine Deaminases Alter HIV-1 Provirus
Integration Site Profiles. *bioRxiv*, 833475 (2019).

40.   F. D. Bushman, Tethering human immunodeficiency virus 1 integrase to a DNA site directs

506   integration to nearby sequences. *Proc.Natl.Acad.Sci.USA*. **91**, 9233–9237 (1994).

507   41.   P. M. Pryciak, H. E. Varmus, Nucleosomes, DNA-binding proteins, and DNA sequence
508   modulate retroviral integration target site selection. *Cell*. **69**, 769–80 (1992).

509   42.   X. Wu, Y. Li, B. Crise, S. M. Burgess, D. J. Munroe, Weak palindromic consensus sequences
510   are a common feature found at the integration target sites of many retroviruses. *J. Virol.* **79**,
511   5211–4 (2005).

512   43.   R. G. McAllister *et al.*, Lentivector integration sites in ependymal cells from a model of
513   metachromatic leukodystrophy: non-B DNA as a new factor influencing integration. *Mol. Ther.*
514   *Nucleic Acids*. **3**, e187 (2014).

515   44.   Jungkweon Choi and Tetsuro Majima, Conformational changes of non-BDNA. *Chem. Soc.*
516   *Rev.* **40**, 5893–5909 (2011).

517   45.   A. Bacolla, R. D. Wells, Non-B DNA conformations, genomic rearrangements, and human
518   disease. *J. Biol. Chem.* **279**, 47411–4 (2004).

519   46.   C. M. Venner *et al.*, Infecting HIV-1 Subtype Predicts Disease Progression in Women of Sub-
520   Saharan Africa. *EBioMedicine*. **13**, 305–314 (2016).

521   47.   C. S. Morrison *et al.*, Hormonal Contraceptive Use and HIV Disease Progression Among
522   Women in Uganda and Zimbabwe. *JAIDS J. Acquir. Immune Defic. Syndr.* **57**, 157–164
523   (2011).

524   48.   C. S. Morrison *et al.*, Hormonal contraception and the risk of HIV acquisition. *AIDS*. **21**, 85–95
525   (2007).

526   49.   T. L. Lemonovich *et al.*, Differences in Clinical Manifestations of Acute and Early HIV-1
527   Infection between HIV-1 Subtypes in African Women. *J. Int. Assoc. Provid. AIDS Care*. **14**,
528   415–422 (2015).

529   50.   A. Ciuffi, S. D. Barr, Identification of HIV integration sites in infected host genomic DNA.
530   *Methods*. **53**, 39–46 (2011).

531   51.   R. Z. Cer *et al.*, Non-B DB v2.0: a database of predicted non-B DNA-forming motifs and its
532   associated tools. *Nucleic Acids Res.* **41**, D94–D100 (2013).

533   52.   L. Guelen *et al.*, Domain organization of human chromosomes revealed by mapping of nuclear
534   lamina interactions. *Nature*. **453**, 948–51 (2008).

535   53.   S. D. Barr *et al.*, HIV integration site selection: targeting in macrophages and the effects of
536   different routes of viral entry. *Mol. Ther.* **14**, 218–225 (2006).

537   54.   D. Derse *et al.*, Human T-cell leukemia virus type 1 integration target sites in the human
538   genome: comparison with those of other retroviruses. *J. Virol.* **81**, 6731–41 (2007).

539   55.   A. Faschinger *et al.*, Mouse Mammary Tumor Virus Integration Site Selection in Human and
540   Mouse Genomes. *J. Virol.* **82**, 1360–1367 (2008).

541   56.   B. Marini *et al.*, Nuclear architecture dictates HIV-1 integration site selection. *Nature*. **521**,
542   227–31 (2015).

543   57.   E. Battivelli *et al.*, Distinct chromatin functional states correlate with HIV latency reactivation in
544   infected primary CD4+ T cells. *Elife*. **7** (2018), doi:10.7554/eLife.34655.

58.   V. Poletti, F. Mavilio, *Interactions between Retroviruses and the Host Cell Genome* (2018;

http://www.ncbi.nlm.nih.gov/pubmed/29159201), vol. 8.

59.   F. Maldarelli *et al.*, Specific HIV integration sites are linked to clonal expansion and
persistence of infected cells. *Science*. **345**, 179–83 (2014).

60.   L. B. Cohn *et al.*, HIV-1 Integration Landscape during Latent and Active Infection. *Cell*. **160**,
420–432 (2015).

61.   C. Cattoglio *et al.*, Hot spots of retroviral integration in human CD34+ hematopoietic cells.
Blood. **110**, 1770–8 (2007).

62.   M. Nettling *et al.*, DiffLogo: a comparative visualization of sequence motifs. *BMC*
Bioinformatics. **16**, 387 (2015).

63.   W. P. Wahls, L. J. Wallace, P. D. Moore, The Z-DNA motif d(TG)30 promotes reception of
information during gene conversion events while stimulating homologous recombination in
human cells in culture. *Mol. Cell. Biol.* **10**, 785–93 (1990).

64.   J. Y. Chin, E. B. Schleifman, P. M. Glazer, Repair and recombination induced by triple helix
DNA. *Front. Biosci.* **12**, 4288 (2007).

65.   V. Brázda, R. C. Laister, E. B. Jagelská, C. Arrowsmith, Cruciform structures are a common
DNA feature important for regulating biological processes. *BMC Mol. Biol.* **12**, 33 (2011).

66.   C. M. Holtz, H. A. Sadler, L. M. Mansky, APOBEC3G cytosine deamination hotspots are
defined by both sequence context and single-stranded DNA secondary structure. *Nucleic*
Acids Res. **41**, 6139–48 (2013).

67.   S. J. Ziegler *et al.*, Insights into DNA substrate selection by APOBEC3G from structural,
biochemical, and functional studies. *PLoS One*. **13**, e0195048 (2018).

68.   A. N. Mazumder *et al.*, Inhibition of human immunodeficiency virus type 1 integrase by
guanosine quartet structures. *Biochemistry*. **35**, 13762–13771 (1996).

69.   J.-C. Jiang, K. R. Upton, Human transposons are an abundant supply of transcription factor
binding sites and promoter activities in breast cancer cell lines. *Mob. DNA*. **10**, 16 (2019).

70.   Y. Kondo, J.-P. J. Issa, Enrichment for histone H3 lysine 9 methylation at Alu repeats in human
cells. *J. Biol. Chem.* **278**, 27658–62 (2003).

71.   A. Siddiqui-Jain, C. L. Grand, D. J. Bearss, L. H. Hurley, Direct evidence for a G-quadruplex in
a promoter region and its targeting with a small molecule to repress c-MYC transcription. *Proc.*
Natl. Acad. Sci. U. S. A. **99**, 11593–8 (2002).

72.   A. Verma, V. K. Yadav, R. Basundra, A. Kumar, S. Chowdhury, Evidence of genome-wide G4
DNA-mediated gene expression in human cancer cells. *Nucleic Acids Res.* **37**, 4194–204
(2009).

73.   S. Waga, S. Mizuno, M. Yoshida, Chromosomal protein HMG1 removes the transcriptional
block caused by the cruciform in supercoiled DNA. *J. Biol. Chem.* **265**, 19424–8 (1990).

74.   S. Waga, S. Mizuno, M. Yoshida, Nonhistone protein HMG1 removes the transcriptional block
caused by left-handed Z-form segment in a supercoiled DNA. *Biochem. Biophys. Res.*
Commun. **153**, 334–9 (1988).

75.   A. Jain, M. Magistri, S. Napoli, G. M. Carbone, C. V. Catapano, Mechanisms of triplex DNA-
mediated inhibition of transcription initiation in cells. *Biochimie*. **92**, 317–320 (2010).

76.   L. J. Maher, P. B. Dervan, B. Wold, Analysis of promoter-specific repression by triple-helical
DNA complexes in a eukaryotic cell-free transcription system. *Biochemistry*. **31**, 70–81 (1992).

77.   M. L. Bochman, K. Paeschke, V. A. Zakian, DNA secondary structures: stability and function of
G-quadruplex structures. *Nat. Rev. Genet.* **13**, 770–80 (2012).

78.   J. Delic, R. Onclercq, M. Moisan-Coppey, Inhibition and enhancement of eukaryotic gene
expression by potential non-B DNA sequences. *Biochem. Biophys. Res. Commun.* **180**, 1273–
83 (1991).

79.   S. Tornaletti, S. Park-Snyder, P. C. Hanawalt, G4-forming sequences in the non-transcribed
DNA strand pose blocks to T7 RNA polymerase and mammalian RNA polymerase II. *J. Biol.*
Chem. **283**, 12756–62 (2008).

80.   B. P. Belotserkovskii *et al.*, A triplex-forming sequence from the human c-MYC promoter
interferes with DNA transcription. *J. Biol. Chem.* **282**, 32433–41 (2007).

81.   R. Löwer, J. Löwer, R. Kurth, The viruses in all of us: characteristics and biological significance
of human endogenous retrovirus sequences. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 5177–84
(1996).

82.   G.-Z. Han, M. Worobey, An endogenous foamy-like viral element in the coelacanth genome.
PLoS Pathog. **8**, e1002790 (2012).

83.   W. M. Switzer *et al.*, Ancient co-speciation of simian foamy viruses and primates. *Nature*. **434**,
376–380 (2005).

84.   S. R. Ross, Mouse mammary tumor virus molecular biology and oncogenesis. *Viruses*. **2**,
2000–12 (2010).

85.   H.-C. Chen, J. P. Martinez, E. Zorita, A. Meyerhans, G. J. Filion, Position effects influence HIV
latency reversal. *Nat. Struct. Mol. Biol.* **24**, 47–54 (2017).

86.   L. S. Vranckx *et al.*, LEDGIN-mediated Inhibition of Integrase–LEDGF/p75 Interaction
Reduces Reactivation of Residual Latent HIV. *EBioMedicine* (2016),
doi:10.1016/j.ebiom.2016.04.039.

87.   J. Demeulemeester *et al.*, HIV-1 integrase variants retarget viral integration and are associated
with disease progression in a chronic infection cohort. *Cell Host Microbe*. **16**, 651–662 (2014).

88.   R. E. Myers, D. Pillay, Analysis of Natural Sequence Variation and Covariation in Human
Immunodeficiency Virus Type 1 Integrase. *J. Virol.* **82**, 9228–9235 (2008).

89.   S. Y. Rhee *et al.*, Natural variation of HIV-1 group M integrase: Implications for a new class of
antiretroviral inhibitors. *Retrovirology*. **5**, 1–11 (2008).

FIGURE LEGENDS

Figure 1: Evolutionarily diverse retroviruses exhibit distinct integration site preferences. A,
Heatmaps depicting the fold enrichment or depletion of integration sites near common genomic
features compared to matched random controls. Darker shades represent higher fold-changes in the
ratio of integration sites to matched random control sites. Bins represent the distance of the
integration sites from each genomic feature. Bin 0 = within the feature; Bin 1= 1-499 bp; Bin 2 = 500-
4,999 bp; Bin 3 = 5,000-49,999 bp; Bin 4 = >49,999 bp. Heatmaps of the diverse retrovirus genera

were superimposed on a BioNJ tree constructed using their reverse transcriptase amino acid

sequences using the Dayhoff substitution model with 1000 bootstraps. All branches are scaled

according to number of amino acids changes per site. The phylogenetic tree shows the evolutionary

relatedness of the different retrovirus genera only. Significant differences are denoted by asterisks (*P

< 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001) (Fisher's Exact test). **B**, Proportion of retroviral

integration sites located within genes compared to the random control (blue line). **C**, Pairwise analysis

was performed on the retroviral integration site profile preferences (fold enrichment and depletion

values) using Euclidean distance as the measurement method (Heatmapper, Babicki et al., 2016).

Stronger relationships between retroviral integration site profiles are indicated by darker blue color in

the pairwise distance matrix. **D**, Nuclear localization of integration sites were determined by

quantifying the proportion of total integrations that fell within a lamin-associated domain (LAD) (=1) as

opposed to outside a LAD (=0). *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; Fisher's Exact

test. Infinite number (inf), 1 or more integrations were observed when 0 integrations were expected by

chance. Not a number (nan), 0 integrations were observed and 0 were expected by chance. HIV-1 =

human immunodeficiency virus, SIV_{mne} = simian immunodeficiency virus (isolated from a pig-tailed

macaque), FIV= feline immunodeficiency virus, HTLV-1= human T-lymphotrophic virus Type 1, MLV =

murine leukemia virus, FV = foamy virus, ASLV = avian sarcoma leucosis virus, MMTV = mouse

mammary tumor virus and ERVs = endogenous retroviruses.

Figure 2: Evolutionarily diverse retroviruses target non-B DNA for integration. A, Heatmaps

illustrating the fold-enrichment or depletion of unique retroviral integration sites near non-B DNA

features compared to matched random controls. Darker shades represent higher fold-changes in the

ratio of integration sites to matched random control sites. Significant differences are denoted by

asterisks *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001) (Fisher's Exact test). **B**, Pairwise

analysis was performed on the retroviral integration site profile preferences using Euclidean distance

as the measurement method (Heatmapper, Babicki et al., 2016). Stronger relationships between

retroviral integration site profiles are indicated by darker blue color in the pairwise distance matrix.

Figure 3: Integration site profiles differ between *in vitro*- and *in vivo*-derived datasets. A,

Heatmaps illustrating the fold-enrichment or depletion of unique integration sites (compared to the

matched random control) near common genomic features from *in vivo*-derived datasets (n= 167,098

sites) and *in* vitro-derived datasets (n= 13,601 sites). Inset numbers represent the fold-change in the

percentage of integration sites. **B,** Comparison of the percentage of integration sites within 5,000 bp

of common genomic features between *in vitro*- and *in vivo*-derived datasets. **C,** Heatmaps illustrating

the fold-enrichment of unique integration sites compared to the matched random control near non-B

DNA from *in vivo*- and *in vitro*-derived datasets. **D,** Comparisons of the percentage of integration sites

within 500 bp of G4, triplex and Z-DNA between *in vitro*- and *in vivo*-derived datasets. Significant

differences are denoted by asterisks *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001) (Fisher's

Exact test).

Figure 4: HIV-1 subtype A, B, C and D have different integration site preferences for genomic

features. A, Comparison of the percentage of integration sites near common genomic features

between HIV-1 subtypes A, B, C and D. **B,** Heatmaps depicting the fold enrichment or depletion of

integration sites near common genomic features compared to the matched random control. Darker

shades represent higher fold-changes in the ratio of integration sites to matched random control sites.

Bins in A and B represent the distance of the integration sites from the genomic feature. **C**,

Comparison of the percentage of integration sites within genes from HIV-1 subtypes A, B, C and D. **D,**

Pairwise analysis was performed on the retroviral integration site profile preferences using Euclidean

distance as the measurement method (Heatmapper, Babicki et al., 2016). Stronger relationships

between retroviral integration site profiles are indicated by darker blue color in the pairwise distance

matrix. Significant differences are denoted by asterisks *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P <$

0.0001 (Fisher's Exact test). **E**, Estimates of Evolutionary Divergence over Sequence Pairs between

Groups. The number of amino acid substitutions per site from averaging over all sequence pairs

between groups are shown. Analyses were conducted using the Poisson correction model. This

analysis involved 486 amino acid sequences. The coding data was translated assuming a Standard

genetic code table. All ambiguous positions were removed for each sequence pair (pairwise deletion

option). There were a total of 265 positions in the final dataset. Evolutionary analyses were conducted

in MEGA X. **F**, Amino acid alignment of the C-terminal domain of HIV-1 integrase from subtypes A, B,

C and D.

Figure 5: HIV-1 subtype A, B, C and D have different integration site preferences for non-B

DNA. A, Comparison of the percentage of integration sites near non-B DNA features between HIV-1

subtypes A, B, C and D. **B,** Heatmaps depicting the fold enrichment or depletion of integration sites

near non-B DNA compared to the matched random control. Darker shades represent higher fold-

changes in the ratio of integration sites to matched random control sites. Bins in A and B represent

the distance of the integration sites from the non-B DNA feature. Comparison of the percentage of

integration sites within genes from HIV-1 subtypes A, B, C and D. **C,** Pairwise analysis was performed

on the retroviral integration site profile preferences using Euclidean distance as the measurement

method (Heatmapper, Babicki et al., 2016). Stronger relationships between retroviral integration site

profiles are indicated by darker blue color in the pairwise distance matrix. Significant differences are

denoted by asterisks *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$ (Fisher's Exact test).

Figure 6: Integration hotspots identified from individuals infected with different HIV-1

subtypes. A, Venn diagram showing the percentage of unique (colored red or light blue) and shared

(colored dark blue) integration site hotspots among the different HIV-1 subtypes. **B** and **C,** circos plots

showing integration site hotspots inside (B) or outside (C) genes. The subtypes whose hotspots were

shared is denoted in capital letters on the outside of the circos plots. For integration site hotspots

outside genes (C), the closest gene and the distance from that gene are shown on the outside of the

circos plots separated by a colon. Hotspots were defined as a 1 kb window in the genome hosting 2

or more integration sites. Note: (+) Due to the large number of subtype B integration site hotspots

(10,762), only those hotspots shared between subtypes A, C and/or D are shown on the circos plots.

No integration site hotspots outside of genes were shared between subtype B and any of the other

subtypes. **D**, Consensus sequences were generated from a window of 100 nucleotides upstream and

100 nucleotides downstream of each integration site. Consensus sequences from integration sites

located in hotspots were compared to consensus sequences from sites not located in hotspots using

DiffLogo. Consensus sequences were analyzed for the presence of non-B DNA motifs (solid colored

lines).