1

Staphylococcal DNA repair confers tolerance of the bactericidal activity of both neutrophils and antibiotics

Kam Pou Ha[1]*, Rebecca S. Clarke[1]*, Jane L. Brittan[2], Jessica E. Rowley[1], Despoina A. I. Mavridou[1,3], Thomas B. Clarke[1], Angela H. Nobbs[2] and Andrew M. Edwards[1#]

[1] MRC Centre for Molecular Bacteriology and Infection, Imperial College London, Armstrong Rd, London, SW7 2AZ, UK.

[2] Bristol Dental School, University of Bristol, Bristol, BS1 2LY, UK.

[3] Department of Molecular Biosciences, University of Texas at Austin, Austin, 78712, Texas, USA

* these authors contributed equally.

[#] For correspondence:

a.edwards@imperial.ac.uk

Keywords: *Staphylococcus* / *Streptococcus* / *Enterococcus* / neutrophil / oxidative burst / antibiotic.

Running title: Neutrophils and antibiotics target *S. aureus* DNA

Abstract

Staphylococcus aureus is a leading cause of chronic and recurrent infections of the skin, bones, joints and bloodstream. During infection, *S. aureus* faces the twin threats of the host immune system and therapeutic antibiotics. Since both host immune cells and antibiotics generate reactive oxygen species, we hypothesised that *S. aureus* may employ a common mechanism to repair damage caused by either threat. We found that staphylococcal DNA is damaged during exposure to both human neutrophils and most bactericidal antibiotics. To understand the nature of this damage and how *S. aureus* repairs it, we screened a panel of transposon mutants defective for various DNA repair processes. This revealed that loss of the *rexBA* operon significantly reduced staphylococcal survival in human blood, during incubation with purified neutrophils, in the peritoneal cavity of mice and during exposure to a large panel of bactericidal antibiotics. We then used biochemical assays to demonstrate that RexAB is a member of the AddAB family of ATP-dependent helicase/nucleases that are required for the repair of DNA double strand breaks. Finally, we found that RexAB homologues in *Enterococcus faecalis* and *Streptococcus gordonii* also promoted survival of these pathogens in human blood, suggesting that DNA repair constitutes a broadly conserved defence against neutrophils. Together, these data demonstrate that DNA is a target of host immune cells and several antibiotics, leading to double-strand breaks, and that repair of this damage by an AddAB-family enzyme enables the survival of Gram-positive pathogens during infection.

Introduction

Staphylococcus aureus is a major human pathogen, responsible for a raft of serious invasive infections including bacteraemia, infective endocarditis and osteomyelitis[1]. Whilst the combination of a potent immune response and the use of antibiotics significantly reduces mortality, many infections become chronic or recurrent[2]. This implies that *S. aureus* has acquired mechanisms by which it can tolerate the damaging effects of both immune cells and antibiotics, and that disruption of such tolerance mechanisms may therefore provide new therapeutic approaches to combat chronic and recurrent infections.

Entry of *S. aureus* into normally sterile tissues triggers a strong immune response, with significant infiltration of neutrophils to control infection[3,4,5]. Neutrophils phagocytose *S. aureus*, exposing them to a bactericidal cocktail of antimicrobial peptides and proteases[3,6], as well as the oxidative burst that consists of reactive oxygen species (ROS) that damages lipids, proteins and DNA [7,8,9,10,11,12]. Whilst the contribution of each ROS to bactericidal activity is the subject of investigation, there is compelling evidence that the oxidative burst is crucial for control of *S. aureus* infection[9,12,13]. For example, individuals with chronic granulomatous disease (CGD) are particularly prone to staphylococcal infections because their neutrophils are defective for the oxidative burst[4,14]. In keeping with this, *S. aureus* survives better in mice defective for the oxidative burst, relative to wild-type animals, whilst the treatment of human neutrophils with an inhibitor of the oxidative burst increased staphylococcal survival relative to untreated immune cells[5,9,15]. However, even when the oxidative burst is functional, there is evidence that some *S. aureus* cells can survive in neutrophils, which contributes to the progression and persistence of infection[16,17].

Despite the importance of the oxidative burst in combatting staphylococcal infection, relatively little is known about how it kills the pathogen. Studies with single oxidants such as H_2O_2 indicate that the molecular targets of the oxidative burst are broad and include proteins, lipids and DNA[18]. To survive this damage, *S. aureus* employs several stress response regulators and repair systems. For example, previous work has shown that DNA damage caused by *S. aureus* exposure to H_2O_2 leads to the initiation of the DNA repair SOS response[19], which facilitates the excision of damaged bases or repair of double-strand breaks (DSBs) by homologous recombination[20,21]. DSBs are particularly dangerous since they are lethal if not repaired, as evidenced by the rapid bactericidal activity of the antibiotic ciprofloxacin which causes this type of DNA damage[21]. However, H_2O_2 is a sub-optimal model for the oxidative burst as it's typically used at concentrations that exceed those generated by the oxidative burst[22]. Therefore, it is unclear whether DNA is damaged by neutrophil-generated ROS, what the nature of this damage is and how *S. aureus* repairs it.

In addition to the threat posed by neutrophil generated ROS during infection, *S. aureus* is also targeted

by therapeutic antibiotics such as the β-lactam oxacillin[23]. In addition to the established molecular

targets of antibiotics, there is an increasingly large body of evidence that bactericidal antibiotics

trigger endogenous ROS production in bacteria, leading to DNA damage [24–31]. However, the degree to

which these ROS contribute to bacterial killing is the subject of debate[32-34].

Since ROS generated by neutrophils and antibiotics have been linked to bactericidal activity against *S.*

aureus, we hypothesised that oxidants from both sources target the same bacterial macromolecules

and that the mechanisms used by staphylococci to survive exposure to host immune defences may

therefore also enable staphylococci to tolerate damage caused by antibiotics. This would explain how

S. aureus is so successful in surviving the twin threats of neutrophils and antibiotics and may identify

targets for novel therapeutics that sensitise *S. aureus* to both host defences and antibiotics, enabling

the efficient clearance of the pathogen from infection sites.

To test this hypothesis, we determined whether staphylococcal DNA is a target of both antibiotic-

induced ROS and the oxidative burst of neutrophils and elucidated the mechanism by which this

pathogen repairs damage caused by both threats. This revealed that a single DNA repair pathway

conferred tolerance of damage caused by both neutrophils and multiple bactericidal antibiotics.

Results

Neutrophils and most antibiotics cause DNA damage in *S. aureus*

DNA damage in most bacteria, including *S. aureus,* triggers activation of the SOS response, which leads

to the transcription of genes whose products contribute to DNA repair. These include *recA*, a crucial

mediator of DNA recombination and a regulator of the SOS response[21]. To determine whether

neutrophils and antibiotics cause DNA damage in *S. aureus*, we constructed a P*recA-gfp* reporter

system that placed the expression of *gfp* under the control of the *recA* promoter in strains from two

distinct genetic backgrounds: SH1000, a methicillin-sensitive *S. aureus* (MSSA) strain, and JE2, a

community-associated methicillin-resistant *S. aureus* (CA-MRSA) strain of the USA300 lineage[35,36]. We

then demonstrated functionality of this system by showing a dose-dependent fluorescent signal in

response to mitomycin C, a well-characterised DNA-damaging antibiotic that triggers the SOS

response (Fig. S1A,B).

S. aureus strains bearing the P*recA-gfp* reporter were incubated with purified human neutrophils to

measure both phagocytosis and induction of *recA* expression (Fig. S2). To study the role of the

oxidative burst in this process, some of the neutrophils were pre-treated with the NADPH inhibitor

diphenyleneiodonium (DPI), whilst others were incubated with the DMSO solvent alone. Incubation

of wild-type SH1000 and JE2 strains containing the *recA-gfp* reporter with neutrophils resulted in rapid

phagocytosis, with >90% bacteria internalised within 5 min and >99% by 30 min (Fig. 1A). During the

first 30 min, there was an increase in GFP-mediated fluorescence for both SH1000 and JE2 strains

incubated with neutrophils treated with DMSO (Fig. 1B). There was also a GFP signal from bacteria

incubated with DPI-treated neutrophils, but to a lesser degree than that seen for DMSO only (Fig. 1B).

Therefore, phagocytosis of *S. aureus* by neutrophils leads to induction of the SOS response, partly but

not exclusively due to the oxidative burst.

Previous work from our group has shown that co-trimoxazole triggers the SOS response in *S. aureus*

in a time and dose-dependent manner[37]. To determine whether other antibiotic classes damaged DNA

and thus induced the SOS response, we incubated the *S. aureus* P*recA-gfp* reporter strains with various

antibiotics (Fig. S3). We found that ciprofloxacin and nitrofurantoin strongly triggered SOS induction

in both the SH1000 and JE2 strains (Fig. 1C,D). DNA damage was also apparent during exposure to

oxacillin, daptomycin, chloramphenicol or linezolid, albeit to varying degrees, but not when exposed

to gentamicin (Fig. 1E-I).

Taken together, these data demonstrated that both neutrophils and almost all clinically relevant

antibiotics, including bacteriostatic agents and those that do not directly target bacterial DNA

replication triggered the SOS response in *S. aureus*, which is indicative of DNA damage.

Figure 1. Induction of *S. aureus* SOS response by neutrophils and antibiotics. (A) Phagocytosis of
SH1000 and JE2 WT *S. aureus recA-gfp* reporter strains by neutrophils and **(B)** resulting induction of
the SOS response measured by GFP expression (RFU) in the presence of the oxidative burst inhibitor
DPI (grey) or an identical volume of the DMSO solvent alone (black) (n = 4, with blood from different
donors). **(C-I)** Induction of SOS measured by GFP expression upon exposure to a range of
concentrations of various antibiotics. Antibiotic concentrations are labelled by multiple of the MIC of
the individual strain (x MIC). GFP fluorescence was normalised by OD_{600} to determine induction of SOS
relative to cell density (n = 3). OD_{600} measurements alone are shown in Fig. S3. Data in **B** were analysed
by two-way ANOVA with Sidak's post-hoc test, for DPI-treated versus DMSO-treated (*, $P < 0.05$).
Where shown, error bars represent standard deviation of the mean. Error bars were omitted in **C-I** for
clarity.

The RexAB DNA repair complex is required for the survival of *S. aureus* during infection

To understand the nature of the DNA damage caused by neutrophils and how it was repaired by *S. aureus*, we assembled a panel of mutants from the ordered NARSA transposon library[35], each of which was deficient for a different protein involved in DNA repair. To understand which of these proteins were important for repairing damage caused by neutrophils, we measured the survival of wild-type and mutant *S. aureus* in an *ex vivo* whole human blood model of infection. Previous characterisation of this model system by our group and others has shown that *S. aureus* is rapidly phagocytosed by neutrophils and exposed to the oxidative burst and other killing mechanisms[38,39,40].

Our targeted screen revealed that survival of mutants defective for *rexA* or *rexB* was <5% of the inoculum, compared with >60% for the wild type (Fig. S4). The *rexA* and *rexB* genes form a two gene operon (*rexBA*) on the staphylococcal chromosome, and are proposed to encode an AddAB helicase/nuclease enzyme involved in the processing of DNA DSBs[41–44]. We confirmed a role for RexAB in staphylococcal survival in blood by complementation of mutants with a plasmid containing the *rexBA* operon (p*rexBA*), which restored bacterial survival to wild type levels (Fig. 2A,B). By contrast, the survival of *rexBA* mutants transformed with the vector alone (pEmpty) was not changed (Fig. 2A,B). Having shown that RexAB contributed to staphylococcal tolerance of neutrophil mediated killing in human blood, we then tested its role in staphylococcal survival during exposure to neutrophils *in vivo*. Mice were infected via the peritoneal cavity, which results in the recruitment of neutrophils within 2 h, with wild-type or *rexBA* mutants of *S. aureus* SH1000 or JE2[45]. After 6 h, the mice were sacrificed and the peritoneal cavity washed with PBS to recover bacteria, which were quantified by CFU counts. This revealed that *rexBA* mutants in both genetic backgrounds were significantly attenuated for survival *in vivo* relative to wild type bacteria, confirming that RexAB contributes to staphylococcal resistance to host immune defences and the progression of infection (Fig. 2C,D).

Figure 2. Reduced survival of *rexB* mutants in human blood and in an *in vivo* murine infection model. **(A-B)** Survival of *S. aureus* WT, *rexB* mutant, empty vector (pEmpty) and complemented mutant (p*rexBA*) in SH1000 (A) and JE2 (B) backgrounds after 6 h incubation in whole human blood. Empty vectors and complemented mutants were supplemented with 100 ng μl^{-1} AHT to control *rexBA* expression (n ≥ 3). **(C-D)** Number of CFU recovered from the peritoneal cavities of mice 6 h after inoculation (each circle represents a single mouse; n = 4-5). Data in **A** and **B** were analysed by one-way ANOVA with Tukey's post-hoc test, for *rexB,* pEmpty or p*rexBA* versus WT (*, P < 0.05). Data in **C** and **D** were analysed by Mann-Whitney test (*, P < 0.05). Where shown, error bars represent standard deviation of the mean.

RexAB contributes to staphylococcal tolerance of the oxidative burst

Having confirmed that *rexBA* is important for staphylococcal survival during infection conditions, we wanted to determine whether this was due to repair of DNA damage caused by the oxidative burst of neutrophils. As described above, the survival of *rexBA* mutants in whole human blood was significantly reduced relative to wild-type bacteria (Fig. 2A,B). However, the presence of DPI, which inhibits the NADPH oxidase complex, promoted the survival of *rexBA* mutants to wild-type levels, indicating that the survival deficit observed for bacteria lacking RexAB was due to increased sensitivity to the oxidative burst (Fig. 3A,B). Comparable results were obtained when *S. aureus* was incubated with

neutrophils purified from human blood. The *rexBA* mutants were more susceptible to neutrophil
mediated killing than wild type, but the presence of DPI restored survival of the *rexBA* mutants to wild
type levels (Fig. 3C,D).

These data strongly indicated that RexAB contributed to staphylococcal survival of the oxidative burst
by repairing DNA. However, several different factors have been shown to affect survival of *S. aureus*
incubated in human blood or exposed to oxidative stress such as production of the staphyloxanthin
pigment[38]. However, whilst there was a slight growth defect for the *rexBA* mutant relative to the wild
type, we did not see significant differences between wild type strains, *rexBA* mutants and
complemented strains for toxin production, staphyloxanthin pigment production or catalase activity
(Fig. S5). Together, the data shown in Figures 1-3 demonstrate that neutrophils cause DNA damage in
S. aureus, in part via the oxidative burst, which needs to be repaired by RexAB to enable staphylococcal
survival in the host.

Figure 3. RexAB protects *S. aureus* from the oxidative burst of phagocytes. (A-B) Survival of *S. aureus* WT, *rexB* mutant, empty vector (pEmpty) and complemented mutant (p*rexBA*) in SH1000 (A) and JE2 (B) backgrounds in whole human blood after 6 h of incubation, in the presence of the oxidative burst inhibitor DPI (grey) or an identical volume of the DMSO solvent alone (black). Empty vectors and complemented mutants were supplemented with 100 ng μl^{-1} AHT to induce *rexBA* expression (n = 3). **(C-D)** Survival of *S. aureus* WT and *rexB* mutant when incubated with purified human neutrophils for 3 h (n = 4). Data in **A** and **B** were analysed by one-way ANOVA with Tukey's post-hoc test, for *rexB,* pEmpty or p*rexBA* versus WT (*, $P < 0.05$). Data in **C** and **D** were analysed by Student's t-test, for *rexB* mutant versus WT (*, $P < 0.05$). Error bars represent standard deviation of the mean.

RexAB contributes to the survival of *S. aureus* exposed to bactericidal antibiotics

We have shown previously that RexAB enables staphylococcal tolerance of the combination antibiotic co-trimoxazole[37]. Since most of the DNA damage caused by co-trimoxazole was due to the production of ROS, we hypothesised that RexAB would promote the growth and survival of *S. aureus* when incubated with other bactericidal antibiotics reported to trigger ROS production in bacteria[24–30].

To test this, we exposed wild-type *S. aureus* SH1000 and JE2, along with associated *rexB* mutants, to a panel of bactericidal (ciprofloxacin, nitrofurantoin, oxacillin, daptomycin, gentamicin, vancomycin, fosfomycin) and bacteriostatic (chloramphenicol, linezolid) antibiotics and determined the minimum

inhibitory concentration (MIC) of each (Fig. 4A). The *rexB* mutants in both JE2 and SH1000 strains were

≥2-fold more susceptible to 7 of the 9 antibiotics tested under aerobic conditions, the exceptions being

vancomycin, and gentamicin for the JE2 strain only (Fig. 4A). When assays were done under anaerobic

conditions the lack of oxygen reduced the susceptibility of one or both *S. aureus* wild-type strains to

ciprofloxacin, daptomycin, oxacillin and gentamicin, inferring that antibiotic-triggered ROS

contributed to growth inhibitory activity. Furthermore, under anaerobic conditions, the only

antibiotics that were more active against both *rexB* mutants relative to their parental strains were

ciprofloxacin, linezolid and genatmicin, suggesting that ROS are the primary, but not the only,

mechanism by which antibiotics cause DNA damage.

Next, we exposed wild-type *S. aureus* and *rexB* mutants to the same antibiotics at 10x the MIC and

measured survival after 8 h (Fig. 4B-J). Similar to the MIC assays, most antibiotics (6/9) were more

active against the *rexB* mutant relative to wild type bacteria under aerobic conditions. This resulted in

5-500-fold greater reduction in CFU counts of the *rexB* mutant compared to wild type cells after 8h

exposure to ciprofloxacin, nitrofurantoin, oxacillin, daptomycin, gentamicin and fosfomycin (Fig. 4B-

I).

Next, the role of ROS was investigated by repeating assays under anaerobic conditions. This revealed

that the presence of oxygen contributed to the bactericidal activity of gentamicin but none of the

other antibiotics examined against wild type bacteria (Fig. 4B-I). However, whilst most antibiotics

killed efficiently under anaerobic conditions, in almost all cases they were equally active against both

wild type and *rexB* mutants (Fig. 4B-I). The exception to this was ciprofloxacin which directly damages

DNA leading to DNA DSBs (Fig. 4B).

Taken together, these findings support previous work by showing that several diverse classes of

antibiotics cause DNA damage under aerobic conditions, most likely via ROS generation[24-30,37].

However, this damage does not contribute to killing of wild type bacteria because damage is repaired

via RexAB and the SOS response.

A

Minimum inhibitory concentration (μg/ml)							
	Aerobic				Anaerobic		
	SH1000 WT	SH1000 *rexB*	JE2 WT	JE2 *rexB*	SH1000 WT	SH1000 *rexB*	JE2 WT	JE2 *rexB*
Ciprofloxacin	0.125	0.0078	16	4	0.25	0.03	32	2
Nitrofurantoin	8	4	16	4	8	4	8	8
Oxacillin	0.125	0.06	0.125	0.06	0.125	0.125	0.25	0.125
Daptomycin	0.25	0.125	0.25	0.125	1	1	1	0.5
Chloramphenicol	4	1	4	2	4	4	4	4
Linezolid	1	0.25	1	0.25	1	0.25	1	0.5
Gentamicin	0.0625	0.031	0.125	0.125	16	1	4	2
Vancomycin	1	1	1	1	1	1	1	1
Fosfomycin	16	8	32	16	8	4	16	16

Figure 4. Lack of effective DNA repair increases the sensitivity of *S. aureus* to bactericidal antibiotics in MIC and killing assays. (A) MIC table of *S. aureus* WT and *rexB* mutant in SH1000 and JE2 backgrounds grown under aerobic and anaerobic conditions (n ≥ 3; median MIC is shown). **(B-J)** Survival of *S. aureus* WT and *rexB* mutant in SH1000 and JE2 backgrounds after 8 h incubation at 37°C in TSB supplemented with 10x MIC. Survival was assessed under aerobic (black) or anaerobic (grey) conditions (n = 3). Data in **B-J** were analysed by Student's t-test (*, P < 0.05). Where shown, error bars represent standard deviation of the mean.

RexAB is required for induction of the SOS response during exposure to the oxidative burst and bactericidal antibiotics.

The SOS response facilitates homologous recombination and mutagenic DNA repair, promoting the acquisition of adaptive mutations and antibiotic resistance[20,21,37]. To determine the degree to which RexAB-mediated repair of DNA damage caused by neutrophils and antibiotics triggered SOS, we used the P*recA-gfp* reporter system in JE2 wild-type and *rexB* mutant strains (Fig. S6). Both JE2 wild-type and *rexB* mutant were rapidly phagocytosed by the neutrophils, with similar levels of internalised bacteria (Fig. 5A). We detected GFP expression in wild type bacteria, but this was significantly weaker for the *rexB* mutant, indicating that RexAB-mediated repair is required for full induction of the SOS response in phagocytosed *S. aureus* (Fig. 5B).

Next, induction of *recA* in the JE2 wild-type and *rexB* mutant was monitored during exposure to a panel of antibiotics. We found that the lack of RexAB reduced *recA* induction relative to the wild-type for the DNA-targeting antibiotic ciprofloxacin (Fig. 5C). Similar trends were observed with nitrofurantoin, oxacillin, daptomycin, chloramphenicol and linezolid, albeit to varying degrees (Fig. 5D-H). As expected from previous data (Fig. 1I), no *recA* induction was observed from either wild-type or mutant when exposed to gentamicin (Fig. 5I).

Taken together, these data show that RexAB is required for maximal induction of SOS in response to DNA damage caused by neutrophils or antibiotics, which facilitates repair of DNA damage.

Figure 5. RexAB is required for induction of the SOS response during exposure to the oxidative burst of neutrophils and bactericidal antibiotics. (A) Phagocytosis of JE2 wild-type and *rexB* mutant *recA-gfp* reporter strains by neutrophils and **(B)** resulting induction of the SOS response measured by GFP expression (RFU) after 30 min of exposure (n = 4). **(C-I)** Induction of SOS response of JE2 wild-type and *rexB* mutant measured by GFP expression upon exposure to varying sub-lethal concentrations of bactericidal antibiotic. Concentrations of antibiotic are labelled by multiple of the MIC of the wild-type strain (x MIC). GFP fluorescence was normalised by OD_{600} to determine induction of SOS relative to cell density (n = 3). OD_{600} measurements alone are shown in Fig. S6. Data in **A** and **B** were analysed by Mann-Whitney test (*, P < 0.05). Where shown, error bars represent standard deviation of the mean. Error bars were omitted in **C-I** for clarity.

Staphylococcal RexAB is a member of the AddAB family of ATP-dependent helicase/nucleases

Previous studies have identified staphylococcal RexAB as a member of the AddAB enzyme family via sequence homology[41–44]. In support of this, *in silico* structural analysis of the predicted *rexBA* gene products confirmed that this operon encoded an AddAB-type ATP-dependent helicase/nuclease involved in DSB repair (Fig. S7). However, there is no experimental evidence that RexAB is a functional AddAB homologue.

AddAB enzymes process DSBs to generate a 3' single-stranded DNA overhang that is necessary for RecA-mediated homologous recombination[46]. In support of the structural predictions, phenotypic testing of *rexBA* mutants showed that they were more sensitive than wild-type strains to the DNA-damaging antibiotics ciprofloxacin and mitomycin C (Fig. 6), both of which cause DSBs[47,48].

The generation of the 3' overhang by AddAB proteins results in the formation of a RecA filament, which triggers the SOS response. Therefore, we determined whether RexAB was needed for SOS induction in response to DNA double strand breaks caused by ciprofloxacin or mitomycin C using the *recA-gfp* reporter system described previously. When compared to wild-type bacteria, the JE2 *rexBA* mutant had greatly reduced *recA* expression in response to either antibiotic, providing additional evidence that RexAB is an AddAB-family enzyme (Fig. 5C and S1).

To confirm the ATP-dependent helicase/nuclease activity of the *S. aureus* RexAB complex, recombinant RexAB protein was generated and the helicase and nuclease activities were measured over 1 h. Nuclease activity assays were performed under conditions of high free Mg^{2+}, which has been previously shown to activate nuclease activity in AddAB enzymes[49]. We found that DNA was degraded over time by the recombinant complex in the presence of ATP, whereas this degradation was minimal in its absence, demonstrating that RexAB has ATP-dependent nuclease activity (Fig. 7A).

At conditions of low free Mg^{2+} to measure helicase activity, we observed a loss of dsDNA concomitant with an increase in ssDNA formation over time (Fig. 7B,C). These experiments were repeated in the absence of single-stranded DNA binding protein (SSB), which is required to prevent reannealing of DNA. In the absence of SSB, no DNA unwinding or ssDNA formation were observed, demonstrating that RexAB has ATP-dependent helicase activity (Fig. 7D,E).

Combined, these results show that RexAB is required for the growth of *S. aureus* in the presence of DNA-damaging antibiotics, induction of the SOS response and exhibits ATP-dependent helicase/nuclease activities, demonstrating it is a functional member of the AddAB family of DNA repair complexes.

367 **Figure 6. RexAB is required for growth of *S. aureus* in the presence of DNA-damaging antibiotics.**
368 MIC of *S. aureus* WT, *rexB* mutant, empty vector (pEmpty) and complemented mutant (p*rexBA*) in
369 SH1000 and JE2 backgrounds, for ciprofloxacin (A-B) and mitomycin C (C-D). Empty vectors and
370 complemented mutants were supplemented with 100 ng µl^{-1} AHT to induce *rexBA* expression (n = 3;
371 median MIC is shown). Error bars represent standard deviation of the mean.

Figure 7. RexAB possesses nuclease and helicase activity. (A) Nuclease activity of recombinant *S. aureus* RexAB protein in the presence or absence of ATP (grey and black, respectively) (n = 4). **(B-E)** Helicase activity of recombinant *S. aureus* RexAB protein in the absence (-) or presence (+) of ATP, by measuring unwinding of dsDNA and formation of ssDNA. Single-stranded DNA binding protein (SSB) was added to prevent reannealing of DNA (B-C) (n = 4). Equivalent assays were performed without SSB to confirm helicase activity (D-E) (n = 4). Data in **A-C** were analysed by two-way ANOVA with Sidak's post-hoc test, for without ATP versus with ATP (*, P < 0.05). Error bars represent standard deviation of the mean.

RexAB is required for survival of other Gram-positive pathogens in human blood and during exposure to certain antibiotics

Since homologues of RexAB are present in most Gram-positive bacteria[43,44,46], we next tested whether this repair complex contributes to the survival of other bacteria exposed to neutrophils. Like *S. aureus*, *Enterococcus faecalis* and *Streptococcus gordonii* are frequent causes of infective endocarditis, which brings these species into close contact with neutrophils[50,51]. Therefore, *rexBA* was deleted in representative strains of each species and their sensitivity to the microbicidal activity of neutrophils determined using the *ex vivo* whole human blood model.

Wild-type *E. faecalis* survived at high levels in human blood with ~100% of the inoculum remaining viable during the full 6 h duration of the assay, but the loss of *rexBA* reduced enterococcal survival by ~50%. This indicated that *E. faecalis* suffers DNA damage whilst in blood, but it can be tolerated via DNA repair (Fig. 8A). By contrast to *S. aureus*, DNA damage in *E. faecalis* did not appear to be due to the oxidative burst, since survival of both wild type and *rexB* mutant in blood was unaffected by the presence of DPI (Fig. 8A).

S. gordonii was much more susceptible to host defences in blood than *E. faecalis*, with <5% wild-type bacteria remaining viable after 2 h (Fig. 8A). However, survival of the *rexBA* mutant was still reduced relative to the wild type, with <1% streptococci surviving after 2 h, indicating that DNA repair also contributes to the survival of *S. gordonii* in blood (Fig. 8A). As for *E. faecalis*, inhibition of the oxidative burst with DPI did not promote survival of either the wild type or *rexBA* mutant indicating that DNA damage was not due to the oxidative burst. In support of this, the *rexBA* mutants of both *S. gordonii* and *E. faecalis* were no more susceptible to H_2O_2 than wild-type bacteria (Fig. 8B).

Next, to determine whether loss of RexAB also increased the susceptibility of *E. faecalis* and *S. gordonii* to bactericidal antibiotics, these strains were exposed to a selection of DNA-damaging (ciprofloxacin, co-trimoxazole) and clinically-relevant (daptomycin, linezolid) antibiotics and survival measured after 8 h under aerobic and anaerobic conditions. Consistent with *S. aureus*, *E. faecalis* and *S. gordonii* mutants defective for *rexBA* showed notable reductions in survival during exposure to ciprofloxacin and co-trimoxazole compared with wild type (Fig. 8C,D). However, in keeping with their resistance to ROS, mutants lacking *rexBA* were no more susceptible to daptomycin than wild type (Fig. 8E). Surprisingly, the *E. faecalis rexBA* mutant was killed by the normally bacteriostatic linezolid (Fig. 8F). Similar results were observed under anaerobic conditions (Fig. 8C-F), mimicking the findings in human blood that DNA damage was not caused by the oxidative burst. Therefore, as for *S. aureus*, neutrophils and certain antibiotics cause DNA damage in both *S. gordonii* and *E. faecalis*, the repair of which by RexAB promotes bacterial survival. However, in contrast to *S. aureus*, oxidative stress does not appear to contribute to neutrophil mediated DNA damage in these bacteria.

Figure 8. RexAB protects *Enterococcus faecalis* and *Streptococcus gordonii* from host defences in blood and from antibiotics. (A) Survival of *E. faecalis* OG1X and *S. gordonii* DL1 wild-type and *ΔrexBA* mutant in whole human blood over 3 h in the presence of the oxidative burst inhibitor DPI (dashed lines) or an identical volume of the DMSO solvent alone (solid lines) (n = 3). **(B)** Survival of OG1X and DL1 wild-type and *ΔrexBA* mutant exposed to H_2O_2 for 1 h (n = 3). **(C-F)** Survival of OG1X and DL1 wild-type and *ΔrexBA* mutant in THYB supplemented with antibiotic at 10x MIC of the wild-type. Survival was assessed under aerobic (black) or anaerobic (grey) conditions (n ≥ 3). Data in **A** were analysed by two-way ANOVA with Sidak's post-hoc test, for *ΔrexBA* mutant versus WT (*, P < 0.05). Data in **C-F** were analysed by Student's t-test (*, P < 0.05). Error bars represent standard deviation of the mean.

Discussion

Neutrophils are an essential host defence against *S. aureus*. However, our understanding of the mechanisms by which these immune cells kill staphylococci is limited. Equally, the bactericidal activity of many important anti-staphylococcal antibiotics is only partially understood. The data described above demonstrate that both neutrophils and several different antibiotics cause DNA damage in *S. aureus*, leading to induction of the SOS response. Repair of this DNA damage is dependent upon the RexAB complex, which is a member of the AddAB helicase/nuclease family of enzymes involved in the repair of DSBs. This provides new information on both the target of the oxidative burst and individual antibiotics, and the mechanisms by which *S. aureus* repairs and survives this damage.

The staphylococcal SOS response was triggered within 30 min of phagocytosis by neutrophils, which corresponds with the time at which ROS are maximally generated in these immune cells[52,53] and establishes bacterial DNA as a target of these cells. This is in keeping with previous reports that ROS damage the DNA of *Escherichia coli* and *Coxiella burnetii* phagocytosed by macrophages[54,55]. The SOS response in *S. aureus* was also triggered by several different classes of antibiotics, which indicated DNA damage. Whilst this was expected for DNA-targeting antibiotics such as the fluoroquinolone ciprofloxacin[21], SOS induction also occurred with antibiotics that do not directly target bacterial DNA. It has been shown that certain β-lactam antibiotics induce the SOS response in *E. coli* via the DpiBA two-component system rather than via DNA damage[56]. Although this mechanism has been hypothesised for *S. aureus*[57,58], our data show that *S. aureus rexBA* mutants were more susceptible to killing by oxacillin, demonstrating that DNA damage does occur during exposure to this beta-lactam antibiotic.

Whilst controversial, there is increasing evidence that many classes of antibiotics trigger the production of ROS [24–33]. However, the degree to which these ROS contribute to bactericidal activity is less clear[32,33,34]. Our data provide evidence that many antibiotics cause DNA damage, presumably via ROS. However, this DNA damage appears to be tolerated by wild type bacteria via RexAB-mediated repair.

The production of ROS by bactericidal but not bacteriostatic antibiotics has been proposed to explain their differences in lethality. However, we observed SOS induction during exposure of *S. aureus* to the bacteriostatic antibiotics linezolid and chloramphenicol but not the bactericidal antibiotic gentamicin. Interestingly, despite DNA damage, the absence of RexAB did not sensitise *S. aureus* mutants to linezolid or chloramphenicol but did increase susceptibility to gentamicin. This may be explained by differences in the type of DNA damage caused by each of the antibiotics. Whilst several different types of DNA damage trigger SOS, only those leading to DSBs would be expected to promote susceptibility of the *rexBA* mutant. As such, it is possible that bactericidal antibiotics trigger the potentially lethal

DNA DSBs, whilst bacteriostatic antibiotics trigger non-lethal types of DNA damage. In keeping with this, the absence of RexAB had only a small effect on SOS induction in *S. aureus* caused by linezolid or chloramphenicol.

It is unclear why gentamicin did not trigger SOS during antibiotic exposure, but it may simply be that this antibiotic doesn't trigger endogenous ROS production in *S. aureus*. Combined, our data indicates significant differences between antibiotics in the degree of DNA damage caused, as well as the time required to cause damage. These differences may explain some of the debate around the contribution of ROS to antibiotic-mediated killing.

In addition to repair of DNA damage caused by antibiotics, we also found that RexAB contributed to staphylococcal survival during exposure to purified human neutrophils, incubation in whole human blood and in a murine model of infection. Moreover, we also demonstrated that RexAB contributed to the survival of *E. faecalis* and *S. gordonii* in blood. Previous *in vivo* studies have shown that AddAB is required for the infection of chickens and mice by *Campylobacter jejuni* and *Helicobacter pylori* respectively[59,60]. In place of AddAB, many other Gram-negative bacteria employ the closely related RecBCD helicase/nuclease to process DSBs, which also appears to be important for survival in the host. For example, a *Salmonella enterica* mutant lacking RecBCD was found to be less virulent than the wild type in a murine model of bacteraemia and was also defective for replication in macrophages[61]. Together, these findings indicate that bacterial DNA is a conserved target of host defences, leading to DSBs that must be repaired for pathogen survival.

However, whilst host defences and antibiotics both caused DNA damage, the mechanism seems to vary depending on the species concerned. We found that strains of *E. faecalis* and *S. gordonii* defective for RexAB were more susceptible to host defences in blood, but this was not due to oxidative stress. Neutrophils employ several different antibacterial elements to kill invading pathogens, including reactive oxygen and nitrogen species, proteases and antimicrobial peptides. Several studies have indicated that bacteria vary in their susceptibility to each of these microbicides. In agreement with our findings, Standish *et al.* showed that *S. aureus* but not *Streptococcus pneumoniae* was killed by the oxidative burst[62]. However, whilst *S. aureus* can grow in the presence of nitric oxide, the replication of several other pathogens including *Pseudomonas aeruginosa* and *Streptococcus pyogenes* is inhibited[63]. Since reactive nitrogen species can cause DNA damage, this may provide an explanation for the finding that mutants of *E. faecalis* and *S. gordonii* lacking RexAB are more susceptible to killing by neutrophils in which the oxidative burst is blocked.

The identification of RexAB as important for staphylococcal survival during infection and exposure to antibiotics provides insight into the nature of the damage caused by these twin threats and makes this

complex a potential target for novel therapeutics, particularly as the lack of RexAB homologues in eukaryotes reduces the likelihood of host toxicity[64].

Inhibitors of AddAB and RecBCD have been previously reported in the literature, but problems include limited *in vivo* stability, poor oral bioavailability and a sub-optimal mechanism of action[65–68]. However, Amundsen *et al.* (2012) identified several small-molecule inhibitors of *Helicobacter pylori* AddAB and *E. coli* RecBCD; in particular, ML328[69] indicating that the development of stable, potent inhibitors is possible. More recently, a derivative of ML328 (IMP-1700) was found to be capable of potentiating antibiotic activity such that a resistant *S. aureus* strain was sensitised to ciprofloxacin[70]. However, the ability of IMP-1700 to inhibit AddAB/RecBCD activity under *in vivo* conditions remains to be determined. Further work in this area may lead to broad-spectrum therapeutics that promote bacterial susceptibility to both host defences and antibiotics, and inhibit induction of the mutagenic SOS response.

In summary, staphylococcal, streptococcal and enterococcal DNA is damaged by the host immune system and several classes of bactericidal antibiotics leading to DSBs that are lethal if not repaired. This work demonstrates that DNA repair via RexAB is a common mechanism by which these pathogens repair damage caused by both threats and identifies this complex as a potential target for novel therapeutics, capable of sensitising Gram-positive pathogens to both neutrophils and antibiotics.

Materials and Methods

Bacterial strains and culture conditions

The bacterial strains used in this study are listed in Table 1. *S. aureus* was cultured in Tryptic Soy Broth (TSB) to stationary phase (18 h) at 37 °C, with shaking (180 rpm). *S. gordonii* and *E. faecalis* were grown in Todd Hewitt Broth (THB) supplemented with 1% (w/v) yeast extract (THB-Y) at 37 °C, statically in 5% CO_2. *E. coli* was grown in Lysogeny Broth (LB) or in Terrific Broth (TB) (1.2% (w/v) tryptone, 2.4% (w/v) yeast extract, 0.5% glycerol, 0.17 M KH_2PO_4, 0.72 M K_2HPO_4) for protein expression, at 37 °C with shaking (180 rpm). Media were supplemented with antibiotics as required. When appropriate, bacteria were grown on Columbia Blood Agar (CBA) made with 5% defibrinated sheep blood.

Table 1. Bacterial strains used in this study.

Strain	Description	Source
Staphylococcus aureus		
SH1000	*rsbU+* derivative of the laboratory strain 8325-4	36
SH1000 *rexB*::Tn	SH1000 with a *bursa aurealis* transposon insertion in *rexB,* Ery[r]	37
SH1000 *rexB*::Tn p*itet* empty	SH1000 with a *bursa aurealis* transposon insertion in *rexB* with integrated p*itet* empty plasmid, Ery[r]	37
SH1000 *rexB*::Tn p*itet rexAB*	SH1000 with a *bursa aurealis* transposon insertion in *rexB* with integrated p*itet* with AHT-inducible *rexB*, Ery[r]	37
JE2	A derivative of CA-MRSA USA300 LAC, cured of plasmids	35
JE2 *rexB*::Tn	JE2 with a *bursa aurealis* transposon insertion in *rexB*, Ery[r]	35
JE2 *rexB*::Tn p*itet* empty	JE2 with a *bursa aurealis* transposon insertion in *rexB* with integrated p*itet* empty plasmid, Ery[r]	37
JE2 *rexB*::Tn p*itet rexAB*	JE2 with a *bursa aurealis* transposon insertion in *rexB* with integrated p*itet* with AHT-inducible *rexB*, Ery[r]	37
SH1000 pCN34 *PrecA-gfp*	SH1000 containing pCN34 with *gfp* under the control of the *recA* promoter, Kan[r]	37
JE2 pCN34 *PrecA-gfp*	JE2 containing pCN34 with *gfp* under the control of the *recA* promoter, Kan[r]	37
JE2 *rexB*::Tn pCN34 *PrecA-gfp*	JE2 *rexB*::Tn containing pCN34 with *gfp* under the control of the *recA* promoter, Kan[r]	37
Streptococcus gordonii		
DL1 (Challis)	Wild type	71
DL1 *ΔrexBA*	DL1 with the *rexA* and *rexB* genes deleted	This study
Enterococcus faecalis		
CK111 (pCF10-101)	Conjugative donor strain with pCF10-101	72
OG1X	Gelatinase deficient	73
OG1X *ΔrexBA*	OG1X with the *rexA* and *rexB* genes deleted	This study
Escherichia coli		
EC1000	Cloning host that provides RepA in trans	74
SoluBL21 (DE3)	Derivative of BL21 (DE3) for the expression of challenging proteins	Genlantis

***recA-gfp* fluorescent reporter assay**

As detailed previously[37], promoter-reporter gene constructs in JE2 and SH1000 backgrounds were used to directly assess expression of *recA*. Antibiotic two-fold dilutions were made in flat-bottomed black-walled 96-well plates containing TSB and Kanamycin (90 μg ml^{-1}) and inoculated with 1/10 dilution of a stationary phase culture of the reporter strains. Plates were placed into an Infinite M200-

PRO microplate reader (Tecan) where cultures were grown for 17 h at 37 °C (700 rpm), and both

absorbance at 600 nm (OD_{600}) and GFP relative fluorescence units (RFU) were measured every 30 min.

OD_{600} data was normalised to no antibiotic controls and RFU data was normalised to no antibiotic

controls. To account for differences in cell-density, RFU values were normalised by OD_{600} data at each

time point.

Neutrophil phagocytosis and measurement of DNA damage

Whole human blood (15 ml) was collected from independent healthy donors in EDTA-treated tubes

(BD Biosciences) and layered over 20 ml of room-temperature Polymorph prep (Alere Limited) before

centrifugation at 500 x g for 45-60 min (brake off, 30 °C) until a clear separation of red blood cells

(RBCs), peripheral blood mononuclear cells (PBMCs) and polymophonuclear leukocytes (PMNs, or

neutrophils) was seen. The PBMCs were discarded and the PMNs were transferred to a fresh

centrifuge tube. Hanks' Balanced Salt Solution (HBSS) was added to the PMNs to a total volume of 50

ml and cells were pelleted at 500 x g for 10 min (brake off, 30 °C). The cells were resuspended in 3 ml

of HBSS, counted using a haemocytometer and adjusted to 5×10^6 cells ml^{-1} in HBSS containing 10%

human serum, 0.1 mM calcium and 0.1 mM magnesium. To the neutrophil suspension, 5×10^6 cells

ml^{-1} of bacteria (stationary or exponential-TRITC-stained) were added. The bacterial and neutrophil

suspension was then incubated at 37 °C with tumbling. At each time point (0.5, 1, 2, 3 h) 100-150 µl

was taken and resuspended in 4% PFA in PBS for a minimum of 1 h. Before analysis using flow

cytometry, samples were washed and resuspended in PBS. Samples were analysed on the FACS Aria

or LSRFortessa flow cytometer (BD Biosciences), and at least 100,000 events were captured, except

for bacteria-only samples where at least 50,000 events were captured. Green (GFP bacteria)

fluorescence was detected at 530 (30) nm, and TRITC labelling at 586 (16) nm. Full gating strategies

are detailed in Figures S6 and S7.

Determination of minimum inhibitory concentration (MIC)

Minimum inhibitory concentrations (MICs) were determined using a serial broth dilution protocol as

described previously[75]. Bacteria were diluted to 1×10^5 CFU ml^{-1} and incubated in flat-bottomed 96-

well plates with a range of antibiotic concentrations for 17 h at 37 °C under static conditions (aerobic,

anaerobic or 5% CO_2). Media containing daptomycin was supplemented with 1.25 mM $CaCl_2$. The MIC

was defined as the lowest concentration at which no growth was observed.

Antibiotic survival assay

Bacteria were adjusted to 10^8 CFU ml^{-1} in TSB (*S. aureus*) or THY broth (*S. gordonii* and *E. faecalis*) supplemented antibiotics at 10x MIC, unless otherwise stated. For aerobic incubation, 3 ml of media were inoculated in 30 ml universal tubes and incubated with shaking. For anaerobic conditions 6 ml of pre-reduced media in 7 ml bijou tubes was inoculated and incubated statically in an anaerobic cabinet. Cultures were incubated at 37 °C and bacterial viability determined by CFU counts. Culture media containing daptomycin was supplemented with 1.25 mM $CaCl_2$. Survival was calculated as a percentage of the number of bacteria in the starting inoculum.

Whole blood and hydrogen peroxide survival assays

Bacteria were washed twice with PBS, adjusted to 10^6 CFU ml^{-1} in HBSS and 10^4 CFU (10 µl) used to inoculate 90 µl of freshly donated human blood (collected in EDTA-treated tubes; BD Biosciences) or freshly diluted H_2O_2 (10 mM in PBS) in 96 well plates. Ethical approval for drawing and using human blood was obtained from the Regional Ethics Committee and Imperial NHS Trust Tissue Bank (REC Wales approval no. 12/WA/0196, ICHTB HTA license no. 12275). In some assays blood was pre-treated for 10 min with diphenyleneiodonium (DPI; 50 µM) or an equivalent volume of DMSO as a solvent control. After 6 h incubation, bacterial survival was determined by CFU counts of blood-bacteria mixtures on CBA plates. For H_2O_2 assays, survival was measured after 1 h at 37 °C (static) in the dark Survival for both assays was calculated as a percentage of the number of bacteria in the starting inoculum.

Neutrophil survival assay

Neutrophils were adjusted to 5×10^6 cells ml^{-1} in HBSS containing 10% human serum, 0.1 mM calcium and 0.1 mM magnesium. Stationary phase bacterial cultures were washed in PBS and 1×10^6 CFU was added to the neutrophil suspension (MOI 1:5) to a total volume of 1 ml. Neutrophils were treated for 10 min prior to the addition of bacteria with either diphenyleneiodonium (DPI; 50 µM) or an equivalent volume of DMSO (solvent control) as needed. The bacterial and neutrophil suspension was subsequently incubated at 37 °C with tumbling. At relevant time points (0.5, 1, 2, 3 h), 50 µl of the suspension was transferred to a 96-well plate and serially diluted ten-fold in PBS up to 10^{-3} dilution. All dilutions (including neat) were then plated onto CBA and incubated for 24 h at 37 °C before counting. Survival was calculated as a percentage of the number of bacteria in the starting inoculum.

Murine infection model

Animal work was conducted in accordance with the Animals (Scientific Procedures) Act 1986 outlined by the UK Home Office regulations. Work was approved by the UK Home Office after ethical approval by the Imperial College Animal Welfare and Ethical Review Body (AWERB). Six- to eight-week-old female C57BL/6 mice (Charles River) were infected via the intraperitoneal route with wild-type or *rexB*::Tn mutant in JE2 and SH1000 backgrounds. Stationary phase bacterial cultures were washed twice with PBS and adjusted to 10^7 CFU ml^{-1}. Subsequently, 400 µl (4 x 10^6 CFU) of the washed bacterial suspensions were injected into the peritoneal cavity of each mouse (5 mice for each strain, 20 in total). After 6 h, the mice were humanely sacrificed by cervical dislocation and death confirmed by severing the femoral artery. The peritoneal cavity was washed with PBS to release the bacteria and CFU counts were determined by plating onto TSA. Sample size was determined prior to the experiment using power analysis based on *in vitro* data [9]. Tubes containing the bacterial suspensions were blinded before starting the experiment. Mice were randomly allocated to group cages and each group was randomly allocated to a treatment. Following Home Office regulations, any animals that displayed two or more of the following symptoms were humanely killed using a Schedule 1 method and excluded from the study: shivering, hunched posture, reduced movement, cyanosis, circling or difficulty breathing.

Construction of Δ*rexBA* mutants of *E. faecalis* and *S. gordonii*

A Δ*rexBA* mutant was generated in *S. gordonii* by in-frame allelic replacement with the erythromycin resistance determinant *ermAM*. Flanking regions directly upstream and downstream of the *rexBA* operon were amplified by PCR from *S. gordonii* DL1 genomic DNA with primer pairs Sg.rexAB.F1/Sg.rexAB.R1 and Sg.rexAB.F2/Sg.rexAB.R2 (Table 2), respectively, while the *ermAM* cassette was amplified from plasmid pVA838[76] using primers ermAM.SgF/ermAM.SgR (Table 2). The resulting amplimers were then joined together via 20-bp overlapping regions by stitch PCR using primers Sg.rexAB.F1/Sg.rexAB.R2, and transformed into *S. gordonii*. Erythromycin-resistant transformants were confirmed by sequencing and the strain designated UB3018.

A *rexBA* mutant was generated in *E. faecalis* by markerless exchange using a 2-step homologous recombination approach, as previously described[72]. In brief, flanking regions directly upstream and downstream of the *rexBA* operon were amplified by PCR from *E. faecalis* OG1X genomic DNA with primer pairs Ef.rexAB.pheF1/Ef.rexAB.pheR1 and Ef.rexAB.pheF2/Ef.rexAB.pheR2 (Table 2), respectively, and then joined together via 20-bp overlapping regions by stitch PCR using primers Ef.rexAB.pheF1/Ef.rexAB.pheR2. The resultant amplimer was cloned into donor plasmid pCJK47[72] via unique restriction sites *Pst*I/*Nco*I to generate pCJK47-*rexAB*, propagated in *E. coli* EC1000 and then

introduced into conjugative donor strain *E. faecalis* CK111 (pCF10-101) by electroporation. Plasmid pCJK47-*rexAB* was transferred to *E. faecalis* OG1X by conjugation. Transconjugants carrying the integrated plasmid were confirmed by colony PCR, before counterselection based on the P-*pheS** marker was used to identify secondary recombinants in which the integrated plasmid had been excised and lost, leaving the desired Δ*rexBA* allele. This was confirmed by sequencing and the strain designated UB2948.

Construction of *S. aureus* RexAB expression vector

Cloning of the *rexA* and *rexB* genes from *S. aureus* was achieved by PCR from wild-type genomic JE2 DNA using the "rexB-F BamHI" and "rexA-R SalI" primers listed in Table 2, which allowed the amplification of the *rexBA* operon immediately flanked by suitable restriction endonuclease recognition sequences (BamHI and SalI). The *rexA* and *rexB* genes were inserted into the pET28b$^+$ vector (Novagen) using standard cloning techniques and site-directed mutagenesis (SDM) was performed to insert a StrepII-tag and a thrombin site in front of the *rexA* gene. This enabled RexA and RexB proteins to be detected individually, via an N-terminal His$_6$-tag for RexB (His$_6$ from the pET28b$^+$ vector) and an N-terminal StrepII-tag for RexA. PCR primers for SDM are listed in Table 2. DNA sequencing was used on the "pET28b$^+$ *rexBA*" expression plasmid to confirm that the sequences of the entire *rexA* and *rexB* genes, tags and promoter regions were as expected.

716 **Table 2. Primers used in this study.**

Oligonucleotide	Sequence 5'-3'
rexB-F BamHI	CCAGGATCCGATGACATTACATGCTTATTTAGG
rexA-R SalI	GCGGTCGACCTATAGTTGCAATGTACC
StrepII rexAB SDM-F	ACAATTCCAGAGAAACCACAAGGCGTGATTTGGACTGACGCGCAATGGC
StrepII rexAB SDM-R	AGCGCTTTTTTCGAACTGCGGGTGGCTCCACATCTATTGCTCACCCCC
Thr rexAB SDM-F	CCCGCAGTTCGAAAAAAGCGCTGGCCTGGTGCCGCGCGGCAGCGGCACAATTCCAGAGAAACCACAAGG
Thr rexAB SDM-R	CCTTGTGGTTTCTCTGGAATTGTGCCGCTGCCGCGCGGCACCAGGCCAGCGCTTTTTTCGAACTGCGGG
Sg.rexAB.F1	CAGCAGGACAAGGGAAGTG
Sg.rexAB.R1	<u>TTTTGTTCAT</u>AGCCTGCCTTTTCCTCTAG
Sg.rexAB.F2	<u>AGGTCCCTAG</u>AAGGAAGCATCTGAGTTG
Sg.rexAB.R2	CGTCGAGCACTAGTCTCG
ermAM.SgF	<u>AAGGCAGGCT</u>ATGAACAAAAATATAAAATATTCTCA
ermAM.SgR	<u>ATGCTTCCTT</u>CTAGGGACCTCTTTAGCTCC
Ef.rexAB.pheF1	TGACGTCGACGCGT*CTGCAG*GATCGCTAAAACGCTAGAAGC
Ef.rexAB.pheR1	<u>TAAAATCTAC</u>CTGCACACTCATGGTTTCAC
Ef.rexAB.pheF2	<u>GAGTGTGCAG</u>GTAGATTTTATAAAGTAGAAAAAATTAGAAG
Ef.rexAB.pheR2	GCTTAGCATG*CCATGG*TCTTAATACTTCGGTGATTGG

717 * Underlined regions indicate bp overlaps for stitch PCR; italicised regions indicate restriction

718 endonuclease sites for ligation into pCJK47

Expression and purification of recombinant *S. aureus* RexAB

Cells from single colonies of *E. coli* SoluBL21 (DE3), freshly transformed with pET28b$^+$ *rexAB* expression

plasmid coding for N-terminally StrepII-tagged RexA and N-terminally His$_6$-tagged RexB, were used to

inoculate an overnight starter culture in LB supplemented with 50 µg ml^{-1} of kanamycin. The starter

culture was diluted to an OD$_{600}$ of 0.05 into 4 L of TB containing 50 µg ml^{-1} of kanamycin. Cells were

grown at 37 °C with shaking at 180 rpm until an OD$_{600}$ of 0.5 was reached prior to induction with 1 mM

of isopropyl β-D-thiogalactoside (IPTG). Following induction, the temperature was reduced to 20 °C

and cultures were further incubated for 20 h. Cells were harvested by centrifugation at >10,000 x *g*

and 4 °C for 30 min and pellets were resuspended in 100 ml of 50 mM Tris, 150 mM NaCl (pH 7.5).

For protein purification, cells were disrupted by sonication and cell debris was cleared by

centrifugation at 32,000 x g 4 °C and for 30 min. The resulting supernatant was added to 5 ml of

Chelating Sepharose Fast Flow resin (GE Healthcare), which had been loaded with 0.1 M NiCl$_2$ and

equilibrated with 100 ml of 50 mM Tris, 150 mM NaCl, 20 mM imidazole (pH 7.5). The supernatant-

resin mixture was left at 4 °C overnight with gentle stirring to optimize the binding of the His-tagged

protein to the nickel-charged resin. Then, the mixture was washed 9x with 40 ml of 50 mM Tris, 150

mM NaCl, 20mM imidazole (pH 7.5) and 1x with 20 ml of 50 mM Tris, 150 mM NaCl, 70 mM imidazole

(pH 7.5). His-tagged protein was eluted with 50 mM Tris, 150 mM NaCl, 150 mM imidazole (pH 7.5);

each 10 ml fraction was tested with Bradford reagent (Bio-Rad) for protein content until no more protein could be detected. Fractions containing protein were pulled, buffer exchanged and concentrated using an Amicon 100 kDa cut-off concentrator (Merck Millipore); RexAB is >250 kDa in size. The concentrating device was centrifuged at 2000 x g (4 °C), and the protein solution was exchanged into 50 mM Tris, 150 mM NaCl (pH 7.5) by four serial concentration and re-dilution steps. The total protein concentration was quantified using the Pierce BCA protein assay kit (Thermo Fisher Scientific) following the manufacturer's instructions, and presence of intact recombinant RexAB was confirmed via SDS-PAGE and Western blot analysis.

Nuclease and helicase activity

The nuclease and helicase activity of RexAB was measured to confirm the AddAB-like activity in our recombinant RexAB protein. Staphylococcal DNA was amplified from the JE2 whole genome by colony PCR using the primer pairs "Chi control F" (5'- TCAGTGAATTAGATGATTCGC -3') and "Chi control R" (5'- TTCATACGTATGAATGTTATTTGC-3') where the amplicon lacked a Chi site region to be used as the DNA substrate in these assays.

Reactions were set up in either nuclease assay buffer (25 mM Tris-acetate pH 7.5, 2 mM Mg acetate, 1 mM DTT) or helicase assay buffer (25 mM Tris-acetate pH 7.5, 0.25 mM Mg acetate, 1 mM DTT), along with 5 ng μl^{-1} of DNA, 1 mM ATP or equivalent volume of nuclease-free water and either 20 nM or 50 nM of recombinant RexAB for the nuclease and helicase assay respectively. Additionally for the helicase assay, 2 µM of single-stranded binding proteins were added to each sample. The DMSO content was maintained at 1% (v/v) when compounds were being tested.

Samples were incubated statically at 37 °C and at 0, 5, 15, 30, 60 and 120 min, 5 µl was removed and pipetted into 20 µl of STEB buffer (40 % (w/v) sucrose, 100 mM Tris-HCl, 10 mM EDTA, 0.5 mg/ml bromophenol blue, pH 8) to stop the reaction. 20 µl of chloroform/isoamyl alcohol (24:1) was added to each tube, vortexed for 10 sec and centrifuged for 2 min at 17,000 x g to remove the protein and any compounds used for inhibition. The aqueous (upper blue) phase was loaded onto a 1% (w/v) agarose gel prepared in TBE buffer and electrophoresis was carried out at 85 V for 1 h. The gels were subsequently stained with SYBR Safe DNA gel stain (Invitrogen) at 1/10,000 dilution in TBE buffer for 2 h with rocking and visualised using a Gel Doc EZ Imager (Bio-Rad). Band intensity was quantified using ImageJ software. For nuclease activity, values were normalised to no-ATP controls at 0 h. For helicase activity, values were normalised to a ssDNA control that lacked the RexAB protein, which had been heated at 95 °C for 2 min to denature the dsDNA and allow the SSB protein to bind and stabilise the two ssDNA strands.

Statistical analyses

Data are represented as the mean or median from three or more independent experiments and analysed by Student's t test (two tailed, unpaired, assuming equal variances), one-way ANOVA or two-way ANOVA corrected for multiple comparison, as described in the figure legends. For each experiment, "n" refers to the number of independent biological replicates. CFU counts from murine experiments are presented as the value obtained from each animal and significance was assessed using the Mann-Whitney test. $P < 0.05$ was considered significant between data points (GraphPad Prism 7 for Windows).

Acknowledgements

A.M.E. and R.S.C. acknowledge funding from Shionogi & Co., Ltd. A.M.E. also acknowledges support from the National Institute for Health Research (NIHR) Imperial Biomedical Research Centre (BRC). K.P.H. is supported by a PhD scholarship funded by a Medical Research Council award to the Centre for Molecular Bacteriology and Infection (MR/J006874/1). D.A.I.M acknowledges funding from the Medical Research Council (Career Development Award MR/M009505/1). All authors acknowledge the provision of strains by the Network on Antimicrobial Resistance in Staphylococcus aureus (NARSA) Program: under NIAID/ NIH Contract No. HHSN272200700055C. The funders had no role in the study design, interpretation of the findings or the writing of the manuscript.

References

1. Tong SYC, Davis JS, Eichenberger E, Holland TL, Fowler VG. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev*. 2015;28(3):603-661.

2. Kreisel K, Boyd K, Langenberg P, Roghmann MC. Risk factors for recurrence in patients with *Staphylococcus aureus* infections complicated by bacteremia. *Diagn Microbiol Infect Dis*. 2006;55(3):179-184.

3. Rigby KM, DeLeo FR. Neutrophils in innate host defense against *Staphylococcus aureus* infections. *Semin Immunopathol*. 2012;34(2):237-259.

4. Buvelot H, Posfay-Barbe KM, Linder P, Schrenzel J, Krause K-H. *Staphylococcus aureus* , phagocyte NADPH oxidase and chronic granulomatous disease. Shen A, ed. *FEMS Microbiol Rev*. 2016;41(2): 139-157.

5. Ellson CD, Davidson K, Ferguson GJ, O'Connor R, Stephens LR, Hawkins PT. Neutrophils from
p40$^{phox-/-}$ mice exhibit severe defects in NADPH oxidase regulation and oxidant-dependent
bacterial killing. *J Exp Med*. 2006;203(8):1927-1937.

6. Hirsch J, Cohn Z. Degranulation of polymorphonuclear leucocytes following phagocytosis of
microorganisms. *J Exp Med*. 1960;112(6):1005-1014.

7. Babior BM. The respiratory burst oxidase. *Trends Biochem Sci*. 1987;12:241-243.

8. Fridovich I. Superoxide radical and superoxide dismutases. *Annu Rev Biochem*. 1995;64(1):97-
112.

9. Hampton MB, Kettle AJ, Winterbourn CC. Involvement of superoxide and myeloperoxidase in
oxygen-dependent killing of *Staphylococcus aureus* by neutrophils. *Infect Immun*.
1996;64(9):3512-3517.

10. Repine JE, Fox RB, Berger EM. Hydrogen peroxide kills *Staphylococcus aureus* by reacting with
staphylococcal iron to form hydroxyl radical. *J Biol Chem*. 1981;256(14):7094-7096.

11. Imlay JA, Chin SM, Linn S. Toxic DNA damage by hydrogen peroxide through the Fenton reaction
in vivo and *in vitro*. *Science*. 1988;240(4852):640-642.

12. Lanza F. Clinical manifestation of myeloperoxidase deficiency. *J Mol Med*. 1998;76(10):676-

13. Parry MF, Root RK, Metcalf JA, Delaney KK, Kaplow LS, Richar WJ. Myeloperoxidase deficiency.
Prevalence and clinical significance. *Ann Intern Med*. 1981;95(3):293-301.

14. Ben-Ari J, Wolach O, Gavrieli R, Wolach B. Infections associated with chronic granulomatous
disease: linking genetics to phenotypic expression. *Expert Rev Anti Infect Ther*. 2012;10(8):881-
894.

15. Pollock JD, Williams DA, Gifford MAC, et al. Mouse model of X–linked chronic granulomatous
disease, an inherited defect in phagocyte superoxide production. *Nat Genet*. 1995;9(2):202-209.

16. Gresham HD, Lowrance JH, Caver TE, Wilson BS, Cheung AL, Lindberg FP. Survival of
Staphylococcus aureus inside neutrophils contributes to infection. *J Immunol*. 2000;164(7):
3713-3722.

17. Thwaites GE, Gant V. Are bloodstream leukocytes Trojan Horses for the metastasis of
Staphylococcus aureus? *Nat Rev Microbiol*. 2011; 9(3):215-22.

18. Imlay JA. Pathways of Oxidative Damage. *Annu Rev Microbiol*. 2003;57(1):395-418.

19. Chang W, Small DA, Toghrol F, Bentley WE. Global transcriptome analysis of *Staphylococcus*
aureus response to hydrogen peroxide. *J Bacteriol*. 2006;188(4):1648-1659.

20. Michel B. After 30 years of study, the bacterial SOS response still surprises us. *PLoS Biol*.
2005;3(7):e255.

21. Cirz RT, Jones MB, Gingles NA, et al. Complete and SOS-mediated response of *Staphylococcus*

aureus to the antibiotic ciprofloxacin. *J Bacteriol*. 2007;189(2):531-539.

22. Fang FC. Antimicrobial actions of reactive oxygen species. mBio. 2011;2(5). pii: e00141-11.

23. Liu C, Bayer A, Cosgrove SE, et al. Clinical practice guidelines by the Infectious Diseases Society

of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults

and children. *Clin Infect Dis*. 2011;52(3):e18-e55.

24. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. A common mechanism of cellular

death induced by bactericidal antibiotics. *Cell*. 2007;130(5):797-810.

25. Dwyer DJ, Belenky PA, Yang JH, et al. Antibiotics induce redox-related physiological alterations

as part of their lethality. *Proc Natl Acad Sci U S A*. 2014;111(20).

26. Wang X, Zhao X. Contribution of oxidative damage to antimicrobial lethality. *Antimicrob Agents*

Chemother. 2009;53(4):1395-1402.

27. Shatalin K, Shatalina E, Mironov A, Nudler E. H_2S: A universal defense against antibiotics in

bacteria. *Science*. 2011;334(6058):986-990.

28. Calhoun LN, Kwon YM. The ferritin-like protein Dps protects *Salmonella enterica* serotype

Enteritidis from the Fenton-mediated killing mechanism of bactericidal antibiotics. *Int J*

Antimicrob Agents. 2011;37(3):261-265.

29. Foti JJ, Devadoss B, Winkler JA, Collins JJ, Walker GC. Oxidation of the guanine nucleotide pool

underlies cell death by bactericidal antibiotics. *Science*. 2012;336(6079):315-319.

30. Liu Y, Liu X, Qu Y, Wang X, Li L, Zhao X. Inhibitors of reactive oxygen species accumulation delay

and/or reduce the lethality of several antistaphylococcal agents. *Antimicrob Agents Chemother*.

2012;56(11):6048-6050.

31. Sampson TR, Liu X, Schroeder MR, Kraft CS, Burd EM, Weiss DS. Rapid killing of *Acinetobacter*

baumannii by polymyxins is mediated by a hydroxyl radical death pathway. *Antimicrob Agents*

Chemother. 2012;56(11):5642-5649.

32. Liu Y, Imlay JA. Cell death from antibiotics without the involvement of reactive oxygen species.

Science. 2013;339(6124):1210-3.

33. Keren I, Wu Y, Inocencio J, Mulcahy LR, Lewis K. Killing by bactericidal antibiotics does not

depend on reactive oxygen species. *Science*. 2013;339(6124):1213-6.

34. Imlay JA. Diagnosing oxidative stress in bacteria: not as easy as you might think. *Curr Opin*

Microbiol. 2015;24:124-31.

35. Fey PD, Endres JL, Yajjala VK, et al. A genetic resource for rapid and comprehensive phenotype

screening of nonessential *Staphylococcus aureus* genes. *MBio*. 2013;4(1):e00537-12.

36. Horsburgh MJ, Aish JL, White IJ, Shaw L, Lithgow JK, Foster SJ. sigmaB modulates virulence

determinant expression and stress resistance: characterization of a functional *rsbU* strain

derived from *Staphylococcus aureus* 8325-4. *J Bacteriol*. 2002;184(19):5457-67.

37. Clarke RS, Bruderer MS, Ha KP, Edwards AM. RexAB is essential for the mutagenic repair of DNA

damage caused by co-trimoxazole. *Antimicrob Agents Chemother*. 2019.

38. Liu GY, Essex A, Buchanan JT, et al. *Staphylococcus aureus* golden pigment impairs neutrophil

killing and promotes virulence through its antioxidant activity. *J Exp Med*. 2005;202(2):209-215.

39. Painter KL, Hall A, Ha KP, Edwards AM. The electron transport chain sensitizes *Staphylococcus*

aureus and *Enterococcus faecalis* to the oxidative burst. *Infect Immun*. 2017;85(12):e00659-17.

40. Malachowa N, Whitney AR, Kobayashi SD, et al. Global changes in *Staphylococcus aureus* gene

expression in human blood. Chakravortty D, ed. *PLoS One*. 2011;6(4):e18617.

doi:10.1371/journal.pone.0018617

41. Zuñiga-Castillo J, Romero D, Martínez-Salazar JM. The recombination genes *addAB* are not

restricted to Gram-positive bacteria: genetic analysis of the recombination initiation enzymes

RecF and AddAB in *Rhizobium etli*. *J Bacteriol*. 2004;186(23):7905-7913.

42. Rocha EPC, Cornet E, Michel B. Comparative and Evolutionary Analysis of the Bacterial

Homologous Recombination Systems. *PLoS Genet*. 2005;1(2):e15.

43. Chedin F, Kowalczykowski SC. A novel family of regulated helicases/nucleases from Gram-

positive bacteria: insights into the initiation of DNA recombination. *Mol Microbiol*.

2002;43(4):823-834.

44. Cromie GA. Phylogenetic ubiquity and shuffling of the bacterial RecBCD and AddAB

recombination complexes. *J Bacteriol*. 2009;191(16):5076-5084. doi:10.1128/JB.00254-09

45. Buscher K, Wang H, Zhang X, et al. Protection from septic peritonitis by rapid neutrophil

recruitment through omental high endothelial venules. *Nat Commun*. 2016;7:10828.

46. Wigley DB. Bacterial DNA repair: recent insights into the mechanism of RecBCD, AddAB and

AdnAB. *Nat Rev Microbiol*. 2013;11(1):9-13.

47. Drlica K. Mechanism of fluoroquinolone action. *Curr Opin Microbiol*. 1999;2(5):504-508.

48. Lee Y-J, Park S-J, Ciccone SLM, Kim C-R, Lee S-H. An *in vivo* analysis of MMC-induced DNA

damage and its repair. *Carcinogenesis*. 2006;27(3):446-453.

49. Yeeles JTP, Dillingham MS. A Dual-nuclease mechanism for DNA break processing by AddAB-

type helicase-nucleases. *J Mol Biol*. 2007;371(1):66-78.

50. Selton-Suty C, Célard M, Le Moing V, et al. Preeminence of *Staphylococcus aureus* in infective

endocarditis: A 1-year population-based survey. *Clin Infect Dis*. 2012;54(9):1230-1239.

51. Holland TL, Baddour LM, Bayer AS, Hoen B, Miro JM, Fowler VG. Infective endocarditis. *Nat Rev*

Dis Prim. 2016;2:16059.

52. Voyich JM, Braughton KR, Sturdevant DE, et al. Insights into mechanisms used by

Staphylococcus aureus to avoid destruction by human neutrophils. *J Immunol*.

2005;175(6):3907-3919.

53. Kobayashi SD, Braughton KR, Palazzolo-Ballance AM, et al. Rapid neutrophil destruction

following phagocytosis of *Staphylococcus aureus*. *J Innate Immun*. 2010;2(6):560-575.

54. Schlosser-Silverman E, Elgrably-Weiss M, Rosenshine I, Kohen R, Altuvia S. Characterization of

Escherichia coli DNA lesions enerated within J774 macrophages. *J Bacteriol*. 2000;182(18):5225-

5230.

55. Mertens K, Lantsheer L, Ennis DG, Samuel JE. Constitutive SOS expression and damage-inducible

AddAB-mediated recombinational repair systems for *Coxiella burnetii* as potential adaptations

for survival within macrophages. *Mol Microbiol*. 2008;69(6):1411-1426.

56. Miller C, Thomsen LE, Gaggero C, Mosseri R, Ingmer H, Cohen SN. SOS response induction by

beta-lactams and bacterial defense against antibiotic lethality. *Science*. 2004;305(5690):1629-

1631.

57. Maiques E, Ubeda C, Campoy S, et al. Beta-lactam antibiotics induce the SOS response and

horizontal transfer of virulence factors in *Staphylococcus aureus*. *J Bacteriol*. 2006;188(7):2726-

2729. doi:10.1128/JB.188.7.2726-2729.2006

58. Cuirolo A, Plata K, Rosato AE. Development of homogeneous expression of resistance in

methicillin-resistant *Staphylococcus aureus* clinical strains is functionally associated with a β-

lactam-mediated SOS response. *J Antimicrob Chemother*. 2009;64(1):37-45.

doi:10.1093/jac/dkp164

59. Gourley CR, Negretti NM, Konkel ME. The food-borne pathogen *Campylobacter jejuni* depends

on the AddAB DNA repair system to defend against bile in the intestinal environment. *Sci Rep*.

2017;7(1).

60. Amundsen SK, Fero J, Hansen LM, et al. *Helicobacter pylori* AddAB helicase-nuclease and RecA

promote recombination-related DNA repair and survival during stomach colonization. *Mol*

Microbiol. 2008;69(4):994-1007.

61. Cano DA, Pucciarelli MG, García-Del Portillo F, Casadesús J. Role of the RecBCD recombination

pathway in *Salmonella* virulence. *J Bacteriol*. 2002;184(2):592-595.

62. Standish AJ, Weiser JN. Human neutrophils kill *Streptococcus pneumoniae* via serine proteases. *J*

Immunol. 2009;183(4):2602-2609.

63. Richardson AR, Libby SJ, Fang FC. A nitric oxide-inducible lactate dehydrogenase enables

Staphylococcus aureus to resist innate immunity. *Science*. 2008;319(5870):1672-1676.

64. Jackson SP. Sensing and repairing DNA double-strand breaks. *Carcinogenesis*. 2002;23(5):687-

696.

65. Dziegielewska B, Beerman TA, Bianco PR. Inhibition of RecBCD Enzyme by Antineoplastic DNA
Alkylating Agents. *J Mol Biol*. 2006;361(5):898-919.

66. Karu AE, Linn S. Uncoupling of the *recBC* ATPase from DNase by DNA crosslinked with psoralen.
Proc Natl Acad Sci U S A. 1972;69(10):2855-2859.

67. Wilkinson M, Troman L, Wan Nur Ismah WA, et al. Structural basis for the inhibition of RecBCD
by Gam and its synergistic antibacterial effect with quinolones. *Elife*. 2016;5.

68. Sun D, Hurley LH. Structure-activity relationships of (+)-CC-1065 analogs in the inhibition of
helicase-catalyzed unwinding of duplex DNA. *J Med Chem*. 1992;35(10):1773-1782.

69. Amundsen SK, Spicer T, Karabulut AC, et al. Small-Molecule Inhibitors of Bacterial AddAB and
RecBCD Helicase-Nuclease DNA Repair Enzymes. *ACS Chem Biol*. 2012;7(5):879-891.

70. Lim CSQ, Ha KP, Clarke RS, et al. Identification of a potent small-molecule inhibitor of bacterial
DNA repair that potentiates quinolone antibiotic activity in methicillin-resistant *Staphylococcus*
aureus. *Bioorganic Med Chem*. 2019; 27(20):114962.

71. Pakula, R., and Walczak, W. On the nature of competence of transformable streptococci. *J Gen*
Microbiol. 1963;31:125–133.

72. Kristich CJ, Chandler JR, Dunny GM. Development of a host-genotype-independent
counterselectable marker and a high-frequency conjugative delivery system and their use in
genetic analysis of *Enterococcus faecalis*. Plasmid. 2007;57(2):131-44.

73. Ike Y, Craig RA, White BA, Yagi Y, Clewell DB. Modification of *Streptococcus faecalis* sex
pheromones after acquisition of plasmid DNA. *Proc Natl Acad Sci U S A*. 1983;80(17):5369-73.

74. Leenhouts K, Buist G, Bolhuis A, et al. A general system for generating unlabelled gene
replacements in bacterial chromosomes. *Mol Gen Genet*. 1996;253(1-2):217-24.

75. Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the minimal
inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc* 2008;3:163.

76. Macrina FL, Evans RP, Tobian JA, Hartley DL, Clewell DB, Jones KR. Novel shuttle plasmid
vehicles for *Escherichia-Streptococcus* transgeneric cloning. *Gene* 1983;25(1):145-50.

967
968
969