1 **Eye Movements Do Not Affect Perceived Size**

2 Paul Linton

4 Centre for Applied Vision Research, City, University of London

5 Northampton Square, Clerkenwell, London EC1V 0HB

6 paul@linton.vision

10   **Abstract**

Since Kepler (1604) and Descartes (1637), it's been suggested that 'vergence' (the angular rotation

of the eyes) plays a key role in size constancy (objects appearing to have a constant size despite

changes in distance radically altering the size of the retinal image). However, this has never been

tested divorced from confounding cues such as changes in the retinal image. In our experiment

participants viewed a target which grew or shrank over 5 seconds. At the same time their vergence

was increased from 50cm to 25cm. The question was whether this vergence increase biased the

participants' judgements of whether the target grew or shrank? We found no evidence of any bias,

and therefore no evidence that eye movements affect perceived size. This finding has three

important implications: First, perceived size is much more reliant on cognitive influences than

previously thought. This is consistent with the argument that visual scale is purely cognitive in

nature (Linton, 2017; 2018). Second, vergence modulation of V1 should no longer be thought of

as a candidate mechanism for size constancy. Third, the influence of multisensory integration on

visual size appears to be reliant on subjective knowledge about changes in hand and gaze position.

Introduction

As objects move forwards or backwards in space, the image they cast on the retina varies drastically in size. And yet objects don't appear to change dramatically in size when they move closer or further away. This suggests that there is a neural mechanism ('size constancy') that compensates for the drastic changes in the retinal image caused by changes in distance (for a review see Sperandio & Chouinard, 2015). We can distinguish between two kinds of visual cues for size constancy: 1. Pictorial Cues, which are present in the static monocular retinal image (such as familiar size and perspective), and which account for the impression of size constancy in pictures. And 2. Triangulation Cues, which typically rely on introducing multiple viewpoints either simultaneously (vergence and binocular disparities) or consecutively (motion parallax).

   1. Pictorial Cues: The neural correlates of size constancy are better understood for pictorial cues since 2D pictures are more easily presented to participants in fMRI scanners (e.g. Murray et al., 2006). However, pictorial cues are neither necessary nor sufficient for size constancy. First, pictorial cues are unnecessary because, as we shall discuss below, observers in the Taylor illusion appear to experience something close to full size constancy from vergence alone. Second, pictorial cues are insufficient because, as Sperandio et al. (2012) observe, size constancy in pictures is merely a fraction (10%-45%) of the full size constancy experienced in the real world (Emmert, 1881) (Murray et al., 2006; Leibowitz et al., 1969; Goodale, 2020 estimates 10%-30%; see also Millard et al., 2020 for a recent attempt to disambiguate static monocular and binocular size constancy, although recognising vergence may have influenced their monocular results).

   2. Triangulation Cues: In terms of triangulation cues to size constancy (vergence, binocular disparity, motion parallax), the emphasis has been on vergence. There are two reasons for this. First, motion parallax is neither necessary for size constancy (full size constancy is observed by participants in an fMRI scanner in Sperandio et al., 2012), nor is there particularly strong evidence that motion parallax contributes to size constancy (Combe & Wexler, 2010 only marginally qualify

"the common notion that size constancy emerges as a result of retinal and vergence processing

alone"). Second, binocular disparity is typically regarded as merely providing relative depth, and

not absolute size and distance. Still, considering size constancy only requires relative changes in

distance to be matched by relative changes in apparent size ($distance_1 \div distance_2 = size_1 \div size_2$),

a merely relative depth cue could suffice. The problem is that binocular disparity doesn't even

provide relative depth information until it has been scaled by absolute distance information, which

is typically assumed to come from vergence. As Brenner & van Damme (1998) observe, a 1°

change in retinal disparity could equally reflect a change in distance of 20cm to 21cm (5% increase

in distance) or 2m to 4m (100% increase in distance).

There is also a deeper conceptual point. Although Linton (2017; 2018) explores the

possibility that vision may be divorced from absolute size and distance, orthodox discussions of

size constancy typically articulate it in terms of the visual system using absolute distance to

determine perceived size. For instance, Emmert's Law (Emmert, 1881) is typically articulated as S

= c(R x D), where S is perceived size, c is a constant, R is retinal image size, and D is perceived

distance. But, as we already mentioned, binocular disparity is typically thought of as being a merely

relative depth cue outside very limited circumstances (objects taking up at least 20° of the visual

field; Rogers & Bradshaw, 1995). Instead, vergence is typically cited as being one of our most

important absolute distance cues at near distances (Mon-Williams & Tresilian, 1999; Viguier et al.,

2001; for a review and challenge to this consensus, see Linton, 2020).

Kepler (1604) and Descartes (1637) were the first to suggest that the visual system uses

vergence (the angular rotation of the eyes) to estimate distance, and scale the size of the retinal

image appropriately. For the last three hundred years, evidence for the role of vergence in size

constancy has come from four specific contexts where it has been found that changing the

vergence angle affects the perceived size of objects (so-called 'vergence micropsia'):

1. Wallpaper Illusion: Before the invention of the stereoscope by Wheatstone (1838), the

earliest evidence of vergence micropsia was the 'wallpaper illusion', the observation that if you

cross your eyes whilst looking at a recurring wallpaper pattern, the wallpaper pattern appears

smaller and closer (Smith, 1738; Priestley, 1772; Goethe, 1810; Meyer, 1842, 1852; Brewster, 1844;

Locke, 1849; Lie, 1965; Ono et al., 1971; Kohly & Ono, 2002; see Howard, 2012 for review).

2. Stereoscopic Viewing: The invention of the stereoscope by Wheatstone (1838) (which

presents separate images to each eye) enabled the eyes to be rotated independently of the retinal

image. Wheatstone observed that if eye rotation was increased, the perceived image appeared to

shrink, even though the images shown to each eye remained fixed (Wheatstone, 1852; Helmholtz,

1866, p.313; Judd, 1897; Frank, 1930; Hermans, 1937, 1954; Locke, 1938; Adams, 1955; Von Holst,

1955a, 1955b, 1957; Heinemann et al., 1959; Gogel, 1962; Biersdorf et al., 1963; Wallach &

Zuckerman, 1963; McCready, 1965; Leibowitz & Moore, 1966; Leibowitz et al., 1972; Komoda &

Ono, 1974; Regan et al., 1986; Enright, 1989).

3. Telestereoscopic Viewing: Building on Wheatstone (1838), Helmholtz (1857) invented

the telestereoscope, and observed that if we use mirrors to artificially increase the distance between

the two eyes, the world appears miniaturised. In his *Treatise on Physiological Optics* he observed that

"it will seem as if the observer were not looking at the natural landscape itself, but a very exquisite

and exact model of it, reduced in scale" (Helmholtz, 1866, p.312). This effect has been attributed

to vergence by Helmholtz (1857; 1858; 1866, p.310) and Rogers (2009; 2011), since the eyes need

to rotate more to fixate on the same physical distance (cf. Linton, 2018 for an alternative account),

and has been extensively studied in the military research (where helicopter pilots often view the

world through cameras with increased interpupillary separation, see Newman & Ostler, 2009;

Stuart et al., 2009; and Priot et al., 2010; 2011; 2012; 2018).

4. Taylor Illusion: Vergence is also thought to be central to the multisensory integration of

hand motion and the retinal image in the Taylor illusion. If you make an after-image of your hand

with a bright flash, and then in complete darkness move your hand closer to your face, the after-

image of your hand appears to shrink even though it is fixed on the retina (Taylor, 1941). The best

current explanation for the Taylor illusion is that it is due (Taylor, 1941; Morrison & Whiteside,

1984; Mon-Williams et al., 1997) or almost entirely due (Sperandio et al., 2013) to the increase in

vergence as the eyes track the physical hand in darkness (see also Gregory et al., 1959; Carey &

Allan, 1996; Bross, 2000; Ramsay et al., 2007; Faivre et al., 2017a; and for vergence scaling of after-

images see Urist, 1959; Suzuki, 1986; Lou, 2007; Zenkin & Petrov, 2015). Most notably, when

Sperandio et al. (2013) moved the participant's hand and vergence in opposite directions, they

found that (a) the after-image size changed in the direction of vergence, not the hand movement,

and (b) the magnitude of the size change when vergence and the hand were in conflict was almost

as large as when both the hand and vergence were moving in the same direction.

Surveying the literature on vergence micropsia, two things are striking: First, to our

knowledge, there has never been a report of a failure of vergence micropsia within peripersonal

space (near distances corresponding to arms reach). Even on the rare occasions when a change in

vergence fails to provide an impression of motion-in-depth (for instance, when motion-in-depth

is vetoed by a stimulus that takes up the whole visual field) as in Regan et al. (1986), the authors

still report "apparent size changes as about threefold when convergence changed from about 0

deg to 25 deg". And this is to be expected. As Regan et al. (1986) note: "Changes in size and depth

produced by ocular vergence changes are well known".

Second, the after-image literature appears to suggest that vergence provides something

close to perfect size constancy for distances between 25-50cm. This can be seen for two reasons:

First, because size constancy appears close to perfect for 25-50cm when vergence is the only

distance cue. Apparent size doubled for the representative subject in Sperandio et al. (2013)

(incongruent condition) from 3.3cm at 25cm (suggested by the $y = -0.61x + 3.3$ line of best fit) to

6.3cm at 50cm (average of size estimates after a >3° vergence eye movement) (my analysis of their

Fig.5 using WebPlotDigitizer 4.2; Marin et al., 2017). Second, the same conclusion is arrived at by

a combination of the fact that (a) the Taylor illusion provides near perfect size constancy in this

distance range (Bross, 2000; Ramsay et al., 2007; Sperandio et al., 2013), coupled with the fact that

(b) the Taylor illusion can be attributed almost entirely to vergence (Sperandio et al., 2013).

128   Vergence size constancy is therefore regarded as a fundamental aspect of visual perception.

129   However, we believe that vergence size constancy should be re-evaluated for two reasons:

130   First, our recent work suggests that vergence is an ineffective absolute distance cue once

131   confounding cues have been controlled for. Participants are unable to use vergence to judge

132   absolute distance (Linton, 2020), and we are reluctant to embrace the possibility (raised by Ono &

133   Comerford, 1977 and Bishop, 1989) that vergence might still be an effective size constancy cue

134   even if it proves to be ineffective for absolute distance judgements.

135   Second, one surprising fact is that to the best of our knowledge vergence size constancy

136   has never been tested divorced from confounding cues (such as changes in the retinal image or

137   changes in hand position) which inform the observer about changes in distance. The reason for

138   this is easy to appreciate. Vergence can only be driven in one of two ways. Either participants track

139   the retinal slip of a visual object moving in depth (such as an LED: Mon-Williams et al., 1997;

140   Sperandio et al., 2013), in which case participants are informed about the change in distance by

141   binocular disparity (the apparent retinal slip of the stimulus as it moves in depth). Or participants

142   track their own hand moving in depth (as in the Taylor illusion), but this gives them proprioceptive

143   information about the changing distance instead. The purpose of our experiment was therefore to

144   test vergence size constancy in a context where it is known to be effective (vergence changes over

145   5 seconds from 25cm to 50cm: Sperandio et al., 2013), but in a way that controls for subjective

146   knowledge about the changing distance.

 Materials and Methods

Subjects view a target over 5 seconds that...

...gets smaller — or — ...gets bigger

At the same time their vergence changes from 50cm to 25cm

25cm   50cm

And subjects are asked "Did the target get bigger or smaller?"
Question: Does the change in vergence bias their response?

Figure 1. Experimental Paradigm. Subjects viewed a target getting bigger or smaller over 5 seconds

whilst their vergence changed from 50cm to 25cm, and were asked to judge whether the target got

bigger or smaller? The question was whether vergence biased their response?

This posed a complex technical challenge which we resolved in five ways:

1. First, in order to drive vergence without providing subjective distance information, we

used a visual stimulus that (unlike an LED) provided 'sub-threshold' binocular disparities:

binocular disparities that are visible to the participant's visual system (in order to drive vergence),

but subjectively invisible to the participant themselves. This we achieved with a 3° target moving

in depth from 50cm to 25cm over 5 seconds. The target consisted of two targets on a display: a

left-hand target that only the right eye could see, and a right-hand target than only the left eye

could see. And by increasing the separation between the targets, we could increase the participant's

vergence angle (Fig.1). Rotating metal plates (Fig.3) ensured that each eye only saw the appropriate

target, and participants were asked to confirm that they saw a single fused target when they first

entered the apparatus, and to report if the target ever went double.

2. Second, in order to present a constant retinal image with eye rotation, we rendered the

targets to maintain a constant radius from, and orientation to, the eye. This was achieved in

OpenGL by 'projecting' the target onto the display, so that the correct retinal image was achieved

when the participants viewed the target (Fig.2) (camera set in OpenGL to nodal point of the eye,

and an asymmetric frustum was used so that the far clipping plane matched the distance and

dimensions of the display). A bite bar was used to ensure that the nodal point of the eye remained

fixed during the experiment (Fig.3), and the 6mm difference between the nodal point and the

centre of rotation of the eye was intentionally ignored (cf. Linton, 2019; Konrad et al., 2019).

Figure 2. OpenGL rendering of target achieves the correct retinal image for a target with a constant

radius and orientation to the eye, whilst presenting the target on a fronto-parallel display.

3. Third, another challenge of this display is that it requires the eyes to focus (or

'accommodate') at the distance of the display (160cm), whilst vergence (the angular rotation of the

eyes) is at 25-50cm. This decoupling of vergence and accommodation doesn't happen in normal

viewing conditions, and too much vergence-accommodation conflict can lead to the target going

blurry or double. To solve this problem we had an optometrist fit each participant with contact

lenses (based on the participant's valid UK prescription) so that the optical distance of the display

was 33cm even though its physical distance was 160cm. This ensured a maximum of $+/-1$

dioptres of vergence-accommodation conflict, well within the zone of 'clear single binocular

vision' (Hoffman et al., 2008). Indeed, some of the most dramatic reports of vergence micropsia

have been in the presence of large vergence-accommodation conflicts (e.g. 6.5 dioptres in Regan

et al., 1986), so the presence of $+/-1$ dioptre should not be objectionable.

4. Fourth, we wanted the target to be presented in darkness to exclude any residual visual

cues. An interesting finding from piloting was that the usual technique (having participants view a

CRTs through neutral density filters) wasn't effective in eradicating residual luminance from the

display (it degraded the target before residual luminance was completely eradicated). Instead, we

achieved this in four ways: First, we used an OLED display (LG OLED55C7V) which unlike

normal displays does not produce residual luminance for black pixels. Second, subjects wore a

mask to block out any residual light, which had red eye filters through which the red stimuli were

viewed (blocking out 100% green and ~90% blue light). Third, subjects viewed the stimuli through

a narrow (17°) viewing window of 48cm x 18cm at a distance of 60cm. Fourth, the whole apparatus

was covered by blackout fabric, and before the experiment began subjects pulled a hood of

blackout fabric over their heads and the external lights were turned off.

A. Apparatus

TOP VIEW

Blackout Fabric

Viewing Distance = 160cm

Eyes

48cm

Rotating Metal Plates

Display

SIDE VIEW

Blackout Fabric

Rotating Metal Plates

Eyes

BiteBar

18cm

29cm

Display

B. Simulated Experiment

Population Bias as % Size Change

Probability Density Function

Figure 3. Methods. A. Experimental Apparatus. B. Simulated Experiment. We simulated the experiment 10,000 times in Quest+ (bias = 0, detection threshold = 5%, lapse rate = 2%) to model how increasing the number of participants would improve the accuracy of our hierarchical Bayesian estimate of the true bias (true bias = 0). We determined that we needed n ≥ 5 to rule out an effect greater than our smallest effect size of interest (vergence size constancy > 1.5%).

5. Fifth, rather than ask participants to match their visual experience to (1) a visible chart (Bross, 2000; Lou, 2007; Sperandio et al., 2013) or (2) a memorised chart (Ramsay et al., 2007), or ask for conscious judgements of (3) the physical size of the after-image (Mon-Williams et al., 1997), or (4) the after-image's % size change (Carey & Allan, 1996), we (5) built size change estimation into the stimulus itself (something which cannot be done with an after-image). We increased or decreased the physical size of the target on each trial by between –20% and +20%, and asked the participants to make a forced choice ("did the target get bigger or smaller?"). And we tested

whether the change in vergence biased their response? If there was no vergence size constancy, then participants would be at chance in determining whether there was a size change when the target's physical size didn't change. By contrast, if participants experienced vergence micropsia, then we would have to increase the physical size of the target in order to cancel out this vergence micropsia for them to be at chance. And the degree of the bias (i.e. the amount that we have to increase physical size before participants are at chance) will indicate just how large the vergence micropsia effect size is.

We used a four-parameter maximum likelihood model (Quest+: Watson, 2017; Brainard, 2017) to estimate when participants were at chance. Participants completed 200 trials (10 sets of 20 trials), and on each trial Quest+ tested the size change that would be most informative. In piloting, we found that the author (an experienced psychophysical observer) could not detect size changes over 5 seconds that were smaller than 1.5%. So, if vergence changes the perceived size of the target by less than 1.5%, vergence size constancy can be dismissed as smaller than the smallest effect size of interest under an inferiority test (Lakens et al., 2018; in our actual experiment this was revised down to 1.43%, the detection threshold of our most sensitive observer).

Assuming vergence doesn't bias size judgements, how would the number of participants affect our ability to determine this fact? We can simulate the experiment 10,000 times (bias = 0, detection threshold = 5%, lapse rate = 2%), and fit a hierarchical Bayesian model (discussed below) to the data (Fig.3B). We found that with 5 or more observers we should be able to rule out any size constancy effects greater than 1.5%.

11 observers (8 female, 3 male; age ranges 20-34, average age 24.5) participated in the experiment: the author and 10 participants recruited using an online advertisement (13 were originally recruited, but 1 was excluded because they could not fuse the target, and 2 were excluded because their vision was blurry with the contact lenses). All participants were screened to ensure their accommodation was within normal bounds for their age (tested with a RAF near-point rule), vergence within normal bounds (18D or above on a Clement Clarke prism bar), and stereoacuity

within normal bounds (60 arc secs or less on a TNO stereo test). The author's participation was

required to (a) confirm Quest+ mirrored the pilot data, and (b) provide a criterion for the

minimum effect size. All other subjects were naïve as to the purpose of the experiment, and were

paid £15/hr for 3 hours. The study was approved by the School of Health Sciences Research

Ethics Committee at City, University of London in accordance with the Declaration of Helsinki.

The code for running the experiment is openly available: https://osf.io/5nwaz/ , running

on Matlab 2019a (MathWorks) with PsychToolBox 3.0.15 (Kleiner et al., 2007).

Results

Let us consider what we would expect to find according to (a) the null hypothesis (vergence has

no effect on perceived size) and (b) the alternative hypothesis (vergence has an effect on perceived

size). As we have already discussed, if vergence has no effect on perceived size, then it is obvious

that participants should be at chance at determining whether there is a size change just when we

don't introduce a size change (bias = 0). By contrast, if participants experience something close to

full size constancy, then we would have to increase the size of the target by 100% in order to

cancel out the reduction in perceived size caused by vergence micropsia (which in normal viewing

conditions would be a reduction in size of 50% because halving the distance leads to a doubling

of the retinal image size, assuming the small angle approximation).

These two hypotheses, and various intermediate degrees of size constancy between these

two extremes, are plotted in Fig.4A. What the hierarchical Bayesian model of our results in Fig.4A

show is that the bias we found was ≈ 0, consistent with there being no vergence size constancy.

Figure 4. Results. A. Hierarchical Bayesian model of the population psychometric function in black

(based on 15,000 posterior estimates, 100 representative posterior estimates in red), plotted against

predictions for various degrees of vergence size constancy effect sizes (in grey). B. Probability

density function of 15,000 posterior estimates of the population bias, with a non-significant bias

of –0.2%. C. Individual subject results fitted with Bayesian psychometric functions in black (based

on 15,000 posterior estimates, 100 representative posterior estimates in red). Blue dots indicating

the physical size changes tested by Quest+ (with darkness of the dot indicating the number of

times it was tested). Individual biases cluster around zero (from –2.2% to 1.2%). For each

participant, alpha (a) is the bias of the logistic function, and beta (b) is the slope.

To explain our conclusion, the individual results are plotted in Fig.4C. Each blue dot represents a

physical size change that was tested by the Quest+ maximum likelihood model, and the darkness

of the dot indicates the number of times it was tested. We then fit each of the individual sets of

data with a four-parameter logistic Bayesian psychometric function (indicated with a black line),

using the Palamedes Toolbox 1.10.1 (Prins & Kingdom, 2018) with CmdStan 2.22.0, using the toolbox's standard priors (bias and slope: normal (0,100), upper and lower lapse rates: beta (1,10)), and based on 15,000 posterior estimates (100 posterior estimates illustrated in red). Fig.4C shows individual biases range from –2.2% to +1.2%, but cluster around 0.

To estimate the population level psychometric function illustrated in Fig.4A, we used the Palamedes Toolbox 1.10.1 (Prins & Kingdom, 2018) with CmdStan 2.22.0 to fit a four-parameter logistic hierarchical Bayesian psychometric function, which fits the data with a multilevel model that takes into account the variability of each subject. We used the toolbox's standard multilevel priors which are documented by Prins & Kingdom (2019) and, based on 15,000 posterior estimates (100 posterior estimates illustrated in red), found a population level bias of –0.219% (95% CI: –1.82% to 1.39%) and a population level slope of –0.732 (95% CI: –1.07 to 0.378).

The estimate that particularly interests us is the population bias, so in Fig.4B we provide a probability density function of the 15,000 posterior estimates of the bias. We find no statistically significant bias, and therefore no statistically significant effect of vergence on perceived size. Indeed, the non-significant bias of –0.2% is in the wrong direction for size constancy.

To go beyond the negative claim that we found no statistically significant effect (null hypothesis not rejected) to the positive claim that there is no effect of vergence on perceived size (null hypothesis accepted), we can make two further arguments.

First, from a Bayesian perspective, we can perform a JZS Bayes factor (Rouder et al., 2009). The estimated Bayes factor that we found was 3.99 (±0.03%), which suggests that the data are four times more likely under the null hypothesis (bias = 0) than under the alternative (bias ≠ 0).

Second, from a frequentist perspective, we can perform an inferiority test that tests whether, if there is a vergence size constancy effect, it is at least as large as the smallest effect size of interest (Lakens et al., 2018). You'll remember, we defined our smallest effect size of interest as the detection threshold for our most sensitive observer (which is 1.43%). Put simply, any vergence size constancy effect that's smaller than a 1.43% size change won't be detected by any of our

observers. Since we have a directional hypothesis (vergence micropsia should reduce, rather than

increase, the apparent size of the target), we specifically tested whether there is a bias > 1.43%.

We therefore performed an inferiority test by taking the 90% confidence interval of the population

bias in Fig.4B in the predicted direction, which is 0.96%. Since this is smaller than 1.43% (our

smallest effect size of interest), from a frequentist perspective we can conclude that any vergence

size constancy effect is effectively equivalent to zero (Lakens et al., 2018).

Discussion

According to the literature, "it is well known that vergence is a reliable source of depth information

for size constancy" (Sperandio et al., 2013). But we find no evidence that vergence makes any

contribution to perceived size. To our knowledge, ours is the first study to report a failure of

vergence size constancy at near distances. But ours is also the first study that controls for

confounding perceptual cues (changes in the retinal image) whilst also controlling for confounding

cognitive cues (keeping subjects naïve about changes in absolute distance). Beyond vergence size

constancy, our results have three further important implications:

1. Visual Scale: First, these results substantiate a broader concern about visual scale. Visual

scale is thought to be provided by a number of well-established distance cues, such as vergence,

accommodation, motion parallax, familiar size, and the ground-plane. However, first,

accommodation (Mon-Williams & Tresilian, 2000) and motion parallax (Renner et al., 2013) have

been found to be largely ineffective as absolute distance cues, second, the ground-plane only

applies to limited viewing conditions (Creem-Regehr et al., 2015), and third, familiar size is merely

thought to affect our cognition, rather than our perception, of visual scale (Gogel, 1969; Predebon,

1992). Given these shortcomings, vergence was meant to provide a solid anchor for our size and

distance judgements in near space. But our results challenge this conclusion. Instead, our results

demonstrate that visual scale is much more reliant on cognitive influences than previously thought.

Our results are consistent with our argument in Linton (2017; 2018) that visual scale is based solely

on higher level cognitive processes, where we extend Gogel (1969)'s and Predebon (1992)'s

observations about familiar size to argue that visual scale itself is a purely cognitive process.

2. Neural Mechanisms: Ever since Trotter et al. (1992) found that the large majority of

neurons in the monkey primary visual context (V1) were modulated by vergence, it has been

suggested that processing of the vergence signal in V1 plays an important role in size constancy.

Further evidence for the vergence modulation of V1 is found by Trotter et al. (1993); Trotter et

al. (1996); Masson et al. (1997); Dobbins et al. (1998); Trotter & Celebrini (1999); Cumming &

Parker (1999); Trotter et al. (2004); Cottereau et al. (2014); see also Richards (1968) on LGN;

Gnadt & Mays (1991; 1995); Quinlan & Culham (2007); Culham et al. (2008) on the parietal cortex;

and Lehky et al. (1990) and Pouget & Sejnowski (1994) for an early neural network model that

Trotter et al. (1992) complements.

More recently Chen et al. (2019) have looked at the time course of size constancy, and

found that vergence and the retinal image are not integrated during (a) initial processing in V1

(~50ms), but instead during (b) recurrent processing within V1, and/or (c) re-entrant projections

from higher-order visual areas (e.g. Gnadt & Mays, 1991; 1995), both of which are consistent with

the ~150ms timeframe. And this is consistent with Trotter et al. (1992)'s suggestion that whilst

vergence responsive neurons encode vergence distance, further computations are required to scale

the retinal image, so vergence responsive neurons "constitute an intermediate step in the

computation of true depth, as suggested by neural network models [Lehky et al., 1990]."

However this whole line of research, from Trotter et al. (1992) to the present, is prefaced

on the fact that "psychophysical data suggest an important role for vergence" (Trotter et al., 1992).

But this is exactly what our results in this experiment, and in Linton (2020), question. We therefore

conclude that there is no link between the vergence modulation of neurons in V1 (or indeed

anywhere else) and size perception. According to our alternative account, visual scale is entirely

dependent upon top-down cognitive processing. Indeed, without the vergence signal we appear to

lose any early visual processing of absolute distance (one exception might be vertical disparities,

but Trotter et al., 1992 proceed on the basis that they are ineffective, and the subsequent evidence

in favour is equivocal at best: see Cumming et al., 1991; Sobel & Collett, 1991; Rogers & Bradshaw,

1995). So what are we to make of the results in Trotter et al. (1992)? One possibility is that they

reflect the participant's own purely cognitive knowledge about their changing gaze position, even

though this has no effect on their visual experience. This is consistent with increasing evidence

that V1 is implicated in purely cognitive (non-visual) processing, e.g. the location of rewards

amongst visually identical targets (Saleem et al., 2018).

3. Multisensory Integration: Subjective knowledge of gaze position also appears to play an

important role in multisensory integration. The Taylor illusion (where an after-image of the hand

appears to shrink or grow with physical hand movements) is an important paradigm for recent

discussions of multisensory integration (Faivre et al., 2017a; Grove et al., 2019). The best current

explanation for the Taylor illusion is that it is due (Taylor, 1941; Morrison & Whiteside, 1984;

Mon-Williams et al., 1997) or almost entirely due (Sperandio et al., 2013) to the change in vergence

as the eyes track the hand moving in darkness. However, in light of our results this explanation no

longer seems sustainable, since vergence had no effect on the perceived size of the target once

subjective knowledge about the fixation distance had been controlled for. Nor does this imply that

the Taylor illusion is primarily due to proprioceptive information from hand motion directly

influencing visual perception (Carey & Allan, 1996; Ramsay et al., 2007), since Sperandio et al.

(2013) demonstrate that when vergence and hand motion are in conflict, the Taylor illusion follows

vergence, and the effect is only marginally reduced in size.

Instead, what both accounts are missing is the participant's subjective knowledge about

their own changing hand and gaze positions. This explains why Sperandio et al. (2013) found that

vergence affects perceived size when their participants knew about their changing gaze position

(from their hand or from the motion in depth of an LED), but why we didn't when our participants

were ignorant of this fact. There are two ways in which conscious knowledge about our changing

hand or gaze position could influence size constancy. First, our subjective knowledge could

influence our visual experience (so-called 'cognitive penetration' of perception by cognition). But

we are skeptical of invoking 'cognitive penetration' to explain an effect that could also be explained

as a purely cognitive bias (for further skeptical discussions of 'cognitive penetration' see Fodor,

1983; Pylyshyn, 1999; Firestone & Scholl, 2016). Second, under our alternative cognitive bias

account, the integration of the retinal image and our changing gaze position could be purely

cognitive, rather than perceptual. Our visual experience of the after-image's angular size remains

constant, but our hand movements cognitively bias our interpretation of our constant visual

experience: because we know that our hand is moving towards our face, we interpret the constant

angular size of the after-image as a reduction in physical size.

This purely cognitive interpretation of the Taylor illusion has wide reaching implications

for multisensory integration, specifically the integration of vision and hand movements. The

Taylor illusion is taken as evidence of multisensory integration at the level of perception.

Specifically, that vision "relies on multimodal signals" (Sperandio et al., 2013; Chen et al., 2018)

and that "visual consciousness is shaped by the body" (Faivre et al., 2015; Faivre et al., 2017a;

Faivre et al., 2017b). But if the integration of proprioception and the retinal image could be purely

cognitive in the context of vergence (the major driver of the Taylor illusion, Sperandio et al., 2013),

there's no reason why the integration of proprioception and the retinal image in the context of

integrating vision and hand movements (the minor driver of the Taylor illusion, Sperandio et al.,

2013) couldn't equally be accounted for in purely cognitive terms. This cognitive approach also

suggests a non-perceptual explanation for variants of the Taylor illusion that appear to

demonstrate the integration of vision with the rubber-hand illusion (Faivre et al., 2017a) and tool

use (Grove et al., 2019). And cognitive interpretations of the integration of vision and

proprioception are also advanced in the contexts of vision and touch in slant estimation (Hillis et

al., 2002; Gepshtein et al., 2005) by Linton (2017), pp.37-38 and pp.65-66, and vision and vestibular

cues in self-motion (Fischer & Kornmüller, 1930; Ash et al., 2011) by Linton (2018).

Conclusions

Vergence is thought to provide an essential signal for size constancy. We tested vergence size constancy for the first time without confounding cues, and found no evidence that eye movements make any contribution to perceived size. This suggests that (1) our impression of visual scale is much more reliant on cognitive processing than previously thought, with the further implications that (2) vergence modulation of neurons in V1 cannot be responsible for our impression of visual scale, and (3) the integration of the retinal image with proprioceptive cues from the hand appears to be reliant upon observers having subjective knowledge about their hand and gaze position.

Acknowledgements

This research was conducted under the supervision of Christopher Tyler and Simon Grant. We thank Salma Ahmad, Mark Mayhew, and Deanna Taylor, for fitting the contact lenses, and Jugjeet Bansal, Priya Mehta, and the staff of the CitySight clinic for their assistance. We thank Joshua Solomon, Matteo Lisi, Byki Huntjens, Michael Morgan, Chris Hull, and Pete Jones for their advice. We also thank audiences at the Applied Vision Association (2019), the British Machine Vision Association's '3D worlds from 2D images in humans and machines' meeting (2020): https://youtu.be/6P3EYCEn52A and the (Virtual) Vision Sciences Society (2020): https://osf.io/tb3un/ / http://youtu.be/VhpYjPj5Q80 where this work was presented.

Open Practice Statement

Code for running the experiment, and all the data and analysis scripts, are accessible in an open access repository: https://osf.io/5nwaz/

References

Adams, O. S. (1955). Stereogram decentration and stereo-base as factors influencing the

apparent size of stereoscopic pictures. *The American Journal of Psychology*, *68*(1), 54–68.

Ash, A., Palmisano, S., & Kim, J. (2011). Vection in Depth during Consistent and Inconsistent

Multisensory Stimulation. *Perception*, *40*(2), 155–174. https://doi.org/10.1068/p6837

Biersdorf, W. R., Ohwaki, S., & Kozil, D. J. (1963). The Effect of Instructions and Oculomotor

Adjustments on Apparent Size. *The American Journal of Psychology*, *76*(1), 1–17. JSTOR.

https://doi.org/10.2307/1419994

Bishop, P. O. (1989). Vertical Disparity, Egocentric Distance and Stereoscopic Depth

Constancy: A New Interpretation. *Proceedings of the Royal Society of London. Series B, Biological*

Sciences, *237*(1289), 445–469.

Brainard, D. H. (2017). *mQUESTPlus: A Matlab implementation of QUEST+*.

https://github.com/brainardlab/mQUESTPlus

Brenner, E., & van Damme, W. J. (1998). Judging distance from ocular convergence. *Vision*

Research, *38*(4), 493–498.

Brewster, D. (1844). On the knowledge of distance given by binocular vision. *Transactions of the*

Royal Society of Edinburgh, *15*, 663–674.

Bross, M. (2000). Emmert's law in the dark: Active and passive proprioceptive effects on

positive visual afterimages. *Perception*, *29*(11), 1385–1391. https://doi.org/10.1068/p3038

Carey, D. P., & Allan, K. (1996). A motor signal and "visual" size perception. *Experimental Brain*

Research, *110*(3), 482–486. https://doi.org/10.1007/BF00229148

Chen, J., Sperandio, I., & Goodale, M. A. (2018). Proprioceptive Distance Cues Restore Perfect

Size Constancy in Grasping, but Not Perception, When Vision Is Limited. *Current Biology:*

CB, *28*(6), 927-932.e4. https://doi.org/10.1016/j.cub.2018.01.076

Chen, J., Sperandio, I., Henry, M. J., & Goodale, M. A. (2019). Changing the Real Viewing

Distance Reveals the Temporal Evolution of Size Constancy in Visual Cortex. *Current*

Biology, *29*(13), 2237-2243.e4. https://doi.org/10.1016/j.cub.2019.05.069

Combe, E., & Wexler, M. (2010). Observer Movement and Size Constancy. *Psychological Science*,

21(5), 667–675. JSTOR.

Cottereau, B., Durand, J.-B., Vayssière, N., & Trotter, Y. (2014). The influence of eye vergence

on retinotopic organization in human early visual cortex. *F1000Research*, *5*.

https://doi.org/10.7490/f1000research.1096674.1

Creem-Regehr, S. H., Stefanucci, J. K., & Thompson, W. B. (2015). Chapter Six - Perceiving

Absolute Scale in Virtual Environments: How Theory and Application Have Mutually

Informed the Role of Body-Based Perception. In B. H. Ross (Ed.), *Psychology of Learning*

and Motivation (Vol. 62, pp. 195–224). Academic Press.

https://doi.org/10.1016/bs.plm.2014.09.006

Culham, J., Gallivan, J., Cavina-Pratesi, C., & Quinlan, D. (2008). FMRI investigations of

reaching and ego space in human superior parieto-occipital cortex. In *Embodiment, Ego-*

space and Action (pp. 247–274).

Cumming, B. G., Johnston, E. B., & Parker, A. J. (1991). Vertical Disparities and Perception of

Three-Dimensional Shape. *Nature; London*, *349*(6308), 411–413.

Cumming, B. G., & Parker, A. J. (1999). Binocular Neurons in V1 of Awake Monkeys Are

Selective for Absolute, Not Relative, Disparity. *Journal of Neuroscience*, *19*(13), 5602–5618.

https://doi.org/10.1523/JNEUROSCI.19-13-05602.1999

Descartes, R. (1637). *Dioptrique (Optics). In J. Cottingham, R. Stoothoff, & D. Murdoch (Eds.), The*

Philosophical Writings of Descartes: Volume 1 (1985). Cambridge University Press.

Dobbins, A. C., Jeo, R. M., Fiser, J., & Allman, J. M. (1998). Distance Modulation of Neural

Activity in the Visual Cortex. *Science*, *281*(5376), 552–555.

https://doi.org/10.1126/science.281.5376.552

Emmert, E. (1881). Größenverhältnisse der Nachbilder. *Klinische Monatsblätter Für Augenheilkunde*

Und Für Augenärztliche Fortbildung, *19*, 443–450.

Enright, J. T. (1989). Enright JT (1989). "The eye, the brain and the size of the moon: Toward a

unified oculomotor hypothesis for the moon illusion". In Hershenson, Maurice (ed.).

The Moon illusion. Hillsdale, N.J: L. Erlbaum Associates. Pp. 59–121. In M.

Hershenson, *The Moon Illusion* (pp. 59–121). L. Erlbaum Associates.

Faivre, N., Arzi, A., Lunghi, C., & Salomon, R. (2017b). Consciousness is more than meets the

eye: A call for a multisensory study of subjective experience. *Neuroscience of Consciousness*,

2017(1). https://doi.org/10.1093/nc/nix003

Faivre, N., Dönz, J., Scandola, M., Dhanis, H., Ruiz, J. B., Bernasconi, F., Salomon, R., &

Blanke, O. (2017a). Self-Grounded Vision: Hand Ownership Modulates Visual Location

through Cortical β and γ Oscillations. *Journal of Neuroscience*, *37*(1), 11–22.

https://doi.org/10.1523/JNEUROSCI.0563-16.2016

Faivre, N., Salomon, R., & Blanke, O. (2015). Visual consciousness and bodily self-

consciousness. *Current Opinion in Neurology*, *28*(1), 23–28.

https://doi.org/10.1097/WCO.0000000000000160

Firestone, C., & Scholl, B. J. (2016). Cognition does not affect perception: Evaluating the

evidence for "top-down" effects. *The Behavioral and Brain Sciences*, *39*, e229.

https://doi.org/10.1017/S0140525X15000965

Fischer, M. H., & Kornmüller, A. E. (1930). Optokinetic-Induced Motion Perception and

Optokinetic Nystagmus. *Journal of Psychology and Neurology (Leipzig)*, *41*, 273–308.

Fodor, J. A. (1983). *The Modularity of Mind: An Essay on Faculty Psychology*. MIT Press.

Frank, H. (1930). Ueber den Einfluss inadaquater Konvergenz und Akkommodation auf die

Sehgrosse. *Psychol. Forsch.*, *13*, 135–144.

Gepshtein, S., Burge, J., Ernst, M. O., & Banks, M. S. (2005). The combination of vision and touch depends on spatial proximity. *Journal of Vision*, *5*(11), 7–7. https://doi.org/10.1167/5.11.7

Gnadt, J. W., & Mays, L. E. (1991). Depth tuning in area LIP by disparity and accommodative cues. *Soc Neurosci Abs*, *17*, 1113.

Gnadt, J. W., & Mays, L. E. (1995). Neurons in monkey parietal area LIP are tuned for eye-movement parameters in three-dimensional space. *Journal of Neurophysiology*, *73*(1), 280–297. https://doi.org/10.1152/jn.1995.73.1.280

Goethe, J. W. von. (1810). *Zur Farbenlehre*. in der J.G. Cotta'schen Buchhandlung.

Gogel, W. C. (1962). The Effect of Convergence on Perceived Size and Distance. *The Journal of Psychology*, *53*(2), 475–489. https://doi.org/10.1080/00223980.1962.9916591

Gogel, W. C. (1969). The Sensing of Retinal Size. *Vision Research*, *9*, 1079–1094.

Goodale, M. A. (2020). *How big is that banana? Differences in size constancy for perception and action*. https://drive.google.com/file/d/1LiOvNebMXRwAB6M53_XIc34vzP4qPWW9/view

Gregory, R. L., Wallace, J. G., & Campbell, F. W. (1959). Changes in the size and shape of visual after-images observed in complete darkness during changes of position in space. *Quarterly Journal of Experimental Psychology*, *11*(1), 54–55. https://doi.org/10.1080/17470215908416288

Grove, C., Cardinali, L., & Sperandio, I. (2019). Does Tool-Use Modulate the Perceived Size of an Afterimage During the Taylor Illusion? [Abstract ECVP Leuven 2019]. *Perception*, *48*, 181.

Heinemann, E. G., Tulving, E., & Nachmias, J. (1959). The effect of oculomotor adjustments on apparent size. *The American Journal of Psychology*, *72*, 32–45. https://doi.org/10.2307/1420209

Helmholtz, H. (1857). Das Telestereoskop. *Annalen Der Physik*, *178*(9), 167–175. https://doi.org/10.1002/andp.18571780907

Helmholtz, H. (1866). *Handbuch der Physiologischen Optik, Vol.III (translated by J. P. C. Southall 1925*

Opt. Soc. Am. Section 26, reprinted New York: Dover, 1962).

Helmholtz, P. H. (1858). II. On the telestereoscope. *The London, Edinburgh, and Dublin*

Philosophical Magazine and Journal of Science, 15(97), 19–24.

https://doi.org/10.1080/14786445808642433

Hermans, T. G. (1937). Visual size constancy as a function of convergence. *Journal of Experimental*

Psychology, 21(2), 145–161. https://doi.org/10.1037/h0058367

Hermans, Thomas G. (1954). The relationship of convergence and elevation changes to

judgments of size. *Journal of Experimental Psychology, 48*(3), 204–208.

https://doi.org/10.1037/h0059355

Hillis, J. M., Ernst, M. O., Banks, M. S., & Landy, M. S. (2002). Combining Sensory Information:

Mandatory Fusion Within, but Not Between, Senses. *Science, 298*(5598), 1627–1630.

https://doi.org/10.1126/science.1075396

Hoffman, D. M., Girshick, A. R., Akeley, K., & Banks, M. S. (2008). Vergence–accommodation

conflicts hinder visual performance and cause visual fatigue. *Journal of Vision, 8*(3), 33–33.

https://doi.org/10.1167/8.3.33

Howard, I. P. (2012). *Perceiving in Depth, Volume 3: Other Mechanisms of Depth Perception.* Oxford

University Press.

Judd, C. H. (1897). Some facts of binocular vision. *Psychological Review, 4*(4), 374–389.

https://doi.org/10.1037/h0073701

Kepler, J. (1604). Paralipomena to Witelo. In W. H. Donahue (Trans.), *Optics: Paralipomena to*

Witelo and Optical Part of Astronomy. Green Lion Press, 2000.

Kleiner, M., Brainard, D., & Pelli, D. (2007). Kleiner M, Brainard D, Pelli D, 2007, "What's new

in Psychtoolbox-3?" Perception 36 ECVP Abstract Supplement. *Perception, 36*(ECVP

Abstract Supplement).

Kohly, R. P., & Ono, H. (2002). Fixating on the wallpaper illusion: A commentary on "The role of vergence in the perception of distance: A fair test of Bishop Berkeley's claim" by Logvinenko et al. (2001). *Spatial Vision*, *15*(3), 377–386.

Komoda, M. K., & Ono, H. (1974). Oculomotor adjustments and size-distance perception. *Perception & Psychophysics*, *15*(2), 353–360. https://doi.org/10.3758/BF03213958

Konrad, R., Angelopoulos, A., & Wetzstein, G. (2019). Gaze-Contingent Ocular Parallax Rendering for Virtual Reality. *ArXiv:1906.09740 [Cs]*. http://arxiv.org/abs/1906.09740

Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in Psychological Science*, *1*(2), 259–269. https://doi.org/10.1177/2515245918770963

Lehky, S., Pouget, A., & Sejnowski, T. (1990). Neural Models of Binocular Depth Perception. *Cold Spring Harbor Symposia on Quantitative Biology*, *55*, 765–777. https://doi.org/10.1101/SQB.1990.055.01.072

Leibowitz, H., Brislin, R., Perlmutrer, L., & Hennessy, R. (1969). Ponzo Perspective Illusion as a Manifestation of Space Perception. *Science*, *166*(3909), 1174–1176. https://doi.org/10.1126/science.166.3909.1174

Leibowitz, H., & Moore, D. (1966). Role of changes in accommodation and convergence in the perception of size. *Journal of the Optical Society of America*, *56*(8), 1120–1129. https://doi.org/10.1364/JOSA.56.001120

Leibowitz, H. W., Shiina, K., & Hennessy, R. T. (1972). Oculomotor adjustments and size constancy. *Perception & Psychophysics*, *12*(6), 497–500. https://doi.org/10.3758/BF03210943

Lie, I. (1965). Convergence as a cue to perceived size and distance. *Scandinavian Journal of Psychology*, *6*(4), 109–116. https://doi.org/10.1111/j.1467-9450.1965.tb01059.x

Linton, P. (2017). *The Perception and Cognition of Visual Space*. Palgrave.

Linton, P. (2018). *Do We See Scale?* https://www.biorxiv.org/content/10.1101/371948v1

Linton, Paul. (2019). *Would Gaze-Contingent Rendering Improve Depth Perception in Virtual and*

Augmented Reality? https://arxiv.org/abs/1905.10366v1

Linton, Paul. (2020). Does vision extract absolute distance from vergence? *Attention, Perception, &*

Psychophysics, *82*(6), 3176–3195. https://doi.org/10.3758/s13414-020-02006-1

Locke, J. (1849). XXV. On single and double vision produced by viewing objects with both eyes;

and on an optical illusion with regard to the distance of objects. *The London, Edinburgh,*

and Dublin Philosophical Magazine and Journal of Science, *34*(228), 195–201.

https://doi.org/10.1080/14786444908646207

Locke, N. M. (1938). Some Factors in Size-Constancy. *The American Journal of Psychology*, *51*(3),

514–520. JSTOR. https://doi.org/10.2307/1416151

Lou, L. (2007). Apparent Afterimage Size, Emmert's Law, and Oculomotor Adjustment.

Perception, *36*(8), 1214–1228. https://doi.org/10.1068/p5513

Marin, F., Rohatgi, A., & Charlot, S. (2017). WebPlotDigitizer, a polyvalent and free software to

extract spectra from old astronomical publications: Application to ultraviolet

spectropolarimetry. *ArXiv:1708.02025 [Astro-Ph]*. http://arxiv.org/abs/1708.02025

Masson, G. S., Busettini, C., & Miles, F. A. (1997). Vergence eye movements in response to

binocular disparity without depth perception. *Nature*, *389*(6648), 283–286.

https://doi.org/10.1038/38496

McCready, D. W. (1965). Size-distance perception and accommodation-convergence

micropsia—A critique. *Vision Research*, *5*(4), 189–206. https://doi.org/10.1016/0042-

6989(65)90065-9

Meyer, H. (1842). Ueber einige Täuschungen in der Entfernung u. Grösse der Gesichtsobjecte.

Archiv Für Physiologische Heilkunde.

Meyer, Hermann. (1852). Ueber die Schätzung der Grösse und der Entfernung der

Gesichtsobjecte aus der Convergenz der Augenaxen. *Annalen Der Physik*, *161*(2), 198–

207. https://doi.org/10.1002/andp.18521610204

Millard, A. S., Sperandio, I., & Chouinard, P. A. (2020). The contribution of stereopsis in

614   Emmert's law. *Experimental Brain Research*, *238*(4), 1061–1072.

615   https://doi.org/10.1007/s00221-020-05747-5

616   Mon-Williams, M., & Tresilian, J. R. (1999). Some Recent Studies on the Extraretinal

617   Contribution to Distance Perception. *Perception*, *28*(2), 167–181.

618   https://doi.org/10.1068/p2737

619   Mon-Williams, M., & Tresilian, J. R. (2000). Ordinal depth information from accommodation?

620   *Ergonomics*, *43*(3), 391–404. https://doi.org/10.1080/001401300184486

621   Mon-Williams, M., Tresilian, J. R., Plooy, A., Wann, J. P., & Broerse, J. (1997). Looking at the

622   task in hand: Vergence eye movements and perceived size. *Experimental Brain Research*,

623   *117*(3), 501–506. https://doi.org/10.1007/s002210050246

624   Morrison, J. D., & Whiteside, T. C. (1984). Binocular cues in the perception of distance of a

625   point source of light. *Perception*, *13*(5), 555–566. https://doi.org/10.1068/p130555

626   Murray, S. O., Boyaci, H., & Kersten, D. (2006). The representation of perceived angular size in

627   human primary visual cortex. *Nature Neuroscience*, *9*(3), 429–434.

628   https://doi.org/10.1038/nn1641

629   Newman, D. G., & Ostler, D. (2009, July). *Hyperstereopsis Associated with Helmet-Mounted Sighting*

630   *and Display Systems for Helicopter Pilots* [Text]. Aerospace Medical Association.

631   https://doi.org/info:doi/10.3357/ASEM.21005.2009

632   Ono, H., & Comerford, J. (1977). Stereoscopic depth constancy. In W. Epstein (Ed.), *Stability*

633   *and constancy in visual perception: Mechanisms and process*. Wiley.

634   Ono, H., Mitson, L., & Seabrook, K. (1971). Change in convergence and retinal disparities as an

635   explanation for the wallpaper phenomenon. *Journal of Experimental Psychology*, *91*, 1–10.

636   https://doi.org/10.1037/h0031795

Pouget, A., & Sejnowski, T. J. (1994). A neural model of the cortical representation of egocentric

distance. *Cerebral Cortex (New York, N.Y.: 1991)*, *4*(3), 314–329.

https://doi.org/10.1093/cercor/4.3.314

Predebon, J. (1992). The role of instructions and familiar size in absolute judgments of size and

distance. *Perception & Psychophysics*, *51*(4), 344–354. https://doi.org/10.3758/BF03211628

Priestley, J. (1772). *The History and Present State of Discoveries Relating to Vision, Light, and Colours*. J.

Johnson.

Prins, N., & Kingdom, F. A. A. (2018). Applying the Model-Comparison Approach to Test

Specific Research Hypotheses in Psychophysical Research Using the Palamedes Toolbox.

Frontiers in Psychology, *9*. https://doi.org/10.3389/fpsyg.2018.01250

Prins, N., & Kingdom, F. A. A. (2019).

http://www.palamedestoolbox.org/hierarchicalbayesian.html

Priot, A. E., Neveu, P., Philippe, M., & Roumes, C. (2012). Adaptation to alterations of apparent

distance in stereoscopic displays: From lab to hyperstereopsis. *2012 International Conference*

on 3D Imaging (IC3D), 1–7. https://doi.org/10.1109/IC3D.2012.6615123

Priot, A.-E., Laboissière, R., Plantier, J., Prablanc, C., & Roumes, C. (2011). Partitioning the

components of visuomotor adaptation to prism-altered distance. *Neuropsychologia*, *49*(3),

498–506. https://doi.org/10.1016/j.neuropsychologia.2010.11.028

Priot, A.-E., Laboissière, R., Sillan, O., Roumes, C., & Prablanc, C. (2010). Adaptation of

egocentric distance perception under telestereoscopic viewing within reaching space.

Experimental Brain Research, *202*(4), 825–836. https://doi.org/10.1007/s00221-010-2188-4

Priot, A.-E., Vacher, A., Vienne, C., Neveu, P., & Roumes, C. (2018). The initial effects of

hyperstereopsis on visual perception in helicopter pilots flying with see-through helmet-

mounted displays. *Displays*, *51*, 1–8. https://doi.org/10.1016/j.displa.2017.11.002

Pylyshyn, Z. (1999). Is vision continuous with cognition?: The case for cognitive impenetrability of visual perception. *Behavioral and Brain Sciences*, *22*(3), 341–365. https://doi.org/10.1017/S0140525X99002022

Quinlan, D. J., & Culham, J. C. (2007). FMRI reveals a preference for near viewing in the human parieto-occipital cortex. *NeuroImage*, *36*(1), 167–187. https://doi.org/10.1016/j.neuroimage.2007.02.029

Ramsay, A. I. G., Carey, D. P., & Jackson, S. R. (2007). Visual-proprioceptive mismatch and the Taylor illusion. *Experimental Brain Research*, *176*(1), 173–181. https://doi.org/10.1007/s00221-006-0609-1

Regan, D., Erkelens, C. J., & Collewijn, H. (1986). Necessary conditions for the perception of motion in depth. *Investigative Ophthalmology & Visual Science*, *27*(4), 584–597.

Renner, R. S., Velichkovsky, B. M., & Helmert, J. R. (2013). The Perception of Egocentric Distances in Virtual Environments—A Review. *ACM Comput. Surv.*, *46*(2), 23:1–23:40. https://doi.org/10.1145/2543581.2543590

Richards, W. (1968). Spatial remapping in the primate visual system. *Kybernetik*, *4*(4), 146–156. https://doi.org/10.1007/BF00288548

Rogers, B. (2009). Are stereoscopic cues ignored in telestereoscopic viewing? *Journal of Vision*, *9*(8), 288–288. https://doi.org/10.1167/9.8.288

Rogers, B. J. (2011). Information, illusion, and constancy in telestereoscopic viewing. In L. R. Harris & M. R. M. Jenkin, *Vision in 3D Environments*. Cambridge University Press.

Rogers, B. J., & Bradshaw, M. F. (1995). Disparity Scaling and the Perception of Frontoparallel Surfaces. *Perception*, *24*(2), 155–179. https://doi.org/10.1068/p240155

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*, *16*(2), 225–237. https://doi.org/10.3758/PBR.16.2.225

Saleem, A. B., Diamanti, E. M., Fournier, J., Harris, K. D., & Carandini, M. (2018). Coherent

encoding of subjective spatial position in visual cortex and hippocampus. *Nature*,

562(7725), 124–127. https://doi.org/10.1038/s41586-018-0516-1

Smith, R. (1738). *A Compleat System of Opticks in Four Books, Viz. A Popular, a Mathematical, a*

Mechanical, and a Philosophical Treatise. To which are Added Remarks Upon the Whole. By Robert

Smith. author, and sold there.

Sobel Erik C., & Collett Thomas Stephen. (1991). Does vertical disparity scale the perception of

stereoscopic depth? *Proceedings of the Royal Society of London. Series B: Biological Sciences*,

244(1310), 87–90. https://doi.org/10.1098/rspb.1991.0055

Sperandio, I., & Chouinard, P. A. (2015). The Mechanisms of Size Constancy. *Multisensory*

Research, *28*(3–4), 253–283. https://doi.org/10.1163/22134808-00002483

Sperandio, I., Chouinard, P. A., & Goodale, M. A. (2012). Retinotopic activity in V1 reflects the

perceived and not the retinal size of an afterimage. *Nature Neuroscience*, *15*(4), 540–542.

https://doi.org/10.1038/nn.3069

Sperandio, I., Kaderali, S., Chouinard, P. A., Frey, J., & Goodale, M. A. (2013). Perceived size

change induced by nonvisual signals in darkness: The relative contribution of vergence

and proprioception. *The Journal of Neuroscience: The Official Journal of the Society for*

Neuroscience, *33*(43), 16915–16923. https://doi.org/10.1523/JNEUROSCI.0977-13.2013

Stuart, G., Jennings, S., Kalich, M., Rash, C., Harding, T., & Craig, G. (2009). Flight performance

using a hyperstereo helmet-mounted display: Adaptation to hyperstereopsis. *Proceedings of*

SPIE - The International Society for Optical Engineering, *7326*.

https://doi.org/10.1117/12.820500

Suzuki, K. (1986). Effects of oculomotor cues on the apparent size of afterimages. *Japanese*

Psychological Research, *28*(4), 168–175. https://doi.org/10.4992/psycholres1954.28.168

Taylor, F. V. (1941). Change in size of the afterimage induced in total darkness. *Journal of*

Experimental Psychology, *29*(1), 75–80. https://doi.org/10.1037/h0058125

Trotter, Y., Celebrini, S., Stricanne, B., Thorpe, S., & Imbert, M. (1992). Modulation of neural

stereoscopic processing in primate area V1 by the viewing distance. *Science*, *257*(5074),

1279–1281. https://doi.org/10.1126/science.1519066

Trotter, Y., Celebrini, S., Stricanne, B., Thorpe, S., & Imbert, M. (1996). Neural processing of

stereopsis as a function of viewing distance in primate visual cortical area V1. *Journal of*

Neurophysiology, *76*(5), 2872–2885. https://doi.org/10.1152/jn.1996.76.5.2872

Trotter, Yves, & Celebrini, S. (1999). Trotter, Y. & Celebrini, S. Gaze direction controls response

gain in primary visual-cortex neurons. Nature 398, 239−242. *Nature*, *398*, 239–242.

https://doi.org/10.1038/18444

Trotter, Yves, Celebrini, S., & Durand, J. B. (2004). Evidence for implication of primate area V1

in neural 3-D spatial localization processing. *Journal of Physiology-Paris*, *98*(1), 125–134.

https://doi.org/10.1016/j.jphysparis.2004.03.004

Trotter, Yves, Stricanne, B., Celebrini, S., Thorpe, S., & Imbert, M. (1993). *Neural processing of*

stereopsis as a function of viewing distance. Oxford University Press. https://0-www-

oxfordscholarship-

com.wam.city.ac.uk/view/10.1093/acprof:oso/9780198547853.001.0001/acprof-

9780198547853-chapter-027

Urist, M. J. (1959). Afterimages and ocular muscle proprioception. *A.M.A. Archives of*

Ophthalmology, *61*(2), 230–232. https://doi.org/10.1001/archopht.1959.00940090232006

Viguier, A., Clément, G., & Trotter, Y. (2001). Distance perception within near visual space.

Perception, *30*(1), 115–124. https://doi.org/10.1068/p3119

Von Holst, E. (1955b). 1st der Einfluss der Akkommodation auf die gesehene Dinggrijsse ein

"reflektorischer" Vorgang? *Naturwissenschaften*, *42*, 445–446.

Von Holst, E. (1955a). Die Beteiligung von Konvergenz und Akkommodation an der

wahrgenommenen Grossenkonstanz. *Naturwissenschaften*, *42*, 444–445.

Von Holst, E. (1957). Aktive Leistungen der menschlichen Gesichtswahrnemung. *Studium*

Generale, *10*, 231–243.

Wallach, H., & Zuckerman, C. (1963). The constancy of stereoscopic depth. *The American Journal*

of Psychology, *76*, 404–412.

Watson, A. B. (2017). QUEST+: A general multidimensional Bayesian adaptive psychometric

method. *Journal of Vision*, *17*(3), 10–10. https://doi.org/10.1167/17.3.10

Wheatstone, C. (1838). XVIII. Contributions to the physiology of vision. —Part the first. On

some remarkable, and hitherto unobserved, phenomena of binocular vision. *Philosophical*

Transactions of the Royal Society of London, *128*, 371–394.

https://doi.org/10.1098/rstl.1838.0019

Wheatstone, C. (1852). I. The Bakerian Lecture.—Contributions to the physiology of vision.—

Part the second. On some remarkable, and hitherto unobserved, phenomena of

binocular vision (continued). *Philosophical Transactions of the Royal Society of London*, *142*, 1–

17. https://doi.org/10.1098/rstl.1852.0001

Zenkin, G. M., & Petrov, A. P. (2015). Transformation of the Visual Afterimage Under Subject's

Eye and Body Movements and the Visual Field Constancy Mechanisms. *Perception*, *44*(8–

9), 973–985. https://doi.org/10.1177/0301006615594937